Characterization of cellular, biochemical and genomic features of the diazotrophic plant growth-promoting bacterium *Azospirillum* sp. UENF-412522, a novel member of the *Azospirillum* genus

Gustavo L. Rodrigues[a,*], Filipe P. Matteoli[a,*], Rajesh K. Gazara[a], Pollyanna S. L. Rodrigues[b], Samuel T. dos Santos[b], Alice F. Alves[b,c], Francisnei Pedrosa-Silva[a], Isabella Oliveira-Pinheiro[a], Daniella Canedo-Alvarenga[a], Fabio L. Olivares[b,c,#], Thiago M. Venancio[a,#]

[a] Laboratório de Química e Função de Proteínas e Peptídeos, Centro de Biociências e Biotecnologia, Universidade Estadual do Norte Fluminense Darcy Ribeiro (UENF), Brazil; [b] Núcleo de Desenvolvimento de Insumos Biológicos para a Agricultura (NUDIBA), UENF, Brazil; [c] Laboratório de Biologia Celular e Tecidual, Centro de Biociências e Biotecnologia, UENF, Brazil. [*] Contributed equally to this work.

[#] **Corresponding authors:**
Thiago M. Venancio; Laboratório de Química e Função de Proteínas e Peptídeos, Centro de Biociências e Biotecnologia, UENF; Av. Alberto Lamego 2000, P5 / sala 217; Campos dos Goytacazes, Rio de Janeiro, Brazil. E-mail: thiago.venancio@gmail.com.

Fabio L. Olivares: Laboratório de Biologia Celular e Tecidual, Centro de Biociências e Biotecnologia, UENF, Brazil. E-mail: fabioliv@uenf.br.

ABSTRACT
Given their remarkable beneficial effects on plant growth, several *Azospirillum* isolates currently integrate the formulations of various commercial inoculants. Our research group isolated a new strain, *Azospirillum* sp. UENF-412522, from passion fruit rhizoplane. This isolate uses carbon sources that are partially distinct from closely-related *Azospirillum* isolates. Scanning electron microscopy analysis and population counts demonstrate the ability of *Azospirillum* sp. UENF-412522 to colonize the surface of passion fruit roots. *In vitro* assays demonstrate the ability of *Azospirillum* sp. UENF-412522 to fix atmospheric nitrogen, to solubilize phosphate and to produce indole-acetic acid. Passion fruit plantlets inoculated with *Azospirillum* sp. UENF-41255 showed increased shoot and root fresh matter, as well as root dry matter, further highlighting its biotechnological potential for agriculture. We sequenced the genome of *Azospirillum* sp. UENF-412522 to investigate the genetic basis of its plant-growth promotion properties. We identified the key *nif* genes for nitrogen fixation, the complete PQQ operon for phosphate solubilization, the *acdS* gene that alleviates ethylene effects on plant growth, and the *napCAB* operon, which produces nitrite under anoxic conditions. We also found several genes conferring resistance to common soil antibiotics, which are critical for *Azospirillum sp.* UENF-412522 survival in the rhizosphere. Finally, we also assessed the *Azospirillum* pangenome and highlighted key genes involved in plant growth promotion. A phylogenetic reconstruction of the genus was also conducted. Our results support *Azospirillum sp.* UENF-412522 as a good candidate for bioinoculant formulations focused on plant growth promotion in sustainable systems.

INTRODUCTION

Modern agriculture production strongly relies on synthetic fertilizers and pesticides to achieve high productivity levels, which are often a matter of environmental and health concerns [1]. The development of bioinoculants has been considered an alternative to reduce the use of such synthetic compounds [2]. These inoculants typically have one or more strains of plant growth-promoting rhizobacteria (PGPR), which compose a heterogeneous group of bacteria that exert mutualistic interactions with plants [3], often playing key roles in the root microbiome [4]. The rhizosphere is the fraction of soil directly influenced by roots exudates [5], which modulate bacterial diversity across the endorhizosphere, rhizoplane and ectorhizhosphere (reviewed by Ahkami et al. [6]). Several *Azospirillum* isolates have been reported as PGPR, typically as soil-borne bacteria that are competent rhizosphere colonizers [7, 8].

The genus *Azospirillum* belongs to the *Rhodospirillaceae* family, which is mainly constituted of aquatic genera. Nevertheless, Azospirilla are mostly soil bacteria that coevolved with vascular plants [9]. Following its original description [10], multiple plant-associated *Azospirillum* species have been shown to perform direct nitrogen fixation [11, 12], phosphate solubilization [13], drought and salt stress alleviation [14, 15], root development promotion [16], among other processes [17]. These desirable properties have led several *Azospirillum* strains to be used as part of commercial soil inoculants [18]. Genes associated with these features have been identified in several publicly available *Azospirillum* genomes, such as *nif* [19], ACC-deaminase (*acds*) [20, 21], *pqq* [22], and indole acetic acid biosynthesis genes (e.g., *iaaH*, *iaaM* and *ipdC*) [23, 24].

Recently, our research group has isolated several plant growth-promoting bacterial strains from vermicompost [25, 26] and from the rhizosphere of tropical fruit trees. Among these strains, *Azospirillum sp.* UENF-412522 was isolated from passion fruit rhizoplane. Here we report *in vitro* and *ex vitro* plant growth promotion capabilities, biochemical tests, microscopy analysis of root colonization, whole-genome sequencing, and comparative genomic analyses to investigate the plant growth promotion properties of this isolate, as well as to uncover genes involved in other ecophysiological processes. Together, our results support the potential of *Azospirillum sp.* UENF-412522 to be used in bioinoculant formulations.

RESULTS AND DISCUSSION

Identification of the isolate and assessment of plant growth promotion properties

The strain UENF-412522 was isolated from passion fruit rhizoplane and formed a pellicle typical of diazotrophic bacteria on N-free semisolid NFb medium. It is a Gram negative, slightly curved rod-shaped bacterium with apparently no polymorphic 1.2 x 0.7 μm cells. In order to characterize the isolated strain, a fragment of the 16S rDNA gene was amplified by polymerase chain reaction (PCR); this sequence was deposited in Genbank under the accession KU836626.1. The 16S rDNA maximum likelihood (ML) phylogenetic reconstruction (Figure S1) has not allowed us to classify the strain at the species level. Nevertheless, it is clear that it belongs to the *Azospirillum* genus, leading us to name it *Azospirillum sp.* UENF-412522.

Given the evidence supporting *Azospirillum* sp. UENF-412522 as a novel *Azospirillum* species, we used an API 50 CH system and found that it grows on 18 out of 49 tested carbon sources: glycerol, D-arabinose, L-arabinose, D-ribose, D-xylose, methyl-beta-D-xylopyranoside, D-galactose, D-glucose, D-fructose, D-mannose, L-sorbose, amygdalin, aesculin ferric citrate, D-

melibiose, glycogen, xylitol, D-xylose, D-fucose. On the other hand, this strain was unable to

grow on: erythritol, L-xylose, D-adonitol, L-rhamnose, dulcitol, inositol, D-mannitol, D-sorbitol,

Methyl-alpha-D-mannopyranoside,   Methyl-alpha-D-glucopyranoside,   N-acetylglucosamine,

arbutin, salicin, D-cellobiose, D-maltose, D-lactose (bovine origin), D-saccharose (sucrose), D-

trehalose, inulin, D-melezitose, D-raffinose, amidon (starch), gentiobiose, D-turanose, D-

tagatose, L-fucose, D-arabitol, L-arabitol, potassium gluconate, potassium 2-ketogluconate,

potassium 5-ketogluconate. Comparison of carbon utilization patterns of *Azospirillum* sp. UENF-

412522 and those of the type strains of *Azospirillum lipoferum* ATCC 29707 [10], *Azospirillum*

doebereinerae DSM 13131 [27] and *Azospirillum brasilense* ATCC 29145 showed that four C-

sources (i.e. D-ribose, D-mannitol, D-sorbitol and N-acetylglucosamine) could be used to

discriminate these isolates (Table S1).

Next, we performed *in vitro* tests that would account for the ability of *Azospirillum* sp.

UENF-412522 to promote plant growth. The N_2-fixation was supported by positive growth on N-

free semisolid medium and subsequently confirmed by the amplification of *nifH* (Figure 1A). We

also confirmed the nitrogenase activity by using the acetylene reduction assay, which showed a

rate of 28.3 $\pm$ 5.1 nmol.h^{-1} ethylene (Figure 1B).

Indole-acetic acid (IAA) production was investigated in DYGS medium, with and without

L- tryptophan, resulting in IAA concentrations of 51.34 $\pm$ 1.1 µg.mL^{-1} and 37.35 $\pm$ 3.5 µg.mL^{-1},

respectively (Figure 1C), supporting the capacity of *Azospirillum* sp. UENF-412522 to synthesize

this important plant hormone.

We also tested whether *Azospirillum* sp. UENF-412522 is able to solubilize Araxá P-rock

and calcium phosphate in solid medium. We found that this strain formed a halo with a

solubilization index (SI) of 3.33 $\pm$ 0.05 and 1.87 $\pm$ 0.11 for Araxá P-rock and calcium phosphate,

respectively (Figure 1D). We confirmed the P solubilization activity using a liquid medium assay,

which showed 147.5 $\pm$ 3.9 mg.L^{-1} and 52.2 $\pm$ 8.6 mg.L^{-1} solubilized P from Araxá P-rock and

calcium phosphate (Figure 1E), respectively. Conversely, zinc solubilization was not detected,

likely because of zinc sensitivity in this strain.

The plant growth-promoting effects of a given bacterial strain often result from several

concerted processes. Therefore, we inoculated passion fruit plantlets with *Azospirillum* sp.

UENF-412522 and found a clear growth increment in inoculated versus non-inoculated plantlets.

After 10 days, inoculated plantlets had a significant increase in height (54.5%), root fresh matter

(88.6%), root dry matter (61.4%), shoot fresh matter (13.8%) and root length (40%). No

significant difference was found for shoot dry matter (Figure S2).

Under gnotobiotic conditions, inoculated plantlets showed a larger diazotrophic

bacteria population associated with rhizosphere, rhizoplane, and roots (Figure S3). For

rhizosphere and rhizoplane compartments, the population density was significantly greater than

control plantlets (approximately 1.5 $\log_{10}$ bacteria cell per g of soil or root). The bacterial

population inside the plantlets showed no difference between inoculated and non-inoculated

plantlets, further supporting that *Azospirillum* sp. UENF-412522 preferentially colonizes the

rhizosphere/rhizoplane. In parallel, samples from seedlings inoculated with *Azospirillum* sp.

UENF-412522 at 1 and 7 days after germination (d.a.g.) were viewed under SEM (Figure 2),

confirming the root surface as the preferential colonization site. After radicle emergence, single

cells were noted at the bottom region of the radicle (Figure 2A), and bacteria aggregates can be

seen in the vicinity of the crack zone between the radicle and the tegument (Figure 2C-D).

Interestingly, sparse stomata-like structures were seen at the radicle, which seems to be an

infection point for the bacteria (Figure 2B). Bacteria cells interact with the root surface mainly
by apolar attachment and by fibrillar material anchoring the plant cell-wall, establishing an
aggregate network (Figure 2E). At seven d.a.g., bacteria successfully colonized the root hair zone
(Figure 2F) and the elongation/differentiation zone (Figure 2G) of the main developed root.
139

Genome assembly and strain identification
The promising *in vitro* and *ex vitro* results prompted us to sequence the *Azospirillum sp.* UENF-
412522 genome. We used an Illumina HiSeq 2500 instrument (paired-end mode, 2 × 100 bp
reads). Sequencing reads were quality filtered and assembled with SPAdes (see methods for
details). The assembly consists of 101 contigs (length ≥ 500 bp), encompassing 7,360,543 bp,
with a 67.92% GC content, N50 and L50 of 175,376 bp and 13 respectively. The genome harbors
6,508, 71 and 8 protein-coding, tRNA and rRNA genes, respectively.
Genome relatedness within the *Azospirillum* genus was computed using the average
nucleotide identity (ANI) and digital DNA-DNA hybridization (dDDH) of all *Azospirillum* spp.
genomes deposited in RefSeq (n = 48). ANI values were used to build a genome-to-genome
distance matrix and a neighbor-joining dendrogram (Figure 3A). This analysis clearly shows
distinct clusters comprising *A. brasilense* and *A. tiophilum* isolates. In contrast, none of the
remaining strains formed a consistent clade. Further, *A. lipoferum* RC and 4B isolates did not
cluster together, indicating that they do not belong to the same species. We found greater
genetic diversity in the isolates that do not belong to these clusters (Figure 3B). Among those is
Azospirillum sp. UENF-412522, which does not cluster with any other known strain, suggesting
that it represents a novel *Azospirillum* species.
Even though the *Azospirillum* genus has more than 15 described species, genomic data
are biased towards *A. brasilense*, which accounts for 37% (19 isolates) of the available genomes
(Figure 3). We also noticed that five isolates have similar names and share more than 99.9% and
70% in ANI and dDDH analyses, respectively. We removed these redundant genomes from the
downstream analyses and used GCF_001315015, GCF_003119195, GCF_003119115, and
GCF_004923295 as representatives of *A. brasilense* Sp7, *A. brasilense* Sp245, *Azospirillum* sp.
TSH58 and *Azospirillum* sp. TSH100 respectively. Hence, the dataset used in the next sections
comprises 43 publicly available genomes and that of *Azospirillum* sp. UENF-412522 genome.
165

Pangenome analyses of the *Azospirillum* genus
The pangenome is the complete gene repertoire of a given clade (e.g., a species) [28]. We
employed a minimum identity threshold of 50% for gene family identification in *Azospirillum*.
The pangenome comprises 42,515 genes, including a core genome (i.e. genes present in all
isolates) of only 771 genes, likely because of the high diversity, prevalence of genomic
rearrangements, and low synteny of this genus [9]. We clustered the isolates based on their
gene presence/absence patterns and confirmed the high genomic heterogeneity across the
genus (Figure 4A). Our results also support an open pangenome, as it grows with the addition of
new genomes (Figure 4B). Similar results were previously shown in *Alphaproteobacteria* [29],
who found 220 core genes across 27 in *Novosphingobium* spp. genomes. The *Azospirillum* core
genome reported here is substantially greater than that, which could be at least partially
explained by the fact that the vast majority of the isolates studied here were obtained from soil
and plant tissues. *Azospirillum* sp. RU37 (GCF_900188305) and *Azospirillum* sp. RU38E
(GCF_900188385) share the majority of accessory clusters. The same pattern is observed for

Azospirillum sp. TSH20 (GCF_003115935) and *Azospirillum* sp. TSH7 (GCF_003115945). In both
cases, the genomes with unusually high accessory genome similarity also exhibit 100% of ANI
and dDDH, indicating that they belong to the same strain.
Next, we analyzed the unique gene complement of *Azospirillum* sp. UENF-412522. Out
of the 18,065 unique genes inferred by Roary, 405 belonged to this strain. Among these genes,
we identified creatinine amidohydrolase *crnA* (EI613_12750), an enzyme involved in the
degradation of creatinine [30], which has been reported as replication-impairing molecule in
bacteria [31]. The gene 3-alpha-hydroxysteroid dehydrogenase *hsdA* (EI613_09215), involved in
steroid catabolism, is also exclusive to this strain and might assist *Azospirillum* sp. UENF-412522
to use steroids as a carbon source [32]. We also found the two component system *prsDE*
(EI613_31600-31605) as exclusive to this strain. PrsDE produces a low molecular weight
succinoglycan, which might assist in plant colonization by *Azospirillum sp.* UENF-412522 [33].
192
Plant growth promotion genes
Plant growth modulation by bacteria is a complex process [34]. Several bacterial genes are often
associated with direct beneficial effects, such as those responsible for biological nitrogen
fixation (BFN), root growth enhancement, phosphate solubilization, and deviation of the
ethylene biosynthesis pathway towards ammonia and α-ketobutyrate. We assessed the
presence of genes that are likely involved in direct plant growth promotion across *Azospirillum*
strains, with a particular focus in *Azospirillum sp.* UENF-412522. This procedure was guided by a
maximum likelihood phylogenetic tree built with the protein sequences encoded by the core-
genome, which was combined with a gene presence/absence matrix (Figure 5). Below we
discuss these genes in light of the mechanisms by which they promote plant growth.
BNF is a major plant growth promotion mechanism present in several *Azospirillum*
strains [8, 35]. This process relies on the nitrogenase enzyme complex, which reduces
atmospheric nitrogen (N_2) to ammonium (NH_4). The *nifHDK* genes encode, respectively, a
nitrogenase iron (Fe) protein, a nitrogenase molybdenum-iron (MoFe) protein alpha chain, and
a nitrogenase MoFe protein beta chain. The FeMo co-factor (FeMoCo) present in the MoFe
protein binds N_2, while the Fe protein uses the energy from ATP hydrolysis to drive the reduction
of N_2 to NH_4 by FeMoCo [36]. We used the presence of the *nifHDK* genes to predict the nitrogen
fixation capacity across *Azospirillum* genomes. All the analyzed genomes have these genes,
except for *Azospirillum* sp. RU37A, *Azospirillum* sp. RU38E and *Azospirillum* sp. L-25-5w-1 (Figure
5). *Azospirillum* sp. UENF-412522 possesses the *nifHDK* (EI613_05225-05215) operon and a
nifDK (EI613_05195-05200) operon, resembling the distribution found in most of the other
Azospirillum genomes containing *nifHDK*. Some exceptions to this structure are found in *A.*
brasilense Sp245, which possesses four *nifDK*, including vanadium nitrogenase variants [37];
Azospirillum sp. B506 presents a single *nifK*, and; *Azospirillum* sp. RU37A and *Azospirillum* sp.
RU38E have two *nifD*. The presence of more than one *nifHDK* increase nitrogen fixation to the
bacteria and associated plants. However, given that BNF is energetically expensive, the selective
advantage of such increased activity remains unclear.
Most of the soil phosphate is insoluble and hence unavailable to direct plant
absorption [38]. Several PGPRs promote plant growth by releasing organic acids that chelate
divalent cations (e.g., Ca^{2+}) in poorly soluble mineral forms (e.g., hydroxyapatite), increasing P
availability [39]. Gluconic acid secretion is the best-characterized mechanism of phosphate
solubilization, that is performed by pyrroloquinoline quinone (PQQ)-dependent glucose

dehydrogenase (GDH) [40]. We used the presence of the genes encoding the PQQ co-factor as
a proxy to understand P solubilization through GDH-PQQ in *Azospirillum* spp. The *pqq* genes
found in *Azospirillum* sp. were *pqqABCDE* (EI613_19670-19690). While *Azospirillum* UENF-
412522 and *A. halopraeferens* DSM3675 harbor the complete operon, all other strains lack *pqqA*
(Figure 5). The *pqq* operon is absent in three strains (*Azospirillum* sp. RU37A, *Azospirillum* sp.
RU38E, *Azospirillum* sp. L-25-5w-1). *A. brasilense* FP2 lacks *pqqA* and *pqqD*. Interestingly,
phytase genes were only found in the pqq-lacking strains *Azospirillum* sp. RU37A and
Azospirillum sp. RU38E, indicating that the capacity to mobilize P from organic compounds
displaced PQQ-dependent P solubilization in these bacteria.

Although *pqqA* has been regarded as non-essential for PQQ biosynthesis [41], other
studies support its role as the backbone for PQQ biogenesis [42]. PqqA binds to PqqD, prior to
PqqAD interaction with PqqE [43]. The crucial role of *pqqA* and *pqqB* was also confirmed by
knockout mutants of *Rahnella aquatilis* HX2, which showed decreased biocontrol and mineral P
solubilization [44]. A direct correlation between *pqqB* and *pqqF* expression and PQQ production
has also been shown [45]. Further, the same authors also reported that *Pseudomonas putida*
KT2440 *pqqF* is controlled by an independent promoter and terminator, allowing this gene to
modulate PQQ levels [45]. The presence of incomplete *pqq* operons in many *Azospirillum* strains
allows one to speculate that these isolates have other genes to compensate for the absence of
pqqA, a hypothesis that warrants experimental validation. Further, upstream of *pqqABCDE*, we
found an alcohol dehydrogenase ADH IIB *qbdA* (EI613_19655) and its associated regulator *amgR*
(EI613_19660), which are possibly related to acetic acid production mediated by PQQ [46]. This
operon also has a a gene encoding a hypothetical protein containing a CXXCW motif, present in
several soil bacteria, which might be associated with PQQ catabolism (TIGR03865). Hence, the
pqqABCDE operon might be co-regulated with *qdbA* in *Azospirillum* sp. UENF-412522.

Ethylene is a plant hormone with a central role in senescence and growth of leaves,
flowers, and fruits [47]. Under stress conditions, ethylene can impair plant cell elongation [48].
Ethylene is synthesized by the oxidation of 1-aminocyclopropane-1-carboxylate (ACC) by ACC
oxidase. ACC can be broken down into α-ketobutyrate and NH_3 by ACC deaminase [49], an
enzyme produced by several PGPR that induce plant growth by lowering ethylene levels under
stress conditions. We identified the ACC deaminase gene (*acdS*, EI613_03680) in *Azospirillum*
sp. UENF-412522 and other 11 *Azospirillum* sp. strains, supporting that their association with
plants can alleviate ethylene-mediated growth inhibition. Importantly, we have not found *acdS*
in *A. brasilense* strains.

Considering its gene complement, *Azospirillum sp.* UENF-412522 has a significant
potential to promote root growth. Despite the strong experimental evidence supporting IAA
biosynthesis, we found none of the classic auxin biosynthesis genes (i.e., *ipdC*, *iaaH* and *iaaM*)
[50-52], which prompted us to investigate other IAA biosynthesis pathways [51] in this strain.
Interestingly, we found a gene encoding a nitrilase (*nit*, EI613_03300), which is 74% similar that
encoded by a *nit* that converts indole-3-acetonitrile (IAN) directly into IAA or into indol-3-
acetamide (IAM) in the rhizobacterium *Pseudomonas* sp. UW4 [53, 54]. In the latter pathway,
the conversion of IAM into IAA requires the action of IAM-hydrolases, for which we found two
candidate genes, EI613_04545 e EI613_07015. These genes are similar to amide hydrolases from
Rhodococcus sp. (Q53116) [55, 56] and *Agrobacterium fabacearum* (ADY67766.1) [57],
respectively. The apparent absence of tryptophan monooxygenase genes, which catalyze the
conversion of Trp into IAM, indicates that *Azospirillum* sp. UENF 412522 produces IAA through

an interplay between the IAN and IAM pathways, as reported in *Rhizobium* spp. and
Bradyrhizobium spp. [53, 58]. Nevertheless, the initial steps in the production of IAN in
Azospirillum sp. UENF 412522 are yet to be elucidated. In contrast, all *A. brasilense* strains
possess the gene *ipdC* (Figure 5), supporting the ability of this species to synthesize IAA by the
indole-3-pyruvate pathway, as experimentally demonstrated [59, 60]. Another eight *A.*
brasilense strains possess nitrile hydratase (*nthB*) and *iaaH* genes, which contribute to IAA
production by the conversion of IAN to IAM, and from IAM to IAA, respectively.

We also investigated *Azospirillum sp.* UENF-412522 genes that might promote root
development through the emission of nitrite and nitric oxide. *A. brasilense* is known for secreting
nitrite as a product of nitrate respiration [61, 62]. *Azospirillum sp.* UENF-412522 has a nitrate
reductase operon *napCAB* (EI613_10540-10550). *napA* encodes a periplasmic nitrate reductase,
while *napCB* encodes electron transfer subunits necessary for napA activity. This operon was
proposed as an alternative electron acceptor for oxygen in *A. brasilense* sp. 245 periplasm [63]
and is possibly responsible for nitrate respiration in *Azospirillum sp.* UENF-412522. The nitrite
generated by nitrate respiration can be exported to the environment, absorbed by the plant
root, and reduced to nitric oxide. Under acidic pH, nitrite can be reduced to nitric oxide without
enzymatic action [64] and absorbed by root cells [65]. The downside of nitrate respiration to
plant growth promotion is the reduction of available nitrogen in the system and the production
of greenhouse gases. Hence, we hypothesize that *Azospirillum* sp. UENF-412522 nitrogen
fixation contributes to nitrate respiration and plant absorption.

Most known *Azospirillum* strains are diazotrophs. N_2-fixation requires a microaerophilic
or anoxic environment because nitrogenase metalloclusters are sensitive to oxygen. Previous
works in *Azospirillum* proposed that complete oxygen depletion inhibited nitrogenase activity
[66], while others suggest otherwise [67]. *Azospirillum sp.* UENF-412522 was isolated from the
rhizoplane, where low oxygen conditions are rather common [68]. Given the genomic
investigations reported here, we hypothesize that P solubilization, root growth enhancement by
nitrate respiration, and N_2 fixation are dependent of an anoxic environment and might be
interconnected in the soil-root system (Figure 6). Under low oxygen availability, nitrate is used
as final electron acceptor in anaerobic respiration, producing nitrite, which is reduced to nitric
oxide under low pH in the periplasm. Such acidic environment can be harnessed by the gluconic
acid generated by the PQQ-dependent glucose dehydrogenase. Nitric oxide or nitrite would then
be exported to the rhizosphere, stimulating root growth. Finally, gluconic acid is also exported
to the rhizosphere, solubilizing phosphate for the root cells uptake.

Antibiotic resistance genes
Azospirillum sp. UENF-412522 has several antibiotic resistance genes, such as those conferring
resistance to: fosmidomycin (*fsr*, EI613_04505) [69], bicyclomycin (*bcr*, EI613_29470,
EI613_14405) [70], bacitracin (*uppP/BacA*, EI613_27660) [71], tabtoxin (*ttr*, EI613_22745) [72],
phenazine (*ehpR*, EI613_22490) [73], and tetracycline (*tetA*, EI613_26440) [74]. We also found
multiple drug resistance genes (*mdtABCE*, EI613_19480-19470) [75], *acrAB* (EI613_25875-
25880) [76], and *norM* (EI613_15640) [77]. Interestingly, most of these antibiotics, such as
bicyclomycin [78], phenazine [79], bacitracin, and tetracycline [80], are commonly produced
Streptomyces, which is abundant in the soil, where it plays a key role in plant organic matter
decomposition [81, 82]. The presence of genes conferring resistance against *Streptomyces*

antimicrobials is likely important to the survival of *Azospirillum sp.* UENF-412522 in the rhizosphere.

CONCLUSION

In the present study we described *Azospirillum sp.* UENF-412522, a new *Azospirillum* plant growth-promoting species. We experimentally demonstrated that this bacterium is able to fix nitrogen, produce IAA, solubilize mineral P, and recognize and colonize seed-root surfaces. This strain is equipped with genes involved in BNF, P solubilization, ethylene catabolism, promotion of root development, and antibiotic resistance in the rhizosphere. Collectively, our results support the biotechnological potential of *Azospirillum sp.* UENF-412522 to be part of bioinoculant formulations for sustainable agriculture. We also propose a system involved in the partitioning of the fixed nitrogen between nitrate respiration and plant absorption, which is likely critical in plant-growth promotion under microaerophilic or anoxic conditions. Finally, this study emphasizes the importance of comprehensive studies using genomic, microscopy, microbiological, and biochemical approaches to characterize novel *Azospirillum* species outside of the *A. brasilense* clade, which could reveal key unlock the genus diversity and open novel possibilities agricultural applications.

METHODS

Bacterial isolation and DNA purification

Passion fruit (*Passiflora edulis* f. *edulis*) root samples were obtained at the Integrated Agroecological Production System (SIPA), Embrapa Agrobiologia, Seropédica, Rio de Janeiro, Brazil (coordinates 22.7635°S 43.6886°W) and taken to the laboratory in a sterilized plastic bag kept in ice. Rhizoplane bacteria isolation was performed by an adapted version of Bramwell et al. (1995)[83]. In summary, root axis segments (10 g fresh weight) were sealed from both tip sides with paraffin, gently washed in tap water, and transferred to a 250 mL glass flask containing 90 mL of saline solution (NaCl, 0.9%). Roots were kept under agitation for 5 min at a rotatory shaker (200 rpm at 30 °C) to remove soil particles. After that, the same sealed root segment was carefully transferred to another 250 mL glass flask containing 90 mL of sterile saline (NaCl, 0.9%) solution plus 10 g of autoclaved sieved sand, where remained under agitation for 10 min at rotatory shaker (150 rpm at 30 °C) to detach bacteria cells from root surfaces (rhizoplane bacteria fraction). After sand decantation, 10 mL of the suspension were transferred to 90 mL of sterile saline solution (10^{-1} dilution), which were used to prepare successive dilutions until 10^{-6}. Three independent 20 µL aliquots of all serial dilutions (10^{-2} to 10^{-6}) were inoculated into glass vials (16 mL volume) containing 5 mL NFb nitrogen-free semisolid medium (composition in g L^{-1}): malic acid, 5.0; K_2HPO_4, 0.5; $MgSO_4.7H_2O$, 0.2; NaCl, 0.1; $CaCl_2. 2H_2O$, 0.02; micronutrient stock solution ($CuSO_4.5H_2O$, 0.04; $ZnSO_4.7H_2O$, 0.12; H_3BO_3, 1.40; $Na_2MoO_4.2H_2O$, 1.0; $MnSO_4. H_2O$, 1.175. Complete volume to 1,000 mL with distilled water), 2 mL; bromothymol blue (5 g L^{-1} in 0.2 N KOH), 2 mL; FeEDTA (solution 16.4 g L^{-1}), 4 mL; vitamin stock solution (biotin, 10 mg; pyridoxal-HCl, 20 mg in 100 mL distilled water), 1 mL; KOH, 4.5 g; distilled water to bring the final volume to 1,000 mL and adjust pH to 6.5. A quantity of 1.6 g agar L^{-1} was added to prepare the semisolid medium [84].

Inoculated NFb vials were incubated in a growth chamber at 30 °C for 5 to 7 days. The growth of diazotrophic bacteria was characterized by the formation of a subsurface white pellicle. Using a loopful, a sample of the pellicle from the last positive dilution was inoculated in

a fresh NFb N-free semisolid medium and incubated under the same conditions described
above. This process was subsequently conducted for three times and the last new pellicle
formed was streaked onto DYGS solid medium plates for colony purification. The DYGS
composition (g L^{-1}): glucose, 2.0; malic acid 2.0; peptone, 1.5; yeast extract, 2.0; K_2HPO_4, 0,5;
$MgSO_4.7H_2O$, 0.5; glutamic acid, 1.5, pH is 6.5. The purified isolate was stored in 10% glycerol at
-80 ºC, under the code UENF-412522, as a diazotrophic bacterium associated with the passion
fruit rhizoplane.
For DNA extraction, 100 µL of the stored bacteria were transferred to a glass tube
containing 5 mL DYGS liquid medium and grown in an orbital shaker at 30 °C and 180 rpm for 48
hours. After that, 20 µL were transferred to the same medium and grown at the same conditions
for 24 h. An aliquot of 200 µL was taken, and bacterial genomic DNA was extracted using the
QIAamp kit (QIAGEN) following the manufacturer's instructions. DNA was quantified in a 3000
NanoDrop (Thermo Scientific USA). DNA quality was analyzed using a 0.8% agarose gel stained
with gel red.
To obtain an initial taxonomic assignment, PCR was conducted with the primers 27F (5`-
AGAGTTTGATCMTGGCTCAG 3`) and 1492R (5'-TACGGYTACCTTGTTACGACTT-3`). These
reactions were performed using 50 ng of genomic DNA in 25 µL final reaction volume,
containing: 2 µL of dNTPs (20 mM each) 2.5 µL of 10 x enzyme buffer, 0.75 µL 50 mM $MgCl_2$, 2.5
µL of each primer (5 mM), 0.3 µL Taq polymerase (5 U/µL). Amplification was performed in a 96-
well thermocycler Veriti model (Applied Biosystems), programmed at 9°C for 3 min, 30 cycles of
amplification (94°C for 1 min, 55 °C for 30 s, 72 °C for 30 s) at 72 °C for 10 min. The sequencing
reactions were performed using the Big Dye Terminator Sequencing Kit-Cycle Sequencing Ready
ABI Prism version 3 (Life Technologies, USA), following manufacturer's recommendations.
Sequencing was performed on ABI Sequencer model 3130 (Applied Biosystems).
383
Plant growth promotion traits of UENF-412522 bacterial strain
nifH gene detection
The *nifH* presence were assessed by PCR, with the primers PolF (5'TGCGAYCCSAARGCBGACTC
3`) and PolR (5'ATSGCCATCATYTCRGCCGA 3') [85]. In each reaction, we used 1 µL dNTPs (200
mM each), 5 µL of 10X buffer, 4 µL of 25 mM $MgCl_2$, 0.5 µL of each primer (10 mM), 0.25 µL Taq
polymerase (5 U/ µL) and 50 ng DNA, in a final volume of 50 µL. The amplification conditions
were denaturation at 94 °C for 5 minutes, then 30 cycles (94 °C for 1 min, 55 ºC for 1 min, 72 °C
for 2 minutes) and a final extension step (72 °C for 5 min). PCR products were inspected on 1.5%
agarose gels. Bp-100 DNA ladder (Invitrogen) was used as a molecular weight marker.
393
Acetylene reduction activity assay (ARA)
The *in vitro* nitrogen fixation ability of UENF-412522 was evaluated using the acetylene
reduction assay (ARA), as previously described by Baldani [84]. The strain was grown in 16 mL
glass vials containing 5 mL of N-free NFB semisolid medium, as described above. These flasks
were inoculated with 20 µL of the bacterial inoculum suspended in sterile water, adjusted to an
optical density (OD) of 1.0, and incubated at 30 °C for 48 h. After pellicle formation, the vials
were closed with a sterilized pierceable rubber stopper of the subseal type. Syringes were used
to remove 1 mL of air and to inject 1 mL of acetylene in each vial. The flasks were incubated at
30 °C for 1 h, and 1 mL of the gas phase was analyzed on a gas chromatograph with flame
ionization (Perkin Elmer), to determine the ethylene concentration in the sample.

Phosphorus and zinc solubilization

Phosphate solubilization was independently carried out using 1 g.L^{-1} Araxá rock phosphate [$Ca_{10}(PO_4)_6F_2$] and tricalcium phosphate [$Ca_3(PO_4)_2$]. Bacterial cultures grown in DYGS liquid medium adjusted to an optical density of 1.0 were inoculated (20 μL) at the center of the petri dish with agar containing Pikovskaya medium. After this procedure, they were incubated at 30 °C for 7 days. The ability of the isolates in solubilizing P was evaluated by measuring the translucent halo, according to Kumar and Narula formula [86], in which S.I. (solubility index) = halo Diameter (mm)/colony diameter (mm). We conducted three biological replicates of this assay.

We also assessed P solubilization in Pikovskaya liquid medium supplemented with 1g.L^{-1} of the Araxá rock phosphate or $Ca_{10}(PO_4)_6F_2$ was used to quantify the soluble P. Aliquots of 100 μL of bacterial cultures grown in DYGS liquid medium were transferred to 50 mL tubes containing Pikovskaya medium and kept for 7 days under constant agitation at 150 rpm in a rotatory shaker at 30 °C. After this period, cultures were centrifuged at 3200 rpm for 15 minutes, and the supernatant was used for pH determination and quantification of soluble P.

Zinc solubilization was carried out according to Intorne et al [87]. Aliquots of 20 μL of bacterial strains cultivated for 24 h at 30 °C in DYGS liquid medium, adjusted to an optical density of 1.0, were inoculated at the center of the petri dish containing Saravanan medium with agar supplemented with 1g.L^{-1} of ZnO and incubated at 30 °C for 7 days. After this period, the ability to solubilize ZnO was also performed according to Kumar and Narula [86]. We conducted three biological replicates of this assay.

Indole acetic acid (IAA) production

Bacterial isolates were cultured in liquid DYGS medium for 24 h at 30 °C and transferred (25 μL) to test tubes containing 5 mL of the same medium in the presence and absence of L-tryptophan (100 mg.L^{-1}) before incubation in the dark for 72 h at 30 °C, shaking at 150 rpm. The cultures were then transferred to 2 mL tubes and centrifuged at 10,000 rpm for 10 min, and the supernatant transferred to a test tube with 2 mL of Salkowski reagent [88]. Tubes were incubated for 30 min in the dark. The production of IAA was evaluated by the presence of pink color in the tubes, and the color intensity was determined with a spectrophotometer at a wavelength of 530 nm. IAA concentrations were measured using a calibration curve.

Biochemical API 50 CH/E test

For the metabolic characterization of bacterial isolates, we used the API 50 CH/E test kit (bioMérieux SA Marcy- l'Etoile/France), which evaluates the ability of the isolates to conduct fermentation of 49 carbohydrates and derivatives. To this end, the isolates were cultured in DYGS medium for 24 h at 150 rpm. After this period, the isolates were suspended in autoclaved water and adjusted to optical density 1.0; 2 mL of this suspension were added to the CHL medium (bioMérieux) and transferred to galleries containing different substrates. Each dome was sealed with a sterile drop of mineral oil and incubated in an environmental chamber at 37 °C. Reads were recorded after 24 h and 48 h after inoculation, and eventual production of organic acids during incubation shifts the pH indicator from red to yellow.

Plant growth promotion assay under greenhouse conditions

A greenhouse assay was carried out to evaluate the effect of UENF-412522 inoculation in plants.
To perform it, passion fruit seeds of the cultivar Yellow master FB200 were disinfected in 1%
sodium hypochlorite solution for 20 minutes, followed by three rinses in autoclaved distilled
water and then transferred to germitest paper moistened with autoclaved water. Then, the
seeds were incubated in a growth chamber at 30 °C, for 16 h in the light and 8 h in the dark.
After germinating, these seeds were transferred to styrofoam trays containing autoclaved plant
substrate under greenhouse conditions. After 30 days of transplanting, seedlings of
approximately 6 cm were transferred to a 300 mL plastic bag containing the same autoclaved
plant growth substrate. The inoculum of the UENF-412522 strain was prepared in liquid DYGS
medium after 40 h of growth in a rotatory shaker at 30 °C and 150 rpm. Bacterial suspensions
containing approximately 10^8 cells.mL^{-1} (O.D. = 1.0 at 600 ηm) were inoculated on the region of
the plant's neck, and plants were transferred to the greenhouse for 30 days. The assay was set
up in a randomized block design, two treatments (inoculated and non-inoculated plantlets) and
six replications. The control plants received 1 mL of the liquid DYGS medium containing lysed
UENF-412522 cells.
463

Plant root colonization assay
We evaluated plant-bacteria interaction by determining the population size and root
colonization by scanning electron microscopy (SEM). For this, passion fruit seeds of the Yellow
cultivar master FB200 were disinfected and placed to germinate as described above. After
germination, the seedlings were transferred to a test tube containing autoclaved vermiculite.
Then, 1 ml of bacterial suspension containing approximately 10^8 cells mL^{-1} was inoculated on the
seedling neck. Seedlings were kept for seven days after inoculation (d.a.i) in a culture room
under a temperature of 30ºC (16 h in the light and 8 hours in the dark). For SEM, root samples
were collected at 1 and 7 days after the inoculation of the bacterial suspension. Samples for SEM
were prepared according to Baldotto et al. [89] and viewed at Zeiss DSEM 962 at 15 Kv voltage
in secondary electron detector mode. The bacteria count at rhizosphere, rhizoplane, and roots
was performed at seven d.a.i., as described above. The control consisted of seedlings inoculated
with the DYGS medium.
477

Genome sequencing and assembly
The sequencing procedures and basic data processing were performed, as previously described
[25]. In summary, paired-end libraries were prepared using the TruSeq Nano DNA LT Library Prep
(Illumina) and sequenced on a HiSeq 2500 instrument at the Life Sciences Core Facility (LaCTAD;
UNICAMP, Campinas, Brazil). Sequencing reads (2 × 100 bp) had their quality checked with
FastQC 0.11.5 (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Quality filtering
was performed with Trimmomatic 0.35 [90], and reads with average quality below 30 were
discarded. The genome assembly was performed using SPAdes 3.8 [91] and assembly metrics
evaluated by QUAST 3.0 [92].
487

Genome annotation and phylogenetic analysis
The assembled genome was annotated with the NCBI Prokaryotic Genome Annotation Pipeline
[93]. The *Azospirillum* sp. UENF-412522 genome was deposited on Genbank under the
BioProject PRJNA508400. *Azospirillum* spp. genomes available in RefSeq [94, 95] were
downloaded (n = 48, as of June 03 2019) and protein sequences were predicted using PROKKA

1.13.3 [96]. We searched for proteins encoded by genes potentially involved in plant growth
promotion using a blastp [97] search on the SwissProt database, with minimum coverage and
similarity thresholds of 70% and 60%, respectively. All-against-all average nucleotide identity
(ANI) was computed using pyani 0.28 [98]. Pan-genome analysis was performed with Roary
3.12.0, with a minimum identity threshold of 50% [99]. Multiple sequence alignments were
performed using Muscle 3.8.30 [100] and phylogenetic reconstructions performed with RAxML
8.2.10 [101].
500

501 **AUTHOR CONTRIBUTIONS**
502 Conceived the study: FLO, TMV; Funding and resources: FLO, TMV; Data analysis: GLR, FPM, RKG,
503 FP-S, DC-A, IP-O; Performed experiments: PSLG, STS, AFA; Interpretation of the results: GLR,
504 FPM, FLO, TMV; Wrote the manuscript: GLR, FPM, FLO, TMV.
505

ACKNOWLEDGEMENTS
This work was supported by Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do
Rio de Janeiro (FAPERJ; grants E-26/111.827/2013, E-26/102.259/2013, E-26/201.239/2014, E-
26/203.309/2016 and E-26/203.014/2018), Coordenação de Aperfeiçoamento de Pessoal de
Nível Superior - Brasil (CAPES; Finance Code 001), and Conselho Nacional de Desenvolvimento
Científico e Tecnológico (CNPq; grant 449904/2014-8 and 314263/2018-7). The funding agencies
had no role in the design of the study and collection, analysis, and interpretation of data and in
writing.

REFERENCES
1.   Bourguet D, Guillemaud T: **The hidden and external costs of pesticide use**. In:
Sustainable Agriculture Reviews. Springer; 2016: 35-120.
2.   Maroniche GA, Diaz PR, Borrajo MP, Valverde CF, Creus CM: **Friends or foes in the**
rhizosphere: traits of fluorescent Pseudomonas that hinder Azospirillum brasilense
growth and root colonization. *FEMS Microbiol Ecol* 2018, **94**(12).
3.   Vejan P, Abdullah R, Khadiran T, Ismail S, Nasrulhaq Boyce A: **Role of Plant Growth**
Promoting Rhizobacteria in Agricultural Sustainability-A Review. *Molecules* 2016,
21(5).
4.   Prasad M, Srinivasan R, Chaudhary M, Choudhary M, Jat LK: **Plant Growth Promoting**
Rhizobacteria (PGPR) for Sustainable Agriculture: Perspectives and Challenges. In:
PGPR Amelioration in Sustainable Agriculture. Elsevier; 2019: 129-157.
5.   Brink SC: **Unlocking the Secrets of the Rhizosphere**. *Trends Plant Sci* 2016, **21**(3):169-
170.
6.   Ahkami AH, White III RA, Handakumbura PP, Jansson C: **Rhizosphere engineering:**
Enhancing sustainable plant ecosystem productivity. *Rhizosphere* 2017, **3**:233-243.
7.   Mehnaz S: **Azospirillum: a biofertilizer for every crop**. In: *Plant Microbes Symbiosis:*
Applied Facets. Springer; 2015: 297-314.
8.   Fukami J, Cerezini P, Hungria M: **Azospirillum: benefits that go far beyond biological**
nitrogen fixation. *AMB Express* 2018, **8**(1):73.
9.   Wisniewski-Dye F, Borziak K, Khalsa-Moyers G, Alexandre G, Sukharnikov LO, Wuichet
K, Hurst GB, McDonald WH, Robertson JS, Barbe V *et al*: **Azospirillum genomes reveal**

transition of bacteria from aquatic to terrestrial environments. *PLoS Genet* 2011,
7(12):e1002430.

10.   Tarrand JJ, Krieg NR, Dobereiner J: **A taxonomic study of the Spirillum lipoferum**
group, with descriptions of a new genus, Azospirillum gen. nov. and two species,
Azospirillum lipoferum (Beijerinck) comb. nov. and Azospirillum brasilense sp. nov.
Can J Microbiol 1978, **24**(8):967-980.

11.   Reis VM, Baldani VLD, Baldani JI: **Isolation, Identification and Biochemical**
Characterization of Azospirillum spp. and Other Nitrogen-Fixing Bacteria. In:
Handbook for Azospirillum: Technical Issues and Protocols. Edited by Cassán FD, Okon
Y, Creus CM. Cham: Springer International Publishing; 2015: 3-26.

12.   Pankievicz VC, do Amaral FP, Santos KF, Agtuca B, Xu Y, Schueller MJ, Arisi AC, Steffens
MB, de Souza EM, Pedrosa FO *et al*: **Robust biological nitrogen fixation in a model**
grass-bacterial association. *Plant J* 2015, **81**(6):907-919.

13.   Rodriguez H, Gonzalez T, Goire I, Bashan Y: **Gluconic acid production and phosphate**
solubilization by the plant growth-promoting bacterium Azospirillum spp.
Naturwissenschaften 2004, **91**(11):552-555.

14.   Cohen AC, Bottini R, Pontin M, Berli FJ, Moreno D, Boccanlandro H, Travaglia CN,
Piccoli PN: **Azospirillum brasilense ameliorates the response of Arabidopsis thaliana**
to drought mainly via enhancement of ABA levels. *Physiol Plant* 2015, **153**(1):79-90.

15.   Garcia JE, Maroniche G, Creus C, Suarez-Rodriguez R, Ramirez-Trujillo JA, Groppa MD:
In vitro PGPR properties and osmotic tolerance of different Azospirillum native
strains and their effects on growth of maize under drought stress. *Microbiol Res*
2017, **202**:21-29.

16.   D'Angioli AM, Viani RAG, Lambers H, Sawaya ACHF, Oliveira RS: **Inoculation with**
Azospirillum brasilense (Ab-V4, Ab-V5) increases Zea mays root carboxylate-
exudation rates, dependent on soil phosphorus supply. *Plant and soil* 2017, **410**(1-
2):499-507.

17.   Vacheron J, Renoud S, Muller D, Babalola OO, Prigent-Combaret C: **Alleviation of**
abiotic and biotic stresses in plants by Azospirillum. In: *Handbook for Azospirillum.*
Springer; 2015: 333-365.

18.   Cassán F, Diaz-Zorita M: **Azospirillum sp. in current agriculture: From the laboratory**
to the field. *Soil Biology and Biochemistry* 2016, **103**:117-130.

19.   Hungria M, Ribeiro RA, Nogueira MA: **Draft Genome Sequences of Azospirillum**
brasilense Strains Ab-V5 and Ab-V6, Commercially Used in Inoculants for Grasses and
Legumes in Brazil. *Genome Announc* 2018, **6**(20).

20.   Blaha D, Prigent-Combaret C, Mirza MS, Moënne-Loccoz Y: **Phylogeny of the 1-**
aminocyclopropane-1-carboxylic acid deaminase-encoding gene acdS in
phytobeneficial and pathogenic Proteobacteria and relation with strain
biogeography. *FEMS Microbiology Ecology* 2006, **56**(3):455-470.

21.   Esquivel-Cote R, Ramírez-Gama RM, Tsuzuki-Reyes G, Orozco-Segovia A, Huante PJP,
Soil: **Azospirillum lipoferum strain AZm5 containing 1-aminocyclopropane-1-**
carboxylic acid deaminase improves early growth of tomato seedlings under nitrogen
deficiency. 2010, **337**(1):65-75.

22.   Vikram A, Alagawadi AR, Krishnaraj PU, Mahesh Kumar KSJWJoM, Biotechnology:
581   **Transconjugation studies in Azospirillum sp. negative to mineral phosphate**
582   **solubilization**. 2007, **23**(9):1333-1337.
583   23.   Kaneko T, Minamisawa K, Isawa T, Nakatsukasa H, Mitsui H, Kawaharada Y, Nakamura
584   Y, Watanabe A, Kawashima K, Ono A *et al*: **Complete genomic structure of the**
585   **cultivated rice endophyte Azospirillum sp. B510**. *DNA Res* 2010, **17**(1):37-50.
586   24.   Malhotra M, Srivastava S: **An ipdC gene knock-out of Azospirillum brasilense strain**
587   **SM and its implications on indole-3-acetic acid biosynthesis and plant growth**
588   **promotion**. *Antonie Van Leeuwenhoek* 2008, **93**(4):425-433.
589   25.   Matteoli FP, Passarelli-Araujo H, Reis RJA, da Rocha LO, de Souza EM, Aravind L,
590   Olivares FL, Venancio TM: **Genome sequencing and assessment of plant growth-**
591   **promoting properties of a Serratia marcescens strain isolated from vermicompost**.
592   *BMC Genomics* 2018, **19**(1):750.
593   26.   Matteoli FP, Passarelli-Araujo H, Pedrosa-Silva F, Olivares FL, Venancio TM: **Population**
594   **structure and pangenome analysis of Enterobacter bugandensis uncover the**
595   **presence of blaCTX-M-55, blaNDM-5 and blaIMI-1, along with sophisticated iron**
596   **acquisition strategies**. *Genomics* 2020, **112**(2):1182-1191.
597   27.   Eckert B, Weber OB, Kirchhof G, Halbritter A, Stoffels M, Hartmann A: **Azospirillum**
598   **doebereinerae sp. nov., a nitrogen-fixing bacterium associated with the C4-grass**
599   **Miscanthus**. *Int J Syst Evol Microbiol* 2001, **51**(Pt 1):17-26.
600   28.   Xiao J, Zhang Z, Wu J, Yu J: **A brief review of software tools for pangenomics**.
601   *Genomics Proteomics Bioinformatics* 2015, **13**(1):73-76.
602   29.   Kumar R, Verma H, Haider S, Bajaj A, Sood U, Ponnusamy K, Nagar S, Shakarad MN,
603   Negi RK, Singh Y *et al*: **Comparative Genomic Analysis Reveals Habitat-Specific Genes**
604   **and Regulatory Hubs within the Genus Novosphingobium**. *mSystems* 2017, **2**(3).
605   30.   Beuth B, Niefind K, Schomburg D: **Crystal structure of creatininase from Pseudomonas**
606   **putida: a novel fold and a case of convergent evolution**. *J Mol Biol* 2003, **332**(1):287-
607   301.
608   31.   McDonald T, Drescher KM, Weber A, Tracy S: **Creatinine inhibits bacterial replication**.
609   *J Antibiot (Tokyo)* 2012, **65**(3):153-156.
610   32.   Mobus E, Maser E: **Molecular cloning, overexpression, and characterization of**
611   **steroid-inducible 3alpha-hydroxysteroid dehydrogenase/carbonyl reductase from**
612   **Comamonas testosteroni. A novel member of the short-chain**
613   **dehydrogenase/reductase superfamily**. *J Biol Chem* 1998, **273**(47):30888-30896.
614   33.   York GM, Walker GC: **The Rhizobium meliloti exoK gene and prsD/prsE/exsH genes**
615   **are components of independent degradative pathways which contribute to**
616   **production of low-molecular-weight succinoglycan**. *Mol Microbiol* 1997, **25**(1):117-
617   134.
618   34.   Numan M, Bashir S, Khan Y, Mumtaz R, Shinwari ZK, Khan AL, Khan A, Al-Harrasi A:
619   **Plant growth promoting bacteria as an alternative strategy for salt tolerance in**
620   **plants: A review**. *Microbiol Res* 2018, **209**:21-32.
621   35.   Steenhoudt O, Vanderleyden J: **Azospirillum, a free-living nitrogen-fixing bacterium**
622   **closely associated with grasses: genetic, biochemical and ecological aspects**. *FEMS*
623   *Microbiol Rev* 2000, **24**(4):487-506.

36.   Dixon R, Kahn D: **Genetic regulation of biological nitrogen fixation**. *Nat Rev Microbiol*
2004, **2**(8):621-631.
37.   Wisniewski-Dye F, Lozano L, Acosta-Cruz E, Borland S, Drogue B, Prigent-Combaret C,
Rouy Z, Barbe V, Herrera AM, Gonzalez V *et al*: **Genome Sequence of Azospirillum**
brasilense CBG497 and Comparative Analyses of Azospirillum Core and Accessory
Genomes provide Insight into Niche Adaptation. *Genes (Basel)* 2012, **3**(4):576-602.
38.   Ewel JJ, Schreeg LA, Sinclair TR: **Resources for Crop Production: Accessing the**
Unavailable. *Trends Plant Sci* 2019, **24**(2):121-129.
39.   Sharma SB, Sayyed RZ, Trivedi MH, Gobi TA: **Phosphate solubilizing microbes:**
sustainable approach for managing phosphorus deficiency in agricultural soils.
Springerplus 2013, **2**:587.
40.   Alori ET, Glick BR, Babalola OO: **Microbial Phosphorus Solubilization and Its Potential**
for Use in Sustainable Agriculture. *Front Microbiol* 2017, **8**:971.
41.   Toyama H, Lidstrom ME: **pqqA is not required for biosynthesis of pyrroloquinoline**
quinone in Methylobacterium extorquens AM1. *Microbiology* 1998, **144 (Pt 1)**:183-
191.
42.   Barr I, Latham JA, Iavarone AT, Chantarojsiri T, Hwang JD, Klinman JP: **Demonstration**
That the Radical S-Adenosylmethionine (SAM) Enzyme PqqE Catalyzes de Novo
Carbon-Carbon Cross-linking within a Peptide Substrate PqqA in the Presence of the
Peptide Chaperone PqqD. *J Biol Chem* 2016, **291**(17):8877-8884.
43.   Latham JA, Iavarone AT, Barr I, Juthani PV, Klinman JP: **PqqD is a novel peptide**
chaperone that forms a ternary complex with the radical S-adenosylmethionine
protein PqqE in the pyrroloquinoline quinone biosynthetic pathway. *J Biol Chem*
2015, **290**(20):12908-12918.
44.   Li L, Jiao Z, Hale L, Wu W, Guo Y: **Disruption of gene pqqA or pqqB reduces plant**
growth promotion activity and biocontrol of crown gall disease by Rahnella aquatilis
HX2. *PLoS One* 2014, **9**(12):e115010.
45.   An R, Moe LA: **Regulation of Pyrroloquinoline Quinone-dependent glucose**
dehydrogenase activity in the model rhizosphere-dwelling bacterium Pseudomonas
putida KT2440. *Appl Environ Microbiol* 2016, **82**(16):4955-4964.
46.   Yakushi T, Matsushita K: **Alcohol dehydrogenase of acetic acid bacteria: structure,**
mode of action, and applications in biotechnology. *Appl Microbiol Biotechnol* 2010,
86(5):1257-1265.
47.   Iqbal N, Khan NA, Ferrante A, Trivellini A, Francini A, Khan MIR: **Ethylene Role in Plant**
Growth, Development and Senescence: Interaction with Other Phytohormones.
Front Plant Sci 2017, **8**:475.
48.   Vaseva, II, Qudeimat E, Potuschak T, Du Y, Genschik P, Vandenbussche F, Van Der
Straeten D: **The plant hormone ethylene restricts Arabidopsis growth via the**
epidermis. *Proc Natl Acad Sci U S A* 2018, **115**(17):E4130-E4139.
49.   Glick BR: **Bacteria with ACC deaminase can promote plant growth and help to feed**
the world. *Microbiol Res* 2014, **169**(1):30-39.
50.   Spaepen S, Vanderleyden J, Remans R: **Indole-3-acetic acid in microbial and**
microorganism-plant signaling. *FEMS Microbiol Rev* 2007, **31**(4):425-448.
51.   Spaepen S, Vanderleyden J: **Auxin and plant-microbe interactions**. *Cold Spring Harb*
Perspect Biol 2011, **3**(4).

52. Imada EL, Rolla Dos Santos AA, Oliveira AL, Hungria M, Rodrigues EP: **Indole-3-acetic**
acid production via the indole-3-pyruvate pathway by plant growth promoter
Rhizobium tropici CIAT 899 is strongly inhibited by ammonium. *Res Microbiol* 2017,
168(3):283-292.
53. Duca D, Rose DR, Glick BR: **Characterization of a nitrilase and a nitrile hydratase from**
Pseudomonas sp. strain UW4 that converts indole-3-acetonitrile to indole-3-acetic
acid. *Appl Environ Microbiol* 2014, **80**(15):4640-4649.
54. Duca DR, Rose DR, Glick BR: **Indole acetic acid overproduction transformants of the**
rhizobacterium Pseudomonas sp. UW4. *Antonie Van Leeuwenhoek* 2018, **111**(9):1645-
1660.
55. Mayaux JF, Cerbelaud E, Soubrier F, Yeh P, Blanche F, Petre D: **Purification, cloning,**
and primary structure of a new enantiomer-selective amidase from a Rhodococcus
strain: structural evidence for a conserved genetic coupling with nitrile hydratase. *J*
Bacteriol 1991, **173**(21):6694-6704.
56. Patten CL, Blakney AJ, Coulson TJ: **Activity, distribution and function of indole-3-acetic**
acid biosynthetic pathways in bacteria. *Crit Rev Microbiol* 2013, **39**(4):395-415.
57. Delamuta JRM, Scherer AJ, Ribeiro RA, Hungria M: **Genetic diversity of Agrobacterium**
species isolated from nodules of common bean and soybean in Brazil, Mexico,
Ecuador and Mozambique, and description of the new species Agrobacterium
fabacearum sp. nov. *Int J Syst Evol Microbiol* 2020, **70**(7):4233-4244.
58. Vega-Hernández MC, León-Barrios M, Pérez-Galdona R: **Indole-3-acetic acid**
production from indole-3-acetonitrile in Bradyrhizobium. *Soil Biology and*
Biochemistry 2002, **34**(5):665-668.
59. Vande Broek A, Lambrecht M, Eggermont K, Vanderleyden J: **Auxins upregulate**
expression of the indole-3-pyruvate decarboxylase gene in Azospirillum brasilense. *J*
Bacteriol 1999, **181**(4):1338-1342.
60. El-Khawas H, Adachi KJB, Soils Fo: **Identification and quantification of auxins in**
culture media of Azospirillum and Klebsiella and their effect on rice roots. 1999,
28(4):377-381.
61. Creus CM, Graziano M, Casanovas EM, Pereyra MA, Simontacchi M, Puntarulo S,
Barassi CA, Lamattina L: **Nitric oxide is involved in the Azospirillum brasilense-**
induced lateral root formation in tomato. *Planta* 2005, **221**(2):297-303.
62. Molina-Favero C, Creus CM, Simontacchi M, Puntarulo S, Lamattina L: **Aerobic nitric**
oxide production by Azospirillum brasilense Sp245 and its influence on root
architecture in tomato. *Mol Plant Microbe Interact* 2008, **21**(7):1001-1009.
63. Steenhoudt O, Keijers V, Okon Y, Vanderleyden J: **Identification and characterization**
of a periplasmic nitrate reductase in Azospirillum brasilense Sp245. *Arch Microbiol*
2001, **175**(5):344-352.
64. Bethke PC, Badger MR, Jones RL: **Apoplastic synthesis of nitric oxide by plant tissues**.
Plant Cell 2004, **16**(2):332-341.
65. Molina-Favero C, Mónica Creus C, Lanteri ML, Correa-Aragunde N, Lombardo MC,
Barassi CA, Lamattina L: **Nitric Oxide and Plant Growth Promoting Rhizobacteria:**
Common Features Influencing Root Growth and Development. In: *Advances in*
Botanical Research. vol. 46: Academic Press; 2007: 1-33.

66.   Zhang Y, Burris RH, Ludden PW, Roberts GP: **Regulation of nitrogen fixation in**
Azospirillum brasilense. *FEMS Microbiol Lett* 1997, **152**(2):195-204.

67.   Malinich EA, Bauer CE: **Transcriptome analysis of Azospirillum brasilense vegetative**
and cyst states reveals large-scale alterations in metabolic and replicative gene
expression. *Microb Genom* 2018, **4**(8).

68.   Lecomte SM, Achouak W, Abrouk D, Heulin T, Nesme X, Haichar FeZ: **Diversifying**
Anaerobic Respiration Strategies to Compete in the Rhizosphere. 2018, **6**(139).

69.   Fujisaki S, Ohnuma S, Horiuchi T, Takahashi I, Tsukui S, Nishimura Y, Nishino T,
Kitabatake M, Inokuchi H: **Cloning of a gene from Escherichia coli that confers**
resistance to fosmidomycin as a consequence of amplification. *Gene* 1996, **175**(1-
2):83-87.

70.   Bentley J, Hyatt LS, Ainley K, Parish JH, Herbert RB, White GR: **Cloning and sequence**
analysis of an Escherichia coli gene conferring bicyclomycin resistance. *Gene* 1993,
127(1):117-120.

71.   El Ghachi M, Bouhss A, Blanot D, Mengin-Lecreulx D: **The bacA gene of Escherichia coli**
encodes an undecaprenyl pyrophosphate phosphatase activity. *J Biol Chem* 2004,
279(29):30106-30113.

72.   Anzai H, Yoneyama K, Yamaguchi I: **The nucleotide sequence of tabtoxin resistance**
gene (ttr) of Pseudomonas syringae pv. tabaci. *Nucleic Acids Res* 1990, **18**(7):1890.

73.   Giddens SR, Feng Y, Mahanty HK: **Characterization of a novel phenazine antibiotic**
gene cluster in Erwinia herbicola Eh1087. *Mol Microbiol* 2002, **45**(3):769-783.

74.   Waters SH, Rogowsky P, Grinsted J, Altenbuchner J, Schmitt R: **The tetracycline**
resistance determinants of RP1 and Tn1721: nucleotide sequence analysis. *Nucleic*
Acids Res 1983, **11**(17):6089-6105.

75.   Nagakubo S, Nishino K, Hirata T, Yamaguchi A: **The putative response regulator BaeR**
stimulates multidrug resistance of Escherichia coli via a novel multidrug exporter
system, MdtABC. *J Bacteriol* 2002, **184**(15):4161-4167.

76.   Okusu H, Ma D, Nikaido H: **AcrAB efflux pump plays a major role in the antibiotic**
resistance phenotype of Escherichia coli multiple-antibiotic-resistance (Mar)
mutants. *J Bacteriol* 1996, **178**(1):306-308.

77.   Brown MH, Paulsen IT, Skurray RA: **The multidrug efflux protein NorM is a prototype**
of a new family of transporters. *Mol Microbiol* 1999, **31**(1):394-395.

78.   Vior NM, Lacret R, Chandra G, Dorai-Raj S, Trick M, Truman AW: **Discovery and**
Biosynthesis of the Antibiotic Bicyclomycin in Distantly Related Bacterial Classes.
Appl Environ Microbiol 2018, **84**(9).

79.   Abdelfattah MS, Ishikawa N, Karmakar UK, Yamaku K, Ishibashi M: **New phenazine**
analogues from Streptomyces sp. IFM 11694 with TRAIL resistance-overcoming
activities. *J Antibiot (Tokyo)* 2016, **69**(6):446-450.

80.   Procopio RE, Silva IR, Martins MK, Azevedo JL, Araujo JM: **Antibiotics produced by**
Streptomyces. *Braz J Infect Dis* 2012, **16**(5):466-471.

81.   Chater KF, Biro S, Lee KJ, Palmer T, Schrempf H: **The complex extracellular biology of**
Streptomyces. *FEMS Microbiol Rev* 2010, **34**(2):171-198.

82.   Chater KF: **Recent advances in understanding Streptomyces**. *F1000Res* 2016, **5**:2795.

83. Bramwell PA, Barallon RV, Rogers HJ, Bailey MJ: **Extraction and PCR amplification of**
DNA from the rhizoplane. In: *Molecular microbial ecology manual.* Springer; 1995: 89-
108.

84. Baldani JI, Reis VM, Videira SS, Boddey LH, Baldani VLD: **The art of isolating nitrogen-**
fixing bacteria from non-leguminous plants using N-free semi-solid media: a practical
guide for microbiologists. *Plant and Soil* 2014, **384**(1):413-431.

85. Poly F, Ranjard L, Nazaret S, Gourbiere F, Monrozier LJ: **Comparison of nifH gene pools**
in soils and soil microenvironments with contrasting properties. *Appl Environ*
Microbiol 2001, **67**(5):2255-2262.

86. Kumar V, Narula N: **Solubilization of inorganic phosphates and growth emergence of**
wheat as affected by Azotobacter chroococcum mutants. *Biology and Fertility of Soils*
1999, **28**(3):301-305.

87. Intorne AC, de Oliveira MV, Lima ML, da Silva JF, Olivares FL, de Souza Filho GA:
Identification and characterization of Gluconacetobacter diazotrophicus mutants
defective in the solubilization of phosphorus and zinc. *Arch Microbiol* 2009,
191(5):477-483.

88. Sarwar M, Kremer RJ: **Determination of bacterially derived auxins using a microplate**
method. *Letters in Applied Microbiology* 1995, **20**(5):282-285.

89. Baldotto LEB, Olivares FL, Bressan-Smith R: **Structural interaction between GFP-**
labeled diazotrophic endophytic bacterium Herbaspirillum seropedicae RAM10 and
pineapple plantlets 'Vitória'. *Brazilian Journal of Microbiology* 2011, **42**:114-125.

90. Bolger AM, Lohse M, Usadel B: **Trimmomatic: a flexible trimmer for Illumina**
sequence data. *Bioinformatics* 2014, **30**(15):2114-2120.

91. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,
Nikolenko SI, Pham S, Prjibelski AD *et al*: **SPAdes: a new genome assembly algorithm**
and its applications to single-cell sequencing. *J Comput Biol* 2012, **19**(5):455-477.

92. Gurevich A, Saveliev V, Vyahhi N, Tesler G: **QUAST: quality assessment tool for**
genome assemblies. *Bioinformatics* 2013, **29**(8):1072-1075.

93. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L,
Lomsadze A, Pruitt KD, Borodovsky M, Ostell J: **NCBI prokaryotic genome annotation**
pipeline. *Nucleic acids research* 2016, **44**(14):6614-6624.

94. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, Rajput B, Robbertse
B, Smith-White B, Ako-Adjei D *et al*: **Reference sequence (RefSeq) database at NCBI:**
current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res*
2016, **44**(D1):D733-745.

95. Pruitt KD, Tatusova T, Maglott DR: **NCBI reference sequences (RefSeq): a curated non-**
redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids*
Res 2007, **35**(Database issue):D61-65.

96. Seemann T: **Prokka: rapid prokaryotic genome annotation**. *Bioinformatics* 2014,
30(14):2068-2069.

97. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ: **Gapped**
BLAST and PSI-BLAST: a new generation of protein database search programs.
Nucleic Acids Res 1997, **25**(17):3389-3402.

98.   Pritchard L, Glover RH, Humphris S, Elphinstone JG, Toth IK: **Genomics and taxonomy**
800   **in diagnostics for food security: soft-rotting enterobacterial plant pathogens**.
801   *Analytical Methods* 2016, **8**(1):12-24.
802   99.   Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, Fookes M, Falush D,
803   Keane JA, Parkhill J: **Roary: rapid large-scale prokaryote pan genome analysis**.
804   *Bioinformatics* 2015, **31**(22):3691-3693.
805   100.   Edgar RC: **MUSCLE: multiple sequence alignment with high accuracy and high**
806   **throughput**. *Nucleic Acids Res* 2004, **32**(5):1792-1797.
807   101.   Stamatakis A: **RAxML version 8: a tool for phylogenetic analysis and post-analysis of**
808   **large phylogenies**. *Bioinformatics* 2014, **30**(9):1312-1313.

846   **Figure 1:** Plant growth promotion *in vitro* traits of *Azospirillum* sp. UENF-412522. (A)
847   Amplification of *nifH* using *E. coli* as negative control; molecular weight marker (MM). (B)
848   Nitrogenase activity by the acetylene reduction assay. (C) Indole acetic acid (IAA) production
849   with and without L- tryptophan. Araxá P-rock and calcium phosphate solubilization in solid (D)
850   and liquid (E) medium.
851

852
Figure 2: Scanning electron microscopy (SEM) of passion fruit (*Passiflora edulis* subsp. *edulis*)
seed to seedlings phase inoculated with *Azospirillum* sp. UENF-412522 in an axenic cultivation
system at one day (A-E) and seven days after germination (F-G). (A) General view of the emerged
radicle surface (r) with single bacteria cells attached (arrows). Note the sparse rudimentary
stomata (rectangle magnified in B), bar = 100 μm; (B) Bacteria cells in the vicinity and into the
stomata pore (arrows), bar = 5μm; (C) General view of the emerged radicle (er) breaking the
seed tegument (t), rectangle magnified in D, bar = 1000 μm; (D) Transition zone between broken

tegument (t) and emerged radicle with bacterial aggregates visible in surface tegument (white arrows) and emerged roots (black arrow), rectangle magnified in E, bar = 100 μm; (E) Bacteria aggregates (b) attached by fibrils in the seed surface (black arrow) and linking cell-to-cell (white arrows), bar = 1 μm; (F) Bacteria cells colonizing the root hair (rh) zone as a single-cell by apolar attachment, bar = 20 μm; (G) Aggregated bacteria cells colonizing the elongation-differentiation root zone mainly by apolar attachment and associated to the epidermal cell-wall-junction, bar = 10 μm.

Figure 3: Average nucleotide identity (ANI). (A) Neighbor-joining cladogram. (B) Genome-to-genome distance matrix.

Figure 4: *Azospirillum* spp. pan-genome. (A) Gene presence/absence matrix, with gene presence
depicted with solid blue boxes. (B) Pan- and core-genome sizes versus new genomes additions
to the pan-genome dataset.

Figure 5: Core maximum likelihood phylogenetic tree. Plant growth-promoting genes were searched, grouped according to the mechanism of growth modulation, and mapped onto the tree.

894
Figure 6: Schematic representation of nitrogen fixation, phosphate solubilization, and
nitrite/nitric oxide emission by *Azospirillum sp.* UENF-412522 in the rhizosphere. The identified
genes responsible for nitrogen fixation (*nifHDK*), root enhancement (*napCAB*), and phosphate
solubilization (*pqqABCDE*) are represented within boxes. Nitrogen fixation occurs in the
cytoplasm, while gluconic acid production and nitrate respiration take place at the periplasm.
Ammonium generated by the nitrogenase can undergo anaerobic nitrification to nitrite, which
can be exported to the periplasm. In the periplasm, glucose dehydrogenase, along with the PQQ
co-factor, produces gluconic acid from glucose, lowering the pH. Nitrate is reduced to nitrite by
a periplasmatic nitrate reductase napCAB. At low pH, nitrite can be reduced to nitric oxide.
Acetic acid, nitrite, and nitric oxide can be exported to the rhizosphere. Acetic acid can reduce
the pH in the rhizosphere and solubilize phosphate. The solubilized phosphate, nitrite, and nitric
oxide can be captured by root cells.

907