The neuropeptide receptor NMUR-1 regulates the specificity of *C. elegans* innate

immunity against pathogen infection

Phillip Wibisono[1], Shawndra Wibisono[1], Jan Watteyne[2], Chia-Hui Chen[1], Durai Sellegounder[1],

Isabel Beets[2], Yiyong Liu[1,3*], and Jingru Sun[1*]

[1]Department of Biomedical Sciences, Elson S. Floyd College of Medicine, Washington State

University, Spokane, WA, USA

[2]Department of Biology, KU Leuven, Belgium

[3]Genomics Core, Washington State University, Spokane, WA, USA

*To whom correspondence should be addressed: yiyong.liu@wsu.edu; jingru.sun@wsu.edu

SUMMARY

A key question in current immunology is how the innate immune system generates high levels

of specificity. Like most invertebrates, *Caenorhabditis elegans* does not have an adaptive

immune system and relies solely on innate immunity to defend itself against pathogen attacks,

yet it can still differentiate different pathogens and launch distinct innate immune responses.

Here, we have found that functional loss of NMUR-1, a neuronal GPCR homologous to

mammalian receptors for the neuropeptide neuromedin U, has diverse effects on *C. elegans*

survival against various bacterial pathogens. Transcriptomic analyses and functional assays

revealed that NMUR-1 modulates *C. elegans* transcription activity by regulating the expression

of transcription factors, which, in turn, controls the expression of distinct immune genes in

response to different pathogens. Our study has uncovered a molecular basis for the specificity

of *C. elegans* innate immunity that could provide mechanistic insights into understanding the

specificity of vertebrate innate immunity.

Key words: *Caenorhabditis elegans*, specificity of innate immunity, neuromedin U, G protein-

coupled receptor, the nervous system

INTRODUCTION

The pathogen-triggered host immune response is a multilayered process programed to fight

invading microorganisms and maintain healthy homeostasis. There are two types of immune

systems: the innate immune system and the adaptive immune system. The innate immune

system, comprised of physical barriers as well as inducible physiological and behavioral

responses, serves as the first line of defense for the host and is evolutionarily conserved across

multicellular organisms. Innate immune responses are initiated by recognizing pathogen-

associated molecular patterns (PAMPs) or damage caused by pathogen-encoded virulence

factors (1,2). Such recognition triggers multiple internal defense responses, including the

release of cytokines and chemokines, to fight invading microbes directly or indirectly. In

vertebrates, when the innate immune response is insufficient to control an infection, the highly

specific adaptive immune response is activated as the second line of defense (3). In this

system, antigen-presenting cells (APCs) engulf and break down pathogens and present the

foreign antigens on their cell surface via MHC class II molecules. Subsequently, T cells and B

cells with antigen-specific receptors are activated to destroy infected cells or produce antibodies

to inhibit infection, respectively.

While a large repertoire of T- and B-cell receptors and antibodies in the adaptive immune

system confer remarkable specificity against different pathogens, the innate immune response

is traditionally believed to be non-specific. However, increasing evidence indicates that the

innate immune system can also generate high levels of specificity. In vertebrates, a diverse

array of pattern recognition receptors (PRRs), such as the Toll-like receptors (TLRs), NOD-like

receptors (NLRs), retinoic acid-inducible gene-I-like receptors (RLRs), and C-type lectins

(CLRs), recognize various PAMPs and mediate a broad pattern of specificity (4). The diversity

of ligand binding can be further expanded by the formation of receptor homodimers or

heterodimers through interactions with other PRR co-receptors (5). In addition to controlling

immune activation through specific receptor-ligand binding, innate immunity is also tightly

regulated by the nervous system to ensure that appropriate responses are mounted against

specific pathogens. Such regulation is critical because insufficient immune responses can

exacerbate infection and excessive immune responses can lead to prolonged inflammation,

tissue damage, or even death (6,7). The mechanisms underlying neuro-immune regulation,

however, are not well understood and are difficult to dissect due to the high degree of

complexity of the nervous and immune systems in most model organisms. Overall, the

molecular basis responsible for the specificity of innate immunity in vertebrates remains unclear

and is largely understudied, possibly because such specificity is overshadowed by the

concomitant adaptive immune response.

Unlike vertebrates, most invertebrates do not have an adaptive immune system and rely solely

on innate immunity to defend themselves against pathogen infection, yet they can still

differentiate between different pathogens (4). For example, *Drosophila melanogaster* possesses

a repertoire of receptors that mediate the distinction of different types of pathogens and fungi

(8). Alternative splicing of exons of the Dscam protein in insects generates thousands of

isoforms that are potentially involved in pathogen recognition and resistance (9). Moreover, in

both *D. melanogaster* and *Caenorhabditis elegans*, brief pre-exposure of animals to a pathogen

(termed priming or conditioning) provides strong protection against a subsequent challenge with

an otherwise lethal dose of the same pathogen (10). These studies demonstrate that

invertebrate innate immune systems can generate high levels of specificity, although the

underlying mechanisms remain poorly understood. Nonetheless, this allows for studies of the

specificity of innate immunity without the confounding variability from adaptive immunity. In

invertebrate organisms, the limited genetic diversity of PRRs and/or immune effectors is likely

insufficient to explain the high levels of specificity (9). This is exemplified by the case of *C.*

elegans. The nematode genome does not encode the majority of known PRRs, and its sole TLR

protein, TOL-1, does not seem to play key roles in immunity (11). However, *C. elegans* can still

discern different pathogens and launch distinct immune responses against them (12,13).

Different infections induce both pathogen-specific and shared immune genes in the worm

(13,14). While how the *C. elegans* immune system identifies specific pathogens remains

elusive, several key studies have revealed that neural regulation of immunity has homologous

occurrence in the nematode (15-21), which could control appropriate immune responses to

different pathogens. The characteristics of *C. elegans*, such as the simplicity of its nervous and

immune systems, genetic tractability, invariant lineage, effectiveness of RNA interference

(RNAi), and a transparent body that allows for the monitoring of gene expression, make this

organism uniquely suited for studying neural regulation of immunity. Indeed, such research in *C.*

elegans is at the forefront of the field and has revealed unprecedented details regarding the

molecules, cells, and signaling pathways involved in neural regulation of immunity (reviewed in

(22)). For example, we have described an octopaminergic neuro-immune regulatory circuit in

great detail (19,23-25) as well as a neural-cuticle defense regulatory circuit mediated by the

neuropeptide receptor NPR-8 (26). Others have also revealed serotonergic, dopaminergic,

cholinergic, and neuropeptidergic immunoregulatory pathways that control innate immune

responses to various pathogens (reviewed in (22)). These studies have greatly facilitated our

understanding of complex neuro-immune signaling mechanisms.

While the above-mentioned *C. elegans* studies showed how the nervous system regulates

immunity against certain pathogens, they did not address whether or how such regulation

generates specificity against different pathogens. During our work on neuro-immune regulation

in *C. elegans*, we observed that functional loss of NMUR-1, a neuronal GPCR homologous to

mammalian receptors for the neuropeptide neuromedin U (NMU), had diverse effects on the

survival of *C. elegans* exposed to various bacterial pathogens. Furthermore, an earlier study by

Alcedo and colleagues showed that an *nmur-1* null mutation affected the lifespan of *C. elegans*

fed different *E. coli* food sources, and that such effects were dependent upon the type of *E. coli*

lipopolysaccharide (LPS) structures (27). These findings indicate that NMU/NMUR-1 signaling

might mediate the specificity of immune responses to different pathogens. NMU is known to

have roles in regulating immune responses in addition to its functions regulating smooth muscle

contractions in the uterus, reducing food intake and weight, and modifying ion transport (28,29).

Indeed, three independent studies in mice demonstrated that NMU from enteric neurons directly

activates type 2 innate lymphoid cells through NMUR1 to drive anti-parasitic immunity (30-32).

In the current study, we have discovered that NMUR-1 modulates *C. elegans* transcriptional

activity by regulating the expression of transcription factors, which, in turn, controls the

expression of distinct immune genes in response to different pathogens. Our study has

uncovered a molecular basis for the specificity of *C. elegans* innate immunity that could

potentially provide mechanistic insights into understanding the specificity of innate immune

responses in vertebrates.

RESULTS

Functional loss of NMUR-1 differentially affects *C. elegans* survival against various

bacterial pathogens

Functional loss of NMUR-1 has distinct effects on the lifespan of *C. elegans* fed different *E. coli*

food sources, and such effects are dependent upon the type of *E. coli* LPS structures (27).

Given the toxic and immunogenic nature of LPS (33), these LPS-dependent lifespan

phenotypes suggest that NMUR-1 might mediate distinct immune responses to different

pathogens. To test this notion, we examined the susceptibility of wild-type *N2* and *nmur-1* null

animals (*nmur-1(ok1387)* strain) to six paradigmatic pathogens that included three Gram-

negative bacteria (*Salmonella enterica* strain SL1344, *Pseudomonas aeruginosa* strain PA14,

and *Yersinia pestis* strain kim5) and three Gram-positive bacteria (*Enterococcus faecalis* strain

OG1RF, *Microbacterium nematophilum* strain CBX102, and *Staphylococcus aureus* strain

NCTC8325). Compared to wild-type animals, *nmur-1(ok1387)* animals exhibited enhanced

survival against *S. enterica* and *Y. pestis*, reduced survival against *E. faecalis*, and showed no

differences in survival against *P. aeruginosa*, *M. nematophilum*, and *S. aureus* (Fig. 1 and Fig.

S1). In addition, with two commonly used worm food sources, namely, *E. coli* strains OP50 and

HT115, we found that the *nmur-1* mutation improved worm survival against OP50 but had no

effect against HT115 (Fig. 1 and Fig. S1), an observation in agreement with a previous study by

Alcedo and colleagues (27). These distinct effects of the *nmur-1* mutation on survival, which

appeared to be unrelated to the Gram-negative or Gram-positive groups, indicate that NMUR-1

plays diverse roles in *C. elegans* defense against various pathogens.

There are many factors affecting the pathogenesis of the above-mentioned bacteria besides

being Gram-negative or Gram-positive. For example, each pathogen produces a unique set of

virulence factors that promote pathogenicity through different mechanisms (34). In turn, *C.*

elegans can launch distinct defense responses to specific pathogens (12-14). The diverse

effects that NMUR-1 has on *C. elegans* survival suggest that it could have multiple physiological

functions in the nematode's defense against these bacteria. Given NMUR-1's potential

versatility in function, we sought to gain insights into NMUR-1's functionality by focusing on its

roles in defense against *E. faecalis* and *S. enterica*, two pathogens that have opposite effects

on the survival of *nmur-1* mutant animals (reduced survival against *E. faecalis* but improved

survival against *S. enterica* (Fig. 1)). Below, we describe our study using these two pathogens

as infecting agents.

Pathogen avoidance behavior does not contribute to the altered survival of *nmur-*

1(ok1387)* animals exposed to *E. faecalis* or *S. enterica

Compared to wild-type animals, *nmur-1(ok1387)* animals exhibited altered survival following

exposure to *E. faecalis* or *S. enterica* (Fig. 1). However, no differences in survival were

observed between mutant and wild-type animals when they were fed worm food *E. coli* strain

HT115 (Fig. 1), heat-killed *E. faecalis* (Fig. S2A), or heat-killed *S. enterica* (Fig. S2B). These

results indicate that the *nmur-1* mutation affects *C. elegans* defense against living pathogens

without impacting the nematode's basic lifespan. The altered survival of mutant animals could

be due to pathogen avoidance behavior, a known mechanism of *C. elegans* defense (35). To

test this possibility, we measured animal survival using full-lawn assays in which the agar plates

were completely covered by pathogenic bacteria to eliminate pathogen avoidance. Consistent

with our original (partial-lawn) survival assays, we observed that *nmur-1(ok1387)* animals

exposed to *E. faecalis* died at a faster rate than wild-type controls, whereas mutants exposed to

S. enterica died at a slower rate than wild-type controls (Fig. S3). These results suggest that

lawn avoidance is not the reason for the altered survival observed in *nmur-1* mutants. We next

conducted lawn occupancy behavioral assays to compare the magnitude of pathogen

avoidance between wild-type and *nmur-1(ok1387)* animals. In these assays, a small lawn of

pathogenic bacteria was seeded in the center of an agar plate, and a set number of

synchronized young adult animals were placed on the lawn. The numbers of animals that

stayed on and off the lawn were then counted at five time points over a period of 36 hours.

Compared to wild-type animals, *nmur-1(ok1387)* animals overall did not show significant

differences in the magnitudes of pathogen avoidance (Fig. 2). Taken together, these data

suggest that pathogen avoidance behavior does not play a role in the altered survival of *nmur-*

1(ok1387) animals exposed to *E. faecalis* or *S. enterica*.

Functional loss of NMUR-1 changes intestinal accumulation of *E. faecalis* and *S. enterica*

Certain mutations in *C. elegans* can change pathogen accumulation in the intestine, leading to

altered resistance to infection (36). To examine whether the *nmur-1* mutation affects bacterial

accumulation, we used GFP-tagged *E. faecalis* to visualize accumulated bacteria in the

nematode's intestine. Our results showed that the GFP fluorescence intensity in *nmur-*

1(ok1387) animals was stronger than that seen in wild-type animals (Fig. 3A). Intestinal

bacterial loads were quantified by counting the Colony Forming Units (CFU) of live bacterial

cells recovered from the intestine. The mutant animals had more *E. faecalis* CFUs than wild-

type animals (Fig. 3B). These results indicated that the *nmur*-1 mutation caused increased

intestinal accumulation of *E. faecalis*. The higher bacterial accumulation could be due to altered

pathogen intake, expulsion, or clearance by immunity. These possible causes were further

investigated. *C. elegans* feeds on bacterial food via rhythmic contractions (pumping) of its

pharynx, so to determine whether there were any differences in pathogen intake between wild-

type and *nmur-1(ok1387)* animals, we measured their pharyngeal pumping rates on bacterial

lawns. When feeding on *E. faecalis*, both wild-type and mutant animals showed similar pumping

rates, reflecting similar levels of pathogen intake (Fig. 3C). Bacterial evacuation was assessed

by measuring the defecation rate, defined as the time interval between expulsions of gut

contents. We found that the defecation rate of *nmur-1(ok1387)* animals on *E. faecalis* was

comparable to the rate observed in wild-type animals (Fig. 3D). Overall, these results indicate

that the reduced survival of *nmur-1(ok1387)* animals exposed to *E. faecalis* is likely due to

increased pathogen accumulation in the intestine. Because the *nmur-1* mutation did not alter *E.*

faecalis intake or expulsion, the increased pathogen accumulation likely resulted from the

mutants' weakened innate immunity that was unable to clear the infection from the intestine.

We next performed similar experiments to investigate intestinal accumulation, intake, and

expulsion of *S. enterica*. Our results showed that, compared to wild-type animals, *nmur-*

1(ok1387) animals exhibited slightly weaker GFP-tagged *S. enterica* fluorescence intensities

(Fig. 3E) and had significantly lower CFU counts (Fig. 3F), indicating that the *nmur*-1 mutation

decreased intestinal accumulation of *S. enterica*. Mutant animals also had slightly higher

pharyngeal pumping rates than wild-type animals (Fig. 3G), suggesting that pathogen intake

was increased in the mutants despite their reduced bacterial accumulation. The defecation rate

of *nmur-1(ok1387)* animals on *S. enterica* was comparable to that of wild-type animals (Fig. 3H).

These results indicate that the enhanced survival of *nmur-1(ok1387)* animals exposed to *S.*

enterica is likely due to decreased pathogen accumulation in the intestine. This decrease did not

result from reduced pathogen intake (pathogen intake actually improved) or increased expulsion

but likely resulted from enhanced innate immunity in the mutants that cleared the infection from

the intestine. Taken together, both the *E. faecalis* and *S. enterica* studies suggest that NMUR-1

could regulate innate immunity in *C. elegans* defense against pathogen infection but plays

opposite roles in such potential regulation against the two pathogens.

NMUR-1 regulates *C. elegans* transcriptional activity

To gain insights into how NMUR-1 regulates *C. elegans* defense at the molecular level, we

employed RNA sequencing to profile gene expression in wild-type and *nmur-1(ok1387)* animals

with or without pathogen infection. For these studies, we collected RNA samples from wild-type

and *nmur-1(ok1387)* animals with or without exposure to *E. faecalis* or *S. enterica* (five

replicates per group). The samples were then submitted to the Washington State University

Genomics Core for RNA-seq analysis. The resulting sequence data (FASTQ files) were

deposited in the National Center for Biotechnology Information (NCBI) Sequence Read Archive

(SRA) database through the Gene Expression Omnibus (GEO). The processed gene

quantification files and differential expression files were deposited in the GEO. All of these data

can be accessed through the GEO with the accession number GSE154324.

Using the resulting sequencing data, we first compared the gene expression profiles of *nmur-*

1(ok1387) animals with those of wild-type animals under normal conditions (i.e., without

infection). In total, 14,206 genes were detected and quantified with a false discovery rate (FDR)

of 5%. Among these genes, 1,599 were upregulated and 275 were downregulated at least two-

fold in *nmur-1(ok1387)* animals relative to wild-type animals. The change in expression of such

a large number of genes in the mutant worms indicates that the lack of NMUR-1 profoundly

affects gene expression in *C. elegans*. Gene ontology (GO) analysis of the 1,599 upregulated

genes identified 26 significantly enriched molecular functions, including protein

kinase/phosphatase activity, ribonucleotide binding, and cuticle structure activity (Table S1).

Interestingly, GO analysis of the 276 downregulated genes showed 11 significantly enriched

molecular functions, all of which involve transcription-related DNA binding activities (Table 1A).

Twenty-three downregulated genes possess these activities and contribute to the functional

enrichment, most of which encode transcription factors involved in binding to RNA polymerase II

regulatory regions (Table 1B). These results indicate that NMUR-1 could affect transcription by

regulating the expression of these transcription factors.

Because of the high importance of transcription in the fundamental functions of life and the

potential regulatory role of NMUR-1 in this process, we next examined how NMUR-1, or the lack

thereof, impacts transcription. To this end, we developed a novel *in vitro* transcription system to

measure *C. elegans* transcriptional activity (37). One of the biggest obstacles for biochemical

studies of *C. elegans* is the high difficulty of preparing functionally active nuclear extract due to

its thick surrounding cuticle. In our transcription system, Balch homogenization was employed to

effectively disrupt worms, and functionally active nuclear extracts were obtained through

subcellular fractionation. Subsequently, the nuclear extracts were used to re-constitute *in vitro*

transcription reactions with a linear DNA template containing the worm *Δpes-10* promoter,

followed by PCR-based, non-radioactive detection of transcripts (37). Using this system, we

were able to compare the general transcriptional activity of *nmur-1(ok1387)* animals with that of

wild-type animals at the whole-organism level. PCR and gel analysis of the transcript products

showed substantial increases in transcriptional activity in reactions with mutant nuclear extract

compared to those with wild-type nuclear extract (Fig. 4A). To quantitatively compare the

transcriptional activity of wild-type and mutant nuclear extracts, we next titrated the DNA

template with varying amounts of each extract and used qRT-PCR to quantify transcripts at

each titration point. The resulting data were then fitted with the Michaelis-Menten model and the

non-linear least-squares method, as we described previously (37). The titration curves and two

titration parameters, namely, the maximum yield and the amount of nuclear extract needed to

reach 50% of the maximum yield, are shown in Fig. 4B. Compared to the titrations with wild-type

nuclear extract, those with mutant extract yielded 2.33-fold maximum number of transcripts and

only needed 0.67-fold extract to reach 50% of the maximum yield (Fig. 4B), indicating that the

reactions performed with mutant extract were more robust and had higher transcriptional

activity. These results were surprising given the fact that the expression of many transcription

factors was downregulated in the mutant animals. Taken together, our data suggest that NMUR-

1 suppresses transcriptional activity in wild-type animals and that functional loss of NMUR-1

abrogates this suppression, which explains the large number of genes (1,599 genes) that were

upregulated in *nmur-1(ok1387)* animals relative to wild-type animals.

We also employed *in vivo* transcription assays to evaluate how the lack of NMUR-1 affects

transcriptional activity in *C. elegans*. For these assays, transgenic strain CL2070, which

contains the *hsp-16.2p::GFP* reporter transgene, was used. This strain does not express

detectable GFP under standard conditions but broadly expresses GFP in all tissues after heat

shock, allowing for the assessment of overall transcriptional activity in living animals (38). The

CL2070 animals were outcrossed with our wild-type animals and then crossed with *nmur-*

1(ok1387) animals. The resulting animals (*CL2070(dvIs70)* and *CL2070(dvIs70);nmur-*

1(ok1387)) were grown to the L4 larval stage, heat shocked at 35°C for 2 hours, and then

returned to 20°C for GFP mRNA quantification by qRT-PCR. Our results showed that GFP

expression was significantly increased in *CL2070(dvIs70);nmur-1(ok1387)* animals compared to

CL2070(dvIs70) animals (Fig. 4C), indicating that the *nmur-1* mutation causes higher overall

transcriptional activity, which is consistent with the results of our *in vitro* transcription assays.

NMUR-1 regulates the innate immune response to *E. faecalis* by controlling the

expression of C-type lectins

Next, we asked how NMUR-1 regulates gene expression in *C. elegans* in response to *E.*

faecalis infection. By examining the gene expression profiles of *nmur-1(ok1387)* animals relative

to wild-type animals exposed to *E. faecalis*, we found that 229 genes were upregulated and 106

genes were downregulated at least two-fold. GO analysis of the upregulated genes identified 11

significantly enriched biological processes, all of which involve defense responses to bacteria or

other external stimuli (Table S2). This is consistent with the fact that *E. faecalis* is a pathogenic

agent for *C. elegans* (39). GO analysis of the downregulated genes yielded one significantly

enriched molecular function involving carbohydrate binding activity (Table 2A). Fifteen of the

downregulated genes were related to this activity, 14 of which encode C-type lectins (Table 2B).

C-type lectins are a group of proteins that contain one or more C-type lectin-like domains

(CTLDs) and recognize complex carbohydrates on cells and tissues (40,41). C-type lectins may

act in pathogen recognition or function as immune effectors to fight infection (4,11). Therefore,

downregulation of C-type lectins in *nmur-1(ok1387)* animals might explain the reduced survival

observed in mutant animals following exposure to *E. faecalis*.

To determine whether NMUR-1-regulated C-type lectin genes contribute to the reduced survival

of *nmur-1(ok1387)* animals, we inactivated these genes using RNAi-mediated knockdown in

wild-type and *nmur-1(ok1387)* animals and assayed for their survival against *E. faecalis*-

mediated killing. Five of the most downregulated C-type lectin genes (*clec-94, clec-137, clec-*

157, clec-208, and *clec-263*) were individually targeted in these studies. While RNAi of these

genes did not further reduce the survival of *nmur-1(ok1387)* animals (Fig. 5B), knockdown of

four of the five genes (all except *clec-157*) significantly suppressed the survival of wild-type

animals (Fig. 5A). These results indicate that some of the NMUR-1-regulated C-type lectin

genes are indeed required for *C. elegans* defense against *E. faecalis* infection.

Since knockdown of C-type lectins in wild-type animals recaptured the reduced survival

phenotype of *nmur-1(ok1387)* animals, we next asked whether complementing the expression

of these C-type lectins in mutant animals could rescue their survival phenotype. To this end, we

employed transgenes to express *clec-94*, *clec-208*, and *clec-263*, respectively, in *nmur-*

1(ok1387) and wild-type animals. Survival assays against *E. faecalis* showed that while

overexpression of these C-type lectins had no effect on wild-type animal survival, the

expression of *clec-94* or *clec-208* partially rescued the reduced survival of the mutant animals

(Fig. 5C), confirming that the downregulation of C-type lectins contributes to the reduced

survival of *nmur-1* mutants challenged with *E. faecalis*.

NMUR-1 suppresses the innate immune response to *S. enterica* by inhibiting unfolded

protein response (UPR) genes

We also compared the gene expression profiles of *nmur-1(ok1387)* animals with those of wild-

type animals exposed to *S. enterica* and found that 28 genes were upregulated and 28 genes

were downregulated at least two-fold. While GO analysis of the downregulated genes did not

yield any enriched GO terms, a similar analysis of the upregulated genes identified four

significantly enriched biological processes and two significantly enriched molecular functions

(Table 3A). Among these enrichments, the biological process "protein quality control for

misfolded or incompletely synthesized proteins" was the most significantly enriched (FDR =

7.89E-5) and has a known role in immune responses to pathogen infection (42). Under an

environmental stress, such as pathogen infection, increased demand for protein folding causes

ER stress, which activates the UPR pathways to alleviate the stress and maintain protein

homeostasis. This process is known to be controlled by the nervous system (43). Four of the

upregulated genes (*T20D4.3*, *T20D4.5*, *C17B7.5*, and *npl-4.2*) are related to UPR activity and

contributed to the biological process enrichment (Table 3B). To determine whether these

NMUR-1-regulated UPR genes play any roles in the enhanced survival of *nmur-1(ok1387)*

animals, we inactivated these genes individually using RNAi and then assayed the animals for

survival against *S. enterica*. Because these genes are mainly expressed in neurons

(Wormbase), we employed strain *TU3401*, which is capable of RNAi-mediated gene knockdown

in neurons (44), in our experiments. While RNAi silencing of *T20D4.3* specifically suppressed

the enhanced survival of *nmur-1(ok1387);TU3401* animals following exposure to *S. enterica*,

RNAi of *npl-4.2* suppressed the survival of both TU3401 and *nmur-1(ok1387);TU3401* animals

(Fig. 6). In comparison, RNAi of *T20D4.5* or *C17B7.5* had no effect on animal survival (Fig. 6).

The gene *npl-4.2* has been previously implicated in ER-associated protein degradation.

Specifically, NPL-4 forms a complex with CDC-48 ATPase and UFD-1, which can extract

polyubiquitin-tagged misfolded proteins from the ER membrane and facilitate their degradation

by the proteasome (45,46). Less is known about *T20D4.3*, except that it reportedly encodes a

peptide N glycanase and functions in longevity, dauer formation, and pathogen responses (47).

Nonetheless, our results indicate that these UPR genes are important for *C. elegans* defense

against *S. enterica*, and that NMUR-1 can suppress the innate immune response to *S. enterica*

by inhibiting UPR genes.

The NMUR-1 ligand CAPA-1 is required for *C. elegans* defense against *E. faecalis* but not

S. enterica

CAPA-1, a homolog of vertebrate NMU and insect *capability* peptides, is an endogenous ligand

of NMUR-1 (48,49). To determine whether CAPA-1 plays any roles in *C. elegans* defense, we

examined the survival of a loss-of-function mutant of CAPA-1, *capa-1(ok3065)*, after *E. faecalis*

exposure. Compared to wild-type animals, *capa-1(ok3065)* animals showed reduced survival, a

phenotype also displayed by *nmur-1(ok1387)* animals (Fig. 7A). We then crossed *capa-*

1(ok3065) with *nmur-1(ok1387)* animals to generate *capa-1(ok3065);nmur-1(ok1387)* double

mutants. These double mutants exhibited reduced survival against *E. faecalis*, similar to *nmur-*

1(ok1387) animals (Fig. 7A). These results indicate that CAPA-1 and NMUR-1 likely function in

the same pathway to mediate *C. elegans* defense against *E. faecalis*. To investigate whether

CAPA-1 is a ligand of NMUR-1 in the defense response, we tested whether exogenous

administration of a synthetic CAPA-1 peptide could rescue the reduced survival phenotype of

the mutant animals. To this end, we exposed wild-type and mutant animals to *E. faecalis* on BHI

media containing 1 μg/mL synthetic CAPA-1 peptide and scored their survival over time. Our

results showed that exogenous CAPA-1 peptide rescued the reduced survival phenotype of

capa-1(ok3065) animals but failed to rescue *nmur-1(ok1387)* animals or *capa-1(ok3065);nmur-*

1(ok1387) double mutants (Fig. 7B). These results demonstrate that the function of NMUR-1 in

defense depends on the presence of CAPA-1, and *vice versa*, indicating that CAPA-1/NMUR-1

function as a ligand/receptor pair in mediating *C. elegans* defense against *E. faecalis*.

To determine whether CAPA-1 also functions in the *C. elegans* defense response to *S. enterica*,

we subjected wild-type and mutant animals to survival assays against *S. enterica*. Results

showed that while *capa-1(ok3065)* animals displayed survival at the wild-type level, *capa-*

1(ok3065);nmur-1(ok1387) double mutants exhibited survival comparable to that of *nmur-*

1(ok1387) animals (Fig. 7C). This indicates that lacking CAPA-1 has no influence on *C. elegans*

defense against *S. enterica*, and that CAPA-1 might not be involved in the NMUR-1-mediated

innate immune response to this particular pathogen. Although CAPA-1 is the only NMUR-1

ligand identified so far (48,49), these results suggest that other ligands may exist, and that

NMUR-1 could bind to different ligands to regulate immune responses to different pathogens.

DISCUSSION

In the current study, we have shown that *C. elegans* lacking the neuronal GPCR NMUR-1

exhibited altered survival against various bacterial pathogens. By focusing on two pathogens, *E.*

faecalis and *S. enterica*, as infecting agents, we have demonstrated that the altered survival

was not caused by pathogen avoidance behavior but by differences in intestinal accumulation of

pathogens, likely due to changes in innate immunity. Our RNA-seq analyses showed that

functional loss of NMUR-1 negatively affects the expression of transcription factors involved in

binding to RNA polymerase II regulatory regions, which, in turn, impacts transcriptional activity

in *C. elegans*, as demonstrated by both *in vitro* and *in vivo* transcription assays. We also found

that NMUR-1 regulates the expression of different sets of genes in *C. elegans* in response to *E.*

faecalis and *S. enterica* exposure. In particular, C-type lectin genes were downregulated in

nmur-1(ok1387) animals relative to wild-type animals exposed to *E. faecalis*, whereas UPR

genes were upregulated following exposure to *S. enterica*. These genes have been previously

implicated in conserved immune signaling pathways in response to pathogen infection (11).

Functional assays with selected genes revealed that at least three C-type lectin genes (*clec-94*,

clec-208, and *clec-263*) and two UPR genes (*T20D4.3* and *npl-4.2*) are important for *C. elegans*

defense against *E. faecalis* and *S. enterica*, respectively. Taken together, we have

demonstrated that NMUR-1 modulates *C. elegans* transcriptional activity by regulating the

expression of transcription factors, which, in turn, controls the expression of distinct immune

genes in response to different pathogens. Overall, our study has uncovered a molecular basis

for the specificity of *C. elegans* innate immunity. Given the evolutionary conservation of

NMU/NMUR-1 signaling in immune regulation across multicellular organisms (29), our work

could provide mechanistic insights into understanding the specificity of innate immunity in other

animals, including vertebrates.

In vertebrates, a diverse array of PRRs that recognize various PAMPs have been implicated in

the specificity of the innate immune system (5). Although *C. elegans* does not possess the

majority of known PRRs, it can differentiate between different pathogens and launch distinct

innate immune responses to specific pathogens (12-14). Recent behavior research in *C.*

elegans indicates that its nervous system might play a major role in pathogen detection (50).

The nematode nervous system contains 32 chemosensory neurons (out of a total of 302

neurons in hermaphrodites), many of which have cilia exposed to the external environment that

can directly sense pathogen-derived molecules in the environment (51). Moreover, *C. elegans*

expresses approximately 1,300 GPCRs that may function as chemoreceptors in sensory

neurons to detect pathogen cues, thus contributing to the specificity of innate immunity (22). In

this context, a relevant question would be whether neuronal NMUR-1 mediates the specificity of

innate immunity by directly interacting with bacterial molecules. We have shown that NMUR-1

has diverse effects on *C. elegans* defense against various bacterial pathogens (Fig. 1 and Fig.

S1). If the hypothesis that NMUR-1 directly interacts with pathogens is correct, then this GPCR

should be able to bind to various pathogen-derived molecules. Such scenario seems unlikely

given that PRR-PAMP interactions follow a simple key-lock system, i.e., each receptor matches

one pathogen-associated molecule or molecular pattern (9). Therefore, it is more likely that

NMUR-1 mediates the ability to differentiate between pathogens by controlling neural signaling

than by recognizing bacterial molecules directly. Indeed, we have found that NMUR-1 regulates

C-type lectin genes in response to *E. faecalis* infection and UPR genes in response to *S.*

enterica infection; both of these genes are known to function in innate immune signaling

pathways that fight infection (11).

How does NMUR-1, a single type of neuronal GPCR, distinguish different pathogens so that it

can respond to some attacks with suppression of the innate immune response and amplify the

response to others? A basic principle in neuroscience is that the interaction of multiple neural

circuits mediates the functional output of a neural network and determines the final state of

organ function (52). Accordingly, the net effect of neural control on the immune response

depends on the total sum of the interactions resulting from ligand-receptor signal transduction in

a specific path. NMUR-1 is normally expressed in the spermathecae of the somatic gonad and

several different types of sensory neurons as well as in interneurons (27). By using a

fluorescent reporter transgene, we observed that distinct subsets of NMUR-1-expressing

neurons are activated upon exposure to *E. faecalis* and *S. enterica* (data not shown), indicating

that different neural circuits are involved in regulating the immune responses to these two

pathogens. This notion is further supported by the fact that CAPA-1, the only NMUR-1 ligand

identified so far, is required for the nematode immune response to *E. faecalis* but is dispensable

for its response to *S. enterica* (Fig. 7). Future studies to decipher these different neuro-immune

regulatory circuits would help us understand how signals pass through different components of

the circuits to exert a net effect on immunity. An associative learning circuit mediated by CAPA-

1/NMUR-1 signaling was recently uncovered in *C. elegans* in which sensory ASG neurons

release CAPA-1, which signals via NMUR-1 in AFD sensory neurons to specifically mediate the

retrieval of learned salt avoidance (49). In mice, cholinergic neurons in the intestine can sense

and respond to infection by expressing NMU, which directly activates type 2 innate lymphoid

cells in the gastrointestinal tract through NMUR1 to drive anti-parasitic immunity (30-32). These

studies provide useful information for deciphering NMUR-1-mediated neuro-immune regulatory

circuits in *C. elegans*, which will be critical for further understanding how the nervous system

regulates innate immunity to generate specificity by either suppressing or amplifying immune

responses to different pathogens.

MATERIALS AND METHODS

Nematode strains

The following *C. elegans* strains were cultured under standard conditions and fed *E. coli* HT115.

Wild-type N2 and *nmur-1(ok1387)* were gifts from Dr. Joy Alcedo (Wayne State University,

Detroit, MI). The *capa-1(ok3065)*, *CL2070(dvIs70)*, and *TU3401* strains were obtained from the

Caenorhabditis elegans Genetics Center (University of Minnesota, Minneapolis, MN). The *capa-*

1(ok3065);nmur-1(ok1387), *nmur-1(ok1387);CL2070(dvIs70)*, and *nmur-1(ok1387);TU3401*

mutant strains were constructed using standard genetic techniques.

Bacterial strains

476 The following bacterial strains were grown using standard conditions (53): *Escherichia coli*

477 HT115, *Escherichia coli* OP50, *Salmonella enterica* strain SL1344, *S. enterica SL1344::gfp*,

478 *Pseudomonas aeruginosa* strain PA14, *Yersinia pestis* strain kim5, *Enterococcus faecalis* strain

479 OG1RF, *E. faecalis OG1RF::gfp*, *Microbacterium nematophilum* strain CBX102, and

480 *Staphylococcus aureus* strain NCTC 8325.

Survival assay

C. elegans wild-type *N2* animals and mutant animals were maintained as hermaphrodites at

20°C, grown on modified NGM with 30 units/mL nystatin and 15μg/mL tetracycline, and fed *E.*

coli HT115. Bacterial lawns for the survival assays were prepared by placing a 30μL drop of an

overnight culture of pathogenic bacteria on each 3.5cm plate. Full-lawn survival assays were

prepared by spreading 30μL of overnight bacterial culture over the complete surface of each

3.5cm plate. Plates were incubated at 37°C for 15~16 hours, cooled down to room temperature,

and then seeded with synchronized 65-hour-old young adult animals. The survival assays were

performed at 25°C, and live animals were transferred daily to fresh plates until egg laying

ceased. Animals were scored at the times indicated and were considered dead when they failed

to respond to touch.

Lawn occupancy assay

Wild-type and mutant animals were maintained as hermaphrodites at 20°C, grown on modified

NGM with 30 units/mL nystatin and 15μg/mL tetracycline and fed *E. coli* HT115. The bacterial

lawns for the lawn occupancy assays were prepared by placing a 30µL drop of an overnight

culture of pathogenic bacteria on each 3.5cm agar plate. Plates were incubated at 37°C for

15~16 hours, cooled down to room temperature, and then seeded with synchronized 65-hour-

old young adult animals. The assays were performed at 25°C and animals were scored at the

described time points for lawn occupancy. Dead animals were censored from the results.

Profiling bacterial accumulation in the nematode intestine

Wild-type and *nmur-1(ok1387)* animals were synchronized by egg-laying. The animals were

grown on *E. coli* HT115 at 20°C for 65 hours or until the young adult stage. The animals were

then transferred to plates seeded with GFP-tagged bacteria and incubated for 24 hours at 25°C.

Following incubation, the animals were transferred to an empty NGM plate for 15 minutes to

eliminate any fluorescent bacteria stuck to the body. The animals were then transferred to a

new, empty NGM plate and into a droplet of M9 buffer to further wash away any bacteria

remaining on the body. Animals were anesthetized using 25mM Sodium Azide and visualized

using a Zeiss Axio Imager M2 stereoscopic microscope.

Quantification of intestinal colonization

Wild-type and *nmur-1(ok1387)* animals were synchronized by egg-laying. Sixty-five-hour-old

animals were transferred to plates seeded with GFP-tagged bacteria and incubated for 24 hours

at 25°C. Ten animals were then transferred to a 1.5mL tube containing a solution of 25mM

Sodium Azide with 1mg/mL ampicillin and kanamycin. The animals were soaked for 45 minutes

to remove any external bacteria. After 45 minutes, the animals were washed three times with

PBS containing 0.1% Triton to remove the remaining antibiotics, and ground in 100µL of PBS

with 0.1% Triton. Serial dilutions of the lysates (10^{-1}, 10^{-2}, 10^{-3}, 10^{-4}, and 10^{-5}) were plated with

rifampicin to select for GFP-tagged bacteria and the plates were incubated at 37°C for 24 hours.

Profiling pumping and defecation rates

Wild-type and *nmur-1(ok1387)* animals were synchronized by egg-laying. Sixty-five-hour-old

animals were transferred to plates seeded with pathogenic bacteria and incubated for 24 hours

at 25°C. Pumping rates were counted by observing the contraction/relaxation cycles of the

terminal pharyngeal bulb using a stereoscopic microscope. Cycles were counted for 30 second

durations, three times for ten individual animals per condition. Defecation rates were measured

by timing the span between the contraction of the rectal muscles and subsequent excretion

using a stereoscopic microscope. Cycles were counted five times for ten individual animals per

condition.

RNA interference

RNAi was conducted by feeding synchronized L3 larval *C. elegans E. coli* strain HT115(DE3)

expressing double-stranded RNA (dsRNA) that was homologous to a target gene (54,55). *E.*

coli with the appropriate dsRNA vector were grown in LB broth containing ampicillin (100µg/mL)

at 37°C for 16 hours and plated on modified NGM plates containing 100µg/mL ampicillin and

3mM isopropyl β-D-thiogalactoside (IPTG). The bacteria were allowed to grow for 16 hours at

37°C. The plates were cooled away from direct light before the synchronized L3 larval animals

were placed on the bacteria. The animals were incubated at 20°C for 24 hours or until the

animals were 65 hours old. *unc-22* RNAi was included as a positive control in all experiments to

account for RNAi efficiency.

Synthetic NMUR-1 ligand administration

Wild-type and mutant animals were maintained as hermaphrodites at 20°C, grown on modified

NGM with 30 units/mL nystatin and 15µg/mL tetracycline, and fed *E. coli* HT115. The animals

were allowed to lay eggs for 45 minutes at 25°C on modified NGM plates seeded with *E.coli*

HT115. The synchronized 65-hour young adult animals were transferred to BHI medium

containing 0, 1, or 5µg/mL oligopeptide AFFYTPRI-NH2 and seeded with 30µL of fresh

pathogen broth that had been incubated at 37°C for 16 hours. The survival assays were

performed at 25°C and live animals were transferred daily to fresh plates until egg laying

ceased. The animals were scored at the times indicated and were considered dead when they

failed to respond to touch.

555   RNA isolation

Gravid adult wild-type and *nmur-1(ok1387)* animals were lysed using a solution of sodium

hydroxide and bleach (ratio 5:2), washed, and eggs were synchronized for 22 hours in S-basal

liquid medium at room temperature. Synchronized L1 larval animals were transferred onto NGM

plates seeded with *E. coli* HT115 and grown at 20°C for 48 hours until the animals had reached

the L4 larval stage. The animals were collected and washed with M9 buffer before being

transferred to plates containing *E. coli* HT115 or pathogenic bacteria for 24 hours at 25°C. After

24 hours, animals were collected and washed with M9 buffer, and RNA was extracted using

QIAzol (Qiagen). Total RNA was isolated using the RNeasy mini kit (Qiagen) following the

protocol provided by the manufacturer.

566   Quantitative real-time PCR (qRT-PCR)

Total RNA was obtained as described above. Two µg of RNA were used to generate cDNA

using the Applied Biosystems High-Capacity cDNA Reverse Transcription Kit. qRT-PCR was

conducted by following the prescribed protocol for PowerUp SYBR Green (Applied Biosystems)

on an Applied Biosystems StepOnePlus real-time PCR machine. Ten-µL reactions were set up

following the manufacturer's recommendations, and 20ng of cDNA were used per reaction.

Relative fold-changes for transcripts were calculated using the comparative CT(2-ΔΔCT)

method and were normalized to pan-actin (act-1, -3, -4). Amplification cycle thresholds were

determined by the StepOnePlus software. All samples were run in triplicate.

RNA sequencing

Total RNA samples were obtained as described above and was submitted to the WSU

Genomics Core for RNA-seq analyses. RNA-seq and related bioinformatic analyses were done

following our published protocol (26). The raw sequence data (FASTQ files) were deposited in

NCBI's SRA database through the GEO. The processed gene quantification files and differential

expression files were deposited in GEO. All of these data can be accessed through GEO with

the accession number GSE154324.

In vitro transcription assay

In vitro transcription assays were performed following our published protocol (37). Briefly, worm

nuclear extracts were prepared from synchronized young adult animals using Balch

homogenization. The nuclear extracts were then incubated with PESDNA (a linear DNA

template containing the worm *Δpes-10* promoter) and other transcription components at 30°C

for 30 minutes, followed by RNA purification and reverse transcription. The resulting cDNA was

either amplified by PCR and detected by gel electrophoresis or quantified by qRT-PCR using

the PowerUp SYBR green qPCR kit (Applied Biosystems, catalog # A25918).

Plasmid construction and transgenic animal generation

A *clec-94* genomic DNA fragment (4,247 bp) was amplified by PCR from mixed stage wild-type

C. elegans using the following primers: 5'- ATTGTCGACGTACAGTCCCACTACTTGTTGC -3'

and 5'- CTAGGATCCCTAACATTGGCCGGGAAAGAG -3'. PCR products were then cloned into

the *pPD95.77* vector (Fire Lab *C. elegans* vector kit; Addgene, Cambridge, MA) via the *SalI* and

BamHI sites. The resulting plasmid pPW01 (*clec-94p::clec-94::SL2::gfp*) was microinjected into

wild-type animals at 10ng/μL to generate strain JRS60 (Table S3). The *clec-94* genomic rescue

strain JRS63 (Table S3) was generated by crossing JRS60 with *nmur-1(ok1387)* animals.

A *clec-208* genomic DNA fragment (3,556 bp) was amplified by PCR from mixed stage wild-type

C. elegans using the following primers: 5'- TTTGTCGACTTTCTGTTCTTGCTACTCTCTACC -3'

and 5'- TTTTCTAGACTGGCTCGTTCTTAGAGACC -3'. PCR products were then cloned into

the *pPD95.77* vector via the *SalI* and *XbaI* sites. The resulting plasmid, pPW02 (*clec-208p::clec-*

208::SL2::gfp), was microinjected into wild-type animals at 50ng/μL with *unc-122p::mCherry* at

20ng/μl as a co-injection marker to generate strain JRS61 (Table S3). The *clec-208* genomic

rescue strain JRS64 (Table S3) was generated by crossing JRS61 with *nmur-1(ok1387)*

animals.

A *clec-263* genomic DNA fragment (5,389 bp) was amplified by PCR from mixed stage wild-type

C. elegans using the following primers: 5'- TTTGTCGACGCGATGGGTGGTTGTATTATTC -3'

and 5'- CAAGGATCCAAATCGTATTTCCGTCGTTGGTC -3'. PCR products were then cloned

into the *pPD95.77* vector via the *SalI* and *BamHI* sites. The resulting plasmid, pPW03 (*clec-*

263p::clec-263::SL2::gfp), was microinjected into wild-type animals at 30ng/μL with *unc-*

122p::mCherry at 20ng/μl as a co-injection marker to generate strain JRS62 (Table S3). The

clec-263 genomic rescue strain JRS65 (Table S3) was generated by crossing JRS62 with *nmur-*

1(ok1387) animals.

Statistical analysis

Survival curves were plotted using GraphPad PRISM (version 8) computer software. Survival

was considered different from the appropriate control indicated in the main text when $P < 0.05$.

PRISM uses the product limit or Kaplan-Meier method to calculate survival fractions and the log-

rank test, which is equivalent to the Mantel-Haenszel test, to compare survival curves.

Occupancy assays, CFU assays, and qRT-PCR results were analyzed using two-sample t-tests

for independent samples; P-values <0.05 are considered significant. All experiments were

repeated at least three times, unless otherwise indicated.

ACKNOWLEDGEMENTS

We thank Dr. Joy Alcedo at Wayne State University for constructive discussions and for

providing us with wild-type *N2* and *nmur-1(ok1387)* worm strains. We thank Dr. Danielle Garsin

at the University of Texas Health Science Center at Houston for providing us with the *E. faecalis*

OG1RF::gfp strain. Some worm strains were provided by the *Caenorhabditis* Genetics Center,

which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440).

FUNDING

This work was supported by the Department of Biomedical Sciences, Elson S. Floyd College of

Medicine, WSU-Spokane (Startup to Y.L.), the Research Foundation Flanders (G093419N to

I.B.), and the NIH (R35GM124678 to J.S.). The funders had no role in study design, data

collection and interpretation, or the decision to submit the work for publication.

AUTHOR CONTRIBUTIONS

P.W., S.W., J.W., C.C., D.S., Y.L., and I.B. designed and performed experiments and analyzed

data. J.S. designed experiments and analyzed data. Y.L., P.W., and J.S. wrote the paper.

COMPETING INTERESTS

The authors declare that they have no competing interests.

DATA AND MATERIALS AVAILABILITY

The RNA-seq data have been deposited in NCBI's SRA database through the GEO. The

processed gene quantification files and differential expression files have been deposited in the

GEO. All of the data can be accessed through the GEO with the accession number

GSE154324. All data needed to evaluate the conclusions in the paper are present in the paper

and/or the Supplementary Materials. Additional data related to this paper may be requested

from the authors. The *C. elegans* strains and plasmids constructed by us and primer sequences

used for their construction are available upon request.

REFERENCES

1.   Medzhitov, R. (2009) Approaching the asymptote: 20 years later. *Immunity* **30**, 766-775

2.   Jones, J. D., and Dangl, J. L. (2006) The plant immune system. *Nature* **444**, 323-329

3.   Chaplin, D. D. (2010) Overview of the immune response. *J Allergy Clin Immunol* **125**, S3-23

4.   Pees, B., Yang, W., Zarate-Potes, A., Schulenburg, H., and Dierking, K. (2016) High Innate

Immune Specificity through Diversified C-Type Lectin-Like Domain Proteins in Invertebrates. *J*

Innate Immun **8**, 129-142

5.   Tan, R. S., Ho, B., Leung, B. P., and Ding, J. L. (2014) TLR cross-talk confers specificity to innate

immunity. *Int Rev Immunol* **33**, 443-453

6.   Sternberg, E. M. (2006) Neural regulation of innate immunity: a coordinated nonspecific host

response to pathogens. *Nature reviews. Immunology* **6**, 318-328

7.   Steinman, L. (2004) Elaborate interactions between the immune and nervous systems. *Nature*

immunology **5**, 575-581

8.   Cherry, S., and Silverman, N. (2006) Host-pathogen interactions in drosophila: new tricks from

an old friend. *Nature immunology* **7**, 911-917

9.   Schulenburg, H., Boehnisch, C., and Michiels, N. K. (2007) How do invertebrates generate a

highly specific innate immune response? *Molecular immunology* **44**, 3338-3344

10.   Sharrock, J., and Sun, J. C. (2020) Innate immunological memory: from plants to animals. *Current*

opinion in immunology **62**, 69-78

11.   Sun, J., Aballay, A., and Singh, V. (2016) Cellular Responses to Infections in *Caenorhabditis*

elegans. *Encyclopedia of Cell Biology* **2**, 845-852

12.   Irazoqui, J. E., Troemel, E. R., Feinbaum, R. L., Luhachack, L. G., Cezairliyan, B. O., and Ausubel, F.

M. (2010) Distinct pathogenesis and host responses during infection of C. elegans by P.

aeruginosa and S. aureus. *PLoS pathogens* **6**, e1000982

13.   Zarate-Potes, A., Yang, W., Pees, B., Schalkowski, R., Segler, P., Andresen, B., Haase, D., Nakad,

R., Rosenstiel, P., Tetreau, G., Colletier, J. P., Schulenburg, H., and Dierking, K. (2020) The C.

elegans GATA transcription factor elt-2 mediates distinct transcriptional responses and opposite

infection outcomes towards different Bacillus thuringiensis strains. *PLoS pathogens* **16**,

e1008826

14.   Wong, D., Bazopoulou, D., Pujol, N., Tavernarakis, N., and Ewbank, J. J. (2007) Genome-wide

investigation reveals pathogen-specific and shared signatures in the response of Caenorhabditis

elegans to infection. *Genome biology* **8**, R194

15.   Kawli, T., and Tan, M. W. (2008) Neuroendocrine signals modulate the innate immunity of

Caenorhabditis elegans through insulin signaling. *Nature immunology* **9**, 1415-1424

16.   Styer, K. L., Singh, V., Macosko, E., Steele, S. E., Bargmann, C. I., and Aballay, A. (2008) Innate

immunity in Caenorhabditis elegans is regulated by neurons expressing NPR-1/GPCR. *Science*

322, 460-464

17.   Reddy, K. C., Andersen, E. C., Kruglyak, L., and Kim, D. H. (2009) A polymorphism in npr-1 is a behavioral determinant of pathogen susceptibility in C. elegans. *Science* **323**, 382-384

18.   Anyanful, A., Easley, K. A., Benian, G. M., and Kalman, D. (2009) Conditioning protects C. elegans from lethal effects of enteropathogenic E. coli by activating genes that regulate lifespan and innate immunity. *Cell host & microbe* **5**, 450-462

19.   Sun, J., Singh, V., Kajino-Sakamoto, R., and Aballay, A. (2011) Neuronal GPCR controls innate immunity by regulating noncanonical unfolded protein response genes. *Science* **332**, 729-732

20.   Zhang, Y., Lu, H., and Bargmann, C. I. (2005) Pathogenic bacteria induce aversive olfactory learning in Caenorhabditis elegans. *Nature* **438**, 179-184

21.   Pradel, E., Zhang, Y., Pujol, N., Matsuyama, T., Bargmann, C. I., and Ewbank, J. J. (2007) Detection and avoidance of a natural product from the pathogenic bacterium Serratia marcescens by Caenorhabditis elegans. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 2295-2300

22.   Liu, Y., and Sun, J. (2021) Detection of pathogens and regulation of immunity by the *C. elegans* nervous system. *MBio*

23.   Sun, J., Liu, Y., and Aballay, A. (2012) Organismal regulation of XBP-1-mediated unfolded protein response during development and immune activation. *EMBO reports* **13**, 855-860

24.   Liu, Y., Sellegounder, D., and Sun, J. (2016) Neuronal GPCR OCTR-1 regulates innate immunity by controlling protein synthesis in *Caenorhabditis elegans*. *Sci Rep* **6**, 36832

25.   Sellegounder, D., Yuan, C. H., Wibisono, P., Liu, Y., and Sun, J. (2018) Octopaminergic Signaling Mediates Neural Regulation of Innate Immunity in Caenorhabditis elegans. *MBio* **9**

26.   Sellegounder, D., Liu, Y., Wibisono, P., Chen, C. H., Leap, D., and Sun, J. (2019) Neuronal GPCR NPR-8 regulates *C. elegans* defense against pathogen infection. *Sci Adv* **5**, eaaw4717

27.   Maier, W., Adilov, B., Regenass, M., and Alcedo, J. (2010) A neuromedin U receptor acts with the

719   sensory system to modulate food type-dependent effects on C. elegans lifespan. *PLoS biology* **8**,

720   e1000376

721   28.   Martinez, V. G., and O'Driscoll, L. (2015) Neuromedin U: a multifunctional neuropeptide with

722   pleiotropic roles. *Clin Chem* **61**, 471-482

723   29.   Ye, Y., Liang, Z., and Xue, L. (2021) Neuromedin U: potential roles in immunity and inflammation.

724   *Immunology* **162**, 17-29

725   30.   Cardoso, V., Chesne, J., Ribeiro, H., Garcia-Cassani, B., Carvalho, T., Bouchery, T., Shah, K.,

726   Barbosa-Morais, N. L., Harris, N., and Veiga-Fernandes, H. (2017) Neuronal regulation of type 2

727   innate lymphoid cells via neuromedin U. *Nature* **549**, 277-281

728   31.   Klose, C. S. N., Mahlakoiv, T., Moeller, J. B., Rankin, L. C., Flamar, A. L., Kabata, H., Monticelli, L.

729   A., Moriyama, S., Putzel, G. G., Rakhilin, N., Shen, X., Kostenis, E., Konig, G. M., Senda, T.,

730   Carpenter, D., Farber, D. L., and Artis, D. (2017) The neuropeptide neuromedin U stimulates

731   innate lymphoid cells and type 2 inflammation. *Nature* **549**, 282-286

732   32.   Wallrapp, A., Riesenfeld, S. J., Burkett, P. R., Abdulnour, R. E., Nyman, J., Dionne, D., Hofree, M.,

733   Cuoco, M. S., Rodman, C., Farouq, D., Haas, B. J., Tickle, T. L., Trombetta, J. J., Baral, P., Klose, C.

734   S. N., Mahlakoiv, T., Artis, D., Rozenblatt-Rosen, O., Chiu, I. M., Levy, B. D., Kowalczyk, M. S.,

735   Regev, A., and Kuchroo, V. K. (2017) The neuropeptide NMU amplifies ILC2-driven allergic lung

736   inflammation. *Nature* **549**, 351-356

737   33.   Mazgaeen, L., and Gurung, P. (2020) Recent Advances in Lipopolysaccharide Recognition

738   Systems. *Int J Mol Sci* **21**

739   34.   Peterson, J. W. (1996) Bacterial Pathogenesis. in *Medical Microbiology* (Baron, S. ed.), 4th Ed.,

740   Galveston (TX). pp

35. Meisel, J. D., and Kim, D. H. (2014) Behavioral avoidance of pathogenic bacteria by

Caenorhabditis elegans. *Trends Immunol* **35**, 465-470

36. Fuhrman, L. E., Goel, A. K., Smith, J., Shianna, K. V., and Aballay, A. (2009) Nucleolar proteins

suppress Caenorhabditis elegans innate immunity by inhibiting p53/CEP-1. *PLoS genetics* **5**,

e1000657

37. Wibisono, P., Liu, Y., and Sun, J. (2020) A novel in vitro Caenorhabditis elegans transcription

system. *BMC Mol Cell Biol* **21**, 87

38. Link, C. D., Cypser, J. R., Johnson, C. J., and Johnson, T. E. (1999) Direct observation of stress

response in Caenorhabditis elegans using a reporter transgene. *Cell Stress Chaperones* **4**, 235-

242

39. Yuen, G. J., and Ausubel, F. M. (2018) Both live and dead Enterococci activate Caenorhabditis

elegans host defense via immune and stress pathways. *Virulence* **9**, 683-699

40. Drickamer, K., and Dodd, R. B. (1999) C-Type lectin-like domains in Caenorhabditis elegans:

predictions from the complete genome sequence. *Glycobiology* **9**, 1357-1369

41. Takeuchi, T., Sennari, R., Sugiura, K., Tateno, H., Hirabayashi, J., and Kasai, K. (2008) A C-type

lectin of Caenorhabditis elegans: its sugar-binding property revealed by glycoconjugate

microarray analysis. *Biochemical and biophysical research communications* **377**, 303-306

42. Grootjans, J., Kaser, A., Kaufman, R. J., and Blumberg, R. S. (2016) The unfolded protein response

in immunity and inflammation. *Nature reviews. Immunology* **16**, 469-484

43. Aballay, A. (2013) Role of the nervous system in the control of proteostasis during innate

immune activation: insights from C. elegans. *PLoS pathogens* **9**, e1003433

44. Calixto, A., Chelur, D., Topalidou, I., Chen, X., and Chalfie, M. (2010) Enhanced neuronal RNAi in

C. elegans using SID-1. *Nature methods* **7**, 554-559

45.   Ye, Y., Meyer, H. H., and Rapoport, T. A. (2001) The AAA ATPase Cdc48/p97 and its partners

765   transport proteins from the ER into the cytosol. *Nature* **414**, 652-656

766   46.   Bays, N. W., Wilhovsky, S. K., Goradia, A., Hodgkiss-Harlow, K., and Hampton, R. Y. (2001)

767   HRD4/NPL4 is required for the proteasomal processing of ubiquitinated ER proteins. *Mol Biol*

768   *Cell* **12**, 4114-4128

769   47.   Gaglia, M. M., Jeong, D. E., Ryu, E. A., Lee, D., Kenyon, C., and Lee, S. J. (2012) Genes that act

770   downstream of sensory neurons to influence longevity, dauer formation, and pathogen

771   responses in Caenorhabditis elegans. *PLoS genetics* **8**, e1003133

772   48.   Lindemans, M., Janssen, T., Husson, S. J., Meelkop, E., Temmerman, L., Clynen, E., Mertens, I.,

773   and Schoofs, L. (2009) A neuromedin-pyrokinin-like neuropeptide signaling system in

774   Caenorhabditis elegans. *Biochemical and biophysical research communications* **379**, 760-764

775   49.   Watteyne, J., Peymen, K., Van der Auwera, P., Borghgraef, C., Vandewyer, E., Van Damme, S.,

776   Rutten, I., Lammertyn, J., Jelier, R., Schoofs, L., and Beets, I. (2020) Neuromedin U signaling

777   regulates retrieval of learned salt avoidance in a C. elegans gustatory circuit. *Nat Commun* **11**,

778   2076

779   50.   Kim, D. H., and Flavell, S. W. (2020) Host-microbe interactions and the behavior of

780   Caenorhabditis elegans. *J Neurogenet*, 1-10

781   51.   Bargmann, C. I. (2006) Chemosensation in C. elegans. *WormBook : the online review of C.*

782   *elegans biology*, 1-29

783   52.   Tracey, K. J. (2014) Approaching the next revolution? Evolutionary integration of neural and

784   immune pathogen sensing and response. *Cold Spring Harb Perspect Biol* **7**, a016360

785   53.   Lagier, J. C., Edouard, S., Pagnier, I., Mediannikov, O., Drancourt, M., and Raoult, D. (2015)

786   Current and past strategies for bacterial culture in clinical microbiology. *Clin Microbiol Rev* **28**,

787   208-236

788   54.   Fraser, A. G., Kamath, R. S., Zipperlen, P., Martinez-Campos, M., Sohrmann, M., and Ahringer, J.

789   (2000) Functional genomic analysis of C. elegans chromosome I by systematic RNA interference.

790   *Nature* **408**, 325-330

791   55.   Timmons, L., and Fire, A. (1998) Specific interference by ingested dsRNA. *Nature* **395**, 854

 FIGURE LEGENDS

Fig. 1. Functional loss of NMUR-1 differentially affects *C. elegans* survival against *E. coli*

HT115, *E. faecalis* and *S. enterica*. Wild-type (WT) and *nmur-1(ok1387)* animals were

exposed to *E. coli* HT115 **(A)**, *E. faecalis* **(B)**, or *S. enterica* **(C)** and scored for survival over

time. Each graph is a combination of three independent experiments. Each experiment included

N = 60 animals per strain. *P*-values represent the significance level of mutant survival relative to

WT, *P* = 0.8706 in (A), *P* = 0.0004 in (B), and *P* < 0.0001 in (C).

Fig. 2. Functional loss of NMUR-1 does not affect pathogen avoidance behavior. WT and

nmur-1(ok1387) animals were subjected to lawn occupancy assays in which the animals were

placed on a small lawn of *E. faecalis* **(A)** or *S. enterica* **(B)** in a 3.5cm plate and monitored over

time for their presence on the lawn. Each graph is a combination of three independent

experiments. Each experiment included *N* = 60 animals per strain. Error bars represent the

standard error of the mean (SEM).

Fig 3. Functional loss of NMUR-1 changes intestinal accumulation of *E. faecalis* and *S.*

***enterica*. (A)** WT and *nmur-1(ok1387)* animals were exposed to GFP-expressing *E. faecalis* for

24 hours and then visualized using a Zeiss Axio Imager M2 fluorescence microscope. The scale

bar indicates 500 μm. **(B)** WT and *nmur-1(ok1387)* animals were exposed to *E. faecalis* for 24

hours, and the Colony Forming Units (CFU) of live bacterial cells recovered from their intestines

were quantified. The graph represents the combined results of three independent experiments.

N = 10 animals per strain were used for each experiment. Error bars represent SEM. *P*-value

represents the significance level of mutant gut colonization relative to WT, *P* = 0.0296. **(C)** WT

and *nmur-1(ok1387)* animals were exposed to *E. faecalis* for 24 hours. Pharyngeal pumping

rates of animals were counted as the number of pumps per 30 sec. Counting was conducted in

triplicate and averaged to give a pumping rate. The measurements included N = 10 adult

animals per strain. Error bars represent SEM. P = 0.1068. **(D)** WT and *nmur-1(ok1387)* animals

were exposed to *E. faecalis* for 24 hours. Defecation rates of animals were measured as the

average time of 10 intervals between two defecation cycles. The measurements included N =

10 animals per strain. P = 0.1379. **(E)** WT and *nmur-1(ok1387)* animals were exposed to GFP-

expressing *S. enterica* for 24 hours and then visualized using a Zeiss Axio Imager M2

fluorescence microscope. The scale bar indicates 500 μm. **(F)** WT and *nmur-1(ok1387)* animals

were exposed to GFP-expressing *S. enterica* for 24 hours, and the CFU of live bacterial cells

recovered from their intestines were quantified. The graph represents the combined results of

three independent experiments. N = 10 animals per strain were used for each experiment. Error

bars represent SEM. P = 0.0290. **(G)** WT and *nmur-1(ok1387)* animals were exposed to *S.*

enterica for 24 hours. Pharyngeal pumping rates of animals were counted as the number of

pumps per 30 sec. Counting was conducted in triplicate and averaged to give a pumping rate.

The measurements included N = 10 adult animals per strain. Error bars represent SEM. P =

0.0006. **(H)** WT and *nmur-1(ok1387)* animals were exposed to *S. enterica* for 24 hours.

Defecation rates of animals were measured as the average time of 10 intervals between two

defecation cycles. The measurements included N = 10 animals per strain. P = 0.6310.

Fig. 4. NMUR-1 regulates transcriptional activity. (A) *In vitro* transcription assays were

performed using nuclear extract from WT or *nmur-1(ok1387)* animals. After transcription, RNA

was purified, reverse transcribed, and amplified by PCR, followed by gel analysis. Lane 1,

transcription with WT nuclear extract; lane 2, transcription with *nmur-1(ok1387)* nuclear extract;

lane 3, transcriptional positive control (with HeLa nuclear extract); lane 4, transcriptional

negative control (without any nuclear extract); lane 5, PCR positive control; lane 6, PCR

negative control (without DNA template). **(B)** WT and mutant nuclear extracts were titrated using

the *in vitro* transcription system. The RNA copy number generated from transcription was

plotted against the amount of nuclear extract used in the reaction, followed by fitting with the

Michaelis-Menten model and the non-linear least-squares method to generate the titration

curves and calculate titration parameters. **(C)** qRT-PCR analysis of GFP expression after two-

hour heat shock in *CL2070(dvIs70)* and *nmur-1(ok1387);CL2070(dvIs70)* animals. Asterisk (*)

denotes a significant difference between *CL2070(dvIs70)* and *nmur-1(ok1387);CL2070(dvIs70)*

animals.

Fig. 5. NMUR-1-regulated C-type lectin genes are required for *C. elegans* defense against

***E. faecalis*. (A)** WT and **(B)** *nmur-1(ok1387)* animals grown on dsRNA for C-type lectin genes

or the empty vector (EV) control were exposed to *E. faecalis* and scored for survival over time.

The graphs are a combination of three independent replicates. Each experiment included *N* =

60 animals per strain. *P*-values in (A) represent the significance level of RNAi treatment relative

to WT+EV: WT+*clec-94*, *P* < 0.0001; WT+*clec-137*, *P* = 0.0220; WT+*clec-157*, *P* = 0.3728;

WT+*clec-208*, *P* < 0.0001; WT+*clec-263*, *P* < 0.0001. *P*-values in (B) represent the significance

level of the RNAi treatment relative to *nmur-1(ok1387)*+EV: *nmur-1(ok1387)+clec-94*, *P* =

0.3633; *nmur-1(ok1387)+clec-137*, *P* = 0.1841; *nmur-1(ok1387)+clec-157*, *P* = 0.8617; *nmur-*

1(ok1387)+clec-208, *P* = 0.1348; *nmur-1(ok1387)+clec-263*, *P* = 0.1320. **(C)** WT and *nmur-*

1(ok1387) animals in which C-type lectin expression was rescued were exposed to *E. faecalis*

and scored for survival over time. All C-type lectin genes were rescued using their own

promoters. The graph is a combination of three independent experiments. Each experiment

included *N* = 60 adult animals per strain. *P*-values represent the significance of the survival of

rescue strains relative to *nmur-1(ok1378)* animals: WT, *P* < 0.0001; nmur-1(ok1387)+*clec-94*, *P*

< 0.0001; *nmur-1(ok1387)+clec-208*, *P* = 0.0025; *nmur-1(ok1387)+clec-263*, *P* = 0.9799.

Fig. 6. NMUR-1-regulated UPR genes are required for *C. elegans* defense against *S.*

***enterica*. (A)** *TU3401* and **(B)** *nmur-1(ok1387);TU3401* animals grown on dsRNA for UPR

genes or the empty vector (EV) control were exposed to *S. enterica* and scored for survival over

time. The graphs are a combination of three independent replicates. Each experiment included

N = 60 animals per strain. *P*-values in (A) represent the significance level of RNAi treatment

relative to *TU3401*+EV: *TU3401+T20D4.3*, *P* = 0.4224; *TU3401+T20D4.5*, *P* = 0.6102;

TU3401+C17B7.5, *P* = 0.5905; *TU3401+npl-4.2*, *P* < 0.0001. *P*-values in (B) represent the

significance level of RNAi treatment relative to *nmur-1(ok1387);TU3401*+EV: *nmur-*

1(ok1387);TU3401+T20D4.3, *P* = 0.0007; *nmur-1(ok1387);TU3401+T20D4.5*, *P* = 0.0993;

nmur-1(ok1387);TU3401+C17B7.5, *P* = 0.1216; *nmur-1(ok1387);TU3401+npl-4.2*; *P* < 0.0001.

Fig. 7. CAPA-1 is required for *C. elegans* defense against *E. faecalis* but not *S. enterica*.

(A) WT, *nmur-1(ok1387)*, *capa-1(ok3065)*, and *capa-1(ok3650);nmur-1(ok1387)* animals were

exposed to *E. faecalis* and scored for survival over time. The graph is representative of three

independent experiments. Each experiment included *N* = 60 animals per strain. *P*-values

represent the significance level of mutant survival relative to WT: *nmur-1(ok1387)*, *P* = 0.0005;

capa-1(ok3065), *P* = 0.0134; *capa-1(ok3065);nmur-1(ok1387)*, *P* = 0.0028. **(B)** WT, *nmur-*

1(ok1387), *capa-1(ok3065)*, and *capa-1(ok3650);nmur-1(ok1387)* animals were exposed to *E.*

faecalis on plates containing either 0 or 1µg/mL synthetic ligand and scored for survival over

time. The graph is a combination of three independent experiments. Each experiment included

N = 60 animals per strain. *P*-values represent the significance level of the survival of

mutant+1µg/mL relative to WT+1µg/mL: *nmur-1(ok1387)*+1µg/mL, *P* < 0.0001; *capa-*

1(ok3065)+1µg/mL, *P* = 0.6758; *capa-1(ok3065);nmur-1(ok1387)*+1µg/mL, *P* = 0.0024. **(C)** WT,

nmur-1(ok1387), *capa-1(ok3065)*, and *capa-1(ok3065);nmur-1(ok1387)* animals were exposed

to *S. enterica* and scored for survival over time. The graph is representative of three

independent experiments. Each experiment included *N* = 60 animals per strain. *P*-values

represent the significance level of mutant survival relative to WT: *nmur-1(ok1387)*, *P* < 0.0001;

capa-1(ok3065), *P* = 0.2323; *capa-1(ok3065);nmur-1(ok1387)*, *P* < 0.0001. **(D)** WT, *nmur-*

1(ok1387), *capa-1(ok3065)*, and *capa-1(ok3065);nmur-1(ok1387)* animals were exposed to *S.*

enterica on plates contain either 0 or 1µg/mL of synthetic ligand and scored for survival over

time. The graph is a combination of three independent experiments. Each experiment included

N = 60 animals per strain. *P*-values represent the significance level of the survival of

mutant+1µg/mL relative to WT+1µg/mL: *nmur-1(ok1387)*+1µg/mL, *P* < 0.0001; *capa-*

1(ok3065)+1µg/mL, *P* = 0.1643; *capa-1(ok3065);nmur-1(ok1387)*+1µg/mL, *P* < 0.0001.

Table 1. Gene ontology (GO) analysis of downregulated genes in *nmur-1(ok1387)* animals

relative to wild-type animals.

Table 2. GO analysis of downregulated genes in *nmur-1(ok1387)* animals relative to wild-

type animals exposed to *E. faecalis*.

Table 3. Enrichment of biological process and molecular function revealed by GO

analysis of upregulated genes in *nmur-1(ok1387)* animals relative to wild-type animals

exposed to *S. enterica*.

SUPPLEMENTARY MATERIALS

Fig. S1. Functional loss of NMUR-1 differentially affects *C. elegans* survival against

various bacteria. WT and *nmur-1(ok1387)* animals were exposed to *E. coli* OP50 **(A)**, *P.*

aeruginosa **(B)**, *Y. pestis* **(C)**, *M. nematophilum* **(D)**, or *S. aureus* **(E)** and scored for survival

over time. Each graph is a representation of three independent experiments. Each experiment

included *N* = 60 animals per strain. *P*-values represent the significance level of mutant survival

relative to WT, *P* = 0.0011 in (A), *P* = 0.8084 in (B), *P* < 0.0001 in (C), *P* = 0.2401 in (D), and *P*

= 0.0678 in (E).

Fig. S2. Functional loss of NMUR-1 does not affect *C. elegans* survival against heat-killed

***E. faecalis* or heat-killed *S. enterica*.** WT and *nmur-1(ok1387)* animals were exposed to heat-

killed *E. faecalis* **(A)** or heat-killed *S. enterica* **(B)** and scored for survival over time. Each graph

is a representation of three independent experiments. Each experiment included *N* = 60 animals

per strain. *P*-value represents the significance level of the survival of mutants relative to WT, *P*

= 0.2191.

Fig. S3. Functional loss of NMUR-1 differentially affects *C. elegans* survival against *E.*

***faecalis* and *S. enterica* in full-lawn assays.** WT and *nmur-1(ok1387)* animals were exposed

to a full lawn of *E. faecalis* **(A)** or *S. enterica* **(B)** and scored for survival over time. Each graph is

a combination of three independent experiments. Each experiment included *N* = 60 animals per

strain. *P*-values represent the significance level of the survival of mutants relative to WT, *P* <

0.0001 in (A); *P* < 0.0001 in (B).

Table S1. Enrichment of molecular functions revealed by GO analysis of upregulated

genes in *nmur-1(ok1387)* animals relative to wild-type animals.

Table S2. Enrichment of biological processes revealed by GO analysis of upregulated

genes in *nmur-1(ok1387)* animals relative to wild-type animals exposed to *E. faecalis*.

Table S3. List of transgenic *C. elegans* strains generated in this study.

 FIGURES AND TABLES

946
947
948
949   **Fig. 1**

950
951

952
953
Fig. 1. Functional loss of NMUR-1 differentially affects *C. elegans* survival against *E. coli*
HT115, *E. faecalis* and *S. enterica*. Wild-type (WT) and *nmur-1(ok1387)* animals were
exposed to *E. coli* HT115 **(A)**, *E. faecalis* **(B)**, or *S. enterica* **(C)** and scored for survival over
time. Each graph is a combination of three independent experiments. Each experiment included
N = 60 animals per strain. *P*-values represent the significance level of mutant survival relative to
WT, *P* = 0.8706 in (A), *P* = 0.0004 in (B), and *P* < 0.0001 in (C).

960
961

962   **Fig. 2**
963
964

965
966
Fig. 2. Functional loss of NMUR-1 does not affect pathogen avoidance behavior. WT and
nmur-1(ok1387) animals were subjected to lawn occupancy assays in which the animals were
placed on a small lawn of *E. faecalis* **(A)** or *S. enterica* **(B)** in a 3.5cm plate and monitored over
time for their presence on the lawn. Each graph is a combination of three independent
experiments. Each experiment included *N* = 60 animals per strain. Error bars represent the
standard error of the mean (SEM).
973
974
975

Fig. 3

A

WT · *nmur-1(ok1387)*

E

WT · *nmur-1(ok1387)*

B

F

C

G

D

H

Fig 3. Functional loss of NMUR-1 changes intestinal accumulation of *E. faecalis* and *S. enterica*. (A) WT and *nmur-1(ok1387)* animals were exposed to GFP-expressing *E. faecalis* for

24 hours and then visualized using a Zeiss Axio Imager M2 fluorescence microscope. The scale
bar indicates 500 μm. **(B)** WT and *nmur-1(ok1387)* animals were exposed to *E. faecalis* for 24
hours, and the Colony Forming Units (CFU) of live bacterial cells recovered from their intestines
were quantified. The graph represents the combined results of three independent experiments.
N = 10 animals per strain were used for each experiment. Error bars represent SEM. *P*-value
represents the significance level of mutant gut colonization relative to WT, P = 0.0296. **(C)** WT
and *nmur-1(ok1387)* animals were exposed to *E. faecalis* for 24 hours. Pharyngeal pumping
rates of animals were counted as the number of pumps per 30 sec. Counting was conducted in
triplicate and averaged to give a pumping rate. The measurements included N = 10 adult
animals per strain. Error bars represent SEM. P = 0.1068. **(D)** WT and *nmur-1(ok1387)* animals
were exposed to *E. faecalis* for 24 hours. Defecation rates of animals were measured as the
average time of 10 intervals between two defecation cycles. The measurements included N =
10 animals per strain. P = 0.1379. **(E)** WT and *nmur-1(ok1387)* animals were exposed to GFP-
expressing *S. enterica* for 24 hours and then visualized using a Zeiss Axio Imager M2
fluorescence microscope. The scale bar indicates 500 μm. **(F)** WT and *nmur-1(ok1387)* animals
were exposed to GFP-expressing *S. enterica* for 24 hours, and the CFU of live bacterial cells
recovered from their intestines were quantified. The graph represents the combined results of
three independent experiments. N = 10 animals per strain were used for each experiment. Error
bars represent SEM. P = 0.0290. **(G)** WT and *nmur-1(ok1387)* animals were exposed to *S.
enterica* for 24 hours. Pharyngeal pumping rates of animals were counted as the number of
pumps per 30 sec. Counting was conducted in triplicate and averaged to give a pumping rate.
The measurements included N = 10 adult animals per strain. Error bars represent SEM. P =
0.0006. **(H)** WT and *nmur-1(ok1387)* animals were exposed to *S. enterica* for 24 hours.
Defecation rates of animals were measured as the average time of 10 intervals between two
defecation cycles. The measurements included N = 10 animals per strain. P = 0.6310.
1009
1010

1011 **Fig. 4**

1012

1013

1014

1015

Fig. 4. NMUR-1 regulates transcriptional activity. (A) *In vitro* transcription assays were

performed using nuclear extract from WT or *nmur-1(ok1387)* animals. After transcription, RNA

was purified, reverse transcribed, and amplified by PCR, followed by gel analysis. Lane 1,

transcription with WT nuclear extract; lane 2, transcription with *nmur-1(ok1387)* nuclear extract;

lane 3, transcriptional positive control (with HeLa nuclear extract); lane 4, transcriptional

negative control (without any nuclear extract); lane 5, PCR positive control; lane 6, PCR

negative control (without DNA template). **(B)** WT and mutant nuclear extracts were titrated using

the *in vitro* transcription system. The RNA copy number generated from transcription was

plotted against the amount of nuclear extract used in the reaction, followed by fitting with the

Michaelis-Menten model and the non-linear least-squares method to generate the titration

curves and calculate titration parameters. **(C)** qRT-PCR analysis of GFP expression after two-

hour heat shock in *CL2070(dvIs70)* and *nmur-1(ok1387);CL2070(dvIs70)* animals. Asterisk (*)

denotes a significant difference between *CL2070(dvIs70)* and *nmur-1(ok1387);CL2070(dvIs70)*

animals

1030

1031

1032

1033 **Fig. 5**
1034
1035

1036
1037
1038
Fig. 5. NMUR-1-regulated C-type lectin genes are required for *C. elegans* defense against
***E. faecalis*. (A)** WT and **(B)** *nmur-1(ok1387)* animals grown on dsRNA for C-type lectin genes
or the empty vector (EV) control were exposed to *E. faecalis* and scored for survival over time.
The graphs are a combination of three independent replicates. Each experiment included *N* =
60 animals per strain. *P*-values in (A) represent the significance level of RNAi treatment relative
to WT+EV: WT+*clec-94*, *P* < 0.0001; WT+*clec-137*, *P* = 0.0220; WT+*clec-157*, *P* = 0.3728;
WT+*clec-208*, *P* < 0.0001; WT+*clec-263*, *P* < 0.0001. *P*-values in (B) represent the significance
level of the RNAi treatment relative to *nmur-1(ok1387)*+EV: *nmur-1(ok1387)*+*clec-94*, *P* =
0.3633; *nmur-1(ok1387)*+*clec-137*, *P* = 0.1841; *nmur-1(ok1387)*+*clec-157*, *P* = 0.8617; *nmur-*
1(ok1387)+*clec-208*, *P* = 0.1348; *nmur-1(ok1387)*+*clec-263*, *P* = 0.1320. **(C)** WT and *nmur-*
1(ok1387) animals in which C-type lectin expression was rescued were exposed to *E. faecalis*
and scored for survival over time. All C-type lectin genes were rescued using their own
promoters. The graph is a combination of three independent experiments. Each experiment
included *N* = 60 adult animals per strain. *P*-values represent the significance of the survival of
rescue strains relative to *nmur-1(ok1378)* animals: WT, *P* < 0.0001; nmur-1(ok1387)+*clec-94*, *P*
< 0.0001; *nmur-1(ok1387)*+*clec-208*, *P* = 0.0025; *nmur-1(ok1387)*+*clec-263*, *P* = 0.9799.
1055
1056
1057

1058 **Fig. 6**
1059
1060

1061
1062
Fig. 6. NMUR-1-regulated UPR genes are required for *C. elegans* defense against *S.*
***enterica*. (A)** *TU3401* and **(B)** *nmur-1(ok1387);TU3401* animals grown on dsRNA for UPR
genes or the empty vector (EV) control were exposed to *S. enterica* and scored for survival over
time. The graphs are a combination of three independent replicates. Each experiment included
N = 60 animals per strain. *P*-values in (A) represent the significance level of RNAi treatment
relative to *TU3401*+EV: *TU3401+T20D4.3*, *P* = 0.4224; *TU3401+T20D4.5*, *P* = 0.6102;
TU3401+C17B7.5, *P* = 0.5905; *TU3401+npl-4.2*, *P* < 0.0001. *P*-values in (B) represent the
significance level of RNAi treatment relative to *nmur-1(ok1387);TU3401*+EV: *nmur-*
1(ok1387);TU3401+T20D4.3, *P* = 0.0007; *nmur-1(ok1387);TU3401+T20D4.5*, *P* = 0.0993;
nmur-1(ok1387);TU3401+C17B7.5, *P* = 0.1216; *nmur-1(ok1387);TU3401+npl-4.2*; *P* < 0.0001.
1073
1074

Fig. 7

Fig. 7. CAPA-1 is required for *C. elegans* defense against *E. faecalis* but not *S. enterica*.
(A) WT, *nmur-1(ok1387)*, *capa-1(ok3065)*, and *capa-1(ok3650);nmur-1(ok1387)* animals were
exposed to *E. faecalis* and scored for survival over time. The graph is representative of three
independent experiments. Each experiment included *N* = 60 animals per strain. *P*-values
represent the significance level of mutant survival relative to WT: *nmur-1(ok1387)*, *P* = 0.0005;
capa-1(ok3065), *P* = 0.0134; *capa-1(ok3065);nmur-1(ok1387)*, *P* = 0.0028. **(B)** WT, *nmur-
1(ok1387)*, *capa-1(ok3065)*, and *capa-1(ok3650);nmur-1(ok1387)* animals were exposed to *E.
faecalis* on plates containing either 0 or 1μg/mL synthetic ligand and scored for survival over
time. The graph is a combination of three independent experiments. Each experiment included
N = 60 animals per strain. *P*-values represent the significance level of the survival of
mutant+1μg/mL relative to WT+1μg/mL: *nmur-1(ok1387)*+1μg/mL, *P* < 0.0001; *capa-
1(ok3065)*+1μg/mL, *P* = 0.6758; *capa-1(ok3065);nmur-1(ok1387)*+1μg/mL, *P* = 0.0024. **(C)** WT,
nmur-1(ok1387), *capa-1(ok3065)*, and *capa-1(ok3065);nmur-1(ok1387)* animals were exposed
to *S. enterica* and scored for survival over time. The graph is representative of three
independent experiments. Each experiment included *N* = 60 animals per strain. *P*-values
represent the significance level of mutant survival relative to WT: *nmur-1(ok1387)*, *P* < 0.0001;
capa-1(ok3065), *P* = 0.2323; *capa-1(ok3065);nmur-1(ok1387)*, *P* < 0.0001. **(D)** WT, *nmur-
1(ok1387)*, *capa-1(ok3065)*, and *capa-1(ok3065);nmur-1(ok1387)* animals were exposed to *S.
enterica* on plates contain either 0 or 1μg/mL of synthetic ligand and scored for survival over
time. The graph is a combination of three independent experiments. Each experiment included
N = 60 animals per strain. *P*-values represent the significance level of the survival of
mutant+1μg/mL relative to WT+1μg/mL: *nmur-1(ok1387)*+1μg/mL, *P* < 0.0001; *capa-
1(ok3065)*+1μg/mL, *P* = 0.1643; *capa-1(ok3065);nmur-1(ok1387)*+1μg/mL, *P* < 0.0001.

Table 1. Gene ontology (GO) analysis of downregulated genes in *nmur-1(ok1387)* animals relative to wild-type animals.

(A) Enriched molecular functions

GO term	Description	P-value#	FDR q-value*	Enrichment (N, B, n, b)§
GO:0003690	double-stranded DNA binding	2.37E-08	5.83E-05	5.59 (9751,169,165,16)
GO:0000976	transcription regulatory region sequence-specific DNA binding	3.29E-08	4.05E-05	6.41 (9751,129,165,14)
GO:0000981	DNA-binding transcription factor activity, RNA polymerase II-specific	3.66E-08	3.01E-05	7.79 (9751,91,165,12)
GO:1990837	sequence-specific double-stranded DNA binding	5.88E-08	3.62E-05	6.13 (9751,135,165,14)
GO:0001067	regulatory region nucleic acid binding	3.08E-07	1.52E-04	5.37 (9751,154,165,14)
GO:0044212	transcription regulatory region DNA binding	3.08E-07	1.26E-04	5.37 (9751,154,165,14)
GO:0003700	DNA-binding transcription factor activity	1.10E-06	3.86E-04	3.35 (9751,371,165,21)
GO:0140110	transcription regulator activity	1.97E-06	6.07E-04	3.01 (9751,451,165,23)
GO:0000977	RNA polymerase II regulatory region sequence-specific DNA binding	4.31E-06	1.18E-03	5.56 (9751,117,165,11)
GO:0001012	RNA polymerase II regulatory region DNA binding	5.51E-06	1.36E-03	5.42 (9751,120,165,11)
GO:0043565	sequence-specific DNA binding	7.12E-05	1.59E-02	2.81 (9751,379,165,18)

(B) Downregulated genes related to enriched molecular functions

Genes	Fold change	Adjusted *P* valueΨ	Genes	Fold change	Adjusted *P* valueΨ
dmd-5	3.7	3.09E-05	nhr-25	2.6	4.47E-13
unc-39	3.6	5.68E-09	ceh-43	2.5	1.35E-13
ceh-27	3.6	2.49E-11	tbx-8	2.4	1.64E-04
egl-46	3.6	9.25E-18	unc-130	2.3	1.84E-11
ces-2	3.4	1.14E-19	efl-3	2.3	2.26E-05
cnd-1	3.4	1.32E-05	sepa-1	2.1	4.00E-03
fkh-2	3.3	6.61E-07	ham-1	2.1	7.64E-13
zip-8	3.2	1.13E-06	ham-2	2.0	6.06E-08
zip-7	2.8	5.01E-11	tbx-11	2.0	6.53E-05
ref-2	2.8	1.97E-06	hbl-1	2.0	6.38E-07
eya-1	2.7	3.20E-13	ceh-32	2.0	1.20E-04
hlh-14	2.6	1.66E-07			

P-value is computed according to the mHG model (Eden *et al.* 2007 PLoS Comp Bio 3(3):e39).
* FDR q-value is the correction of the above p-value for multiple testing using the Benjamini and Hochberg method (Benjamini and Hochberg 1995 J R Statist Soc B 57(1):289-300).
§ Enrichment (N, B, n, b) is defined as follows: N - total number of genes; B - total number of genes associated with a specific GO term; n - number of genes in the target set; b - number of genes in the intersection;Enrichment = (b/n) / (B/N).
Ψ Adjusted *P* value is the correction of the P value for multiple testing using the Benjamini and Hochberg method (Benjamini and Hochberg 1995 J R Statist Soc B 57 (1):289–300).

Table 2. GO analysis of downregulated genes in *nmur-1(ok1387)* animals relative to wild-type animals exposed to *E. faecalis*.

(A) Enriched molecular functions

GO term	Description	P-value[#]	FDR q-value[*]	Enrichment (N, B, n, b)[§]
GO:0030246	carbohydrate binding	2.29E-15	5.65E-12	17.45 (10161,182,48,15)

(B) Downregulated genes related to the enriched molecular functions

Genes	Fold change	Adjusted *P* value[Ψ]	Genes	Fold change	Adjusted *P* value[Ψ]
clec-207	5.4	4.16E-15	clec-161	2.4	1.10E-04
clec-94	4.9	1.50E-12	clec-95	2.4	2.91E-04
clec-137	4.3	3.68E-10	clec-99	2.4	6.31E-05
clec-263	3.7	1.68E-08	clec-197	2.4	7.91E-04
clec-208	3.4	2.01E-07	clec-136	1.9	2.03E-02
clec-157	2.9	3.25E-06	clec-174	1.9	2.50E-03
pqn-84	2.8	2.76E-06	clec-190	1.8	1.29E-04
clec-219	2.5	2.92E-04			

[#] P-value is computed according to the mHG model (Eden *et al.* 2007 PLoS Comp Bio 3(3):e39).
[*] FDR q-value is the correction of the above p-value for multiple testing using the Benjamini and Hochberg method (Benjamini and Hochberg 1995 J R Statist Soc B 57(1):289-300).
[§] Enrichment (N, B, n, b) is defined as follows: N - total number of genes; B - total number of genes associated with a specific GO term; n - number of genes in the target set; b - number of genes in the intersection;Enrichment = (b/n) / (B/N).
[Ψ]Adjusted *P* value is the correction of the P value for multiple testing using the Benjamini and Hochberg method (Benjamini and Hochberg 1995 J R Statist Soc B 57 (1):289–300).

Table 3. Enrichment of biological process and molecular function revealed by GO analysis of upregulated genes in *nmur-1(ok1387)* animals relative to wild-type animals exposed to *S. enterica*.

(A) Enriched biological processes and molecular functions

GO term	Description	P-value[#]	FDR q-value[*]	Enrichment (N, B, n, b)[§]
Biological process				
GO:0006515	protein quality control for misfolded or incompletely synthesized proteins	1.35E-8	7.89E-5	137.71 (9812,19,15,4)
GO:0006516	glycoprotein catabolic process	3.45E-7	1.01E-3	196.24 (9812,10,15,3)
GO:0006517	protein deglycosylation	1.04E-6	2.03E-3	140.17 (9812,14,15,3)
GO:0009100	glycoprotein metabolic process	2E-5	2.92E-2	54.51 (9812,36,15,3)
Molecular function				
GO:0000224	peptide-N4-(N-acetyl-beta-glucosaminyl)asparagine amidase activity	2.41E-7	5.96E-4	218.04 (9812,9,15,3)
GO:0016811	hydrolase activity, acting on carbon-nitrogen (but not peptide) bonds, in linear amides	2.98E-5	3.67E-2	47.86 (9812,41,15,3)

(B) Upregulated genes related to the enriched biological process protein quality control

Genes	Fold change	Adjusted *P* value[Ψ]
C17B7.5	2.9	2.57E-14
T20D4.5	2.8	2.40E-15
npl-4.2	2.8	1.22E-22
T20D4.3	2.6	1.60E-12

[#] P-value is computed according to the mHG model (Eden *et al.* 2007 PLoS Comp Bio 3(3):e39).
[*] FDR q-value is the correction of the above p-value for multiple testing using the Benjamini and Hochberg method (Benjamini and Hochberg 1995 J R Statist Soc B 57(1):289-300).
[§] Enrichment (N, B, n, b) is defined as follows: N - total number of genes; B - total number of genes associated with a specific GO term; n - number of genes in the target set; b - number of genes in the intersection;Enrichment = (b/n) / (B/N).
[Ψ]Adjusted *P* value is the correction of the P value for multiple testing using the Benjamini and Hochberg method (Benjamini and Hochberg 1995 J R Statist Soc B 57 (1):289–300).

1152 **SUPPLEMENTARY MATERIALS**
1153
1154 **Fig. S1**
1155

1156
1157
Fig. S1. Functional loss of NMUR-1 differentially affects *C. elegans* survival against
various bacteria. WT and *nmur-1(ok1387)* animals were exposed to *E. coli* OP50 **(A)**, *P.*
aeruginosa **(B)**, *Y. pestis* **(C)**, *M. nematophilum* **(D)**, or *S. aureus* **(E)** and scored for survival
over time. Each graph is a representation of three independent experiments. Each experiment
included $N = 60$ animals per strain. *P*-values represent the significance level of mutant survival
relative to WT, $P = 0.0011$ in (A), $P = 0.8084$ in (B), $P < 0.0001$ in (C), $P = 0.2401$ in (D), and *P*
$= 0.0678$ in (E).
1165
1166

1167 **Fig. S2**
1168

1169
1170
Fig. S2. Functional loss of NMUR-1 does not affect *C. elegans* survival against heat-killed
***E. faecalis* or heat-killed *S. enterica*.** WT and *nmur-1(ok1387)* animals were exposed to heat-
killed *E. faecalis* **(A)** or heat-killed *S. enterica* **(B)** and scored for survival over time. Each graph
is a representation of three independent experiments. Each experiment included *N* = 60 animals
per strain. *P*-value represents the significance level of the survival of mutants relative to WT, *P*
= 0.2191.
1177
1178

 Fig. S3

1180

1181

1182
1183
Fig. S3. Functional loss of NMUR-1 differentially affects *C. elegans* survival against *E.*
***faecalis* and *S. enterica* in full-lawn assays.** WT and *nmur-1(ok1387)* animals were exposed
to a full lawn of *E. faecalis* **(A)** or *S. enterica* **(B)** and scored for survival over time. Each graph is
a combination of three independent experiments. Each experiment included N = 60 animals per
strain. *P*-values represent the significance level of the survival of mutants relative to WT, $P <$
0.0001 in (A); $P < 0.0001$ in (B).
1190
1191
1192

Table S1. Enrichment of molecular functions revealed by GO analysis of upregulated genes in *nmur-1(ok1387)* animals relative to wild-type animals.

GO term	Description	P-value[#]	FDR q-value*	Enrichment (N, B, n, b)[§]
GO:0004721	phosphoprotein phosphatase activity	1.21E-40	3.08E-37	5.37 (9948,161,921,80)
GO:0004725	protein tyrosine phosphatase activity	6.28E-37	7.97E-34	6.95 (9948,87,921,56)
GO:0042302	structural constituent of cuticle	1.05E-34	8.86E-32	5.62 (9948,125,921,65)
GO:0140096	catalytic activity, acting on a protein	1.97E-33	1.25E-30	2.17 (9948,1133,921,228)
GO:0004672	protein kinase activity	2.29E-30	1.16E-27	3.19 (9948,376,921,111)
GO:0016791	phosphatase activity	4.24E-30	1.79E-27	4.08 (9948,212,921,80)
GO:0042578	phosphoric ester hydrolase activity	3.37E-25	1.22E-22	3.53 (9948,245,921,80)
GO:0016773	phosphotransferase activity, alcohol group as acceptor	9.08E-25	2.88E-22	2.78 (9948,431,921,111)
GO:0004715	non-membrane spanning protein tyrosine kinase activity	2.07E-21	5.83E-19	7.36 (9948,44,921,30)
GO:0004674	protein serine/threonine kinase activity	2.35E-21	5.96E-19	3.12 (9948,277,921,80)
GO:0016301	kinase activity	2.42E-21	5.59E-19	2.52 (9948,485,921,113)
GO:0016772	transferase activity, transferring phosphorus-containing groups	1.34E-15	2.84E-13	2.13 (9948,579,921,114)
GO:0016788	hydrolase activity, acting on ester bonds	1.19E-14	2.33E-12	2.22 (9948,471,921,97)
GO:0004713	protein tyrosine kinase activity	9.57E-13	1.74E-10	4.19 (9948,80,921,31)
GO:0004722	protein serine/threonine phosphatase activity	2.86E-11	4.84E-09	4.63 (9948,56,921,24)
GO:0005198	structural molecule activity	4.50E-09	7.14E-07	2.08 (9948,342,921,66)
GO:0005524	ATP binding	5.16E-08	7.71E-06	1.61 (9948,798,921,119)
GO:0032559	adenyl ribonucleotide binding	1.31E-07	1.85E-05	1.58 (9948,820,921,120)
GO:0030554	adenyl nucleotide binding	1.60E-07	2.14E-05	1.57 (9948,823,921,120)
GO:0008144	drug binding	1.52E-06	1.93E-04	1.49 (9948,912,921,126)
GO:0035639	purine ribonucleoside triphosphate binding	7.87E-05	9.51E-03	1.38 (9948,960,921,123)
GO:0032555	purine ribonucleotide binding	1.46E-04	1.68E-02	1.36 (9948,983,921,124)
GO:0017076	purine nucleotide binding	1.74E-04	1.92E-02	1.36 (9948,987,921,124)
GO:0005102	signaling receptor binding	1.89E-04	1.99E-02	1.92 (9948,186,921,33)
GO:0032553	ribonucleotide binding	2.89E-04	2.94E-02	1.34 (9948,999,921,124)
GO:0042329	structural constituent of collagen and cuticulin-based cuticle	3.51E-04	3.43E-02	5.40 (9948,12,921,6)
GO:0030246	carbohydrate binding	4.64E-04	4.37E-02	1.94 (9948,156,921,28)

[#] P-value is computed according to the mHG model (Eden *et al.* 2007 PLoS Comp Bio 3(3):e39).
* FDR q-value is the correction of the above p-value for multiple testing using the Benjamini and Hochberg method (Benjamini and Hochberg 1995 J R Statist Soc B 57(1):289-300).
[§] Enrichment (N, B, n, b) is defined as follows:
  N - total number of genes
  B - total number of genes associated with a specific GO term
  n - number of genes in the target set
  b - number of genes in the intersection
  Enrichment = (b/n) / (B/N)

1207 **Table S2. Enrichment of biological processes revealed by GO analysis of upregulated**
1208 **genes in *nmur-1(ok1387)* animals relative to wild-type animals exposed to *E. faecalis*.**
1209

GO term	Description	P-value#	FDR q-value*	Enrichment (N, B, n, b)§
GO:0050830	defense response to Gram-positive bacterium	1.24E-11	7.49E-08	15.32 (10357,61,133,12)
GO:0042742	defense response to bacterium	3.97E-09	1.20E-05	7.57 (10357,144,133,14)
GO:0009617	response to bacterium	3.97E-09	7.98E-06	7.57 (10357,144,133,14)
GO:0098542	defense response to other organism	2.70E-08	4.07E-05	6.53 (10357,167,133,14)
GO:0043207	response to external biotic stimulus	2.91E-08	3.52E-05	6.49 (10357,168,133,14)
GO:0009607	response to biotic stimulus	2.91E-08	2.93E-05	6.49 (10357,168,133,14)
GO:0051707	response to other organism	2.91E-08	2.51E-05	6.49 (10357,168,133,14)
GO:0051704	multi-organism process	2.02E-07	1.52E-04	5.56 (10357,196,133,14)
GO:0009605	response to external stimulus	9.53E-06	6.39E-03	4.02 (10357,271,133,14)
GO:0042338	cuticle development involved in collagen and cuticulin-based cuticle molting cycle	1.13E-05	6.83E-03	16.22 (10357,24,133,5)
GO:0006952	defense response	3.61E-05	1.98E-02	3.38 (10357,346,133,15)

1210
P-value is computed according to the mHG model (Eden *et al.* 2007 PLoS Comp Bio 3(3):e39).
* FDR q-value is the correction of the above p-value for multiple testing using the Benjamini and
Hochberg method (Benjamini and Hochberg 1995 J R Statist Soc B 57(1):289-300).
§ Enrichment (N, B, n, b) is defined as follows:
N - total number of genes
B - total number of genes associated with a specific GO term
n - number of genes in the target set
b - number of genes in the intersection
Enrichment = (b/n) / (B/N)
1220
1221

1222 **Table S3. List of transgenic *C. elegans* strains generated in this study.**
1223

Strain	Plasmid and genetic background	Host	Note
JRS60	pPW01 *clec-94p(4.2kb)::clec-94 gDNA::SL2::gfp*	*N2*	*clec-94* gene over-expression in *N2* animals
JRS61	pPW02 *clec-208p(3.6kb)::clec-208 gDNA::SL2::gfp*	*N2*	*clec-208* gene over-expression in *N2* animals
JRS62	pPW03 *clec-263p(5.4kb)::clec-263 gDNA::SL2::gfp*	*N2*	*clec-263* gene over-expression in *N2* animals
JRS63	pPW01 *clec-94p(4.2kb)::clec-94 gDNA::SL2::gfp*	*nmur-1(ok1387)*	*clec-94* genomic rescue driven by *clec-94* promoter
JRS64	pPW02 *clec-208p(3.6kb)::clec-208 gDNA::SL2::gfp*	*nmur-1(ok1387)*	*clec-208* genomic rescue driven by *clec-208* promoter
JRS65	pPW03 *clec-263p(5.4kb)::clec-263 gDNA::SL2::gfp*	*nmur-1(ok1387)*	*clec-263* genomic rescue driven by *clec-263* promoter

1224
1225
1226
1227