Spatial control of cell division by GA-OsGRF7/8 module in a leaf explains the leaf length variation between cultivated and wild rice

Vikram Jathar[1], Kumud Saini[1], Ashish Chauhan[1], Ruchi Rani[1], Yasunori Ichihashi[2], Aashish Ranjan[1*]

[1]National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi-110067, India.

[2]RIKEN BioResource Research Center, 3-1-1 Koyadai, Tsukuba, Ibaraki, 305-0074, Japan.

*Address for correspondence

Aashish Ranjan

Tel: 91-11-26735117

Email: aranjan@nipgr.ac.in

Total word count for the main body of the text: 6,448

Introduction: 830

Materials and Methods: 1,618

Results: 2,684

Discussion: 1,316

8 figures (4 color figures)

2 tables

Supporting information: 10 figures, 6 tables, 7 datasets

Summary

- Cellular and genetic understanding of rice leaf size regulation is limited, despite rice being the staple food of more than half of the global population. We investigated the mechanism controlling the rice leaf length using cultivated and wild rice accessions that remarkably differed for leaf size.

- Comparative transcriptomics, Gibberellic Acid (GA) quantification, and leaf kinematics of the contrasting accessions suggested the involvement of GA, cell cycle, and Growth-Regulating Factors (GRFs) in the rice leaf size regulation. Zone-specific expression analysis and VIGS established the functions of specific GRFs in the process.

- The leaf length of the selected accessions was strongly correlated with GA levels. Higher GA content in wild rice accessions with longer leaves and GA-induced increase in the leaf length via an increase in cell division confirmed a GA-mediated regulation of division zone in rice. Downstream to GA, *OsGRF7* and *OsGRF8* function for controlling cell division to determine the rice leaf length.

- Spatial control of cell division to determine the division zone size mediated by GA and downstream OsGRF7 and OsGRF8 explains the leaf length differences between the cultivated and wild rice. This mechanism to control rice leaf length might have contributed to optimizing leaf size during domestication.

Keywords: Cell division, Cultivated and wild rice, Gibberellic acid (GA), Growth-Regulating Factors (GRFs), Leaf Kinematics, Leaf length, Leaf growth zones.

Introduction

The basic leaf development program includes leaf primordia initiation on the flanks of

shoot apical meristem that follows growth and patterning by multiple transcription factor

families (Townsley & Sinha, 2012; Bar & Ori, 2014; Du *et al*., 2018). While leaf initiation

and patterning are regulated by a suite of transcription factors and complex hormonal

interactions, the final leaf size is determined by the coordination of cell division and

expansion (Li *et al*., 2007; Gonzalez *et al*., 2012; Chitwood & Sinha 2016; Sarvepalli *et al*.,

2019). Considering the leaf as the major plant organ for light-harvesting and perception of

environmental signals, understanding the genetic basis of leaf size regulation is imperative

for crop improvement.

The regulation of leaf size is extensively studied for simple leaves of Arabidopsis and

compound leaves of tomato, however detailed mechanistic studies are relatively scarce for

monocot cereals except for maize (Efroni *et al*., 2010; Rodriguez *et al*., 2013; Vercruysse *et*

al., 2020a). The spatial and temporal regulation of cell proliferation and cell elongation to

determine the final leaf size varies between monocots and dicots. Cell proliferation and

expansion are temporally separated in dicots as cell proliferation predominates in young

developing leaf, thereafter cells expand before reaching maturity (Beemster *et al*., 2005;

Piazza *et al*., 2005; Andriankaja *et al*., 2012). In contrast, cell proliferation and expansion are

spatially separated in monocots, such as maize, where division, elongation, and maturation

zones are present together at steady-state leaf growth (Tardieu & Garnier, 2000; Granier &

Tardieu, 2009; Fina *et al*., 2017). Monocot leaves with such spatial resolution of the growth

zones allow the systematic study of the zones and their contribution in the leaf size control.

Changes in the size of the division and/or elongation zone alter the final leaf size in

Arabidopsis and maize (Tardieu & Granier, 2000; Tardieu *et al*., 2000; Gonzalez *et al*., 2010;

Kalve *et al*., 2014; Gazquez & Beemster, 2017; Nelissen *et al*., 2018). Several

phytohormones play important roles in the regulation of cell division, elongation, and

transition between division and elongation (Inze & De Veylder, 2006; Su *et al.*. 2011;

Avramova *et al*., 2015; Takatsuka & Umeda, 2014). GA is classically known for mediating

cell-elongation response, however recent studies in Arabidopsis and maize have shown its

key roles in determining cell division (Achard *et al*., 2009; Nelissen *et al*., 2012). GA, after

being perceived by a receptor protein GIBBERELLIN INSENSITIVE DWARF1 (GID1),

promotes degradation of DELLA repressor proteins as well as suppresses expression of

DELLA-encoding *SLR1* in feedback regulation (Willige *et al*., 2007; Davière & Achard,

2013; Shan *et al*., 2014). GA-mediated suppression of DELLA proteins results in activation of GA-responsive transcription factors, such as GROWTH-REGULATING FACTORs (GRFs) (Omidbakhshfard *et al*., 2015). GRFs are reported to directly regulate cyclins to control cell division in different plant species (van der Knaap *et al*., 2000; Choi *et al*., 2004; Nelissen *et al.*, 2015; Fina *et al*., 2017). Arabidopsis plants overexpressing *AtGRF1*, *AtGRF2*, and *AtGRF3* develop longer leaves and larger organ sizes, whereas *grf* mutants produce smaller and narrower leaves (Kim *et al*., 2003; Beltramino *et al*., 2018). Moreover, *GRF* expression modulation in maize alters leaf length with associated changes in division zone size (Wu *et al*., 2014; Nelissen *et al*., 2015)

A detailed cellular basis and underlying genetic and hormonal regulation of rice leaf length control are not systematically investigated. Rice domestication has significantly reduced the genetic variability in the cultivated varieties (Huang *et al*., 2012; Chen *et al*., 2021). Therefore, a comparative study of leaf size and associated cellular differences across cultivated and wild rice would provide a comprehensive genetic understanding of the rice leaf size regulation. Several wild relatives of rice grow taller and accumulate higher biomass associated with larger organ size including leaf size (Sanchez *et al*., 2013; Mathan *et al*., 2021a). GA is known to regulate stem elongation in rice, and manipulation of GA-biosynthesis and signaling genes have optimized plant height during the green revolution (van der Knaap *et al*., 2000). Several GRFs downstream to GA are known to regulate various aspects of rice development and architecture (Luo *et al*., 2005; Kuijt *et al*., 2014; Gao *et al*., 2015; Tang *et al*., 2018; Chen *et al*., 2020). OsGRF1, the founding member of the rice GRF transcription factor family, is expressed in intercalary meristems of internodes and influences rice leaf growth (van der Knaap *et al*., 2000; Lu *et al*., 2020). However, the contribution of cell division and its regulation by GA and specific downstream GRFs for rice leaf length regulation is not systematically elucidated. Further, the signaling module that explains the leaf length differences between the contrasting cultivated and wild rice is also not investigated. Our study has systematically integrated GA with the rice leaf length control via cell cycle regulation and identified specific GRFs, OsGRF7 and OsGRF8, downstream to GA that not only explains the leaf length differences between the selected cultivated and wild rice but also would provide a way to optimize rice leaf length for improved physiological performance.

Materials and Methods

Plant materials and growth conditions

The genotypes used for this study are summarized in Table S1. Rice seeds were first
germinated on Petri plates containing wet germination paper, followed by transfer of
seedlings to a hydroponic solution containing one-fourth MS Media (Himedia) in standard
concentrations. The nutrient solution was changed every third day during the seedling
growth. Seedlings were grown at 28 ℃ day and 23 ℃ night temperature with 14:10 hours of
light and dark cycle. For exogenous treatments, 10µM of gibberellic acid (GA$_3$), and 1µM of
gibberellic acid biosynthesis inhibitor Paclobutrazol (PAC) were added in hydroponic
solution.

Phenotyping of leaf morphological traits and leaf growth analysis
The quantification of leaf length and width was performed on each fully-grown leaf from the
second to eight leaves of all the selected rice accessions. Leaf length and width were
quantified manually using a ruler. Leaf-sheath length was measured as a distance between the
seedling base to ligule. The leaf-blade length was quantified as the distance between the
ligule to the leaf tip. Leaf-blade width was measured in the central widest part of the fully
expanded leaf blade. Total mature leaf length was calculated by adding leaf sheath and leaf
blade length. The leaf growth analysis was done on the fourth leaf, when the leaf emerged
from the sheath of the third leaf, as explained by Sprangers *et al.* (2016). Leaf length was
measured daily from the day of emergence up to the seventh day by considering the seedling
base as the starting point of the leaf. Leaf elongation rate (LER) was calculated as the leaf
growth in one day divided by the time interval.

RNA-seq library preparation and sequencing
The fourth leaves of the selected accessions were collected on the second day after leaf
emergence for RNA-seq library preparation as described by Sprangers *et al.*, 2016. Leaves
were in the primordia 5 (P5) stage and were growing at a steady growth rate at the time of
collection. Libraries were prepared in three biological replicates using YourSeq Full
Transcript RNAseq Library Kit (AMARYLLIS NUCLEICS) as per the manufacturer's
protocol. These RNA-seq libraries were sequenced on the Illumina HiSeq platform, and reads
were generated in 150bp paired-end format.

Quality filtering, mapping of reads, and differential gene expression analysis
The reads were quality filtered using the NGSQC toolkit that involved the removal of adapter
sequences and low-quality reads (Phred score <30) (Patel *et al.*, 2012). The quality-filtered

reads were mapped to reference cDNA sequences of rice from the Rice Genome Annotation
Project database (RGAP 7) using default parameters of bowtie2 (Langmead & Salzberg,
2012). SAMtools were used to generate bam alignment files that were used to tabulate raw
counts for each gene (Li *et al*., 2009). The raw counts were normalized to library size and
corrected for library composition bias using the Trimmed Mean of M values normalization
approach (Robinson and Oshlack, 2010). edgeR was used for differential gene expression
analysis (Robinson *et al*., 2010). glm approach with quasi-likelihood F-test was used for
differential gene expression analysis. Significant differentially expressed genes for each pair-
wise comparison were determined based on FDR cut-off < 0.05 and logFC value >1 and <-1.
179

Principle Component Analysis with Self Organizing Map (PCA-SOM) clustering
Normalized read counts (Dataset S7) were used for a gene expression clustering method
(Chitwood *et al*., 2013). For clustering, the genes in the upper 50% of the coefficient of
variation for expression across different accessions were used. The scaled expression values
were used to generate accession-specific gene clusters (Wehrens & Buydens, 2007). Three by
three hexagonal SOM clusters and 100 training interactions were used during clustering. The
SOM clusters were visualized in PCA space, where PC values were calculated based on gene
expression across the accessions with the help of the prcomp R package. The heatmaps were
generated for each cluster using superheat R package.
189

GO Enrichment Analysis
191 GO enrichment analysis was performed using the Plant Transcriptional Regulatory Map GO
192 enrichment tool of the Plant Transcription Factor Database (Tian *et al*., 2019).
193

Gene regulatory network analysis
An unsigned correlation network was constructed using the weighted gene correlation
network (WGCNA) package version 1.68 (Langfelder *et al*., 2008). Data were z-score
normalized (R scale function). The soft-thresholding power was chosen based on a scale-free
topology with a fit index of 0.8. An adjacency matrix with the selected soft-thresholding
power was calculated and transformed into a topological overlap matrix (TOM). Using the
TOM, network properties such as strength were calculated and the network was visualized
using the igraph package version 1.2.5 (Csardi *et al*., 2006) and visNetwork 2.0.9
(https://datastorm-open.github.io/visNetwork/). A Fast-Greedy modularity optimization
algorithm was selected to define modules in the integrated network.

Leaf kinematics analysis

To estimate the division zone size, the basal 4 cm segment of the fourth leaf was collected at different days of emergence up to day five as described by Sprangers *et al.* (2016). Samples were placed in 3:1 (v/v) absolute ethanol: acetic acid, rinsed for 30 min in a buffer containing 50 mM NaCl, 5 mM EDTA, and 10 mM Tris-HCl, pH 7, and nuclei were stained by incubating the samples in the dark for 2 min in the same buffer solution containing 1 mg/ml 49,6-diamino-phenylindole (DAPI). Fluorescent nuclei were observed with a microscope with an epi-fluorescent condenser (Nikon 80i-epifluorescence Microscope), and the images obtained were used to measure the division zone size as the distance between the base of the leaf and the most distal mitotic cell. To determine cell length profiles, fourth leaves at the P5 developmental stage were collected on the second day after leaf emergence at the steady-state growth. The leaves were kept in absolute ethanol for 48 h (with one change at 24 h), then transferred to lactic acid, and stored at 4°C. Samples were mounted on microscope slides using glycerol. The epidermal cells at each millimeter length were visualized using a phase-contrast microscope (LMI, UK) at 40X magnification. The various growth parameters from the leaf kinematics experiments were quantified using formulae explained in Sprangers *et al.* (2016).

GA quantification

The fourth leaves at the P5 developmental stage were collected on the second day after leaf emergence at the steady-state growth, similar to leaf kinematics and transcriptomics experiments, for GA quantification in full leaves. For zone-specific GA quantification, the samples were collected from two different zones of fourth leaves. Zone I was up to 0.8 cm from the leaf base and zone II was from 0.8 cm to 1.6 cm. The samples were lyophilized for 72 hr and finely ground in liquid nitrogen. 25 mg of each sample was extracted in ice-cold buffer (MeOH: H_2O: HCOOH, 15:4:0.1) on a shaker for overnight at 4°C. The internal standard was added to the extract and vortexed for 10 min. The extract was centrifuged for 15 min at 16,000 rcf at 4°C, the supernatant was passed onto a pre-conditioned (MeOH and 0.1% of HCOOH) C18 RP SPE column, and the column was washed with 0.1% HCOOH and 5% MeOH. Finally, elution was done with ice-cold 0.1% HCOOH in Acetonitrile. After evaporation of the solvent, the pellet was re-suspended in 5% MeOH. GA content was then analyzed using LC-MS/MS.

RNA isolation, cDNA synthesis, and qRT-PCR analysis

For zone-specific expression profiling, samples from zone I and II of the fourth leaves were collected as explained above for zone-specific GA quantification. Total RNA was isolated using Trizol reagent (Invitrogen) and treated with RNase-free DNase I (Thermo Fisher Scientific). 1.5 µg of total RNA was converted to first-strand cDNA using RevertAid First-Strand cDNA Synthesis Kit (Thermo Fisher Scientific). The quantitative real-time-PCR analysis was performed on the CFX Connect Real-Time System (Bio-rad) using PowerUp™ SYBR® Green fluorescence dye (Thermo Fisher Scientific). The primer pairs for qRT-PCR were designed based on gene sequences obtained from RGAP 7 (Table S6). The primer efficiency was analyzed using a standard curve method. From each species five different cDNA concentration (25,50,100,200 and 400ng) used to generate standard curve and primer efficiency was calculated using the formula E = (10[-1/slope] 257 -1) *100. The primer specificity of amplified gene products across the selected rice genotypes was viewed through dissociation curve analysis. Relative expressions of the target genes were analyzed using actin (LOC_Os03g50885.1) as the internal control applying the $2^{-\Delta\Delta Ct}$ method for fold change and $2^{-\Delta Ct}$ method for relative transcript levels quantification (Livak & Schmittgen, 2001). The expression pattern of each gene was confirmed by three independent biological experiments.

Transient gene silencing line generation

Virus-induced gene silencing (VIGS) mediated transient silencing lines for *OsGA20OX2, OsSLR1*, *OsGRF7,* and *OsGRF8* genes were generated using the mVIGS vector as previously described by Kant et al. (2017). Recombinant vector constructs were transformed into Agrobacterium strain EHA105, and a single Agrobacterium colony was used for primary culture. After overnight incubation at 28 ℃, a secondary culture was grown to an OD600 of 0.6 to 0.8 with 200 µM acetosyringone. Cells were harvested after centrifugation and resuspended in 10 mM MES, 10 mM MgCl2, and 200 µM acetosyringone. The meristematic region located at the basal region (junction of shoot and root) of 6-day old rice seedlings were injected with 50 µL of bacterial suspension using a vertically positioned syringe as described by Kant et al., 2015. After inoculation seedlings were transferred onto sterile Whatman No. 1 filter paper immersed in MS medium placed on a solid support with its ends dipped into a reservoir containing the nutrient medium. To avoid drying, plant roots were covered with moist tissue paper and transferred to tubes containing MS medium 24 h post-inoculation and were maintained at 28°C under conditions described above. As positive controls, we generated transient silencing lines for chlorophyll biosynthesis gene (*ChlH*) and confirmed

positive transgenic plants by observing leaves' chlorosis phenotype and gene expression
analysis (Fig. S9a-d).

Results

Leaf size variation across the selected cultivated and wild rice accessions

Quantification of the leaf length and the maximum leaf blade width of the fully-grown
second to eighth leaves of the twelve rice accessions showed a strong variation in leaf length
and width (Tables 1, S2). The length of the fully-grown eighth leaf varied from 44.8 cm in a
cultivated rice *Oryza sativa ssp. indica* cv. IR64 to 87.8 cm in a wild rice *Oryza australiensis*
(Table 1). The length of the fully-grown eighth leaf was significantly higher in the five wild
rice accessions *O. australiensis, O. rufipogon, O. latifolia, O. glumaepatula*, and *O. punctata*
than the Asian cultivated rice varieties and the African cultivated rice *O. glaberrima*.
Significant variation in leaf width was also observed across the cultivated and wild rice
accessions (Table S2). We then selected four genetically diverse accessions, which captured
the leaf length variations across the investigated accessions, for detailed phenotypic and
genetic characterization. The four selected accessions were: IR64 (an *indica* variety of the
cultivated species *O. sativa*) and Nipponbare (a *japonica* variety of the cultivated species *O.*
sativa) with smaller leaves, the wild rice *O. australiensis* with the longest leaves, and *O.*
glaberrima (the African cultivated rice) with intermediate leaf length (Table 1; Fig. 1a). The
leaf phenotype of second to tenth leaves of the four accessions further confirmed the variation
in leaf length pattern (Fig. 1b, S1).

Quantification of the fourth leaf length of the selected four accessions consistently
showed longer leaf in the wild rice *O. australiensis* compared to the cultivated accessions at
different days of emergence (Fig. 1c, d). *O. glaberrima* showed intermediate leaf length
between the wild rice *O. australiensis* and cultivated Asian rice varieties at different days of
emergence. Consistent with the leaf length, the leaf elongation rate (LER) was higher for the
wild rice than the cultivated accessions at different days of emergence (Fig. 1e). Taken
together, the wild rice *O. australiensis* showed remarkably longer leaves than the cultivated
varieties, and *O. glaberrima* also developed significantly longer leaves than IR64 and
Nipponbare.

Transcriptome profiling identified cell cycle, GA, and GRFs as the key regulators of
rice leaf length

To identify genes and pathways controlling the leaf length differences, we compared
the transcriptomes of the developing fourth leaves of the selected accessions. The maximum
number of differentially expressed genes (DEGs) was found between Nipponbare and *O.*
australiensis, whereas a lower number of DEGs was found between cultivated accessions
(Table S3, Dataset S1). Since the leaves of the wild rice *O. australiensis* were remarkably
longer than IR64 and Nipponbare, we first compared the gene expression profiles between
the leaves of *O. australiensis* and two cultivated varieties of *O. sativa*. 2,384 genes were
expressed at higher levels and 3,870 genes were expressed at lower levels in growing leaves
of *O. australiensis* compared to both IR64 and Nipponbare (Fig. 2a). The genes expressed at
higher levels showed enrichment of GO terms 'cell cycle', 'development process', 'cell
growth', and 'cell morphogenesis' (Dataset S2). The enriched GO terms for the genes
expressed at lower levels included 'photosynthesis', 'biosynthetic process', and 'signaling'.
We also compared the gene expression differences between the significantly longer leaves of
O. glaberrima and the two cultivated varieties of *O. sativa*. The 1,125 genes expressed at
higher levels showed enrichment of GO terms 'cell cycle', 'development process', 'growth',
and 'hormone transport', whereas 1,290 genes expressed at lower levels were enriched for the
term 'metabolic process' (Fig. S2a; Dataset S2). We then analyzed the differentially
expressed genes in the wild rice *O. australiensis* compared to all three cultivated accessions.
The 1,707 shared genes expressed at higher levels in *O. australiensis* compared to all three
cultivated accessions showed enrichment of GO terms "development process", "cell cycle",
and "growth", whereas the GO term "biosynthetic process" was enriched for genes expressed
at lower levels (Fig. S2b; Dataset S2).
Principle component 1 (PC1), explaining 39% variation in the dataset, primarily
explained the expression differences of the wild rice *O. australiensis* compared to the three
cultivated accessions (Fig. S3a, b). PC2, explaining 31% of the overall variation, explained
the expression differences of *O. glaberrima* compared to the other three accessions. The
clusters generated by PCA-SOM provided the accession-specific resolution of the transcripts
(Fig. 2b; Dataset S3). Since *O. australiensis* and *O. glaberrima* had significantly longer
leaves compared to IR64 and Nipponbare, we analyzed the genes and enriched pathways in
the clusters specific to either *O. australiensis* or *O. glaberrima* or both. The cluster 4 that had
genes specific to both *O. australiensis* and *O. glaberrima* showed enrichment of terms 'cell
cycle', 'development process', 'response to gibberellin', and 'response to auxin' (Fig. 2b;
Dataset S4). The transcripts in *O. australiensis* specific cluster 7 and *O. glaberrima* specific
cluster 1 were enriched for 'development process' and 'cell cycle'.

Since transcription factors, phytohormones, and cell division and expansion are
crucial for leaf size regulation, we compared the expression of the related genes from the
relevant clusters across the selected accessions (Dataset S5). Several cell cycle genes, such as
CYCLINB, CYCLINA, and *CDKD1*, as well as cell cycle regulators, such as *E2F2* and *RB1*,
were expressed at higher levels in *O. australiensis* and *O. glaberrima* than IR64 and
Nipponbare (Fig. 2c). Further, transcription factor genes known to regulate leaf growth and
development, such as *GRF*-family, *TCP*-family, *YABBY*-family, *NAM*-family, and *ANT*, were
expressed at higher levels in *O. australiensis* and *O. glaberrima* (Fig. 2d). Among the *GRFs*,
OsGRF7 and *OsGRF8* specifically showed higher expression in *O. australiensis* leaves
compared to the cultivated varieties. GA biosynthesis and signaling genes that included *CPS,*
KAO, and *GID1* were expressed at higher levels in *O. australiensis* and *O. glaberrima* than
IR64 and Nipponbare (Fig. 2e). Moreover, differential expressions of auxin and cytokinin
biosynthesis and signaling genes were also observed across the selected accessions (Fig.
S3c).

The coexpression network of genes from the relevant clusters showed 3 major
modules including more than 20 nodes (Fig. 2f; Dataset S6). The network showed significant
connections for GA biosynthesis and signaling genes *CPS* and *GID1L2,* cell cycle-related
genes *CYCLIN* and *CDKD1*, as well as *GRF7*. Investigation of the hub genes in module 2
showed *CDKD1* as one of the highly connected genes, which was connected to GA receptor
GID1L2, and *GRF7* (Fig. S3d). Taken together, transcript profiling along with gene co-
expression network highlighted cell cycle, GA, and transcription factors, such as OsGRF7
and OsGRF8, as key contributors of rice leaf length regulation.

GA-levels correlated with the leaf length variation across the cultivated and wild rice
accessions

Since the transcript profiling indicated the involvement of GA for the leaf length
differences across the accessions, we quantified GA in the fourth leaves of the cultivated and
wild rice accessions that showed significant variation for leaf length (Fig. S4). We detected
significantly higher levels of GA_4 in the leaves of wild rice accessions with longer leaves
compared to the cultivated rice accessions (Fig. 3a). A strong significant positive correlation
between leaf length and GA_4 levels was observed for the selected cultivated and wild rice
accessions, suggesting a key role of GA in rice leaf length regulation (Fig. 3b). We further
detected a higher level of GA_7 in addition to GA_4 in the leaves of *O. australiensis* (wild rice

with longest leaves among all the investigated accessions) compared to Nipponbare
(cultivated rice variety with smaller leaves) (Fig. 3c).

To confirm the GA-mediated rice leaf length regulation, we examined the effects of
exogenous treatment of GA$_3$ and GA-biosynthesis inhibitor Paclobutrazol (PAC) on leaf
length of the two contrasting accessions, *O. australiensis* and Nipponbare. GA$_3$ treatment
promoted the leaf length of both Nipponbare and *O. australiensis* (Fig. 3d, e, and S5a, b). The
extent of increase in leaf length was more in Nipponbare (~1.6-fold) than *O. australiensis*
(~1.3-fold). Exogenous treatment of PAC drastically reduced the leaf length of both the
accessions. PAC inhibited the leaf length of Nipponbare by ~2-fold, and of *O. australiensis*
by ~2.5-fold. The higher extent of leaf length increase in Nipponbare by GA$_3$ treatment and
higher extent of leaf length decrease in *O. australiensis* under PAC along with a general
higher GA content in wild rice accessions with longer leaves suggested that GA levels could
be determining for the differences in the leaf length across the cultivated and wild rice
accessions.

Leaf kinematics confirmed the GA-mediated control of division zone to regulate rice
leaf length

Since transcriptomic comparison showed higher expression of cell-cycle related genes
in the wild rice accessions than the cultivated varieties, we performed DAPI staining and
kinematics analyses on the growing fourth leaves to investigate the differences in cell
division and growth zones across the selected accessions. *O. australiensis*, with longer leaves
than the cultivated varieties, showed dividing cells for longer length from the leaf base, and
thus longer division zone at different days of emergence (Fig. S6a). *O. glaberrima* also
showed a longer division zone compared to IR64 and Nipponbare. Detailed kinematics
analyses on the fourth leaves of each selected accession confirmed the differences in the
division zone (Fig. 4a, b, and S6; Table 2). All the four selected accessions showed a
sigmoidal curve for epidermal cell length profile, consisting of a division zone at leaf base
followed by elongation zone and maturation zone (Fig. 4a, b and S6f, g). The cultivated
varieties Nipponbare and IR64 showed division zone extending up to 7.8mm and 7.5mm
from the leaf base, respectively (Fig. 4a and S5b, d, f). *O. australiensis*, with the longest
leaves, showed a remarkably longer division zone that extended to 16.7mm from the leaf base
(Fig. 4b and S5c, e). The longer division zone in *O. australiensis* leaves was accompanied by
a faster cell production rate and leaf elongation rate (Table 2). *O. glaberrima* also showed a

longer division zone as well as a higher cell production rate compared to IR64 and
Nipponbare (Fig. S5g).

407    We then asked the question of whether GA regulates rice leaf length by controlling
cell proliferation and size of the division zone. To this end, we performed leaf kinematics
analysis on the fourth leaves of *O. australiensis* and Nipponbare under GA_3 and PAC
treatment. We observed that the leaf length promotion in response to GA treatment was due
to an increase in the size of the division zone in both the accessions (Fig. 4c, d, and S5c, d).
However, the impact of GA on cell division was more striking in Nipponbare than *O.*
australiensis in terms of the size of the division zone, cell production rate, and the number of
cells in the meristem (Tables S4 and S5). The inhibitory effect of PAC on the leaf length was
also clearly due to the reduced size of the division zone in both the accessions (Fig. 4c, d, and
S5c, d). Interestingly, the kinematics profile of GA-treated Nipponbare leaves resembled that
of *O. australiensis* leaves under control conditions (Fig. 4b, c). Together, the leaf kinematics
profile of the contrasting rice accessions and the effects of exogenous GA and PAC
treatments on the division zone confirmed that the higher GA levels promoted the rice leaf
length by increasing the size of the division zone.

Growth-zone-specific expression analysis of GA-related genes and associated GA levels

423    Next, we asked if zone-specific expressions of GA-biosynthesis and signaling genes
along with associated GA levels explain the leaf size differences. To this end, we quantified
the expression of GA biosynthesis genes in two leaf zones of Nipponbare and *O.*
australiensis: zone I (leaf base – 0.8 cm; division zone in both the accessions) and zone II
(0.8 – 1.6cm; elongation zone in Nipponbare, where division extends in *O. australiensis*).
The transcript levels of GA biosynthetic genes *OsGA20OX2* and *OsGA3OX2* were higher in
the zone I of *O. australiensis*, which showed longer leaves and had higher GA levels in their
leaves than Nipponbare (Fig. 5a, b). Furthermore, the expression level of the GA catabolic
gene *OsGA2OX4* was lower in zone II of *O. australiensis* than Nipponbare (Fig. 5c).
Consistent with the expression of GA-biosynthesis and catabolic genes, more GA_4 and GA_7
levels were detected in both zones of *O. australiensis* than Nipponbare (Fig. 5e, f). The
transcript levels of *OsSLR1*, encoding a GA-signalling repressor DELLA protein, was also
higher in both the regions of Nipponbare leaves (Fig. 5d). Taken together, higher expression
of GA biosynthetic genes *OsGA20OX2* and *OsGA3OX2* in zone I, lower expression of GA
catabolic gene *OsGA2OX4* in zone II, and lower expression of GA-signalling repressor

OsSLR1 in *O. australiensis* promoted more zone-specific GA levels, and thus higher GA

effects in *O. australiensis* than Nipponabre.

Higher GA levels expanded the expression domain of *GRFs* and cell-cycle-related genes

Similar to the expression of GA-biosynthesis and signaling genes, we observed

remarkably higher transcript levels of *OsGRF7* and *OsGRF8* in both zones I and II of *O.*

australiensis leaves than Nipponbare (Fig. 6a, b). Further to corroborate the involvement of

OsGRF7 and *OsGRF8* in rice length regulation, we analyzed the transcripts level of the two

*GRF*s in the two growth zones of cultivated and wild rice accessions initially used for leaf

phenotyping and GA quantification. Wild rice accessions with longer leaves and higher GA

content had a higher abundance of *OsGRF7* and *OsGRF8* transcripts in division zones (Fig.

S7a, b). The transcript levels of *OsCYCB1;4* and *OsCYCA3;2* were also higher in both zone I

and II of the *O. australiensis* than Nipponbare (Fig. 6c, d). Limited transcript levels of

OsCYCB1;4 and *OsCYCA3;2* in zone II of Nipponbare leaves matched with the negligible

cell division activity beyond 0.8cm in the accession. The significantly lower expression level

of cell cycle inhibitor *OsKRP4* and a higher level of cell cycle activator *OsE2F2* in basal

regions of *O. australiensis* leaves than Nipponbare further reflected on the differences in cell

cycle between the two accessions (Fig. S7c, d). The exogenous GA treatment remarkably

induced the expression levels of *OsGRF7* and *OsGRF8* as well as *OsCYCB1;4* and

OsCYCA3;2 in zone II of Nipponbare (Fig. 6e-h). Moreover, the expression of *OsSLR1* was

suppressed in both the zones in response to GA treatment (Fig. S8a). PAC treatment led to

significant suppression of the expressions of *GRF*s and *CYCLIN*s in the division zone of

Nipponbare (Fig. S8b). Together, gene expression analyses in different leaf zones of the

selected cultivated and wild rice accessions as well as in response to the exogenous GA

treatment strongly supported our hypothesis of GA-mediated cell cycle control through

OsGRF7 and OsGRF8 to determine the rice leaf length.

Gene silencing confirmed the regulation of division zone size by GA and GRFs to

control rice leaf length

To confirm the GA-mediated leaf length differences between the contrasting

cultivated and wild rice accessions via specific GRFs, we performed transient silencing of

genes encoding a GA biosynthesis enzyme GA20OX2, a GA signaling repressor SLR1, and

Growth-Regulating Factors GRF7 and GRF8 in *O. australiensis* and Nipponbare. Since

extended division zone and longer leaf length in the wild rice *O. australiensis* was associated

with higher GA levels in leaves, we checked the effects of reducing GA levels by silencing
GA20OX2 in the wild rice. The silenced *O. australiensis* lines showed a remarkable reduction
in the leaf length associated with decreased division zone size and lower expression of
OsGRF7, OsGRF8, and *CYCLIN*s compared to the control plants (Fig. 7a-c and S10a).
Similar to *OsGA20OX2*, silencing of *OsGRF7* and *OsGRF8* in *O. australiensis* reduced the
leaf length along with the smaller size of the division zone and reduced expression of
downstream *CYCLIN* genes (Fig. 7a-c, and S10c). Silencing of *OsGA20OX2*, *OsGRF7*, and
OsGRF8 in Nipponbare also led to a reduction in leaf length, similar to exogenous PAC
treatment, with reduced size of the division zone as well as lower expression levels of
OsGRF7, *OsGRF8,* and *CYCLIN*s (Fig. 7d-f and S10b, d). In contrast, silencing of *OsSLR1* in
both the accessions led to a longer leaf, increased size of division zone, and higher expression
levels of *OsGRF7, OsGRF8,* and *CYCLIN*s compared to the control (Fig. S9e-h and S10e-f).
Together, leaf phenotypes of *O. australiensis* and Nipponbare silencing lines confirmed the
regulation of cell division zone size by GA and downstream OsGRF7 and OsGRF8
transcription factors to determine the rice leaf length.
487
Discussion
Natural variation for the leaf size in a crop species and related wild accessions is an
excellent resource for understanding the genetic and hormonal basis of leaf size regulation.
Studies have reported remarkable variations in the leaf morphological and anatomical
features across cultivated and wild species of genus *Oryza*, suggesting a strong genetic
control on leaf features of rice (Chatterjee *et al*., 2016; Mathan *et al*., 2021b). We performed
detailed leaf phenotyping of seven cultivated and five wild rice accessions followed by GA
quantification in leaves. Four accessions were then selected for detailed mechanistic studies
that not only represented a significant genetic diversity but also covered the entire range of
leaf length variations among the investigated accessions.
Differences in the leaf length are attributed to basic cellular features and leaf
developmental programing (Gonzalez *et al*., 2012; Kierzkowski *et al*., 2019). Consistent with
this, cell cycle and development were the obvious enriched terms for the genes expressed at
higher levels in the accessions with longer leaves. Detailed leaf kinematics analyses for the
contrasting cultivated and wild rice accessions precisely established the predominant role of
cell division activity, and thus the size of the division zone, for defining the rice leaf length.
Higher cell production rate and expansion of cell division to a longer length from leaf base
drove the longer leaf phenotype in the wild rice *O. asutraliensis* than the cultivated varieties

(Fig. 4; Table 2). Cell cycle duration and cell division rate were comparable across the
accessions, suggesting that the longer cell division zone and increased cell production rate at
the leaf base resulted in the production of a higher number of cells that contributed to longer
leaves in the wild rice (Gazquez & Beemster, 2017). Higher expression levels and expanded
expression domains of genes encoding cyclins and cyclin-dependent kinases in the growing
leaves of *O. asutraliensis* than Nipponbare further supported the key importance of the cell
division activity for the rice leaf length regulation (Fig. 2c and 6c, d). *O. asutraliensis*,
besides showing the expansion of the division zone, showed longer epidermal cells compared
to cultivated varieties, suggesting that the cell elongation also contributes to the longer leaf
length of the wild rice (Fig. 4b).

Spatial and temporal regulation of cell division activity in plants is under complex
regulation of phytohormones and transcription factors (Nelissen *et al.*, 2012; Takatsuka &
Umeda, 2014). We detected auxin-, cytokinin-, and GA-related genes among the
differentially expressed genes for the selected rice accessions. Besides, genes encoding
transcription factors known to regulate cell division, such as GRFs and TCPs, were also
differentially expressed between the contrasting rice accessions (Vercruysse *et al.*, 2020b).
GA is shown to control the cell division activity via the involvement of GRF transcription
factors in Arabidopsis and maize (Lantzouni *et al.*, 2020; Fina *et al.*, 2017). Our study shows
that this mechanism could have been instrumental in optimizing rice leaf length during the
domestication as wild rice species with longer leaves had higher endogenous GA levels than
the cultivated varieties. GA- and GRF- mediated spatial control of cell division activity
explain the difference in the size of the division zone leading to leaf length diversity in rice.
Comparative transcriptome profiling on full leaves showed higher expression of GA-
biosynthesis and signaling genes as well as GRF transcription factors in the longer leaves of
O. australiensis and *O. glaberrima* than IR64 and Nipponbare, suggesting the key functions
of GA and downstream GRFs to regulate the division zone size (Fig. 2d, e). Given the spatial
genetic regulation for the leaf size control, functions of these factors at the leaf basal regions
with cell division activity could be critical to the final leaf size. Consistent with this, our
results on zone-specific expression profiling supported the key role of local differences in the
expression of the relevant genes at the leaf basal region for leaf size regulation.

In support of the GA-mediated regulation of rice leaf length by expanding cell
division activity, growing leaves of *O. australeinsis* had higher levels of endogenous GAs in
the basal division zone than Nipponbare (Fig. 5 e, f). Higher expression levels of GA
biosynthesis genes *OsGA20OX2* and *OsGA3OX2* in the basal division zone of *O.*

australeinsis leaves than Nipponbare corroborated the higher GA content in the wild rice

(Fig. 5a, b). In addition, lower expression of GA-catabolic gene *OsGA2OX4* in the transition

zone of the wild rice than Nipponbare likely maintained higher GA levels in the zone.

Consistent with these, exogenous GA-treatment increased the rice leaf length by extending

the zone where cell division operates. GA is known to induce cell elongation response (Jones

& Kaufman, 1983). Exogenous GA-treatment, however, increased the leaf length without

altering the mature epidermal cell length, suggesting that GA primarily affects rice leaf length

via regulation of cell division (Tables S4 and S5). Finally, reduction in leaf length of both

cultivated rice Nipponbare and wild rice *O. australiensis* by silencing *OsGA20OX2* that

involved reduced division zone and downregulation of *OsGRF7, OsGRF8,* and *CYCLIN*s

confirmed the central role of GA in rice leaf length regulation through the control of the size

of division zone (Fig. 7a-f and S10a, b).

GRF transcription factors promote cell division activity by directly binding to the

promoters of *CYCA1;1* and *CDC20* (Li *et al*., 2018). OsGRF7 regulation of GA-biosynthesis

and auxin-signaling genes control the rice plant architecture (Chen *et al*., 2020). In addition,

OsGRF1 is shown to influence rice leaf growth (van der Knaap *et al*., 2000). We identified

specific OsGRFs regulating cell division activity to control the size of the division zone for

final rice leaf size determination by genetic comparison of wild rice with longer leaves with

the cultivated varieties. Higher *OsGRF7* and *OsGRF8* expression in growing leaves of *O.*

australiensis with longer leaves from RNAseq analysis along with more *OsGRF7* and

OsGRF8 transcripts in division zone of wild rice accessions than cultivated varieties

suggested that *Os*GRF7 and *Os*GRF8 may connect GA-levels and size of division zone for

rice leaf length regulation (Fig. 6a, b and S7a, b). Interestingly, *OsGRF7* and *OsGRF8* were

found to be closely related to *AtGRF1* and *AtGRF2*, which are shown to influence

Arabidopsis leaf size, in phylogenetic analysis (Kim *et al*., 2003; Zhang et al., 2008; Chen *et*

al., 2020). GA-treatment, increasing the leaf length by increasing the domain of cell division,

induced the *OsGRF7* and *OsGRF8* expression in the division zone, further, suggesting the

OsGRF7 and OsGRF8 functions downstream to GA (Fig. 6e-h). Consistent with this,

silencing of *OsGRF7* and *OsGRF8* resulted in reduced rice leaf length with an associated

decrease in the expression of the *CYCLIN*s (Fig. 7 and S10c, d). Silencing of *OsSLR1*, a

DELLA protein that inhibits GRF functions to repress GA signaling and whose expression is

suppressed by exogenous GA treatment, led to increased leaf length along with higher

expression of *OsGRF7* and *OsGRF8* further supporting the functions of OsGRF7 and

OsGRF8 to integrate GA levels and size of division zone to regulate rice leaf length (Fig.
S9e-h).

In conclusion, we present a detailed cellular basis and underlying genetic mechanism
for leaf length regulation in rice, a staple diet for more than half of the world's population.
We showed that the size of the cell division zone regulated by GA and downstream OsGRF7
and OsGRF8 transcription factors determines the final rice leaf length and explains the
variation in leaf length across the cultivated and wild rice accessions (Fig. 8). Higher GA
levels and increased abundance of *OsGRF7* and *OsGRF8* resulting in expanded cell division
zone explained the longer leaves of the wild rice accessions compared to the cultivated
varieties (Fig. 8). The established GA-OsGRF7/OsGRF8 module to control the division zone
size could not only be a general regulator of rice leaf length but might also have contributed
to the optimization of leaf size during domestication. The module could, further, be a way for
plants to achieve cellular plasticity in response to their growth environment.

586
587
588

Acknowledgments
This work was supported by the core funding from the National Institute of Plant Genome
Research as well as Ramalingaswamy Re-entry Fellowship (BT/RLF/Re-entry/05/2013) and
Rice Network Project (BT/Ag/Network/Rice/2019-20) from the Department of
Biotechnology, Ministry of Science and Technology, India. V.J. and K.S. acknowledge their
UGC-SRF and SERB-NPDF fellowships, respectively. We thank the NIPGR Metabolomics
Facility for Gibberellic Acid (GA) quantification and NIPGR Central Instrumentation Facility
for their support. Seeds of wild rice species and *Oryza glaberrima* were kindly provided by
Dr. Kuldeep Singh and Dr. Kumari Neelam, Punjab Agricultural University, Ludhiana, India.

Author contributions: V.J. and A.R, conceptualized the study. V.J., K.S., and A.R. designed
experiments. V.J. and A.C. performed experiments. V.J., R.R., Y.I., and A.R. analyzed
transcriptomic data. V.J. and K.S. analyzed leaf kinematics data. V.J. and A.R. compiled
figures and wrote the manuscript. V.J., K.S. and A.R. edited the final manuscript with the
contribution of all the authors.

Data Availability Statement:

The data that supports the findings of this study are available in the supplementary material
of this article. The quality-filtered reads, which were used to get the normalized read counts
and for DE analysis, are deposited to the NCBI Short Read Archive under accessions
SRR15144741, SRR15144745, SRR15144747, and SRR15144751.

Tables

Table 1 Quantification of leaf length (cm) of fully-developed second leaf to the eighth leaf of the seven cultivated and five wild rice accessions. Each value represents mean ± SD, where n = 15 data points from different plants. Different letters indicate statistically significant differences according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P < 0.05$.

Oryza species	Cultivar/ accession number	Leaf number						
		Leaf 2	Leaf 3	Leaf 4	Leaf 5	Leaf 6	Leaf 7	Leaf 8
Oryza sativa ssp. japonica	Nipponbare	3.94 ± 0.35 (a)	11.80± 1.15 (a)	21.96 ± 1.53 (a)	26.92 ± 2.00 (a)	38.11 ± 1.73 (a)	44.37 ± 1.40 (a)	48.71 ± 2.84 (a)
Oryza sativa ssp. indica	IR 64	5.60 ± 0.22(b)	11.75 ± 0.85 (a)	20.95 ± 1.91 (a)	26.25 ± 1.72 (a)	34.05 ± 1.07 (a)	41.56 ± 1.48 (b)	44.85 ± 2.75 (b)
Oryza sativa ssp. indica	Vandana	4.54 ± 0.62 (a)	11.50 ± 1.43 (a)	21.12 ± 1.55 (a)	29.06 ± 1.34 (a)	32.59 ± 1.71 (b)	37.36 ± 1.77 (c)	45.59 ± 1.74 (ab)
Oryza sativa ssp. indica	Swarna	4.70 ± 0.44 (a)	14.93 ± 1.24 (b)	27.92 ± 2.52 (b)	39.88 ± 2.65 (b)	46.90 ± 1.87 (c)	52.76 ± 2.86 (d)	58.72 ± 3.03 (c)
Oryza sativa ssp. indica	Pusa Basmati1	6.25 ± 0.54(b)	14.66 ± 2.36 (b)	28.48 ± 3.11 (b)	37.11 ± 2.01 (c)	42.98 ± 1.76 (d)	48.30 ± 2.26 (a)	55.17 ± 1.94 (d)
Oryza sativa ssp. indica	Nagina 22	4.21 ± 0.38 (a)	14.55 ± 1.52 (b)	27.98 ± 1.30 (b)	38.50 ± 2.29 (bc)	44.56 ± 2.14 (de)	53.41 ± 2.59 (d)	58.77 ± 1.79 (c)
Oryza glaberrima	IR 102925	6.00 ± 0.30(b)	14.95 ± 1.65 (bc)	26.90 ± 1.18 (b)	38.20 ± 1.19 (c)	45.90 ± 2.42 (ce)	53.70 ± 2.40 (d)	59.50 ± 2.82 (c)
Oryza glumaepatula	IR104151	8.01 ± 0.56 (c)	23.24 ± 1.68 (e)	37.26 ± 3.10 (ce)	43.30 ± 1.30 (d)	46.76 ± 1.51 (ce)	53.35 ± 2.05 (d)	63.91 ± 3.07 (e)
Oryza punctata	IRGC105137	6.08 ± 0.63(b)	17.67 ± 1.19 (cd)	34.38 ± 1.27 (cd)	47.52 ± 1.39 (e)	51.92 ± 1.31 (f)	58.26 ± 1.23 (e)	70.67 ± 2.04 (f)
Oryza latifolia	IRGC 99596	10.15 ± 0.62(d)	18.32 ± 1.42 (d)	31.52 ± 2.05 (d)	46.13 ± 1.19 (e)	53.12 ± 1.10 (f)	60.63 ± 2.26 (e)	70.93 ± 1.68 (f)
Oryza rufipogon	IRGC 99562	8.57 ± 1.57(c)	23.75 ± 3.20 (e)	37.55 ± 2.80 (e)	47.00 ± 4.47 (e)	66.40 ± 6.29 (g)	74.72 ± 3.55 (f)	85.56 ± 1.67 (g)
Oryza australiensis	IRGC 105272	20.57 ± 2.68 (e)	34.19 ± 3.44 (f)	50.66 ± 4.55 (f)	56.57 ± 2.60 (f)	64.74 ± 4.36 (g)	69.50 ± 0.71 (f)	87.80 ± 5.27 (g)

Table 2 Quantification of leaf kinematics parameters of the growing fourth leaf of the selected cultivated and wild rice accessions. Each value represents mean ± SD (n = 15 for mature leaf length and leaf elongation rate, n = 5 for all other parameters). Different letters indicate statistically significant differences according to one-way ANOVA followed by post-hoc Tukey HSD calculation at P < 0.05.

Growth parameters	*O. sativa* cv. IR 64	*O. sativa* cv. Nipponbare	*O. glaberrima*	*O. australiensis*
Mature leaf length (cm)	20.06 ± 1.38(a)	21.46 ± 0.81(a)	26.98 ± 1.01(b)	41.38 ± 4.25(c)
Leaf elongation rate (mm/h)	1.32 ± 0.01(a)	1.34 ± 0.01(a)	2.01 ± 0.01(b)	2.41 ± 0.01(b)
Meristem length (mm)	7.52 ± 0.14(a)	7.81 ± 0.34(a)	12.32 ± 0.34(b)	16.66 ± 0.64(c)
Length of the growth zone (mm)	24.00 ± 1.00(a)	31.33 ± 0.58(b)	24.67 ± 1.53(a)	34.33 ± 0.58(c)
Mature cell length (µm)	70.61 ± 0.15(a)	64.26 ± 1.41(b)	66.00 ± 1.07(b)	84.57 ± 1.32(c)
Cell production rate (cells/h)	18.76 ± 0.12(a)	20.88 ± 0.40(b)	30.44 ± 0.51(c)	28.49 ± 0.42(d)
Number of cells in the meristem	738.29 ± 27.50(a)	715.92 ± 17.05(a)	842.73 ± 31.50(b)	904.75 ± 52.90(b)
Number of cells in the growth zone	1193.63 ± 70.37(a)	1328.80 ± 26.33(b)	1100.53 ± 43.44(ac)	1211.09 ± 38.48(abc)
Number of cells in the elongation zone	455.35 ± 68.50(a)	612.88 ± 41.50(b)	257.80 ± 38.30(c)	306.34 ± 19.50(c)
Average cell division rate (cell. cell^{-1}.h^{-1})	0.03 ± 0.001(a)	0.03 ± 0.001(ab)	0.04 ± 0.001(bc)	0.03 ± 0.001(ab)
Cell cycle duration (h)	27.29 ± 1.15(a)	23.78 ± 0.99(b)	19.20 ± 1.03(c)	22.00 ± 0.98(b)
Time in the elongation zone (h)	24.26 ± 3.53(a)	29.34 ± 1.62(a)	8.47 ± 1.29(b)	10.76 ± 0.79(b)
Time in division zone (h)	260.01 ± 12.40(a)	225.54 ± 10.10(b)	186.66 ± 11.05(c)	216.13 ± 11.45(bc)
Length of cells leaving meristem (µm)	15.15 ± 0.37(a)	17.79 ± 0.41(a)	31.33 ± 1.41(b)	39.93 ± 3.36(c)
Average cell expansion rates (µm. µm^{-1}.h^{-1})	0.06 ± 0.01(ac)	0.04 ± 0.003(a)	0.09 ± 0.01(bc)	0.07 ± 0.01(ac)

References

Achard P, Gusti A, Cheminant S, Alioua M, Dhondt S, Coppens F, Beemster GT, Genschik P. **2009**. Gibberellin signaling controls cell proliferation rate in Arabidopsis. *Current Biology* **19:** 1188–1193

Andriankaja M, Dhondt S, De Bodt S, Vanhaeren H, Coppens F, De Milde L, Mühlenbock P, Skirycz A, Gonzalez N, Beemster GTS, et al. 2012. Exit from proliferation during leaf development in Arabidopsis thaliana: a not-so-gradual process. *Development Cell* **22:** 64–78

Avramova, V, Sprangers, K, and Beemster, GTS. 2015. The Maize Leaf: Another Perspective on Growth Regulation. *Trends in Plant Science* **20:** 787-797

Bar M, Ori N. **2014**. Leaf development and morphogenesis. *Development* **141**(22)**:** 4219-30

Beemster GT, De Veylder L, Vercruysse S, West G, Rombaut D, Van Hummelen P, Galichet A, Gruissem W, Inze D, and Vuylsteke M. **2005**. Genome-wide analysis of gene expression profiles associated with cell cycle transitions in growing organs of Arabidopsis. *Plant Physiology* **138:** 734–743

Beltramino M, Ercoli M, Debernardi J, Goldy C, Rojas A, Nota F, Alvarez M, Vercruyssen L, Inzé D, Palatnik J, Rodriguez R. **2018.** Robust increase of leaf size by Arabidopsis thaliana GRF3-like transcription factors under different growth conditions. *Scientific Report* **8:** 13447

Chatterjee J, Dionora J, Elmido-Mabilangan A, Wanchana S, Thakur V, Bandyopadhyay A, Brar D, Quick W. **2016**. The evolutionary basis of naturally diverse rice leaves anatomy. *PLoS ONE* **11:** e0164532

Chen Q, Li W, Tan L, Tian F. **2021**. Harnessing knowledge from maize and rice domestication for new crop breeding. *Molecular Plant* **14:** 9-26

Chen Y, Dan Z, Gao F, Chen P, Fan F, Li S. 2020. Rice GROWTH-REGULATING FACTOR7 Modulates Plant Architecture through Regulating GA and Indole-3-Acetic Acid Metabolism. *Plant Physiology* **184** (1)**:** 393-406

Chitwood D, Kumar R, Headland L, Ranjan A, Covington M, Ichihashi Y, Fulop D, Jiménez-Gómez J, Peng J, Maloof J, Sinha N. 2013. A quantitative genetic basis for leaf morphology in a set of precisely defined tomato introgression lines. *The Plant Cell* **25:** 2465–2481

Chitwood DH, Sinha NR. 2016. Evolutionary and environmental forces sculpting leaf development. *Current Biology* **26**(7)**:** 297-306

Choi J, Kende H. 2004. Whole genome analysis of the OsGRF gene family encoding plant-specific putative transcription activators in rice (*Oryza sativa* L.). *Plant Cell Physiology* **45:** 897-904

Csardi G, Nepusz T. 2006. The igraph software package for complex network research. *InterJournal, Complex Systems* **1695:** 1–9

Davière JM, Achard P. 2013. Gibberellin signaling in plants. *Development* **140:** 1147-51

Du F, Guan C, Jiao Y. 2018. Molecular mechanisms of leaf morphogenesis. *Molecular Plant* **11**(9)**:** 1117-1134

Efroni I, Eshed Y, Lifschitz E. 2010. Morphogenesis of simple and compound leaves: a critical review. *The Plant Cell* **4:** 1019-32

Fina J, Casadevall R, AbdElgawad H, Prinsen E, Markakis M, Beemster G, Casati P. 2017. UV-B Inhibits Leaf Growth through Changes in Growth Regulating Factors and Gibberellin Levels. *Plant Physiology* **174**(2)**:** 1110-1126

Gao F, Wang K, Liu Y, Chen Y, Chen P, Shi Z, Luo J, Jiang D, Fan F, Zhu Y , et al. 2015. Blocking miR396 increases rice yield by shaping inflorescence architecture. *Nature Plants* **2:** 15196

Gazquez A, Beemster GTS. 2017. What determines organ size differences between species? A meta-analysis of the cellular basis. *New Phytologist* **215**: 299– 308

Gonzalez N, De Bodt S, Sulpice R, Jikumaru Y, Chae E, Dhondt S, Van Daele T, De Milde L, Weigel D, Kamiya Y, Stitt M, Beemster GT, Inzé D. 2010. Increased leaf size: different means to an end. *Plant Physiology* **153:** 1261-79

Gonzalez N, Vanhaeren H, Inzé D. 2012. Leaf size control: complex coordination of cell division and expansion. *Trends in Plant Science* **6:** 332-40

Granier C, Tardieu F. 2009. Multi-scale phenotyping of leaf expansion in response to environmental changes: the whole is more than the sum of parts. *Plant, Cell and Environment* **32:** 1175- 1184

Huang X, Kurata N, Wei X, Wang ZX, Wang A, Zhao Q, Zhao Y, Liu K, Lu H, Li W, et al. **2012**. A map of rice genome variation reveals the origin of cultivated rice. *Nature* **490:** 497–501

Inze D, De Veylder L. 2006. Cell cycle regulation in plant development. *Annual Review of Genetics* **40:** 77–105

Jones R, Kaufman P. 1983. The role of gibberellins in plant cell elongation. *Critical Reviews in Plant Sciences* **1:** 23-47

Kalve S, de Vos D, Beemster GTS. 2014. Leaf development: a cellular perspective. *Frontiers in Plant Science* **5:** 362

Kant R, Sharma S, Dasgupta I. 2015. Virus-induced gene silencing (VIGS) for functional genomics in rice using Rice Tungro bacilliform virus (RTBV) as a vector. *Methods in Molecular Biology* **1287:**201–217

Kant R, Dasgupta I. 2017. Phenotyping of VIGS-mediated gene silencing in rice using a vector derived from a DNA virus. *Plant Cell Reports* **36**(7)**:** 1159-1170

Kierzkowski D, Runions A, Vuolo F, Strauss S, Lymbouridou R, Routier-Kierzkowska AL, Wilson-Sánchez D, Jenke H, Galinha C, Mosca G, Zhang Z, Canales C, Dello Ioio R, Huijser P, Smith RS, Tsiantis M. 2019. A Growth-Based Framework for Leaf Shape Development and Diversity. *Cell* **177**(6)**:** 1405-1418

Kim J, Choi D, Kende H. 2003. The AtGRF family of putative transcription factors is involved in leaf and cotyledon growth in Arabidopsis. *The Plant Journal* **36:** 94–104

van der Knaap E, Kim J, Kende H. 2000. A novel gibberellin-induced gene from rice and its potential regulatory role in stem growth. *Plant Physiology* **122**(3)**:** 695-704

Kuijt SJ, Greco R, Agalou A, Shao J, 't Hoen CC, Overnäs E, Osnato M, Curiale S, Meynard D, van Gulik R , et al. 2014. Interaction between the GROWTH-REGULATING FACTOR and KNOTTED1-LIKE HOMEOBOX families of transcription factors. *Plant Physiology* **164:** 1952–1966

Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* **9:** 559

Langmead B, Salzberg S. 2012. Fast gapped-read alignment with Bowtie 2. *Nature Methods* **9** (4)**:** 357-9

Lantzouni O, Alkofer A, Falter-Braun P, Schwechheimer C. 2020. GROWTH-REGULATING FACTORS Interact with DELLAs and Regulate Growth in Cold Stress. *The Plant Cell* **32**(4)**:** 1018-1034

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. 1000 Genome Project Data Processing Subgroup, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**(16)**:** 2078–2079

Li LC, Kang DM, Chen ZL, Qu LJ. 2007. Hormonal Regulation of Leaf Morphogenesis in *Arabidopsis*. *Journal of Integrative Plant Biology* **49:** 75-80

Li S, Tian Y, Wu K, Ye Y, Yu J, Zhang J, Liu Q, Hu M, Li H, Tong Y, Harberd N, Fu X. 2018. Modulating plant growth-metabolism coordination for sustainable agriculture. *Nature* **560**(7720)**:** 595-600

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**(4)**:** 402-8

Lu Y, Meng Y, Zeng J, Luo Y, Feng Z, Bian L, Gao S. 2020. Coordination between GROWTH-REGULATING FACTOR1 and GRF-INTERACTING FACTOR1 plays a key role in regulating leaf growth in rice. *BMC Plant Biology*. **20**(1)**:** 200

Luo AD, Liu L, Tang ZS, Bai XQ, Cao SY, Chu CC. 2005. Down-regulation of OsGRF1 gene in rice rhd1 mutant results in reduced heading date. *Journal of Integrative Plant Biology* **47:** 745–752

Mathan J, Singh A, Ranjan A. 2021a. Sucrose transport and metabolism control carbon partitioning between stem and grain in rice. *Journal of Experimental Botany* **72**(12)**:** 4355-4372

Mathan J, Singh A, Jathar V, Ranjan A. 2021b. High photosynthesis rate in two wild rice species is driven by leaf anatomy mediating high Rubisco activity and electron transport rate. *Journal of Experimental Botany* **72** (20): 7119–7135

Nelissen H, Eeckhout D, Demuynck K, Persiau G, Walton A, van Bel M, Vervoort M, Candaele J, De Block J, Aesaert S, et al. 2015. Dynamic changes in ANGUSTIFOLIA3 complex composition reveal a growth regulatory mechanism in the maize leaf. *The Plant Cell* **27:** 1605–1619

Nelissen H, Rymen B, Jikumaru Y, Demuynck K, Van Lijsebettens M, Kamiya Y, Inzé D, Beemster GTS. 2012. A local maximum in gibberellin levels regulates maize leaf growth by spatial control of cell division. *Current Biology* **22:** 1183–1187

Nelissen H, Sun X, Rymen B, Jikumaru Y, Kojima M, Takebayashi Y, Abbeloos R, Demuynck K, Storme V, Vuylsteke M, De Block J, Herman D, Coppens F, Maere S,

Kamiya Y, Sakakibara H, Beemster G and Inzé D. 2018. The reduction in maize leaf growth under mild drought affects the transition between cell division and cell expansion and cannot be restored by elevated gibberellic acid levels. *Plant Biotechnology Journal* **16:** 615-627

Omidbakhshfard MA, Proost S, Fujikura U, Mueller-Roeber B. 2015. Growth-Regulating Factors (GRFs): A Small Transcription Factor Family with Important Functions in Plant Biology. *Molecular Plant* **8:** 998-1010

Patel RK, Jain M. 2012. NGS QC Toolkit: A Toolkit for Quality Control of Next Generation Sequencing Data. *PLoS ONE* **7**(2)**:** e30619

Piazza P, Jasinski S, Tsiantis M. 2005. Evolution of leaf developmental mechanisms. *New Phytologist* **167:** 693–710

Robinson MD, Oshlack A. 2010. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biology* **11:** R25

Robinson M, McCarthy D, Smyth G. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**(1)**:** 139–140

Rodriguez RE, Debernardi JM, Palatnik JF. 2013. Morphogenesis of simple leaves: regulation of leaf size and shape. *Wiley Interdisciplinary Reviews: Developmental Biology* **3**(1)**:** 41-57

Sanchez PL, Wing RA, and Brar DS. 2013. The Wild Relative of Rice: *Genomes and Genomics*. In: Zhang Q and Wing RA (eds.) *Plant Genetics and Genomics*: Crops and Models 5 Genetics and Genomics of Rice, Springer, New York, 2–25

Sarvepalli K, Das Gupta M, Challa KR, Nath U. 2019. Molecular cartography of leaf development□-□role of transcription factors. *Current Opinion in Plant Biology* **47:** 22-31

Shan C, Mei ZL, Duan JL, Chen HY, Feng HF, Cai WM. 2014. OsGA2ox5, a Gibberellin Metabolism Enzyme, Is Involved in Plant Growth, the Root Gravity Response and Salt Stress. *Plos One* **9:** 1

Sprangers K, Avramova V, Beemster GT. 2016. Kinematic Analysis of Cell Division and Expansion: Quantifying the Cellular Basis of Growth and Sampling Developmental Zones in Zea mays Leaves. *Journal of Visualized Experiments* **118:** 54887

Su YH, Liu YB, Zhang XS. 2011. Auxin-cytokinin interaction regulates meristem development. *Molecular Plant* **4:** 616-25

Takatsuka H, Umeda M. 2014. Hormonal control of cell division and elongation along differentiation trajectories in roots. *Journal of Experimental Botany* **65:** 2633–2643

Tang Y, Liu H, Guo S, Wang B, Li Z, Chong K, Xu Y. **2018**. OsmiR396d affects gibberellin and brassinosteroid signaling to regulate plant architecture in rice. *Plant Physiology* **176:** 946–959

Tardieu F, Granier C. 2000. Quantitative analysis of cell division in leaves: methods, developmental patterns and effects of environmental conditions. *Plant Molecular Biology* **43:** 555-67

Tardieu F, Reymond M, Hamard P, Granier C, Muller B. 2000. Spatial distributions of expansion rate, cell division rate and cell size in maize leaves: a synthesis of the effects of soil water status, evaporative demand and temperature. *Journal of Experimental Botany* **51:** 1505-1514

Tian F, Yang DC, Meng YQ, Jin JP and Gao G. 2019. PlantRegMap: charting functional regulatory maps in plants. *Nucleic Acids Research* **48**(D1)**:** D1104-D1113

Townsley BT, Sinha NR. 2012. A new development: evolving concepts in leaf ontogeny. *Annual Review of Plant Biology* **63:** 535-62

Vercruysse J, Baekelandt A, Gonzalez N, Inzé D. 2020b. Molecular networks regulating cell division during Arabidopsis leaf growth. *Journal of Experimental Botany* **71**(8)**:** 2365-2378

Vercruysse J, van Bel M, Osuna-Cruz CM, Kulkarni SR, Storme V, Nelissen H, Gonzalez N, Inzé D, Vandepoele K. 2020a. Comparative transcriptomics enables the identification of functional orthologous genes involved in early leaf growth. *Plant Biotechnol Journal* **2:** 553-567

Wehrens R, Buydens L. 2007. Self- and super-organizing maps in R: the Kohonen package. *Journal of Statistical Software* **21:** 1–19

Willige B, Ghosh S, Nill C, Zourelidou M, Dohmann E, Maier A, Schwechheimer C. 2007. The DELLA domain of GA INSENSITIVE mediates the interaction with the GA INSENSITIVE DWARF1A gibberellin receptor of Arabidopsis. *The Plant Cell* **19**(4)**:** 1209–1220

Wu L, Zhang D, Xue M, Qian J, He Y and Wang S. 2014. Overexpression of the maize GRF10, an endogenous truncated GRF protein, leads to reduction in leaf size and plant height. *Journal of Integrative Plant Biology* **56:** 1053-1063

Zhang DF, Li B, Jia GQ, Zhang TF, Dai JR, Li JS, Wang SC. 2008. Isolation and characterization of genes encoding GRF transcription factors and GIF transcriptional coactivators in Maize (*Zea mays L.*). *Plant Science* **175:** 809–817

Figure Legends

Fig. 1 Wild rice *Oryza australiensis* grows longer leaf compared to the selected cultivated rice accessions. (a) Representative photographs of the selected cultivated and wild rice accessions showing leaf size variations. (b) Quantification of mature leaf length of the second leaf to the tenth leaf. (c) Representative fourth leaf images at the second day of emergence. (d) Quantification of fourth leaf length at the different days of emergence. (e) Leaf elongation rate of the selected rice accessions. Values in graphs represent mean±SD (n=15). Different letters indicate statistically significant differences according to one-way ANOVA followed by post-hoc Tukey HSD at $P < 0.05$.

Fig. 2 Transcriptome profiling reveals the key contribution of the cell cycle and Gibberellic Acid (GA) for differential leaf length across the selected rice accessions. (a) The number of differentially expressed transcripts and enriched GO terms in *O. australiensis* compared to *O. sativa* cv. IR64 and *O. sativa* cv. Nipponbare. * show enriched GO terms. (b) Gene expression clusters showing genotype-specific resolution for the selected four accessions derived from PCA-SOM analysis. Enriched GO terms are shown for *O. australiensis* and *O. glaberrima* specific clusters 1, 4, and 7. (c-e) Expression patterns of cell cycle-related genes (c), transcription factors and leaf developmental genes (d), and GA biosynthesis and signaling genes (e) present in clusters 1, 4 and 7 across the four accessions. (f) Gene regulatory network generated for the genes from the relevant clusters of (b). Three different modules of the network are shown in different colors.

AUS= *O. australiensis,* GLA= *O. glaberrima,* IR= *O. sativa* cv. IR64, NIP= *O. sativa* cv. Nipponbare.

Fig. 3 GA levels correlate with the leaf length variation across the cultivated and wild rice accessions. (a) GA_4 quantification in the fourth leaf of the cultivated and wild rice accessions on the second day of emergence. Data show mean ±SD (n=5). (b) Correlation between GA_4 and leaf length. (c) GA_4 and GA_7 quantification in the fourth leaf of *O. sativa* cv. Nipponbare and *O. australiensis* on the second day of emergence. Data show mean ±SD (n=5),***

P □< □ 0.001 by Student's t-test. (d, e) Effects of GA_3 (10µM) and PAC (1µM) treatments on leaf length of *O. sativa* cv. Nipponbare (d) and *O. australiensis* (e). Data show mean ±SD (n=15). Different letters indicate statistically significant differences according to one-way ANOVA followed by post-hoc Tukey HSD at $P < 0.05$.

Fig. 4 Leaf kinematics analyses confirm the GA-mediated control of cell division for rice leaf length regulation. (a, b) Different growth zones on the fourth leaf of *O. sativa* cv. Nipponbare (a) and *O. australiensis* (b) at the second day of emergence as quantified by epidermal cell length profiling and DAPI staining. Vertical lines of different shades represent the separation of different leaf-growth zones. (c, d) Different growth zones on the fourth leaf of *O. sativa* cv. Nipponbare (c) and *O. australiensis* (d) in response to GA_3 and PAC treatments at the second day of emergence as quantified by epidermal cell length profiling and DAPI staining. Vertical lines of different shades represent the size of division zones under different treatments. Values in graphs represent mean±SD (n=5).

Fig. 5 Zone-specific expression analysis of GA-related genes and GA quantification. (a - d) Expression pattern of GA biosynthesis genes *OsGA20OX2* and *OsGA3OX2* (a, b), GA catabolic gene *OsGA2OX4* (c), and GA-signalling repressor *OsSLR1* (d) in the two growth zones of fourth leaves of *O. sativa* cv. Nipponbare and *O. australiensis* on the second day of emergence. Data shown are mean ±SD of three biological replicates, *** P □< □ 0.001 by Student's t □ test. (e, f) Zone specific GA_4 (e) and GA_7 (f) quantification in the two growth zones of fourth leaves of *O. sativa* cv. Nipponbare and *O. australiensis* on the second day of emergence. Data shown are mean ±SD (n=5),*** P □< □ 0.001 by Student's t □ test. Zone I represents the region from leaf base to 0.8cm and zone II represents 0.8cm to 1.6cm from the leaf base.

Fig. 6 GA treatment affects zone-specific expression patterns of *GRF*s and cell cycle genes. (a - d) Expression pattern of genes encoding GRF transcription factors *OsGRF7* and *OsGRF8* (a, b), and cell cycle genes *OsCYCB1;4* and *OsCYCA3;2* (c, d) in the two growth zones of fourth leaves of *O. sativa* cv. Nipponbare and *O. australiensis* on the second day of emergence. (e - h) Effect of GA_3 treatment on the expression of *OsGRF7* (e), *OsGRF8* (f), *OsCYCB1;4* (g), and *OsCYCA3;2* (h) in the two growth zones of fourth leaves of *O. sativa* cv. Nipponbare. Data shown are mean ±SD of three biological replicates, *** P □< □ 0.001 by

Student's t‑test. Zone I represents the region from leaf base to 0.8cm and zone II represents 0.8cm to 1.6cm from the leaf base.

Fig. 7 Silencing of GA-related genes confirms the GA-mediated regulation of division zone size to control rice leaf length. (a) Shown are the leaf phenotypes of *OsGA20OX2, OsGRF7,* and *OsGRF8* silencing in *O. australiensis* along with the control plants. (b, c) Effect of *OsGA20OX2, OsGRF7,* and *OsGRF8* silencing on leaf length (b) and size of division zone (c) in *O. australiensis*. (d) Shown are the leaf phenotypes of *OsGA20OX2, OsGRF7,* and *OsGRF8* silencing in *O. sativa* cv. Nipponbare along with the control plants. (e, f) Effect of *OsGA20OX2, OsGRF7,* and *OsGRF8* silencing on leaf length (e) and size of division zone (f) in *O. sativa* cv. Nipponbare. Data represent mean ±SD of at least three biological replicates, *** $P < 0.001$ by Student's t‑test. White arrow indicates fourth leaf and scale bar = 2cm in (a) and (d).

Fig. 8 GA-OsGRF7/8 module is a key regulator of the rice leaf length via controlling the size of the division zone. GA-mediated induction in the expression domain of OsGRF7 and OsGRF8 transcription factors expands the region with cell division activity at the leaf base, thus promoting the longer division zone with a higher cell production rate leading to longer leaves in rice. Higher GA levels and downstream signaling resulting in increased cell division activity explained the longer leaves of the wild rice *O. australiensis* compared to the cultivated varieties.

Supporting Information

Fig. S1 Leaf length and width phenotyping of the four selected accessions.

Fig. S2 Differentially expressed transcripts and enriched GO terms in *O. glaberrima* and *O. australiensis*.

Fig. S3 PCA-SOM clustering of gene expression across the selected rice accessions.

Fig. S4 Quantification of mature fourth leaf length across the selected cultivated and wild rice accessions used for GA quantification.

Fig. S5 Effects of GA and paclobutrazol treatments on leaf length and division zone size.

Fig. S6 Cellular basis of leaf size differences across the selected rice accessions.

Fig. S7 Expression analysis of *GRFs* and cell cycle-related genes in different growth zones of cultivated and wild rice accessions.

Fig. S8 Expression analysis of GA- and cell cycle-related genes in response to GA and PAC treatment.

Fig. S9 Silencing of positive control *OsChlH* and GA signaling repressor *OsSLR1*.

Fig. S10 Expressions analysis of *GRFs* and *CYCLINs* in silencing lines of *OsGA20OX2 OsGRF7, OsGRF8,* and *OsSLR1*.

Table S1 List of cultivated and wild rice accessions used in the present study

Table S2 Quantification of maximum leaf width of fully-developed leaves of the seven cultivated and four wild rice accessions.

Table S3 The total number of differentially expressed genes along with genes expressed at higher and lower levels for each pair-wise comparison.

Table S4 Quantification of leaf kinematics parameters of the growing fourth leaf of *O. sativa* cv. Nipponbare under exogenous GA_3 and paclobutrazol treatments.

Table S5 Quantification of leaf kinematics parameters of the growing fourth leaf of *O. australiensis* under exogenous GA_3 and paclobutrazol treatments.

Table S6 List of primers used for gene expression analysis and transient silencing experiments.

Dataset S1 List of statistically significant differentially expressed genes for each pair-wise comparison.

Dataset S2 List of statistically significant GO-terms enriched for genes expressed at higher and lower levels in Fig. 2 and Supplemental Fig. S3.

Dataset S3 List of genes present in each cluster derived from PCA-SOM analysis.

Dataset S4 List of statistically significant GO-terms enriched for the genes present in the PCA-SOM clusters.

Dataset S5 List of cell cycle-related genes, transcription factors and leaf developmental genes, and phytohormone-related genes present in clusters 1, 4, and 7 that were used for generating heatmaps in Fig. 2c-e and Fig. S3c.

Dataset S6 List of genes present in the expression network shown in Fig. 2f with their network properties and module belongings.

Dataset S7 Normalized read counts used for generating gene coexpression network.

Fig. 1 Wild rice *Oryza australiensis* grows longer leaf compared to the selected cultivated rice accessions. (a) Representative photographs of the selected cultivated and wild rice accessions showing leaf size variations. (b) Quantification of mature leaf length of the second leaf to the tenth leaf. (c) Representative fourth leaf images at the second day of emergence. (d) Quantification of fourth leaf length at the different days of emergence. (e) Leaf elongation rate of the selected rice accessions. Values in graphs represent mean±SD (n=15). Different letters indicate statistically significant differences according to one-way ANOVA followed by post-hoc Tukey HSD at P < 0.05.

(a)

Genes expressed at higher levels in *O. australiensis*

O. sativa cv. IR64   *O. sativa* cv. Nipponbare

*Enriched GO Terms
Cell cycle, Cell growth, Development process, Cell morphogenesis

1152   2384 *   1633

Genes expressed at lower levels in *O. australiensis*

O. sativa cv. IR64   *O. sativa* cv. Nipponbare

*Enriched GO Terms
Photosynthesis, Biosynthetic process, Signalling

1657   3870 *   1895

(b)

O. glaberrima (Cluster1)

*Enriched GO Terms
Cell cycle, Development process, Response to auxin

Cluster_1   Cluster_2   Cluster_3

O. australiensis + *O. glaberrima* (Cluster 4)

*Enriched GO Terms
Cell cycle, Development process, Response to gibberellin, Response to auxin

Cluster_4   Cluster_5   Cluster_6

O. australiensis (Cluster 7)

*Enriched GO Terms
Cell cycle, Development process, Developmental growth, Cell morphogenesis, Response to cytokinin

Cluster_7   Cluster_8   Cluster_9

(c) Cell-cycle-related genes

CYCA2, CYCD6, CYCB2, CYCB1

CDK D1, ROOT HAIRLESS, WEE1, AML1, EXPANSIN10

CYCB1, CYCD4, CYCB2, RB1, E2F2, CDT1A, EXPANSIN4

(d) Transcription factors and leaf developmental genes

YABBY6, SHORT-ROOT, NAM-family, KNOX1, GRF10

YABBY-family, MYB-family, ZF-HD-family, GATA-family, GRF7, GRF8, PINHEAD, HOTHEAD, LEAFBLADELESS1

AP2-family, bZIP-family, TCP6, MYB-family, CLAVATA1, GRF6, AINTEGUMENTA

(e) Gibberellic acid biosynthesis and signalling genes

GID1

CPS, KAO, SPY, GID1

WRKY51, MYB-family, CYP450

(f)

GID1 L2 — GA signalling
ent-kaurene synthase (CPS) — GA biosynthesis
Mitotic checkpoint protein — Cell cycle regulation
CDK D1 — Cell cycle regulation
MEL1 ARGONAUTE — progression of premeiotic mitosis and meiosis during sporogenesis
Cyclin — Cell cycle regulation
AML1 — Meiosis and vegetative growth
AINTEGUMENTA — Leaf growth and development
Growth regulatory factor 7 — Cell proliferation related TF
Leafbladeless1 — Leaf development

Fig. 2 Transcriptome profiling reveals the key contribution of the cell cycle and Gibberellic Acid (GA) for differential leaf length across the selected rice accessions. (a) The number of differentially expressed transcripts and enriched GO terms in *O. australiensis* compared to *O. sativa* cv. IR64 and *O. sativa* cv. Nipponbare. * show enriched GO terms. (b) Gene expression clusters showing genotype-specific resolution for the selected four accessions derived from PCA-SOM analysis. Enriched GO terms are shown for *O. australiensis* and *O. glaberrima* specific clusters 1, 4, and 7. (c-e) Expression patterns of cell cycle-related genes (c), transcription factors and leaf developmental genes (d), and GA biosynthesis and signaling genes (e) present in clusters 1, 4 and 7 across the four accessions. (f) Gene regulatory network generated for the genes from the relevant clusters of (b). Three different modules of the network are shown in different colors.

AUS= *O. australiensis,* GLA= *O. glaberrima,* IR= *O. sativa* cv. IR64, NIP= *O. sativa cv.* Nipponbare.

Fig. 3 GA levels correlate with the leaf length variation across the cultivated and wild rice accessions. (a) GA$_4$ quantification in the fourth leaf of the cultivated and wild rice accessions on the second day of emergence. Data show mean $\pm$SD (n=5). (b) Correlation between GA$_4$ and leaf length. (c) GA$_4$ and GA$_7$ quantification in the fourth leaf of *O. sativa* cv. Nipponbare and *O. australiensis* on the second day of emergence. Data show mean $\pm$SD (n=5), *** $P < 0.001$ by Student's t-test. (d, e) Effects of GA$_3$ (10μM) and PAC (1μM) treatments on leaf length of *O. sativa* cv. Nipponbare (d) and *O. australiensis* (e). Data show mean $\pm$SD (n=15). Different letters indicate statistically significant differences according to one-way ANOVA followed by post-hoc Tukey HSD at $P < 0.05$.

Fig. 4 Leaf kinematics analyses confirm the GA-mediated control of cell division for rice leaf length regulation. (a, b) Different growth zones on the fourth leaf of *O. sativa* cv. Nipponbare (a) and *O. australiensis* (b) at the second day of emergence as quantified by epidermal cell length profiling and DAPI staining. Vertical lines of different shades represent the separation of different leaf-growth zones. (c, d) Different growth zones on the fourth leaf of *O. sativa* cv. Nipponbare (c) and *O. australiensis* (d) in response to GA_3 and PAC treatments at the second day of emergence as quantified by epidermal cell length profiling and DAPI staining. Vertical lines of different shades represent the size of division zones under different treatments. Values in graphs represent mean±SD (n=5).

Fig. 5 Zone-specific expression analysis of GA-related genes and GA quantification. (a - d) Expression pattern of GA biosynthesis genes *OsGA20OX2* and *OsGA3OX2* (a, b), GA catabolic gene *OsGA2OX4* (c), and GA-signalling repressor *OsSLR1* (d) in the two growth zones of fourth leaves of *O. sativa* cv. Nipponbare and *O. australiensis* on the second day of emergence. Data shown are mean ±SD of three biological replicates, *** $P < 0.001$ by Student's t-test. (e, f) Zone specific GA_4 (e) and GA_7 (f) quantification in the two growth zones of fourth leaves of *O. sativa* cv. Nipponbare and *O. australiensis* on the second day of emergence. Data shown are mean ±SD (n=5),*** $P < 0.001$ by Student's t-test. Zone I represents the region from leaf base to 0.8cm and zone II represents 0.8cm to 1.6cm from the leaf base.

Fig. 6 GA treatment affects zone-specific expression pattern of *GRF*s and cell cycle genes. (a - d) Expression pattern of genes encoding GRF transcription factors *OsGRF7* and *OsGRF8* (a, b), and cell cycle genes *OsCYCB1;4* and *OsCYCA3;2* (c, d) in the two growth zones of fourth leaves of *O. sativa* cv. Nipponbare and *O. australiensis* on the second day of emergence. (e - h) Effect of GA_3 treatment on the expression of *OsGRF7* (e), *OsGRF8* (f), *OsCYCB1;4* (g), and *OsCYCA3;2* (h) in the two growth zones of fourth leaves of *O. sativa* cv. Nipponbare. Data shown are mean ±SD of three biological replicates, *** P < 0.001 by Student's t-test. Zone I represents the region from leaf base to 0.8cm and zone II represents 0.8cm to 1.6cm from the leaf base.

Fig. 7 Silencing of GA-related genes confirms the GA-mediated regulation of division zone size to control rice leaf length. (a) Shown are the leaf phenotypes of *OsGA20OX2, OsGRF7,* and *OsGRF8* silencing in *O. australiensis* along with the control plants. (b, c) Effect of *OsGA20OX2, OsGRF7,* and *OsGRF8* silencing on leaf length (b) and size of division zone (c) in *O. australiensis*. (d) Shown are the leaf phenotypes of *OsGA20OX2, OsGRF7,* and *OsGRF8* silencing in *O. sativa* cv. Nipponbare along with the control plants. (e, f) Effect of *OsGA20OX2, OsGRF7,* and *OsGRF8* silencing on leaf length (e) and size of division zone (f) in *O. sativa* cv. Nipponbare. Data represent mean ±SD of at least three biological replicates, *** $P < 0.001$ by Student's t-test. White arrow indicates fourth leaf and scale bar = 2cm in (a) and (d).

Fig. 8 GA-OsGRF7/8 module is a key regulator of the rice leaf length via controlling the size of the division zone. GA-mediated induction in the expression domain of OsGRF7 and OsGRF8 transcription factors expands the region with cell division activity at the leaf base, thus promoting the longer division zone with a higher cell production rate leading to longer leaves in rice. Higher GA levels and downstream signaling resulting in increased cell division activity explained the longer leaves of the wild rice *O. australiensis* compared to the cultivated varieties.