1

Reconstruction of dynamic regulatory networks reveals signaling-induced topology

changes associated with germ layer specification

Emily Y. Su[1,2*], Abby Spangler[1*], Qin Bian[1,2], Jessica Y. Kasamoto[2], Patrick Cahan[1,2,3]

[1]Institute for Cell Engineering, Johns Hopkins School of Medicine, Baltimore MD 21205 USA

[2]Department of Biomedical Engineering, Johns Hopkins School of Medicine, Baltimore MD

21205 USA

[3]Department of Molecular Biology and Genetics, Johns Hopkins School of Medicine, Baltimore

MD 21205 USA

* These authors made equal contributions.
15

17 Correspondence to: patrick.cahan@jhmi.edu

19 Declarations of interest: none

24 **Keywords**: Single cell RNA-seq, Gene regulatory network, Network inference, Directed

25 differentiation, Gastrulation, Embryoid body

26
27

Abstract

Elucidating regulatory relationships between transcription factors (TFs) and target genes is fundamental to understanding how cells control their identity and behavior. Computational gene regulatory network (GRN) reconstruction methods aim to map this control by inferring relationships from transcriptomic data. Unfortunately, existing methods are imprecise, may be computationally burdensome, and do not illustrate how networks transition from one topology to another. Here we present Epoch, a computational network reconstruction tool that leverages single cell transcriptomics to infer dynamic network structures. Epoch performs favorably when benchmarked on reconstruction of synthetically generated, *in vivo*, and *in vitro* data. To illustrate the usefulness of Epoch, we used it to identify the dynamic networks underpinning directed differentiation of mouse ESC guided by multiple primitive streak induction treatments. Our analysis demonstrates that modulating signaling pathways drives topological network changes that shape cell fate potential. We also find that Peg3 is a central contributor to the rewiring of the pluripotency network to favor mesoderm specification. By integrating signaling pathways with GRN structures, we traced how Wnt activation and PI3K suppression govern mesoderm and endoderm specification, respectively. The methods presented here are available in the R package Epoch, and provide a foundation for future work in understanding the biological implications of dynamic regulatory structures.

Introduction

Gene regulatory networks (GRNs) model the regulatory relationship between a set of regulators, or transcription factors (TFs), and their target genes. The topology of these networks, defined by edges that map the total existing regulatory interactions between TFs and targets, offers a molecular-level view of a controlled system in which genes work together as part of a framework to accomplish specific cell functions. Uncovering topology within these GRNs is fundamental in answering a number of questions including: understanding how cellular identity is maintained and established, elucidating mechanisms of disease caused by dysfunctional gene regulation, and finding novel drug targets among others. In the context of cell fate engineering, accurate mapping of such GRNs would thus enable the identification of TFs required to activate expression of target genes of interest such that we could more accurately control cell fate transitions or cell behavior. Unfortunately, how best to map these relationships remains both an experimental and computational challenge.

Experimental approaches, including ChIP-ChIP and ChIP-seq, have provided valuable insights into identifying specific regulatory relationships. However, experimental methods are generally limited in scope and scalability. For this reason, computational methods to efficiently infer GRN structures are needed. Existing methods have leveraged advances in data collection and machine learning approaches that have allowed for the statistical inference of GRNs via gene expression data. These include tools that rely on information theoretic measures (Faith et al., 2007; Margolin et al., 2006; Meyer et al., 2007), ensemble methods (Huynh-Thu et al., 2010), Bayesian approaches (Yu et al., 2004; Hartemink, 2005), or differential equation-based (ODE) approaches (di Bernardo et al., 2005). Unfortunately, computational methods developed to date suffer from several drawbacks including low precision and low sensitivity, often stemming from bloated false positives due to indirect interactions and a high degree of homogeneity from bulk RNA-sequencing data that does not offer enough perturbation for true relationships to be detected. These limitations ultimately result in imprecise networks (Marbach et al., 2012).

With the advent of droplet-based systems for single-cell RNA-sequencing (Zheng et al., 2017), and the corresponding rapidly growing body of single-cell transcriptomic data, computational analysis techniques have seen widespread advancement in harnessing the single-cell resolution to identify cell types and states (Villani et al., 2017) and to trace developmental trajectories (Treutlein et al., 2014). Of particular interest, trajectory inference (TI) methods, or pseudotemporal ordering methods, such as the transcriptomic-based methods Monocle (Trapnell et al., 2014), Slingshot (Street et al., 2018), and DPT (Haghverdi et al.,

2016), as well as cytometry-based methods such as SPADE (Qiu et al., 2011), have allowed for

computational modelling of dynamic processes like cell differentiation. Such methods order cells

in an unbiased fashion along a trajectory following a linear, bifurcating, or more complex graph

structure. In doing so, TI methods have proved valuable in the identification of novel cell

subpopulations (Chen et al., 2016) and the elucidation of differentiation trees (Velten et al.,

2017).

Unsurprisingly, several recently published tools have aimed to take advantage of the

resolution offered by single-cell transcriptomics to infer GRN structures (Aibar et al., 2017;

Matsumoto et al., 2017; Qiu et al., 2020), and can be categorized similarly to their bulk

counterparts into information theoretic, ensemble, Bayesian, and ODE approaches.

Unfortunately, they also suffer from similar limitations, and aside from low precision and

sensitivity, many of these methods are computationally burdensome, limiting users' ability to

hone analysis through iterative application. Many of these methods, though they employ

pseudotemporal analysis to aid in reconstruction, are limited in their ability to explain how

network topology evolves over time. Further, it has been shown that the dynamic and

noncommutative nature of such complex networks (e.g. via altering chromatin accessibility),

permits independent control of targets (or genetic programs) that would otherwise be

simultaneously activated with non-sequential, or combinatorial, logic (Letsou and Cai, 2016).

Thus, it's necessary for GRN models to encapsulate this dynamic behavior in order to gain a

more nuanced view of how a changing GRN topology generates distinct cell fates in

development or disease progression.

For those reasons, we developed a computational GRN reconstruction tool called

Epoch, which leverages single-cell transcriptomics to efficiently reconstruct dynamic network

structures. Three key strategies exist at the center of Epoch: first, reconstruction is limited to

dynamically expressed genes. Second, Epoch employs an optional "cross-weighting" strategy to

reduce false positive interactions. Third, Epoch divides pseudotime into "epochs", or time

periods, and extracts a dynamic network and predicted top regulators influential in driving

topology changes. Finally, Epoch includes a number of functionalities to aid in network analysis

and comparison, including integration of the reconstructed GRN with major signaling pathways,

and the subsequent tracing through the GRN of shortest paths from signaling effectors toward

activation or repression of target genes.

To demonstrate the utility of our tool we benchmarked Epoch's performance against

commonly used computational GRN reconstruction tools on synthetically generated data, *in vivo*

mouse muscle development data, and *in vitro* directed differentiation data. We demonstrate that

Epoch performs well in precision-recall analysis and in runtime comparison. We further highlight

how Epoch's stepwise-structured pipeline allows for user flexibility in adapting it to other GRN

reconstruction methods.

Finally, we applied Epoch to answer open questions in gene regulation and directed

differentiation. In particular, as pluripotent cells differentiate and undergo gastrulation, their

transcriptomic profiles diverge, ultimately specifying and committing to distinct germ layers.

Pluripotent cells undergoing *in vitro* differentiation often begin by undergoing embryoid body

(EB) formation, which mimics *in vivo* gastrulation, and can be guided by the introduction of

specific molecules to induce signaling pathway activity. However, it is unclear if the activation of

these pathways influences cell fate by simply activating distinct targets, or by impacting GRN

topology, thus altering the fate decision landscape and changing the capacity for, and

dependence of, regulatory control over differing target states.

To this end, we performed scRNA-seq on day 0 through day 4 *in vitro* mouse embryonic

stem cell (mESC) directed differentiation guided by four separate primitive streak induction

treatments, following a protocol similar to one previously established (Spangler et al., 2018). We

utilized Epoch reconstructed networks to quantify the extent of target gene differences amongst

active TFs in all four treatments. We found that there were remarkably large differences

between target genes of the same TF across treatments, and further applied gene set

enrichment analysis to understand the function consequences of these differences. Overall, our

results suggest that these topological differences may broadly alter cell fate potential.

We next applied Epoch to more specifically evaluate cell fate potential differences seen

in the data, and focused on understanding why cells treated with different induction treatments

exhibited differing mesodermal fate potential, the lineage that was overall least likely chosen by

differentiating cells. Our analysis revealed a number of topological features affecting

mesodermal fate potential, including a central role for Peg3 in regulating the exit of pluripotency

and transition to mesodermal fate. Additionally, we used Epoch to integrate signal transduction

pathways with reconstructed network topologies to demonstrate that cells treated with Wnt3a,

ActivinA, and GSK inhibitor developed a network topology that allows for more effective

mesodermal specification through activation of Wnt signaling. This was in direct contrast to

other treatments in which network topologies more easily specified endodermal fate through

suppression of PI3K signaling and subsequent activation of Foxo1. We believe that these

results and similar analyses can be used to hone more efficient differentiation of mesodermal-

fated cells. Further, we expect that analogous analyses can be performed to explain fate

decisions in other dynamic biological processes.

Results

Epoch workflow relies on single-cell analysis techniques to infer dynamic network structures

Epoch takes as minimum input processed single-cell transcriptomic data and

pseudotime or equivalent annotation from any trajectory inference method a user chooses to

utilize (**Figure 1**). From here, Epoch employs its first step: limiting reconstruction to dynamically

expressed genes in an effort to both focus on genes playing an active role in cell state changes,

and to ensure that identified interactions are ones that can be inferred from perturbations in

transcriptomic data. To accomplish this, Epoch employs a method reminiscent of Slingshot

(Street et al., 2018), in which individual genes are modeled across pseudotime using a

generalized additive model. Upon filtering the data for dynamically expressed genes, Epoch

reconstructs an initial static network via a CLR-like (Context Likelihood of Relatedness) method,

employing either Pearson correlation or mutual information to infer interactions (Faith et al.,

2007).

In the second step, which we have termed "cross-weighting", Epoch employs an optional

network refinement step that relies on a cross-correlation-based weighting scheme. This helps

significantly reduce false positives that may result from indirect interactions or that may

represent non-logical interactions, ultimately improving precision. In this step, the expression

profiles over pseudotime for each TF-target pair is aligned and progressively shifted to

determine an average estimated offset value at which maximum correlation is achieved

between the two profiles. A graded-decline weighting factor is computed based on this offset,

and used to negatively weight interactions that are unlikely to be true positives.

Once a static network has been inferred, Epoch extracts a dynamic network from the

static network, and further identifies the top regulators driving changes in network topology and

thus cell state. Specifically, this begins with Epoch breaking down pseudotime into "epochs", or

essentially "time periods", based on pseudotime, cell ordering, k-means clustering, or user-

defined assignment. From here, genes are assigned to specific epochs based on their activity

along pseudotime. Epoch then fractures the static network into a dynamic one, composed of

"epoch networks", representing active interactions within a particular epoch, and possible

"transition networks", describing how an epoch network transitions into a subsequent epoch

network.

As Epoch was initially built with the improvement of cell fate engineering protocols in

mind, we included in the Epoch framework a number of network analysis and network

comparison capabilities. For example, Epoch will identify influential transcription factors within a

given static or dynamic network by extracting top regulators that appear to have the most

influence in driving or maintaining topology. Specifically, Epoch will compute and rank TFs

according to their PageRank in the context of their epoch networks, and will further look for TFs

that simultaneously score high in betweenness and degree centralities compared to all other

TFs. Additionally, Epoch can integrate major signal transduction pathways with reconstructed

networks and determine the shortest paths through the networks to functional groups of

specified target genes. This allows users to identify and trace topologies capable of activating or

repressing specific groups of genes.

Finally, Epoch is designed in a stepwise workflow such that it can be broken into

individual steps and flexibly merged and used with any trajectory inference, network

reconstruction, and even network refinement tools that currently exist or that may be developed

in the future.

Epoch performs favorably benchmarked on synthetically generated data

To assess the performance of Epoch in generating static networks, we began by

benchmarking its performance reconstructing networks from synthetically generated data. To

this end, we generated fifteen synthetic datasets using the Dyngen R package (Cannoodt et al.,

2020). Networks were designed to encompass varying regulatory motifs and spanned 100-420

genes including 20-70 transcription factors. Synthetic single-cell datasets generated from these

networks varied in size, up to 1000 cells. We benchmarked Epoch against original versions of

GENIE3, which is the GRN reconstruction engine of SCENIC (Aibar et al., 2017), and CLR, as

well as variations on these methods that incorporated them within the Epoch framework. Thus,

we assessed a total of 11 methods: CLR using mutual information (original CLR), CLR using

Pearson correlation, CLR using mutual information with cross-weighting, CLR using Pearson

correlation with cross-weighting, GENIE3 (original GENIE3), GENIE3 limited to dynamically

expressed genes, GENIE3 limited to dynamically expressed genes with cross-weighting, Epoch

with Pearson correlation (with and without cross-weighting), Epoch with mutual information (with

and without cross-weighting).

We reconstructed individual networks across all methods for each dataset, using

normalized simulation time as the pseudotemporal annotation for Epoch. To assess

performance, we compared each resulting network to a set of five random networks, unique to

each reconstructed network, that were generated by permuting edge weights of targets for each

transcription factor in the reconstructed network (**Figure 2a**). As a gold standard, all networks

were compared against the original network used to generate the synthetic single-cell data.

Based on fold improvement in area under the precision-recall curve over the random networks,

all variations of Epoch outperformed original versions of CLR and GENIE3 (**Figure 2b,**

Supplementary Figure 1a). Additionally, we found that Epoch using mutual information with or

without cross-weighting resulted in higher fold improvement over random compared to any other

method. In all cases it was apparent that key steps in the Epoch framework improved overall

static reconstruction. Specifically, limiting reconstruction to dynamically expressed genes and

applying the cross-weighting scheme both led to increased fold improvement in AUPR over

random, demonstrating the utility of Epoch's framework.

Epoch performs favorably benchmarked on real world data

While the results from assessment of networks reconstructed from synthetic data

demonstrated Epoch's favorable performance as a network inference tool, we acknowledge that

there are caveats associated with synthetic data, including limited stochasticity and reliance on

simulations that may not encompass the full extent of potential cell state responses seen in real

biological systems. Thus, we next turned our attention to benchmarking on real world data. To

assess Epoch's performance, we began by assessing the networks reconstructed by the same

11 methods listed above from data of E12.5 mouse muscle development (**Figure 2c**). For this

dataset, we used DPT to compute pseudotime for Epoch reconstruction. We used Chip-X data

from the Enrichr database (which includes ChIP-Seq, ChIP-ChIP, ChIP-PET, DamID used to

profile transcription factor binding to DNA, covering 220 TFs and >35,000 genes) as a gold

standard network (Lachmann et al., 2010). Based on precision recall, we found that all

variations of Epoch once again outperformed original versions of CLR and GENIE3.

Interestingly, we found that the two variations of GENIE3 embedded within the Epoch

framework performed best overall in this assessment, followed by the two variations of Epoch

using mutual information (with and without cross-weighting). It was again apparent that limiting

reconstruction to dynamically expressed genes and cross-weighting significantly improved

overall reconstruction (**Figure 2d, Supplementary Figure 1b**).

To determine if these results were consistent across other real world datasets, we turned

to single-cell transcriptomic data collected from day 4 in vitro mESC directed differentiation

(Spangler et al., 2018). We limited our reconstruction to cells in earlier stages of differentiation

that had not yet transitioned into any committed germ layer state, as determined by

pseudotemporal analysis, RNA velocity analysis, expression of Zfp42, marking naive pluripotent

cells, and expression of Fgf5, marking cells in a primed state (**Supplementary Figure 1c**). The

filtering ensured the trajectory remained linear, which, while absolutely not a requirement for

Epoch reconstruction, was employed to minimize performance variability of trajectory inference

methods in which branched structures may be more difficult to order. This allowed us to narrow

our focus on reconstruction performance while minimizing impact from potentially inaccurate

pseudotemporal ordering. Based on precision-recall, we found that Epoch variations using

crossweighting and GENIE3 embedded within the Epoch framework performed best

(**Supplementary Figure 1d**). Overall, these benchmarking results demonstrate Epoch's utility

as a reconstruction tool, and further demonstrate the usefulness of Epoch as a flexible

framework that can improve reconstruction of other methods.

Epoch elucidates regulatory networks driving lineage specification in early in vitro mouse ESC

differentiation

We next wanted to apply Epoch to elucidate the underlying dynamic networks involved

in mesodermal fate specification in mESC directed differentiation. We collected scRNA-seq data

from day 0 through day 4 *in vitro* mESC directed differentiation optimized to induce primitive

streak formation (**Figure 3a**). Briefly, cells were allowed to differentiate in serum-free

differentiation media before being treated with one of four treatments on day 2: Wnt3a and

Activin A alone (WA), or with one of Bmp4 (WAB), GSK inhibitor (WAG), or Noggin (WAN).

Multiple rounds of differentiation were staggered such that we could harvest samples

representative of d0-d4 at one time for sequencing using the MULTI-seq protocol (McGinnis et

al., 2019). After barcode classification, samples were preprocessed using the SCANPY pipeline

(Wolf et al., 2018), and RNA velocity analysis was performed (Bergen et al., 2019; La Manno et

al., 2018) (**Figure 3b,c**). We found the presence of three distinct lineages, which we confirmed

to be representative of neuroectoderm, mesoderm, and endoderm based on gene expression

and further verified by singleCellNet analysis (Tan and Cahan, 2019) (**Figure 3d,**

Supplementary Figure 2a-c, 3a). A majority of the differentiating cells transitioned toward the

neuroectoderm (clusters labeled 0,2, 0,3, 4, 2, 5, 9) fate with smaller populations transitioning

toward mesoderm (cluster labeled 8) and endoderm fate (clusters labeled 7,1 and 7,0).

Interestingly, there were differences between treatments in terms of the distribution of
cells amongst the different populations (**Figure 3e,f, Supplementary Figure 3b**). Notably, cells
treated with WAG exhibited a stunted trajectory toward neuroectodermal fate but a fuller
trajectory toward mesodermal fate. This was in direct contrast to cells treated with the remaining
three treatments (WAB, WAN, WA), in which very few cells differentiated toward mesodermal
fate, but instead exhibited strong neuroectodermal or endodermal commitment. Further, while
WAB and WA cells tended to transition into cluster 7,0 (furthest along endodermal development
according to RNA velocity analysis), WAN and WAG cells were more likely to stay in cluster 7,1.
We then applied Epoch to this data, implementing latent time (Bergen et al., 2020) in
place of pseudotime, and analyzed the networks underlying each of the three lineages. Latent
time was divided into three epochs, and a dynamic network was extracted for each lineage
(**Figure 4a-c, Supplementary Figure 4**). Both top regulator prediction methods, PageRank and
Betweenness-Degree, had similar results (**Figure 4d, Supplementary Figure 4d**). In a more
nuanced analysis, we limited our scope to early latent time, corresponding to the inner cell mass
(ICM) to epiblast transition in an *in vivo* context, and bootstrapped the network reconstruction
(**Supplementary Figure 5**). In doing so, we were able to predict top regulators via a consensus
approach in which regulators were ranked by their frequency as a top regulator (PageRank) or
average importance (Betweenness-Degree) in the bootstrapped networks. The top regulators
recovered by both PageRank and Betweenness-Degree were largely overlapping, and included
TFs such as Zfp42, Klf2, Sox11, Pou3f1, and Klf4.
318
319   *Peg3 is a central regulator in mesodermal WAG networks and may drive differences in gene*
320   *module expression amongst treatments*
321
As noted above, cell fate potential amongst the differentiating cells appeared to be
dependent on treatment. We hypothesized that underlying these differences were treatment-
specific GRNs. To this end, we reconstructed and compared dynamic networks for each
treatment along the mesodermal path (which we refer to as the treatment-specific networks:
WAG, WAB, WAN, and WA networks), and further compared these against the full
reconstructed mesoderm network described in the previous section (which is reconstructed from
all cells regardless of treatment, and which we refer to as the mesoderm network). We then
aimed to understand why WAG treated cells were able to specify mesodermal fate while WAB,
WAN, and WA treated cells did not.

Because differences in state were largely apparent toward the end of latent time, we
performed community detection on epoch 2, epoch 3, and their transition, of the mesoderm
network and identified distinct TF communities, or modules, within the topology (**Figure 5a,**
Supplementary Figure 7). We assessed the activity of each module by looking at the average
expression of member genes (excluding genes predicted to be repressed in a given module)
across latent time for each treatment (**Figure 5b Supplementary Figure 8a**). Importantly, we
found the existence of certain modules that were strongly activated in the WAG treatment that
failed to activate or were weakly activated in the remaining three treatments. Three such
modules (one activated in each of the second epoch, third epoch, and transition) were
particularly interesting as they exhibited strong activity in the WAG treatment along not only the
mesoderm path, but also the full dataset. The TFs in these modules were Peg3, Lhx1, Hoxb2,
Foxc1, Tshz1, Meis2, Mesp1, Tbx6, Foxc2, Prrx2, Meox1, and Notch1. Upon extracting the
differential network (which we define as the network containing the interactions, or edges, that
have large weight differences between two treatments, and thus represent interactions most
likely to be unique to a treatment network; see methods) between WAG and WA, we recovered
the same modules and further identified the top differential regulators of these TFs themselves
(**Figure 5c, Supplementary Figure 8c**). Interestingly Peg3 was predicted to have a role in
activating many of these TFs in the WAG treatment networks, but had a notably absent role in
the WA treatment.
Peg3's apparent influence in the WAG networks suggested to us the potential existence
of a pathway linking GSK3 inhibition, Peg3 expression, and mesodermal gene expression.
Indeed, previous reports have demonstrated a role for GSK3 in maintaining DNA methylation at
imprinted loci, including Peg3, in mESCs (Meredith et al., 2015). Additional literature has
demonstrated that methylation at imprinted loci may be maintained by Dnmt3a2, which has
been suggested to be regulated by the GSK3-dependent transcription factor N-Myc (Popkie et
al., 2010). Unfortunately, our data was only able to capture Dnmt3a dynamics (representing
composite Dnmt3a and Dnmt3a2 expression), and not specifically Dnmt3a2, which is
transcribed from an alternative promoter within the Dnmt3a locus. However, since Dnmt3a2 is
the isoform predominant in ESCs (Chen et al., 2002), we roughly compared Dnmt3a temporal
dynamics and Peg3 expression between the different treatments (**Supplementary Figure 8b**).
Notably, Dnmt3a and Peg3 expression were largely mutually exclusive, and Dnmt3a expression
was low to none in the WAG treated cells. We then asked if this was a phenomenon observed in
in vivo gastrulation. In the single-cell mouse gastrulation dataset from Chan et al. (Chan et al.,
2019), we found relatively high expression of Dnmt3a in epiblast cells which was tapered in

primitive streak and mesodermal cells (**Figure 5d**). In contrast, Peg3 expression was low in

epiblast and increased in the primitive streak, mirroring our *in vitro* data.

Target gene differences amongst TFs confirm network topology restructuring dependent on

treatment

Given Peg3's apparent role in restructuring GRN topology, we next aimed to understand

the extent to which broad, network-wide topology changes are responsible for differences in cell

fate potential between treatments. To answer this, we performed two edge-level analyses

focusing on differences in targets of transcription factors across the four treatment networks.

First, we asked if transcription factors actively expressed in all four treatments regulated

the same set of target genes. To this end, we focused on the treatment-specific networks along

the mesodermal path. We narrowed our analysis to the 72 TFs that were active in all four

treatments during day 3 or day 4 along the mesodermal path. We quantified overlap between

targets of each TF in a pairwise comparison of the treatments using the Jaccard similarity

(**Figure 6a**). As a baseline, we implemented a bootstrapping method in which we reconstructed

10 networks for each treatment along the mesodermal path (using 400 sampled cells each); for

each TF we calculated the average Jaccard similarity of its targets amongst pairwise

comparisons of the reconstructed networks in a treatment. This gave us a baseline, or

expected, Jaccard similarity for each TF in each treatment. Overall, we found that

overwhelmingly, target differences between treatments were significantly greater than the

baselines, indicating vast network topology differences amongst the treatment networks.

Next, we asked if these differences in predicted targets had functional roles in impacting

fate potential. To this end, we focused on the 13 TFs amongst the 72 mentioned above that

exhibited the largest differences in predicted targets between treatments (**Figure 6b**). We

performed gene set enrichment analysis on the predicted targets of these TFs, and compared

the results between treatments (**Figure 6c**). Interestingly, we found enrichment for heart

development and mesoderm formation in only the WAG treatment, despite the fact that the

networks we analyzed were reconstructed along the mesodermal path. Of note, all treatments

showed enrichment for embryonic digestive tract morphogenesis. Our enrichment results thus

supported our hypothesis that modulating signaling results in a topological restructuring of the

GRN, ultimately guiding cell fate potential.

Restrictive network topologies impede paths toward mesoderm gene activation and limit
mesodermal fate potential

We next asked if we could trace pathways from signaling effectors to the activation of meseodermal fate. To determine which signal transduction pathways were active at any given point in latent time along the mesodermal lineage, we used ChIP-seq data from ChIP-Atlas (Oki et al., 2018) to compile a list of binding targets for 18 signaling effector TFs (available as part of the Epoch framework). We then computed average expression of these targets across latent time broken down by treatment, and unsurprisingly found differences in the signaling pathways that were activated in each treatment (**Figure 7a**). Of note, Notch signaling was activated in WAB, WAN, and WA treatments, but not in the WAG treatment along the mesodermal lineage (and weakly activated along the neuroectodermal lineage), consistent with literature demonstrating that Notch signaling plays a role in specifying neuroectodermal fate in the neuroectoderm-mesendoderm fate decision (Androutsellis-Theotokis et al., 2006; Aubert et al., 2002).

Because Wnt signaling is essential for mesodermal fate, we looked for paths connecting Wnt effectors to the TFs in the WAG-specific modules from the previous analysis. Specifically, we computed the shortest paths from targets of these effectors to the module TFs in each treatment-specific dynamic network (using edge lengths inversely proportional to the cross-weighted score) (**Figure 7b**). Paths were assigned infinite length if such a path did not exist. We normalized these path lengths against the average path length in a given network, and found that for the WAB, WAN, and WA treatments, paths from Wnt to many of the module TFs were non-existent. This was again consistent with our hypothesis that differing network topologies underlie the treatment-specific differences we see in cell fate. Further, this analysis provides a basis for identifying which signaling pathways must be targeted for directing certain fate transitions, for example through an exhaustive search of effector targets.

Suppression of PI3K signaling encourages endodermal fate

Our signaling effector target analysis also revealed increased Foxo1 activity in WAB, WAN, and WA treated cells as compared to WAG treated cells, indicating the suppression of PI3K signaling along the WAB, WAN, and WA trajectories (**Figure 7a**). Importantly, it is known that the establishment of definitive endoderm (DE) requires suppression of PI3K signaling (McLean et al., 2007; Yu et al., 2015). Further, it has been recently shown in hiPSC to DE

differentiation that Foxo1 binds to DE-formation-related genes and that inhibition of Foxo1

during DE induction impedes DE establishment (Nord et al., 2020). We therefore hypothesized

that a second fate choice between mesodermal and endodermal fates further exacerbated the

uneven mesodermal fate preference between the treatments. Indeed, our analysis indicates that

in the mesodermal-endodermal fate choice, the majority of WAG treated cells differentiate

toward mesodermal fate. This is in contrast to the WAN and WA treatment in which cells

preferentially differentiate toward definitive endoderm. WAB treated cells exhibited mixed

potential, and split roughly equally along the two paths.

We sought to elucidate a possible explanation for Foxo1's role in this fate choice by

searching for the shortest paths toward a set of endodermal genes. Specifically, similar to the

previous mesodermal analysis, we applied community detection to the full endodermal network

and identified the cluster containing Sox17 and Foxa2, known master regulators of endodermal

fate (**Figure 7c**). This cluster included the transcription factors Sox17, Foxa2, Bmp2, Cited1,

Foxa1, Gata6, Hhex, Lhx1, and Tcf7l2. For each treatment-specific network, we computed the

shortest paths from Foxo1 targets (as determined in the same manner as the previous section)

to these genes (**Figure 7d**). Importantly, we found that many of these regulators, including both

Sox17 and Foxa2, were not reachable from Foxo1 in the WAG network, consistent with the

observation that WAG treated cells preferentially differentiated toward mesodermal fate over

endodermal fate. Interestingly, Foxa2 was not reachable in the WAN network, though Sox17

was. We believe this may explain the differences in distribution of cells between the endodermal

clusters 7,1 and 7,0. Namely, of the cells that specify endodermal fate, WA and WAB treated

cells more fully transitioned into cluster 7,0. In contrast, WAN treated cells remained mostly in

cluster 7,1 and were less likely to differentiate further into 7,0. Previous studies in directed

differentiation of human ESCs toward DE have implicated Sox17's regulatory role in

establishing DE to be upstream of Foxa2, the loss of which impairs foregut and subsequent

hepatic endoderm differentiation (Genga et al., 2019). This is consistent with our cell type

annotations of cluster 7,1 (definitive endoderm) and 7,0 (gut endoderm), and offers a possible

mechanism of the discrepancy in cell fate preferences between treatments.

Discussion

Methods of inferring regulatory network structures are essential for understanding cell

identity transitions, uncovering mechanisms of disease progression due to dysfunctional gene

regulation, and answering other regulatory questions in biology. Unfortunately, experimental

methods are unscalable and existing computational methods suffer from poor performance or

high computational burden. Here, we present a novel GRN reconstruction tool, Epoch, which

leverages single-cell transcriptomics and efficiently infers dynamic network structures. We

demonstrate that Epoch out performs commonly used methods CLR and GENIE3 in both

synthetically generated and real-world datasets. Furthermore, Epoch's relative efficiency allows

it to be easily applied iteratively for applications requiring fine-tuning of network topology or

iterative simulations of changes in network structure.

Further, the Epoch framework is flexible. There are no strict requirements on

pseudotemporal input as we demonstrate here using synthetic simulation time, diffusion

pseudotime, and RNA velocity-based latent time. We further validate on the in vivo muscle

development dataset that precision-recall is largely comparable for different pseudotime

methods (**Supplementary Figure 1f,g**). Additionally, Epoch's workflow is structured in discrete

steps allowing users to pick-and-choose as well as substitute portions of the workflow. We

anticipate that this flexibility will also be useful as advances in sequencing technology, such as

multi-modal methods, and analysis techniques, such as novel cell ordering or reconstruction

methods, can be easily adapted into Epoch's framework. Importantly, because of single-cell

transcriptomic data's inherent noisiness, we also asked if smoothing gene expression over time

could alleviate noise-related drops in reconstruction performance. We compared performance

between unsmoothed and kernel-smoothed data and found that the there was no significant

difference (**Supplementary Figure 1e**). However, smoothing functions remain in Epoch's code

base to aid in cleaner visualizations of gene dynamics.

Finally, we applied Epoch to scRNA-seq data from day 0 through day 4 *in vitro* mESC

directed differentiation to uncover underlying dynamic networks driving mesodermal lineage

specification. Our data consisted of cells transitioning into all three germ layers, and Epoch was

able to recover a number of key lineage drivers. Because our data consists of cells undergoing

four different treatments (WAG, WAB, WAN, WA), we were additionally able to reconstruct

treatment-specific GRNs. In direct contrast to the cells in the WAB, WAN, and WA treatment

groups, the cells in the WAG treatment group had higher mesodermal fate potential and

exhibited a relatively stunted neuroectodermal trajectory. To understand this observation we

applied a number of network comparison strategies to elucidate differences in network topology

between treatments responsible for this fate choice. Importantly, we found three sets of

topological differences that may play a role in specifying mesodermal fate.

First, we identified a set of TF modules that were preferentially activated in WAG treated

cells compared to cells undergoing the other three treatments along the mesodermal fate.

Specifically, these modules encompassed the transcription factors Peg3, Lhx1, Hoxb2, Foxc1,

Tshz1, Meis2, Mesp1, Tbx6, Foxc2, Prrx2, Meox1, and Notch1. Upon further analysis, we

observed that the transcription factor Peg3 was predicted by Epoch to have a role in regulating

a large fraction of these mesodermal TFs. This influence was specific to WAG treated-cells, and

indeed, prior studies have implicated a GSK3-dependent pathway for the epigenetic regulation

of imprinted loci including Peg3 (Meredith et al., 2015). Peg3 expression has been detected in

early somites and other mesodermal tissues (Kuroiwa et al., 1996). Interestingly, it has been

separately demonstrated that knockdown of Peg3 in mouse embryonic fibroblasts and neural

stem cells increases expression of pluripotency genes and enhances overall reprogramming

efficiency (Theka et al., 2017). Taken together these results suggest a role for Peg3 in

modulating the exit of pluripotency and mesodermal lineage specification. If so, this finding

could paint Peg3 as a highly enticing transcription factor to target with the goal of improving

efficiency of directed differentiation protocols toward certain mesodermal fates like somites.

Next, we compared topological differences in the treatment-specific networks by

examining the existence of, and computing the length of, shortest paths. Specifically, we sought

to trace paths from signaling effectors to the activation of mesodermal genes within these

dynamic networks, thereby providing possible mechanisms of how genetic programs are

activated from signal transduction pathways. A number of studies have pointed to the influence

of "Wntch" signaling on the neuroectoderm-mesendoderm cell fate decision in gastrulation, as

reviewed by (Hayward et al., 2008). Our data was consistent with these reports: WAB, WAN,

WA treated cells exhibited activated Notch pathways and exhibited full trajectories toward

neuroectoderm fate; WAG treated cells exhibited markedly diminished or non-existent Notch

signaling and reached mesodermal fates while seemingly displaying a stunted capacity for

neuroectodermal fate.

Our analysis indicated the existence of paths from Wnt signaling effectors to the

aforementioned mesodermal TFs within the WAG network. Analogous paths were markedly

sparser in WAB, WAN, and WA networks, and some mesodermal TFs were entirely

unreachable. This suggests topological network differences caused by treatment, though the

nature of the emergence of these differences remains unclear. One possibility is that the

network structures are dynamically altered by epigenetic events initiated through the various

signaling pathways activated in treating the cells. We point to GSK3's reported effect over Peg3

methylation as an example of such a phenomenon. Further studies to elucidate differences in

chromatin accessibility between mESCs undergoing directed differentiation by way of distinct

treatments, or activation of distinct signaling pathways, will aid in clarifying if such a mechanism

is at play.

Finally, we focused our analysis on cells specifying mesendoderm fate, and outlined
topological differences that could lead to differences in mesoderm vs. endoderm fate potential in
cells undergoing different treatments. We applied an analogous shortest path analysis along the
endodermal fated path to identify the link between suppression of PI3K signaling and endoderm
specification. We again found topological differences between WAG and the remaining three
treatments, this time indicating that many endodermal TFs, such as essential endodermal
regulators Sox17 and Foxa2, were unreachable from PI3K suppression in the WAG network,
explaining the preference of these cells to transition into mesodermal rather than endodermal
fates. Interestingly, we found that Foxa2 was not reachable from PI3K suppression in the WAN
network, which was consistent with our finding that of the WAN treated cells that specified
endodermal fate, the majority tended to remain in earlier endodermal stages based on RNA
velocity. This was contrast to WAB and WA treated cells which tended to transition more fully
into a later endodermal stage. Taken together, these results offer explanation of differences in
mesodermal vs. endodermal fate potential between the treatments.
We believe the single-cell directed differentiation dataset collected in this study will be a
useful resource for further analysis of decision-making in early gastrulation. Moreover, it will be
illuminating to compare the networks reconstructed from this dataset against networks
reconstructed from *in vivo* gastrulation. Furthermore, the methodologies presented here,
available in the R package Epoch and the Python module PyEpoch, are broadly applicable to
any biological question that may benefit from uncovering dynamic regulatory network structures
and the comparison of such networks. In particular, we can apply Epoch to understand cell fate
transitions and decisions at branch points in lineage trajectories, uncover key regulators driving
these decisions, and identify paths from the signal transduction level to the transcriptional
regulation level. Such an approach provides a powerful strategy to not only elucidate dynamic,
multi-scale processes in development, but also to identify signaling pathways to dysregulate for
the purposes of directing cell identity transitions *in vitro*.
559
560

561   **Acknowledgements**
562   We thank Chris McGinnis and the Gartner Lab for their help implementing the MULTI-seq
563   protocol and for providing the required reagents.

564
565
566

Methods

mESC maintenance and differentiation

GFP-Brychyury mESC cells (Gadue et al., 2006) were maintained and differentiated through day 2 of the directed differentiation protocol according to Spangler et al. (Spangler et al., 2018). At day 2, four different primitive-streak induction treatments were established by adding growth factors Wnt3a (25ng/mL) and Activin A (WA; 9ng/mL) along with one of the following additional growth factors: Noggin (150ng/mL), Gsk inhibitor (10mM), or BMP4 (0.5ng/mL). On the day of sequencing, EBs and monolayers were dissociated to a single cell suspension through the use of TrypLE and 40uM cell strainers.

Multi-Seq protocol

We utilized the MULTI-seq sample barcoding and library preparation protocol from the McGinnis lab (McGinnis et al., 2019) in order to sequence 27 samples in two 10x capture runs. In short, the protocol involved tagging cell membranes with sample specific barcodes using a lipid-modified oligonucleotide (LMO). The LMOs (reagents obtained from the McGinnis lab) anchor into the cell membrane and allow for the attachment of a normal ssDNA oligonucleotide (MULTI-Seq barcode). A unique MULTI-seq barcode was added to each sample after which, all samples were pooled and sequenced as if they were one sample. An equal number of cells from each sample were combined to make up the pooled samples and ensure equal representation of each sample after down-stream sequencing. MULTI-Seq barcodes were used down-stream to identify which cells were from which samples.

Library preparation and sequencing

After cells were successfully tagged and pooled, they were submitted to the sequencing core facility for 10x capture and library preparation. A Truseq library preparation was performed with the necessary adjustments made to accommodate the MULTI-Seq platform. Libraries were sequenced on Illumina NovaSeq.

Single-cell data processing

Sequencing alignment to the mm10 reference genome was performed using CellRanger

(version 3.1), and spliced, unspliced, and ambiguous read counts were extracted using

Velocyto. Sample barcode classification was performed using the deMULTIplex R package

(https://github.com/chris-mcginnis-ucsf/MULTI-seq). Subsequent quality control filtering,

normalization, clustering, and differential gene expression analysis was performed using

Scanpy (version 1.4.5) (Wolf et al., 2018). Specifically, doublets and negatives from the sample

barcode classification were removed, along with 271 cells we identified as residual fibroblasts

(Thy1+). Next, genes were excluded if they were detected in less than 5 cells; cells were

excluded if their mitochondrial gene content exceeded 10% of their total reads or if they had

fewer than 500 unique genes. This left us with 5530 cells and 17528 genes. The data was then

normalized and log transformed before highly variable genes were identified (1938 genes). The

data was scaled and PCA was performed. Leiden clustering was performed and visualized on a

UMAP embedding. We further removed a cluster of possible dying cells expressing relatively

high mitochondrial gene content and low total read count, leaving a dataset with 5228 cells and

1938 variable genes. Leiden clustering was refined, and RNA velocity was computed using

scvelo (version 0.2.2) (Bergen et al., 2020). Latent time was computed with scvelo, using root

and end states that were computed through CellRank (version 1.0.0-rc.0). Cell type

classification was performed via SingleCellNet (Tan and Cahan, 2019), using complied embryo

data for training the classifier. For the E12.5 muscle development data and day 4 early directed

differentiation data, diffusion maps and diffusion pseudotime were computed through Scanpy.

Synthetic benchmarking

15 synthetic datasets were generated using the Dyngen package in R (Cannoodt et al., 2020).

Specifically networks were generated with 100-420 genes (including 20-70 TFs), containing

varied topological motifs, and cell trajectories were simulated over time. Simulated single-cell

sequencing was performed, resulting in datasets that included up to 1000 cells. CLR was

implemented with the minet package in R (Meyer et al., 2008), and GENIE3 was implemented

with the GENIE3 package in R (Aibar et al., 2017; Huynh-Thu et al., 2010). Variations of these

two methods were carried out using the Epoch framework. Performance (AUPR) was compared

against sets of 5 random networks generated for each reconstructed network via the random

permutation of edge weights across targets.

Bootstrapped network reconstruction of early MULTI-seq data

We isolated early latent time clusters 1 and 3 and uniformly sampled 400 cells, 10 times to

reconstruct 10 networks. For each reconstructed network, we extracted top regulators by

PageRank and identified TFs that were most frequently ranked in the top 10 regulators. We

further computed the average normalized betweenness and average normalized degree across

the 10 reconstructed networks to identify top regulators.

Differential network analysis and assessment

As one method of network comparison, we used a naïve edge weight-based approach to extract

the most differentially weighted edges between networks. Such an approach resulted in

"differential networks" which are representative of unique topological features existing in each

treatment-specific network. To assess the feasibility of this simple method in extracting real and

important topological differences between networks, we first applied the same method to

compare the three lineage networks in the dataset. The ground truth on which we assessed

these differential networks was a set of lineage-unique "pseudo-ChIP-seq" networks we

compiled similar to TF-promoter binding predictions generated by (Lu and Mar, 2020)

(**Supplementary Figure 6a**). In short, TF-target interactions were acquired by mapping the

PWM of a TF (Weirauch et al., 2014) to the promoter region sequence of potential target genes

using the Find Individual Motif Occurrences (FIMO) software (described in more detail below).

These ground truth networks were filtered for dynamically expressed genes along given paths

(for lineage-specificity), and further filtered for unique edges to mimic extraction of unique

topological features (lineage-unique). Using this set of networks, we compared AUPR of our

path-differential networks against random networks (generated by shuffling edge weights of

targets of each TF in the differential networks). On average the differential networks had a 1.7

fold improvement over random AUPR, with many comparisons falling in the 1.8-3.0 range

(**Supplementary Figure 6b**).

Construction of path-unique TF-target pseudo-gold standard

Transcription star site (TSS) annotations were downloaded from the USCS Genome Browser

(mouse GRCm38/mm10 genome assembly) (Haeussler et al., 2019). Promoter regions were

defined as -750 to +250 around the TSSs. We then acquired position weight matrices (PWMs)

of transcription factors from the Catalog of Inferred Sequence Binding Preferences (CIS-BP

Database build version 2.00) (Weirauch et al., 2014). We used the Find Individual Motif

Occurrences tool (Grant et al., 2011) to obtain potential TF-promoter binding pairs for 648 TFs.

We kept all interactions for which p-value < 5 x10^{-5}. To generate path-unique gold standards, for

each lineage (mesoderm, endoderm, neuroectoderm) we filtered interactions for dynamically

expressed genes along each path as computed by Epoch. Further, when assessing the

"differential network" pairwise comparisons, we filtered out interactions that were redundant

between the gold standards, and only kept interactions unique to each path.

TF-target level comparison

To quantify the extent of topology differences between treatments along the mesodermal path,

we performed pairwise comparison on the four treatment-specific networks and computed

Jaccard similarity between predicted targets of TFs. The comparison was carried out on 72 TFs

that were selected based on their active expression on day 3 or day 4 (i.e. after induction

treatments). We computed baseline similarities for each TF in each treatment by bootstrapping

the network reconstruction (10 network reconstructions per treatment, 400 sampled cells per

network reconstruction) and averaging their pairwise Jaccard similarities. We selected the 13

TFs exhibiting the greatest differences in targets amongst the treatments and performed gene

set enrichment analysis using the Enrichr R package (Chen et al., 2013; Kuleshov et al., 2016)

and the GO Consortium Biological Processes database (Ashburner et al., 2000; Gene Ontology

Consortium, 2021) on the targets of each TF. We further filtered out terms that did not meet the

criteria of adjusted p-value<0.05. Finally, we summarized the results between the 13 TFs by

counting the frequency (out of 13) that a term was considered enriched and ranked the results

accordingly.

Comprehensive signaling effector TF targets list

We acquired binding score (MACS2) data for 18 signaling effector TFs from the ChIP-Atlas (Oki

et al., 2018), and processed each accordingly: target genes were ranked by maximum binding

score and the top 2000 targets were retained (or all retained, if less than 2000 targets). To

compute average effector activity, target lists were filtered for those appearing in the MULTI-seq

directed differentiation data, and not considered in the analysis if the effector had less than 10

targets after filtering.

Epoch Workflow

Epoch takes as input processed (normalized and log-transformed) single-cell transcriptomic

data and accompanying pseudotime (or equivalent) annotation. The Epoch workflow is based in

three strategies: (1) the extraction of dynamically expressed genes and subsequent static

reconstruction with CLR, (2) network refinement using a cross-correlation based strategy

termed "crossweighting," (3) the extraction of a dynamic network and subsequent identification

of top regulators. Epoch is available as a package in R, and code and tutorials can be found at

https://github.com/pcahan1/epoch.

Identification of dynamically expressed genes

Limiting network reconstruction to dynamically expressed genes serves two purposes.

Importantly, it focuses the network on interactions that are more likely playing a role in the

observed biological process. Second, it reduces the instances of false positive interactions and

improves precision by limiting possible edges between temporally-variable genes. To select for

such genes, Epoch models individual genes across the annotated pseudotime using a

generalized additive model. Specifically, Epoch uses the 'gam' package (gaussian family, loess

smooth pseudotime) to fit a model for each gene using the backfitting algorithm. Genes are

considered dynamic based on significance of the smooth term (pseudotime).

CLR

The Context Likelihood of Relatedness (CLR) algorithm was first proposed in 2007 by Faith et

al. as a method for inferring gene regulatory networks from expression data. In short, CLR

computes the mutual information (MI) between expression levels of every potential transcription

factor and target gene. From here the method scores the MI values in the context of the network

by computing a z-score relying on (1) the distribution of MI values for all potential regulators of

the target gene, z_i, and on (2) the distribution of MI values for all potential targets of the

transcription factor, z_j:

$$z_{ij} = \sqrt{z_i^2 + z_j^2}$$

The result is a weighted adjacency matrix describing the likelihood of TF-target gene

interactions that can be thresholded to derive a network structure. For Epoch, a version of CLR

is applied to dynamically expressed genes to reconstruct an initial static network via the 'minet'

package in R. Users can choose between implementing the MI version or a version using

Pearson correlation in place of MI. In either case, Epoch at this step will return a GRN table.

Each row outlines an interaction in the initial network identifying the transcription factor, target

gene, correlation, and CLR-based z-score.

Cross-weighting

After inferring an initial network structure, Epoch can apply a network refinement step we call

cross-weighting. The objective of this step is to negatively weight edges in the initial network

that are unlikely to be true interactions, that may, for example, be representative of indirect

interactions. To this end, for every TF-target pair in the initial network, Epoch computes cross-

correlation across a given lag time (defaults to a fifth of total psuedotime). After ordering the lag

times by decreasing correlation, Epoch computes an offset value where maximum correlation is

achieved, defined by default as the top third of the ranked lag times. Finally, Epoch scores the

offset values of each interactions:

753
$$weight = \begin{cases} 1 & x \leq w_{min} \\ 0 & x \geq w_{max} \\ -x/(w_{max} - w_{min}) + 1 & else \end{cases}$$

where the offset, x, is computed as described, and maximum and minimum windows can be

altered by the user. Z-scores from the initial network are weighted accordingly, ultimately

filtering out false positives. Epoch, at this step will return an appended GRN table, including for

each interaction, the offset value and new weighted score.

Dynamic Network Extraction

Epoch will extract a dynamic network from the reconstructed static network. Specifically, the

process begins by breaking pseudotime into "epochs," or time periods. A number of options are

available to users to accomplish this. Briefly, Epoch can define the epochs based on

pseudotime, equal cell ordering (resulting in equal number of cells per epoch), k-means

clustering, or user-defined manual assignment. The number of epochs is specified by the user,

and can be determined by examining the heatmap of gene expression across pseudotime and

estimating the rough number of expression states represented in the data. After epochs are

defined, Epoch will assign genes to epochs, based on their activity along pseudotime. In brief,

this is either based on activity (i.e. genes active in any epochs will be assigned to those

epochs), or based on differential expression (i.e. genes are assigned based on if they are

differentially expressed in an epoch). Specifically, if genes are assigned by activity, Epoch will

compute, for each gene, a threshold against which average expression of the gene in an epoch

is compared. This threshold can be modified by user input. If instead genes are assigned by

differential expression, a p-value threshold is used to determine assignments. Finally, "orphan

genes," which we define as dynamically expressed genes that do not get assigned to any

epoch, are assigned to the epoch in which their average expression is maximum.

Based on these assignments, Epoch will fracture the static network into a dynamic one

composed of "epoch networks" and "transition networks". Specifically, an edge between

regulator and target gene appears in an epoch network if the regulator is assigned to that

epoch. Further, an edge will appear in a transition subnetwork under two conditions: (1) For an

activating edge, the target is not active in the source epoch, is active in the subsequent epoch,

and regulator is active or (2) For a repressive edge, the target is active in the source epoch, is

not active in the subsequent epoch, and the regulator is active. Following this step, Epoch will

return a dynamic network represented by a list of individual GRN tables. This includes the

epoch networks or essentially the dynamic network, as well as transition networks that describe

how an epoch network may transition into a subsequent epoch network.

Top Regulator Prediction

Epoch employs two graph theoretic methods to predict "top regulators," the transcription factors

that appear to have the most influence in driving changes in or maintaining topology. First,

Epoch will rank regulators by weighted PageRank. In brief, the PageRank centrality measures

the importance of nodes in a network based on the number and quality of links of which a node

is a part. In essence, the most influential nodes are likely to interact with many influential nodes.

Second, Epoch will rank regulators by the product of normalized betweenness and normalized

degree. Here, the assumption is that the most influential nodes are likely to be traversed by

many shortest paths (and thus have high betweenness) and interact with many other nodes

(and thus have high degree). In Epoch, PageRank, betweenness, and degree centralities are

implemented through the 'igraph' package. By default, cross-weighted edge weights are used,

but can be further specified by the user. In both top regulator prediction methods Epoch will

return ranked lists of nodes, and further specify their corresponding PageRank or normalized

betweenness, normalized degree, and betweenness-degree product.

Acknowledgements

We would like to thank Yuqi Tan, Ray Cheng, and David Johanson for helpful discussions and

testing Epoch code. This work was supported by the National Institutes of Health under grant

R35GM124725 to PC and by the National Science Foundation Graduate Research Fellowship

under Grant No. DGE-1746891 to EYS.

Bibliography

Aibar, S., González-Blas, C.B., Moerman, T., Huynh-Thu, V.A., Imrichova, H., Hulselmans, G., Rambow, F., Marine, J.-C., Geurts, P., Aerts, J., van den Oord, J., Atak, Z.K., Wouters, J., Aerts, S., 2017. SCENIC: single-cell regulatory network inference and clustering. Nat. Methods 14, 1083–1086. doi:10.1038/nmeth.4463

Androutsellis-Theotokis, A., Leker, R.R., Soldner, F., Hoeppner, D.J., Ravin, R., Poser, S.W., Rueger, M.A., Bae, S.-K., Kittappa, R., McKay, R.D.G., 2006. Notch signalling regulates stem cell numbers in vitro and in vivo. Nature 442, 823–826. doi:10.1038/nature04940

Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., Harris, M.A., Hill, D.P., Issel-Tarver, L., Kasarskis, A., Lewis, S., Matese, J.C., Richardson, J.E., Ringwald, M., Rubin, G.M., Sherlock, G., 2000. Gene Ontology: tool for the unification of biology. Nat. Genet. 25, 25–29. doi:10.1038/75556

Aubert, J., Dunstan, H., Chambers, I., Smith, A., 2002. Functional gene screening in embryonic stem cells implicates Wnt antagonism in neural differentiation. Nat. Biotechnol. 20, 1240–1245. doi:10.1038/nbt763

Bergen, V., Lange, M., Peidli, S., Wolf, F.A., Theis, F.J., 2019. Generalizing RNA velocity to transient cell states through dynamical modeling. BioRxiv. doi:10.1101/820936

Bergen, V., Lange, M., Peidli, S., Wolf, F.A., Theis, F.J., 2020. Generalizing RNA velocity to transient cell states through dynamical modeling. Nat. Biotechnol. 38, 1408–1414. doi:10.1038/s41587-020-0591-3

Cannoodt, R., Saelens, W., Deconinck, L., Saeys, Y., 2020. dyngen: a multi-modal simulator for spearheading new single-cell omics analyses. BioRxiv. doi:10.1101/2020.02.06.936971

Chan, M.M., Smith, Z.D., Grosswendt, S., Kretzmer, H., Norman, T.M., Adamson, B., Jost, M., Quinn, J.J., Yang, D., Jones, M.G., Khodaverdian, A., Yosef, N., Meissner, A., Weissman, J.S., 2019. Molecular recording of mammalian embryogenesis. Nature 570, 77–82. doi:10.1038/s41586-019-1184-5

Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., Ma'ayan, A., 2013. Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics 14, 128. doi:10.1186/1471-2105-14-128

Chen, J., Schlitzer, A., Chakarov, S., Ginhoux, F., Poidinger, M., 2016. Mpath maps multi-branching single-cell trajectories revealing progenitor cell progression during development. Nat. Commun. 7, 11988. doi:10.1038/ncomms11988

Chen, T., Ueda, Y., Xie, S., Li, E., 2002. A novel Dnmt3a isoform produced from an alternative promoter localizes to euchromatin and its expression correlates with active de novo methylation. J. Biol. Chem. 277, 38746–38754. doi:10.1074/jbc.M205312200

di Bernardo, D., Thompson, M.J., Gardner, T.S., Chobot, S.E., Eastwood, E.L., Wojtovich, A.P., Elliott, S.J., Schaus, S.E., Collins, J.J., 2005. Chemogenomic profiling on a genome-wide scale using reverse-engineered gene networks. Nat. Biotechnol. 23, 377–383. doi:10.1038/nbt1075

Faith, J.J., Hayete, B., Thaden, J.T., Mogno, I., Wierzbowski, J., Cottarel, G., Kasif, S., Collins, J.J., Gardner, T.S., 2007. Large-scale mapping and validation of Escherichia coli transcriptional regulation from a compendium of expression profiles. PLoS Biol. 5, e8. doi:10.1371/journal.pbio.0050008

Gadue, P., Huber, T.L., Paddison, P.J., Keller, G.M., 2006. Wnt and TGF-beta signaling are required for the induction of an in vitro model of primitive streak formation using embryonic stem cells. Proc. Natl. Acad. Sci. USA 103, 16806–16811. doi:10.1073/pnas.0603916103

Gene Ontology Consortium, 2021. The Gene Ontology resource: enriching a GOld mine. Nucleic Acids Res. 49, D325–D334. doi:10.1093/nar/gkaa1113

Genga, R.M.J., Kernfeld, E.M., Parsi, K.M., Parsons, T.J., Ziller, M.J., Maehr, R., 2019. Single-
Cell RNA-Sequencing-Based CRISPRi Screening Resolves Molecular Drivers of Early
Human Endoderm Development. Cell Rep. 27, 708–718.e10.
doi:10.1016/j.celrep.2019.03.076
Grant, C.E., Bailey, T.L., Noble, W.S., 2011. FIMO: scanning for occurrences of a given motif.
Bioinformatics 27, 1017–1018. doi:10.1093/bioinformatics/btr064
Haeussler, M., Zweig, A.S., Tyner, C., Speir, M.L., Rosenbloom, K.R., Raney, B.J., Lee, C.M.,
Lee, B.T., Hinrichs, A.S., Gonzalez, J.N., Gibson, D., Diekhans, M., Clawson, H.,
Casper, J., Barber, G.P., Haussler, D., Kuhn, R.M., Kent, W.J., 2019. The UCSC
Genome Browser database: 2019 update. Nucleic Acids Res. 47, D853–D858.
doi:10.1093/nar/gky1095
Haghverdi, L., Büttner, M., Wolf, F.A., Buettner, F., Theis, F.J., 2016. Diffusion pseudotime
robustly reconstructs lineage branching. Nat. Methods 13, 845–848.
doi:10.1038/nmeth.3971
Hartemink, A.J., 2005. Reverse engineering gene regulatory networks. Nat. Biotechnol. 23,
554–555. doi:10.1038/nbt0505-554
Hayward, P., Kalmar, T., Arias, A.M., 2008. Wnt/Notch signalling and information processing
during development. Development 135, 411–424. doi:10.1242/dev.000505
Huynh-Thu, V.A., Irrthum, A., Wehenkel, L., Geurts, P., 2010. Inferring regulatory networks from
expression data using tree-based methods. PLoS One 5.
doi:10.1371/journal.pone.0012776
Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev, S.,
Jenkins, S.L., Jagodnik, K.M., Lachmann, A., McDermott, M.G., Monteiro, C.D.,
Gundersen, G.W., Ma'ayan, A., 2016. Enrichr: a comprehensive gene set enrichment
analysis web server 2016 update. Nucleic Acids Res. 44, W90-7.
doi:10.1093/nar/gkw377
Kuroiwa, Y., Kaneko-Ishino, T., Kagitani, F., Kohda, T., Li, L.L., Tada, M., Suzuki, R.,
Yokoyama, M., Shiroishi, T., Wakana, S., Barton, S.C., Ishino, F., Surani, M.A., 1996.
Peg3 imprinted gene on proximal chromosome 7 encodes for a zinc finger protein. Nat.
Genet. 12, 186–190. doi:10.1038/ng0296-186
La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V., Lidschreiber,
K., Kastriti, M.E., Lönnerberg, P., Furlan, A., Fan, J., Borm, L.E., Liu, Z., van Bruggen,
D., Guo, J., He, X., Barker, R., Sundström, E., Castelo-Branco, G., Cramer, P.,
Adameyko, I., Linnarsson, S., Kharchenko, P.V., 2018. RNA velocity of single cells.
Nature 560, 494–498. doi:10.1038/s41586-018-0414-6
Lachmann, A., Xu, H., Krishnan, J., Berger, S.I., Mazloom, A.R., Ma'ayan, A., 2010. ChEA:
transcription factor regulation inferred from integrating genome-wide ChIP-X
experiments. Bioinformatics 26, 2438–2444. doi:10.1093/bioinformatics/btq466
Letsou, W., Cai, L., 2016. Noncommutative biology: sequential regulation of complex networks.
PLoS Comput. Biol. 12, e1005089. doi:10.1371/journal.pcbi.1005089
Lu, T., Mar, J.C., 2020. Investigating transcriptome-wide sex dimorphism by multi-level analysis
of single-cell RNA sequencing data in ten mouse cell types. Biol. Sex Differ. 11, 61.
doi:10.1186/s13293-020-00335-2
Marbach, D., Costello, J.C., Küffner, R., Vega, N.M., Prill, R.J., Camacho, D.M., Allison, K.R.,
DREAM5 Consortium, Kellis, M., Collins, J.J., Stolovitzky, G., 2012. Wisdom of crowds
for robust gene network inference. Nat. Methods 9, 796–804. doi:10.1038/nmeth.2016
Margolin, A.A., Nemenman, I., Basso, K., Wiggins, C., Stolovitzky, G., Dalla Favera, R.,
Califano, A., 2006. ARACNE: an algorithm for the reconstruction of gene regulatory
networks in a mammalian cellular context. BMC Bioinformatics 7 Suppl 1, S7.
doi:10.1186/1471-2105-7-S1-S7
Matsumoto, H., Kiryu, H., Furusawa, C., Ko, M.S.H., Ko, S.B.H., Gouda, N., Hayashi, T.,

Nikaido, I., 2017. SCODE: an efficient regulatory network inference algorithm from
single-cell RNA-Seq during differentiation. Bioinformatics 33, 2314–2321.
doi:10.1093/bioinformatics/btx194
McGinnis, C.S., Patterson, D.M., Winkler, J., Conrad, D.N., Hein, M.Y., Srivastava, V., Hu, J.L.,
Murrow, L.M., Weissman, J.S., Werb, Z., Chow, E.D., Gartner, Z.J., 2019. MULTI-seq:
sample multiplexing for single-cell RNA sequencing using lipid-tagged indices. Nat.
Methods 16, 619–626. doi:10.1038/s41592-019-0433-8
McLean, A.B., D'Amour, K.A., Jones, K.L., Krishnamoorthy, M., Kulik, M.J., Reynolds, D.M.,
Sheppard, A.M., Liu, H., Xu, Y., Baetge, E.E., Dalton, S., 2007. Activin a efficiently
specifies definitive endoderm from human embryonic stem cells only when
phosphatidylinositol 3-kinase signaling is suppressed. Stem Cells 25, 29–38.
doi:10.1634/stemcells.2006-0219
Meredith, G.D., D'Ippolito, A., Dudas, M., Zeidner, L.C., Hostetter, L., Faulds, K., Arnold, T.H.,
Popkie, A.P., Doble, B.W., Marnellos, G., Adams, C., Wang, Y., Phiel, C.J., 2015.
Glycogen synthase kinase-3 (Gsk-3) plays a fundamental role in maintaining DNA
methylation at imprinted loci in mouse embryonic stem cells. Mol. Biol. Cell 26, 2139–
2150. doi:10.1091/mbc.E15-01-0013
Meyer, P.E., Kontos, K., Lafitte, F., Bontempi, G., 2007. Information-theoretic inference of large
transcriptional regulatory networks. EURASIP J. Bioinform. Syst. Biol. 79879.
doi:10.1155/2007/79879
Meyer, P.E., Lafitte, F., Bontempi, G., 2008. minet: A R/Bioconductor package for inferring large
transcriptional networks using mutual information. BMC Bioinformatics 9, 461.
doi:10.1186/1471-2105-9-461
Nord, J., Schill, D., Pulakanti, K., Rao, S., Cirillo, L.A., 2020. The transcription factor FoxO1 is
required for the establishment of the human definitive endoderm. BioRxiv.
doi:10.1101/2020.12.22.423976
Oki, S., Ohta, T., Shioi, G., Hatanaka, H., Ogasawara, O., Okuda, Y., Kawaji, H., Nakaki, R.,
Sese, J., Meno, C., 2018. ChIP-Atlas: a data-mining suite powered by full integration of
public ChIP-seq data. EMBO Rep. 19. doi:10.15252/embr.201846255
Popkie, A.P., Zeidner, L.C., Albrecht, A.M., D'Ippolito, A., Eckardt, S., Newsom, D.E., Groden,
J., Doble, B.W., Aronow, B., McLaughlin, K.J., White, P., Phiel, C.J., 2010.
Phosphatidylinositol 3-kinase (PI3K) signaling via glycogen synthase kinase-3 (Gsk-3)
regulates DNA methylation of imprinted loci. J. Biol. Chem. 285, 41337–41347.
doi:10.1074/jbc.M110.170704
Qiu, P., Simonds, E.F., Bendall, S.C., Gibbs, K.D., Bruggner, R.V., Linderman, M.D., Sachs, K.,
Nolan, G.P., Plevritis, S.K., 2011. Extracting a cellular hierarchy from high-dimensional
cytometry data with SPADE. Nat. Biotechnol. 29, 886–891. doi:10.1038/nbt.1991
Qiu, X., Rahimzamani, A., Wang, L., Ren, B., Mao, Q., Durham, T., McFaline-Figueroa, J.L.,
Saunders, L., Trapnell, C., Kannan, S., 2020. Inferring Causal Gene Regulatory
Networks from Coupled Single-Cell Expression Dynamics Using Scribe. Cell Syst. 10,
265–274.e11. doi:10.1016/j.cels.2020.02.003
Spangler, A., Su, E.Y., Craft, A.M., Cahan, P., 2018. A single cell transcriptional portrait of
embryoid body differentiation and comparison to progenitors of the developing embryo.
Stem Cell Res. 31, 201–215. doi:10.1016/j.scr.2018.07.022
Street, K., Risso, D., Fletcher, R.B., Das, D., Ngai, J., Yosef, N., Purdom, E., Dudoit, S., 2018.
Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. BMC
Genomics 19, 477. doi:10.1186/s12864-018-4772-0
Tan, Y., Cahan, P., 2019. SingleCellNet: A Computational Tool to Classify Single Cell RNA-Seq
Data Across Platforms and Across Species. Cell Syst. 9, 207–213.e2.
doi:10.1016/j.cels.2019.06.004
Theka, I., Sottile, F., Aulicino, F., Garcia, A.C., Cosma, M.P., 2017. Reduced expression of

Paternally Expressed Gene-3 enhances somatic cell reprogramming through
mitochondrial activity perturbation. Sci. Rep. 7, 9705. doi:10.1038/s41598-017-10016-7
Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N.J., Livak,
K.J., Mikkelsen, T.S., Rinn, J.L., 2014. The dynamics and regulators of cell fate
decisions are revealed by pseudotemporal ordering of single cells. Nat. Biotechnol. 32,
381–386. doi:10.1038/nbt.2859
Treutlein, B., Brownfield, D.G., Wu, A.R., Neff, N.F., Mantalas, G.L., Espinoza, F.H., Desai, T.J.,
Krasnow, M.A., Quake, S.R., 2014. Reconstructing lineage hierarchies of the distal lung
epithelium using single-cell RNA-seq. Nature 509, 371–375. doi:10.1038/nature13173
Velten, L., Haas, S.F., Raffel, S., Blaszkiewicz, S., Islam, S., Hennig, B.P., Hirche, C., Lutz, C.,
Buss, E.C., Nowak, D., Boch, T., Hofmann, W.-K., Ho, A.D., Huber, W., Trumpp, A.,
Essers, M.A.G., Steinmetz, L.M., 2017. Human haematopoietic stem cell lineage
commitment is a continuous process. Nat. Cell Biol. 19, 271–281. doi:10.1038/ncb3493
Villani, A.-C., Satija, R., Reynolds, G., Sarkizova, S., Shekhar, K., Fletcher, J., Griesbeck, M.,
Butler, A., Zheng, S., Lazo, S., Jardine, L., Dixon, D., Stephenson, E., Nilsson, E.,
Grundberg, I., McDonald, D., Filby, A., Li, W., De Jager, P.L., Rozenblatt-Rosen, O.,
Lane, A.A., Haniffa, M., Regev, A., Hacohen, N., 2017. Single-cell RNA-seq reveals new
types of human blood dendritic cells, monocytes, and progenitors. Science 356.
doi:10.1126/science.aah4573
Weirauch, M.T., Yang, A., Albu, M., Cote, A.G., Montenegro-Montero, A., Drewe, P.,
Najafabadi, H.S., Lambert, S.A., Mann, I., Cook, K., Zheng, H., Goity, A., van Bakel, H.,
Lozano, J.-C., Galli, M., Lewsey, M.G., Huang, E., Mukherjee, T., Chen, X., Reece-
Hoyes, J.S., Govindarajan, S., Shaulsky, G., Walhout, A.J.M., Bouget, F.-Y., Ratsch, G.,
Larrondo, L.F., Ecker, J.R., Hughes, T.R., 2014. Determination and inference of
eukaryotic transcription factor sequence specificity. Cell 158, 1431–1443.
doi:10.1016/j.cell.2014.08.009
Wolf, F.A., Angerer, P., Theis, F.J., 2018. SCANPY: large-scale single-cell gene expression
data analysis. Genome Biol. 19, 15. doi:10.1186/s13059-017-1382-0
Yu, J., Smith, V.A., Wang, P.P., Hartemink, A.J., Jarvis, E.D., 2004. Advances to Bayesian
network inference for generating causal networks from observational biological data.
Bioinformatics 20, 3594–3603. doi:10.1093/bioinformatics/bth448
Yu, J.S.L., Ramasamy, T.S., Murphy, N., Holt, M.K., Czapiewski, R., Wei, S.-K., Cui, W., 2015.
PI3K/mTORC2 regulates TGF-β/Activin signalling by modulating Smad2/3 activity via
linker phosphorylation. Nat. Commun. 6, 7212. doi:10.1038/ncomms8212
Zheng, G.X.Y., Terry, J.M., Belgrader, P., Ryvkin, P., Bent, Z.W., Wilson, R., Ziraldo, S.B.,
Wheeler, T.D., McDermott, G.P., Zhu, J., Gregory, M.T., Shuga, J., Montesclaros, L.,
Underwood, J.G., Masquelier, D.A., Nishimura, S.Y., Schnall-Levin, M., Wyatt, P.W.,
Hindson, C.M., Bharadwaj, R., Wong, A., Ness, K.D., Beppu, L.W., Deeg, H.J.,
McFarland, C., Loeb, K.R., Valente, W.J., Ericson, N.G., Stevens, E.A., Radich, J.P.,
Mikkelsen, T.S., Hindson, B.J., Bielas, J.H., 2017. Massively parallel digital
transcriptional profiling of single cells. Nat. Commun. 8, 14049.
doi:10.1038/ncomms14049

Figure legends

Figure 1. Epoch workflow. Epoch relies on a three-step process to reconstruct dynamic network structures: (1) extraction of dynamic genes and CLR, (2) application of crossweighting, and (3) extraction of the dynamic network.

Figure 2. Benchmarking Epoch. (A) Synthetic dataset generation, reconstruction, and assessment. (B) Fold improvement in AUPR over random of three versions of Epoch (Pearson, MI, and MI with crossweighting) against CLR and GENIE3 on GRN reconstruction from synthetic data. (C) Clustering and diffusion pseudotime for E12.5 muscle development data used in benchmarking. (D) Runtimes for reconstruction. Runtimes were measured on a MacBook Pro laptop with 2.9 GHz Intel Core i7 processor, 16GB 2133 MHz DDR3 of RAM). (E) Precision-recall curves for three versions of Epoch (Pearson, MI, and MI with crossweighting), CLR, and GENIE3 based on reconstruction from muscle development dataset.

Figure 3. mESC in vitro directed differentiation. (A) Directed differentiation protocol. Cells were harvested from d0 through d4 samples for MULTI-seq (B) Clustering for MULTI-seq data. (C) RNA velocity for MULTI-seq data. (D) Select marker gene expression. (E) Cell populations by treatment. (F) Quantification of cell fate distribution based on treatment comparing: mesendoderm vs. neuroectoderm, endoderm vs. mesoderm, cluster 7,0 vs. 7,1 along endoderm fate, and cluster distribution along neuroectoderm fate.

Figure 4. MULTI-seq directed differentiation network reconstruction. (A) Latent time annotation for MULTI-seq data. (B) population breakdown by cluster of epoch assignment along each germ layer path. (C) Reconstructed dynamic networks along mesoderm, endoderm, and neuroectoderm paths. (D) Top regulators as predicted by Epoch via betweenness and degree for the third epochs along each germ layer path.

Figure 5. Mesodermal network analysis. (A) The epoch 3 subnetwork of the mesodermal dynamic network. TFs are colored by community and faded by betweenness. Blue and red edges represent activating and repressive edges respectively. (B) Average community expression over time by treatment along the mesodermal path. Communities shown (each row) are from the epoch 2 subnetwork (yellow), transition (green), and epoch 3 subnetworks (aqua). (C) The epoch 3 differential network between WAG (on) and WA (off). Interactions in this network represent edges present in the WAG mesodermal network that are not present in the

1054 WA mesodermal network. TFs are colored by community and faded by betweenness. Blue and

1055 red edges represent activating and repressive edges respectively. (D) Peg3 and Dnmt3a

1056 expression in a sampled portion of relevant cell types in gastrulation data from Chan et al.

1057

Figure 6. Similarity of targets of TFs in different treatments. (A) Jaccard similarity of targets of a

given TF (each row) between treatments (as indicated by treatment 1 and treatment 2).

Similarity is normalized by intra-treatment baseline (expected) similarity, such that values <1

(green) indicate more-than-expected differences. (B) The 13 TFs selected for GSEA and their

average pairwise Jaccard similarity amongst the treatment-specific networks. (C) Summarized

results of performing GSEA on targets of TFs in B. Heatmap shows rankings (top 10 shown)

based on frequency a term is considered enriched amongst the 13 TFs (1 = most frequent

enriched term).

1066

Figure 7. Integrating signaling effectors and tracing paths to target genes. (A) Average

expression of targets of signaling effectors by treatment over time. Pathway and effector are

labled on each row. (B) Shortest path analysis showing length of shortest paths from Wnt

effector targets (each row) to mesodermal target genes (each column) within treatment-specific

networks. Path lengths are normalized against average path length within a network. Blue and

red indicate paths that activate and repress the target gene respectively. If such a path does not

exist, it has length of infinity and is therefore white. (C) The epoch 3 subnetwork of the

endodermal dynamic network. TFs are colored by community and faded by betweenness. Blue

and red edges represent activating and repressive edges respectively. (D) Shortest path

analysis showing length of shortest paths from PI3K suppression (targets of Foxo1 activation)

(each row) to endodermal genes (each column) within treatment-specific networks. As before,

path lengths are normalized against average path length within a network. Blue and red indicate

activation and repression of the target gene respectively. If a path does not exist, it has length of

infinity and is therefore white.

1081

1082 **Supplementary materials**

1083

Supplementary Figure 1. Epoch benchmarking continued. (A) Full results of benchmarking

reconstruction of synthetic data. Four versions of CLR (Pearson vs. MI, crossweighting vs. no

crossweighting), four versions of Epoch (same as CLR), and three versions of GENIE3 (original,

limited to dynamic genes, and with crossweighting) are compared. (B) Full results of

benchmarking reconstruction of the muscle development data. Methods are the same as in A.

(C) Early mESC directed differentiation data (subsetted from Spangler et al. 2018). Zfp42

expression and velocity pseudotime are shown. (D) Full results of benchmarking reconstruction

of the early Spangler et al. data. Methods are the same as in A. (E) Effect of smoothing vs.

unsmoothing on AUPR of reconstruction using various methods. (F) Clustering, diffusion

pseudotime, velocity pseudotime, and latent time of the E12.5 muscle development data. (G)

Precision recall curves of four variations of Epoch using three different pseudotime annotations.

1095

1096   **Supplementary Figure 2.** MULTI-seq directed differentiation cell typing. (A) timepoint,

1097   treatment, total counts, and percent mitochondrial transcripts content of the MULTI-seq data. (B)

1098   Select marker gene expression of the MULTI-seq data. (C) Differential expression analysis of

1099   the MULTI-seq data by cluster.

1100

1101   **Supplementary Figure 3.** MULTI-seq cell type classification and cluster transitions. (A)

1102   SingleCellNet classification results. The classifier was trained from mouse embryo data found in

1103   literature, curated by members in the lab. (B) Sankey diagrams depicting cluster transitions (and

1104   cell fate transition proportions) as broken down by treatment.

1105

1106   **Supplementary Figure 4.** MULTI-seq network reconstruction and top regulators. Heatmaps

1107   depicting TF expression across time and Epoch definitions along the (A) mesoderm path, (B)

1108   endoderm path, and (C) neuroectoderm path. (D) Top 10 regulators in each epoch network

1109   along each germ layer path as predicted by Epoch via PageRank.

1110

1111   **Supplementary Figure 5.** Bootstrapped top regulator prediction for inner cell mass to epiblast-

1112   like transition. (A) Workflow for bootstrapping network reconstruction in order to predict top

1113   regulators. (B) Only cells from early time were used. Clusters used and corresponding diffusion

1114   pseudotime shown here. (C) Top regulators for this early fate transition as predicted using

1115   PageRank and Betweenness-Degree.

1116

Supplementary Figure 6. Proof of concept for differential network analysis. (A) Path-unique

gold standard creation. (B) Performance (AUPR) over random of differential network in

recovering path-unique gold standard edges. Each row is a comparison between two paths.

Each column represents unique edges recovered for a given path. As an example, the top left

box indicates that in the endoderm vs. neuroectoderm comparison, in the first epoch the

endoderm differential network is 1.82 times better than a shuffled network in recovering edges

that are specific to endoderm (and not present in neuroectoderm).

1124

1125   **Supplementary Figure 7.** Mesodermal network and differential networks. The left column ("Full

1126   Reconstruction") depicts the dynamic mesodermal network. The middle and right columns

1127   depict the WAG vs. WA differential network. Middle is the network unique to WAG, right is the

1128   network unique to WA. TFs are colored by community. Blue and red edges represent activating

1129   and repressive edges respectively.

1130

Supplementary Figure 8. Mesodermal network analysis continued. (A) Average community

expression over time by treatment along the full dataset. Communities shown (each row) are

from the epoch 2 subnetwork (yellow), transition (green), and epoch 3 subnetworks (aqua) of

the mesodermal network. (B) Peg3 and Dnmt3a expression across time along the mesodermal

path of the MULTI-seq data. (C) Top differential regulators of mesodermal genes. Blue and red

represent activators and repressors respectively. Y-axis is plotting difference in edge weights

between the WAG and WA networks. Thus, TFs appearing above the 0-line are more unique to

the WAG network, and those below are more unique to the WA network.

1139

1140   **Supplementary Table 1.** Cell type annotation by cluster.

Figure 1

pseudotime

Compute cross correlation

Find dynamically expressed genes, define and assign epochs

Apply CLR to dynamically expressed genes

Weight interactions based on max correlation offset

Extract dynamic networks and top regulators

Epoch 1

Epoch 2

transition network

t

corr

offset 0

Figure 2

Figure 3

A

Day 0, 1, 2, 3, 4: Harvest for MULTI-seq

| DMEM, LIF, Serum, MEFs | SFD | SFD + [WAG, WAB, WAN, WA] |

WAG: { Wnt3a + Activin A + GSKi }
WAB: { Wnt3a + Activin A + Bmp4 }
WAN: { Wnt3a + Activin A + Noggin }
WA: { Wnt3a + Activin A }

Day 0: Remove LIF, serum and MEFs.

Day 2: Introduce factors to induce primitive streak

B

C

D Zfp42, Fgf5, T, Mesp1, Foxa2, Sox1

E WAG, WAB, WAN, WA

F

Figure 4

Figure 5

Figure 6

B

TF	avg(JS)	TF	avg(JS)	TF	avg(JS)
Prrx2	0.06175	Hes7	0.12431	Hey1	0.21341
Tbx6	0.06835	Cited1	0.12985	Lmo2	0.25975
Mesp1	0.09984	Gata6	0.19159	Mixl1	0.28015
Gata4	0.11422	Isl1	0.20265	Lhx1	0.28297
				Ets1	0.29662

C

treatment

negative regulation of transcription, DNA-templated (GO:0045892)
embryonic digestive tract morphogenesis (GO:0048557)
endoderm formation (GO:0001706)
heart development (GO:0007507)
mesoderm formation (GO:0001707)
positive regulation of cell proliferation (GO:0008284)
positive regulation of gene expression (GO:0010628)
positive regulation of macromolecule metabolic process (GO:0010604)
positive regulation of non-canonical Wnt signaling pathway (GO:2000052)
positive regulation of nucleic acid-templated transcription (GO:1903508)
columnar/cuboidal epithelial cell differentiation (GO:0002065)
digestive tract morphogenesis (GO:0048546)
embryonic cranial skeleton morphogenesis (GO:0048701)
embryonic digestive tract development (GO:0048566)
endoderm development (GO:0007492)
gastrulation (GO:0007369)
positive regulation of multicellular organismal process (GO:0051240)
regulation of cell cycle (GO:0051726)
regulation of ubiquitin-protein ligase activity involved in mitotic cell cycle (GO:0051439)
DNA metabolic process (GO:0006259)
embryonic organ morphogenesis (GO:0048562)
anterior/posterior pattern specification (GO:0009952)
epithelial tube morphogenesis (GO:0060562)
negative regulation of cellular macromolecule biosynthetic process (GO:2000113)
negative regulation of gene expression (GO:0010629)

treatment
WAG
WAB
WAN
WA

Figure 7