Changes in wheat rhizosphere microbiota in response to chemical inputs, plant genotype and phenotypic plasticity

Samuel Jacquiod[1], Tiffany Raynaud[1], Eric Pimet[1], Chantal Ducourtieux[1], Leonardo Casieri[2, 3]

Daniel Wipf[2], Manuel Blouin[1*]

[1]Agroécologie, AgroSup Dijon, INRAE, Univ. Bourgogne, Univ. Bourgogne Franche-Comté, F-21000 Dijon, France

[2]Agroécologie, AgroSup Dijon, CNRS, INRAE, Univ. Bourgogne, Univ. Bourgogne Franche-Comté, F-21000 Dijon, France

[3]Mycorrhizal Applications LLC, c/o BRDG Park at the Danforth Center. St. Louis, Missouri (USA)

[*]Corresponding author

Correspondance:

Manuel Blouin

manuel.blouin@agrosupdijon.fr

Running title: Phenotypic plasticity of plant-microbiota interactions

ABSTRACT

Since modern wheat varieties are grown with chemical inputs, we ignore if changes observed in rhizosphere microorganisms between ancient and modern varieties are due to i) breeding-induced changes in plant genotype, ii) modifications of the environment via synthetic chemical inputs, or (iii) phenotypic plasticity, defined as the interaction between the genotype and the environment. In the field, we evaluated the effects of various wheat varieties (modern and ancient) grown with or without chemical inputs (N-fertilizer, fungicide and herbicide together) in a crossed factorial design. We analysed rhizosphere bacteria and fungi by amplicons sequencing and mycorrhizal association by microscopic observations. When considered independently of plant genotype, chemical inputs were responsible for an increase in dominance for bacteria and decrease in evenness for bacteria and fungi. Independently of inputs, modern varieties had richer and more even bacterial communities compared to ancient varieties. Phenotypic plasticity had a significant effect: bacterial and fungal diversity decreased when inputs were applied in ancient varieties but not in modern ones. Mycorrhiza were more abundant in modern than ancient varieties, and less abundant when using chemical inputs. Although neglected, phenotypic plasticity is important to understand the evolution of plant-microbiota associations and a relevant target in breeding programs.

Keywords: ancient varieties; bacteria; fungi; genotype; inputs; landraces; microbial community; modern varieties; mycorrhiza; phenotypic plasticity; rhizosphere; wheat

INTRODUCITON

Rhizosphere microbiota is involved in several key functions for the plant, such as the regulation of nutrient access, environmental disturbance tolerance and disease resistance (Rodriguez et al., 2008; Mei & Flinn, 2010; Farrar et al., 2014). There is abundant evidence that host genotypes have a significant influence on rhizosphere microbial community composition (Lundberg et al., 2012; Peiffer et al., 2013) and its function, with consequences on plant growth, development and immunity (Lemanceau et al., 2017). In agriculture, understanding factors that favour crop's associations with beneficial bacteria and fungi could help in maintaining high crop yields without using synthetic chemical inputs. Breeding and application of synthetic chemical inputs are two interconnected agricultural practices, which can modify plant-microbiota interactions. Therefore, studying the effect of domestication on the rhizosphere microbiota without considering chemical inputs does not allow an accurate understanding of the evolution of plant-microbiota interactions. The effects of breeding and inputs can be described by adopting the formalism of quantitative genetics $P = G + E + G{\times}E$ (Falconer, 1989). In our case, the plant-microbiota interactions can be considered as a phenotypical trait "P", which can be determined by: (i) the plant genotype "G" (either modern or ancient varieties; microorganisms genotypes will not be considered explicitly here), (ii) the environment "E" (modified by agricultural practices such as the application of inputs) and (iii) the interaction between crop genotype and inputs "GxE" (defined as plant phenotypic plasticity). Indeed, previously reported observations call for considering these three important drivers of plant-microbiota interactions, which we will briefly present in the following order: E, G and G{\times}E.

First, synthetic chemical inputs applied in the fields since the Green Revolution can be considered as a modification of the environment (E), with direct modifications of the soil and rhizosphere microbial communities. In a meta-analysis, long term mineral fertilizer application has been shown to increase microbial biomass by 15.1% (Geisseler and Scow,

2014). Fertilization can change microbial community composition, by promoting copiotroph

organisms such as specific members from Actinobacteria and Firmicutes, whereas decreasing

the oligotroph organisms such as specific members from Acidobacteria and Verrucomicrobia

(Ramirez et al. 2012). These modifications of the soil microbial communities can reverberate

on the rhizosphere community. For example, nitrogen application selects less mutualistic

rhizobia, with less benefit to the host (Weese et al. 2015). In a long-term experiment, Ai et al.

(2015) demonstrated that inorganic fertilization decreased the rhizospheric dependence on

root derived carbon for Actinobacteria members. Similarly, plants-AMF (arbuscular

mycorrhizal fungi) symbiotic association can also be altered by inorganic fertilization

(Lamber et al., 2009), with changes in the diversity of AMF (Egerton☐Warburton et al.,

2007).

Second, some changes in plant-microbiota interactions can arise from modifications in the

plant genome (G), through mutation, hybridization or allele fixation. Domesticated plants

changed their interactions with soil microorganisms compared to their wild relatives (Garcia-

Palacios et al., 2013; Milla et al., 2015; Turcotte et al., 2015). Artificial selection for

improved yield in high-input agriculture could unintentionally lead to a reduction of root

microbiota members involved in nutrient acquisition or plant immunity under low input

conditions (Perez-Jaramillo et al., 2016). The effect of domestication (from wild relatives to

cultivated crops) on rhizosphere microbiota has been observed in several crops such as barley

(Bulgarelli et al., 2015), maize (Szoboszlay et al. 2015), foxtail millet (Chaluvadi and

Bennetzen, 2018) and common bean (Perez-Jaramillo et al., 2017). In wheat, the rhizosphere

bacterial communities of ancient varieties was more diverse than modern varieties (Germida

and Siciliano, 2001). The general pattern is a rhizospheric enrichment of Actinobacteria and

Proteobacteria members and a decrease in Bacteroidetes members in modern varieties

compared to wild relatives (Perez-Jaramillo et al., 2018). Changes in plant-microbes

interactions between wild relatives, ancient varieties and modern varieties were also observed

for mycorrhizal associations. Many studies showed that the mycorrhizal association and

responsiveness of modern wheat varieties in terms of yield gain was lower than that of the

ancient varieties or wild relatives (Kapulnik and Kushnir 1991; Hetrick et al., 1992; Zhu et al.,

2001; Leiser et al., 2016). However, a meta-analysis showed that modern varieties were less

intensely colonized but were more mycorrhiza-responsive compared to ancestral genotypes,

concluding on the absence of evidence that agricultural and breeding practices are responsible

for a lack of response to mycorrhiza in new crop genotypes (Lehmann et al., 2012).

Third, artificial selection may have influenced the way plant genotypes respond to inputs in

terms of plant-microbiota interaction (G×E), i.e. the phenotypic plasticity of plant-microbiota

interaction. Phenotypic plasticity denotes the ability of a genotype to exhibit changes in a

specific trait across different environments (Laitinen et al., 2019). Artificial selection of

modern varieties has been very efficient in providing agriculture with productive cultivars

displaying stable performances across diverse environmental conditions, but it is not clear if

these stable yields are linked to the phenotypic plasticity of other traits (Gage et al., 2017),

such as those involved in plant-microbiota interaction. Several authors thus suggest that

understanding G×E interactions could be a novel breeding strategy, coping better with

changing environments while securing stable yields. Since genes responsible for yield (i.e.

mean trait value) and phenotypic plasticity (i.e. variance) are distinct, breeders should

theoretically be able to select both at the same time to generate plastic varieties that adapt

better to a changing environment, while maintaining a decent yield (Kusmec et al., 2017).

Therefore, it is necessary to evaluate the relative importance of E, G and the G×E interaction

for a more accurate understanding of the plant-microbiota interactions in the rhizosphere of

selected crops. Specifically, evaluating the contribution of these parameters can help to

determine if artificial selection has played an important role in the evolution of plant-

microbiota interactions through genetic effects, being either independent (G) or dependent (G×E) of the environment. In this aim, we studied the structure of rhizosphere microbial community of ancient and modern varieties of wheat, in the presence or absence of inputs (N-fertilizer, fungicide and herbicide). We hypothesized that i) the presence of inputs decreases the microbial diversity in the rhizosphere, ii) modern genotypes have lost their ability to establish interactions with some microbial species, iii) the lower plant-microbiota association in modern genotypes is amplified in the presence of inputs. We used an integrated approach coupling the analysis of microscopic observations of mycorrhiza, and amplicon sequencing of the bacterial (16S rRNA gene) and fungal (ITS2) communities.

MATERIALS AND METHODS

Field site description

The field experiment was conducted on dedicated plots at AgroSup Dijon, the Institut National Supérieur des Sciences Agronomiques de l'Aliment et de l'Environnement (47 ° 18'32 "N 5 ° 04'02" E, Dijon, France). The climate of the experimental area is semi-continental, with an average annual temperature of 11 $^{\circ}$C ($\pm$4.5°C). Average precipitation per year is 760.5 mm, monthly sunshine hours are 1848.8 h. The clay loam soil characteristics in the 0-22 cm horizon were as follow: pH_{H2O} = 8.0; 34.6% clay, 36.2% loam, 29.2% sand; 26.7 g.kg^{-1} organic carbon (46.2 g.kg^{-1} organic matter), 2.11 g.kg^{-1} total nitrogen; 294.0 g.kg^{-1} total Ca; 0.020 g.kg^{-1} P_2O_5; 24.30 cmol$^+$.kg^{-1}. The preceding culture was a field bean (*Vicia faba*) for all the experimental plots.

Experimental design and sampling

Two kinds of genotypes, hereafter called "breeding types" consisting of five modern and five ancient wheat varieties were sown in the 3th and 4th of November 2016. Modern varieties were

selected after the 60's in agrosystems concomitantly receiving high levels of inputs: Rubisko (R, 2012) provided by RAGT Semences, Descartes (D, 2014) and Sherlock (S, 2015) provided by Secobra, Alixan (A, 2005) provided by Limagrain and Tulip (T, 2011) provided by Saaten Union (http://www.fiches.arvalis-infos.fr/). Ancient varieties were provided by the non-governmental organization "Graines de Noé" (http://www.graines-de-noe.org/), which promotes the conservation of wheat landraces. Among their 200 varieties, all grown without inputs, we selected some with a local origin: Barbu du Mâconnais (BM, XIXth–beginning XXth century), Blé de la Saône (BS, before 1960), Automne Rouge (AR, XIXth century). To diversify the panel of wheat grown without synthetic inputs, we also selected Alauda (AL), a variety dedicated for organic agriculture or biodynamic, obtained in 2013 by crossing the varieties Probus (1948) and Inntaler (before 1960) and einkorn wheat (*Triticum monococcum*, EW) which is increasingly grown by farmers interested in ancient varieties (https://urgi.versailles.inra.fr/Projects/Achieved-projects/Siregal). Two growing conditions were tested for each variety: i) with inputs (w) and ii) without inputs (w/o), with three replicates of each condition distributed in three blocks. Inputs included fertilizer (CAN 27% Granulé, Dijon Céréales, France) for a total of 150 kg N.ha^{-1}, applied as 50 kg N.ha^{-1} in three times, on week 18 after sowing (09/03/2017), week 25 (24/04/2017) and week 30 (30/05/2017)); herbicide (Bofix™, Dow Agro Science, supplied once at 0.3 l.ha^{-1} on week 23, the 10/04/2017); and fungicide (Bell Star™, Dow Agro Science), applied once at 2.5 kg.ha^{-1} on week 26, the 05/05/2017. The experiment thus consisted in 60 plots (five modern and five ancient varieties, with or without inputs, replicated three times) of 1m² each, separated from the other by 0.8m. In each plot, 300 seeds of each variety were sown manually in seven rows.

Sampling of rhizosphere microbial communities

In each plot, we sampled randomly two wheat rhizospheres. After loosening the soil with a fork, the root system of the plant with its root-adhering soil was extracted from the bulk soil.

In the laboratory, rhizosphere soil was gently removed from the roots by brushing. The two

samples from the same plot were pooled together to obtain a representative sample. All

samples were frozen at -20°C until further processing for DNA extraction.

Microbial community analysis

Total bacterial and fungal diversity and composition from rhizosphere soil samples were

respectively analyzed by sequencing the 16S rRNA gene V3-V4 region, and the ITS2 region

via Illumina Miseq 2x 250 bp paired-end analysis. Total DNA was extracted from 250 mg of

rhizospheric soil using the DNeasy PowerSoil-htp 96 well DNA isolation kit (Qiagen,

France). In two steps, 16S rRNA gene and ITS2 amplicons were generated for all extracts. In

the first step, the V3-V4 hypervariable region of the bacterial 16S rRNA gene was amplified

by polymerase chain reaction (PCR) using the fusion primers U341F (5'-

CCTACGGGRSGCAGCAG-3') and 805R (5'-GACTACCAGGGTATCTAAT-3'), with

overhang adapters (forward: TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG,

reverse: GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG) to allow the successive

addition of multiplexing index-sequences. Fungal ITS2 was amplified using the primers

ITS3F (5'- GCATCGATGAAGAACGCAGC-3') and ITS4R (5'-

TCCTCCGCTTATTGATATGC-3') with overhang adapters (forward:

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGNNNNGCATCGATGAAGAACGC

AGC, reverse:

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGNNNNTCCTCSSCTTATTGATA

TGC) to allow the successive addition of multiplexing index-sequences. First step PCR and

their thermal cycling conditions were as follows: 98°C for 3 min followed by 98°C for 30 s,

55°C for 30 s and 72°C for 30 s (25 and 30 cycles for 16S rRNA and ITS genes, respectively)

and a final extension for 10 min at 72°C. The PCR products of the first step were used as a

template for the second step of PCR. The second PCR amplification added multiplexing index

sequences to the overhang adapters using a unique combination of primers for each sample.

Thermal cycling conditions were as follows: 98°C for 3 mn followed by 98°C for 30 s, 55°C

for 30 s and 72°C for 30 s (8 and 10 cycles for 16S rRNA and ITS genes, respectively) and a

final extension for 10 min at 72°C. PCR products from the second step were deposited on a

2% agarose gel to validate amplification and amplicons size. Amplicon products were purified

using HighPrep™ PCR Clean Up System (AC-60500, MagBio Genomics Inc., USA)

paramagnetic beads using a 0.65:1 (beads:PCR product) volumetric ratio to eliminate DNA

fragments below 100 bp in size and primers. Samples were normalized using SequalPrep

Normalization Plate (96) Kit (Invitrogen, Maryland, MD, USA) and pooled using a 5 µl

volume for each sample. The pooled sample library was concentrated using DNA Clean and

Concentrator™-5 kit (Zymo Research, Irvine, CA, USA). The pooled library concentration

was determined using the Quant-iT™ High-Sensitivity DNA Assay Kit (Life Technologies).

The final pool concentration was adjusted to 4 nM before library denaturation and

sequencing. Amplicon sequencing was performed on an Illumina MiSeq platform using

Reagent Kit v2 [2 x 300 cycles] (Illumina Inc., CA, US). Demultiplexing and trimming of

Illumina adapters and barcodes was done with Illumina MiSeq Reporter software (version

2.5.1.3).

The 16S rRNA gene and ITS amplicon sequences were analyzed internally using a Python

notebook (available upon request). Briefly, using PEAR with default settings, forward and

reverse sequences were assembled. Additional quality checks were conducted using the

QIIME pipeline and short sequences were removed (< 400 bp for 16S and < 300 bp for ITS).

Reference based and *de novo* chimera detection, as well as clustering into operational

taxonomic units (OTUs, hereafter called taxa) were executed using VSEARCH and the

adequate reference databases (Greengenes' representative set of sequences for 16S rRNA

gene and UNITE's ITS2 reference dynamic dataset for ITS). We deliberately choose the OTU

analysis over the Amplicon Sequence Variant analysis (ASVs) for consistency reasons, since

the latter pipeline is currently not available for fungal ITS sequences. The identity thresholds

were appropriately set at 94 % for 16S and 97 % for ITS based on our routine internal

calibration controls (mock communities) in order to get the most accurate resolution. For

simplicity reasons, we refer only to 16S rRNA gene amplicon sequences as "bacterial", since

archaeal sequences were extremely rare. Taxonomy was allocated using UCLUST and the

latest released Greengenes database (v.05/2013). For ITS, the taxonomy assignment was

executed using BLAST algorithm and the UNITE reference database (v.7-08/2016). A

summary table is provided in supporting information to present the sequenced samples and

the number of sequences recovered (Table S1). Sequences have been submitted to the

Sequence read Archive public repository (SRA, https://www.ncbi.nlm.nih.gov/sra), under the

following accession numbers for the 16S rRNA gene amplicon dataset: SUB9063594; and for

the ITS2 amplicon dataset: SUB9104701

Alpha-diversity analysis of microbial communities

To explore the sequencing completeness in terms of diversity recovery per sample, individual

raw rarefaction curves were calculated using the "*vegan*" package in R (Fig. S1). After careful

evaluation of the rarefaction curves, we identified a series of problematic samples coming

from the same column in the 96-well plate, harboring significantly higher sequences counts

but relatively lower OTU discovery rates than the others ($p < 0.001$). In order to avoid biasing

our conclusions, we decided not to consider those samples for further analyses. Due to the

unevenness of sequencing depth and its consequences on diversity assessment, we applied

random re-sampling for normalization of samples in each data set. The samples were rarefied

at 14,000 for both bacteria and fungi. Those levels are considered to be appropriated for

accurate alpha-diversity estimations based on best practices guidelines (Schöler et al., 2017).

Alpha-diversity analysis was perform using the following indices: observed taxa richness (S),

estimated richness (Chao-1), ACE (Abundance-based Coverage Estimator), Simpson index

(1-D, D = Dominance), Shannon index (H) and Equitability (J = H/ln(S)). Alpha-diversity

indices were exported for bacteria and fungi. An additional PERMANOVA on the six

diversity indices taken together was also performed to detect the overall effect of factors on

bacterial and fungal diversity.

Beta-diversity analysis of microbial communities

For the beta diversity analysis, we used the rarefied dataset. In order to consistently analyze

both bacterial and fungal profiles with the same method, we choose the Bray-Curtis

dissimilarity index to generate the distance matrices (package '*vegan*', Dixon, 2003). The

Bray-Curtis dissimilarity index was preferred over other metrics (e.g. UniFrac) as ITS

sequences cannot be aligned to obtain a phylogenetic distance. We first assessed with

PERMANOVA the significance of each factor on the structure of bacterial and fungal

community. As only the breeding type and the block had a significant effect, a partial

distance-based redundancy analysis (db-RDA) was used with the following model: breeding

type + Condition (block) ('*capscale*' function, package '*vegan*', 10,000 permutations).

Network analysis

We investigated the structure of the modern and ancient wheat varieties rhizosphere

microbiota using a network approach based on edge arithmetic (Jacquiod et al. 2020) to focus

on OTU correlations that are specific of each type of breeding. We first calculated two

separated partial correlation matrices amongst dominant rhizosphere OTUs (total summed

counts = 100 minimum, occurrence = 25/51 samples), one for the modern and one for the

ancient varieties using Poisson Log Normal models (PLN, package '*PLNmodels*', Chiquet et

al. 2019). This method allows the combining of several datasets, thus enabling the integration

of both bacterial and fungal sequencing data together using the 'TSS' offset criteria (Total

Sum Scaling). The following model was built to account for the block effect for both modern

and ancient varieties "~ 1 + block + offset". The PLN models were validated by using the

Bayesian Information Criterion (BIC) to determine the most appropriated sparsifying penalty

levels (BIC, R^2 ancient = 0.98 ; R^2 modern = 0.97). We then applied edge arithmetic with the

'*igraph*' package functions (Csardi and Nepusz, 2006). Briefly, we intersected both modern

and ancient networks to systematically determine whether correlations were common to both

breeding types or specific of one breeding type. For visualization, the two networks were then

merged and opened in the '*Cytoscape*' software (Shannon et al. 2003). The complexity of

networks was investigated by means of the degree index, the node betweenness and the edge

betweenness.

Mycorrhizal colonization

Two root systems per plot were collected and pooled together 29 weeks after sowing, from

the 22[th] to the 24[th] of May 2017. Root material was washed thoroughly and prepared for

staining as described in Vierheilig et al. (1998). Roots were cleared at 90°C for 5 to 10 min in

10% KOH, placed in black ink (5% in acetic acid) for 5 min at 90°C for coloration, rinsed

with water and put for 25 min in 8% acetic acid at room temperature. Roots were rinsed again

with water, covered with pure glycerol and stored at 4°C. For microscopic observation, roots

were cut into 1 cm fragments and 15 fragments were placed in glycerol between slide and slip

cover four times per sample. There were two batches of microscope slides for convenience of

counting, called "Series" in the statistical analysis. Mycorrhization rates were assessed

according to Trouvelot et al. (1986) and expressed as mycorrhizal colonization i.e. frequency

of root fragments with mycorrhizal structures at the root system scale (F), intensity of the

mycorrhizal colonization at the root system scale (M) or restricted to myccorhizal root

fragments (m), arbuscule abundance at the root system scale (A) or restricted to mycorrhizal

root fragments (a).

Univariate statistical analysis

The statistical analysis of univariate data, including alpha-diversity indices and mycorrhiza

traits, was performed in Rstudio software (RStudio Team, 2020). Normality and

homoscedasticity of the data was assessed using Shapiro and Bartlett test respectively using R

default functions. Normally distributed data were analyzed with construction of an ANOVA

model, significance was assessed using the D'Agostino test of skewedness on the residual

variance (package '*moments*', Komsta and Novomestky, 2015), followed by a post-hoc

Tukey'Honest Significant Detection test (Tukey's HSD, p<0.05, package '*agricolae*', De

Mendiburu, 2017). Non-parametric data were analyzed with a Kruskal-Wallis test followed by

a post-hoc False Discovery Rate test correction to account for multiple testing (FDR, $p < 0.05$,

package '*agricolae*'). We present the percentage of variance explained and associated *p*-

values from the ANOVA models if variables are normally distributed, and only the *p*-value

from the Kruskal-Wallis test if not normally distributed.

RESULTS

Effects of breeding types and inputs on bacterial community

Breeding type (ancient *vs* modern varieties) had a significant effect on all bacterial alpha

diversity indices, including the richness (explaining 15.56% of the total variance, $p < 0.001$),

Chao-1 (6.9%, $p = 0.025$), ACE (5.9%, $p = 0.044$), Simpson reciprocal ($p = 0.044$), Shannon

($p = 0.002$) and Equitability ($p = 0.012$) (Tables 1, Table S2). The presence/absence of inputs

had a barely significant effect on community evenness, including the Simpson reciprocal ($p =$

0.048) and Equitability ($p = 0.057$) (Tables 1 and S2). The interaction between breeding type

and inputs had a significant effect on richness (7.7%, $p = 0.011$), ACE (6.8%, $p = 0.031$),

Simpson reciprocal ($p = 0.013$), Shannon ($p = 0.001$) and Equitability ($p = 0.006$). The

different varieties inside each breeding type was the main source of variation for richness

(46.1%, $p = 0.009$), Chao1 (49.9%, $p = 0.015$) and ACE (47.5%, $p = 0.028$) indices. When the

six diversity indices were analyzed simultaneously in a PERMANOVA (Table 2), we found

an overall significant effect of the breeding type (7.4%, $p = 0.013$) and its interaction with

chemical inputs (5.7, $p = 0.034$). The variety error term inside breeding type was also

significant (48.8%, $p = 0.015$). The overall multivariate model explained 63.5% of the

variance.

Regarding the direction of the effects of factors on bacterial diversity, we found that,

considering both treatments with and without inputs together, modern varieties had richer and

more even communities as compared to ancient varieties (Fig. 1). Considering both ancient

and modern varieties together, the presence of inputs had significant effects only on evenness

indices, with a slight increase in dominance of some taxa. However, the addition of inputs

was responsible for a strong and significant decrease in diversity for ancient varieties for

richness, Simpson reciprocal, Shannon and Equitability, but it had no effect on modern

varieties (Fig. 1).

The PERMANOVA on bacterial community beta diversity (Table 3) showed that the breeding

type (2.7%, $p = 0.036$) and the block (6.5%, $p < 0.001$) but not the inputs had a significant

effect (1.8%, $p = 0.701$). From this result, we profiled the structural changes in bacterial

communities using a distance-based partial redundancy analysis on Bray-Curtis dissimilarities

for breeding type only (the block effect was added as an error term). The total variance

explained was 2.3% ($p = 0.016$) and the first constrained component (CAP1) and first non-

constrained component (MDS1) were explaining 2.5 and 4.2% of variance, respectively (Fig.

2A). Ordination plots indicated that bacterial community structure differed between breeding

types.

Effects of breeding types and inputs on fungal community

Breeding type had no significant effect on fungal diversity (Table 4, Table S3). The

presence/absence of inputs showed a significant effect on the Shannon diversity (6.7%, p =

0.044). The interaction between breeding type and inputs had a significant effect on the Chao-

1 estimation (13.9%, p = 0.025). The effect of the different varieties inside each breeding type

had no significant effect, despite a relatively high percentage of variance (Table S3). When

the six diversity indices were analyzed simultaneously in a PERMANOVA (Table 2), we

found no significant effects of inputs (0.4%, p = 0.765), breeding type (0.02%, p = 0.995),

variety (11.8%, p = 1.000) and block (4.6%, p = 0.444) alone, but the breeding x inputs

interaction was responsible for an overall effect on fungal diversity indices, with 10.5% of

explained variance (p = 0.053). The overall multivariate model explained 27.3% of the

variance.

Regarding the direction of effects, we found no effect of the breeding type or of the presence

of inputs alone (Fig. 3). Ancient varieties in the absence of inputs had the highest observed

fungal diversity (not significant after post-hoc correction for multiple comparison, likely due

to the interference of the block effect). For most of the diversity indices, the addition of inputs

induced a decrease in fungal diversity for ancient varieties and, conversely, an increase in

diversity in modern varieties (Fig. 3). The significant interaction was illustrated by the

crossing of all reaction norms, which was only supported statistically for the Chao-1 index.

The PERMANOVA on fungal community beta diversity showed that the breeding type

(2.5%, p = 0.052), the block (5.4%, p =0.011) and variety (34.7%, p = 0.020) had significant

effects, but not the inputs (2.2%, $p = 0.14$). Based on this result, we profiled the structural

changes in fungal communities using a distance-based partial redundancy analysis based on

Bray-Curtis dissimilarities for breeding types alone (the block effect was added as an error

term). The total variance explained was 3.0% ($p = 0.057$), and the first constrained

component (CAP1) and first non-constrained component (MDS1) were explaining 2.76 and

13.50% of variance, respectively (Fig. 2B). As for bacterial community structure, ordination

plots showed a differentiation trend in the fungal community structure between modern and

ancient varieties while inputs addition had no effect.

Network analysis

The PLN models successfully converged into stable networks for the modern varieties (BIC

$R^2 = 0.97$) and ancient varieties (BIC $R^2 = 0.98$). The modern network featured 178 OTUs

(nodes) and 452 edges (correlations) of which 245 were positive and 207 negative. The

ancient network featured 185 OTUs (nodes) and 402 edges (correlations) of which 238 were

positive and 164 negative. The comparison between the two networks revealed a similar level

of node degree (Fig. 4A), but a significantly higher level of centrality in the ancient network

based on average betweenness (either on nodes: $p = 0.05$; and on edges $p = 4.93.10^{-14}$; Fig.

4BC). The positive-to-negative edge ratio was in favour of more positive correlations in the

ancient network (1.40 vs 1.14). When combining both networks and highlighting the position

of common and unique edges found either in modern or ancient varieties, a clear distinction

was found, with a clearly different interaction structure depending on the breeding type (Fig.

4A). In total, this combined network featured 252 unique OTUs, amongst which 108 were

common to both breeding type, while 76 were unique of the ancient varieties and 68 of the

modern varieties. Most OTUs were affiliated to Ascomycota (55%), which clearly dominated

both networks.

Mycorrhizal colonization response to breeding types and inputs

The frequency of mycorhization, F, (number of mycorrhizal root fragments divided by the

total number of observed fragments) was affected neither by the breeding type, the inputs nor

their interaction (Table 5 and S4). Breeding types had an impact on the intensity of

mycorrhizal colonization in root system M (explaining 4.3% of total variance, $p = 0.005$), in

mycorrhizal root fragments m (4.5%, $p = 0.009$), as well as on arbuscule abundance in root

system A ($p = 0.018$) and in mycorrhizal root fragments a (5.6%, $p= 0.002$). Inputs had an

impact only on the intensity of mycorrhizal colonization in root system M (explaining 2.6% of

the total variance, $p= 0.027$) and in mycorrhizal root fragments m (2.6%, $p= 0.046$). The

interaction between breeding type and inputs was only significant on arbuscular abundance in

root system A ($p = 0.032$, Table 5). When performing a PERMANOVA on all mycorrhiza

indices, a significant breeding effect was detected, in favor of higher index values for the

modern varieties (2.8%, $p = 0.008$, Table 2). The overall multivariate model explained % of

the variance. Regarding the direction of effects, mycorrhiza colonization was higher in

modern varieties compared to ancient ones (Fig. 5ABCD) and the application of inputs

decreased the intensity of mycorrhizal colonization in the root system and root fragment (Fig.

5EF).

DISCUSSION

Modern varieties of wheat were selected and generally grown with synthetic chemical inputs

whereas ancient varieties were selected and grown without this kind of inputs, in organic

farming systems. This correlation between variation in genotype and environment prevent to

assess the relative importance of individual factors (E and G) and their interaction (G×E) in

plant-microbiota relationships. In addition, comparison between modern and ancient varieties

are generally made in controlled conditions in the absence of inputs or in nutrient depleted

soils, which prevents any realistic conclusion on the evolution of plant-microbiota

relationships in the field, since modern varieties are grown with inputs. Our results are a first

attempt to quantify these environmental and genotypic effects independently, in the field.

Effects of inputs on plant-microbiota interactions

The presence/absence of inputs – here considered as our environmental conditions "E" – had

a low impact on plant-microbiota relationships. When considered independently of the

breeding type, inputs had only a slight significant effect on microbial alpha-diversity, via a

lowering of specific evenness indices when applied, thus indicating an increased dominance

of some microbial OTUs (Fig. 1 & 3; Tables 2 & 4; Tables S2 & S4). The intensity of

mycorrhizal colonization was higher in the absence than in the presence of inputs (M and m,

Table 5 and Fig. 5). However, this had no consequence on the intensity of arbuscule

development (A and a, Table 5 and Fig. 5), and thus likely no functional impact. The

presence/absence of inputs also had no effect on the overall composition of fungal and

bacterial communities in the rhizosphere (Fig. 2). Taken together, these results support our

first hypothesis that the presence of inputs decreases the microbial diversity in the

rhizosphere. This decrease in diversity in the presence of inputs was expected for several

reasons. First, the addition of N fertilizer can induce a shift in bacterial communities. In line

with our results, Grunert et al. (2019) reported that the addition of an inorganic fertilizer

(struvite) reduced the diversity (Pielou equitability, Shannon and Simpson reciprocal) of

tomato rhizosphere microbiota. In a study with two sites where N was added for 27 and 8

years, the authors found that bacterial community structure was highly responsive to N

additions, but the diversity of bacterial community did not have a consistent response

(Ramirez et al., 2010). As a matter of fact, the shift in bacterial community composition is

mainly observed in soils with low C and N concentrations (Ramirez et al., 2012). The C and

N concentrations in our soil presented a medium value, which could explain the weak effect

of nitrogen fertilization effect. Second, the effect of fungicide could have suppressed some

fungal taxa. It has been shown that commonly used fungicides with foliar application had

moderate but significant effect on the composition of fungal communities in the wheat

phyllosphere (Karlsson et al., 2014). While fungicides are supposed to target specific fungal

pathogens, the impact on fungal communities has been already observed (Esmaeili Taheri et

al., 2015). Third, the effect of herbicide could have modified the weed community, which in

turn could have influenced microbial community, although this mechanism has not been

studied yet, to our knowledge. Our weak observations can also be explained by the fact that

inputs are often reported to affect *bulk soil* microbial communities, whereas *rhizosphere*

microbial communities are more dependent on plant factors than on soil properties (Grunert et

al., 2019).

Effects of breeding on plant-microbiota interactions

The breeding type – here considered as the genotype G – had a strong effect on rhizosphere

microbial communities, which was dependent on the microbial taxa (bacteria or fungi),

meaning that plant-microbiota relationships were differently influenced by breeding from

ancient to modern varieties. When considering the impact of the genotype independently of

the use of inputs, we found that breeding type affected bacterial diversity, with a percentage

of explained variance from 5.9 to 15.6% of the variance according to the index (Table S2) and

7.4% of the variance when all indices were analyzed together (Table 2). The variety inside

each breeding type was the strongest determinant of bacterial diversity, explaining 46.1 to

49.9% of the total variance, and 48.8% when all indices were analyzed together, stressing the

importance of genotype. Conversely, the breeding type had no effect on fungal diversity

(Table S3). However, all mycorrhiza indices (except F) were higher with modern varieties

compared to ancient ones (Table 5, Table 2, Fig. 5). The breeding type also affected slightly

the overall composition of fungal and bacterial communities (Table 3, Fig. 2), but its strongest

effect was noticed on the structure of the rhizosphere microbiota network (Fig. 4). Indeed, the

different breeding types featured completely distinct co-occurrence links amongst the

dominant microbial OTUs, which resulted in a significantly less complex network for the

modern varieties compared to the ancient ones. Therefore, our results partially invalidated the

second hypothesis that modern genotypes have lost their ability to establish interactions with

microbial species, as we evidenced that this interaction still exists and may even be reinforced

in the case of mycorrhiza. However, the interaction with bacteria and other fungal members

has been completely restructured into a simplified form.

Although this hypothesis is supported by several studies in the literature, there is no

consensus. Among supporting results, Valente et al. (2020) found that ancient wheat varieties

were more capable of interacting with beneficial plant growth rhizobacteria than modern

varieties. Moreover, a decrease in the diversity of symbiotic rhizobia associations has been

observed in domesticated legumes compared to wild relatives (Kim et al. 2014; Sangabriel-

Conde et al. 2015). Mutch and Young (2004) reported that the ability to interact with

symbionts was limited for modern pea and broad bean as compared to wild relatives of the

Vicia and *Lathyrus* genera in a non-agricultural soil without inputs. Other studies found more

nuanced results: Leff et al. (2016) reported that neither root nor rhizosphere bacterial

communities were affected by sunflower domestication, but domestication did affect the

composition of rhizosphere fungal communities. Brisson et al. (2019) reported similar

Shannon index for prokaryotic or fungal communities in teosinte, inbred and modern

varieties; they observed that co-occurrence network of microbiota of inbred maize lines'

rhizosphere were significantly closer from those of the teosintes than to the modern hybrids.

Opposed results also exist: Kinnunen-Grubb et al. (2020) reported that bacterial colonization

in the roots of modern cultivars of wheat is faster than in ancestors. This could be explained

by the fact that modern varieties could have lost their ability to specifically select for

beneficial microbial partners compared to ancient varieties (Kiers and Denison, 2008).

Indeed, with six soybean cultivars representing 60 years of breeding, Kiers et al. (2007)

showed that ancient varieties were more performant than modern varieties in maintaining a

high fitness (seed production) when infected with a mixture of effective and ineffective

rhizobia strains. This lower selectivity of modern varieties could explain the associated higher

diversity in rhizosphere microbiota that we observed in the presence of inputs. However, a

meta☐analysis on the effects of breeding on mycorrhizal responsiveness found that varieties

released after 1950 were more mycorrhizal☐responsive (in terms of increased biomass

production) than old varieties (1900-1950) and ancestors (before 1900) (Lehmann et al.,

2012). So the fact that modern varieties establish either adaptive or non-adaptive interactions

with soil microbiota remains an open question (see Ghalambor et al., 2007 for in-depth

discussion). The aim of our experiment was not to evaluate the fitness gain due to plant-

microbiota interactions, but it would be an interesting perspective.

Effects of phenotypic plasticity on plant-microbiota interactions

We speculate that these contradictory results regarding the effect of breeding could be due to

the interaction between plant genotype and the environment. Our experimental design allowed

to assess whether variations in plant-microbiota interaction are affected by the interaction

between the breeding type and the presence of inputs, called phenotypic plasticity. Namely,

phenotypic plasticity describe the capability of a genotype to produce different phenotypes in

response to variation in environmental conditions (Gause, 1947; Bradshaw, 1965). It is an

ubiquitous aspect of organisms. The profile of phenotypes produced by a genotype across

environments is the "norm of reaction" (Schmalhausen, 1949; Stearns, 1989).

Indeed, phenotypic plasticity – here considered as the G×E interaction – had an important

impact on plant-microbiota relationships. It affected several indices of bacterial and fungal

diversity (Tables 1 and 3, Tables S2 and S3, Figs. 1 and 3). For the bacterial community, this

effect of phenotypic plasticity (5.7%) was in the same range as the effect of the breeding type

(7.4%, Table 2). For the fungal community, the effect of the breeding type was not

significant, whereas the effect of phenotypic plasticity was significant and relatively

important when considering all indices together (10.5%, Table 2). However, phenotypic

plasticity had no impact on the intensity of mycorrhizal colonization (Table 5) or on the

overall structure of bacterial and fungal communities (Table 3). This G×E interaction was key

in understanding the variations of diversity: ancient varieties had a decreased bacterial and

fungal diversity in the presence compared to the absence of inputs, whereas modern varieties

kept diverse rhizosphere microbial community in the presence of inputs. This contradicts our

third hypothesis that the lower plant-microbiota association with modern genotypes is

amplified in the presence of inputs. The literature reports many cases of interaction between

genotype and environment for other plant traits and considering plasticity as a trait among

others allows finding out some genetic determinant of plasticity. For example, grain yield

variation in maize was found to result from the environment (a large geographic and climatic

transect of North American) for 43%, genotype for 7% and G×E for 6% (see Gage et al. 2017,

Supp. Fig. 4). The authors found that genomic regions that have experienced changes in allele

frequency due to selection for productivity in temperate conditions explain less G×E variation

than regions in which allele frequency was unaffected by selection. Loci associated with G×E

interaction were mainly located in the regulatory regions of the genome (Gage et al., 2017).

Regarding plant-microbiota interactions, it has been observed in *Boechera stricta*

(Brassicaceae), a perennial wild mustard grown in various sites in North America, that the

G×E effect on microbial community was stronger for the phyllosphere than the rhizosphere,

with an impact on the Shannon index greater than the impact of the genotype itself (but the

opposite was true for Chao1) (Wagner et al., 2016). We found one experiment considering

both wild relatives and cultivated varieties and the manipulation of inputs as an environmental

variation (Shi et al. 2019). Using one single ancient and modern varieties of soybean and rice,

they reported that the rhizosphere microbiota of wild relatives are more affected by a

fungicide than the cultivated ones and that the G×E interaction had a significant effect on

fungal community. However, they did not discuss their results in terms of phenotypic

plasticity.

CONCLUSION

The evolution of the phenotypic plasticity of plant-microbiota interactions has been neglected

up to now by microbial ecologists and geneticists. This knowledge gap makes difficult any

robust conclusion on the fact that modern varieties have lost or improved their ability to

interact with soil microbiota. Our results suggest that modern varieties are less sensitive to the

negative effect of inputs on plant-microbiota interactions than ancient varieties. This could be

explained by a loss of ability to regulate plant-microbiota interactions according to soil

fertility in modern varieties, in accordance the results of Gage et al. (2017) showing that G×E

variation is disproportionately controlled by regulatory mechanisms as compared with other

traits. This is coherent with the fact that modern varieties were not exposed to huge variations

of soil fertility such as ancient ones, as they have been selected and grown in the presence of

inputs. So there were less selective constraints on the modern regarding the ability to fit with

the heterogeneity of soil fertility. However, environmental pollution by synthetic chemical

inputs will likely lead to grow crops with less inputs, so in environments with more

heterogeneous levels of fertility. Integrating phenotypic plasticity in plant-microbiota

interactions as a target in future breeding programs could be very useful to obtain varieties

with high yield, but able to finely adapt their interactions with microbiota to variations in soil

fertility.

ACKNOWLEDGEMENTS

We thank AgroSup Dijon for experimental facilities. We are grateful to Jean-Philippe

Guillemin, Wilfried Queyrel and Etienne Gaujour for their advices in wheat culture and the

management of the experimental site. We thank Tariq Shah for his contribution to the

bibliographical search. We also thank Graines de Noé, especially Hélène Montaz, for

providing the seeds of ancient varieties, as well as RAGT Semences, Secobra, Saaten Union

and Limagrain for providing the modern varieties.

AUTHOR CONTRIBUTIONS

MB conceived the research, MB wrote the paper, CD, EP and MB carried out the experiment,

SJ dealt with sample preparation for sequencing and bioinformatics analysis, SJ did the

statistical analyses and edited the figures and tables, DW, LC and TR did the mycorrhiza

observations, SJ, TR, LC and DW edited and commented the paper.

FUNDING

Samuel Jacquiod was funded by the University of Bourgogne Franche-Comté via the ISITE-

BFC International Junior Fellowship award (AAP3:RA19028.AEC.IS).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

Ai, C., Liang, G., Sun, J., Wang, X., He, P., Zhou, W., & He, X. (2015). Reduced dependence of rhizosphere microbiome on plant-derived carbon in 32-year long-term inorganic and organic fertilized soils. Soil Biology and Biochemistry, 80, 70-78.

Bradshaw, A. D. (1965). Evolutionary significance of phenotypic plasticity in plants. Advances in Genetics, 13, 115–155.

Brisson, V. L., Schmidt, J. E., Northen, T. R., Vogel, J. P., & Gaudin, A. C. (2019). Impacts of maize domestication and breeding on rhizosphere microbial community recruitment from a nutrient depleted agricultural soil. Scientific reports, 9(1), 1-14.

Bulgarelli, D., Garrido-Oter, R., Münch, P. C., Weiman, A., Dröge, J., Pan, Y., ... & Schulze-Lefert, P. (2015). Structure and function of the bacterial root microbiota in wild and domesticated barley. Cell host & microbe, 17(3), 392-403.

Chaluvadi, S., & Bennetzen, J. L. (2018). Species-associated differences in the below-ground microbiomes of wild and domesticated Setaria. Frontiers in plant science, 9, 1183.

Chiquet, J., Robin, S., & Mariadassou, M. (2019). Variational inference for sparse network reconstruction from count data. In International Conference on Machine Learning (pp. 1162-1171). PMLR.

Csardi, G., & Nepusz, T. (2006). The igraph software package for complex network research. InterJournal, complex systems, 1695(5), 1-9.

De Mendiburu, F. (2017). Package 'Agricolae': Statistical Procedures for Agricultural Research, R Package Version 1.2–4.

Dixon, P. (2003). VEGAN, a package of R functions for community ecology. Journal of Vegetation Science, 14(6), 927-930.

Egerton-Warburton, L. M., Johnson, N. C., & Allen, E. B. (2007). Mycorrhizal community

dynamics following nitrogen fertilization: a cross□site test in five grasslands. Ecological

monographs, 77(4), 527-544.

Esmaeili Taheri, A., Hamel, C., & Gan, Y. (2015). Pyrosequencing reveals the impact of

foliar fungicide application to chickpea on root fungal communities of durum wheat in

subsequent year. Fungal Ecology, 15, 73–81. https://doi.org/10.1016/j.funeco.2015.03.005

Falconer, D.S., 1989. Introduction to quantitative genetics. Longman, London.

Farrar, K., Bryant, D., & Cope□Selby, N. (2014). Understanding and engineering beneficial

plant–microbe interactions: plant growth promotion in energy crops. Plant biotechnology

journal, 12(9), 1193-1206.

Gage, J. L., Jarquin, D., Romay, C., Lorenz, A., Buckler, E. S., Kaeppler, S., … de Leon, N.

(2017). The effect of artificial selection on phenotypic plasticity in maize. Nature

Communications, 8, 1348.

García□Palacios, P., Maestre, F. T., Kattge, J., & Wall, D. H. (2013). Climate and litter

quality differently modulate the effects of soil fauna on litter decomposition across biomes.

Ecology letters, 16(8), 1045-1053.

Gause, G. F. (1947). Problems of evolution. Trans. Conn. Acad. Sci., 37, 17–68.

Geisseler, D., & Scow, K. M. (2014). Long-term effects of mineral fertilizers on soil

microorganisms - A review. Soil Biology and Biochemistry, 75, 54–63.

https://doi.org/10.1016/j.soilbio.2014.03.023

Germida, J., & Siciliano, S. (2001). Taxonomic diversity of bacteria associated with the roots

of modern, recent and ancient wheat cultivars. Biology and Fertility of Soils, 33(5), 410-415.

Ghalambor, C.K., McKay, J.K., Carroll, S.P., Reznick, D.N., 2007. Adaptive versus non-

adaptive phenotypic plasticity and the potential for contemporary adaptation in new

environments. Functional Ecology 21, 394–407. https://doi.org/10.1111/j.1365-

2435.2007.01283.x

Grunert, O., Robles-Aguilar, A. A., Hernandez-Sanabria, E., Schrey, S. D., Reheul, D., Van

Labeke, M. C., ... & Temperton, V. M. (2019). Tomato plants rather than fertilizers drive

microbial community structure in horticultural growing media. Scientific reports, 9(1), 1-15.

Hetrick, BAD, Wilson, GWT, & Cox, TS (1992). Mycorrhizal dependence of modern wheat

varieties, landraces, and ancestors. Canadian Journal of Botany, 70 (10), 2032-2040.

Jacquiod, S., Puga-Freitas, R., Spor, A., Mounier, A., Monard, C., Mougel, C., ... & Blouin,

M. (2020). A core microbiota of the plant-earthworm interaction conserved across soils. Soil

Biology and Biochemistry, 144, 107754.

Kapulnik, Y., & Kushnir, U. (1991). Growth dependency of wild, primitive and modern

cultivated wheat lines on vesicular-arbuscular mycorrhiza fungi. Euphytica, 56(1), 27-36.

Karlsson, I., Friberg, H., Steinberg, C., & Persson, P. (2014). Fungicide effects on fungal

community composition in the wheat phyllosphere. PLoS ONE, 9(11), e111786.

https://doi.org/10.1371/journal.pone.0111786

Kiers, E. T., & Denison, R. F. (2008). Sanctions, cooperation, and the stability of plant-

rhizosphere mutualisms. Annual Review of Ecology, Evolution, and Systematics, 39, 215-

236.

Kiers, E. T., Hutton, M. G., & Denison, R. F. (2007). Human selection and the relaxation of

legume defences against ineffective rhizobia. Proceedings of the Royal Society B: Biological

Sciences, 274(1629), 3119-3126.

Kim, B. H., Ramanan, R., Cho, D. H., Oh, H. M., & Kim, H. S. (2014). Role of Rhizobium, a

plant growth promoting bacterium, in enhancing algal biomass through mutualistic

interaction. Biomass and Bioenergy, 69, 95-105.

Kinnunen-Grubb, M., Sapkota, R., Vignola, M., Nunes, I. M., & Nicolaisen, M. (2020).

Breeding selection imposed a differential selective pressure on the wheat root-associated

microbiome. FEMS Microbiology Ecology, 96(11), fiaa196.

Komsta, L., & Novomestky, F. (2015). Moments, cumulants, skewness, kurtosis and related

tests. R package version, 14.

Kusmec, A., Srinivasan, S., Nettleton, D., & Schnable, P. S. (2017). Distinct genetic

architectures for phenotype means and plasticities in Zea mays. Nature Plants, 3, 715–723.

Laitinen, R. A. E., & Nikoloski, Z. (2019). Genetic basis of plasticity in plants. Journal of

Experimental Botany, 70(3), 739–745.

Lambers, H., Mougel, C., Jaillard, B., & Hinsinger, P. (2009). Plant-microbe-soil interactions

in the rhizosphere: an evolutionary perspective. Plant and soil, 321(1-2), 83-115.

Leff, J. W., Lynch, R. C., Kane, N. C., & Fierer, N. (2017). Plant domestication and the

assembly of bacterial and fungal communities associated with strains of the common

sunflower. New Phytologist, 214(1), 412-423.

Lehmann, A., Barto, E. K., Powell, J. R., & Rillig, M. C. (2012). Mycorrhizal responsiveness

trends in annual crop plants and their wild relatives-a meta-analysis on studies from 1981 to

2010. Plant and Soil, 355(1–2), 231–250.

Leiser, W. L., Olatoye, M. O., Rattunde, H. F. W., Neumann, G., Weltzien, E., & Haussmann,

B. I. (2016). No need to breed for enhanced colonization by arbuscular mycorrhizal fungi to

improve low-P adaptation of West African sorghums. Plant and soil, 401(1-2), 51-64.

Lemanceau, P., Blouin, M., Muller, D., & Moënne-Loccoz, Y. (2017). Let the core microbiota

be functional. Trends in Plant Science, 22(7), 583–595.

Lundberg, D. S., Lebeis, S. L., Paredes, S. H., Yourstone, S., Gehring, J., Malfatti, S., ... &

Edgar, R. C. (2012). Defining the core Arabidopsis thaliana root microbiome. Nature,

488(7409), 86-90.

Mei, C., & Flinn, B. S. (2010). The use of beneficial microbial endophytes for plant biomass

and stress tolerance improvement. Recent Patents on Biotechnology, 4(1), 81-95.

Milla, R., Osborne, C. P., Turcotte, M. M., & Violle, C. (2015). Plant domestication through

an ecological lens. Trends in ecology & evolution, 30(8), 463-469.

Mutch, L. A., & Young, J. P. W. (2004). Diversity and specificity of *Rhizobium*

leguminosarum biovar viciae on wild and cultivated legumes. Molecular Ecology, 13(8),

2435-2444.

Peiffer, J. A., Spor, A., Koren, O., Jin, Z., Tringe, S. G., Dangl, J. L., ... & Ley, R. E. (2013).

Diversity and heritability of the maize rhizosphere microbiome under field conditions.

Proceedings of the National Academy of Sciences, 110(16), 6548-6553.

Pérez-Jaramillo, J. E., Carrión, V. J., Bosse, M., Ferrão, L. F., de Hollander, M., Garcia, A.

A., ... & Raaijmakers, J. M. (2017). Linking rhizosphere microbiome composition of wild and

domesticated Phaseolus vulgaris to genotypic and root phenotypic traits. The ISME journal,

11(10), 2244-2257.

Pérez-Jaramillo, J. E., Carrión, V. J., de Hollander, M., & Raaijmakers, J. M. (2018). The

wild side of plant microbiomes. Microbiome, 6(1), 143.

Pérez-Jaramillo, J. E., Mendes, R., & Raaijmakers, J. M. (2016). Impact of plant

domestication on rhizosphere microbiome assembly and functions. Plant molecular biology,

90(6), 635-644.

Ramirez, K. S., Lauber, C. L., Knight, R., Bradford, M. A., & Fierer, N. (2010). Consistent

effects of nitrogen fertilization on soil bacterial communities in contrasting systems. Ecology,

91(12), 3463–3470. https://doi.org/10.1890/10-0426.1

Ramirez, K. S., Craine, J. M., & Fierer, N. (2012). Consistent effects of nitrogen amendments

on soil microbial communities and processes across biomes. Global change biology, 18(6),

1918-1927.

Rodriguez, R., & Redman, R. (2008). More than 400 million years of evolution and some

plants still can't make it on their own: plant stress tolerance via fungal symbiosis. Journal of

experimental botany, 59(5), 1109-1114.

RStudio Team (2020). RStudio: Integrated Development for R. RStudio, PBC, Boston, MA

URL http://www.rstudio.com/.

Sangabriel-Conde, W., Maldonado-Mendoza, I. E., Mancera-López, M. E., Cordero-Ramírez,

J. D., Trejo-Aguilar, D., & Negrete-Yankelevich, S. (2015). Glomeromycota associated with

Mexican native maize landraces in Los Tuxtlas, Mexico. Applied Soil Ecology, 87, 63-71.

Schmalhausen, I. I. (1949). Factors of Evolution: The Theory of Stabilizing Selection.

Blakiston, Philadelphia, PA

Schöler, A., Jacquiod, S., Vestergaard, G., Schulz, S., Schloter, M., 2017. Analysis of soil

microbial communities based on amplicon sequencing of marker genes. Biology and Fertility

of Soils 53, 485.

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., ... & Ideker, T.

(2003). Cytoscape: a software environment for integrated models of biomolecular interaction

networks. Genome research, 13(11), 2498-2504.

Shi, S., Tian, L., Xu, S., Ji, L., Nasir, F., Li, X., ... & Tian, C. (2019). The rhizomicrobiomes

of wild and cultivated crops react differently to fungicides. Archives of microbiology, 201(4),

477-486.

Stearns, S. C. (1989). The Evolutionary Significance of Phenotypic Plasticity. BioScience,

39(7), 436–445. https://doi.org/10.2307/1311135

Szoboszlay, M., Lambers, J., Chappell, J., Kupper, J. V., Moe, L. A., & McNear Jr, D. H.

(2015). Comparison of root system architecture and rhizosphere microbial communities of

Balsas teosinte and domesticated corn cultivars. Soil Biology and Biochemistry, 80, 34-44.

Trouvelot, A., Kough, J.L., and Gianinazzi-Pearson, V. (1986). "Mesure du taux de

mycorhization VA d'un système radiculaire. Recherche de methodes d'estimation ayant une

signification fonctionnelle." In Physiological and Genetical Aspects of Mycorrhizae.

Proceedings of the 1st European Symposium on Mycorrhizae, eds V.Gianinazzi-Pearson and

S.Gianinazzi (Paris: Institut National de la Recherche Agronomique), 217–221.

Turcotte, M. M., Lochab, A. K., Turley, N. E., & Johnson, M. T. (2015). Plant domestication

slows pest evolution. Ecology letters, 18(9), 907-915.

Valente, J., Gerin, F., Le Gouis, J., Moënne□Loccoz, Y., & Prigent–Combaret, C. (2020).

Ancient wheat varieties have a higher ability to interact with plant growth□promoting

rhizobacteria. Plant, Cell & Environment, 43(1), 246-260.

Vierheilig, H., Coughlan, A.P., Wyss, U., Piché, Y., 1998. Ink and vinegar, a simple staining

technique for arbuscular-mycorrhizal fungi. Appl. Environ. Microbiol. 64, 5004–5007.

https://doi.org/10.1128/aem.64.12.5004-5007.1998

Wagner, M. R., Lundberg, D. S., Tijana, G., Tringe, S. G., Dangl, J. L., & Mitchell-Olds, T.

(2016). Host genotype and age shape the leaf and root microbiomes of a wild perennial plant.

Nature communications, 7(1), 1-15.

Weese, D. J., Heath, K. D., Dentinger, B. T., & Lau, J. A. (2015). Long□term nitrogen

addition causes the evolution of less□cooperative mutualists. Evolution, 69(3), 631-642.

Zhu, Y., Christie, P., & Laidlaw, A. S. (2001). Uptake of Zn by arbuscular mycorrhizal white

clover from Zn-contaminated soil. Chemosphere, 42(2), 193-199.

 FIGURE LEGENDS

 Figure 1: Alpha-diversity of the rhizosphere bacterial community (16S rRNA gene amplicon

 sequencing), presented as reaction norms of ancient and modern wheat genotypes to the

 environment modification by inputs. Panel are respectively showing: the richness (A), the

 Chao-1 estimator (B), the ACE estimator (C), the Simpson reciprocal index (D), the Shannon

 index (E) and the Equitability of Pielou (F). Data is shown as the interaction between the

 breeding type and the presence/absence of inputs (w/o: without inputs; w: with inputs).

 Significance was inferred with an ANOVA under the Tukey's HSD post-hoc test for normally

 distributed data (Honest Significant Detection, $p < 0.05$). Non-parametric data were analyzed

 with a Kruskal-Wallis test under False Discovery Rate post-hoc correction (FDR, $p < 0.05$).

 Figure 2: Beta-diversity analysis of the bacterial (A) and fungal (B) communities using

 partial distance-based redundancy analysis (db-RDA). Partial db-RDA were used to output

 the following constrained models: breeding type + Condition (block) (10,000 permutations).

 Figure 3: Alpha-diversity of the rhizosphere fungal community (ITS2 gene amplicon

 sequencing), presented as reaction norms of ancient and modern wheat genotypes to the

 environment modification by inputs. Panel are respectively showing: the richness (A), the

 Chao-1 estimator (B), the ACE estimator (C), the Simpson reciprocal index (D), the Shannon

 index (E) and the Equitability of Pielou (F). Data is shown as the interaction between the

 breeding type and the presence/absence of inputs (w/o: without inputs; w: with inputs).

 Significance was inferred with an ANOVA under the Tukey's HSD post-hoc test for normally

 distributed data (Honest Significant Detection, $p < 0.05$). Non-parametric data were analyzed

 with a Kruskal-Wallis test under False Discovery Rate post-hoc correction (FDR, $p < 0.05$).

 Figure 4: Network analysis of the rhizosphere microbiota associated to the modern and

 ancient wheat varieties. The networks were constructed using dominant OTUs (sum min =

100 counts, occurrence = 25/51 for bacteria, and 25/50 for fungi) and based on partial

correlations obtained with Poisson Log Normal models combining the bacterial and fungal

datasets together (~1 + block + offset). For visualization, both networks were merged into

one, and shown twice with edges that are unique of modern varieties highlighted on the left

network (pink), and edges that are unique of the ancient varieties highlighted on the right

network (khaki). Common link are highlighted in both networks in deep blue. Hidden links

are corresponding to modern/ancient edges respectively not shown in each network. Barcharts

represent average network complexity indices estimated from edges or nodes (Kruskal-Wallis

test, $p < 0.05$).

Figure 5: Mycorrhizal colonization for breeding types and inputs. Panel are respectively

showing: the intensity of mycorrhizal colonization in root system (A), the intensity of

mycorrhizal colonization in root fragment (B), the arbuscular abundance in root system (C),

the arbuscular abundance in root fragment (D) in response to the breeding types (whatever the

presence of inputs) and the intensity of mycorrhizal colonization in root system (E) and the

intensity of mycorrhizal colonization in root fragment (F) in response to inputs (whatever the

breeding type). Significance was inferred with an ANOVA under the Tukey's HSD post-hoc

test for normally distributed data (Honest Significant Detection, $p < 0.05$). Non-parametric

data were analyzed with a Kruskal-Wallis test under False Discovery Rate post-hoc correction

(FDR, $p < 0.05$).

TABLES LEGENDS

Table 1: Analysis of variance and Kruskal-Wallis tests for bacterial alpha diversity indices. F

and chi-square (χ^2) values are given, with asterisks indicating the significance of effects.

"Block" is the factor identifying the three repeated blocks in the experimental design.

"Inputs:Breeding type:Variety" is an error term for the variance explained by the variety

inside each breeding type. For non-parametric data, Kruskal-Wallis tests were done

individually on each factor or factor combinations. The directions of significant effects are

indicated. w, presence of inputs; w/o, absence of inputs; Mod., modern varieties; Anc.,

ancient varieties; ˙, $p < 0.10$; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Table 2: Permutational analysis of variance for bacterial and fungal alpha diversity based on

the six diversity indices (Richness, Chao-1, ACE, Simpson reciprocal, Shannon and

Equitability) and for parameters describing mycorrizal colonization (F, M, m, A and a, see

Materials and Methods for a description). F values are given, with asterisks indicating the

significance of effects. ˙, $p < 0.10$; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Table 3: Permutational analysis of variance for bacterial (16S) and fungal (ITS) beta

diversity. F values are given, with asterisks indicating the significance of effects. The

directions of effects are indicated. Mod., modern varieties; Anc., ancient varieties; ˙, $p < 0.10$;

*, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Table 4: Analysis of variance and Kruskal-Wallis tests for fungal alpha diversity indices. F

and chi-square (χ^2) values are given, with asterisks indicating the significance of effects.

"Block" is the factor identifying the three repeated blocks in the experimental design.

"Inputs:Breeding type:Variety" is an error term for the variance explained by the variety

inside each breeding type. For non-parametric data, Kruskal-Wallis tests were done

individually on each factor or factor combinations. The directions of significant effects are

indicated. w, presence of inputs; w/o, absence of inputs; Mod., modern varieties; Anc.,

ancient varieties; ˙, $p < 0.10$; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Table 5: Analysis of variance and Kruskal-Wallis tests for myccorhiza indices. F and chi-

square (χ^2) values are given, with asterisks indicating the significance of effects. "Block" is

the factor identifying the three repeated blocks in the experimental design. "Series" refers to

the two different batches of staining of mycorrhiza. "Inputs:Breeding type:Variety" is an error

term for the variance explained by the variety inside each breeding type. For non-parametric

data, Kruskal-Wallis tests were done individually on each factor or factor combinations. The

directions of significant effects are indicated. w, presence of inputs; w/o, absence of inputs;

M, modern varieties; A, ancient varieties; ˙, $p < 0.10$; *, $p < 0.05$; **, $p < 0.01$; ***, $p <$

0.001.

Figure 1: Bacterial alpha-diversity analysis

Figure 2: Rhizosphere microbiota beta-diversity analysis

Figure 3: Fungal alpha-diversity analysis

Figure 4: Rhizosphere microbiota network analysis

852   **Figure 5**: Mycorrhiza analysis

855 TABLES

856 **Table 1:** Bacterial alpha-diversity univariate analysis

Factors	Richness (F)	Chao-1 (F)	ACE (F)	Simpson reciprocal (χ^2)	Shannon (χ^2)	Equitability (χ^2)
Inputs	0.063	1.688	0.207	3.914*	1.278	3.622 .
Breeding type	14.81***	5.586*	4.436*	4.077*	9.578**	6.300*
Block	0.043	0.444	0.437	5.233 .	2.611	4.972 .
Inputs x Breeding type	7.363*	3.412 .	5.151*	10.83*	15.64**	12.50**
Inputs x Breeding type/Variety	2.740**	2.528*	2.247*	10.10	14.31	11.64
Direction of effects	Mod. > Anc.	Mod. > Anc. w/o > w in Anc.		w/o > w Mod. > Anc. w/o > w in Anc.	Mod. > Anc. w/o > w in Anc.	w/o > w in Anc.
R^2 (variance explained)	0.695	0.642	0.616	na	na	na

858 **Table 2:** Alpha-diversity and mycorrhiza multivariate analysis

PERMANOVA	R^2 (Bacteria)	R^2 (Fungi)	R^2 (Myco)
Input	0.014	0.004	0.008
Breeding	0.074*	1.7E-05	0.028**
Input x Breeding	0.057*	0.105*	0.011
Input x Breeding/variety	0.478*	0.118	0.188**
Block	0.012	0.046	0.028*
Residual	0.365	0.727	0.737
R^2 (variance explained)	0.635	0.273	0.263

861   **Table 3:** Rhizosphere microbiota beta-diversity analysis

Factors	Bacteria (16S rRNA gene)	Fungi (ITS2 fragment)
Inputs	0.018	0.022
Breeding type	0.027*	0.025*
Block	0.065***	0.054*
Inputs x Breeding type	0.019	0.017
Inputs x Breeding type/Variety	0.308	0.347*
Residual	0.564	0.535

Table 4: Fungal alpha-diversity univariate analysis

Factors	Richness (F)	Chao-1 (F)	ACE (F)	Simpson reciprocal (χ^2)	Shannon (F)	Equitability (F)
Inputs	1.002	0.005	0.048	0.746	4.457*	3.601 .
Breeding type	0.001	0.012	0.004	1.141	0.899	1.124
Block	0.924	1.148	0.677	4.557 .	6.571**	5.987**
Inputs x Breeding type	2.845	5.624*	3.097 .	2.590	3.035 .	1.413
Inputs x Breeding type/Variety	0.298	0.282	0.282	10.40	1.049	1.277
Direction of effects		w/o < w in Mod. w/o > w in Anc.	w/o < w in Mod. w/o > w in Anc.		w/o > w Mod. < Anc. in w/o	w/o>w
R^2 (explained variance)	0.272	0.308	0.243	na	0.578	0.579

865 **Table 5:** Mycorrhiza univariate analysis

Factors	M (F)	m (F)	a (F)	A (χ^2)	F (χ^2)
Inputs	5.03*	4.06*	0.18	0.628	0.872
Breeding type	8.08**	7.12**	5.75*	5.591*	0.122
Series	25.93***	6.97**	13.62***	16.14***	31.70***
Block	3.91*	3.03*	0.97	2.771	4.487
Inputs x Breeding type	0.99	0.62	2.19	8.824*	2.838
Inputs x Breeding type/Variety	2.80***	2.21**	1.86*	10.70	3.211
Direction of effects	Mod. > Anc. w/o > w	Mod. > Anc. w/o > w	Mod. > Anc.	Mod. > Anc.	
R^2 (explained variance)	0.489	0.383	0.355	na	na