Massive export of diazotrophs across the South Pacific tropical Ocean

Sophie Bonnet*, Mar Benavides, Frédéric A.C. Le Moigne, Mercedes Camps, Antoine Torremocha, Olivier Grosso, Céline Dimier, Dina Spungin, Ilana Berman-Frank, Laurence Garczarek, Francisco M. Cornejo-Castillo

*corresponding author
Sophie.bonnet@mio.osupytheas.fr

Abstract

Diazotrophs are widespread microorganisms that alleviate nitrogen limitation in 60% of our oceans, regulating marine productivity. Yet, their contribution to organic matter export has not been quantified, making an assessment of their impact on the biological carbon pump impossible. Here, we demonstrate that cyanobacterial and non-cyanobacterial diazotrophs are massively exported down to 1000 m-depth in the western subtropical South Pacific Ocean (WTSP), accounting for up to 52-100% of the total particulate nitrogen export fluxes. We further demonstrate that small size unicellular diazotrophs (UCYN, 1-8 μm) are exported more efficiently than filamentous diazotrophs (>100-1000 μm) under the form of large (>50 μm) aggregates linked by an extracellular organic matrix. Beyond the WTSP, our data are supported by analysis of the Tara Oceans metagenomes collected in other ocean basins, showing that diazotrophs are systematically detected in mesopelagic waters when present at the surface. We thus conclude that diazotrophs are key players in carbon sequestration in the ocean and need to be considered in future studies to improve the accuracy of current regional and global estimates of export.

Introduction

Nitrogen (N) availability limits primary productivity throughout much of the surface low-latitude ocean[1]. In such nitrogen (N)-limited waters, microbial dinitrogen (N_2) fixation by diazotrophic plankton provides the major source of new N to surface waters[2] that maintains the ocean fertility and, on appropriate timescales, equates export production to the deep ocean[3]. However, the fate of this production remains obscure[4]. No consensus as whether it is exported to the deep ocean or stimulates remineralization in surface waters currently exists. An increasing number of studies have shown that diazotroph-derived N is quickly translocated to non-diazotrophic plankton such as diatoms[5, 6], which eventually contribute to secondary export of organic matter out of the photic zone. Yet, the direct gravitational settling of diazotrophs themselves to the deep ocean has rarely been assessed.

Diazotrophs may associate with sinking particles and contribute to direct export by different mechanisms. The most direct ones include gravitational settling of individual cells/filaments or aggregates. According to Stokes' Law, particle sinking velocity scales with the square of particle size. Therefore, large particles should sink faster and are more likely to reach the deep ocean before being remineralized by bacteria[7]. Aggregation is thus a crucial step for the transport of small phytoplankton species which could export particulate organic carbon (POC) and N (PON) in similar proportion to their production in surface waters[8]. Diazotrophs have diverse morphologies and their size spans several orders of magnitude. Some types such as the free-living unicellular diazotrophic cyanobacteria (UCYN from groups B and C) are small (2-8 μm in diameter), while others such as *Trichodesmium sp.* are filamentous and can form large-sized colonies (>100-1000 μm). In addition*,* some diazotrophs live in symbiosis with calcified (UCYN-A, ~1 μm) or silicified eukaryotes (*Richelia* sp., *Calothrix* sp., >20 μm, forming Diatom-Diazotrophic Associations or DDAs). These dense biominerals may provide ballast enhancing the downward export of these symbioses into the deep ocean. Therefore, the presence of different diazotrophs in surface waters may result in drastically different POC and PON export fluxes via the biological pump. Yet, no field observations have related individual diazotroph groups to the magnitude of downward particles fluxes to date.

Thanks to their inherently ballasted character, DDAs are well known to contribute to particulate matter export[9, 10], and are involved in seasonal peaks of POC export to the deep sea (4000 m) in the North Pacific subtropical gyre[11]. *Trichodesmium* is one of the major contributors to global N_2 fixation[12]. However, *Trichodesmium* are thought to have a limited export capacity and to be preferentially remineralized in the surface layers due to the presence of gas vesicles providing them buoyancy[4, 13, 14]. Yet, some studies have reported the presence of *Trichodesmium* in sediment traps material in the Kuroshio Current[15], the tropical North Atlantic[16], North[17] and Pacific Oceans[18]. Intact filaments and colonies of *Trichodesmium* sp. have also been reported as deep as 3000-4000 m in the

tropical Atlantic, Pacific and Indian Oceans[16, 19], but their contribution to organic matter export has yet
to be quantified.
Theoretically, UCYN may not contribute importantly to POC export fluxes either due to their
small size. Yet, Berthelot et al.[20] reported in experimental mesocosms that primary production
supported by UCYN was twice as more efficient in promoting POC export than production supported
by DDA. During the same study, Bonnet et al.[21] showed that the sinking of UCYN-B and C was driven
by the aggregation of small individual cells (5-6 μm) into larger aggregates (100 to >500 μm) that were
eventually exported and accounted for up to ~20% of the POC flux. Recently, Caffin et al.[18] confirmed
the presence of UCYN-B in sinking particles collected with sediment traps (330 m) throughout the
western tropical South Pacific Ocean (WTSP) (10^3 to 10^7 nifH gene copies L^{-1} of trap material). Finally,
Farnelid et al.[22] observed nifH gene sequences in exported material (150 m) in the North Pacific
subtropical gyre and found that all the diazotroph groups above as well as non-cyanobacterial
diazotrophs were present in the samples.
Taken together, these studies suggest that all diazotroph groups, small or large, free-living or
symbiotic, ballasted or not, have been detected below the photic layer. However, a detailed
examination relating types of diazotrophs to the magnitude of downward particles fluxes and their
contribution is needed to refine our understanding of their role in the magnitude and mechanisms
controlling the biological carbon pump. This is even more timely as climate models predict an
expansion of the oligotrophic gyres[23, 24], where diazotrophs represent a major component of the
plankton biomass and sustain most new primary production[25].
Here, we examine the species-specific fate of diazotrophs in the mesopelagic ocean. We used
an innovative approach consisting of the combined deployment of surface-tethered drifting sediment
traps, Marine Snow Catcher (MSC), and Bottle-net, in which we performed nifH sequencing and
quantitative PCR on major diazotroph groups across the WTSP in parallel with biogenic element
export fluxes quantification (Fig. 1a). This study builds up on prior investigations that have
characterized this region as a hotspot for marine N_2 fixation[26]. We show that all globally-significant N_2-
fixing cyanobacteria and non-cyanobacterial diazotrophs are systematically present in sinking particles
down to 1000 m, where they account for a large fraction of exported PON (52-100%). Small size
UCYN (1-8 μm) are exported more efficiently than large filamentous diazotrophs under the form of
large (>50 μm) aggregates linked by an extracellular matrix. Globally, our analysis of the Tara Oceans
metagenomes confirms that diazotrophs are always detected in mesopelagic waters when present in
surface waters, revealing that the gravitational settling of diazotrophs is an important pathway for
diazotroph-derived export production, influencing the ability of our ocean to sequester carbon.
92
93 ## Results and discussion
94
Surface conditions
The study region was characterized by typical oligotrophic waters (chlorophyll concentrations <0.15 μg
L^{-1}, DCM 100-180 m) east and west of Tonga, corresponding to the first group of stations S01, S02,
S07 and S08 (Fig. 1a). The second group of stations (S03, S04, S05, S10, S11) was located in
mesotrophic waters (chlorophyll >0.15 μg L^{-1}, DCM 70-90 m) in the vicinity of the Tonga volcanic arc
(Fig. 1a). Surface (0-50 m) nitrate concentrations were consistently close or below the detection limit
(0.05 μmol L^{-1}) throughout the transect, while phosphate concentrations were typically 0.1 μmol L^{-1} at
oligotrophic stations and depleted down to detection limit (0.05 μmol L^{-1}) at mesotrophic stations, likely
due to consumption by higher plankton stocks (Fig. S1). Seawater temperature ranged from 23.1 to
27.3 °C in the mixed layer (0-15 m to 0-60 m) determined according to de Boyer Montégut et al.
(2004)[27] (Fig. S1).
The abundances of key diazotroph groups indicated that UCYN-A1 were present at high
abundances throughout the photic layer (0-130 m) of the transect (average 3.2 x 10^8 nifH gene copies
L^{-1}) (Fig. 1b), peaking at oligotrophic stations (average 1.1 x 10^9 nifH gene copies L^{-1}). Trichodesmium
was the second most abundant group, particularly in the vicinity of Tonga (average 1.3 x 10^7 nifH gene
copies L^{-1}), followed by UCYN-B (5.7 x 10^5 nifH gene copies L^{-1}). UCYN-C and Gamma-A (a non-
cyanobacterial diazotroph) were also detected throughout the transect albeit at lower abundances (10^2
to 10^4 nifH gene copies L^{-1}).
113
How efficient are different diazotrophs groups to export?
We first examined the nifH gene community composition of particles collected in drifting sediment
traps located at 170 m, 270 m and 1000 m at stations S05M and S10M (Fig. 1a). On average, 32% of
the retrieved Amplicon Sequence Variants (ASV) corresponded to cyanobacteria genera over both
stations (Fig. 2). The most abundant genus was Trichodesmium sp., but ASVs related to

Crocosphaera watsonii (UCYN-B), *Candidatus* Atelocyanobacterium thalassa (UCYN-A) and
Katagnymene spp. were also identified (Fig. 2). Genera related to DDAs (*Richelia* spp., *Calothrix* spp.)
contributed little to the percentage reads. Besides autotrophic diazotrophs, 68% of ASVs were
affiliated to non-cyanobacterial diazotrophs including Alpha-, Beta-, and Gammaproteobacteria at both
stations.
Amplicon sequencing provides relative abundances, which are not necessarily indicative of
the abundance of the target microorganism in the natural environment[28, 29]. Hence, to assess the
export capacity of individual diazotroph groups, we quantified the abundance of five groups spanning
different forms, sizes, lifestyle (symbiotic or not), using qPCR assays (UCYN-A1 symbiosis, UCYN-B,
UCYN-C, *Trichodesmium* and Gamma-A; see Methods). DDAs were not quantified as they were
almost not detected on microscopy inspections, pointing to their rarity in those waters at the time of
the cruise. All five groups were detected at abundances ranging from 10^3 to 10^8 *nifH* gene copies L^{-1} of
traps material, the highest being reported for UCYN-A1 and the lowest for UCYN-C (Fig. 3a,b). The
diazotrophs assemblage exported to sediment traps generally reflected that of the photic layer,
although some diazotrophs seemed to sink more efficiently than others: the highest diazotroph export
fluxes were measured for UCYN-A1 at both sites and all depths of traps deployment ($5.0 \pm 1.1 \times 10^8$
to $8.5 \pm 2.0 \times 10^9$ *nifH* gene copies m^{-2}d^{-1}), followed either by UCYN-B ($2.2 \pm 0.9 \times 10^7$ to $7.5 \pm 1.4 \times$
10^7 *nifH* gene copies m^{-2}d^{-1}) or *Trichodesmium* ($2.5 \pm 1.9 \times 10^6$ to $1.2 \pm 0.3 \times 10^8$ *nifH* gene copies m^{-2}
) depending on station and depth (Fig. 3a,b). Gamma-A and UCYN-C were also exported, albeit at
lower rates (~10^5 and ~10^6 *nifH* gene copies m^{-2}d^{-1}, respectively).
Specific export turnover rates provide information on the rate at which diazotrophs are "lost"
from the photic layer due to export (Turnover rate (d^{-1}) = export flux/abundance in the photic layer).
The range of calculated rates ($0.7 \pm 1.2 \times 10^{-5}$ to $11.2 \pm 1.9 \times 10^{-3}$ d^{-1}) spans through two orders of
magnitude and varied depending on the diazotroph group, depth and station considered. On average
(over all stations and depths), the export turnover rate of UCYN ($5.5 \pm 3.1 \times 10^{-3}$ d^{-1}) was ca. four
times higher than that of *Trichodesmium* ($1.3 \pm 2.2 \times 10^{-3}$ d^{-1}) (Fig. 3c), pointing to preferential export
of UCYN groups relative to the filamentous *Trichodesmium*. Among UCYN groups, the highest export
turnover rate was measured for UCYN-B, followed by UCYN-C and UCYN-A1, although differences
were not significant between groups (Mann-Whitney test, $p < 0.05$). The export turnover rate of
Gamma-A (average $1.8 \pm 1.5 \times 10^{-3}$ d^{-1}) was intermediate between that of *Trichodesmium* and UCYN
groups.
Epifluorescence microscopy confirmed the presence of phycoerythrin-containing UCYN-B and
UCYN-C-like cells in sediment trap samples (Fig. 4a-f). Scanning electron microscopy revealed that
they were recurrently found embedded in large organic aggregates, or organized into clusters of tens
to hundreds of cells linked by an extracellular matrix (Fig. 4g-l), which was further confirmed by Alcian
blue straining (Fig. S2). *Trichodesmium* was also observed in all samples, mostly as free filaments,
but intact colonies were observed in sediment trap samples especially at 1000 m at both stations.
Gamma-A and UCYN-A1 cannot be visualized by these techniques and are thus assessed solely on
the basis of qPCR counts (above).
Pigment concentrations measured in sediment trap samples at both stations indicate that total
pigment concentrations and diversity decreased with depth. Apart from the degradation pigments
phaeophorbide and phaeophytin, zeaxanthin, a biomarker of cyanobacteria predominated at all depths
(Fig. S3). The Chlorophyll *a* : phaeopigments ratios were elevated (average 1.8, range 0.8-3.),
indicative of fresh organic matter in sediment trap samples.
163
Contribution of diazotrophs to PON export fluxes
Sediment trap-based PON export fluxes at 170 m, 270 m and 1000 m ranged between 1.6 ± 0.2 and
6.1 ± 1.2 mg N m^{-2} d^{-1} at S05M and 5.2 ± 0.1 and 13.4 ± 1.3 mg N m^{-2} d^{-1} at S10M (Table 1). Fluxes
attenuated with depth, with transfer efficiencies of 0.02 and 0.03 between 170 m and 1000 m at both
stations. The N export efficiency (Ne-ratio, the amount of PON export at the base of the euphotic zone
over the integrated N_2 fixation rate) was 18% and 53% at stations S05M and S10M. The C:N ratios in
exported material was lower (5.2) than that of the surface (7.5, integrated value over the photic layer)
at station S10M, pointing to a preferential export of PON relative to POC.
The proportion of PON attributed to diazotrophs and collected by sediment traps was
evaluated based on qPCR results and estimated PON content per diazotroph (see Methods). The
PON flux attributed to diazotrophs ranged from 0.9-2.3 mg N m^{-2} d^{-1} at S05M and from 1.2-2.7 mg N
m^{-2} d^{-1} at S10M, accounting for 18% (170 m) to 100% (1000 m) and 9% (170 m) to 52% (1000 m) of
the total PON export flux at stations S05M and S10M, respectively (Table 1). *Trichodesmium* was the
major contributor of the PON attributed to diazotrophs at 1000 m at both stations, whereas UCYN
dominated at shallower depths (Table S1). Despite non-cyanobacterial diazotrophs were present in

exported material (68% of sequences), their C and N composition remains unknown due to the poor
availability of cultured representatives. Thus, quantifying their contribution to export fluxes is not
possible at this stage. In addition, sequencing data also revealed the presence of *Katagnymene* spp.,
which are large (>100-1000 μm) and likely have high C and N content. Hence the contribution of
diazotrophs to exported PON reported here likely represent a lower-end estimate.
184
The fate of diazotrophs in the mesopelagic ocean
We further assessed the fate of the exported diazotroph community by collecting fresh particles in
mesopelagic waters using the Marine Snow Catcher (MSC, see Methods). In essence, the MSC
allows separating non-sinking particles from slow sinking and fast sinking particles[30] (see Methods).
These different fractions can hence be studied individually. MSC samples were collected at the same
depths as those of sediment traps (170, 270, 1000 m) at stations S05M and S10M and in addition at
seven other stations (Fig. 1a).
The diazotroph community composition (based on *nifH* amplicon sequencing) in the various
fractions generally mirrored that of traps at stations S05M and S10M with 32-54% of sequences
affiliated with cyanobacteria and a prevalence of *Trichodesmium* (Fig. 5). This was also the case at
stations S03, S04, S07 and S11, but not at stations S01, S02 and S08, where sequences assigned to
Candidatus Atelocyanobacterium thalassa and *Crocosphaera watsonii* were generally more abundant
than *Trichodesmium* sequences. As in sediment traps, a large number of *nifH* sequences (46-58%)
were affiliated to non-cyanobacterial diazotrophs, whose composition was overall consistent with that
of the traps, although some classes not detected in sediment traps (Epsilonproteobacteria and
Clostridia) were present in all MSC fractions. In general, the total number of *nifH* gene reads was
consistently 10-25% higher in the slow and fast sinking fractions compared to the suspended fraction
at all stations. This trend was mainly driven by non-cyanobacterial sequences that are suspected to be
attached to particles[31], and to a lesser extent by sequences affiliated with cyanobacteria.
As in sediment traps, five diazotroph groups were quantified in the suspended, slow and fast
sinking fractions to more finely assess their sinking dynamics over short time scales (h). Overall,
UCYN-A1 and *Trichodesmium* were the most abundant groups in the mesopelagic MSC samples (170
m to 1000 m), and their numbers varied in parallel with the abundance of their population in the photic
layer (Fig. 6). UCYN-A1 were the most abundant at oligotrophic stations (S01, S02, S07, S08,
chlorophyll <0.15 μg L^{-1}), while *Trichodesmium* peaked at mesotrophic stations (Chlorophyll a >0.15
μg L^{-1}, Fig. 1). Conversely, the abundance of UCYN-B, UCYN-C and Gamma-A in mesopelagic waters
was poorly correlated with their photic layer abundance.
The MSC suspended fraction accounted for the most abundant pool of diazotrophs at all
stations, followed by either the slow or the fast sinking pools, depending on groups (Fig. 6a-e). At the
mooring stations S05M and S10M, the total concentration of diazotrophs and the diazotroph
community composition remained generally constant at the three sampled depths (170 m, 270 m and
1000 m), consistent with the results observed in sediment traps. To evaluate the export potential of
each individual diazotroph group, we calculated the ratio of sinking versus non-sinking cells (the sum
of the slow and fast sinking fractions over the total). The proportion of sinking cells varied widely
across the transect, but was the highest for UCYN-A1 (32 ± 33 %), UCYN-C (31 ± 38 %), Gamma-A
(29 ± 19 %) and UCYN-B (27 ± 11 %), and the lowest for *Trichodesmium* (18 ± 20 %) (Fig. 6f). Among
the sinking fractions, UCYN-B and Gamma-A were generally equally distributed among the slow and
fast sinking fractions, whereas the majority of *nifH* gene copies were found in the fast sinking fraction
for UCYN-A1 (59%), UCYN-C (62%) and *Trichodesmium* (67%). Taken together, these results
indicate that over short time scales (2 h, the conventional settling of particles following MSC
deployment[30]): i) small UCYN (few μm) generally sink more efficiently than large *Trichodesmium*
(>100 μm), ii) when sinking, *Trichodesmium* sinks fast, iii) UCYN-A1 and UCYN-C sink faster than
UCYN-B and Gamma-A. A detailed imaging study performed on 170 m MSC samples at the mooring
stations S05M and S10M indicates that the 50-80% of phycoerythrin-containing UCYN in the fast
sinking fraction were organized into aggregates of tens to >250 cells measuring 30 to >100 μm, while
the majority (60-95%) of UCYN were free living in the suspended fraction (Figure S4). This indicates
that under minimum turbulent agitation, UCYN quickly (within 2 h) form aggregates large/dense
enough to sink.
Concentrations of PON averaged across all stations 1.85 ± 0.49 μg L^{-1}, 0.25 ± 0.13 μg L^{-1} and
0.26 ± 0.08 μg L^{-1} within the suspended, slow sinking and fast sinking fractions (Fig. S5a). PON in the
fast sinking fraction contributed 11 ± 2 %, while the slow sinking and suspended fractions contributed
11 ± 4 % and 79 ± 5 % in terms of total PON. Thus, ~22% of the PON was sinking out of the upper
part of the MSC within 2h during our study. We converted transparent expolymeric particles into C
(TEP-C), that revealed higher concentrations in the suspended fraction compared to the fast sinking

fraction at all stations (TEP-C concentrations were null in the SS fraction) (Fig. S5b). The TEP-C:POC
ratio was generally higher in the suspended fraction compared to the fast sinking. This indicates that,
despite UCYN were embedded in TEPs in FS samples, cells rather than TEPs were the major
contributor of the POC pool in the FS fraction.
Finally, we quantified diazotrophs on vertical profiles spanning the water column between 200
m and 2000 m by using a Bottle-net mounted on the CTD rosette frame at six stations across the
transect (Fig. 1a). The bottle-net consists of a 20-µm conical plankton net housed in a cylindrical PVC
pipe[19]. The top cover is opened at the desired bottom depth (2000 m) of the tow, remains opened
during the ascent of the rosette, and closed again at the upper depth (200 m) of the water column to
be sampled. This results in one integrated sample of 200 m to 2000 m per deployment. Overall, bottle-
net tows confirm that diazotrophs are consistently present in this deep ocean layer, with
concentrations averaging 1.4×10^7 *nifH* gene copies m^{-2}. Among the groups targeted by qPCR, the
community was primarily dominated by UCYN-A1 (63% on average over all sampled stations) and
Trichodesmium (27%), and secondarily by UCYN-B (9%) (Fig. 7), generally mirroring the diazotroph
community in surface water (these numbers are conservative and may be underestimated as some
individual UCYN pass through the 20 µm mesh net of the bottle-net). The 2000-200 m stock was
relatively constant among stations, except at the most oligotrophic station S08, where it was lower by
two orders of magnitude than that of other stations.
257
Beyond subtropical south Pacific waters
To assess whether the sinking of globally-distributed diazotrophs down to mesopelagic waters is a
widespread feature, we explored the presence of diazotrophic cyanobacterial genomes using *Tara*
Oceans metagenomes collected from other ocean basins[32, 33]. Among all the stations for which
metagenomic samples from both surface and mesopelagic were available, we selected those ones
showing significant abundances of *nifH* metagenomic reads in surface waters (5 m)[32]. This led to a
total of five stations located in the South Atlantic and North and South Pacific oceans where we also
explored the presence of diazotrophs, at the genome level, in the mesopelagic (300 to 800 m) (Fig. 8).
Recruitment of metagenomic reads against representative genomes of the diazotrophic community
(*Trichodesmium, Richelia,* UCYN-B and UCYN-A1 and UCYN-A2; Table S2) across different size-
fractions in surface and mesopelagic waters show that diazotrophs in general were systematically
detected in mesopelagic waters at all five stations (Fig. 8; Table S3 ; see Methods). As in the WTSP,
every specific group was detected at depth when present in surface, except when abundances in
surface were very low (<6.7 reads/total 100,000 reads). Accordingly with their cell size,
Trichodesmium, *Richelia*, UCYN-B and UCYN-A2 reads were recovered in the size fraction >3 µm at
mesopelagic depths, whereas the UCYN-A1 symbiosis was recovered in the 0.2-3 µm and 0.8-3 µm
size fractions. Overall, this metagenomic analysis shows that the export of diazotrophs through
gravitational settling is not restricted to the WTSP and is a widespread phenomenon in the tropical
ocean.
277
Biogeochemical implications
Excluding silicified DDAs[11], diazotrophs have seldom been regarded as important contributors to
organic matter export. Yet, our results provide clear evidence that they are present and ubiquitous in
the mesopelagic ocean and they contribute significantly to the PON export fluxes in the WTSP. They
present, however different behaviors.
We showed that diazotroph cell sizes is not necessarily a key variable controlling the
diazotrophs' ability to sink out the euphotic zone. Small UCYN (1-8 µm) displayed the highest export
turn-over rates ~10^{-3} d^{-1}, in the range or higher than those reported for ballasted phytoplankton groups
such as diatoms and coccolithophores[34]. Within UCYN, UCYN-A1 clearly dominated the diazotroph
community targeted by qPCR in mesopelagic waters of the WTSP. Although the UCYN-A1 symbiosis
has been detected once in mesopelagic waters of the South Atlantic Ocean[32, 35], their capacity to
leave the photic layer and sink throughout the mesopelagic zone down to 1000 m (Fig. 3, 8) has not
been thoroughly explored before. As UCYN-A rank among the most abundant diazotrophs in the
global ocean[12] and span from tropical to polar waters[36, 37], we suggest that they potentially play a
significant contribution to ocean organic matter sequestration. Some UCYN-A ecotypes (e.g. UCYN-
A2) live in obligate symbiosis with a calcified coccolithophore[38], hence their export is likely enhanced
by this ballast mineral increasing the density of sinking particles to which they are associated. In future
studies, better constrains on UCYN-A sedimentation rates and aggregation processes would be of
primary importance to assess their role in particle fluxes and cycling.
UCYN-B and -C are free-living unicellular cyanobacteria and their export pathway therefore
does not rely on mineral ballasted hosts. Our results indicate that the sinking of such small cells is

made possible through the aggregation of UCYN into large-sized aggregates (30->100 μm) of tens to
hundreds of cells, large/dense enough to sink, in accordance with a previous mesocosm study[21]. The
presence of UCYN-B in sediment traps is also in accordance with previous reports of *Crocospheara*
watsonii sequences at station ALOHA either just below the euphotic zone[22], or as deep as 4000 m[39].
The visualization of our sediment trap samples shows that they can be subsequently embedded in
mixed aggregates with other cells and debris (Fig. 4, S2). The UCYN-B large (>4 μm) ecotype, which
was dominating in this study (data not shown), produces large amounts of C-rich transparent
exopolymeric particles (TEP) at rates one to two order of magnitude higher than that of diatoms and
coccolithophorids[40], probably in response to nutrient limitation and excess light[41]. In cultures of
Crocosphaera watsonii, TEP account for ~22% of the particulate C pool[40]. Hence, TEP produced by
UCYN-B do not only provide a matrix for the formation of large aggregates, but may also account for a
significant fraction of C export. TEP are indeed greatly enriched in C relative to N (C:N ~25[40]).
Trichodesmium was generally the second contributor to the diazotroph community targeted by
qPCR in mesopelagic waters after UCYN-A1, which contradicts the common assertion that they are
entirely remineralized in the euphotic layer[4, 13]. Several potential mechanisms could explain the
presence of *Trichodesmium* in mesopelagic waters. *Trichodesmium* colonies can migrate vertically to
exploit the deep phosphate stock (Villareal and Carpenter 2003) (the phosphacline was around 200 m
in our study region at the time of the cruise). According to this theory, they overcome their positive
buoyancy by fixing C that result in carbohydrate ballasting. However, Walsby (1978)[42] observed that
100% of the gas vacuoles of *Trichodesmium erythraeum* (the most abundant species in this study)
collapse at depths between 105 m and 120 m, resulting in a loss of their buoyancy. Hence,
Trichodesmium could be locked in a persistent and irreversible downward trajectory. Alternatively,
Berman-Frank et al.[43, 44] have shown that programmed cell death induces internal cellular degradation,
gas-vacuole loss and increased production of TEP, resulting in an increase in the vertical flux of
Trichodesmium[45]. Whatever the mechanism behind the vertical flux of *Trichodesmium*, we report here
higher abundances at 1000 m compared to those at shallower depths at both stations, consistent with
microscopic observations showing intact colonies at 1000 m. This likely results from a spatio-temporal
decoupling between production and export[46]. Despite sinking less efficiently than UCYN in the
aggregates, they are much larger and contain more C and N per filament than UCYN[12], and
accounted for a significant fraction of the diazotroph-attributed PON export in this study, especially at
1000 m (Table S1), which is in accordance with previous studies reporting intact *Trichodesmium*
colonies down to 3000-4000m[16, 19].
More than 90% of the organic matter sinking below the euphotic zone is respired before it
reaches a depth of 1000 m[7]. Fast sinking particles will therefore theoretically make a greater
contribution to the deep ocean flux than slow sinking particles, since the latter will be rapidly recycled
at shallow depths. The relative similarity of the taxonomic diazotroph community composition in
mesopelagic waters sampled by three independent sampling approaches (traps, bottle-net, and MSC)
compared to the diazotrophic composition of the euphotic layer suggests a rapid export mode of
diazotrophs. This is further confirmed by i) the high proportion of diazotroph groups quantified in the
MSC fast sinking fraction, ii) the whole cells and colonies and high Chlorophyll *a* : pheopigments ratios
in sediment traps, indicative of healthy phytoplankton (Fig. 4; S2). Finally, iii) while PON export fluxes
were attenuated with depth in our study, which is a classical feature in the oligotrophic ocean[47], the
diazotroph export fluxes were not, suggesting a high transfer efficiency to the deep ocean, and thus
sinking velocities high enough to escape short-term remineralization. Indeed, a parallel study
performed during the same cruise[48] reveals significant N$_2$ fixation rates in sediment traps and MSC
samples, suggesting that part of the diazotroph community in mesopelagic waters sunk fast enough to
remain alive at mesopelagic depths. Bar-Zeev et al.[45] reported sinking velocities of *Trichodesmium*
erythraeum of ~200 m d^{-1}. More recently, Ababou et al.[49] reported sinking velocities of *Trichodesmium*
erythraeum and UCYN aggregates formed in roller tanks of 92 ± 37 m d^{-1} and 333 ± 176 m d^{-1},
respectively. We calculated that *Trichodesmium* sinking at those rates (92-200 m d^{-1}) would take 5 to
10 days to reach 1000 m, whereas UCYN would take 3 days, which would be compatible with finding
active cells observed at those depths[48].
351
352　Conclusion
Our findings challenge the common assumption that the fate of diazotroph-derived production is
constrained to the surface layer. They provide a significant advance in defining the role of diazotrophs
on an important planetary carbon flux, the biological carbon pump, and bring new insights into the
species-specific export of diazotrophs in the ocean, revealing the previously-unseen role of UCYN and
Trichodesmium into the overall export fluxes. Diazotrophs being not restricted to the tropical ocean, it

would be interesting in future studies to explore the direct gravitational export of diazotrophs in
temperate and polar waters.
Direct export through gravitational settling of diazotrophs is most likely only the tip of the
iceberg and diazotrophs can potentially be exported through secondary pathways: diazotrophs release
in seawater 10–50% of recently fixed N_2 (referred to as Diazotroph Derived, DDN) as NH_4^+ and
dissolved organic N (DON)[50, 51, 52]. This DDN is potentially available for assimilation by the surrounding
phytoplanktonic communities, supporting their growth and leading a potential secondary (indirect)
export pathway of diazotroph-derived OC[6, 21, 53]. Moreover, DDN is also transferred to zooplankton,
which produces dejections termed fecal pellets, that sink fast and play a major role in OC export to the
deep ocean[54]. In future studies, there is an urgent need to develop appropriate approaches to
decipher diazotroph export pathways (direct *vs* indirect) if we are to understand the role of N_2 fixation
in the biological carbon pump. This is an especially pressing question given that current climate
models predict an expansion of the oligotrophic gyres (60% of our oceans)[23, 24], where diazotrophs
thrive. N_2 fixation will thus likely be crucial to supporting primary productivity and export in the future
ocean.
373
Methods
Samples were collected during the TONGA cruise (Fig. 1, DOI: 10.17600/18000884) in the tropical
South Pacific Ocean onboard the R/V *L'Atalante* from November 1 to December 5 2019. We collected
suspended and sinking particles in the mesopelagic zone using three complementary devices:
surface-tethered drifting sediment traps[55], MSC and Bottle-net. Additionally, water column samples
were collected from the euphotic layer using Go-Flo bottles mounted on a trace metal clean rosette to
quantify the stocks of the major groups of diazotrophs in the photic layer.
381
Satellite products and photic layer sampling. Satellite-derived surface chlorophyll *a* concentration
during the TONGA cruise (November 2019) was accessed at http://oceancolor.gsfc.nasa.gov (MODIS
Aqua, 4 km, 8-days composite, level 3 product). Vertical 0-150 m depth profiles were performed at
each station using a trace metal clean titanium rosette of Go-Flo bottles equipped with a fluorometer
and temperature, conductivity and oxygen sensors. Seawater samples were collected from 5 depths
(75, 50, 20, 10, 1% surface irradiance levels) to quantify the stock of major groups of diazotrophs in
the photic layer by quantitative PCR (qPCR) as described below. Additionally, primary production and
N_2 fixation rates were measured in triplicates at the same six depths at the locations of the traps
deployments (S05M and S10M), as described in [56, 57]. N_2 fixation rates were measured at two
occasions (on day 1 and day 3 of the traps deployment). The average value between both profiles was
used to calculate the e-ratios (POC export:primary production) and the contribution of N_2 fixation to
primary production rates.
394
Sediment trap deployment and sample analyses. A surface tethered mooring line (~1000 m long)
was deployed at stations S05M (21.157°S;175.153°W, 5 days) and S10M (19.423°S;175.133°W, 4
days). The line was equipped with sediment traps (KC Denmark®) at 3 depths: 170 m (corresponding
to the base of the photic layer), 270 m and 1000 m. Each trap was composed of four particle
interceptor tubes (PITs) mounted on a cross frame (collecting area of 0.0085 m^2, aspect ratio of 6.7).
Two of the tubes were used for this study: one tube was used for biogeochemical analyses (hereafter
referred to as 'Biogeo' tube), and one for microbiological analyses ('Microbio' tube). Prior to
deployment, the tubes were filled with 0.2 µm filtered seawater with added saline brine (50 g L^{-1}).
Borate-buffered formalin (5%) was also added to the Biogeo tube to prevent in situ microbial
decomposition[55]. After recovery, the density gradient was visually verified, and the PITs were allowed
to settle for 2 h before the supernatant seawater was carefully removed with a peristaltic pump. The
remaining water containing the sinking material was transferred to a chlorohydric acid-washed
container, while being screened with a 500 µm mesh to remove swimmers (zooplankton that actively
entered the traps)[58]. Subsequently, samples were split into 12 aliquots. A triplicate set of aliquots were
filtered onto 25-mm diameter Supor filters for *nifH* sequencing and *nifH* qPCR as described below.
Another triplicate set of aliquots were filtered onto 25-mm diameter combusted (4h, 450°C) glass
microfiber filters (Whatman GF/F), which were subsequently dried for 24 h at 60°C, pelleted and from
which particulate N (PON) and C (POC) were analyzed by EA-IRMS (Elemental Analyzer-Isotope
Ratio Mass Spectrometry) using an Integra 2 (Sercon) mass spectrometer. Lastly, a triplicate set of
aliquots was filtered on GF/F filters for further pigment analyses[59], and another triplicate filtered on 1
µm polycarbonate filters for microscopic analyses (see below for methods).
416
MSC deployment

Suspended and sinking particles were sampled using a MSC at 3 depths at the mooring stations
S05M and S10M (170 m, 270 m and 1000 m), and at 200 m, 400 m and 1000 m at S8. Additionally,
the MSC was deployed at 200 m at S01, S02, S03, S04, S05V, S07, S10V and S11 (Fig. 1). The MSC
is a large volume water sampler (100 L) that collects sinking particles with minimal turbulent
agitation[60]. Upon recovery, the MSC is placed on deck for 2 h while any particles present settle onto
the base of the bottom 7 L chamber. After each recovery, the MSC is conventionally left on deck for 2
h allowing particles to settle[30]. Fast sinking particles were thereby deposited collected in a dedicated
plate at the bottom of the MSC, slow sinking particles were collected from the 7-litre compartment
above the plate, and the non-sinking (or suspended) fraction was collected from the upper part of the
MSC[30]. All three fractions were collected separately (4 L for the suspended, 1.2 L for the slow sinking).
For the fast sinking fraction, the entire volume of the plate was sampled (280 to 310 mL). Subsamples
from each fraction were filtered for PON and POC quantification, TEP quantification according to
Passow and Alldredge (1995)[61], microscopic observations, and *nifH* sequencing, *nifH* qPCR as
described below. We used the formula described in Riley et al., (2012)[30] to obtain values associated
with fast or slow sinking particles only, quantitatively removing the contribution of other fractions.
433
Bottle-net deployment
Vertical 2000-200 m profiles were done using a bottle-net mounted on the CTD rosette frame at
stations S03, S04, S05M, S07, S08, S10M, S11 and S12. The bottle-Net consists of a 20-µm conical
plankton net housed in a cylindrical PVC pipe[19]. It is lowered with the top cover closed, which is
opened at the desired bottom depth (Db, 2000 m) of the tow, remains opened during the ascension of
the rosette, and closed again at the upper depth (Du, 200 m) of the water column to be sampled. This
results in one integrated sample of 2000 to 200 m per deployment. Once on deck, the bottle-nets were
gently rinsed with filtered seawater before retrieving the sample from the collector. At each station,
samples were split into aliquots that were processed for were analyzed for *nifH* qPCR. Sampled
volume was estimated as the product between the cross-sectional area of the mouth of the bottle-Net
(7.5 cm, aspect ratio of 4) and the vertical distance covered by the device from the start of the
ascension to the closure of the top cover (1800 m). Blank casts were performed with the bottle-net
closed during the entire cast to assess for potential contaminations, and blanks were subtracted.
447
nifH **gene sequencing and bioinformatics**
The *nifH* gene was sequenced from a total of 71 samples: 18 samples collected from sediment traps
deployed at 170 m, 270 m and 1000 m at S05M and S10M, and 53 samples collected from the
suspended, slow sinking and fast sinking fractions of particles collected with the MSC. To that end,
DNA was extracted using the DNeasy Plant Mini Kit (Qiagen, Courtaboeuf, France) with additional
freeze-thaw bead beating and proteinase K steps before the kit purification[62]. Triplicate nested PCR
reactions were conducted using degenerate *nifH* primers[63]. The PCR mix was composed of 5 µL of 5X
MyTaq red PCR buffer (Bioline), 1.25 µL of 25 mM $MgCl_2$, 0.5 µl of 20 µM forward and reverse
primers, 0.25 µL Platinum Taq and 5 µL of DNA extract (1 µL on second round). The reaction volume
was adjusted to 25 µL with PCR grade water. Triplicate PCR products were pooled and purified using
the Geneclean Turbo kit (MP Biomedicals). Partial adapters were added by ligation at the sequencing
facility (Genewiz) and llumina MiSeq 2x300 paired end sequenced. Demultiplexed paired-end
sequences were dereplicated, denoised, assembled and chimeras discarded using the DADA2
pipeline[64]. Fifty-thousand to 60,000 reads were obtained per sample. In total, >2.9 millions of high
quality *nifH* sequences were obtained resulting in 14925 ASVs (submitted to NCBI with accession
numbers SAMN19796776-SAMN19796835
). ASVs were annotated down to the genus level using a DADA2 formatted *nifH* gene database
(https://github.com/moyn413/nifHdada2). Sequences were grouped into 17 genus according to the
database, and sequences not identified to the genus level were grouped as "others". The *nifH* gene
was successfully amplified from all samples.
468
Abundance of diazotrophs et contribution to N export fluxes
The abundance of diazotrophs was determined using TaqMan qPCR assays and previously published
primer-probe sets for *Trichodesmium,* UCYN-A1, UCYN-B, UCYN-C and γ-24474A11 (the latter
hereafter referred to as Gamma-A) targeting the *nifH* gene[65, 66, 67]. The qPCR was run in 25 µL
reactions consisting of 12.25 µL TaqMan PCR Master Mix (Applied Biosystems, Villebon Sur Yvette,
France), 1 µL of the forward and reverse primers at 10 µM (HPLC purified, Eurofins, Nantes, France),
0.25 µL probe at 10 µM, 8.25 µL PCR grade water, 0.25 µL bovine serum albumin at 10.08 µg µL^{-1},
and 2 µL standard or template sample. The qPCR program was run on a CFX96 Real-Time System
thermal cycler (BioRad, Marnes-la-Coquette, France) and consisted of 2 min at 50ºC, 10 min at 95ºC

continued by 45 cycles of 15 s at 95ºC and 1 min at 64ºC. The annealing temperature was changed to
60ºC for UCYN-A1 qPCR runs[67]. Standard dilutions (10^7-10^1 gene copies) were run in duplicate, and
samples and no-template controls (NTCs) in triplicate. NTCs did not show any amplification. The
efficiency was 98-113%. Inhibition tests were carried out on all samples and each primer-probe set by
adding the 2 µL of the 10^5 copies standard to each sample. No inhibition was observed. The limit of
detection and detected but not quantifiable limits were 1 and 8 gene copies per reaction, respectively.
The diazotroph turnover rate representing the fraction of surface diazotrophs exported out of
the photic layer per day was calculated as follows: Turnover rate (d^{-1}) = export flux/abundance in the
photic layer. The proportion of PON collected by sediment traps attributed to diazotrophs was
calculated based on estimated PON content per cell. For nanoplanktonic UCYN, cell dimensions were
directly measured and their biovolume calculated[68]. The C content per cell was estimated from the
biovolume according to Verity et al. (1992)[69] and the N content calculated based on C:N ratios of 5 for
UCYN-B [70, 71] and 8.5 for UCYN-C[50]. For the UCYN-A1-host symbiosis, we considered a size of 1 µm
for the UCYN-1 and 2 µm for the host and a C:N ratio of 6.3[72]. For *Trichodesmium,* we considered the
average value of 10 ng N per *Trichodesmium* filament from the same region[73]. To circumvent the
polyploidy of *Trichodesmium*[74], we divided the qPCR-based abundances by 12. This ratio was
determined for this study by dividing the qPCR-based abundances by the number of *Trichodesmium*
filaments counted at S05M and S10M stations at 6 depths in the photic layer (we considered 100 cells
per filament of *Trichodesmium*).
497
Particle imaging
Seawater samples from traps, MSC and bottle-nets were gently filtered on 0.2 µm (for scanning
electron microscopy, SEM) and 2 µm polycarbonate filters (for epifluorescence microscopy) at very
low pressure to preserve the particle structure. For epifluorescence microscopy, filters were fixed with
paraformaldehyde (2% prepared in filtered seawater) for 10 minutes at ambient temperature and
stored at-80°C until visualized using a Zeiss Axioplan (Zeiss, Jena, Germany) microscope fitted with a
green (510–560 nm) excitation filter, which targeted the phycoerythrin-rich cells. For SEM, samples
were fixed with 2.5% glutaraldehyde and 1.6% PFA for 1 h at room temperature. Subsequently, the
filters were rinsed twice in 0.2 µm filtered in seawater during 15 min, rinsed in osmium during 30 min,
and rinsed thrice in filtered seawater to eliminate excess osmium. Next, the filters passed through a
series of ethanol drying solutions (50, 70, 95 and 100%, 10 min each), and a series of HDMS solutions
(30, 50, 80 and 100%, 10 min each). Finally, the filters were air-dried and stored at room temperature
until SEM analyses. The filters were visualized onshore using a Phenom-Pro benchtop scanning
electron microscope at 10 kV.
512
***Tara* Oceans sampling and reads recruitments in metagenomes**
23 metagenomes, collected from five stations along the *Tara* Oceans expedition transect and
corresponding to a subset of the data presented in (Karlusich JJP*, et al.* 2020), were selected for this
study since these stations were the only ones for which : i) diazotrophs were present in surface waters
and ii) metagenomes samples from both surface (5m) and mesopelagic waters were collected and
sequenced (200-1000 m ; Supplementary Table S2). Briefly, the plankton were separated into discrete
organismal size fractions using a serial filtration system[75] corresponding to picoplankton size (0.2-3
µm), nanoplankton (0.8-3 µm or 0.8-5 µm) and a 'bulk' size fraction corresponding to the fraction >0.8
µm or >3 µm. Metagenomes were sequenced as Illumina overlapping paired reads of 100-108 bp,
which were merged and trimmed based on quality, resulting in 100-215 bp fragments[76]. Metagenomic
reads were recruited against a database of 9 representative genomes of cyanobacterial diazotrophs
(Table S1) using BLASTN (v2.9.0; (Altschul *et al.*, 1990[77]) with default parameters but limiting the
results to one target sequence (--max_target_seqs 1) and keeping only results with an E-value below
1e-30 (-evalue 1e-30) and with a query coverage of, at least, 90% of the read length (-qcov_hsp_perc
90). Following the criteria proposed by Caro-Quintero & Konstantinidis[78] for metagenome-based
prokaryotic genome classification, reads mapping to diazotroph genomes with a percentage of
nucleotide identity equal or greater than 95% were taxonomically assigned to each diazotroph genome
according to their best hit, except reads that mapped to the ribosomal operon of any of the diazotroph
genomes that were filtered out. The number of reads recruited by each diazotroph genome was
normalized to the sequencing depth of each sample (Table S2). For Fig. 8, only samples for which
genome coverage was higher than 1% were taken into account for each organism.
534
535
536
537

547 Table 1. Particulate nitrogen (PON) export fluxes at stations S05M and S10M at 170 m, 270 m and
548 1000 m, and contribution of diazotrophs to total PON export.
549

	Station/Depth (m)	170 m	270 m	1000 m
PON export flux (mg N m^{-2} d^{-1})		4.9±0.6	6.1±1.2	1.6±0.2
Diazotroph-PON (mg N m^{-2} d^{-1})	S05M	0.9	2.3	1.6
Contribution diazotrophs (%)		18	38	100
PON export flux (mg N m^{-2} d^{-1})		13.4±1.3	9.7±0.6	5.2±0.1
Diazotroph-PON (mg N m^{-2} d^{-1})	S10M	1.2	1.2	2.7
Contribution diazotrophs (%)		9	12	52

550
551
552
Figure 1. a. Surface chlorophyll MODIS composite averaged over the time period corresponding to the
TONGA cruise (1 November-6 December 2019) at a resolution of 4 km. Black triangles correspond to
stations where surface-tethered drifting sediment traps were deployed (170 m, 270 m, 1000 m). Grey
dots correspond to stations where Marine Snow Catcher (MSC) casts were performed at three depths
(see Methods), and white dots to MSC casts performed at one depth (200 m). Black circles
correspond to stations where the bottlenet profiles were performed between 2000 m and 200 m. b.
Abundances of the five *nifH* phylotypes (Log10 *nifH* gene copies L^{-1}) studied during the cruise over
the transect (averages over the photic layer, ~0-130 m).

561
562 Figure 2. Relative abundance of diazotroph genera on sediment trap samples. The dendrogram
563 corresponds to the clustering of samples generated by Bray-Curtis similarity. The legend indicates the
564 *nifH* genus affiliation.

565
566 Figure 3. a. Export flux (*nifH* gene copies m^{-2} d^{-1}) of the five diazotroph groups targeted by qPCR
567 (UCYN-A1 symbiosis, UCYN-B, UCYN-C, *Trichodesmium* and Gamma-A) in sediment trap samples at
568 170 m, 270 m and 1000 m at stations S05M and b. at S10M. Error bars represent strandard deviations
569 from triplicate aliquote analyzed in duplicates. c. Export turnover rates (d^{-1}) of the same diazotroph
570 groups (average of the three depths and the two stations for each group).

571
572 Figure 4. Microscopy images showing examples of phycoerythrin-containing UCYN-like cells and
573 *Trichodesmium* in sediment trap samples collected at 170 m, 270 m, and 1000 m at stations S05M
574 and S10M. a-f. Images taken by epifluorescence microscopy (green excitation 510–560 nm, scale bar:
575 50 μm). k-l. Image taken by scanning electron microscopy (SEM).

576
577 Figure 5. Relative abundance of diazotroph genera from marine snow catcher samples. The
578 dendrogram corresponds to the clustering of samples generated by Bray-Curtis similarity. The legend
579 indicates the *nifH* genus affiliation.

580
Figure 6. a-e. Quantification of the five diazotroph groups targeted by qPCR in Marine Snw catcher
samples: UCYN-A1 symbiosis: *Trichodesmium*, UCYN-B, UCYN-C, and Gamma-A in the suspended
(dark color), slow sinking (medium color) and fast sinking (light color) pools of the MSC samples
across the transect (left Y axis, *nifH* gene copies L^{-1}). Dots represent the integrated pool of each
phylotype (right Y axix, *nifH* gene copies m^{-2}) in the photic (~0-130 m) layer. f. Percentage of sinking
versus non-sinking diazotrophs for each group.
587

Figure 7. Abundance (*nifH* gene copies m^{-2}) of the five diazotroph groups targeted by qPCR (UCYN-A1 symbiosis, *Trichodesmium*, UCYN-B, UCYN-C, and Gamma-A) in the bottle-net samples across the transect, representing an integrated sample from 200 m to 2000 m.

Figure 8. Distribution of diazotrophic cyanobacteria in surface and mesopelagic waters during the *Tara* Oceans expedition. A. Geographical location of the *Tara* oceans stations used in this study. Only stations in which diazotrophic cyanobacteria were present in surface and metagenomes were available both from surface and mesopelagic waters were selected. B. Abundance of metagenomic reads recruited against different diazotrophic cyanobacteria genomes in surface and mesopelagic samples. See Table S2 for the complete dataset. Note that to better visualize the data, read abundance was expressed as number of reads per 100,000 total reads for surface samples (left axis) and per 1,000,000 total reads for mesopelagic samples (right axis).

Acknowledgements
This research is a contribution of the TONGA project (Shallow hydroThermal sOurces of trace elemeNts: potential impacts on biological productivity and the bioloGicAl carbon pump; TONGA cruise DOI: 10.17600/18000884) funded by the Agence Nationale de la Recherche (grant TONGA ANR-18-CE01-0016 and grant CINNAMON ANR-17-CE2-0014-01), the LEFE-CyBER program (CNRS-INSU), the A-Midex foundation, the Institut de Recherche pour le Développement (IRD). F.M.C-C. and L.G acknowledge funding from the European Union's Horizon 2020 research programme under the Marie Sklodowska-Curie grant agreement No. 749380 (UCYN2PLAST). The authors warmly thank the crew of the R/V L'Atalante for outstanding shipboard operations. Nagib Bhairy is warmly thanked for his efficient help with MSC deployment and clean CTD rosette management and Vincent Taillandier is tanked for CTD data processing.

Author contributions
SB designed the experiments and SB, MB and IBF carried them out at sea, with advice from FLM; SB, MB, MC, OG, AT, DS analyzed the samples; FMCC and LG analyzed the *Tara* Oceans metagenomes. SB analyzed the data with the help of MB for the sequencing data. SB prepared the manuscript with contributions of all co-authors.

References
1.   Moore CM*, et al.* Processes and patterns of oceanic nutrient limitation. *Nature Geoscience* **6**, 701–710 (2013).

2.   Gruber N. The marine nitrogen cycle: overview and challenges. In: *Nitrogen in the marine environment*) (2008).

3.   Eppley RW, Peterson BJ. The flux of particulate organic matter to the deep ocean and its relation to planktonic new production. *Nature* **282**, 677-680 (1979).

4.   Mulholland MR. The fate of nitrogen fixed by diazotrophs in the ocean. *Biogeosciences* **4**, 37-51 (2007).

5.   Adam B*, et al.* N2-fixation, ammonium release and N-transfer to the microbial and classical food web within a plankton community. *The ISME journal* **10**, 450-459 (2016).

6.   Bonnet S*, et al.* Diazotroph derived nitrogen supports diatom growth in the South West Pacific: A quantitative study using nanoSIMS. *Limnology and Oceanography* **61**, 1549-1562 (2016).

7.   Robinson C*, et al.* Mesopelagic zone ecology and biogeochemistry - a synthesis. *Deep-Sea Research Part Ii-Topical Studies in Oceanography* **57**, 1504-1518 (2010).

8.   Richardson TL, Jackson GA. Small phytoplankton and carbon export from the surface ocean. *Science* **315**, 838-840 (2007).

9.   Subramaniam A*, et al.* Amazon River enhances diazotrophy and carbon sequestration in the tropical North Atlantic Ocean. *Proceedings of the National Academy of Sciences* **105**, 10460–10465 (2008).

10.   White AE*, et al.* Nitrogen fixation in the Gulf of California and the Eastern Tropical North Pacific. *Progess in Oceanography* **109**, 1-17 (2012).

11.   Karl DM, Church MJ, Dore JE, Letelier R, Mahaffey C. Predictable and efficient carbon sequestration in the North Pacific Ocean supported by symbiotic nitrogen fixation. *Proceedings of the National Academy of Sciences* **109**, 1842–1849 (2012).

12.   Luo YW*, et al.* Database of diazotrophs in global ocean: abundances, biomass and nitrogen fixation rates. *Earth System Science Data* **5**, 47-106 (2012).

13.   Sharek RM, Tupas LM, Karl DM. Diatom fluxes to the deep sea in the oligotrophic North Pacific gyre at Station ALOHA. *Marine and Ecological Progress Series* **82**, 55-67 (1999).

14.   Walsby AE. The gas vesicles and buoyancy of *Trichodesmium. Marine Pelagic Cyanobacteria: Trichodesmium and other Diazotrophs*, 141-161 (1992).

15.   Chavez FP, Ryan J, Lluch-Cota SE, Niquen M. From anchovies to sardines and back: Multidecadal change in the Pacific Ocean. *Science* **299**, 217-221 (2003).

16.   Pabortsava K*, et al.* Carbon sequestration in the deep Atlantic enhanced by Saharan dust. *Nature Geoscience* **10**, 189-U141 (2017).

17.   Guidi L*, et al.* Does eddy-eddy interaction control surface phytoplankton distribution and carbon export in the North PacificSubtropical Gyre? *Journal of Geophysical Research* **117**, (2012).

18.   Caffin M*, et al.* N2 fixation as a dominant new N source in the western tropical South Pacific Ocean (OUTPACE cruise). *Biogeosciences* **15**, 2565-2585 (2018).

19.   Agusti S*, et al.* Ubiquitous healthy diatoms in the deep sea confirm deep carbon injection by the biological pump. *Nat Commun* **6**, 7608 (2015).

20.   Berthelot H*, et al.* Dinitrogen fixation and dissolved organic nitrogen fueled primary production and particulate export during the VAHINE mesocosm experiment (New Caledonia lagoon). *Biogeosciences* **12**, 4099-4112 (2015).

21.   Bonnet S*, et al.* Dynamics of N2 fixation and fate of diazotroph-derived nitrogen in a low nutrient low chlorophyll ecosystem: results from the VAHINE mesocosm experiment (New Caledonia) *Biogeosciences* **13**, 2653-2673 (2016).

692   22.   Farnelid H, *et al.* Diverse diazotrophs are present on sinking particles in the North
693   Pacific Subtropical Gyre. *Isme Journal* **13**, 170-182 (2019).
694
695   23.   Moran XAG, Lopez-Urrutia A, Calvo-Diaz A, Li WKW. Increasing importance of
696   small phytoplankton in a warmer ocean. *Global Change Biology* **16**, 1137-1144
697   (2010).
698
699   24.   Polovina JJ, Howell EA, Abecassis M. Ocean's least productive waters are expanding.
700   *Geophysical Research Letters* **35**, (2008).
701
702   25.   Karl D, Michaels A, Bergman B, Capone DG, Carpenter EJ, Letelier R. Dinitrogen
703   fixation in the world's oceans. *Biogeochemistry* **57/58**, 47-98 (2002).
704
705   26.   Bonnet S, Caffin M, Berthelot H, Moutin T. Hot spot of N2 fixation in the western
706   tropical South Pacific pleads for a spatial decoupling between N2 fixation and
707   denitrification. *Proceedings of the National Academy of Sciences* **114**, E2800-E2801
708   (2017).
709
710   27.   de Boyer Montégut C, Madec G, Fischer AS, Lazar A, Iudicone D. Mixed layer depth
711   over the global ocean: An examination of profile data and a profile-based climatology.
712   *Journal of Geophysical Research: Oceans* **109**, (2004).
713
714   28.   Hewson I, Moisander PH, Morrison AE, Zehr JP. Diazotrophic bacterioplankton in a
715   coral reef lagoon: phylogeny, diel nitrogenase expression and response to phosphate
716   enrichment. *Isme Journal* **1**, 78-91 (2007).
717
718   29.   Turk-Kubo KA, Frank IE, Hogan ME, Desnues A, Bonnet S, Zehr JP. Diazotroph
719   community succession during the VAHINE mesocosms experiment (New Caledonia
720   Lagoon). *Biogeosciences* **12**, 7435-7452 (2015).
721
722   30.   Riley JS, Sanders R, Marsay C, Le Moigne FAC, Achterberg EP, Poulton AJ. The
723   relative contribution of fast and slow sinking particles to ocean carbon export. *Global
724   Biogeochemical Cycles* **26**, (2012).
725
726   31.   Cornejo-Castillo FM, Zehr JP. Intriguing size distribution of the uncultured and
727   globally widespread marine non-cyanobacterial diazotroph Gamma-A. *The ISME
728   journal* **15**, 124-128 (2021).
729
730   32.   Karlusich JJP, *et al.* Global distribution patterns of marine nitrogen-fixers by imaging
731   and molecular methods. *BioRxiv*, (2020).
732
733   33.   Salazar G, *et al.* Gene Expression Changes and Community Turnover Differentially
734   Shape the Global Ocean Metatranscriptome. *Cell* **179**, 1068-1083 e1021 (2019).
735
736   34.   Durkin CA, Van Mooy BAS, Dyhrman ST, Buesseler KO. Sinking phytoplankton
737   associated with carbon flux in the Atlantic Ocean. *Limnology and Oceanography* **61**,
738   1172-1187 (2016).
739
740   35.   Cornejo-Castillo FM. Diversity, ecology and evolution of marine diazotrophic
741   microorganisms.). Polytechnic Univ. of Catalonia (UPC) (2017).

742
743   36. Harding K, Turk-Kubo KA, Sipler RE, Mills MM, Bronk DA, Zehr JP. Symbiotic
744   unicellular cyanobacteria fix nitrogen in the Arctic Ocean. *Proceedings of the*
745   *National Academy of Sciences of the United States of America* **115**, 13371-13375
746   (2018).
747
748   37. Shiozaki T, *et al.* Diazotroph community structure and the role of nitrogen fixation in
749   the nitrogen cycle in the Chukchi Sea (western Arctic Ocean). *Limnology and*
750   *Oceanography* **63**, 2191-2205 (2018).
751
752   38. Thompson AW*, et al.* Unicellular Cyanobacterium Symbiotic with a Single-Celled
753   Eukaryotic Alga. *Science* **337**, 1546-1550 (2012).
754
755   39. Poff KE, Leu AO, Eppley JM, Karl DM, DeLong EF. Microbial dynamics of elevated
756   carbon flux in the open ocean's abyss. *Proceedings of the National Academy of*
757   *Sciences of the United States of America* **118**, (2021).
758
759   40. Sohm JA, Edwards BR, Wilson BG, Webb EA. Constitutive Extracellular
760   Polysaccharide (EPS) Production by Specific Isolates of *Crocosphaera watsonii.*
761   *Frontiers in microbiology* **2**, 229 (2011).
762
763   41. Berman-Frank I, Dubinsky Z. Balanced growth in aquatic plants: Myth or reality?
764   Phytoplankton use the imbalance between carbon assimilation and biomass production
765   to their strategic advantage. *Bioscience* **49**, 29-37 (1999).
766
767   42. Walsby AE. The properties and buoyancy-providing role of gas vacuoles in
768   Trichodesmium Ehrenberg. *European Journal of Phycology* **13**, 103-116 (1978).
769
770   43. Berman-Frank I, Bidle KD, Haramaty L, Falkowski PG. The demise of the marine
771   cyanobacterium, Trichodemsium spp., via an autocatalyzed cell death pathway.
772   *Limnology and Oceanography* **49**, 997-1005 (2004).
773
774   44. Berman-Frank I, Rosenberg G, Levitan O, Haramaty L, X. M. Coupling between
775   autocatalytic cell death and transparent exopolymeric particle production in the marine
776   cyanobacterium *Trichodesmium. Environmental microbiology* **9**, 1415-1422 (2007).
777
778   45. Bar-Zeev E, Avishay I, Bidle KD, Berman-Frank I. Programmed cell death in the
779   marine cyanobacterium Trichodesmium mediates carbon and nitrogen export. *ISME*
780   *Journal* **7**, 2340-2348 (2013).
781
782   46. Buesseler KO. The decoupling of production and particulate export in the surface
783   ocean. *Global Biogeochemical Cycles* **12**, 297-310 (1998).
784
785   47. Buesseler KO, Boyd PW, Black EE, Siegel DA. Metrics that matter for assessing the
786   ocean biological carbon pump. *Proceedings of the National Academy of Sciences of*
787   *the United States of America* **117**, 9679-9687 (2020).
788
789   48. Benavides M*, et al.* Staying alive: fast sinking Trichodesmium remains active in the
790   twilight zone. (In prep.).
791

49. Ababou FE, Le Moigne F, Bonnet S. Mechanistic understanding of diazotroph aggregation: a rolling tank aproach. *Targetted journal: Limnology and Oceanography*, (In prep.).

50. Berthelot H, Bonnet S, Camps M, Grosso O, Moutin T. Assessment of the dinitrogen released as ammonium and dissolved organic nitrogen by unicellular and filamentous marine diazotrophic cyanobacteria grown in culture. *Frontiers in Marine Science* **2**, (2015).

51. Glibert PM, Bronk D. Release of dissolved organic nitrogen by marine diazotrophic cyanobacteria, *Trichodesmium* spp. *Applied and Environmental Microbiology* **60**, 3996-4000 (1994).

52. Mulholland MR, Bernhardt PW, Heil CA, Bronk DA, O'Neil JM. Nitrogen fixation and regeneration in the Gulf of Mexico. *Limnology and Oceanography* **51**, 176-177 (2006).

53. Capone DG, *et al.* An extensive bloom of the N_2-fixing cyanobacterium, *Trichodesmiumn erythraeum*, in the Central Arabian Sea. *Marine Ecology Progress Series* **172**, 281-292 (1998).

54. Steinberg DK, Landry MR. Zooplankton and the Ocean Carbon Cycle. *Annu Rev Mar Sci* **9**, 413-444 (2017).

55. Engel A, Wagner H, Le Moigne FAC, Wilson ST. Particle export fluxes to the oxygen minimum zone of the eastern tropical North Atlantic. *Biogeosciences* **14**, 1825-1838 (2017).

56. Berthelot H, Benavides M, Moisander PH, Grosso O, Bonnet S. High nitrogen fixation rates in the particulate and dissolved pools in the Western Tropical Pacific (Solomon and Bismarck Seas). *Geophysical Research Letters*, (2017).

57. Bonnet S, *et al.* In-depth characterization of diazotroph activity across the western tropical South Pacific hotspot of N-2 fixation (OUTPACE cruise). *Biogeosciences* **15**, 4215-4232 (2018).

58. Conte MH, Ralph N, Ross EH. Seasonal and interannual variability in deep ocean particle fluxes at the Oceanic Flux Program (OFP)/Bermuda Atlantic Time Series (BATS) site in the western Sargasso Sea near Bermuda. *Deep-Sea Research Part Ii-Topical Studies in Oceanography* **48**, 1471-1505 (2001).

59. Ras J, Claustre H, Uitz J. Spatial variability of phytoplankton pigment distributions in the Subtropical South Pacific Ocean: comparison between in situ and predicted data. *Biogeosciences* **5**, 353-369 (2008).

60. Lampitt RS, Wishner KF, Turley CM, Angel MV. Marine Snow Studies in the Northeast Atlantic-Ocean - Distribution, Composition and Role as a Food Source for Migrating Plankton. *Marine Biology* **116**, 689-702 (1993).

841   61.   Passow U, Alldredge AL. A dye-binding assay for the spectrophotometric
842   measurement of transparent exopolymer particles (TEP). *Limnology and*
843   *Oceanography* **40**, 1326-1335 (1995).
844
845   62.   Moisander PH, Beinart RA, Voss M, Zehr JP. Diversity and abundance of
846   diazotrophic microorganisms in the South China Sea during intermonsoon. *Isme*
847   *Journal* **2**, 954-967 (2008).
848
849   63.   Zehr JP, Turner PJ. Nitrogen fixation: Nitrogenase genes and gene expression. In:
850   *Methods in Marine Microbiology* (ed Paul JH). Academic Press (2001).
851
852   64.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
853   High-resolution sample inference from Illumina amplicon data. *Nat Methods* **13**, 581-
854   + (2016).
855
856   65.   Church MJ, Jenkins BD, Karl DM, Zehr JP. Vertical distributions of nitrogen-fixing
857   phylotypes at Stn ALOHA in the oligotrophic North Pacific Ocean. *Aquatic Microbial*
858   *Ecology* **38**, 3-14 (2005).
859
860   66.   Moisander AM, Beinart A, Voss M, Zehr JP. Diversity and abundance of diazotrophs
861   in the South China Sea during intermonsoon. *The ISME journal* **2**, 954-967 (2008).
862
863   67.   Thompson A, Carter BJ, Turk-Kubo K, Malfatti F, Azam F, Zehr JP. Genetic diversity
864   of the unicellular nitrogen-fixing cyanobacteria UCYN-A and its prymnesiophyte
865   host. *Environmental microbiology* **16**, 3238–3249 (2014).
866
867   68.   Sun J, Liu D. Geometric models for calculating cell biovolume and surface area for
868   phytoplankton. *Journal of Plankton Research* **25**, 1331-1346 (2003).
869
870   69.   Verity PG, Robertson CY, Tronzo CR, Andrews MG, Nelson JR, Sieracki ME.
871   Relationships between cell volume and the carbon and nitrogen content of marine
872   photosynthetic nanoplankton. *Limnology and Oceanography* **37**, 1434–1446 (1992).
873
874   70.   Dekaezemacker J, Bonnet S. Sensitivity of N_2 fixation to combined nitrogen forms
875   (NO_3^- and NH_4^+) in two strains of the marine diazotroph *Crocosphaera watsonii*
876   (Cyanobacteria). *Marine Ecology Progress Series* **438**, 33-46 (2011).
877
878   71.   Knapp AN, Dekaezemacker J, Bonnet S, Sohm JA, Capone DG. Sensitivity of
879   *Trichodesmium erythraeum* and *Crocosphaera watsonii* abundance and N_2 fixation
880   rates to varying NO_3^- and PO_4^{3-} concentrations in batch cultures. *Aquatic Microbial*
881   *Ecology* **66**, 223-236 (2012).
882
883   72.   Martinez-Perez C, *et al.* The small unicellular diazotrophic symbiont, UCYN-A, is a
884   key player in the marine nitrogen cycle. *Nature Microbiology* **1**, 16163 (2016).
885
886   73.   Rodier M, Le Borgne R. Population and trophic dynamics of Trichodesmium
887   thiebautii in the SE lagoon of New Caledonia. Comparison with T. erythraeum in the
888   SW lagoon. *Marine Pollution Bulletin* **61**, 349-359 (2010).
889

890   74.   Sargent EC*, et al.* Evidence for polyploidy in the globally important diazotroph
891   Trichodesmium. *FEMS microbiology letters* **363**, (2016).
892

893   75.   Pesant S*, et al.* Open science resources for the discovery and analysis of Tara Oceans
894   data. *Sci Data* **2**, (2015).
895

896   76.   Alberti A*, et al.* Viral to metazoan marine plankton nucleotide sequences from the
897   Tara Oceans expedition. *Sci Data* **4**, (2017).
898

899   77.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic Local Alignment
900   Search Tool. *Journal of molecular biology* **215**, 403-410 (1990).
901

902   78.   Caro-Quintero A, Konstantinidis KT. Bacterial species may exist, metagenomics
903   reveal. *Environmental microbiology* **14**, 347-355 (2012).
904
905

UCYN-A   UCYN-B   UCYN-C   Trichodesmium   Gamma-A