Massive export of diazotrophs across the tropical south Pacific Ocean

Sophie Bonnet, Mar Benavides, Mercedes Camps, Antoine Torremocha, Olivier Grosso, Dina Spungin, Ilana Berman-Frank, Frédéric A.C. Le Moigne

Abstract

Diazotrophs are widespread microorganisms that regulate marine productivity in 60% of our oceans by alleviating nitrogen limitation. Yet, their contribution to organic carbon and nitrogen export fluxes has never been quantified, making an assessment of their impact on the biological carbon pump impossible. Here, we examine species-specific fates of several groups of globally-distributed unicellular (UCYN) and filamentous diazotrophs in the mesopelagic ocean. We used an innovative approach consisting of the combined deployment of surface-tethered drifting sediment traps, Marine Snow Catcher, and Bottle-net, in which we performed *nifH* sequencing and quantitative PCR on major diazotroph groups across the subtropical South Pacific Ocean. *nifH* sequencing data from sediment traps deployed at 170 m, 270 m and 1000 m provide clear evidence that cyanobacterial and non-cyanobacterial diazotrophs are systematically present in sinking particles down to 1000 m, with export fluxes being the highest for the UCYN-A1 symbiosis, followed by UCYN-B or *Trichodesmium* (depending on station and depth), Gamma A and UCYN-C. Specific export turnover rates (a metric similar to the export efficiency adapted to organisms) point to a more efficient export of UCYN groups relative to the filamentous *Trichodesmium*. This is further confirmed by Marine Snow catcher data showing that the proportion of sinking cells was significantly higher for UCYN compared to *Trichodesmium*. Phycoerythrin-containing UCYN-B and UCYN-C-like cells were indeed recurrently found embedded in large (> 50 µm) seemingly organic aggregates, or organized into clusters of tens to hundreds of cells linked by an extracellular matrix, facilitating the export. Overall, diazotrophs accounted for 6-13% (170 m) to 45-100% (1000 m) of the total particulate nitrogen export fluxes in our study. We thus conclude that diazotrophs are important contributors to carbon sequestration in the subtropical South Pacific Ocean and need to be considered in future studies to improve the accuracy of current regional and global estimates of export.

Introduction

Nitrogen (N) availability limits primary productivity throughout much of the surface low-latitude ocean[1]. In such nitrogen (N)-limited waters, microbial dinitrogen (N_2) fixation by diazotrophic plankton provides the major source of new N to surface waters[2] that on appropriate timescales, equates export production[3]. However, the fate of this production, whether it is exported to the deep ocean or stimulates remineralization in surface waters remains obscure[4]. An increasing number of studies have shown that diazotroph-derived N is quickly translocated to non-diazotrophic plankton such as diatoms[5, 6], which eventually contribute to secondary export of organic matter out of the photic zone. Yet, the direct gravitational settling of diazotrophs themselves to the deep ocean has rarely been assessed. The few existing observations (see below) conclude that diazotrophs provide a minor contribution to the total amount of N in the exported material.

Diazotrophs may associate with sinking particles to directly contribute to export by different mechanisms. The two most direct include gravitational settling of individual cells/filaments or aggregates. According to Stokes' Law, particle sinking velocity scales with the square of particle size. Therefore, large particles should sink faster and are more likely to reach the deep ocean before being remineralized by bacteria[7]. Aggregation is thus a crucial step for the transport of small phytoplankton species which could export particulate organic carbon (POC) and N (PON) in similar proportion to their production in surface waters[8]. Diazotrophs have diverse morphologies and their size spans several orders of magnitude. Some types such as the free-living unicellular diazotrophic cyanobacteria (UCYN from groups B and C) are small (2-8 µm in diameter), while others such as *Trichodesmium sp.* are filamentous and can form large-sized colonies (>100-1000 µm). In addition, some diazotrophs live in symbiosis with calcified (UCYN-A, ~1 µm) or silicified eukaryotes (*Richelia* sp., *Calothrix* sp., >20 µm, forming Diatom-Diazotrophic Associations or DDAs). These dense biominerals may provide ballast enhancing the downward export of these symbioses into the deep ocean. Therefore, the presence of different diazotrophs in surface waters may result in drastically different C and N export fluxes via the biological pump. Yet, no field observations have related individual diazotroph groups to the magnitude of downward particles fluxes to date.

Thanks to their inherently ballasted character, DDAs are well known to contribute to particulate matter export[9, 10], and are involved in seasonal peaks of POC export to the deep sea (4000 m) in the North Pacific subtropical gyre[11]. *Trichodesmium* is one of the major contributors to global N_2 fixation[12]. However, it is thought to have a limited export capacity and to be preferentially remineralized in the

surface layers due to the presence of gas vesicles potentially providing buoyancy[4, 13, 14]. Yet, some
studies have reported the presence of *Trichodesmium* in sediment traps material in the Kuroshio
Current[15], the tropical North Pacific[16] and the South Pacific Oceans[17]. Intact filaments and colonies of
Trichodesmium sp. have also been reported as deep as 3000-4000 m in the tropical Atlantic, Pacific
and Indian Oceans[18, 19], but their contribution to organic matter export has to be quantified.
A priori, UCYN may not contribute importantly to POC export fluxes either due to their small
size. Yet, Berthelot et al.[20] reported in experimental mesocosms that primary production supported by
UCYN was twice as more efficient in promoting POC export than production supported by DDA. In the
same experiment, Bonnet et al.[21] showed that the sinking of UCYN-B and C was driven by the
aggregation of small individual cells (5-6 μm) into larger aggregates (100 to >500 μm) that were
eventually exported and accounted for up to ~20% of the POC flux. Recently, Caffin et al.[17] confirmed
the presence of UCYN-B in sinking particles collected with sediment traps (330 m) throughout the
tropical South Pacific Ocean (10^3 to 10^7 *nifH* gene copies L^{-1} of trap material). Finally, Farnelid et al.[22]
observed sequenced *nifH* genes of exported material (150 m) in the North Pacific subtropical gyre and
found that all the diazotroph groups above as well as non-cyanobacterial diazotrophs were present in
the exported material.
Taken together, these studies suggest that all diazotroph groups, small or large, free-living or
symbiotic, ballasted or not, are present below the photic layer. However, a detailed examination relating
types of diazotrophs to the magnitude of downward particles fluxes and their contribution is needed to
refine our understanding of their role in the magnitude and mechanisms controlling the biological pump.
This is all the more topical as climate models predict an expansion of the oligotrophic gyres[23, 24], where
diazotrophs represent a major component of the plankton biomass and sustain most new primary
production.
Here, we examine species-specific fates of diazotrophs in the mesopelagic ocean. We used an
innovative approach consisting of the combined deployment of surface-tethered drifting sediment traps,
Marine Snow Catcher (MSC), and Bottle-net, in which we performed *nifH* sequencing and quantitative
PCR on major diazotroph groups across the subtropical South Pacific (Fig. 1a). This study builds up on
prior investigations that have characterized this region as a hotspot for marine N$_2$ fixation[25]. We show
that all globally-significant N$_2$-fixing cyanobacteria and non-cyanobacterial diazotrophs are
systematically present in sinking particles down to 1000 m. Small size UCYN (1-8 μm) are exported
more efficiently than large filamentous diazotrophs as large (>50 μm) aggregates, but both are major
contribution to PON export fluxes. We thus conclude that diazotrophs are important contributors to
carbon sequestration in the ocean.
93
Results and discussion
95
Surface conditions
The study region was characterized by typical oligotrophic waters (chlorophyll concentrations <0.15 μg
L^{-1}, DCM 100-180 m) east and west of Tonga, corresponding to the first group of stations S01, S02, S07
and S08 (Fig. 1a). The second group of stations (S03, S04, S05, S10, S11) was located in mesotrophic
waters (chlorophyll >0.15 μg L^{-1}, DCM 70-90 m) in the vicinity of the Tonga volcanic arc (Fig. 1a). Surface
(0-50 m) nitrate concentrations were consistently close or below the detection limit (0.05 μmol L^{-1})
throughout the transect, while phosphate concentrations were typically 0.1 μmol L^{-1} at oligotrophic
stations and depleted down to detection limit (0.05 μmol L^{-1}) at mesotrophic stations, likely due to
consumption by higher plankton stocks (Fig. S1). Seawater temperature ranged from 23.1 to 27.3 °C in
the mixed layer (0-15 m to 0-60 m) determined according to de Boyer Montégut et al. (2004)[26] (Fig. S1).
The abundances of key diazotroph groups indicated that UCYN-A1 were present at high
abundances throughout the photic layer (0-130 m) of the transect (average 3.2 x 10^8 *nifH* gene copies
L^{-1}) (Fig. 1b), peaking at oligotrophic stations (average 1.1 x 10^9 *nifH* gene copies L^{-1}). *Trichodesmium*
was the second most abundant group, particularly in the vicinity of Tonga (average 1.3 x 10^7 *nifH* gene
copies L^{-1}), followed by UCYN-B (5.7 x 10^5 *nifH* gene copies L^{-1}). UCYN-C and Gamma-A (a non-
cyanobacterial diazotroph) were also detected throughout the transect albeit at lower abundances (10^2
to 10^4 *nifH* gene copies L^{-1}).
113
How efficient are different diazotrophs groups to export?
We first examined the *nifH* gene community composition of particles collected in sediment traps moored
at 170 m, 270 m and 1000 m at stations S05M and S10M (Fig. 1). 32% of the retrieved Amplicon
Sequence Variants (ASV) corresponded to cyanobacteria genera on average over both stations (Fig.
2). The most abundant genus was *Trichodesmium* sp., but ASVs related to *Crocosphaera watsonii*
(UCYN-B), *Candidatus Atelocyanobacterium thalassa* (UCYN-A) and *Katagnymene* spp. were also

identified (Fig. 2). Genera related to DDAs (*Richelia* spp., *Calothrix* spp.) contributed little to the
percentage reads. Besides autotrophic diazotrophs, 68% of ASVs were affiliated to non-cyanobacterial
diazotrophs including Alpha-, Beta-, and Gammaproteobacteria at both stations.
Amplicon sequencing provides relative abundances, which are not necessarily indicative of the
abundance of the target microorganism in the natural environment[27, 28]. Hence, to assess the export
capacity of individual diazotroph groups, we quantified the abundance of five groups spanning different
forms, sizes, lifestyle (symbiotic or not), using qPCR assays (UCYN-A1 symbiosis, UCYN-B, UCYN-C,
Trichodesmium and Gamma-A; see Methods). DDAs were not quantified as they were almost not
detected on microscopic filters, pointing to their rarity in those waters at the time of the cruise. All five
groups were detected at abundances ranging from 10^3 to 10^8 *nifH* gene copies L^{-1} of trap material, the
highest being reported for UCYN-A1 and the lowest for UCYN-C (Fig. 3a). The diazotrophs assemblage
exported in sediment traps generally reflected that of the photic layer, although some diazotrophs seem
to sink more efficiently than others: the highest diazotroph export fluxes were measured for UCYN-A1
at both sites and all depths of traps deployment ($9.5 \pm 1.9 \times 10^8$ to $1.3 \pm 1.0 \times 10^9$ *nifH* gene copies m^{-}
^{2}d^{-1}), followed either by UCYN-B ($1.3 \pm 0.6 \times 10^7$ to $4.7 \pm 1.2 \times 10^7$ *nifH* gene copies m^{-2}d^{-1}) or
Trichodesmium ($1.6 \pm 1.2 \times 10^6$ to $7.6 \pm 1.7 \times 10^8$ *nifH* gene copies m^{-2}) depending on station and depth
(Fig. 3). Gamma-A and UCYN-C were also exported, albeit at lower rates (~10^5-10^6 and ~10^5 *nifH* gene
copies m^{-2}d^{-1}, respectively).
Specific export turnover rates provide information on the rate of diazotrophs "lost" from the
photic layer per day due to export. This metric is similar to the export efficiency (e-ratio=POC export/POC
produced by Primary Production), but adapted to organisms. The range of calculated rates ($0.8 \pm 0.6 \times 10^{-5}$
to $7.0 \pm 1.2 \times 10^{-3}$ d^{-1}) spans through two orders of magnitude and vary depending on the diazotroph
group, depth and station considered. On average (over all stations and depths), the export turnover rate
of UCYN ($3.5 \pm 1.9 \times 10^{-3}$ d^{-1}) was ca. four times higher than that of *Trichodesmium* ($0.8 \pm 1.3 \times 10^{-3}$ d^{-}
^{1}) (Fig. 3c), pointing to preferential export of UCYN groups relative to the filamentous *Trichodesmium*.
Among UCYN groups, the highest export turnover rate was measured for UCYN-B, followed by UCYN-
C and UCYN-A1, although differences were not significant between groups (Mann-Whitney test,
$p<0.05$). The export turn-over rate of Gamma-A (average $1.1 \pm 0.9 \times 10^{-3}$ d^{-1}) was intermediate between
that of *Trichodesmium* and UCYN groups.
Epifluorescence microscopy confirmed the presence of phycoerythrin-containing UCYN-B and
UCYN-C-like cells in sediment trap samples (Fig. 4a-f). Scanning electron microscopy revealed that
they were recurrently found embedded in large seemingly organic aggregates, or organized into clusters
of tens to hundreds of cells linked by an extracellular matrix (Fig. 4g-l), which was further confirmed by
alcian blue straining (Fig. S2). *Trichodesmium* was also observed in all samples, mostly as free
filaments, but intact colonies were observed in sediment trap samples especially at 1000 m at both
stations. Gamma-A and UCYN-A1 cannot be visualized by these techniques and are thus assessed
solely on the basis of qPCR counts (above).
Pigment concentrations measured in sediment trap samples at both stations indicate that total
pigment concentrations and diversity decreased with depth. Apart from the degradation pigments
phaeophorbide and phaeophytin, zeaxanthin, a biomarker of cyanobacteria predominated at all depths
(Fig. S2). The Chlorophyll *a* : pheopigments ratios were elevated (average 1.8, range 0.8-3.), indicative
of fresh organic matter in sediment trap samples.
162
Contribution of diazotrophs to PN export fluxes
Sediment trap-based export fluxes of PN at 170 m, 270 m and 1000 m ranged between 1.6 ± 0.2 and
6.1 ± 1.2 mg N m^{-2} d^{-1} at S05M and 5.2 ± 0.1 and 13.4 ± 1.3 mg N m^{-2} d^{-1} at S10M (Table 1). Fluxes
attenuated with depth, with transfer efficiencies of 0.02 and 0.03 between 170 m and 1000 m at both
stations. The N export efficiency (the amount of PN export at the base of the euphotic zone over the
integrated N_2 fixation rate) was 18% and 53% at stations S05M and S10M. The C:N ratios in exported
material was lower (5.2) than that of the surface (7.5, integrated value over the photic layer) at station
S10M, pointing to a preferential export of N relative to C.
The proportion of PN attributed to diazotrophs and collected by sediment traps was evaluated
based on qPCR results and estimated PN content per diazotroph (see Methods). The PN flux attributed
to diazotrophs ranged from 0.6-12.3 mg N m^{-2} d^{-1} at S05M and 2.3-7.8 mg N m^{-2} d^{-1} at S10M, accounting
for 13% (170 m) to 100% (1000 m) and 6% (170 m) to 45% (1000 m) of the total PN export flux,
respectively at stations S05M and S10M (Table 1). *Trichodesmium* was the major contributor of the PN
attributed to diazotrophs at 1000 m at both stations, whereas UCYN dominated at shallower depths
(Table S1). Despite non-cyanobacterial diazotrophs were present in exported material (68% of
sequences), their C and N composition remains unknown due to the poor availability of cultured
representatives. Thus, quantifying their contribution to export fluxes is not possible at this stage. In

addition, sequencing data also revealed the presence of *Katagnymene* spp., which are large (>100-
1000 μm) and likely have high C and N content. Hence the contribution of diazotrophs to exported PN
reported here represent a lower-end estimate.
183
The fate of diazotrophs in the mesopelagic ocean
We further assessed the fate of the exported diazotroph community by collecting fresh particles in
mesopelagic waters using the Marine Snow Catcher (MSC, see Methods). MSC samples were collected
at the same depths as those of sediment traps (170, 270, 1000 m) at stations S05M and S10M and in
addition at seven other stations (Fig. 1). In essence, the MSC allows the separating non-sinking particles
from slow sinking and fast sinking particles[29] (see Methods). These different fractions can hence be
studied individually.
The *nifH* composition in the various fractions generally mirrored that of traps at stations S05M
and S10M with 32-54% of sequences affiliated with cyanobacteria and a prevalence of *Trichodesmium*
(Fig. 5). This was also the case at stations S03, S04, S07 and S11, but not at stations S01, S02 and
S08, where sequences assigned to *Candidatus Atelocyanobacterium thalassa* and *Crocosphaera*
watsonii were generally more abundant than *Trichodesmium* sequences. As in sediment traps, a large
number of *nifH* sequences (46-58%) were affiliated to non-cyanobacterial diazotrophs, which
composition was overall consistent with that of the traps, although some classes not detected in
sediment traps (Epsilonproteobacteria and Clostridia) were present in all MSC fractions. In general, the
total number of reads was consistently 10-25% higher in the slow and fast sinking fractions compared
to the suspended fraction at all stations. This trend was mainly driven by non-cyanobacterial sequences
that are suspected to be attached to particles[30], and to a lesser extent by sequences affiliated with
cyanobacteria.
As in sediment traps, five diazotroph groups were quantified in the suspended, slow and fast
sinking fractions to more finely assess their sinking dynamics over short time scales (h). Overall, UCYN-
A1 and *Trichodesmium* were the most abundant groups in the mesopelagic MSC samples (170 m to
1000 m), and their numbers varied in parallel with the abundance of their population in the photic layer
(Fig. 6). UCYN-A1 were the most abundant at oligotrophic stations (S01, S02, S07, S08, chlorophyll
<0.15 μg L^{-1}), while *Trichodesmium* peaked at mesotrophic stations (Chlorophyll a >0.15 μg L^{-1}, Fig. 1).
Conversely, the abundance of UCYN-B, UCYN-C and Gamma-A in mesopelagic waters was poorly
correlated with their photic layer abundance.
The MSC suspended fraction accounted for the most abundant pool of diazotrophs at all
stations, followed by either the slow or the fast sinking pools, depending on groups (Fig. 6a-e). At the
mooring stations S05M and S10M, the total concentration of diazotrophs and the diazotroph community
composition remained generally constant at the three sampled depths (170 m, 270 m and 1000 m),
consistent with the results observed in sediment traps. To evaluate the export potential of each individual
diazotroph group, we calculated the ratio of sinking versus non-sinking cells (the sum of the slow and
fast sinking fractions over the total). The proportion of sinking cells varied widely across the transect,
but was the highest for UCYN-A1 (32 ± 33 %), UCYN-C (31 ± 38 %), Gamma-A (29 ± 19 %) and UCYN-
B (27 ± 11 %), and the lowest for *Trichodesmium* (18 ± 20 %) (Fig. 6f). Among the sinking fractions,
UCYN-B and Gamma-A were generally equally distributed among the slow and fast sinking fractions,
whereas the majority of *nifH* gene copies were found in the fast sinking fraction for UCYN-A1 (59%),
UCYN-C (62%) and *Trichodesmium* (67%). Taken together, these results indicate that over short time
scales (2 h, the conventional settling of particles following MSC deployment[29]): i) small UCYN (few μm)
generally sink more efficiently than large *Trichodesmium* (>100 μm), ii) when sinking, *Trichodesmium*
sinks fast, iii) UCYN-A1 and UCYN-C sink faster than UCYN-B and Gamma-A. A detailed imaging study
performed on 170 m MSC samples at the mooring stations S05M and S10M indicates that the 50-80%
of phycoerythrin-containing UCYN in the fast sinking fraction were organized into aggregates of tens to
>250 cells measuring 30 to >100 μm, while the majority (60-95%) of UCYN were free living in the
suspended fraction (Figure S3). This indicates that under minimum turbulent agitation, UCYN quickly
(within 2 h) form aggregates large/dense enough to sink.
Concentrations of PN averaged across all stations 1.85 ± 0.49 μg L^{-1}, 0.25 ± 0.13 μg L^{-1} and
0.26 ± 0.08 μg L^{-1} within the suspended, slow sinking and fast sinking fractions (Fig. S4). PON in the
fast sinking fraction contributed 11 ± 2 %, while the slow sinking and suspended fractions contributed
11 ± 4 % and 79 ± 5 % in terms of total PON. Thus, ~22% of the PON was sinking out of the upper part
of the MSC within 2h during our study. We converted transparent expolymeric particles into C (TEP-C),
that revealed higher concentrations in the suspended fraction compared to the fast sinking fraction at
all stations (TEP-C concentrations were null in the SS fraction) (Fig. S4). The TEP-C:POC ratio was
generally higher in the suspended fraction compared to the fast sinking, indicating a larger contribution
of TEP to the POC pool.

Finally, we quantified diazotrophs on vertical profiles spanning the water column between 200
m and 2000 m by using a Bottle-net mounted on the CTD rosette frame at six stations across the transect
(Fig. 1). The bottle-net consists of a 20-µm conical plankton net housed in a cylindrical PVC pipe[18]. It is
lowered with the top cover closed, which is opened at the desired bottom depth (2000 m) of the tow,
remains opened during the ascent of the rosette, and closed again at the upper depth (200 m) of the
water column to be sampled. This results in one integrated sample of 200 m to 2000 m per deployment.
The concentration of diazotrophs in this deep layer averaged 1.4 x 10^7 *nifH* gene copies m^{-2} and was
primarily dominated by UCYN-A1 (63% on average over all sampled stations) and *Trichodesmium*
(27%), and secondarily by UCYN-B (9%) (Fig. 7), generally mirroring the diazotroph community in
surface water (these numbers are conservative and may be underestimated as some individual UCYN
pass through the 20 µm mesh net of the bottle-net). The 2000-200 m stock was relatively constant
among stations, except at the most oligotrophic station S08, where it was lower by two orders of
magnitude than that of other stations.

Biogeochemical implications
Apart from silicified DDAs[11], diazotrophs have been seldom regarded as important contributors to
organic matter export. Yet, our results provide clear evidence that they are present and ubiquitous in
the mesopelagic ocean and they contribute significantly to the PN export fluxes in the sub-tropical south
Pacific Ocean. They have different behaviors however.
We showed that diazotroph cell sizes is not necessarily a key variable controlling their ability to
sink out. Small UCYN (1-8 µm) displayed the highest export turn-over rates ~10^{-3} d^{-1}, in the range or
higher than those reported for diatoms and coccolithophores[31]. Within UCYN, UCYN-A1 was clearly
dominating the diazotroph community targeted by qPCR in mesopelagic waters. Although the UCYN-
A1 symbiosis has been detected once in mesopelagic waters of the South Atlantic ocean[32, 33], their
capacity to leave the photic layer and sink throughout the mesopelagic zone down to 1000 m (Fig. 3)
has not been thoroughly explored before. As UCYN-A rank among the most abundant diazotrophs in
the global ocean[12] and spans from tropical to polar waters[34, 35], we suggest that they potentially play a
significant contribution to ocean organic matter sequestration. In future studies, better constraining
UCYN-A sedimentation rates and aggregation processes would be of primary importance to assess
their role in particle fluxes and cycling.
UCYN-B and -C are free-living unicellular cyanobacteria and their export pathway therefore
does not rely on mineral ballasted hosts. Our results indicate that the sinking of such small cells is made
possible thanks to aggregation of UCYN into large-size aggregates (30->100 µm) of tens to hundreds
of cells, large enough to sink, in accordance with a previous mesocosm study[21]. The presence of UCYN-
B in sediment traps is also in accordance with previous reports of *Crocospheara watsonii* sequences at
station ALOHA either just below the euphotic zone[22], or as deep as 4000 m[36]. The visualization of our
sediment trap samples shows that they can be subsequently embedded in mixed aggregates with other
cells and debris (Fig. 4, S2). The UCYN-B large (>4 µm) ecotype, which was dominating in this study
(data not shown), produces large amounts of C-rich transparent exopolymeric particles (TEP) at rates
one to two order of magnitude higher than that of diatoms and coccolithophorids[37], probably in response
to nutrient limitation and excess light[38]. In cultures of *Crocosphaera watsonii*, TEP account for ~22% of
the particulate C pool[37]. Hence, TEP produced by UCYN-B do not only provide a matrix for the formation
of large aggregates, but may also account for a significant fraction of C export. TEP are indeed greatly
enriched in C relative to N (C:N ~25[37]), which is confirmed by the high TEP-C:PC ratios in the fast sinking
fraction of the MSC.
Trichodesmium was generally the second contributor to the diazotroph community targeted by
qPCR in mesopelagic waters after UCYN-A1, which contradicts the common assertion that they are
entirely remineralized in the euphotic layer[4, 13]. Several potential mechanisms could explain the
presence of *Trichodesmium*'s in mesopelagic waters. *Trichodesmium* colonies are able to perform
vertical migrations for exploiting the deep phosphate stock (Villareal and Carpenter 2003) (the
phosphacline was around 200 m in our study region at the time of the cruise). According to this theory,
they overcome their positive buoyancy by fixing C that result in carbohydrate ballasting. However,
Walsby (1978)[39] observed that 100% of the gas vacuoles of *Trichodesmium erythraeum* (the most
abundant species in this study) collapse at depths between 105 m and 120 m, resulting in a loss of their
buoyancy. Hence, *Trichodesmium* could be locked in a persistent and irreversible downward trajectory.
Alternatively, Berman-Frank et al.[40, 41] have shown that programmed cell death induces internal cellular
degradation, vacuole loss and increased production of TEP, resulting in an increase in the vertical flux
of *Trichodesmium*[42]. Whatever the mechanism behind the vertical flux of *Trichodesmium*, we report here
higher abundances at 1000 m compared to those at shallower depths at both stations, consistent with
microscopic observations showing intact colonies at 1000 m. This likely results from a spatio-temporal

decoupling between production and export[43]. Despite sinking less efficiently than UCYN in the
aggregates, they are much larger and contain more C and N per filament than UCYN[12], and accounted
for a significant fraction of the diazotroph-attributed PN export in this study, especially at 1000 m (Table
S1), which is in accordance with previous studies reported intact *Trichodesmium* colonies down to 3000-
4000m[18, 19].
305   More than 90% of the organic matter sinking below the euphotic zone is respired before it
reaches a depth of 1000 m[7]. Fast sinking particles will therefore theoretically make a greater contribution
to the deep ocean flux than slow sinking particles, since the latter will be rapidly recycled at shallow
depths. The relative similarity of the taxonomic diazotroph community composition in mesopelagic
waters sampled by three independent sampling approaches (traps, bottle-net, and MSC) compared to
the diazotrophic composition of the euphotic layer suggests a rapid export mode of diazotrophs. This is
further confirmed by i) the high proportion of diazotroph groups quantified in the MSC fast sinking
fraction, ii) the whole cells and colonies and high Chlorophyll *a* : pheopigments ratios in sediment traps,
indicative of healthy phytoplankton (Fig. 4; S2). Finally, iii) while PN export fluxes were attenuated with
depth in our study, which is a classical feature in the oligotrophic ocean[44], the diazotroph export fluxes
were not, suggesting a high transfer efficiency to the deep ocean, and thus sinking velocities high
enough to escape short-term remineralization. Indeed, a parallel study performed during the same
cruise[45] reveals significant N_2 fixation rates in sediment traps, MSC and bottle-net samples, suggesting
that part of the diazotroph community in mesopelagic waters sunk fast enough to remain alive at
mesopelagic depths. Bar-Zeev et al.[42] reported sinking velocities of *Trichodesmium erythraeum* of ~200
m d^{-1}. More recently, Ababou et al.[46] reported sinking velocities of *Trichodesmium erythraeum* and
UCYN aggregates formed in roller tanks of 92 ± 37 m d^{-1} and 333 ± 176 m d^{-1}, respectively. We
calculated that *Trichodesmium* sinking at those rates (92-200 m d^{-1}) would take 5 to 10 days to reach
1000 m, whereas UCYN would take 3 days, which would be compatible with finding active cells observed
at those depths[45].
325

Conclusion

These findings challenge the common assumption that the fate of diazotrophic new nitrogen is
constrained to the surface layer. We show that diazotrophs other than DDAs get exported below the
euphotic zone through direct gravitational settling, and that small diazotrophs are the most efficiently
exported thanks to aggregation processes in large particle. However, direct export through gravitational
settling of diazotrophs is most likely only the tip of the iceberg and diazotrophs can potentially be
exported through secondary pathways: diazotrophs release in seawater 10–50% of recently fixed N_2
(referred to as Diazotroph Derived, DDN) as NH4+ and dissolved organic N (DON)[47, 48, 49]. This DDN is
potentially available for assimilation by the surrounding phytoplanktonic communities, supporting their
growth and leading a potential secondary (indirect) export pathway of diazotroph-derived OC[6, 21, 50].
Moreover, DDN is also transferred to zooplankton, which produces dejections termed fecal pellets, that
sink fast and play a major role in OC export to the deep ocean[51]. In future studies, there is an urgent
need to develop appropriate approaches to decipher diazotroph export pathways if we are to understand
the role of diazotroph of the biological carbon pump. This is an especially pressing question given that
current climate models predict an expansion of the oligotrophic gyres (60% of our oceans), where
diazotrophs thrive and sustain most new primary productivity[23, 24].
342

Methods

Samples were collected during the TONGA cruise (Fig. 1, DOI: 10.17600/18000884) in the tropical
South Pacific Ocean onboard the R/V *L'Atalante* from November 1 to December 5 2019. We collected
suspended and sinking particles in the mesopelagic zone using three complementary devices: surface-
tethered drifting sediment traps[52], MSC and Bottle-net. Additionally, water column samples were
collected from the euphotic layer using Go-Flo bottles mounted on a trace metal clean rosette to quantify
the stock of the major groups of diazotrophs in the photic layer.
350

Satellite products and photic layer sampling. Satellite-derived surface chlorophyll *a* concentration
during the TONGA cruise (November 2019) was accessed at http://oceancolor.gsfc.nasa.gov (MODIS
Aqua, 4 km, 8-days composite, level 3 product). Vertical 0-150 m depth profiles were performed at each
station using a trace metal clean titanium rosette of Go-Flo bottles equipped with a fluorometer and
temperature, conductivity and oxygen sensors. Seawater samples were collected from 5 depths (75, 50,
20, 10, 1% surface irradiance levels) to quantify the stock of major groups of diazotrophs in the photic
layer by quantitative PCR (qPCR) as described below. Additionally, primary production and N_2 fixation
rates were measured in triplicates at the same six depths at the locations of the traps deployments
(S05M and S10M), as described in [53, 54]. N_2 fixation rates were measured at two occasions (on day 1

and day 3 of the traps deployment). The average value between both profiles was used to calculate the
e-ratios (POC export:primary production) and the contribution of N_2 fixation to primary production rates.
362
Sediment trap deployment and sample analyses. A surface tethered mooring line (~1000 m long)
was deployed at stations S05M (21.157°S;175.153°W, 5 days) and S10M (19.423°S;175.133°W, 4
days). The line was equipped with sediment traps (KC Denmark®) at 3 depths: 170 m (corresponding
to the base of the photic layer), 270 m and 1000 m. Each trap was composed of four particle interceptor
tubes (PITs) mounted on a cross frame (collecting area of 0.0085 m^2, aspect ratio of 6.7). Two of the
tubes were used for this study: one tube was used for biogeochemical analyses (hereafter referred to
as 'Biogeo' tube), and one for microbiological analyses ('Microbio' tube). Prior to deployment, the tubes
were filled with 0.2 µm filtered seawater with added saline brine (50 g L^{-1}). Borate-buffered formalin (5%)
was also added to the Biogeo tube to prevent in situ microbial decomposition[52]. After recovery, the
density gradient was visually verified, and the PITs were allowed to settle for 2 h before the supernatant
seawater was carefully removed with a peristaltic pump. The remaining water containing the sinking
material was transferred to a chlorohydric acid-washed container, while being screened with a 500 µm
mesh to remove swimmers (zooplankton that actively entered the traps)[55]. Subsequently, samples were
split into 12 aliquots. A triplicate set of aliquots were filtered onto 25-mm diameter Supor filters for *nifH*
sequencing and *nifH* qPCR as described below. Another triplicate set of aliquots were filtered onto 25-
mm diameter combusted (4h, 450°C) glass microfiber filters (Whatman GF/F), which were subsequently
dried for 24 h at 60°C, pelleted and from which particulate N (PN) and C (PC) were analyzed by EA-
IRMS (Elemental Analyzer-Isotope Ratio Mass Spectrometry) using an Integra-CN (Sercon) mass
spectrometer. Lastly, a triplicate set of aliquots were filtered on GF/F filters for further pigment
analyses[56], and another triplicate filtered on 1 µm polycarbonate filters for microscopic analyses (see
below for methods). The turnover rate of diazotrophs in the photic layer due to vertical export was
determined by dividing cell fluxes (*nifH* gene copies m^{-2} d^{-1}) by their integrated abundance in the photic
later (*nifH* gene copies m^{-2}).
386
MSC deployment
Suspended and sinking particles were sampled using a MSC at 3 depths at the mooring stations S05M
and S10M (170 m, 270 m and 1000 m), and at 200 m, 400 m and 1000 m at S8. Additionally, the MSC
was deployed at 200 m at S01, S02, S03, S04, S05V, S07, S10V and S11 (Fig. 1). The MSC is a large
volume water sampler (100 L) that collects sinking particles with minimal turbulent agitation[57]. Upon
recovery, the MSC is placed on deck for 2 h while any particles present settle onto the base of the
bottom 7 L chamber. After each recovery, the MSC is conventionally left on deck for 2 h allowing particles
to settle[29]. Fast sinking particles were thereby deposited collected in a dedicated plate at the bottom of
the MSC, slow sinking particles were collected from the 7-litre compartment above the plate, and the
non-sinking (or suspended) fraction was collected from the upper part of the MSC[29]. All three fractions
were collected separately (4 L for the suspended, 1.2 L for the slow sinking). For the fast sinking fraction,
the entire volume of the plate was sampled (280 to 310 mL). Subsamples from each fraction were filtered
for PN and PC quantification, TEP quantification according to Passow and Alldredge (1995)[58],
microscopic observations, and *nifH* sequencing, *nifH* qPCR as described below. We used the formula
described in Riley et al., (2012)[29] to obtain values associated with fast or slow sinking particles only,
quantitatively removing the contribution of other fractions.
403
Bottle-net deployment
Vertical 2000-200 m profiles were done using a bottle-net mounted on the CTD rosette frame at stations
S03, S04, S05M, S07, S08, S10M, S11 and S12. The bottle-Net consists of a 20-µm conical plankton
net housed in a cylindrical PVC pipe[18]. It is lowered with the top cover closed, which is opened at the
desired bottom depth (Db, 2000 m) of the tow, remains opened during the ascension of the rosette, and
closed again at the upper depth (Du, 200 m) of the water column to be sampled. This results in one
integrated sample of 2000 to 200 m per deployment. Once on deck, the bottle-nets were gently rinsed
with filtered seawater before retrieving the sample from the collector. At each station, samples were split
into aliquots that were processed for were analyzed for *nifH* qPCR. Sampled volume was estimated as
the product between the cross-sectional area of the mouth of the bottle-Net (7.5 cm, aspect ratio of 4)
and the vertical distance covered by the device from the start of the ascension to the closure of the top
cover (1800 m). Blank casts were performed with the bottle-net closed during the entire cast to assess
for potential contaminations, and blanks were subtracted.
417
418
419

nifH gene sequencing and bioinformatics

The *nifH* gene was sequenced from a total of 71 samples: 18 samples collected from sediment traps deployed at 170 m, 270 m and 1000 m at S05M and S10M, and 53 samples collected from the suspended, slow sinking and fast sinking fractions of particles collected with the MSC. To that end, DNA was extracted using the DNeasy Plant Mini Kit (Qiagen, Courtaboeuf, France) with additional freeze-thaw bead beating and proteinase K steps before the kit purification[59]. Triplicate nested PCR reactions were conducted using degenerate *nifH* primers[60]. The PCR mix was composed of 5 µL of 5X MyTaq red PCR buffer (Bioline), 1.25 µL of 25 mM MgCl$_2$, 0.5 µl of 20 µM forward and reverse primers, 0.25 µL Platinum Taq and 5 µL of DNA extract (1 µL on second round). The reaction volume was adjusted to 25 µL with PCR grade water. Triplicate PCR products were pooled and purified using the Geneclean Turbo kit (MP Biomedicals). Partial adapters were added by ligation at the sequencing facility (Genewiz) and llumina MiSeq 2x300 paired end sequenced. Demultiplexed paired-end sequences were dereplicated, denoised, assembled and chimeras discarded using the DADA2 pipeline[61]. Fifty-thousand to 60,000 reads were obtained per sample. In total, >2.9 millions of high quality *nifH* sequences were obtained resulting in 14925 ASVs (submitted to NCBI with accession numbers xx-xx). ASVs were annotated down to the genus level using a DADA2 formatted *nifH* gene database (https://github.com/moyn413/nifHdada2). Sequences were grouped into 17 genus according to the database, and sequences not identified to the genus level were grouped as "others". The *nifH* gene was successfully amplified from all samples.

Abundance of diazotrophs et contribution to N export fluxes

The abundance of diazotrophs was determined using TaqMan qPCR assays and previously published primer-probe sets for *Trichodesmium,* UCYN-A1, UCYN-B, UCYN-C and γ-24474A11 (the latter hereafter referred to as Gamma-A) targeting the *nifH* gene[62, 63, 64]. The qPCR was run in 25 µL reactions consisting of 12.25 µL TaqMan PCR Master Mix (Applied Biosystems, Villebon Sur Yvette, France), 1 µL of the forward and reverse primers at 10 µM (HPLC purified, Eurofins, Nantes, France), 0.25 µL probe at 10 µM, 8.25 µL PCR grade water, 0.25 µL bovine serum albumin at 10.08 µg µL^{-1}, and 2 µL standard or template sample. The qPCR program was run on a CFX96 Real-Time System thermal cycler (BioRad, Marnes-la-Coquette, France) and consisted of 2 min at 50ºC, 10 min at 95ºC continued by 45 cycles of 15 s at 95ºC and 1 min at 64ºC. The annealing temperature was changed to 60ºC for UCYN-A1 qPCR runs[64]. Standard dilutions (10^7-10^1 gene copies) were run in duplicate, and samples and no-template controls (NTCs) in triplicate. NTCs did not show any amplification. The efficiency was 98-113%. Inhibition tests were carried out on all samples and each primer-probe set by adding the 2 µL of the 10^5 copies standard to each sample. No inhibition was observed. The limit of detection and detected but not quantifiable limits were 1 and 8 gene copies per reaction, respectively.

The diazotroph turnover rate representing the fraction of surface diazotrophs exported out of the photic layer per day was calculated as follows: Turnover rate (d^{-1}) = export flux/abundance in the photic layer. The proportion of PN collected by sediment traps attributed to diazotrophs was calculated based on estimated PN content per cell. For nanoplanktonic UCYN, cell dimensions were directly measured and their biovolume calculated[65]. The C content per cell was estimated from the biovolume according to Verity et al. (1992)[66] and the N content calculated based on C:N ratios of 5 for UCYN-B [67, 68] and 8.5 for UCYN-C[47]. For the UCYN-A1-host symbiosis, we considered a size of 1 µm for the UCYN-1 and 2 µm for the host and a C:N ratio of 6.3[69]. For *Trichodesmium,* we considered the lower-end value of 10 ng N per *Trichodesmium* filament from the same region[70]. To circumvent the polyploidy of *Trichodesmium*[71], we divided the qPCR-based abundances by 12. This ratio was determined for this study by dividing the qPCR-based abundances by the number of *Trichodesmium* filaments counted at S05M and S10M stations at 6 depths in the photic layer (we considered 100 cells per filament of *Trichodesmium*).

Particle imaging

Seawater samples from traps, MSC and bottle-nets were gently filtered on 0.2 µm (for scanning electron microscopy, SEM) and 2 µm polycarbonate filters (for epifluorescence microscopy) at very low pressure (xxx) to preserve the particle structure. For epifluorescence microscopy, filters were fixed with paraformaldehyde (2% prepared in filtered seawater) for 10 minutes at ambient temperature and stored at-80°C until visualized using a Zeiss Axioplan (Zeiss, Jena, Germany) microscope fitted with a green (510–560 nm) excitation filter, which targeted the phycoerythrin-rich cells. For SEM, samples were fixed with 2.5% glutaraldehyde and 1.6% PFA for 1 h at room temperature. Subsequently, the filters were rinsed twice in 0.2 µm filtered in seawater during 15 min, rinsed in osmium during 30 min, and rinsed thrice in filtered seawater to eliminate excess osmium. Next, the filters passed through a series of ethanol drying solutions (50, 70, 95 and 100%, 10 min each), and a series of HDMS solutions (30, 50,

480 80 and 100%, 10 min each). Finally, the filters were air-dried and stored at room temperature until SEM
481 analyses. The filters were visualized onshore using a Phenom-Pro benchtop scanning electron
482 microscope at 10 kV.
483
484 Table 1. Particulate nitrogen (PN) export fluxes at stations S05M and S10M at 170 m, 270 m and 1000
485 m, and contribution of diazotrophs to total PN export.

Station	S05M			S10M		
Depth of deployment	PN export flux	Diazotroph-PN	%Contribution diazotrophs	PN export flux	Diazotroph-PN	%Contribution diazotrophs
	mg N m^{-2} d^{-1}	mg N m^{-2} d^{-1}	%	mg N m^{-2} d^{-1}	mg N m^{-2} d^{-1}	%
170 m	4.9±0.6	0.6	13	13.4±1.3	7.6	6
270 m	6.1±1.2	1.9	30	9.7±0.6	7.8	8
1000 m	1.6±0.2	12.3	100	5.2±0.1	2.3	45

486
487
Figure 1. a. Surface chlorophyll MODIS composite averaged over the time period corresponding to the
TONGA cruise (1 November-6 December 2019) at a resolution of 4 km. Black triangles correspond to
stations where surface-tethered drifting sediment traps were deployed (170 m, 270 m, 1000 m). Grey
dots correspond to stations where Marine Snow Catcher (MSC) casts were performed at three depths
(see Methods), and white dots to MSC casts performed at one depth (200 m). Black circles
correspond to stations where the bottlenet profiles were performed between 2000 m and 200 m. b.
Abundances of the five *nifH* phylotypes (Log10 *nifH* gene copies L^{-1}) studied during the cruise over
the transect (averages over the photic layer, ~0-130 m).
496
497 Figure 2. Relative abundance of diazotroph genera on sediment trap samples. The dendrogram
498 corresponds to the clustering of samples generated by Bray-Curtis similarity. The legend indicates the
499 *nifH* genus affiliation.
500
501 Figure 3. a. Export flux (*nifH* gene copies m^{-2} d^{-1}) of the five diazotroph groups targeted by qPCR (UCYN-
502 A1 symbiosis, UCYN-B, UCYN-C, *Trichodesmium* and Gamma-A) in sediment trap samples at 170 m,
503 270 m and 1000 m at stations S05M and b. at S10M. Error bars represent strandard deviations from
504 triplicate aliquote analyzed in duplicates. c. Export turnover rates (d^{-1}) of the same diazotroph groups
505 (average of the three depths and the two stations for each group).
506
507 Figure 4. Microscopy images showing examples of phycoerythrin-containing UCYN-like cells and
508 *Trichodesmium* in sediment trap samples collected at 170 m, 270 m, and 1000 m at stations S05M and
509 S10M. a-f. Images taken by epifluorescence microscopy (green excitation 510–560 nm, scale bar: 50
510 μm). k-l. Image taken by scanning electron microscopy (SEM).
511
512 Figure 5. Relative abundance of diazotroph genera from marine snow catcher samples. The
513 dendrogram corresponds to the clustering of samples generated by Bray-Curtis similarity. The legend
514 indicates the *nifH* genus affiliation.
515
Figure 6. a-e. Quantification of the five diazotroph groups targeted by qPCR in Marine Snw catcher
samples: UCYN-A1 symbiosis: *Trichodesmium*, UCYN-B, UCYN-C, and Gamma-A in the suspended
(dark color), slow sinking (medium color) and fast sinking (light color) pools of the MSC samples across
the transect (left Y axis, *nifH* gene copies L^{-1}). Dots represent the integrated pool of each phylotype
(right Y axix, *nifH* gene copies m^{-2}) in the photic (~0-130 m) layer. f. Percentage of sinking versus non-
sinking diazotrophs for each group.
522
523 Figure 7. Abundance (*nifH* gene copies m^{-2}) of the five diazotroph groups targeted by qPCR (UCYN-A1
524 symbiosis, *Trichodesmium*, UCYN-B, UCYN-C, and Gamma-A) in the bottle-net samples across the
525 transect, representing an integrated sample from 200 m to 2000 m.
526

Acknowledgements
This research is a contribution of the TONGA project (Shallow hydroThermal sOurces of trace elemeNts:
potential impacts on biological productivity and the bioloGicAl carbon pump; TONGA cruise DOI:
10.17600/18000884) funded by the Agence Nationale de la Recherche (grant ANR-18- CE01-0016),
the LEFE-CyBER program (CNRS-INSU), the A-Midex foundation, the Institut de Recherche pour le
Développement (IRD). The authors thank the crew of the R/V L'Atalante for outstanding shipboard
operations. Nagib Bhairy is warmly thanked for their efficient help with MSC deployment and clean CTD
rosette management and Vincent Taillandier is tanked for CTD data processing.

Author contributions

SB designed the experiments and SB, MB and IBF carried them out at sea, with advice from FLM; SB, MB, MC, OG, AT, DS analyzed the samples; SB analyzed the data with the help of MB for the sequencing data. SB prepared the manuscript with contributions of all co-authors.

References

1. Moore CM*, et al.* Processes and patterns of oceanic nutrient limitation. *Nature Geoscience* **6**, 701–710 (2013).

2. Gruber N. The marine nitrogen cycle: overview and challenges. In: *Nitrogen in the marine environment*) (2008).

3. Eppley RW, Peterson BJ. The flux of particulate organic matter to the deep ocean and its relation to planktonic new production. *Nature* **282**, 677-680 (1979).

4. Mulholland MR. The fate of nitrogen fixed by diazotrophs in the ocean. *Biogeosciences* **4**, 37-51 (2007).

5. Adam B*, et al.* N2-fixation, ammonium release and N-transfer to the microbial and classical food web within a plankton community. *The ISME journal* **10**, 450-459 (2016).

6. Bonnet S*, et al.* Diazotroph derived nitrogen supports diatom growth in the South West Pacific: A quantitative study using nanoSIMS. *Limnology and Oceanography* **61**, 1549-1562 (2016).

7. Robinson C*, et al.* Mesopelagic zone ecology and biogeochemistry - a synthesis. *Deep-Sea Research Part Ii-Topical Studies in Oceanography* **57**, 1504-1518 (2010).

8. Richardson TL, Jackson GA. Small phytoplankton and carbon export from the surface ocean. *Science* **315**, 838-840 (2007).

9. Subramaniam A*, et al.* Amazon River enhances diazotrophy and carbon sequestration in the tropical North Atlantic Ocean. *Proceedings of the National Academy of Sciences* **105**, 10460–10465 (2008).

10. White AE*, et al.* Nitrogen fixation in the Gulf of California and the Eastern Tropical North Pacific. *Progess in Oceanography* **109**, 1-17 (2012).

11. Karl DM, Church MJ, Dore JE, Letelier R, Mahaffey C. Predictable and efficient carbon sequestration in the North Pacific Ocean supported by symbiotic nitrogen fixation. *Proceedings of the National Academy of Sciences* **109**, 1842–1849 (2012).

12. Luo YW*, et al.* Database of diazotrophs in global ocean: abundances, biomass and nitrogen fixation rates. *Earth System Science Data* **5**, 47-106 (2012).

13. Sharek RM, Tupas LM, Karl DM. Diatom fluxes to the deep sea in the oligotrophic North Pacific gyre at Station ALOHA. *Marine and Ecological Progress Series* **82**, 55-67 (1999).

14. Walsby AE. The gas vesicles and buoyancy of *Trichodesmium*. *Marine Pelagic Cyanobacteria: Trichodesmium and other Diazotrophs*, 141-161 (1992).

15. Chavez FP, Ryan J, Lluch-Cota SE, Niquen M. From anchovies to sardines and back: Multidecadal change in the Pacific Ocean. *Science* **299**, 217-221 (2003).

16. Guidi L*, et al.* Does eddy-eddy interaction control surface phytoplankton distribution and carbon export in the North PacificSubtropical Gyre? *Journal of Geophysical Research* **117**, (2012).

17. Caffin M*, et al.* N2 fixation as a dominant new N source in the western tropical South Pacific Ocean (OUTPACE cruise). *Biogeosciences* **15**, 2565-2585 (2018).

595   18.   Agusti S, *et al.* Ubiquitous healthy diatoms in the deep sea confirm deep carbon injection by the
596   biological pump. *Nat Commun* **6**, 7608 (2015).
597
598   19.   Pabortsava K, *et al.* Carbon sequestration in the deep Atlantic enhanced by Saharan dust.
599   *Nature Geoscience* **10**, 189-U141 (2017).
600
601   20.   Berthelot H, *et al.* Dinitrogen fixation and dissolved organic nitrogen fueled primary production
602   and particulate export during the VAHINE mesocosm experiment (New Caledonia lagoon).
603   *Biogeosciences* **12**, 4099-4112 (2015).
604
605   21.   Bonnet S, *et al.* Dynamics of N2 fixation and fate of diazotroph-derived nitrogen in a low nutrient
606   low chlorophyll ecosystem: results from the VAHINE mesocosm experiment (New Caledonia)
607   *Biogeosciences* **13**, 2653-2673 (2016).
608
609   22.   Farnelid H, *et al.* Diverse diazotrophs are present on sinking particles in the North Pacific
610   Subtropical Gyre. *Isme Journal* **13**, 170-182 (2019).
611
612   23.   Moran XAG, Lopez-Urrutia A, Calvo-Diaz A, Li WKW. Increasing importance of small
613   phytoplankton in a warmer ocean. *Global Change Biology* **16**, 1137-1144 (2010).
614
615   24.   Polovina JJ, Howell EA, Abecassis M. Ocean's least productive waters are expanding.
616   *Geophysical Research Letters* **35**, (2008).
617
618   25.   Bonnet S, Caffin M, Berthelot H, Moutin T. Hot spot of N2 fixation in the western tropical South
619   Pacific pleads for a spatial decoupling between N2 fixation and denitrification. *Proceedings of*
620   *the National Academy of Sciences* **114**, E2800-E2801 (2017).
621
622   26.   de Boyer Montégut C, Madec G, Fischer AS, Lazar A, Iudicone D. Mixed layer depth over the
623   global ocean: An examination of profile data and a profile-based climatology. *Journal of*
624   *Geophysical Research: Oceans* **109**, (2004).
625
626   27.   Hewson I, Moisander PH, Morrison AE, Zehr JP. Diazotrophic bacterioplankton in a coral reef
627   lagoon: phylogeny, diel nitrogenase expression and response to phosphate enrichment. *Isme*
628   *Journal* **1**, 78-91 (2007).
629
630   28.   Turk-Kubo KA, Frank IE, Hogan ME, Desnues A, Bonnet S, Zehr JP. Diazotroph community
631   succession during the VAHINE mesocosms experiment (New Caledonia Lagoon).
632   *Biogeosciences* **12**, 7435-7452 (2015).
633
634   29.   Riley JS, Sanders R, Marsay C, Le Moigne FAC, Achterberg EP, Poulton AJ. The relative
635   contribution of fast and slow sinking particles to ocean carbon export. *Global Biogeochemical*
636   *Cycles* **26**, (2012).
637
638   30.   Cornejo-Castillo FM, Zehr JP. Intriguing size distribution of the uncultured and globally
639   widespread marine non-cyanobacterial diazotroph Gamma-A. *The ISME journal* **15**, 124-128
640   (2021).
641
642   31.   Durkin CA, Van Mooy BAS, Dyhrman ST, Buesseler KO. Sinking phytoplankton associated with
643   carbon flux in the Atlantic Ocean. *Limnology and Oceanography* **61**, 1172-1187 (2016).
644
645   32.   Cornejo-Castillo FM. Diversity, ecology and evolution of marine diazotrophic microorganisms.).
646   Polytechnic Univ. of Catalonia (UPC) (2017).
647
648   33.   Karlusich JJP, *et al.* Global distribution patterns of marine nitrogen-fixers by imaging and
649   molecular methods. *BioRxiv*, (2020).
650
651   34.   Harding K, Turk-Kubo KA, Sipler RE, Mills MM, Bronk DA, Zehr JP. Symbiotic unicellular
652   cyanobacteria fix nitrogen in the Arctic Ocean. *Proceedings of the National Academy of*
653   *Sciences of the United States of America* **115**, 13371-13375 (2018).
654

35. Shiozaki T*, et al.* Diazotroph community structure and the role of nitrogen fixation in the nitrogen
cycle in the Chukchi Sea (western Arctic Ocean). *Limnology and Oceanography* **63**, 2191-2205
(2018).

36. Poff KE, Leu AO, Eppley JM, Karl DM, DeLong EF. Microbial dynamics of elevated carbon flux
in the open ocean's abyss. *Proceedings of the National Academy of Sciences of the United*
States of America **118**, (2021).

37. Sohm JA, Edwards BR, Wilson BG, Webb EA. Constitutive Extracellular Polysaccharide (EPS)
Production by Specific Isolates of *Crocosphaera watsonii*. *Frontiers in microbiology* **2**, 229
(2011).

38. Berman-Frank I, Dubinsky Z. Balanced growth in aquatic plants: Myth or reality? Phytoplankton
use the imbalance between carbon assimilation and biomass production to their strategic
advantage. *Bioscience* **49**, 29-37 (1999).

39. Walsby AE. The properties and buoyancy-providing role of gas vacuoles in Trichodesmium
Ehrenberg. *European Journal of Phycology* **13**, 103-116 (1978).

40. Berman-Frank I, Bidle KD, Haramaty L, Falkowski PG. The demise of the marine
cyanobacterium, Trichodemsium spp., via an autocatalyzed cell death pathway. *Limnology and*
Oceanography **49**, 997-1005 (2004).

41. Berman-Frank I, Rosenberg G, Levitan O, Haramaty L, X. M. Coupling between autocatalytic
cell death and transparent exopolymeric particle production in the marine cyanobacterium
Trichodesmium. *Environmental microbiology* **9**, 1415-1422 (2007).

42. Bar-Zeev E, Avishay I, Bidle KD, Berman-Frank I. Programmed cell death in the marine
cyanobacterium Trichodesmium mediates carbon and nitrogen export. *ISME Journal* **7**, 2340-
2348 (2013).

43. Buesseler KO. The decoupling of production and particulate export in the surface ocean. *Global*
Biogeochemical Cycles **12**, 297-310 (1998).

44. Buesseler KO, Boyd PW, Black EE, Siegel DA. Metrics that matter for assessing the ocean
biological carbon pump. *Proceedings of the National Academy of Sciences of the United States*
of America **117**, 9679-9687 (2020).

45. Benavides M*, et al.* Staying alive: fast sinking Trichodesmium remains active in the twilight zone.
(In prep.).

46. Ababou FE, Le Moigne F, Bonnet S. Mechanistic understanding of diazotroph aggregation: a
rolling tank aproach. *Targetted journal: Limnology and Oceanography*, (In prep.).

47. Berthelot H, Bonnet S, Camps M, Grosso O, Moutin T. Assessment of the dinitrogen released
as ammonium and dissolved organic nitrogen by unicellular and filamentous marine
diazotrophic cyanobacteria grown in culture. *Frontiers in Marine Science* **2**, (2015).

48. Glibert PM, Bronk D. Release of dissolved organic nitrogen by marine diazotrophic
cyanobacteria, *Trichodesmium* spp. *Applied and Environmental Microbiology* **60**, 3996-4000
(1994).

49. Mulholland MR, Bernhardt PW, Heil CA, Bronk DA, O'Neil JM. Nitrogen fixation and
regeneration in the Gulf of Mexico. *Limnology and Oceanography* **51**, 176-177 (2006).

50. Capone DG*, et al.* An extensive bloom of the N_2-fixing cyanobacterium, *Trichodesmiumn*
erythraeum, in the Central Arabian Sea. *Marine Ecology Progress Series* **172**, 281-292 (1998).

51. Steinberg DK, Landry MR. Zooplankton and the Ocean Carbon Cycle. *Annu Rev Mar Sci* **9**,
413-444 (2017).

52. Engel A, Wagner H, Le Moigne FAC, Wilson ST. Particle export fluxes to the oxygen minimum zone of the eastern tropical North Atlantic. *Biogeosciences* **14**, 1825-1838 (2017).

53. Berthelot H, Benavides M, Moisander PH, Grosso O, Bonnet S. High nitrogen fixation rates in the particulate and dissolved pools in the Western Tropical Pacific (Solomon and Bismarck Seas). *Geophysical Research Letters*, (2017).

54. Bonnet S, *et al.* In-depth characterization of diazotroph activity across the western tropical South Pacific hotspot of N-2 fixation (OUTPACE cruise). *Biogeosciences* **15**, 4215-4232 (2018).

55. Conte MH, Ralph N, Ross EH. Seasonal and interannual variability in deep ocean particle fluxes at the Oceanic Flux Program (OFP)/Bermuda Atlantic Time Series (BATS) site in the western Sargasso Sea near Bermuda. *Deep-Sea Research Part Ii-Topical Studies in Oceanography* **48**, 1471-1505 (2001).

56. Ras J, Claustre H, Uitz J. Spatial variability of phytoplankton pigment distributions in the Subtropical South Pacific Ocean: comparison between in situ and predicted data. *Biogeosciences* **5**, 353-369 (2008).

57. Lampitt RS, Wishner KF, Turley CM, Angel MV. Marine Snow Studies in the Northeast Atlantic-Ocean - Distribution, Composition and Role as a Food Source for Migrating Plankton. *Marine Biology* **116**, 689-702 (1993).

58. Passow U, Alldredge AL. A dye-binding assay for the spectrophotometric measurement of transparent exopolymer particles (TEP). *Limnology and Oceanography* **40**, 1326-1335 (1995).

59. Moisander PH, Beinart RA, Voss M, Zehr JP. Diversity and abundance of diazotrophic microorganisms in the South China Sea during intermonsoon. *Isme Journal* **2**, 954-967 (2008).

60. Zehr JP, Turner PJ. Nitrogen fixation: Nitrogenase genes and gene expression. In: *Methods in Marine Microbiology* (ed Paul JH). Academic Press (2001).

61. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods* **13**, 581-+ (2016).

62. Church MJ, Jenkins BD, Karl DM, Zehr JP. Vertical distributions of nitrogen-fixing phylotypes at Stn ALOHA in the oligotrophic North Pacific Ocean. *Aquatic Microbial Ecology* **38**, 3-14 (2005).

63. Moisander AM, Beinart A, Voss M, Zehr JP. Diversity and abundance of diazotrophs in the South China Sea during intermonsoon. *The ISME journal* **2**, 954-967 (2008).

64. Thompson A, Carter BJ, Turk-Kubo K, Malfatti F, Azam F, Zehr JP. Genetic diversity of the unicellular nitrogen-fixing cyanobacteria UCYN-A and its prymnesiophyte host. *Environmental microbiology* **16**, 3238–3249 (2014).

65. Sun J, Liu D. Geometric models for calculating cell biovolume and surface area for phytoplankton. *Journal of Plankton Research* **25**, 1331-1346 (2003).

66. Verity PG, Robertson CY, Tronzo CR, Andrews MG, Nelson JR, Sieracki ME. Relationships between cell volume and the carbon and nitrogen content of marine photosynthetic nanoplankton. *Limnology and Oceanography* **37**, 1434–1446 (1992).

67. Dekaezemacker J, Bonnet S. Sensitivity of N_2 fixation to combined nitrogen forms (NO_3^- and NH_4^+) in two strains of the marine diazotroph *Crocosphaera watsonii* (Cyanobacteria). *Marine Ecology Progress Series* **438**, 33-46 (2011).

68. Knapp AN, Dekaezemacker J, Bonnet S, Sohm JA, Capone DG. Sensitivity of *Trichodesmium erythraeum* and *Crocosphaera watsonii* abundance and N_2 fixation rates to varying NO_3^- and PO_4^{3-} concentrations in batch cultures. *Aquatic Microbial Ecology* **66**, 223-236 (2012).

775
776   69.   Martinez-Perez C, *et al.* The small unicellular diazotrophic symbiont, UCYN-A, is a key player
777   in the marine nitrogen cycle. *Nature Microbiology* **1**, 16163 (2016).
778
779   70.   Rodier M, Le Borgne R. Population and trophic dynamics of Trichodesmium thiebautii in the SE
780   lagoon of New Caledonia. Comparison with T. erythraeum in the SW lagoon. *Marine Pollution*
781   *Bulletin* **61**, 349-359 (2010).
782
783   71.   Sargent EC, *et al.* Evidence for polyploidy in the globally important diazotroph Trichodesmium.
784   *FEMS microbiology letters* **363**, (2016).
785
786

a

Traps ▲
MSC ⬤
Bottle-net ⭕

New Caledonia

Fiji

S11 S10V
S10M
S04 S07 S08
S01 S02 S03
Tonga
S05V S05M

Longitude

Latitude

Chl (mg m⁻³)

Chl (mg m^{-3})

b

Log NifH copies L⁻¹

S01 S02 S03 S04 S05V S05M S11 S10M S10V S07 S08

● UCYN-A1 ● Trichodesmium ● UCYN-B ● UCYN-C ● Gamma-A

Log10 *nifH* gene copies m^{-2} d^{-1}

a

| | 1,0E+00 | 2,0E+02 | 4,0E+04 | 8,0E+06 | 1,6E+09 |

b

| | 1,0E+00 | 2,0E+02 | 4,0E+04 | 8,0E+06 | 1,6E+09 |

c

Depth (m)

170 m

270 m

1000 m

Diazotroph group

■ UCYN-A ■ Trichodesmium ■ UCYN-B ■ UCYN-C ■ Gamma-A