BAYESIAN ADDITIVE REGRESSION TREES FOR GENOTYPE BY ENVIRONMENT INTERACTION MODELS

BY DANILO A. SARTI[1], ESTEVÃO B. PRADO[1,2], ALAN N. INGLIS[1,2], ANTÔNIA A. L. DOS SANTOS[1], CATHERINE B. HURLEY[1], RAFAEL A. MORAL[1], ANDREW C. PARNELL[1,2]

[1]*Hamilton Institute, Department of Mathematics and Statistics, Maynooth University, Ireland*

[2]*Insight Centre for Data Analytics, Maynooth University, Ireland*

We propose a new class of models for the estimation of genotype by environment (GxE) interactions in plant-based genetics. Our approach, named AMBARTI, uses semi-parametric Bayesian additive regression trees to accurately capture marginal genotypic and environment effects along with their interaction in a fully Bayesian model. We demonstrate that our approach is competitive or superior to similar models widely used in the literature via both simulation and a real world data set. Furthermore, we introduce new types of visualisation to properly assess both the marginal and interactive predictions from the model. An R package that implements our approach is available at `https://github.com/ebprado/ambarti`.

1. Introduction. The interaction between genotypes and environments (GxE) is a key parameter in plant breeding (Allard and Bradshaw, 1992). Poor understanding of GxE can lead to sub-optimal selection of new genotypes and inbred lines. Understanding the GxE interactions is crucial for germplasm management, having strong genetic and economic impacts on seed production and crop yield (Sarti, 2013). Many models have been proposed for studying GxE in the context of multi-environmental experiments (METs) (Sarti, 2019). One special case is the additive main effects multiplicative interactions model (AMMI) (Mandel, 1971).

The classical AMMI models combine features of analysis of variance (ANOVA) with a bilinear term to represent GxE interactions. Such interactions will be named here bilinear interactions. In addition, AMMI models allow for estimation of main effects of genotypes and environments as well as the decomposition of the interaction through a bilinear term. Many extensions to the AMMI models have been proposed, including robust AMMI (Rodrigues, Monteiro and Lourenço, 2016) and weighted AMMI (Sarti, 2019).

Bayesian additive regression trees (BART) (Chipman et al., 2010) is a non-parametric Bayesian model that generates a set of trees and uses random splitting rules to produce predictions for a univariate response. Given its flexibility to deal with non-linear structures and richer, non-multiplicative interactions terms, the use of BART and its extensions has increased with applications in many areas including proteomic studies (Hernández et al., 2018), hospital performance evaluation (Liu et al., 2015), credit scores (Zhang and Härdle, 2010) among many others.

In this paper, we extend the AMMI model to allow for richer GxE interactions, and similarly sidestep the model choice complexity term present in all AMMI-type approaches. We achieve this goal by including a new variant of BART, which we term 'double-grow' BART. The new proposed method, named AMBARTI, provides a fully Bayesian joint model, where the 'double-grow' BART component is solely responsible for GxE interactions.

Keywords and phrases: Bayesian non-parametric regression, Bayesian additive regression trees, additive main effects multiplicative interactions model, genotype-by-environment interactions.

We compare our newly proposed model with the traditional AMMI approaches and other competing interaction detection models, and we show that its performance is superior (judged on out-of-sample error) in both simulated and real-world example data. The real data set we use is taken from the value of cultivation and usage (VCU) experiments of the Irish Department of Agriculture, which were conducted in the years between 2010 and 2019. Furthermore, the output of AMBARTI leads us to suggest several new forms of visualisation that are easier to interpret for non-specialists.

The paper is structured as follows. In Section 2, we describe the framework used to collect evidence from METs, including the classic genetic equation to describe the relationship between phenotypes, genotypes, and environments. We also outline the formulation of the AMMI model in its classic form. In Section 2.3, we briefly describe the standard Bayesian additive regression trees model. In Section 2.4, we present the structure of our novel AMBARTI approach. Sections 3 and 4 contain the main results from the simulation experiments and real data sets, respectively. In Section 5, we conclude with a discussion and outline further opportunities.

2. Methods.

2.1. *GxE interactions and MET.* The phenotypic expression of a genetic character can be theoretically decomposed in terms of genetic factors, environmental factors, and the interactions between them as

$$p = g + e + (ge),$$

where p is the phenotypic response, g is the genetic factor, e is the environmental factor, and (ge) is the interaction between genotypes and environments. The last term is necessary due to the different response of genotypes across different environments. The presence of GxE interactions implies that each genotype may have different phenotypic responses across a set of environments. If we produce a rank that orders the performance of each genotype into each environment, we will notice that the order of the best to worst genotype is different across the environments. The presence of GxE interactions is known to be capable of having large effects on the phenotypic response (Falconer and Mackay, 1996; Dias, 2005).

The (ge) terms can be estimated in a MET design where several environments and genotypes are evaluated for a given phenotype (Isik, Holland and Maltecca, 2017). In plant breeding, the need for METs is constant given the fact that the germplasm generates new genotypes every year and the pressure of diseases and other factors are dynamic. Such experiments require a complex set of logistical activities, leading to high costs of implementation. These trials thus have strong regulatory appeal in the seed and biotech industries around the world (Sarti, 2013).

Reliable information about GxE can help breeders make decisions on cultivar recommendations. Thus, models for the study of GxE need to be able to answer questions such as which genotypes can perform well across a set of environments and which are specifically recommended for a given environment. The answers to these questions are crucial both to broad breeding strategies, i.e., to obtain one or more genotypes that perform well in a set of environments, and to target breeding, where we determine the best genotype for a given environment (Sarti, 2019).

2.2. *Traditional AMMI models.* A simple statistical linear model can be used to estimate GxE effects from METs. The model can be written as

$$(1) \qquad y_{ij} = \mu + g_i + e_j + (ge)_{ij} + \epsilon_{ij},$$

where $\epsilon_{ij} \sim \mathrm{N}(0, \sigma^2)$, $i = 1, \ldots, I$, $j = 1, \ldots, J$, g_i is the effect of genotype i, e_j is the effect of environment j, and $(ge)_{ij}$ represents the interaction between genotype i and environment j.

In the specification of the Equation 1, the term (ge) can be thought of as representing a decomposition of the residual from a more basic linear model. Gollob (1968) and Mandel (1971) proposed a method to decompose the residual term as a sum of multiplicative factors that includes the (ge) term. This yields the decomposition:

$$(ge)_{ij} = \sum_{q=1}^{Q} \lambda_q \gamma_{iq} \delta_{jq}, \tag{2}$$

where Q is the number of components to be considered in the analysis, λ_q is the strength of the interaction of component q, γ_{iq} represents the importance of genotype i in component q, and δ_{jq} represents the importance of environment j in component q; see Appendix B for the restrictions imposed on γ_{iq}, δ_{jq} and λ_q to make the model identifiable. Hence, the complete AMMI model is written as

$$y_{ij} = \mu + g_i + e_j + \sum_{q=1}^{Q} \lambda_q \gamma_{iq} \delta_{jq} + \epsilon_{ij}. \tag{3}$$

The interaction terms in (3) are estimated by a singular value decomposition (SVD) of the matrix of means of genotypes into environments *(GE)*. Here λ_q is the q-th eigenvalue of the matrix *(GE)*, γ_{ik} is the *i*-th element of the left singular vector and δ_{jk} is *j*-th element of the right singular vector obtained in the SVD (Good, 1969). In practice, the classical AMMI model can be run in R (R Core Team, 2020) using the package `agricolae` (de Mendiburu, 2019) or via functions programmed by the user as in Onofri and Ciriciofolo (2007).

The protocol for estimation of the terms in a standard AMMI model is given by Gauch Jr (2013). This involves the following steps:

1. Obtain the grand mean and principal effects of the genotypes and environments using ANOVA with two factors based on a matrix of means containing the means of each genotype within each environment;
2. Obtain the residuals from the model above that will comprise the interaction matrix, where each row is an environment and each column a genotype;
3. Choose an appropriate value for the number of components Q;
4. Form the multiplicative terms that represent the reduced-dimension interactions via an SVD of the matrix of interaction residuals.

The rank of the matrix *(GE)* is assumed to be $r = \min(I - 1, J - 1)$. Thus, the number of components Q may vary from $1, \ldots, r$. The term $\min(I - 1, J - 1)$ establishes the minimum number of non-zero eigenvectors to be obtained in the SVD. Taking $Q = r$, the AMMI model would capture all the variance related to the interaction, and it would result in over-fitting. This problem is ameliorated by using a limited number of components Q. The choice of Q is related to the amount of total variability captured by the principal components (PCs) and, in general, is recommended to use a number of PCs that captures at least 80% of the total variability (Lal et al., 2020; Shafii and Price, 1998; Love et al., 2004; Tyagi et al., 2016; Dias and Krzanowski, 2006). Usually, the value of Q varies from 1 to 3.

AMMI models have been extensively used for evaluation of phenotype performance of cultivars. Nachit et al. (1992) used AMMI models to assess the performance of wheat germplasm from International Maize and Wheat Improvement Center. Farshadfar and Sutka (2003) explored Quantitative Trait Loci (QTL) related to adaptation in Wheat. Rad et al. (2013) studied GxE for Wheat in the context of drought and normal conditions. Brancourt-Hulmel and

Lecomte (2003) evaluated the impact of environmental conditions in the stability of winter Wheat. Badu-Apraku et al. (2012) used AMMI to study the stability of early Maize genotypes in Africa. Mitroviaã et al. (2012) evaluated the performance of experimental maize hybrids using AMMI models, and Sarti (2019) studied the performance of the AMMI model in the context of simulated data. The applications of AMMI models can also be found in several other species including: a) rice (Mahalingam et al., 2006), b) barley (Mahalingam et al., 2006; Romagosa et al., 1996; Sato and Takeda, 1993; Anbessa et al., 2009), and c) sugarcane Silveira et al. (2013).

2.3. *Tree-based methods and BART.* Introduced by Chipman et al. (2010), BART is a Bayesian model that uses a sum of trees to approximate a univariate response. In BART, each tree works as a weak learner that yields a small contribution to the final prediction. Based on a design matrix $\mathbf{X}$, BART is able to capture interactions and non-linear relationships. The BART model can be written as

$$(4) \qquad y_i|\mathbf{x}_i, \mathcal{M}_t, \mathcal{T}_t, \sigma^2 \sim \mathrm{N}\left(\sum_{t=1}^{T} h(\mathbf{x}_i, \mathcal{M}_t, \mathcal{T}_t), \sigma^2\right), \ i = 1, \ldots, n,$$

where $\mathbf{x}_i$ is the i-th row of the design matrix $\mathbf{X}$, $\mathcal{M}_t$ denotes the set of terminal node parameters of tree t, $\mathcal{T}_t$ is the set of binary splitting rules that define the tree t, and $h(\cdot) = \mu_{t\ell}$ is a function that assigns the predicted values $\mu_{t\ell} \in \mathcal{M}_t$ based on the design matrix $\mathbf{X}$ and tree structure $\mathcal{T}_t$. The number of trees T can be chosen so that non-linear and interaction effects are properly estimated, but it can also be selected through cross-validation; Chipman et al. (2010) recommends $T = 200$ as a default.

Unlike other tree-based methods where a loss function is optimised to grow the trees, in BART the trees are grown using Markov Chain Monte Carlo (MCMC) (Gamerman and Lopes, 2006; Robert and Casella, 2013). The trees are either accepted or rejected via a Metropolis-Hastings step. In addition, the trees can be learned by four moves: grow, prune, change or swap. It is important to highlight that in all moves the splitting rule is defined by randomly selecting one covariate and one split point. In the grow move, a terminal node is selected and then two children nodes are created below it. When pruning, a parent of two terminal nodes is selected and its children nodes are removed. During the change process, a parent node is picked and its splitting rule (i.e., covariate and split point) is redefined. In the swap move, a pair of parents of terminal nodes is chosen and their splitting rules are swapped.

As a fully Bayesian model, BART assumes prior distributions on all quantities of interest. First, the node-level parameters $\mu_{t\ell}$ are assumed to be i.i.d $\mathrm{N}(0, \sigma_\mu^2)$, where $\sigma_\mu = 0.5/(k\sqrt{T})$ and $k \in [1, 3]$. Second, the residual variance σ^2 is assumed to be distributed as $\mathrm{IG}(\nu/2, \nu\lambda/2)$, where $\mathrm{IG}(\cdot)$ denotes an Inverse Gamma distribution. Third, to control how shallow/deep a tree may be, each non-terminal node has a prior probability of $\alpha(1+d)^\beta$ of being observed, where $\alpha \in (0, 1)$, $\beta \geq 0$, and d corresponds to the depth of the node; Chipman et al. (2010) recommends $\alpha = 0.95$ and $\beta = 2$ as default values. These hyperparameter values tend to select trees which are not too deep.

We remark that the priors above are a crucial element of the BART model to identify σ^2 as they control how deep or shallow the trees can be, and how large the terminal node predictions are. Were these priors to be set too vague (e.g., so that deep trees were favoured), then the value of σ^2 would shrink towards zero and we would be left with an over-fitted model. Thus, the priors force the trees to be shallow, and such that each tree only explains a small component of the data, which leave some variation in the residuals and resolves the identifiability issue of σ^2.

The identification of the interaction effects in BART is powered by the full conditional of the trees, which we denote by $p(\mathcal{T}_t|R_t, \sigma^2)$, where $R_t = \mathbf{y} - \sum_{k\neq t}^{T} h(\mathbf{X}; \mathcal{T}_k, \mathcal{M}_k)$ represents

the vector of the partial residuals excluding tree t; see Appendix A for details. As in BART the trees are learned by using splitting rules that are created by randomly selecting a covariate and a split point, $p(\mathcal{T}_t | R_t, \sigma^2)$ is utilised to select only 'good' trees (i.e., trees that help reduce the residual variance).

Finally, the structure of the BART model for a continuous response can be summarised as follows. First, all $\mathcal{T}_t$ are initialised as stumps. Then, each tree is learned, one at a time, using one of the four moves previously described (grow, prune, change or swap). Next, the newly proposed tree $\mathcal{T}_t^*$ is compared to its previous version $\mathcal{T}_t$ via a Metropolis-Hastings step taking into account the partial residuals R_t and the structure/depth of $\mathcal{T}_t$ and $\mathcal{T}_t^*$. After that, the predicted values for each terminal node ℓ of the tree t are generated and then σ^2 is updated. For a binary outcome, the data augmentation strategy of (Albert and Chib, 1993) can be used; see Tan and Roy (2019) and Prado et al. (2021) for more details.

Due to its flexibility and excellent performance on regression and classification problems, BART has been applied and extended to credit modelling (Zhang and Härdle, 2010), survival analysis (Sparapani et al., 2016; Linero et al., 2021; Basak et al., 2021), proteomic biomarker analysis (Hernández et al., 2015), polychotomous response (Kindo et al., 2016) and large data sets (Hernández et al., 2018; Linero and Yang, 2018a,b). More recently, papers exploring the theoretical aspects of BART have been developed by Jeong and Ročková (2020); Ročková and van der Pas (2020); Ročková and Saha (2019); Linero and Yang (2018b). In practice, there are many R packages to fit BART, such as BartMachine (Kapelner and Bleich, 2016), BART (McCulloch et al., 2020) and dbarts (Dorie, 2020).

2.4. *AMBARTI.* To insert the BART model inside an AMMI approach, we make some fundamental changes to the way the trees are grown and structured. As a first step, we can write the sum of trees inside the Bayesian version of the AMMI model

$$(5) \qquad y_{ij} | \mathbf{x}_{ij}, \Theta \sim \mathrm{N} \left(\mu + g_i + e_j + \sum_{t=1}^{T} h(\mathbf{x}_{ij}, \mathcal{M}_t, \mathcal{T}_t), \sigma^2 \right),$$

where y_{ij} denotes the response for genotype i and environment j, $\Theta = (\mu, g_i, e_j, \mathcal{M}_t, \mathcal{T}_t, \sigma^2)$, μ is the grand mean, and g_i and e_j denote the effects of genotypes and environments, respectively. The component $\sum_{t=1}^{T} h(\mathbf{x}_i, \mathcal{M}_t, \mathcal{T}_t)$ is the same as presented in (4) and $\mathbf{x}_{ij}$ contains dummy variables that represent combinations of g_i and e_j. In order to get the posterior distribution of the new parameters, we assume that $\mu \sim \mathrm{N}(m, \sigma_m^2)$, $g_i \sim \mathrm{N}(0, \sigma_g^2)$ and $e_j \sim \mathrm{N}(0, \sigma_e^2)$ as well as that $\sigma_g^2 \sim \mathrm{IG}(a_g, b_g)$ and $\sigma_e^2 \sim \mathrm{IG}(a_e, b_e)$.

At first look our model is similar to the semi-parametric BART proposed by Zeldow et al. (2019). However, our approach differs in that i) we do not partition the covariates into two distinct subsets, as the dummy variables (g_i and e_j) that are used in the linear predictor are also contained in $\mathbf{x}_i$; ii) most importantly, we replace the growing move with a 'double grow' (and equivalently 'double prune') step so that we guarantee the trees will include at least one g_i and one e_j as splitting criteria and; iii) unlike Zeldow et al. (2019), we do not use the full residuals $\mathbf{r} = \mathbf{y} - \sum_{t=1}^{T} h(\mathbf{X}, \mathcal{M}_t, \mathcal{T}_t)$ to update the linear predictor estimates, but rather the response variable itself, which is analogous to the two-stage estimation idea from the classical AMMI model, and equivalent to a cut feedback model, following Plummer (2015).

In Zeldow et al. (2019), the trees use the partial and full residuals rather than the response variable to both generate the node-level predictions and update the main effects. However, in the classical AMMI, the estimation of the main effects (g and e) is carried out taking into account the response variable and then the bilinear/interaction term is estimated via an SVD by using the residuals. That is, in the AMMI model the estimation of the model components is carried out in two stages, which does not occur in Zeldow et al. (2019). As

Fig 1: An example of a tree generated from a double grow move based on the sets of dummy variables generated for g and e. $g_{1,3}$ denotes a dummy variable that combines genotypes 1 and 3, and $e_{4,5,6}$ represents a dummy variable that binds environments 4, 5, and 6.

AMBARTI is a combination of BART and AMMI, we estimate the main effects as AMMI and the interactions via BART also using cut-feedback estimation.

The rationale of the 'double' moves is to force the trees to exclusively work on the interactions between g_i and e_j. This removes the chance that they split on a single g_i or e_j variable, which would lead to confounding with the main marginal genomic or environment effects. For example, in the double grow, rather than randomly selecting one covariate and one split point when growing a stump, a variable g^* is chosen and then another variable e^* is randomly selected and both define the splitting rules of the corresponding tree. Here, the dummy variables g^* and e^* are sampled from the sets of all possible combinations of g_i and e_j, respectively.

We allow for extra flexibility in the identification of the interaction effects by setting up the design matrix of the BART model to contain all possible combinations of genotype and environment effects. For instance, when $I = 5$ genotypes and $J = 6$ environments, $2^{I-1} - 1 = 15$ and $2^{J-1} - 1 = 31$ unique dummy variables are generated for g and e, respectively; see Wright and König (2019) for splitting methods on categorical covariates. In this case, there are 46 dummy variables available so that BART can (double) split on them to identify possible interaction between genotypes and environments. To illustrate this, in Figure 1 a dummy variable is sampled from the set of 15 dummy variables related to g along with a dummy variable from the set of 31 dummy variables associated with e.

An appealing advantage of AMBARTI over AMMI is that it does not require the specification of the number of components Q in the bilinear sum and does not require complex orthonormality constraints on the interaction structure; see Appendix B for the constraints of the AMMI model. In a Bayesian context, these constraints can lead to complex prior distribution choices for implementation of AMMI (as in Josse et al., 2014; Crossa et al., 2011). Furthermore, although AMBARTI adds a computational cost to the BART model, we have found this to be negligible for standard MET data sets that usually have values of I and J up to the low tens or hundreds.

An additional advantage of using a fully Bayesian approach as in AMBARTI is that we have access to full posterior distributions of each parameter. As the model is fitted jointly, we are thus able to ascertain the general levels of uncertainty in each g_i or e_j component, which may assist with future experimental designs. Similarly, the interaction term is also estimated probabilistically, and so may avoid interpretation errors associated with, e.g., biplots from a traditional AMMI model.

In terms of estimation, the AMBARTI model can be fitted as follows. First, the parameter estimates g_i and e_j are sampled taking into account the response variable $\mathbf{y}$ (not the residuals). Then, one at a time, the trees are updated via partial residuals $r_t = \mathbf{y} - \hat{\mu} - \hat{g}_i - \hat{e}_j - \sum_{k \neq t}^{T} h(\mathbf{X}, \mathcal{M}_k, \mathcal{T}_k)$. Hence, the terminal node parameters are generated and the sample variance is updated. In the end, posterior samples associated with μ, g_i, e_i, σ_g, σ_e, $\mathcal{T}_t$,

$\hat{\mathbf{y}}$ are available, which allow for the calculation of credible intervals and evaluation of the significance of the parameter estimates; see Algorithm 1 in Appendix A for more details.

2.4.1. *Validity of the posterior sampling in AMBARTI.* The AMBARTI algorithm estimates the genotype and environment main effects taking into account only the response and disregarding the BART component. In contrast, when fitting the BART model, the effects of g and e are taken into account.

We set up AMBARTI in this way as we aimed to compare its results to the frequentist AMMI model. We recall that in the classical AMMI the estimation of g and e are first carried out and then the interactions are estimated based on a singular value decomposition. As stated, AMBARTI is an implementation of the 'naive cut algorithm' described in Plummer (2015). Nonetheless, we show below that our model, under the naive cut algorithm, generates valid samples following Plummer (2015).

A simple way of thinking about our model is, conditioned on knowing the tree structures:

(6)
$$\mathbf{y} = \underbrace{\mathbf{W}\phi}_{\text{g + e}} + \underbrace{\mathbf{X}\theta}_{\text{BART}} + \epsilon,$$

where $\epsilon \sim \mathrm{N}(0, \sigma^2)$, $\mathbf{W}$ is a binary matrix of values allocating $\mathbf{y}$ to the correct genotype/environment, ϕ are the g and e parameters, $\mathbf{X}$ is a matrix of binary interaction identifiers, and θ are parameters of the interactions. Under this formulation, we have that the quantities of interest ϕ and θ can be estimated via their full conditional distributions:

(7)
$$p(\phi|\mathbf{Y}, \mathbf{W}, \theta, \sigma^2) \text{ and}$$

$$p(\theta|\mathbf{Y}, \mathbf{X}, \phi, \sigma^2).$$

Figure 2 presents the AMBARTI model under the naive cut algorithm perspective. The likelihood function depends on θ and ϕ, and both parameters are of interest. The graph is divided in two parts (G_1 and G_2). The idea of the naive cut algorithm is that when constructing the full conditionals for parameters in G_1, likelihood terms involving random variables in G_2 are ignored (Plummer, 2015). The 'cut' conditional distributions for ϕ and θ are given by:

(8)
$$p(\phi|\mathbf{Y}, \mathbf{W}, \sigma^2) \text{ and}$$

$$p(\theta|\mathbf{Y}, \mathbf{X}, \phi, \sigma^2).$$

The full conditional in (7) considers the *full* Bayesian model, whilst in (8) there is no dependence/influence of θ over ϕ. Plummer (2015) describes three situations when the feedback from one component may not be helpful to the other. First, when the relation between the response and θ is speculative. Second, when there is conflict between the sources such that $p(\phi|\mathbf{Y}, \mathbf{W}, \theta, \sigma^2)$ is very different from $p(\phi|\mathbf{Y}, \mathbf{W}, \sigma^2)$. Third, when there are computational issues in terms of convergence and mixing.

Plummer (2015) states that the cut in the feedback of θ into the conditional distribution for ϕ may lead to lack of convergence to the (cut) posterior distribution. However, for the cut algorithm to draw approximate samples from the cut model correctly, at least one of the two conditions below need to be verified.

1. The transition from ϕ to ϕ' cannot cause significant changes in the conditional distribution for θ, i.e., the limit of

$$\frac{p(\theta'|\mathbf{Y}, \mathbf{X}, \phi', \sigma^2)}{p(\theta'|\mathbf{Y}, \mathbf{X}, \phi, \sigma^2)} \to 1$$

Fig 2: Graphical representation of the AMBARTI model.

2. The probability of moving from θ to θ', denoted by $\theta \to \theta'$, does not depend on θ, i.e., the limit of

(9)
$$\frac{p(\theta \to \theta'|\phi)}{p(\theta'|\mathbf{Y},\mathbf{X},\phi,\sigma^2)} \to 1.$$

We remark the AMBARTI model attains the condition in (9), as the probability of $\theta \to \theta'$ depends exclusively on the conditional distribution $p(\theta|\mathbf{Y},\mathbf{X},\phi,\sigma^2)$, which in turn does not depend on the previous θ. To make our point clearer, recall expression (6) and that in BART, a priori, $\mu_{t\ell} \sim \mathrm{N}(0,\sigma_\mu^2)$, which is completely specified with no dependence on its previous values. In addition, the conditional distribution of the predicted values $\mu_{t\ell}$, conditioned on ϕ, has a closed-form expression, which is a Normal distribution presented in Appendix A, that relies only on the partial residuals, number of observations in a node, and the residual variance.

3. Simulation Study. In this Section, we compare AMMI, the Bayesian version of AMMI (B-AMMI) proposed by Josse et al. (2014), and AMBARTI using the root mean square error (RMSE) for predicted values $\hat{y}$ and for the interaction term on out-of-sample data. Our simulation experiment was carried out considering two scenarios. In the first, we simulated from the AMMI model with $Q = \{1,2,3\}$ and then fitted AMBARTI, B-AMMI, and AMMI. In the second scenario, we simulated from the AMBARTI equation and then fitted AMBARTI and three AMMI (and B-AMMI) models with different number of components to describe the interactions (i.e., $Q = \{1,2,3\}$). In both scenarios, we fitted the models to a training set with $I \times J$ observations and evaluated the performance on an out-of-sample test set of the same size.

For both scenarios, we set $I = J = \{10,25\}$, $\mu = 100$ and generated g_i and e_j from $\mathrm{N}(0,\sigma_g^2)$ and $\mathrm{N}(0,\sigma_e^2)$, respectively, where $\sigma_g = \sigma_e = \{1,5\}$. The parameters γ_{ik} and δ_{jk} were generated from $\mathrm{N}(0,1)$ and then the orthornormality constraints were applied following the results presented in Appendix B. In addition, for $Q = 1$, we consider two values for λ (i.e., $\lambda = \{8,12\}$); for $Q = 2$, $\lambda = (\{12,8\},\{12,10\})$ and; for $Q = 3$, $\lambda = \{12,10,8\}$. In the simulation from AMBARTI, we set $T = 200$ trees and generated each tree by using the 'double grow' move considering $2^{I-1} - 1$ possible covariates for g_i and $2^{J-1} - 1$ for e_j.

Finally, the AMMI model used in the simulations is presented in Equation 3, which represents a completely randomised trial design. The AMBARTI model used is shown in Equation 5.

3.1. *Simulation results.* We start simulating synthetic data from the AMMI system, which is the harshest test for the AMBARTI model. Figure 3 shows the RMSE values for $\hat{y}$ based on the out-of-sample sets of three models considering 10 Monte Carlo repetitions. The data sets considered in this Figure were simulated considering $I = 10$ genotypes and $J = 10$ environments, with different values of $Q = \{1, 2, 3\}$, and two values for the genotypic and environmental standard errors $\sigma_g = \{1, 5\}$ and $\sigma_e = \{1, 5\}$, respectively.

As the data were simulated from the AMMI equation, we would expect that the AMMI model would perform exceedingly well, and this is what we see in general considering all the results of Figure 3. More specifically, we can see in the first upper panel that AMBARTI has higher RMSEs compared to AMMI for all values of Q. Further, we see that B-AMMI has similar performance to the frequentist AMMI only for $Q = 1$. As the number of components in the bilinear term increases, the results from B-AMMI deteriorate compared to AMMI and AMBARTI. In addition, it is possible to note that there is no clear effect of σ_g or σ_e on the RMSEs. However, even with the AMMI model presenting the best results, AMBARTI demonstrates highly competitive performance, with RMSE values around 17% higher than that of the AMMI model.

Fig 3: Out-of-sample RMSE for $\hat{y}$ based on the results of AMMI, B-AMMI, and AMBARTI for data simulated from the AMMI model with $I = J = 10$. The different panels contain 10 Monte Carlo repetitions and represent different combinations of the simulated parameters for the creation of the data set. Unsurprisingly, AMMI performs very well here, with AMBARTI having RMSE values around 17% higher.

Figure 4 shows the results of the second simulation scenario, where the data were simulated from the AMBARTI equation. Again, different combinations of parameters were used in the simulation of the training and out-of-sample sets. The upper panels show results for $I = 10$ genotypes and $J = 10$ environments; the lower ones for $I = 25$ and $J = 25$. Furthermore, three AMMI and three B-AMMI models were fitted considering Q from 1 to 3. In this

case, the AMMI model, even with high values of Q, performs very poorly with RMSE values 3 times higher on average than that of AMBARTI for $I = 10$. For $I = 25$, the AMMI model with $Q = 3$ is competitive with AMBARTI, with the latter being slightly better. In addition, we can see that AMBARTI presents better results when compared with B-AMMI, regardless of the value of Q. In this comparison, it is worth mentioning that more complex possibilities of interactions may be obtained when simulating from AMBARTI compared to AMMI.

Fig 4: Out-of-sample RMSE for $\hat{y}$ based on the results of AMMI (with varying Q), B-AMMI, and AMBARTI for data simulated from the AMBARTI model with $I = J = 10$ and $I = J = 25$. The boxplots contain 10 Monte Carlo repetitions. The AMMI RMSE values are on average 3 times higher than that of AMBARTI for $I = 10$.

The next important comparison to be made between AMBARTI and AMMI is related to the interaction term (i.e., the bilinear term for AMMI and the BART component for AM-BARTI). Such tests are shown in Figures 5 and 6, where we show the RMSE performance only for the interaction component.

Figure 5 presents the RMSEs associated solely with the interaction terms from AMMI, B-AMMI, and AMBARTI when the data are simulated from AMMI (which has a bilinear structure for the interactions). The results are presented considering 10 genotypes and 10 environments with different combinations of genotypic and environmental variances. The performance of AMMI is optimal compared to AMBARTI, though the difference between the two is lessened with more complex AMMI model structures (i.e., $Q = 3$). In Figure 6, the values of RMSE are presented for data sets simulated from AMBARTI. In the margins of the figure, the parameters used in the simulations can be found. The RMSE values show that AMMI performs worse than AMBARTI in all scenarios, and in the same cases AMMI RMSEs are three times higher on average than those of AMBARTI for the scenario with $I = 10$ genotypes.

In summary, the information presented in Figures 5 and 6 shows that the AMMI model fails for the complex interactions that can be obtained in the AMBARTI simulated data sets. From a quantitative genetics/biological perspective, there is no reason for the structure of

Fig 5: Out-of-sample RMSE related to the interaction term of AMMI models for data simulated from AMMI. The different panels show the different parameter values used in the simulation. The performance of AMMI here is optimal, with AMBARTI performing slightly worse than AMMI when $Q = 3$.

interactions between genotypes and environments to be modelled strictly by a bilinear structure, as more complex structures can be assumed to be present in nature. Thus, AMBARTI may be a more suitable model to estimate the interaction structure in real-world applications.

4. Case study: Irish VCU InnoVar wheat data.   In addition to the simulation study, real data sets were used to evaluate the performance of AMBARTI. We compare our new approach not only to AMMI and B-AMMI, but also to more sophisticated interaction detection models including smoothing splines ANOVA models (SS-ANOVA Gu, 2014) and Bayesian multivariate adaptive regression splines (B-MARS Denison, Mallick and Smith, 1998). To run SS-ANOVA and B-MARS, we used the R packages `gss` (Francom and Sansó, 2020) and `BASS` (Gu, 2014), respectively. A set of value of cultivation and usage (VCU) experiments conducted in Ireland between the years of 2010 and 2019 were considered, and such experiments evaluated the performance of genotypes of wheat *Triticum aestivum L.* across the country for regulatory purposes (i.e., registration of new varieties). Here, our phenotypic response variable is the production of wheat in tonnes per hectare. The design of experiments used was that of a randomised complete block design with 4 replicates. VCUs alongside distinctness, uniformity and stability (DUS) are the most important kind of regulatory multi environmental trials conducted around the world.

The data were kindly provided by the Irish Department of Agriculture, Food, and Marine. Both genotypes and environments were anonymised. These historical Irish VCUs form part of the Horizon2020 EU InnoVar project database (`www.h2020innovar.eu`). The project aims to build and improve technical solutions for cultivar recommendation based on genomic and phenomic parameters. The models were fitted for all years available (summarised in Table 1), but for brevity we show detailed plots only for the year 2015.

Fig 6: Out-of-sample RMSE related to interaction term of AMBARTI and AMMI models for data simulated from AMBARTI. The different panels show the different parameter values used for the simulation. It appears that the AMMI structure, even with $Q = 3$ cannot capture the interaction behaviour present in the AMBARTI model for $I = 10$.

We compare the models by evaluating the estimated values of the genotype and environment effects, and the predictions of interaction behaviour evaluated as: $(\hat{ge})_{ij} = y_{ij} - \hat{g}_i - \hat{e}_j$. As the Irish data from the InnoVar project correspond to a block design with 4 replicates, to fit all models we randomly selected two replicates and then averaged the response variable across them. This is a common practice in the analysis of GxE experiments with AMMI and B-AMMI models (Josse et al., 2014; Crossa et al., 2011) as these models cannot deal with replicates in an experiment. However, this pre-processing is not needed for AMBARTI. To validate the models, we use the remaining two replicates to calculate the RMSE for $\hat{y}$.

The estimates of the genotype effects g_i and environment effects e_j are shown in Figure 7 and 8, respectively. The results for AMBARTI show the samples obtained from the posterior distributions, while for AMMI, as it is a frequentist method, we adopted the approach of Goodman and Haberman (1990) where the results correspond to samples from the estimator distributions of g_i and e_j. The rationale of using samples from the estimator distribution is to be able to compare AMBARTI and AMMI results not only in terms of point estimates but also in terms of the uncertainty associated to the point estimates. Although we have tested different number of components Q for AMMI, we remark the parameter estimates g_i and e_j do not change regardless of Q (i.e., only the bilinear term depends on Q; see Equation (2)).

In Figure 7, we can see that the credible intervals associated with the main effects of genotypes for AMBARTI are narrower than the confidence intervals from the AMMI model. This is somewhat expected and occurs as BART is able to capture the interactions between g_i and e_j along the MCMC process, which in turn decreases the residual variance σ^2. Hence, as the full conditionals of g_i's and e_j's depend on σ^2, their estimates become less uncertain as σ^2 gets smaller; see Equations (10) and (11) in Appendix A. For the sake of visualisation, we decided to omit the B-AMMI results as they presented too many values that exceeded -10 and 10.

Fig 7: For AMBARTI, the boxplots represent the posterior distribution of genotype effects for the Irish VCU Innovar data set for 2015. As the parameters estimates from the AMMI model are obtained under the frequentist paradigm, their boxplots summarise samples from the estimator distribution of the genotype effects as presented in Goodman and Haberman (1990).

A more complete comparison across all years is shown in Table 1. In this table, we calculated the predicted values $\hat{y}$ on the out-of-sample data. We can see that RMSEs obtained with AMBARTI are smaller than the ones returned by the AMMI model for all years, thus highlighting that the AMBARTI model can more accurately estimate the marginal effects along with interaction component. Further, we can note that the B-AMMI presents the worst results among all methods considered. One could expect the results from AMMI and B-AMMI would be similar, but the Bayesian version is different in spirit to its frequentist counterpart, as it does not estimate the main effects and bilinear term in a two-stage approach. Instead, B-AMMI obtains the parameters through a one-stage procedure and assumes priors that do not take into account the orthormality constraints, which are fundamental for the identifiability of the frequentist AMMI model. In the B-AMMI, the orthormality constraints are applied via a post-processing on the parameter estimates after the model is fitted. However, we observed the B-AMMI model with and without the post-processing may have very different performance.

Although the results for SS-ANOVA and B-MARS are competitive with AMBARTI, they work differently and do not return results in the format needed to fit the purpose of the analysis. For instance, SS-ANOVA returned interactions that do not make sense (e.g., interactions between environments or interactions between genotypes). In contrast, B-MARS is similar to BART in the sense that it does not require any specification but with the advantage that it decomposes the impact of individual and interaction effects on the response.

Regarding the computational time, AMBARTI took about 6 minutes on average to run, considering 50 trees, 1000 iterations as burn-in and 1000 iterations as post burn-in. This time was registered in a MacBook Pro 2.3 GHz Dual-Core Intel Core i5 with 8GB memory. AMMI took just seconds. This difference could be reduced by optimising the AMBARTI implementation using routines in C++ similar to those for BART implementations in R packages BART

Fig 8: For AMBARTI, the boxplots represent the posterior distribution of environment effects for the Irish VCU Innovar data set for 2015. As the parameters estimates from the AMMI model are obtained under the frequentist paradigm, their boxplots summarise samples from the estimator distribution of the environment effects as presented in Goodman and Haberman (1990).

Year	AMBARTI	AMMI	B-AMMI	SS-ANOVA	B-MARS
2010	**0.81**	0.83	2.21	0.82	0.87
2011	**0.62**	0.63	2.66	0.65	0.69
2012	**0.45**	0.47	2.53	0.60	0.65
2013	0.57	0.57	1.48	**0.55**	0.56
2014	0.55	0.54	1.93	**0.53**	0.54
2015	**0.46**	0.46	1.75	0.52	0.55
2016	**0.44**	0.45	1.98	0.50	0.50
2017	**0.51**	0.53	2.24	0.60	0.62
2018	0.74	0.73	1.09	**0.70**	0.70
2019	**0.56**	0.58	1.26	0.72	0.70

TABLE 1

RMSE for $\hat{y}$ on out-of-sample data considering all years in the historical Irish VCU Innovar data. The values of RMSE obtained with AMBARTI are smaller than the ones obtained via AMMI and Bayesian AMMI models (both with $Q = 3$) for all years considered. The values in bold correspond to the smallest RMSE within each year.

(McCulloch et al., 2020) and `dbarts` (Dorie, 2020). However, we believe AMBARTI's superior performance and posterior estimation of uncertainties outweighs the longer computational time.

4.1. *New visualisations for AMBARTI main effects and interactions.* One of the key outputs of the standard AMMI model is the biplot (Gabriel, 1971), which assists in the determination of important GxE interactions and may be used for cultivar recommendation. However, these plots display only the interaction measure, thus missing the key marginal effects that may also come into play. For example, a certain genotype and environment may have a strong positive interaction, but if the genotype is consistently poor in all environments

this may not be clear in the biplot. Instead, we introduce new types of plots that give this full consideration.

Our first new plot is based on a heat map adapted to display both the GxE interactions (along with the marginal effects) and the predicted yields from the AMBARTI model. In Figure 9, we display the GxE interactions in the centre of the plot and the marginal effects for both environment and genotype as separate bars in the margins. The ordering applied to Figure 9 is in terms of the marginal effects for both environment and genotype, and displays low values in red to high values in blue. As both the GxE interactions and marginal effects are on the same scale and are centred around zero, we display them using only one legend and use a divergent colour palette. A diverging colour palette uses two diverging hues to represent the extremes and highlights the midpoint with a light colour. This allows for quick identification of the GxE interactions and to observe which of the environments or genotypes are the most or least optimal.

Fig 9: GxE interactions and main effects for the AMBARTI model sorted by the main effects for the Irish VCU InnoVar data in 2015. We can clearly see that environments 1, 4, 6, and 5 provide superior yields for many of the genotypes studied. Furthermore, environment 1, for example, seems to interact particularly strongly in a negative way with genotype 2. The grand mean μ is not included in this plot for ease of identification of marginal and interacting effects.

In Figure 10, we show the ordered heat map of the predicted yields (as opposed to their component parts shown in Figure 9) for each combination of environment and genotype for the AMBARTI model. In this case, we use the same ordering as that in Figure 9 with high values being generally displayed in the top left, moving to low values at the bottom right, with the units for the plot being the same as that of the phenotype (i.e., yield/production of grains in tonnes per hectare). For this plot, we use the same diverging colour palette as in Figure 9 as the scale is centred around the mean, and when combined with the ordering, this gives a clear identification as to which environment and genotype produce high or low yields. Additionally, the AMBARTI methodology makes it possible to use tree methods to identify which combinations of genotypes and environments are similar considering a given phenotypic characteristic, e.g., yield. For example, the combination of genotype 1 and environment 1 and genotype 17 and environment 1 are similar in terms of predicted yield, as

shown by the colours in Figure 10. Similarly, genotype 2 has similar yields for environments 4 and 5. Thus, we can consider these environments similar and group them in a set named mega-environment 4-5 considering genotype 2. Such a concept is crucial for cultivar recommendations. To visualise the uncertainty associated with Figures 9 and 10, we provide plots which show the median, 5%, and 95% quantiles for the predicted response and GxE interactions and main effects in Appendix C.

Fig 10: Predicted yields from the AMBARTI model for the Irish VCU InnoVar data in 2015. Values are sorted by the main effects. We can see, for example, a high value for the predicted yield for environment 1 with genotype 5 and a low value between environment 2 and genotype 11.

In Figure 11, we show a bipartite plot of the information displayed in Figure 9, but showing only the extremes of the high and low values. In this case, we display just the top 2% and the lowest 2% of the interactions. We employ the same diverging colour palette as Figure 9 except in this case the colour of the nodes represent the marginal effects and the size of each node represents the absolute value of the marginal effects. Similarly, the colour of the connecting edges represent the interaction values and the width of each edge represents the absolute interaction value. That is, larger magnitudes of the marginal effects will result in larger nodes (and vice-versa), and larger magnitudes of the interactions will result in thicker edges (and vice-versa). The aim of this plot is to allow the reader to easily and quickly identify which of the environments are the most and least optimal for each genotype and to also identify where there are clear interactions. Quantile versions of Figure 11 could also be plotted to assess uncertainty.

The visualisation perspective proposed here helps construct easily interpretable agronomic recommendations. Figure 9 can help users with no background in statistics identify that the best genotypes considering yield are the ones in the top left corner: g_{17}, g_{16}, g_5, g_2. These genotypes will have a tendency to have a better acceptance by farmers, considering solely the yield in tonnes per hectare assuming higher yields are economically preferred. Figure 9 shows us that environments e_1, e_4, e_6, e_5 are related to higher marginal effects and should be considered preferential to crop the list of wheat genotypes evaluated.

Fig 11: Bipartite network plot showing the top (in blue) and bottom (in red) 2% GxE interactions and main effects from the AMBARTI model for the Irish VCU InnoVar data in 2015.. We can see that environment 3 has strong positive and negative interactions with genotypes 12 and 13, respectively.

Figures 9 and 11 are also useful to establish combinations of genotypes and environments that should be avoided when the interaction is negative, indicating that a given genotype does not perform well in a given environment. This negative interaction increases the risk of low yield and consequent economic impacts. Combinations to be avoided exist even for environments and genotypes with high marginal effects. For instance, the combination of g_2, e_1 should be avoided even though g_2 and e_1 have high marginal effects. This is an important information for regulators who may be responsible for a variety's commercialisation approval or agents that promote credit or insurance for farmers given to risks that the negative interaction implies. Farmers who produce a genotype not indicated for their environment can end having a worse score or risk. On the other hand, Figure 10 is also useful to spot the combinations of genotypes and environments that should be encouraged once the signal of the interactions is positive.

In adaptability breeding, the breeder seeks to find the best genotype for a specific environment or a small set of environments. In broad target strategies, the aim is to find genotypes that perform well across several environments. For example, in Figure 9, g_5 has high marginal effect and performs well (and interacts positively) with environments e_1, e_4, e_6, e_9. Similarly g_{16}, the second best genotype considering marginal effects, performs well in environments e_4, e_5, e_9, e_7. Genotypes which present better performance across several environments are classified as high stability genotypes. They tend to be preferred by breeders because they allow optimisation of processes in the chain of seed production.

5. Discussion. We have introduced a new model named additive main effects Bayesian additive regression trees interaction (AMBARTI). AMBARTI is a fully Bayesian semi-parametric machine learning approach that estimates main effects of genotypes and environments and interactions with an adapted regression tree-like structure. This approach to interactions allows the treatment of more complex structures than the ones considered by traditional models.

Given the fact that GxE interactions are the result of a tangled myriad of genetics, proteomics, biochemical, environmental and additional factors, the flexibility of AMBARTI in dealing with more complex interactions can be seen as an important improvement in the understanding of the complexities associated to GxE phenomenon. In practice, the choice between a low-rank model (AMMI) and a model that is able to deal with a sparse interaction structure (AMBARTI) is a modelling choice. However, in the real data examples upon we have tested the models, AMBARTI performed slightly or much better than AMMI, which suggests that a bilinear term is perhaps an oversimplification to study the relations that arise from GxE interactions. We believe that AMBARTI is a useful candidate to expand the understanding of experimental data in quantitative genetics.

The main novelty in AMBARTI comes from its semi-parametric structure which enables the uncertainty to be shared between the main effects and the interaction trees. More specifically, we design the trees so that they can only incorporate the interaction terms by forcing each branch of a tree to split on both a genotype and an environment. We have shown in simulation experiments that this yields similar estimates to traditional models for the marginal effects and superior estimates for the interaction terms, which are no longer restricted to be linear in a restricted dimensional space. This removes the need for, e.g., the arbitrary selection of the Q parameter in a standard AMMI formulation.

A second novelty is that we have introduced new displays that simultaneously allow for interpretation of the marginal and joint effects. We have created both a heatmap and a bipartite network-style plot of the results, which are not limited for use with an AMBARTI model and could be applied to any suitable model (for example, AMMI, B-AMMI, etc.) and are a useful tool for deciphering complex model structures. From these plots, we hope to enable those using the output of AMBARTI models to make more informative decisions about which genotype and environments are most compatible.

We believe that there are many possible extensions of the AMBARTI approach. Other more advanced methods, such as PARAFAC (Basford, Kroonenberg and DeLacy, 1991; Harshman and Lundy, 1994), are available for higher dimensional interactions (such as with time). These are different versions of tensor regression (Guhaniyogi, Qamar and Dunson, 2017) and, in theory, there is no reason why the AMBARTI approach cannot be used for higher dimensional tensor-type interactions, though this is not currently possible in our code. Similar enhancements for multivariate outputs and time-series like-structure seem promising, and we hope to explore these in future work.

Acknowledgments. Danilo and Estevão are joint first authors. We are very grateful to John Joe Byrne at the Department of Agriculture for providing us with the data set for our case study. Danilo Sarti and Antônia A. L. dos Santos received funding for their work from the European Union's Horizon 2020 research and innovation programme under grant agreement No 818144. Alan Inglis and Estevão Prado's work was supported by the Science Foundation Ireland Career Development Award grant number 17/CDA/4695 and SFI research centre 12/RC/2289_P2. Andrew Parnell's work was supported by: the Science Foundation Ireland Career Development Award (17/CDA/4695); an investigator award (16/IA/4520); the Marine Research Programme funded by the Irish Government, co-financed by the European Regional Development Fund (Grant-Aid Agreement No. PBA/CC/18/01); European Union's Horizon 2020 research and innovation programme under grant agreement No 818144; SFI Centre for Research Training 18CRT/6049, and SFI Research Centre awards 16/RC/3872 and 12/RC/2289_P2. For the purpose of Open Access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission'.

REFERENCES

ALBERT, J. H. and CHIB, S. (1993). Bayesian analysis of binary and polychotomous response data. *Journal of the American statistical Association* **88** 669–679.

ALLARD, R. W. and BRADSHAW, A. D. (1992). Implications of genotype environmental interactions in applied plant breeding. *Crop Science* **4** 503-508.

ANBESSA, Y., JUSKIW, P., GOOD, A., NYACHIRO, J. and HELM, J. (2009). Genetic variability in nitrogen use efficiency of spring barley. *Crop Science* **49** 1259–1269.

BADU-APRAKU, B., OYEKUNLE, M., OBENG-ANTWI, K., OSUMAN, A., ADO, S., COULIBAY, N., YAL-LOU, C., ABDULAI, M., BOAKYEWAA, G. and DIDJEIRA, A. (2012). Performance of extra-early maize cultivars based on GGE biplot and AMMI analysis. *The Journal of Agricultural Science* **150** 473.

BASAK, P., LINERO, A., SINHA, D. and LIPSITZ, S. (2021). Semiparametric analysis of clustered interval-censored survival data using soft Bayesian additive regression trees (SBART). *Biometrics*.

BASFORD, K., KROONENBERG, P. and DELACY, I. (1991). Three-way methods for multiattribute genotype× environment data: an illustrated partial survey. *Field Crops Research* **27** 131–157.

BRANCOURT-HULMEL, M. and LECOMTE, C. (2003). Effect of environmental variates on genotype× environment interaction of winter wheat: A comparison of biadditive factorial regression to AMMI. *Crop Science* **43** 608–617.

CHIPMAN, H. A., GEORGE, E. I., MCCULLOCH, R. E. et al. (2010). BART: Bayesian additive regression trees. *The Annals of Applied Statistics* **4** 266–298.

CROSSA, J., PEREZ-ELIZALDE, S., JARQUIN, D., COTES, J. M., VIELE, K., LIU, G. and CORNELIUS, P. L. (2011). Bayesian estimation of the additive main effects and multiplicative interaction model. *Crop Science* **51** 1458–1469.

DE MENDIBURU, F. (2019). Package 'agricolae'. *R Package, Version* 1–2.

DENISON, D. G., MALLICK, B. K. and SMITH, A. F. (1998). Bayesian mars. *Statistics and Computing* **8** 337–346.

DIAS, C. (2005). Métodos para escolha de componentes em modelo de efeito principal aditivo e interação multiplicativa (AMMI). 2005. 73p, PhD thesis, Tese (Livre Docência)–Escola Superior de Agricultura Luiz de Queiroz, Piracicaba.

DIAS, C. T. D. S. and KRZANOWSKI, W. J. (2006). Choosing components in the additive main effect and multiplicative interaction (AMMI) models. *Scientia Agricola* **63** 169–175.

DORIE, V. (2020). dbarts: Discrete Bayesian Additive Regression Trees Sampler. R package version 0.9-19.

FALCONER, D. and MACKAY, T. (1996). Introduction to quantitative genetics. 1996. *Harlow, Essex, UK: Longmans Green* **3**.

FARSHADFAR, E. and SUTKA, J. (2003). Locating QTLs controlling adaptation in wheat using AMMI model. *Cereal Research Communications* **31** 249–256.

FRANCOM, D. and SANSÓ, B. (2020). BASS: An R Package for Fitting and Performing Sensitivity Analysis of Bayesian Adaptive Spline Surfaces. *Journal of Statistical Software* **94** 1–36.

GABRIEL, K. R. (1971). The biplot graphic display of matrices with application to principal component analysis. *Biometrika* **58** 453–467.

GAMERMAN, D. and LOPES, H. F. (2006). *Markov chain Monte Carlo: stochastic simulation for Bayesian inference*. CRC Press.

GAUCH JR, H. G. (2013). A simple protocol for AMMI analysis of yield trials. *Crop Science* **53** 1860–1869.

GOLLOB, H. F. (1968). A statistical model which combines features of factor analytic and analysis of variance techniques. *Psychometrika* **33** 73–115.

GOOD, I. J. (1969). Some applications of the singular decomposition of a matrix. *Technometrics* **11** 823–831.

GOODMAN, L. A. and HABERMAN, S. J. (1990). The analysis of nonadditivity in two-way analysis of variance. *Journal of the American Statistical Association* **85** 139–145.

GU, C. (2014). Smoothing Spline ANOVA Models: R Package gss. *Journal of Statistical Software* **58** 1–25.

GUHANIYOGI, R., QAMAR, S. and DUNSON, D. B. (2017). Bayesian tensor regression. *The Journal of Machine Learning Research* **18** 2733–2763.

HARSHMAN, R. A. and LUNDY, M. E. (1994). PARAFAC: Parallel factor analysis. *Computational Statistics & Data Analysis* **18** 39–72.

HERNÁNDEZ, B., PENNINGTON, S. R., PARNELL, A. C. et al. (2015). Bayesian methods for proteomic biomarker development. *EuPA Open Proteomics* **9** 54–64.

HERNÁNDEZ, B., RAFTERY, A. E., PENNINGTON, S. R., PARNELL, A. C. et al. (2018). Bayesian additive regression trees using Bayesian model averaging. *Statistics and computing* **28** 869–890.

ISIK, F., HOLLAND, J. and MALTECCA, C. (2017). Multi environmental trials. In *Genetic data analysis for plant and animal breeding* 227–262. Springer.

JEONG, S. and ROČKOVÁ, V. (2020). The art of BART: On flexibility of Bayesian forests. *arXiv preprint arXiv:2008.06620*.

JOSSE, J., VAN EEUWIJK, F., PIEPHO, H.-P. and DENIS, J.-B. (2014). Another look at Bayesian analysis of AMMI models for genotype-environment data. *Journal of Agricultural, Biological, and Environmental Statistics* **19** 240–257.

KAPELNER, A. and BLEICH, J. (2016). bartMachine: Machine Learning with Bayesian Additive Regression Trees. *Journal of Statistical Software, Articles* **70** 1–40.

KINDO, B. P., WANG, H., PEŇA, E. A. et al. (2016). Multinomial probit Bayesian additive regression trees. *Stat* **5** 119–131.

LAL, R., CHANOTIYA, C., DHAWAN, S., GUPTA, P., MISHRA, A., SRIVASTAVA, S., SHUKLA, S. and MAURYA, R. (2020). Estimation of intra-specific genetic variability and half-sib family selection using AMMI (Additive Main Effects and Multiplicative Interactions) model in menthol mint (Mentha arvensis L.). *J. Med. Arom. Plant Sci* **42** 102–113.

LINERO, A. R. and YANG, Y. (2018a). Bayesian regression tree ensembles that adapt to smoothness and sparsity. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **80** 1087–1110.

LINERO, A. R. and YANG, Y. (2018b). Bayesian regression tree ensembles that adapt to smoothness and sparsity. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **80** 1087–1110.

LINERO, A. R., BASAK, P., LI, Y. and SINHA, D. (2021). Bayesian Survival Tree Ensembles with Submodel Shrinkage. *Bayesian Analysis* **1** 1–24.

LIU, Y., TRASKIN, M., LORCH, S. A., GEORGE, E. I. and SMALL, D. (2015). Ensemble of trees approaches to risk adjustment for evaluating a hospital's performance. *Health care management science* **18** 58–66.

LOVE, S. L., SALAIZ, T., SHAFII, B., PRICE, W. J., MOSLEY, A. R. and THORNTON, R. E. (2004). Stability of expression and concentration of ascorbic acid in North American potato germplasm. *HortScience* **39** 156–160.

MAHALINGAM, L., MAHENDRAN, S., BABU, R. C. and ATLIN, G. (2006). AMMI analysis for stability of grain yield in rice (Oryza sativa L.). *International Journal of Botany*.

MANDEL, J. (1971). A new analysis of variance model for non-additive data. *Technometrics* **13** 1–18.

MCCULLOCH, R., SPARAPANI, R., SPANBAUER, C., GRAMACY, R. and PRATOLA, M. (2020). BART: Bayesian Additive Regression Trees R package version 2.8.

MITROVIAĂ, B., TRESKI, S., STOJAKKOVĂ, M., IVANOVIĂ, M., BEKAVAC, G. et al. (2012). Evaluation of experımental maize hybrids tested in multi-location trials using AMMI and GGE biplot analyses. *Turkish Journal of Field Crops* **17** 35–40.

NACHIT, M., NACHIT, G., KETATA, H., GAUCH, H. and ZOBEL, R. (1992). Use of AMMI and linear regression models to analyze genotype-environment interaction in durum wheat. *Theoretical and Applied genetics* **83** 597–601.

ONOFRI, A. and CIRICIOFOLO, E. (2007). Using R to perform the AMMI analysis on agriculture variety trials. *R News* **7** 14–19.

PLUMMER, M. (2015). Cuts in Bayesian graphical models. *Statistics and Computing* **25** 37–43.

PRADO, E. B., MORAL, R. A., PARNELL, A. C. et al. (2021). Bayesian additive regression trees with model trees. *Statistics and Computing* **31** 1–13.

RAD, M. N., KADIR, M. A., RAFII, M., JAAFAR, H. Z., NAGHAVI, M. and AHMADI, F. (2013). Genotype environment interaction by AMMI and GGE biplot analysis in three consecutive generations of wheat (Triticum aestivum) under normal and drought stress conditions. *Australian Journal of Crop Science* **7** 956.

ROBERT, C. and CASELLA, G. (2013). *Monte Carlo statistical methods*. Springer Science & Business Media.

RODRIGUES, P. C., MONTEIRO, A. and LOURENÇO, V. M. (2016). A robust AMMI model for the analysis of genotype-by-environment data. *Bioinformatics* **32** 58–66.

ROMAGOSA, I., ULLRICH, S. E., HAN, F. and HAYES, P. M. (1996). Use of the additive main effects and multiplicative interaction model in QTL mapping for adaptation in barley. *Theoretical and Applied Genetics* **93** 30–37.

ROČKOVÁ, V. and SAHA, E. (2019). On theory for BART. In *The 22nd International Conference on Artificial Intelligence and Statistics* 2839–2848. PMLR.

ROČKOVÁ, V. and VAN DER PAS, S. (2020). Posterior concentration for Bayesian regression trees and forests. *Annals of Statistics* **48** 2108–2131.

SARTI, D. A. (2013). Uncertainty management through decision analysis: applications to production optimization and uncertain demands, Master's thesis, University of São Paulo.

SARTI, D. A. (2019). The statistical paradigm: probabilistic and multivariate analysis applied through computational simulation in the interaction between genotype x environment, PhD thesis, Universidade de São Paulo.

SATO, K. and TAKEDA, K. (1993). Pathogenic variation of Pyrenophora teres isolates collected from Japanese and Canadian spring barley. *Report by the Institute of Resource Biological Sciences, Okayama University* **1** 147–158.

SHAFII, B. and PRICE, W. J. (1998). Analysis of genotype-by-environment interaction using the additive main effects and multiplicative interaction model and stability estimates. *Journal of Agricultural, Biological, and Environmental Statistics* 335–345.

SILVEIRA, L. C. I. D., KIST, V., PAULA, T. O. M. D., BARBOSA, M. H. P., PETERNELLI, L. A. and DAROS, E. (2013). AMMI analysis to evaluate the adaptability and phenotypic stability of sugarcane genotypes. *Scientia Agricola* **70** 27–32.

SPARAPANI, R. A., LOGAN, B. R., MCCULLOCH, R. E. and LAUD, P. W. (2016). Nonparametric survival analysis using Bayesian additive regression trees (BART). *Statistics in medicine* **35** 2741–2753.

TAN, Y. V. and ROY, J. (2019). Bayesian additive regression trees and the General BART model. *Statistics in medicine* **38** 5048–5069.

R CORE TEAM (2020). R: A Language and Environment for Statistical Computing R Foundation for Statistical Computing, Vienna, Austria.

TYAGI, B., SINGH, M., SINGH, G., KUMAR, R., VERMA, A., SHARMA, I. et al. (2016). Genetic variability and AMMI bi-plot analysis in bread wheat based on multi-location trials conducted under drought conditions across agro-climatic zones of India. *Triticeae Genomics and Genetics* **7**.

WRIGHT, M. N. and KÖNIG, I. R. (2019). Splitting on categorical predictors in random forests. *PeerJ* **7** e6339.

ZELDOW, B., RE III, V. L., ROY, J. et al. (2019). A semiparametric modeling approach using Bayesian additive regression trees with an application to evaluate heterogeneous treatment effects. *The Annals of Applied Statistics* **13** 1989.

ZHANG, J. L. and HÄRDLE, W. K. (2010). The Bayesian additive classification tree applied to credit risk modelling. *Computational Statistics & Data Analysis* **54** 1197–1205.

APPENDIX A: AMBARTI IMPLEMENTATION

In this Section, we detail the AMBARTI model. Firstly, the likelihood function associated to y_{ij} is

$$y_{ij}|\mathbf{x}_{ij}, \Theta \sim \mathrm{N}\left(\mu + g_i + e_j + \sum_{t=1}^{T} h(\mathbf{x}_{ij}, \mathcal{M}_t, \mathcal{T}_t), \sigma^2\right),$$

where y_{ij} denotes the response for genotype i and environment j, $\Theta = (\mu, g_i, e_j, \mathcal{M}_t, \mathcal{T}_t, \sigma^2)$, μ is the grand mean, $\mathbf{x}_{ij}$ is the row of the design matrix $\mathbf{X}$ associated to observations with genotype i and environment j, and $h(\cdot) = \mu_{t\ell}$ is a function that assigns the predicted values $\mu_{t\ell} \in \mathcal{M}_t$ to observations that belong to $\mathcal{P}_{t\ell}$, with $\mathcal{P}_{t\ell}$ denoting the set of rules that define the node ℓ of the tree t. In order to obtain the posterior distributions needed for the model, we assume the following prior distributions:

$$\mu \sim \mathrm{N}(m, \sigma_m^2),$$
$$\mu_{t\ell}|\mathcal{T}_t \sim \mathrm{N}(0, \sigma_\mu^2),$$
$$g_i|\mathcal{T}_t \sim \mathrm{N}(\mu_g, \sigma_g^2),$$
$$e_j|\mathcal{T}_t \sim \mathrm{N}(\mu_e, \sigma_e^2),$$
$$\sigma_g^2 \sim \mathrm{IG}(a_g, b_g),$$
$$\sigma_e^2 \sim \mathrm{IG}(a_e, b_e),$$
$$\sigma^2 \sim \mathrm{IG}(a, b).$$

The prior distribution on the tree structure depends on the number of terminal and internal nodes, and is given by

$$p(\mathcal{T}_t) = \prod_{\ell \in \mathcal{A}_I} \left[\alpha(1 + d_{t\ell})^{-\beta}\right] \times \prod_{\ell \in \mathcal{A}_T} \left[1 - \alpha(1 + d_{t\ell})^{-\beta}\right],$$

where $\mathcal{A}_I$ and $\mathcal{A}_T$ denote the sets of indices of the internal and terminal nodes, respectively, and $d_{t\ell}$ represents the depth of the node ℓ of the tree t. Furthermore, let $R_t = \mathbf{y} - \left(\mu + \mathbf{g} + \mathbf{e} + \sum_{k \neq t}^{T} h(\mathbf{X}; \mathcal{T}_k, \mathcal{M}_k)\right)$ denote the vector of the partial residuals, where $\mathbf{g}$ and $\mathbf{e}$ are vectors containing the main effects g_i and e_j for all i and j. Below, we present the full conditional of μ.

$$p(\mu|-) \propto p(\mathbf{y}|g_i, e_j, \mathbf{x}_{ij}, \mathcal{M}_t, \mathcal{T}_t, \sigma^2)p(\mu),$$

$$\propto \exp\left(-\frac{1}{2\sigma_{m*}^2}(\mu - \mu^*)^2\right),$$

which is a

$$N\left(\frac{\sum_i \sum_j \left[y_{ij} - \hat{y}_{ij}^*\right]/\sigma^2 + m/\sigma_m^2}{n/\sigma^2 + 1/\sigma_m^2}, \frac{1}{n/\sigma^2 + 1/\sigma_m^2}\right),$$

where $\hat{y}_{ij}^* = g_i + e_j + \sum_{t=1}^{T} h(\mathbf{x}_{ij}; \mathcal{T}_t, \mathcal{M}_t)$. Hence, the full conditional of g_i is given by

$$p(g_i|-) \propto p(\mathbf{y}|g_i, e_j, \mathbf{x}_{ij}, \mathcal{M}_t, \mathcal{T}_t, \sigma^2)p(g_i),$$

$$\propto \exp\left(-\frac{1}{2\sigma_{g*}^2}(g_i - g_i^*)^2\right),$$

which is a

(10) $$N\left(\frac{\sum_j [y_{ij} - \hat{\mu} - \hat{e}_j - \hat{\mu}_{ij}]/\sigma^2}{n_{g_i}/\sigma^2 + 1/\sigma_g^2}, \frac{1}{n_{g_i}/\sigma^2 + 1/\sigma_g^2}\right),$$

where $\hat{\mu}_{ij} = \sum_{t=1}^{T} h(\mathbf{x}_{ij}, \mathcal{T}_t, \mathcal{M}_t)$ and n_{g_i} is the number of observations that belong to g_i; similarly to n_{e_j}. Analogously, the full conditional of e_j can be written as

$$p(e_j|-) \propto p(\mathbf{y}|g_i, e_j, \mathbf{x}_{ij}, \mathcal{M}_t, \mathcal{T}_t, \sigma^2)p(e_j),$$

$$\propto \exp\left(-\frac{1}{2\sigma_{e*}^2}\left(e_j - e_j^*\right)^2\right),$$

which is a

(11) $$N\left(\frac{\sum_i [y_{ij} - \hat{\mu} - \hat{g}_i - \hat{\mu}_{ij}]/\sigma^2}{n_{e_j}/\sigma^2 + 1/\sigma_e^2}, \frac{1}{n_{e_j}/\sigma^2 + 1/\sigma_e^2}\right).$$

The full conditional of σ_g^2 is given by

$$p(\sigma_g^2|-) \propto p(\mathbf{g}|\sigma_g^2)p(\sigma_g^2),$$

which is an

$$IG\left(\frac{I}{2} + a_g, \frac{\sum_{i=1}^{I} g_i^2}{2} + b_g\right).$$

The full conditional of σ_e^2 is written as

$$p(\sigma_e^2|-) \propto p(\mathbf{e}|\sigma_e^2)p(\sigma_e^2),$$

which is an

$$IG\left(\frac{J}{2} + a_e, \frac{\sum_{j=1}^{J} e_j^2}{2} + b_e\right).$$

In addition, we present the full conditional of the trees. This distribution is utilised to compare the previous tree to the current one, as in BART the splitting rules are created by randomly selecting a covariate and a split point. Below, we present the full conditional of $\mathcal{T}_t$ as

$$p(\mathcal{T}_t|R_t, \sigma^2) \propto p(\mathcal{T}_t) \int p(R_t|\mathcal{M}_t, \mathcal{T}_t, \sigma^2) p(\mathcal{M}_t|\mathcal{T}_t) d\mathcal{M}_t,$$

$$\propto p(\mathcal{T}_t) p(R_t|\mathcal{T}_t, \sigma^2),$$

$$\propto p(\mathcal{T}_t) \prod_{\ell=1}^{b_t} \left[\left(\frac{\sigma^2}{\sigma_\mu^2 n_{t\ell} + \sigma^2} \right)^{1/2} \exp \left(\frac{\sigma_\mu^2 \left[n_{t\ell} \bar{R}_\ell \right]^2}{2\sigma^2 (\sigma_\mu^2 n_{t\ell} + \sigma^2)} \right) \right],$$

where $\bar{R}_\ell = \sum_{(i,j) \in \mathcal{P}_{t\ell}} (r_{ij}^{(t)} - \hat{\mu} - \hat{g}_i - \hat{e}_j)/n_{t\ell}$, $r_{ij}^{(t)} \in R_t$ and $n_{t\ell}$ is the number of observations that belong to $\mathcal{P}_{t\ell}$. To sample from this expression, the Metropolis-Hastings algorithm is used, because a closed-form distribution is not obtained in this case.

As all $\mu_{t\ell}$ are i.i.d, it is possible to write $p(\mathcal{M}_t|\mathcal{T}_t, R_t, \sigma^2) = \prod_{\ell=1}^{b_t} p(\mu_{t\ell}|\mathcal{T}_t, R_t, \sigma^2)$. Similarly to the original BART, the full conditional of $\mu_{t\ell}$ in the AMBARTI model also depends only on the information provided by all trees, except by $\mathcal{T}_t$, via partial residual as R_t. Hence, the full conditional of $\mu_{t\ell}$ can be written as

$$p(\mu_{t\ell}|-) \propto p(R_t|\mathcal{M}_t, \mathcal{T}_t, \sigma^2) p(\mu_{t\ell}),$$

$$\propto \exp \left(-\frac{1}{2\sigma_*^2} (\mu_{t\ell} - \mu_{t\ell}^*)^2 \right),$$

which is a

$$\mathrm{N} \left(\frac{\sigma^{-2} \sum_{(i,j) \in \mathcal{P}_{t\ell}} r_{ij}^{(t)}}{n_{t\ell}/\sigma^2 + \sigma_\mu^{-2}}, \frac{1}{n_{t\ell}/\sigma^2 + \sigma_\mu^{-2}} \right).$$

Finally, after generating all predicted values for all trees, σ^2 can be updated based on

$$p(\sigma^2|-) \propto p(\mathbf{y}|\mathbf{g}, \mathbf{e}, \mathbf{X}, \mathcal{M}_t, \mathcal{T}_t, \sigma^2) p(\sigma^2)$$

(12)
$$\propto (\sigma^2)^{-\left(\frac{n+\nu}{2} + 1 \right)} \exp \left(-\frac{S + \nu\lambda}{2\sigma^2} \right),$$

where $S = \sum_{i=1}^{I} \sum_{j=1}^{J} (y_{ij} - \hat{y}_{ij})^2$ and $\hat{y}_{ij} = \mu + g_i + e_j + \sum_{t=1}^{T} h(\mathbf{x}_{ij}; \mathcal{T}_t, \mathcal{M}_t)$. The expression in (12) is an $\mathrm{IG}((n+\nu)/2, (S+\nu\lambda)/2)$. In algorithm 1, the full structure of the AMBARTI model is presented for one MCMC iteration.

Algorithm 1: AMBARTI Algorithm

Update μ, g_i and e_j;

for *t in 1:T* **do**

    Compute $R_t = \mathbf{y} - \left(\mu + \mathbf{g} + \mathbf{e} + \sum_{k \neq t}^{T} h(\mathbf{X}; \mathcal{T}_k, \mathcal{M}_k) \right)$;

    Propose a new tree $\mathcal{T}_t^*$ based on $\mathcal{T}_t$ by (double) growing, (double) pruning, changing or swapping;

    Accept the proposed tree with probability $\alpha(\mathcal{T}_t, \mathcal{T}_t^*) = \min \left\{ 1, \frac{p(R_t|\mathcal{T}_t^*, \sigma^2) p(\mathcal{T}_t^*)}{p(R_t|\mathcal{T}_t, \sigma^2) p(\mathcal{T}_t)} \right\}$;

    Update the node-level parameters $\mu_{t\ell}$ from $p(\mu_{t\ell}|-)$;

end

Update σ^2 sampling from $p(\sigma^2|-)$.

APPENDIX B: ORTHONORMALITY CONSTRAINTS OF THE AMMI MODEL

We recall the AMMI model is overparameterised, so constraints need to be imposed so that the parameters can be estimated (Josse et al., 2014). In this Section, we show how to apply the orthonormality constraints on γ_{iq} and δ_{jq} when simulating from the AMMI model.

Let $\boldsymbol{\gamma}$ be an $I \times Q$ matrix, $\boldsymbol{\delta}$ a $J \times Q$ matrix, and consider that γ_{iq} and δ_{jq} are elements in row i and column q of the corresponding matrices. The following constraints are considered: i) $\sum_{i=1}^{I} \gamma_{iq} = \sum_{j=1}^{J} \delta_{jq} = 0$, for $q = 1, \ldots, Q$; ii) $\boldsymbol{\gamma}^{\top} \boldsymbol{\gamma} = \boldsymbol{\delta}^{\top} \boldsymbol{\delta} = I_q$, where I_q represents an identity matrix of dimension q; iii) $\lambda_1 \geq \lambda_2 \geq \cdots \lambda_{q-1} \geq \lambda_q \geq 0$ and; iv) $\gamma_{1q} \geq 0$, for all $q = 1, \ldots, Q$.

To illustrate our the strategy to meet the constraints presented above, we take the γ_{iq} as an example, but this also works for δ_{jq}. First, we create S an $I \times Q$ matrix, where $s_{iq} \sim$ N$(0, \sigma_x^2 = 1)$. Here, s_{iq} could be sampled from other distributions, such as Gamma or Beta. In addition, we define M as an $I \times Q$ matrix with each element being the mean of the corresponding q column of S. Hence, we know that

$$B = S - M$$

$$\Rightarrow \mathbf{1}_I^{\top} B = 0$$

$$\not\Rightarrow B^{\top} B = \mathbb{I},$$

where $\mathbf{1}_I$ is a column vector of dimension I containing ones, B is, by construction, a full rank matrix and $B^{\top} B$ is symmetric. However, we find a matrix A such that $C = BA \Rightarrow C^{\top} C = \mathbb{I}$. That is, we know that

$$D = C^{\top} C = \mathbb{I}$$

$$\Rightarrow (BA)^{\top} BA = \mathbb{I}$$

$$\Rightarrow A^{\top} B^{\top} BA = \mathbb{I}$$

$$\Rightarrow B^{\top} B = A^{-\top} A^{-1}$$

$$\Rightarrow B^{\top} B = (AA^{\top})^{-1}$$

$$\Rightarrow (B^{\top} B)^{-1} = AA^{\top}$$

$$\Rightarrow (B^{\top} B)^{-1} = A^2 \text{ (by symmetry)}$$

$$\Rightarrow (B^{\top} B)^{-1/2} = A.$$

In the end, we have that $\boldsymbol{\gamma} = B(B^{\top} B)^{-1/2}$.

APPENDIX C: VISUALISING THE UNCERTAINTIES OF THE PARAMETER ESTIMATES IN THE AMBARTI MODEL

(a) 5% quantile

(b) median

(c) 95% quantile

Fig 12: 5%, median, and 95% quantiles for predicted yields from the AMBARTI model for the Irish VCU InnoVar data in 2015. The three parts of the graph allow us to address the uncertainties associated with the predicted response in yields described in Figure 10.

(a) 5% quantile

(b) median

(c) 95% quantile

Fig 13: 5%, median, and 95% quantiles for predicted yields from the AMBARTI model for the Irish VCU InnoVar data in 2015. The three parts of the graph allow us to address the uncertainties associated with the GxE interactions and the main effects obtained by AMBARTI. We can see that the credible interval associated to g_{18}/e_2, g_{10}/e_7, g_{13}/e_1, g_{11}/e_1, and g_{13}/e_3 (to list a few) do not contain zero.