Striatin 3 and MAP4K4 cooperate towards oncogenic growth and tissue invasion

in medulloblastoma

Jessica Migliavacca[1], Buket Züllig[1], Charles Capdeville**[1]**, Michael Grotzer[2] and Martin Baumgartner[1,*]

[1] Division of Oncology, Children's Research Center, University Children's Hospital Zürich, Zürich,

Switzerland

[2] Division of Oncology, University Children's Hospital Zürich, Zürich, Switzerland

*e-mail: Martin.Baumgartner@kispi.uzh.ch

The authors declare that they have no conflict of interest."

Abstract

MAP4K4 has been correlated with increased cell motility and reduced proliferation in mammalian cells.

Consequences and regulation of this dichotomous functionality of MAP4K4 in tumor cells remained

elusive. We find that MAP4K4 interacts with the STRN3 protein in medulloblastoma and that STRN3

and MAP4K4 exert opposing functions in Hippo tumor suppressor signaling and clonal growth. However,

depletion of either STRN3 or MAP4K4 reduces migration and tissue invasion, and concomitant loss of

both proteins halts tumor cell growth in the cerebellar tissue. Mechanistically, STRN3 couples MAP4K4

to the protein phosphatase 2A, which inactivates growth repressing activities of MAP4K4. In parallel,

STRN3 enables growth factor-induced PKCθ activation and the direct phosphorylation of VASP$_{S157}$ by

MAP4K4, which are necessary for efficient migration and tissue invasion. Thus, STRN3 controls

MAP4K4 functionality in growth factor-activated tumor cells, and the cooperation of the two proteins

promotes invasiveness through nPKCs and VASP and increases proliferation via inactivation of Hippo

signaling. Targeting of the STRN3-MAP4K4 cooperation could reactivate Hippo signaling and repress

invasiveness in growth factor-driven tumors.

Introduction

Medulloblastoma (MB) is the most common malignant brain tumor in children[1]. Four distinct molecular subgroups (Wingless (WNT), Sonic Hedgehog (SHH), group 3, and group 4) can be discriminated with a total of 12 subtypes[2–4]. The characteristic tendency of MB cells to locally infiltrate the cerebrospinal fluid (CSF) via leptomeningeal dissemination reduces the efficacy of current treatments[5]. One key driver of tissue invasion and growth in MB is the Fibroblast Growth Factor Receptor (FGFR) signaling pathway[6,7]. The molecular mechanisms that balance FGFR-driven tissue infiltration and growth in MB and other tumors remain incompletely understood, and no specific therapy is available to subvert this balance towards tumor suppression[8]. Tissue infiltration and growth rely on the complex orchestration of parallel, potentially opposing signaling pathways to overcome the incompatible concomitant induction of cell migration and growth. Improved anti-dissemination therapies should specifically repress both the pro-migratory and the proliferation promoting signaling in tumor cells. In this study we address the concomitant regulation of proliferation and invasiveness downstream of the activated FGFR, and we delineate the underlying molecular mechanisms and identify novel drug targets in MB.

The serine/threonine mitogen-activated protein kinase kinase kinase kinase 4 (MAP4K4/HGK/NIK), is upregulated in MB patient samples and acts as a critical promoter of receptor tyrosine kinase (RTK)-induced MB cell dissemination[9,10]. MAP4K4 controls a wide range of biological processes[11–16], which can be explained by the interaction of MAP4K4 with a number of structurally and functionally different effectors. The pro-migratory function of MAP4K4 is in part due to its capability to directly phosphorylate the ezrin, radixin, moesin (ERM) proteins, which mediate lamellipodium formation in response to growth-factor (GF) stimulation[17]. MAP4K4 couples GF signaling to actin polymerization through phosphorylation of the actin-related protein 2 (Arp2) subunit of the Arp2/3 complex, which increases its actin nucleation activity and triggers membrane protrusion in response to EGF[18]. In MB, MAP4K4 promotes turnover and activation of the receptor tyrosine kinase c-MET and of the β-1 integrin adhesion receptor, which is necessary for membrane protrusion at the leading edge of invading cells[10]. The pleiotropic functionality of MAP4K4 and its high expression in the healthy brain tissue argue against global pharmacological repression of its catalytic activity. Rather, more subtle interference strategies that target only subsets of its functions would be needed to repress MAP4K4 oncogenic activities specifically.

The related kinase Misshapen-like kinase (MINK)[19] and MAP4K4[20,21] were identified as

components of the Striatin-interacting phosphatase and kinase (STRIPAK) complex[22–24]. The

STRIPAK complex is a supramolecular scaffold assembled around the striatin family proteins (STRN,

STRN3, STRN4). STRNs act as regulatory subunits of the protein phosphatase PP2A and are

associated with STRN-interacting proteins (STRIP1/2) and with members of the germinal center kinase

(GCK) families [22,25]. The STRIPAK complex controls Hippo tumor suppressor signaling towards

YAP/TAZ transcriptional activation and proliferation by repressing associated Hippo-activating kinases

including MAP4K4. In this context, STRIPAK-MAP4K4 interaction contributes to small T antigen-

mediated cell transformation[21]. MAP4K4 interacts with STRN4, which on the one hand orchestrates

MAP4K4 repression and thereby causes decreased Hippo signaling[20]. On the other hand, STRN4

was also proposed to repress malignant characteristics of cancer cells by blocking MAP4K4 functions.

However, the mechanistic details how STRIPAK-controlled MAP4K4 contributes to tumorigenesis, in

particular migration and tissue invasion, remain unresolved[26].

Using proximity-dependent biotin identification (BioID), we identified MAP4K4 as a component of the

STRIPAK complex in MB cells. We explored whether MAP4K4-STRIPAK complex components are

implicated in FGFR-driven cell dissemination, tissue invasion and growth in MB, and we identified

underlying changes in relevant signaling pathways by kinase activity profiling. Our study identified the

MAP4K4-STRIPAK complex as a central hub and bifurcation point, which promotes tissue invasion and

growth in MB tumors. It provides relevant mechanistic insights in the regulation of these central

processes in MB tumor cells, and it runs against the recently established dogma of STRIPAK-MAP4K4

antagonism by demonstrating a cooperative activity of STRN3 and MAP4K4 towards increased

invasiveness.

Results

MAP4K4 is a component of the STRIPAK complex in MB cells

We used BioID to assess the MAP4K4 interactome and identified proteins biotinylated by FLAG-

tagged N- or C-terminally biotin ligase-fused (BioID2[27]) full-length MAP4K4 (BioID2-MAP4K4 and

MAP4K4-BioID2) or BioID2-FLAG alone through mass spectrometry (MS)[28,29] in HEK-293T cells

(Figure S1A). The expression of 3xFLAG-tagged MAP4K4-BioID2 was confirmed by anti-FLAG

immunoblot (IB) analysis and of biotinylation activity by horseradish peroxidase (HRP)-coupled

streptavidin binding (Figure S1B). Biotinylated proteins were enriched by affinity-capturing using

streptavidin-conjugated beads and subjected to MS. We identified a total of 156 MAP4K4-specific

proteins (Figure S1C, Table S1). Gene ontology (GO) analysis with these proteins predicted functions

such as endosomal transport[10], actin polymerization[18], or regulation of cell adhesion[16] (Figure

1A), and found enrichment of components of the STRIPAK complex (Figure 1A-C). We confirmed the

interaction of 3xFLAG-BioID2-MAP4K4 with STRN3, STRN4, and STRIP1 in HEK-293T cells by co-

immunoprecipitation analysis (Figure 1D). In agreement with other studies[30], these data confirm

MAP4K4 association with the STRIPAK complex in cells, possibly through direct interaction with STRN4

and STRN3.

We generated MB cell lines stably expressing either the 3XFLAG-BioID2-MAP4K4 fusion proteins or

BioID2 alone to test MAP4K4-STRIPAK complex interaction in these cells (Figure S1D). BioID2-

MAP4K4 localized to the cytoplasm, the primary residency of MAP4K4, where it co-localized with

biotinylated proteins (Figure S1E). In contrast, BioID2 distributes throughout the cells, with enrichment

in the nucleus. Streptavidin affinity purification-MS and anti-FLAG immunoprecipitation confirmed the

association of MAP4K4 with the STRIPAK complex in MB cells (Figure 1E,F, Figure S1F, Table S2).

MAP4K4 is dispensable for STRNs interaction with STRIP1, while STRN3 is necessary for MAP4K4

association with the complex (Figure 1G). Unlike other germinal center kinases that have been reported

to interact with STRIP1 in an activity-dependent manner[31], inhibition of MAP4K4 by GNE-495 did not

impact its constitutive association with STRN4, STRN3, and STRIP1 (Figure S1G). We generated DAOY

cells stably expressing 3XFLAG-tagged single domains or truncated versions of MAP4K4 (Figure 1H).

We found that N-terminally truncated MAP4K4 (ID1-ID2-CNH, 290-1273) pulled down STRN4 and

STRIP1 more effectively than C-terminally truncated KD-ID1-ID2 (1-954) (Figure 1H,I). The kinase

domain (KD, 1-289) or the intermediate domains (ID1, 290-634 or ID2, 635-954) alone are not sufficient

for pulling down STRN4 or STRIP1. Combined, these results suggest that the citron homology domain

(CNH, 954-1273) is required for MAP4K4-STRIPAK interaction.

STRIPAK complex controls bFGF-induced collagen I invasion of MB cells

STRIPAK complex members STRN4, STRN3, and STRIP1 are highly expressed in MB, and the

expression of STRNs correlates positively with site-specific phosphorylation of MAP4K4 in ID1 and

negatively with phosphorylation in ID2 *in vivo* ([32] and Figure S2A-D), suggesting a potential

contribution of these proteins to MAP4K4 regulation and MB pathogenesis. MAP4K4 promotes GF-

induced MB cell motility[9,10]. Hence, we tested the implication of STRIPAK complex proteins in

invasive motility in DAOY (SHH) MB cell lines by stable knockout (KO) of STRN4, STRN3, or STRIP1

(Figure S3A,B). Using the spheroid collagen I invasion assay (SIA[33]), we observed that STRN4 KO

abrogated bFGF-induced cell invasion in DAOY cells, while KO of either STRIP1 or STRN3 had little or

no effect (Figure S3C).

We next determined the effect of simultaneous depletion of various STRIPAK components on GF-

induced migration using siRNA (Figure S3D,E) in DAOY cells. siSTRN4 confirmed STRN4 implication

in bFGF-induced DAOY collagen I invasion, and siSTRN3 also reduced bFGF-induced invasion (Fig.

2A). In contrast, siSTRIP1 caused a marked increase in both EGF- and bFGF-induced dissemination.

Consistent with a functional redundancy of STRN3 and STRN4, we observed that combined depletion

of STRN3 and STRN4 had an additive effect on EGF- or bFGF-induced collagen I invasion. Thus,

STRN3 and STRN4 are, together with MAP4K4, required for GF-induced collagen I invasion in MB. We

confirmed the implication of STRIPAK components in migration control with another cell line, UW228,

an found that depletion of either MAP4K4, STRN4 or STRN3 reduced collagen I invasion (Figure 2B,

Figure S3F,G).

STRIPAK complex can regulate associated kinases by enabling dephosphorylation through the

PP2A catalytic subunit[34,35]. Inhibition of PP2A sensitizes tumors to radiation and/or chemotherapy

and attenuates cell migration[36,37]. We found that the PP2A inhibitor LB-100 decreased bFGF-induced

collagen I invasion in a dose-dependent manner (Figure 2C) without affecting cell viability (Figure S3H).

Thus, PP2A is not acting as a negative regulator of STRIPAK complex-associated MAP4K4 towards

GF-induced migration in MB cells, and STRN3/4 enable or facilitate the pro-invasive phenotype in GF-

stimulated MB cells.

STRN3 controls GF-induced tissue expansion of tumor cell

We used an *ex vivo* organotypic cerebellum slice culture model (OCSCs) to test STRIPAK implication

in MB tissue invasion (Figure S4A). OCSCs maintain key features of the normal cerebellum

cytoarchitecture and constitute a physiologically relevant microenvironment model[38], where FGFR

signaling contributes to MB growth and tissue invasion[6]. We first tested whether the depletion of

MAP4K4 or STRIPAK components prevented growth and tissue invasion of DAOY cells. We found that

STRN3 depletion caused a significant reduction of tumor area (Figure 3A). Treatment of the co-culture

with EGF enhanced brain tissue invasion of the tumor cells (Figure S4B,C). Depletion of MAP4K4

caused rounded cell morphology and moderately reduced EGF-induced invasion (Figure S4B-D).

Depletion of STRN3 caused compaction of the tumor cell spheroids and prevented tissue invasion also

under EGF-stimulation (Figure S4C,D). Conversely, depletion of STRIP1 induced a more infiltrative

behavior, similar to what was observed *in vitro* (Figure 2A)*.* The combined depletion of MAP4K4 and

STRN3 in DAOY cells caused a dramatic reduction of tumor area, both under basal condition and under

EGF stimulation (Figure 3A,B, Figure S4B-D). The reduced tumor area in cells with combined depletion

is not the result of impaired proliferation, as remaining cells display proportionally similar EdU

incorporation as the control (Fig. S4E).

To confirm STRN3 implication in tissue invasion, we implanted HD-MBO3 group 3 MB cell lines KO

for either MAP4K4, STRN4, STRN3, or STRIP1 (Figure S3B) in OCSCs. HD-MBO3 cells are invasive

and display a high growth rate[38]. We observed a moderate for MAP4K4 KO and a significant reduction

in the expansion for STRN3 KO cell-derived tumors, both after five days in basal or GF-stimulated

conditions (Figure 3C,D, Figure S5). Moreover, the number of EdU-positive nuclei was considerably

reduced in sgSTRN3 and sgSTRN4 but not in sgMAP4K4 cells (Figure 3E).

This data supports a model of STRIPAK complex control of growth and invasion of MB tumor cells

in the cerebellum tissue environment.

Opposing functions of STRN3/4 and MAP4K4 in cell proliferation and YAP/TAZ target gene regulation

Depletion of STRN3 and STRN4 but not of MAP4K4 reduced MB cell growth in the cerebellum tissue.

We therefore tested the implication of MAP4K4 and of STRIPAK complex components in colony-

formation and clonal growth. Depletion of MAP4K4 caused a marked increase in colony number

independent of growth factor stimulation in DAOY cells (Figure 4A). Conversely, depletion of STRN3

impaired colony formation and clonal growth. Consistently, depletion of either STRN4 or STRN3 caused

a significant decrease in the number of HD-MB03 colonies (Figure 4B), demonstrating that an intact

STRIPAK complex and in particular STRN3 was necessary for clonal growth and colony expansion in

MB cells. The effect of STRN3 depletion was particularly striking under bFGF stimulation in HD-MB03

cells, which promoted massive colony-forming activity in this group 3 MB model (Figure 4B).

MAP4K4 is an activator of the Hippo tumor suppressor pathway, thereby repressing YAP/TAZ target

gene expression and growth[39,40]. By antagonizing these growth-repressive MAP4K4 functions, the

STRIPAK complex halts Hippo pathway activation and promotes YAP/TAZ transcriptional

activity[30,41,42]. We found that MAP4K4 depletion significantly increased the expression of the

YAP/TAZ target genes *CTGF*, *CYR61*, and *ANKRD1*, which supports a Hippo-pathway activating

function of MAP4K4 in MB cells (Figure 4C,D). Conversely, suppression of STRN3 or STRN4 decreased

transcription of these genes and also reverted the transcriptional activation induced by MAP4K4

depletion (Figure 4C). Thus, MAP4K4 blocks MB cell proliferation, possibly by promoting the activation

of the Hippo pathway, whereas the STRIPAK complex and, in particular, STRN3, antagonizes Hippo

activation and is necessary for proliferation of MB cells (Figure S6A).

STRIPAK phosphatase PP2A antagonizes different Hippo kinases and thereby suppresses the

Hippo tumor-suppressing activity and promotes a YAP/TAZ transcriptional program[30,43].

Consistently, PP2A inhibitor LB-100 repressed transcription of YAP/TAZ target genes (Fig. 4E).

However, depletion of MAP4K4 restored YAP/TAZ target gene expression under LB-100 treatment,

indicating that PP2A acts as a negative regulator of MAP4K4 function towards Hippo pathway activation

(Figure 4E, Figure S6B). Furthermore, we found that bFGF stimulation promotes YAP/TAZ target genes

expression in DAOY and HD-MBO3 cells. This increased YAP/TAZ target gene expression requires

active PP2A and STRN3 (Figure 4F,G, Figure S6C,D), suggesting that STRN3 functions as Hippo

pathway antagonist in FGFR-activated MB cells by enabling PP2A-mediated repression of MAP4K4

(Figure 4E-G, Figure S6B-D).

Novel protein kinases C are downstream effectors of MAP4K4 and STRN3 in promigratory signaling

To identify effectors of MAP4K4 and of STRIPAK complex components downstream of FGFR

activation, we profiled changes in global kinase activities using the PamGene array technology. FGFR

signaling caused significantly increased activities of CDKs, PKC, SRC kinases, and FAK (Figure 5A,

Figure S7A). Several of these and additional bFGF-induced kinase activities require the expression of

MAP4K4 or of STRIPAK complex components (Figure 5A, Figure S7A). The most differently regulated

Ser/Thr (STK) kinases cluster in the AGC (cAMP-dependent, cGMP-dependent, and PKC) family of

kinases (Figure S7B-E). The activities of several AGC kinases were reduced when MAP4K4, STRN4,

or STRN3 were downregulated, and increased in STRIP1-depleted cells. Moreover, STRIP1 depletion

increased the activity of a MAP4K4 signature (Figure 5A), indicating that STRIP1 represses MAP4K4 in

MB cells. Interestingly, depletion of MAP4K4, STRN3 or STRN4, but not of STRIP1, caused a striking

decrease in bFGF-induced activity signature of "novel" protein kinases C (nPKCs, PKCδ, ε, η and θ)

(Figure 5A). This suggested that MAP4K4 and STRIPAK components are necessary for bFGF-induced

activation of nPKCs. Indeed, by IB analysis we could demonstrate that depletion of MAP4K4 or of

STRN3 reduced T538 phosphorylation of the catalytic domain of PKCθ (Figure 5B), which suggested

that MAP4K4 and STRN3 promote the activation of PKCθ[44].

Activated nPKCs could constitute druggable effector kinases to repress MB cell dissemination. We

tested this possibility using inhibitors specific for PKCθ (PKCθi[45] and Darovasertib[46]). We observed

a dose-dependent reduction of bFGF-induced cell dissemination in DAOY cells (Figure 5C, Figure S8A).

A similar effect was obtained using the PKCδ inhibitor Rottlerin (Figure S8B)[47]. Notably, the inhibition

of classical PKCα, β, γ kinases by Bisindolylmaleimide I (BIM) did not significantly repress MB cell

dissemination (Figure S8C). PKCθ or PKCδ inhibition also resulted in the compaction of DAOY and HD-

MBO3 tumor spheroids in the OCSC model and in the reduction of cerebellum tissue invasion, without

affecting MB cell proliferation (Figure 5D-F, Figure S8D-F). Depletion of PKCθ by siRNA (Figure S8G,H)

confirmed its implication in bFGF-induced DAOY collagen I invasion (Figure 5G). Moreover, PKCθ

depletion abolished bFGF-induced cerebellum tissue invasion of DAOY and HD-MBO3 cells (Figure

5H,I, Figure S8I,J).

Collectively these data demonstrate that MAP4K4 and STRN3 promote a pro-migratory phenotype

in MB cells through the activation of common downstream effectors, which include novel but not classical

PKC family members.

MAP4K4 and STRN3 enable bFGF-induced VASP Ser157 phosphorylation

Depletion of MAP4K4 in bFGF-stimulated DAOY cells resulted in an overall decrease in the

phosphorylation of multiple STK and PTK consensus phosphorylation sites (Figure 6A, Figure S9A,B,

Table S3). Many of these alterations were phenocopied by the depletion of STRN3 (Figure 6A).

Conversely, depletion of STRIP1 caused a global increase in the phosphorylation of STK substrates

(Figure 6A, Figure S9A). Phosphorylation profiles of PTK substrates were similar under all conditions

(Figure 6A, Figure S9B), with 23 substrates uniquely altered by MAP4K4 or STRN3 depletion (Figure

6B), including the phospholipase C-γ (PLC-γ)-binding site[48] of FGFR1 (FGFR$_{Y766}$), and the SRC

phosphorylation site of AKT (AKT$_{Y326}$)[49] (Figure 6C). Additionally, we observed the reduction of

FAK1$_{Y570/576}$ phosphorylation in siMAP4K4 and siSTRN3, supporting the previously described role of

MAP4K4 in mediating FAK protein activation and focal adhesion assembly[10] (Figure 6C).

Exploring potential direct substrates of MAP4K4, we found that both MAP4K4 and STRN3 depletion

decreased the phosphorylation of VASP$_{S157}$ (Figure 6C), an F-actin elongation promoting factor[50,51].

Phosphorylation of S157 enables VASP translocation to the plasma membrane at the leading edge of

migrating cells[52,53] and regulates anti-capping activity[51]. In bFGF-stimulated cells, VASP

accumulated in the cortical cytoskeleton and the plasma membrane (Figure 6D), and in cells invading

the brain tissue, we observed filopodia-like protrusions only in control but not in MAP4K4 or STRN3-

depleted cells (Figure S9C). Consistently, depletion of MAP4K4 or of STRN3 caused a robust reduction

of bFGF-induced $VASP_{S157}$ phosphorylation in cells seeded on collagen I matrix (Figure 6E). Under the

same condition, $VASP_{S157}$ phosphorylation is repressed by the MAP4K4 inhibitor GNE-495 in DAOY

and HD-MBO3 cells (Figure S9D,E), supporting the notion that MAP4K4 directly phosphorylates this

residue both under basal condition as well as in response to FGFR activation. Overexpression of full-

length MAP4K4 increased bFGF-induced $VASP_{S157}$ phosphorylation (Figure 6F). No increased

$VASP_{S157}$ phosphorylation was observed in cells overexpressing a C-terminally truncated MAP4K4 (KD-

ID1-ID2-FLAG, Figure 1H) or depleted of STRN3 (Figure 6F,G). To test whether the MAP4K4-STRN3

complex is able to directly phosphorylate recombinant VASP, we performed an *in vitro* kinase assay

using bacterially expressed VASP protein. MAP4K4 immunoprecipitated from bFGF stimulated cells

caused increased $VASP_{S157}$ phosphorylation *in vitro* (Figure 6G). We observed no increase in $VASP_{S157}$

phosphorylation when MAP4K4 is immunoprecipitated from STRN3-depleted cells, suggesting that

STRN3 is necessary to either activate MAP4K4 or to couple it to the VASP protein (Figure 6G). To test

the implication of VASP in bFGF-induced invasiveness in MB cells, we assessed siRNA-depletion of

VASP using the SIA (Figure S9F,G). We found that VASP depletion causes a significant but only partial

reduction of bFGF-induced collagen invasion (Figure 6H).

Taken together, these data identified $VASP_{S157}$ phosphorylation as a novel direct effector mechanism

of MAP4K4 and STRN3 downstream of FGFR signaling and highlighted VASP as a mediator of bFGF-

induced MB cell invasion.

Discussion

We found that MAP4K4 interacts with the STRIPAK complex, and that this interaction orchestrates

growth and invasion control downstream of FGFR signaling in MB cells (Fig. 7). Mechanistically,

MAP4K4 interaction with STRN3 and STRN4 represses MAP4K4 activity towards Hippo tumor

suppressor pathway activation by bringing the kinase in proximity of PP2A, which stalls MAP4K4

repression of YAP/TAZ target gene expression. Conversely, STRN3 and MAP4K4 interaction is

necessary for the activation of PKCθ and VASP, which we found to promote invasion downstream of

activated FGFR in MB. This axis of STRIPAK complex function is not repressed by PP2A. We furthermore revealed direct phosphorylation of VASP by MAP4K4 and found that VASP phosphorylation by MAP4K4 requires its interaction with STRN3. Thus, MAP4K4-mediated activation of PKCθ and VASP constitute a molecular link between the STRIPAK complex and the control of cytoskeleton dynamics. With this, the STRIPAK complex may balance the dichotomous functionality of MAP4K4 towards increased invasion and reduced growth repression in MB and other FGFR-driven tumors.

The interaction of the STRIPAK complex with MAP4K4 confirms previous findings describing a direct interaction of STRIPAK with MST1/2, MST3/4, and MAP4Ks[19,25,34,35] in a brain tumor, where MAP4K4 is highly expressed. MAP4K4 is a Hippo pathway kinase that through the direct phosphorylation of LATS1/2 caused repression of YAP/TAZ transcriptional activity. The STRIPAK complex promotes PP2A-mediated inactivation of MAP4K4 and thereby antagonizes Hippo signaling[30,39,42]. We confirmed this canonical function of MAP4K4 and of STRIPAK [30,41,42,54] and furthermore demonstrated that MAP4K4 depletion increases colony formation and clonal growth in two different MB tumor cell models. From these findings, we conclude that MAP4K4 and STRN3/4 have opposing functions towards Hippo signaling in MB, and that the STRIPAK complex exerts a growth-promoting function by suppressing the corresponding activity of MAP4K4 through the phosphatase PP2A in MB cells.

Parallel to antagonistic proliferation regulation of MAP4K4 and STRN3/4, we discovered that MAP4K4 and STRN3 display a co-operative function towards bFGF-induced invasion control. We identified AGC kinases and, in particular, novel PKCs (nPKCs), which are known promoters of tumorigenesis[55], as downstream effectors of MAP4K and STRN3 in invasion control, complementing previous reports of tumor invasion control by STRIPAK with mechanistic insight[26,56]. bFGF-driven invasion in MB may additionally involve the activation of adhesion and cytoskeleton-associated proteins like ezrin- radixin-moesin (ERM), β1 integrin, and focal adhesion kinase (FAK)[57–61], the activity of latter we also found increased by bFGF in a MAP4K4 and STRN3/4-dependent manner. The activation of nPKCs can be induced downstream of RTKs through phospholipase Cγ (PLCγ) signaling[62]. PLCγ is activated by its association with the phosphorylated Y766 residue of the activated FGFR[48]. Depletion of either MAP4K4 or of STRN3 not only decreased the nPKC activity signature but also decreased the phosphorylation of the Y766 residue on FGFR1. This indicates that regulation of Y766 phosphorylation could mechanistically link MAP4K4-STRIPAK function to PKC activation downstream

of bFGF stimulation. Specific targeting of PKCδ and PKCθ may thus contribute to an effective blockade

of FGFR-driven invasiveness in MB and other FGFR-driven tumors.

STRN3 and MAP4K4 are necessary for maintaining an elongated, mesenchymal motility mode in the

brain tissue, and cells devoid of STRN3 or of MAP4K4 display a rounded morphology that expresses

fewer filopodia-like protrusions. The involvement of MAP4K4 and STRIPAK complex members in

regulating the actomyosin cytoskeleton has been addressed in several studies[11,24]. However, the

interplay of MAP4K4 and the STRIPAK complex in cytoskeleton remodeling towards cancer cell

dissemination has not been addressed and a direct substrate of MAP4K4 in this process not identified.

Therefore, we consider the phosphorylation of $VASP_{S157}$ by MAP4K4 particularly relevant, because

phosphorylation of this residue could couple STRIPAK complex function to VASP localization,

cytoskeleton regulation, and cell migration[51,52]. MAP4K4-dependent $VASP_{Ser157}$ phosphorylation may

contribute to GF-induced MB cell motility and invasion by promoting cytoskeleton remodeling and the

formation of membrane protrusions. In support of this is our observation in previous studies[6] and herein

that tissue-invading MB tumor cells display filopodia-like protrusions, and that loss of these protrusions

is associated with reduced invasion. MAP4K4 phosphorylation of $VASP_{Ser157}$ *in vitro* depends on

MAP4K4 co-purification with STRN3, implying that VASP is a direct substrate of MAP4K4 and that

STRN3 is necessary for productive VASP phosphorylation by MAP4K4. It is possible that STRN3 brings

VASP and MAP4K4 in proximity. VASP was pulled down specifically by N-BioID2-M4K4 but not C-

BioID2-M4K4, indicating that VASP localizes in the proximity of the kinase domain of MAP4K4 in cells.

MAP4K4 and STRN3 also promote the activity of several AGC kinases that are implicated in $VASP_{Ser157}$

phosphorylation, like PKG and PKD1[63], which could act as additional VASP kinases downstream of

MAP4K4 in MB cells. Depletion of VASP caused a moderate reduction (-25%) of bFGF-induced collagen

I invasion. This indicates that VASP is unlikely the main driver of invasion control in MB. However,

additional functions of VASP in tumorigenesis argue for further clarification of the implication of $VASP_{S157}$

phosphorylation by MAP4K4 in MB.

Depletion of STRIP1 moderately increased GF-induced migration and tissue invasion in MB. The

increase in the MAP4K4 kinase activity signature after STRIP1 depletion further supports the notion of

an antagonizing function of STRIP1 towards MAP4K4. However, the lack of a clear phenotype after

STRIP1 depletion also indicates only a minor contribution to migration and growth control in MB tumor

cells. STRIP1 depletion has been associated with a contractile motility mode, supporting migration in

confined, stiff spaces[64]. Rounded, blebbing motility may contribute to brain invasion, albeit at lower

efficacy, and we previously observed a similar phenomenon in cells treated with the SRC-BCR/ABL

inhibitor dasatinib, which represses mesenchymal migration[65].

Together, our data indicate that the cooperation of MAP4K4 and STRN3 promotes motility in a 3D

environment and in the brain tissue. This motility depends in part on the activation of novel PKCs and

on the phosphorylation of VASP, which in concert with the previously identified regulation of ERM

proteins and the Arp2/3 complex by MAP4K4[10,17,18], may promote invasive motility in environments

of variable complexity and stiffness.

In conclusion, we found that MAP4K4 exerts pro-invasive and anti-proliferative activities in MB cells

and that STRN3 contributes to and modulates these MAP4K4 functions downstream of FGFR signaling.

This dichotomous functionality of MAP4K4 in tumor cells provides an explanation why MAP4K4

depletion alone is not sufficient to shrink the tumor volume *ex vivo,* despite the anti-invasive activity of

this treatment. Remarkably, however, co-depletion of MAP4K4 and STRN3 caused a near-complete

eradication of the tumor cells and completely abrogated cell dissemination *in vitro*. These findings argue

strongly for the development of STRN3 targeting strategies as therapeutic interventions to block MB cell

proliferation and invasion. Peptide disruption of the STRN3-PP2A interaction has anti-tumor effects by

impairing the phosphatase inactivation of the kinase in the complex and concomitant hippo pathway

activation[66]. However, PP2A inhibition repressed invasion in MB cells in a dose-dependent manner,

indicating that STRIPAK-associated PP2A is rather unlikely to repress MAP4K4 pro-invasive activity.

Rather, targeting of the cooperation of STRN3 and MAP4K4 to repress the pro-invasive activities of

MAP4K4 and to unleash its anti-proliferative functions through Hippo pathway activation could represent

a successful strategy to impair both MB cell proliferation and spreading. YAP/TAZ target gene

expression and the phosphorylation status of their common downstream effectors PKCθ and VASP in

patient samples may inform clinicians in the future about the potential susceptibility to therapies targeting

this pathway.

Material and methods

Reagents

Protein phosphatase PP2A inhibitor LB-100 (S7537), PKC-θ inhibitor (S6577), Darovasertib (S6723), Rottlerin (S7862), and Bisindolylmaleimide I (S7208) were purchased from Selleckchem, Houston, TX, USA. MAP4K4 inhibitor GNE-495 was kindly provided by Genentech Inc.[67]. EGF (100-47) and bFGF (100-18B) were purchased from PeproTech, London, UK. If not otherwise indicated, the cells were treated with growth factors at these concentrations: EGF: 30 ng/ml, bFGF: 100 ng/ml.

Cells and cell culture

DAOY human MB cells were purchased from the American Type Culture Collection (ATCC, Rockville, MD, USA) and cultured in Iscove's Modified Dulbecco's Medium (IMEM, Sigma). UW228 cells were generously provided by John Silber (Seattle, USA) and cultured in Dulbecco's Modified Eagle Medium (DMEM, Sigma). HD-MBO3 group 3 MB cells were generously provided by Till Milde (DKFZ, Germany) and cultured in RPMI medium (Sigma). Human embryonic kidney HEK-293T cells were cultured in Dulbecco's modified Eagle's medium (DMEM, Sigma). Cell culture media were supplemented with 10% fetal bovine serum (FBS, Sigma), 1% Penicillin-Streptomycin (ThermoFisher Scientific), and 1% GlutaMAX (Gibco). DAOY LA-EGFP and HD-MBO3 LA-EGFP cells were produced by lentiviral transduction of DAOY and HD-MBO3 cells with pLenti-LA-EGFP. The cells were maintained at 37°C in a humidified atmosphere containing 5% CO_2. All cells were confirmed to be mycoplasma negative. Cell line authentication and cross-contamination testing were performed by Multiplexion by single nucleotide polymorphism (SNP) profiling.

Plasmids

Lentiviral constructs were ordered from VectorBuilder (Santa Clara, CA, USA). For BioID experiments, 3xFLAG-tagged BioID2 biotin ligase was fused to either the N-terminus (3xFLAG-BioID2-MAP4K4) or the C-terminus (MAP4K4-BioID2-3xFLAG) of human MAP4K4 cDNA (NM_145686.4). An extended flexible linker consisting of 13 repeats of GGGGS was inserted between MAP4K4 and the BioID2 ligase[27]. As negative controls, only BioID2-FLAG was used. Details of the vectors can be found at https://en.vectorbuilder.com/vector/VB180410-1463qqf.html (N-BioID-MAP4K4), https://en.vectorbuilder.com/vector/VB180410-1356dmp.html (C-BioID2-MAP4K4), and https://en.vectorbuilder.com/vector/VB180411-1121jky.html (BioID2-CTRL). See also Figure S1. For

the analysis of MAP4K4 domains, we designed different constructs encoding 3xFLAG-tagged truncated

versions of MAP4K4 cDNA (Figure 1j). Details of the vector can be found at:

https://en.vectorbuilder.com/vector/VB190724-1021fsg.html (full-length MAP4K4),

https://en.vectorbuilder.com/vector/VB190724-1072aza.html (ID1-ID2-CNH, residues 290-1273),

https://en.vectorbuilder.com/vector/VB190724-1048drb.html (KD, residues 1-289),

https://en.vectorbuilder.com/vector/VB190724-1089brg.html (KD-ID1-ID2, residues 1-954),

https://en.vectorbuilder.com/vector/VB190724-1058sct.html (ID1, residues 290-634), and

https://en.vectorbuilder.com/vector/VB190724-1062xrk.html (ID2, residues 635-954). All plasmids were

verified by Sanger sequencing.

Mouse maintenance

Mouse protocols for organotypic brain slice culture were approved by the Veterinary Office of the Canton

Zürich (Approvals ZH134/17, ZH116/20). Wild type C57BL/6JRj pregnant females were purchased from

Janvier Labs and were kept in the animal facilities of the University of Zürich Laboratory Animal Center.

Immunoblotting (IB)

To analyze total cell lysates by immunoblot, 1×10^5 DAOY wild type or 2.5×10^5 HD-MBO3 cells were

lysed using RIPA buffer supplemented with protease (Complete Mini, Roche) and phosphatase

inhibitors (PhosSTOP, Roche) and cleared by centrifugation. Protein concentration was assessed using

the Pierce BCA Protein Assay Kit (Thermo Fisher Scientific) according to the manufacturer's

instructions. Protein separation was performed on Mini-Protean TGX (4-15% or 4-20%) SDS-PAGE gel

(Bio-Rad) and transferred to PVDF membranes (Bio-Rad). After 1 h of blocking with 5% non-fat milk,

membranes were probed with primary antibodies listed in Supplementary material (Table 1). GAPDH or

β-tubulin were used as internal loading control. HRP-linked secondary antibodies (1:5000) were used

to detect the primary antibodies. Chemiluminescence detection was carried out using ChemiDoc Touch

Gel and Western Blot imaging system (Bio-Rad). The integrated density of immuno-reactive bands was

quantified using ImageJ (National Institutes of Health, USA).

To assess PKCθ T538 and VASP S157 phosphorylation, the cells were seeded on a collagen-coated

plate and starved for 16 h before treatment with growth factors and inhibitors. Protein extraction was

performed by directly incubating the cells at 95°C for 5 minutes with a lysis buffer containing 2.5% SDS

and 125 mM Tris-HCl pH 6.8 supplemented with protease and phosphatase inhibitors. The lysates were

resolved by SDS-PAGE and analyzed by IB as described above.

Biotin-streptavidin affinity purification

The protocol for isolating proteins biotinylated by biotin ligase was adapted from the BioID method

[27,29]. For proximity-labeling experiments with HEK-293T, 7 x 10[6] cells were transiently transfected

with N-BioID2-M4K4, C-BioID2-M4K4, or BioID2-CTRL plasmids using jetPRIME reagent (Polyplus

Transfection) according to manufacturers' instructions. Non-transfected WT HEK-293T cells were used

as a negative control (to exclude biotin-independent binders). Twenty-four hours after the transfection,

the cells were incubated with 50 µM biotin for 16 h. For proximity-labeling experiments with DAOY MB

cells, we generated cells stably transfected with N-BioID2-M4K4, C-BioID2-M4K4, or BioID2-CTRL

lentivirus. 5 x 10[6] cells were incubated with biotin as above. Parental WT DAOY cells were used as a

negative control. After 16 h of biotin incubation and two PBS washes, the cells were lysed at 4°C with 1

ml of lysis buffer containing 50 mM Tris-HCl (pH 7.5), 150 nM NaCl, 1 mM EDTA, 1 mM EGTA, 1%

Triton X-100, 0.1% SDS and protease inhibitor tablet (Roche). The protein concentrations were

normalized among the samples. Two mg of protein samples were incubated with 75 µl of pre-equilibrated

Dynabeads MyOne Streptavidin T1 magnetic beads (65602, Thermo Fisher Scientific) in low-binding

Eppendorf tubes. The mixtures were incubated for 3 h at 4 °C with end-over-end rotation. The beads

were collected using a magnetic rack and washed twice with lysis buffer and twice with 50 mM

ammonium bicarbonate (pH 8.3). 10% of the streptavidin-conjugated beads were used for

immunoblotting. Proteins were eluted by heating the beads at 95°C for 5 min in Laemmli Sample Buffer

(1610747, Bio-Rad) with 50 mM DTT followed by magnetic separation. The remaining beads with bound

proteins were transferred to a fresh centrifuge tube, resuspend in 150 µl of 50 mM ammonium

bicarbonate, and used for MS analysis. For each condition, three biological replicates were performed

and analyzed by MS.

Mass spectrometry (MS)

Tryptic digestion of proteins isolated by streptavidin pull-down was performed based on the protocol

described in[28]. Data acquisition was performed by the Functional Genomics Center Zurich (FGCZ). In

brief, 50 µl of 8 M urea were added to 150 µl of bead-protein suspensions in 50 mM ammonium

bicarbonate (pH 8.3). Beads were reduced by adding 4 µl of 100 mM Tris (2-carboxyethyl) phosphine

(TCEP), alkylated with 4 µl of 500 mM chloroacetamide (CAA) and incubated at 30°C for 1 h in the dark

using an Eppendorf Thermomixer at 700 rpm. Sample volume was adjusted by adding 200 µl of digest

buffer (10 mM Tris-HCl, 2 mM CaCl$_2$, pH 8.2) to dilute the 8 M urea to 1 M before trypsin digestion. The

mixture was treated with 1 µg/sample trypsin and incubated for overnight digestion in the dark at 30°C

using an Eppendorf Thermomixer at 1000 rpm. The supernatant was collected in a fresh Eppendorf tube

after magnetic separation of the beads. The beads were washed twice with 100 µl 10 mM Tris-HCl pH

8.2/10% acetonitrile. These washes were pooled with the first eluate and acidified with 60 µl 5%

trifluoroacetic acid (TFA, final concentration 0.5%). The samples were desalted on Sep-Pack C18

columns (Thermo Scientific), completely dried using speed vac centrifugation, and dissolved in 20 µl

0.1% formic acid (FA). The samples were diluted 10-fold and transferred to autosampler vials for liquid

chromatography-tandem mass spectrometry (LC-MS/MS). 2 µl of samples were injected by an Easy-

nLC 1000 system (Thermo Scientific) and separated on an EasySpray-column (75 µm x 500 mm)

packed with C18 material (PepMap, C18, 100 Å, 2 µm, Thermo Scientific). The column was equilibrated

with 100% solvent A (0.1% FA in water). The column was equilibrated with 100% solvent A (0.1% FA in

water). Peptides were eluted using the following gradient of solvent B (0.1% FA in acetonitrile): 5-25%

B in, 60 min; 25-35% B in 10 min; 35-99% B in 5 min at a flow rate of 0.3 µl/min. All precursor signals

were recorded in the Orbitrap using quadrupole transmission in the mass range of 300-1500 m/z.

Spectra were recorded with a resolution of 120 000 at 200 m/z, a target value of 5E5 and the maximum

cycle time was set to 3 seconds. Data dependent MS/MS were recorded in the linear ion trap using

quadrupole isolation with a window of 1.6 Da and HCD fragmentation with 30% fragmentation energy.

The ion trap was operated in rapid scan mode with a target value of 8E3 and a maximum injection time

of 80 ms. Precursor signals were selected for fragmentation with a charge state from +2 to +7 and a

signal intensity of at least 5E3. A dynamic exclusion list was used for 25 seconds. The mass

spectrometry proteomics data were handled using the local laboratory information management system

(LIMS).

Proteomic data analysis

The MS/MS raw data of three biological replicates were combined and searched against a SwissProt

database with *Homo sapiens* taxonomy using Mascot (version 2.6.2, London, UK), with a parent ion

tolerance of 10 ppm and a fragment ion mass tolerance of 0.030 Da. The database was also expanded

with frequently observed contaminants, such as bovine serum albumin, trypsin, and keratins

carbamidomethylation of cysteine residues (C) was specified as a fixed modification and oxidation of methionine (M) was set as a variable modification. Scaffold (version 4.8.9, Proteome Software Inc., Portland, OR) was used to validate MS/MS-based peptide and protein identifications. 1% and 0.1% false discovery rate (FDR) cutoff on peptide and protein, respectively, was used to control for false identifications. Protein identifications were accepted if they contained at least 2 identified peptides per protein. Proteins that contained similar peptides and could not be differentiated based on MS/MS analysis alone were grouped to satisfy the principles of parsimony. Proteins sharing significant peptide evidence were grouped into clusters. Proteins detected in the parental control samples (WT cells lacking BioID2 expression) were subtracted from the results. The significance of enrichment was tested using a two-tailed student's *t*-test with equal variances. Hits were considered significantly enriched in MAP4K4-BioID2 cells compared the unspecific background (BioID2-CTRL) if they had > 2-fold enrichment and p-values < 0.05. Gene ontology (GO) enrichment analysis of MAP4K4-interacting partners was performed using Metascape (http://metascape.org/) [68]. All relevant data have been deposited to the ProteomeXchange Consortium via the PRIDE (http://www.ebi.ac.uk/pride) partner repository with the data set identifier PXD031863 and 10.6019/PXD031863 for the MAP4K4 interactome in HEK293T cells and PXD031870 and 10.6019/PXD031870 for for the MAP4K4 interactome in DAOY cells.

Co-immunoprecipitation (Co-IP)

2.5×10^6 DAOY or 3.5×10^6 HEK-293T cells were lysed at 4°C with 400 µl of buffer containing 50 mM Tris-HCl (pH 7.5), 150 nM NaCl, 0.3% NP-40 with protease and phosphatase inhibitor cocktail. Lysates were cleared by centrifugation, and protein concentrations were normalized among the samples. For immunoprecipitation, 30 µl of Dynabeads Protein G magnetic beads (10004D, Thermo Fisher Scientific) were coated with 3 µg of specific antibody or control IgG by resuspending them in 60 µl PBS containing 0.02% Tween-20 (P1379, Sigma), and rotated for 45 min at 4°C. 1 mg of protein samples was incubated with the antibody-coated Dynabeads for 3 h at 4 °C with end-over-end rotation. The beads were washed four times with TBS buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl). The proteins were eluted by incubating the beads with 35 µl 100 µg/ml 3xFLAG-Peptide (F4799, Sigma Aldrich) at room temperature for 20 min or by adding Laemmli Sample Buffer (1610747, Bio-Rad) containing 50 mM DTT and denaturation at 95°C for 5 min. The beads were then magnetically separated from the eluates and the samples analyzed by IB as described above.

 Immunofluorescence analysis (IFA)

To visualize MAP4K4-BioID2 proteins, stably transduced DAOY cells were plated on glass coverslips

and incubated with 50 µM biotin for 16 h. The cells were fixed and permeabilized as described in[69].

The fixed cells were incubated with Streptavidin Alexa Fluor 594 conjugate (Invitrogen) for 1.5 h at RT,

followed by incubation with primary antibodies overnight at 4°C. Secondary antibodies were incubated

for 2 h at RT.

For VASP localization analysis, 2000 LA-EGFP DAOY cells were seeded in a 384 well plate (781090,

Greiner Bio-One) coated overnight with 0.07 µg/µl collagen I (5005-B, Cell systems) diluted in 70%

EtOH. After seeding, the cells were starved for 16 h and treated with 100 ng/ml bFGF for 15 min. Cells

were fixed and incubated with primary and secondary antibodies as described above.

Imaging acquisition was performed using an SP8 confocal microscope (Leica). The list of the primary

and secondary antibodies used is provided in Supplementary material (Table 1). IFA Metadata can be

found in table S4.

Lentivirus Production and Transduction

Lentiviral particles were produced by transfection of HEK-293T cells with transfer plasmids, psPAX2

(#12260, Addgene) and pCMV-VSV-G plasmids (#8454, Addgene)[70] in a ratio of 5:3:2 using

polyethylenimine (24765-2, Polysciences). Virus supernatant was harvested 30 h after transfection,

filtered, and concentrated using Amicon Ultra 15 ml Centrifugal Filters (Millipore). For the transduction,

human MB cells were incubated for 24 hours with supernatant containing the viral particles and 8

µg/ml polybrene. Two days after transduction, cells were selected with puromycin (DAOY: 2 µg/ml, HD-

MBO3: 1 µg/ml).

Generation of CRISPR/Cas9-mediated knockout cells

Gene-specific single-guide RNA (sgRNAs) were designed using the online design tool at

https://design.synthego.com. Only highly specific target sites were selected, and the respective

sequences are listed in Supplementary material (Table 2). Oligonucleotides specific for the target sites

were synthesized with BsmBI restriction site overhangs by Microsynth (Balgach, Switzerland) and

cloned into the LentiCRISPRv2 transfer plasmid (Addgene plasmid #52961) with a single tube restriction

and ligation method as described in[71]. Production of lentiviral vectors and cell transduction was

performed as described above. The efficiency of the knockouts was tested by immunoblot.

RNA interference

DAOY, UW228, or HD-MBO3 cells at approximately 70% confluency were transfected with DharmaFECT Transfection Reagent 4 (T-2004-03, Dharmacon) or Lipofectamine RNAiMAX Transfection Reagent (13778075, Thermo Fisher Scientific) with 10 nM siRNAs following the manufacturer's instruction. The list of the siRNA used is provided in Supplementary material (Table 3). Non-Targeting Control siRNA (siCTRL, D-001210-02-05, Dharmacon) was used as a negative control for siRNA transfection. After 48 or 72 hours, RNA and proteins were isolated from the cells to determine gene expression by qRT-PCR and to evaluate protein expression by IB. For SIA or organotypic cerebellum slice culture experiements, the cells were re-seeded to form spheroids in ultra-low adhesion plates 24 h after the transfection.

RNA extraction and quantitative Real-Time PCR (qRT-PCR)

Total RNA was isolated from cells with the RNeasy mini kit (74106, Qiagen). cDNA was obtained by retro-transcription of 1 µg total RNA using High-capacity cDNA Reverse Transcription Kit (4368813, Applied Biosystems). qRT-PCR was performed using PowerUp Syber Green (A25776, Thermo Scientific) under conditions optimized for the 7900HT Fast Real-Time PCR System (Applied Biosystems). The primers used were synthesized by Microsynth AG (Balgach, Switzerland). Primer sequences are detailed in Supplementary material (Table 4). The relative expression levels of each gene of interest were calculated using the $2^{-\Delta\Delta Ct}$ method normalized to 18s.

Spheroid invasion assay (SIA)

SIA was performed and quantified according to[33]. In brief: 2500 cells/100 µl per well were seeded in cell-repellent 96 well microplates (650970, Greiner Bio-One, Kremsmünster, Austria). The cells were incubated at 37°C for 24 h to form spheroids. 70 µl medium was removed and replaced with 70 µl of a solution containing 2.5% bovine collagen I (CellSystems, Troisdorf, Germany). Polymerized collagen I hydrogels were overlaid with 100 µl of serum-free media containing growth factors and/or inhibitors 2x concentrated. The cells were allowed to invade the collagen matrix for 24 h, after which they were stained with Hoechst (1:2000, B2883, Sigma Aldrich) for 3-4 h. Images were acquired with an AxioObserver 2 mot plus fluorescence microscope (Zeiss, Munich, Germany) or with an Operetta high-content imaging system (PerkinElmer) at 5x magnification. When acquired with the Operetta microscope, the spheroids were localized in the well by acquiring 3 z-planes with 250 µm z-spacing, re-

scanned by acquiring 16 z-planes with 25 µm z-spacing and a maximum intensity projection was

computed and used for further analysis. Spheroids and invading cells were delineated based on

fluorescence threshold. Cell invasion was determined as the sum of the distance invaded by the cells

from the center of the spheroid as quantified using an automated cell dissemination counter (aCDc)[33]

or using Harmony 4.5 software (PerkinElmer).

Colony formation assay

DAOY (400 cells/well) or HD-MBO3 (600 cells/well) cells were seeded in a 6-well plate in complete

growth medium containing 10% FBS. The cells were allowed to adhere for 48 h, and the medium was

replaced with fresh growth medium containing 1% FBS ± bFGF. The cells were cultured at 37°C for 10

days (DAOY) or 15 days (HD-MBO3) with medium changes every third or fourth day. The colonies were

fixed in methanol for 10 min at -20°C and then stained with 0.5% crystal violet for 15 min. The number

and the area of the colonies were quantified by ImageJ using the plugin colony area.

CellTox green cytotoxicity assay

Cytotoxicity was determined using the CellTox^{TM} Green Cytotoxicity Assay (G8742, Promega,

Dübendorf, Switzerland), according to manufacturer's instructions. 2500 DAOY cells were seeded in

100 µl per well of complete medium in 96-well Corning black wall microplates with clear round bottom

(Corning Incorporated, NY, USA). The cells were incubated at 37°C for 24 h to form spheroids. 50 µl of

the medium were then removed from each well and replaced with 50 µL of fresh medium containing

1:500 of CellTox^{TM} Green Dye and different drug concentrations. Wells with medium but without cells

were used to measure background fluorescence. The fluorescence representing nonviable cells was

measured after 24, 48, and 72 h of incubation using Cytation 3 imaging reader (BioTek, Sursee,

Switzerland) at Ex/Em 485/520 nm.

Ex vivo organotypic cerebellum slice culture (OCSC)

Cerebellar slices were prepared from P8-10 C57BL/6JRj mouse pups as described in[38]. Briefly, Wild

type C57BL/6JRj mice pups were sacrificed at postnatal day (PND) 8–10 by decapitation. Cerebella

were dissected and kept in ice-cold Geys balanced salt solution containing kynurenic acid (GBSSK) and

then embedded in 2% low melting point agarose gel. Solidified agarose blocks were glued onto the

vibratome (VT 1200S, Leica, Wetzlar, Germany), and 350 µm thick sections were cut. Slices were

transferred on inserts (PICM 03050, Merck Millipore, Burlington, VT, USA) for further *in vitro* culture and

were kept in culture for 15 days. Medium was changed daily, and cultures were monitored for signs of

cell death. Tumor spheroids were formed from 2500 DAOY LA-EGFP or HD-MBO3 LA-EGFP cells

seeded in ultra-low adhesion plate for 48 h before implantation on OCSCs. Just before implantation,

spheroid size and integrity were evaluated under the microscope to confirm viability, integrity, and similar

spheroid size among the different conditions. The slice-spheroid co-cultures were treated with EGF (30

ng/ml), bFGF (12.5 ng/ml), PKCθ inhibitor (5 µM) or Rottlerin (5 µM) for 5 days where indicated. For

experiments with siRNA transfected cells, the cells were re-seeded to form spheroids in ultra-low

adhesion plate, implanted on the slices 48 h after transfection, and then grown on the slices for other

48 h with or without GF treatment. Following the treatment, the co-cultures were fixed with 4%

paraformaldehyde and stained for calbindin, GFAP, or anti-human nuclei, and analyzed using IFA as

described in[38]. To evaluate the extent of proliferation both in the tumor cells and in the slice culture,

we performed 5-ethynyl-2′-deoxyuridine (EdU) staining using the Click-iT EdU Alexa Fluor 647 imaging

kit (C10340, Invitrogen). Four-color image acquisition of three replicates per condition was performed

on an SP8 confocal microscope (Leica). The area of the infiltrating spheroids and the number of EdU-

positive nuclei were measured with ImageJ (Fiji)[72] using the tool analyzed particles after adjusting the

threshold to the same value for all the compared conditions.

In vitro VASP kinase assay

2×10^{6} DAOY cells expressing FLAG-tagged MAP4K4 were transfected with siCTRL or siSTRN3 as

described above. 48 h after the transfection and O/N starvation, cells were stimulated with 100 ng/ml

bFGF for 15 minutes and lysed at 4°C with 400 µl of buffer containing 50 mM Tris-HCl (pH 7.5), 150 nM

NaCl, 0.3% NP-40 supplemented with protease and phosphatase inhibitors. Parental WT DAOY cells

were used as a negative control. Lysates were cleared by centrifugation, and protein concentrations

were normalized among the samples. The supernatants were immune precipitated with 30 µl of

Dynabeads Protein G magnetic beads (10004D, Thermo Fisher Scientific) coated with anti-FLAG

monoclonal antibody (F1804, Sigma) for 3 h at 4°C. The beads were washed twice in lysis buffer and

twice with kinase assay buffer (20 mM Tris [pH 7.4], 200 mM NaCl, 0.5 mM dithiothreitol, 10 mM MgCl2).

The washed beads were resuspended in 25 µl of kinase assay buffer with 100 µM ATP (A7699, Sigma)

supplemented with phosphatase inhibitors and incubated for 20 min in an Eppendorf Thermomixer at

30°C at maximal mixing velocity. Then 4 µg of recombinant human VASP protein (ab105601, Abcam)

was added, and the reactions were allowed to continue for 10 min at 30°C. The kinase reactions were

stopped by adding Laemmli Sample Buffer (1610747, Bio-Rad) containing 50 mM DTT and boiling the

samples at 95°C for 5 min. The beads were then magnetically separated from the supernatant, and the

samples were analyzed by IB as described above.

Patient gene expression and proteomic data

Gene expression and proteomic data of pediatric brain tumor patients were obtained from the ProTrack

data portal (http://pbt.cptac-data-view.org/) from the Clinical Proteomic Tumor Analysis Consortium

(CPTAC) and the Children's Brain Tumor Tissue Consortium (CBTTC). The dataset used includes a

cohort of 218 tumor samples representing seven distinct histological diagnoses, including

medulloblastoma (MB), low-grade glioma (LGG), high-grade glioma (HGG), ependymoma (EP),

craniopharyngioma (CP), ganglioglioma (GG), and atypical teratoid rhabdoid tumor (ATRT)[32].

Protein kinase activity profiling

4.5×10^5 DAOY cells were transfected with siRNA as described above. 72 h after the transfection and

O/N starvation, cells were stimulated with 100 ng/ml bFGF for 15 minutes and lysed in M-PER protein

extraction reagent (Thermo Fischer Scientific) supplemented with Halt Phosphatase Inhibitor Cocktail

and Halt Protease Inhibitor Cocktail EDTA free (Thermo Fischer Scientific). Lysates were cleared by

centrifugation, and protein concentrations were determined by Pierce BCA Protein Assay Kit (Thermo

Fisher Scientific). Aliquots of 1 mg/ml of lysates were prepared and snap-frozen in liquid nitrogen. Kinase

activity profiles were performed using the PamChip® serine/threonine (STK) and protein tyrosine (PTK)

peptide microarray system from PamGene ('s-Hertogenbosch, The Netherlands), according to standard

manufacturer's instructions. The PamChip® consists of 4 identical peptide arrays, each array containing

144 (STK) or 196 (PTK) peptides. The technology measures the phosphorylation of the arrayed STK

and PTK peptides and compares derived peptide phosphorylation signatures to known consensus

signatures of kinases to predict kinase activities. Three PamChip® microarrays were processed at the

same time on the PamStation®12 (PamGene). One and five µg of protein extract were used for the STK

array and the PTK array protocol, respectively, and the enzymatic assay was started by adding ATP

(final concentration 100 µM). For the PTK array, fluorescein isothiocyanate (FITC)–labeled pY20

antibody was incubated on the chip, and the phosphorylation of the individual Tyr peptides was followed

by fluorescence detection in real-time. For the STK array, the primary antibody mixture was incubated

with the chip, and the secondary FITC- conjugated antibody was used to quantify the signal. For each

condition, three biological replicates were performed and analyzed in three different runs.

Software and data analysis

The fluorescent signal intensity for each peptide on the PamChip® was quantitated by BioNavigator

software (PamGene) and represented as linear regression slope as a function of exposure time. A local

background correction was used per spot. Nominal CV (Coefficient of variation) was calculated per

peptide with a 2-component error fit model, using the overall mean as input. Only peptides that showed

nominal CV <0.5 were included in the analysis. Additionally, Combat correction/normalization was

performed to correct for differences in batch effects. After averaging signal intensities across the 3

biological replicates per condition, the signal ratio between siTarget and siCTRL was used to calculate

fold change (FC) values. Peptides with a FC > 1.67 or FC < 0.6 were considered differentially

phosphorylated. To generate the peptide phosphorylation heatmap and compare the global

phosphorylation levels, the linear regression slope of each peptide was log_2 transformed. Peptides

differently phosphorylated in siTarget vs. siCTRL were identified by ANOVA and postHoc Dunnett's

tests. Venn diagram analyses were performed in Venny 2.1 (https://bioinfogp.cnb.csic.es/tools/venny/).

An upstream kinase prediction tool based on the PamApp on BioNavigator Analysis software was used

to generate a putative list of kinases responsible for phosphorylating the phosphosites on the PamChip.

The prediction of the differentially activated upstream kinases for each experimental group (siTarget vs.

siCTRL, or bFGF vs. UT) is carried out by a comparison of the phosphorylation sites of the peptides on

the array with databases containing experimental and literature-based protein modifications such as

HPRD, PhosphoELM (http://phospho.elm.eu.org), PhosphoSite Plus (https://www.phosphosite.org),

Reactome (http://reactome.org), UNIPROT (https://www.uniprot.org/) and in silico computational

predictions database such as PhosphoNet (http://www.phosphonet.ca/). The prediction tool functional

rank-orders the top kinases differentially activated between two compared groups. The mean kinase

statistic represents the difference in the predicted protein kinase activity between the two compared

groups. A positive value indicates an activation, while a negative value an inhibition. The specificity

score indicates the specificity of the normalized kinase statistics with respect to the number of peptides

used for predicting the corresponding kinase. The higher the specificity score, the higher is the power

of the prediction. Values of the specificity score >0.9 were considered as statistically relevant. To group

the kinases into sequence families, phylogenetic trees were created using the web-based kinome tool

CORAL (http://phanstiel-lab.med.unc.edu/CORAL/).

Statistical analysis

Statistical analyses were performed using GraphPad Prism 8 software (San Diego, California). Statistical significance of differences between groups was determined using Student's *t*-test (unpaired, two-tailed) or one-way ANOVA repeated-measures test with Bonferroni's correction. The experiments were performed in at least three independent biological replicates (*n*=3) unless differently indicated, and results are shown as mean ± standard deviation (SD) or standard error of the mean (SEM). The results were considered significant when $p < 0.05$ (*$p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$). Where indicated, asterisks show statistical significances between control and test sample.

References

1. Hatten, M. E. & Roussel, M. F. (2011) Development and cancer of the cerebellum. *Trends Neurosci.* **34**, 134–142.

2. Cavalli, F. M. G. *et al.* (2017) Intertumoral Heterogeneity within Medulloblastoma Subgroups. *Cancer Cell* **31**, 737--754.e6.

3. Schwalbe, E. C., Lindsey, J. C., Nakjang, S., Crosier, S., Smith, A. J., Hicks, D., Rafiee, G., Hill, R. M., Iliasova, A., Stone, T., Pizer, B., Michalski, A., Joshi, A., Wharton, S. B., Jacques, T. S., Bailey, S., Williamson, D. & Clifford, S. C. (2017) Novel molecular subgroups for clinical classification and outcome prediction in childhood medulloblastoma: a cohort study. *Lancet Oncol.* **18**, 958–971.

4. Hovestadt, V., Ayrault, O., Swartling, F. J., Robinson, G. W., Pfister, S. M. & Northcott, P. A. (2020) Medulloblastomics revisited: biological and clinical insights from thousands of patients. *Nat. Rev. Cancer* **20**, 42–56.

5. Wu, X. *et al.* (2012) Clonal selection drives genetic divergence of metastatic medulloblastoma. *Nature* **482**, 529–533.

6. Santhana Kumar, K., Neve, A., Guerreiro Stucklin, A. S., Kuzan-Fischer, C. M., Rushing, E. J., Taylor, M. D., Tripolitsioti, D., Behrmann, L., Kirschenbaum, D., Grotzer, M. A. & Baumgartner,

697   M. (2018) TGF-β Determines the Pro-migratory Potential of bFGF Signaling in Medulloblastoma.

698   *Cell Rep.* **23**, 3798-3812.e8.

699   7.   Neve, A., Migliavacca, J., Capdeville, C., Schönholzer, M. T., Gries, A., Ma, M., Kumar, K. S.,

700   Grotzer, M. & Baumgartner, M. (2019) Crosstalk between SHH and FGFR signaling pathways

701   controls tissue invasion in medulloblastoma. *Cancers (Basel).* **11**, 1985.

702   8.   Juraschka, K. & Taylor, M. D. (2019) Medulloblastoma in the age of molecular subgroups: A

703   review. *J. Neurosurg. Pediatr.* **24**, 353–363.

704   9.   Santhana Kumar, K., Tripolitsioti, D., Ma, M., Grählert, J., Egli, K. B., Fiaschetti, G., Shalaby, T.,

705   Grotzer, M. A. & Baumgartner, M. (2015) The Ser/Thr kinase MAP4K4 drives c-Met-induced

706   motility and invasiveness in a cell-based model of SHH medulloblastoma. *Springerplus* **4**, 19.

707   10. Tripolitsioti, D., Kumar, K. S., Neve, A., Migliavacca, J., Capdeville, C., Rushing, E. J., Ma, M.,

708   Kijima, N., Sharma, A., Pruschy, M., McComb, S., Taylor, M. D., Grotzer, M. A. & Baumgartner,

709   M. (2018) MAP4K4 controlled integrin β1 activation and c-Met endocytosis are associated with

710   invasive behavior of medulloblastoma cells. *Oncotarget* **9**, 23220–23236.

711   11. Tripolitsioti, D., Grotzer, M. A. & Baumgartner, M. (2017) The Ser/Thr Kinase MAP4K4 Controls

712   Pro-Metastatic Cell Functions. *J. Carcinog. Mutagen.* **08**, 1–8.

713   12. Flach, R. J. R., Skoura, A., Matevossian, A., Danai, L. V, Zheng, W., Cortes, C., Bhattacharya,

714   S. K., Aouadi, M., Hagan, N., Yawe, J. C., Vangala, P., Menendez, L. G., Cooper, M. P.,

715   Fitzgibbons, T. P., Buckbinder, L. & Czech, M. P. (2015) Endothelial protein kinase MAP4K4

716   promotes vascular inflammation and atherosclerosis. *Nat. Commun.* **6**, 8995.

717   13. Danai, L. V, Roth Flach, R. J., Virbasius, J. V, Garcia Menendez, L., Jung, D. Y., Kim, J. H., Kim,

718   J. K. & Czech, M. P. (2015) Inducible Deletion of Protein Kinase Map4k4 in Obese Mice Improves

719   Insulin Sensitivity in Liver and Adipose Tissues. *Mol. Cell. Biol.* **35**, 2356–2365.

720   14. Collins, C. S., Hong, J., Sapinoso, L., Zhou, Y., Liu, Z., Micklash, K., Schultz, P. G. & Hampton,

721   G. M. (2006) A small interfering RNA screen for modulators of tumor cell motility identifies

722   MAP4K4 as a promigratory kinase. *Proc. Natl. Acad. Sci.* **103**, 3775–3780.

723   15. Prolo, L. M., Li, A., Owen, S. F., Parker, J. J., Foshay, K., Nitta, R. T., Morgens, D. W., Bolin, S.,

724   Wilson, C. M., L, J. C. M. V., Luo, E. J., Nwagbo, G., Waziri, A., Li, G., Reimer, R. J., Bassik, M.

C. & Grant, G. A. (2019) Targeted genomic CRISPR-Cas9 screen identifies MAP4K4 as

essential for glioblastoma invasion. *Sci. Rep.* **9**, 1420.

16.   Vitorino, P., Yeung, S., Crow, A., Bakke, J., Smyczek, T., West, K., Mcnamara, E., Eastham-

anderson, J., Gould, S., Harris, S. F., Ndubaku, C. & Ye, W. (2015) MAP4K4 regulates integrin-

FERM binding to control endothelial cell motility. *Nature* **519**, 425–430.

17.   Baumgartner, M., Sillman, A. L., Blackwood, E. M., Srivastava, J., Madson, N., Schilling, J. W.,

Wright, J. H. & Barber, D. L. (2006) The Nck-interacting kinase phosphorylates ERM proteins for

formation of lamellipodium by growth factors. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 13391–13396.

18.   LeClaire, L. L., Rana, M., Baumgartner, M. & Barber, D. L. (2015) The Nck-interacting kinase

NIK increases Arp2/3 complex activity by phosphorylating the Arp2 subunit. *J. Cell Biol.* **208**,

161–170.

19.   Hyodo, T., Ito, S., Hasegawa, H., Asano, E., Maeda, M., Urano, T., Takahashi, M., Hamaguchi,

M. & Senga, T. (2012) Misshapen-like kinase 1 (MINK1) is a novel component of striatin-

interacting phosphatase and kinase (STRIPAK) and is required for the completion of cytokinesis.

J. Biol. Chem. **287**, 25019–25029.

20.   Seo, G., Han, H., Vargas, R. E., Yang, B., Li, X. & Wang, W. (2020) MAP4K Interactome Reveals

STRN4 as a Key STRIPAK Complex Component in Hippo Pathway Regulation. *Cell Rep.* **32**,

107860.

21.   Kim, J. W., Berrios, C., Kim, M., Schade, A. E., Adelmant, G., Yeerna, H., Damato, E., Iniguez,

A. B., Florens, L., Washburn, M. P., Stegmaier, K., Gray, N. S., Tamayo, P., Gjoerup, O., Marto,

J. A., DeCaprio, J. & Hahn, W. C. (2020) STRIPAK directs PP2A activity toward MAP4K4 to

promote oncogenic transformation of human cells. *Elife* **9**, 1–27.

22.   Goudreault, M. *et al.* (2009) A PP2A Phosphatase High Density Interaction Network Identifies a

Novel Striatin-interacting Phosphatase and Kinase Complex Linked to the Cerebral Cavernous

Malformation 3. *Mol. Cell. proteomics* **8**, 157–171.

23.   Chen, C., Shi, Z., Zhang, W., Chen, M., He, F., Zhang, Z., Wang, Y., Feng, M., Wang, W., Zhao,

Y., Brown, J. H., Jiao, S. & Zhou, Z. (2014) Striatins Contain a Noncanonical Coiled Coil That

Binds Protein Phosphatase 2A A Subunit to Form a 2 : 2 Heterotetrameric Core of Striatin-

interacting Phosphatase and Kinase (STRIPAK) Complex *. *J. Biol. Chem.* **289**, 9651–9661.

24.   Kück, U., Radchenko, D. & Teichert, I. (2019) STRIPAK , a highly conserved signaling complex

755   , controls multiple eukaryotic cellular and developmental processes and is linked with human

756   diseases Discovery of striatin. *Biol. Chem.* **400**, 1–18.

757   25.   Couzens, A. L., Knight, J. D. R., Kean, M. J., Teo, G., Weiss, A., Dunham, W. H., Lin, Z. Y.,

758   Bagshaw, R. D., Sicheri, F., Pawson, T., Wrana, J. L., Choi, H. & Gingras, A. C. (2013) Protein

759   interaction network of the mammalian hippo pathway reveals mechanisms of kinase-

760   phosphatase interactions. *Sci. Signal.* **6**, rs15.

761   26.   Wong, M., Hyodo, T., Asano, E., Funasaka, K., Miyahara, R., Hirooka, Y., Goto, H., Hamaguchi,

762   M. & Senga, T. (2014) Silencing of STRN4 suppresses the malignant characteristics of cancer

763   cells. *Cancer Sci.* **105**, 1526–1532.

764   27.   Kim, D. I., Jensen, S. C., Noble, K. A., KC, B., Roux, K. H., Motamedchaboki, K. & Roux, K. J.

765   (2016) An improved smaller biotin ligase for BioID proximity labeling. *Mol. Biol. Cell* **27**, 1188–

766   1196.

767   28.   Kim, D. I., Kc, B., Zhu, W., Motamedchaboki, K., Doye, V. & Roux, K. J. (2014) Probing nuclear

768   pore complex architecture with proximity-dependent biotinylation. *Proc. Natl. Acad. Sci.* **111**,

769   E2453-61.

770   29.   Roux, K. J., Kim, D. I., Raida, M. & Burke, B. (2012) A promiscuous biotin ligase fusion protein

771   identifies proximal and interacting proteins in mammalian cells. *J. Cell Biol.* **196**, 801–810.

772   30.   Kim, J. W. *et al.* (2020) STRIPAK directs PP2A activity toward MAP4K4 to promote oncogenic

773   transformation of human cells. *Elife* **9**, 1–27.

774   31.   Tang, Y., Chen, M., Zhou, L., Ma, J., Li, Y., Zhang, H., Shi, Z., Xu, Q., Zhang, X., Gao, Z., Zhao,

775   Y., Cheng, Y., Jiao, S. & Zhou, Z. (2019) Architecture, substructures, and dynamic assembly of

776   STRIPAK complexes in Hippo signaling. *Cell Discov.* **5**, 3.

777   32.   Petralia, F. *et al.* (2020) Integrated Proteogenomic Characterization across Major Histological

778   Types of Pediatric Brain Cancer. *Cell* **183**, 1962-1985.e31.

779   33.   Kumar, K. S., Pillong, M., Kunze, J., Burghardt, I., Weller, M., Grotzer, M. A., Schneider, G. &

780   Baumgartner, M. (2015) Computer-assisted quantification of motile and invasive capabilities of

cancer cells. *Sci. Rep.* **5**, 15338.

34.   Gordon, J., Hwang, J., Carrier, K. J., Jones, C. A., Kern, Q. L., Moreno, C. S., Karas, R. H. &

Pallas, D. C. (2011) Protein phosphatase 2a (PP2A) binds within the oligomerization domain of

striatin and regulates the phosphorylation and activation of the mammalian Ste20-Like kinase

Mst3. *BMC Biochem.* **12**, 1–18.

35.   Zheng, Y., Liu, B., Wang, L., Lei, H., Pulgar Prieto, K. D. & Pan, D. (2017) Homeostatic Control

of Hpo/MST Kinase Activity through Autophosphorylation-Dependent Recruitment of the

STRIPAK PP2A Phosphatase Complex. *Cell Rep.* **21**, 3612–3623.

36.   Ho, W. S., Feldman, M. J., Maric, D., Amable, L., Matthew, D., Feldman, G. M., Ray-chaudhury,

A., Lizak, M. J., Robison, R. A., Zhuang, Z. & Heiss, J. D. (2016) PP2A inhibition with LB100

enhances cisplatin cytotoxicity and overcomes cisplatin resistance in medulloblastoma cells.

Oncotarget **7**, 12447–12463.

37.   Yang, D., Okamura, H., Morimoto, H., Teramachi, J. & Haneji, T. (2016) Protein phosphatase 2A

C α regulates proliferation , migration , and metastasis of osteosarcoma cells. *Lab. Investig.*

96, 1050–1062.

38.   Neve, A., Kumar, K. S., Tripolitsioti, D., Grotzer, M. A. & Baumgartner, M. (2017) Investigation

of brain tissue infiltration by medulloblastoma cells in an ex vivo model. *Sci. Rep.* **7**, 1–12.

39.   Meng, Z., Moroishi, T., Mottier-Pavie, V., Plouffe, S. W., Hansen, C. G., Hong, A. W., Park, H.

W., Mo, J. S., Lu, W., Lu, S., Flores, F., Yu, F. X., Halder, G. & Guan, K. L. (2015) MAP4K family

kinases act in parallel to MST1/2 to activate LATS1/2 in the Hippo pathway. *Nat. Commun.* **6**, 1–

13.

40.   Zheng, Y., Wang, W., Liu, B., Deng, H., Uster, E. & Pan, D. (2015) Identification of

Happyhour/MAP4K as Alternative Hpo/Mst-like Kinases in the Hippo Kinase Cascade. *Dev. Cell*

34, 642–655.

41.   Ribeiro, P. S., Josué, F., Wepf, A., Wehr, M. C., Rinner, O., Kelly, G., Tapon, N. & Gstaiger, M.

(2010) Combined Functional Genomic and Proteomic Approaches Identify a PP2A Complex as

a Negative Regulator of Hippo Signaling. *Mol. Cell* **39**, 521–534.

42.   Chen, R., Xie, R., Meng, Z., Ma, S. & Guan, K. L. (2019) STRIPAK integrates upstream signals

to initiate the Hippo kinase cascade. *Nat. Cell Biol.* **21**, 1565–1577.

43.   Bae, S. J., Ni, L., Osinski, A., Tomchick, D. R., Brautigam, C. A. & Luo, X. (2017) SAV1 promotes

Hippo kinase activation through antagonizing the PP2A phosphatase STRIPAK. *Elife* **6**, 1–28.

44.   Liu, Y., Graham, C., Li, A., Fisher, R. J. & Shaw, S. (2002) Phosphorylation of the protein kinase

C-theta activation loop and hydrophobic motif regulates its kinase activity, but only activation

loop phosphorylation is critical to in vivo nuclear-factor-κB induction. *Biochem. J.* **361**, 255–265.

45.   Cywin, C. L. *et al.* (2007) Discovery of potent and selective PKC-θ inhibitors. *Bioorganic Med.*

Chem. Lett. **17**, 225–230.

46.   Frey, C. R., Wagle, M.-C., Vaidya, K., Hambleton, J., Lackner, M. & Mounir, Z. (2020) Analysis

of drug combinations with the PKC inhibitor IDE196 support dual MEK and PKC inhibition as a

rational combination in metastatic uveal melanoma. *Cancer Res.* **80**, 5337 LP – 5337.

47.   Gschwendt, M., Müller, H. J., Kielbassa, K., Zang, R., Kittstein, W., Rincke, G. & Marks, F. (1994)

Rottlerin, a novel protein kinase inhibitor. *Biochemical and Biophysical Research*

Communications **199**, 93–98.

48.   Mohammadi, M., Honegger, A. M., Rotin, D., Fischer, R., Bellot, F., Li, W., Dionne, C. A., Jaye,

M., Rubinstein, M. & Schlessinger, J. (1991) A tyrosine-phosphorylated carboxy-terminal peptide

of the fibroblast growth factor receptor (Flg) is a binding site for the SH2 domain of phospholipase

C-gamma 1. *Mol. Cell. Biol.* **11**, 5068–5078.

49.   Chen, R., Kim, O., Yang, J., Sato, K., Eisenmann, K. M., McCarthy, J., Chen, H. & Qiu, Y. (2001)

Regulation of Akt/PKB Activation by Tyrosine Phosphorylation. *J. Biol. Chem.* **276**, 31858–

31862.

50.   Skoble, J., Auerbuch, V., Goley, E. D., Welch, M. D. & Portnoy, D. A. (2001) Pivotal role of VASP

in Arp2/3 complex-mediated actin nucleation, actin branch-formation, and Listeria

monocytogenes motility. *J. Cell Biol.* **155**, 89–100.

51.   Barzik, M., Kotova, T. I., Higgs, H. N., Hazelwood, L., Hanein, D., Gertler, F. B. & Schafer, D. A.

(2005) Ena/VASP proteins enhance actin polymerization in the presence of barbed end capping

proteins. *J. Biol. Chem.* **280**, 28653–28662.

52.   Benz, P. M., Blume, C., Seifert, S., Wilhelm, S., Waschke, J., Schuh, K., Gertler, F., Münzel, T.

& Renné, T. (2009) Differential VASP phosphorylation controls remodeling of the actin
cytoskeleton. *J. Cell Sci.* **122**, 3954–3965.

53.   Harbeck, B., Hüttelmaier, S., Schlüter, K., Jockusch, B. M. & Illenberger, S. (2000)
Phosphorylation of the vasodilator-stimulated phosphoprotein regulates its interaction witn actin.
J. Biol. Chem. **275**, 30817–30825.

54.   Tang, Y. *et al.* (2020) Selective Inhibition of STRN3-Containing PP2A Phosphatase Restores
Hippo Tumor-Suppressor Activity in Gastric Cancer Article Selective Inhibition of STRN3-
Containing PP2A Phosphatase Restores Hippo Tumor-Suppressor Activity in Gastric Cancer.
Cancer Cell **38**, 115-128.e9.

55.   Garg, R., Benedetti, L. G., Abera, M. B., Wang, H., Abba, M. & Kazanietz, M. G. (2014) Protein
kinase C and cancer: What we know and what we do not. *Oncogene* **33**, 5225–5237.

56.   Du, Q. Y., Yao, J. H., Zhou, Y. C., Xu, L. J., Zhao, F. Y. & Yang, Y. (2020) High STRN Expression
Promotes HCC Invasion and Migration but Not Cell Proliferation or Apoptosis through Facilitating
Epithelial-Mesenchymal Transition. *Biomed Res. Int.* **2020**, 6152925.

57.   Lee, T. H., Chen, J. L., Liu, P. S., Tsai, M. M., Tsai, M. M., Wang, S. J., Hsieh, H. L. & Hsieh, H.
L. (2020) Rottlerin, a natural polyphenol compound, inhibits upregulation of matrix
metalloproteinase-9 and brain astrocytic migration by reducing PKC-δ-dependent ROS signal.
J. Neuroinflammation **17**, 1–13.

58.   Sarkar, S. & Yong, V. W. (2010) Reduction of protein kinase C delta attenuates tenascin-C
stimulated glioma invasion in three-dimensional matrix. *Carcinogenesis* **31**, 311–317.

59.   Brenner, W., Greber, I., Gudejko-Thiel, J., Beitz, S., Schneider, E., Walenta, S., Peters, K.,
Unger, R. & Thüroff, J. W. (2008) Migration of renal carcinoma cells is dependent on protein
kinase Cδ via β1 integrin and focal adhesion kinase. *Int. J. Oncol.* **32**, 1125–1131.

60.   Cannon, J. L., Asperti-Boursin, F., Letendre, K. A., Brown, I. K., Korzekwa, K. E., Blaine, K. M.,
Oruganti, S. R., Sperling, A. I. & Moses, M. E. (2013) PKCθ regulates T cell motility via ezrin-
radixin-moesin localization to the uropod. *PLoS One* **8**, e78940.

61.   Madaro, L., Marrocco, V., Fiore, P., Aulino, P., Smeriglio, P., Adamo, S., Molinaro, M. & Bouché,
M. (2011) PKCθ signaling is required for myoblast fusion by regulating the expression of

caveolin-3 and β1D integrin upstream focal adhesion kinase. *Mol. Biol. Cell* **22**, 1409–1419.

62.   Parker, P. J. & Murray-Rust, J. (2004) PKC at a glance. *J. Cell Sci.* **117**, 131–132.

63.   Döppler, H. & Storz, P. (2013) Regulation of VASP by phosphorylation Consequences for cell
868   migration. *Cell Adhes. Migr.* **7**, 482–486.

869   64.   Madsen, C. D., Hooper, S., Tozluoglu, M., Bruckbauer, A., Fletcher, G., Erler, J. T., Bates, P. A.,
870   Thompson, B. & Sahai, E. (2015) STRIPAK components determine mode of cancer cell migration
871   and metastasis. *Nat. Cell Biol.* **17**, 68–80.

872   65.   Schönholzer, M. T., Migliavacca, J., Alvarez, E., Santhana Kumar, K., Neve, A., Gries, A., Ma,
873   M., Grotzer, M. A. & Baumgartner, M. (2020) Real-time sensing of MAPK signaling in
874   medulloblastoma cells reveals cellular evasion mechanism counteracting dasatinib blockade of
875   ERK activation during invasion. *Neoplasia (United States)* **22**, 470–483.

876   66.   Tang, Y. *et al.* (2020) Selective Inhibition of STRN3-Containing PP2A Phosphatase Restores
877   Hippo Tumor-Suppressor Activity in Gastric Cancer. *Cancer Cell* **38**, 115-128.e9.

878   67.   Ndubaku, C. O. *et al.* (2015) Structure-Based Design of GNE-495, a Potent and Selective
879   MAP4K4 Inhibitor with Efficacy in Retinal Angiogenesis. *ACS Med. Chem. Lett.* **6**, 913–918.

880   68.   Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., Benner, C. &
881   Chanda, S. K. (2019) Metascape provides a biologist-oriented resource for the analysis of
882   systems-level datasets. *Nat. Commun.* **10**, 1523.

883   69.   Ma, M. & Baumgartner, M. (2014) Intracellular Theileria annulata Promote Invasive Cell Motility
884   through Kinase Regulation of the Host Actin Cytoskeleton. *PLoS Pathog.* **10**, e1004003.

885   70.   Stewart, S. A., Dykxhoorn, D. M., Palliser, D., Mizuno, H., Yu, E. Y., An, D. S., Sabatini, D. M.,
886   Chen, I. S. Y., Hahn, W. C., Sharp, P. A., Weinberg, R. A. & Novina, C. D. (2003) Lentivirus-
887   delivered stable gene silencing by RNAi in primary cells. *Rna* **9**, 493–501.

888   71.   McComb, S., Aguadé-Gorgorió, J., Harder, L., Marovca, B., Cario, G., Eckert, C., Schrappe, M.,
889   Stanulla, M., Von Stackelberg, A., Bourquin, J. P. & Bornhauser, B. C. (2016) Activation of
890   concurrent apoptosis and necroptosis by SMAC mimetics for the treatment of refractory and
891   relapsed ALL. *Sci. Transl. Med.* **8**, 339ra70.

72.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,

S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K.,

Tomancak, P. & Cardona, A. (2012) Fiji: An open-source platform for biological-image analysis.

Nat. Methods **9**, 676–682.

73.   Jeong, B., Bae, S. J., Ni, L., Zhang, X., Bai, X. & Luo, X. (2021) Cryo-EM structure of the Hippo

signaling integrator human STRIPAK. *Nat. Struct. Mol. Biol.* **28**, 290–299.

Acknowledgments

The authors thank Dr. Savithri Rangarajan (PamGene International) for scientific support and analysis

of PamChip imaging data and Dr. Jens Sobek (Functional Genomics Center Zurich) for assistance with

the experiments performed on PamStation®12. We would like to thank members of the Functional

Genomics Center Zurich for mass spectrometry support. Imaging was performed with equipment

maintained by the Centre for Microscopy and Image Analysis, University of Zurich. This study was

supported by grants from the Swiss National Science foundation (SNF_31003A_165860/1,

SNF_310030_188793) and the Swiss Cancer League (SCL_KLS-3834-02-2016) to MB, the University

of Zürich (Candoc) to J.M. and from the Childhood Cancer Foundation to MG.

Author contributions

909   J.M. contributed to designing the study, planned and conducted all the experiments, prepared the

910   figures, and wrote the manuscript. B.Z. generated DAOY cells stably expressing different MAP4K4

911   domains. C.C. provided experimental support for image acquisition and quantification with the Operetta

912   microscope. M.G. helped to draft the study. M.B. designed and drafted the study and wrote the

913   manuscript.

Figures

Figure 1: BioID identifies MAP4K4-STRIPAK complex interaction in MB cells. (A) Gene ontology
(GO) classifications of MAP4K4-interacting protein from BioID experiments conducted in HEK-293T.
Mass spectrometry was performed after immunoprecipitation (IP) with streptavidin-conjugated beads of
N- or C- BioID2-MAP4K4 and BioID2 alone overexpressing cells. **(B)** Volcano-plots of interacting

proteins in N–BioID2–MAP4K4 (left) or C–BioID2–MAP4K4 (right) HEK-293T cells compared to BioID2-CTRL. Log$_2$ ratios of cumulated peptide detection values plotted against negative Log$_{10}$ p-values of Student's *t*-test. Enriched proteins are highlighted in red (p < 0.05 and > 2-fold enrichment) and members of the STRIPAK complex indicated. *n*=3. **(C)** Architecture of the STRIPAK complex, adapted from[73]. **(D)** Immunoblot (IB) detection of BioID2-MAP4K4-FLAG and components of STRIPAK complex in anti-FLAG-IP HEK-293T cells. **(E)** Volcano-plots of protein interactions identified by streptavidin affinity purification-MS in N–BioID2–MAP4K4 (left) or C–BioID2–MAP4K4 (right) DAOY cells compared to BioID2-CTRL. Members of the STRIPAK complex are highlighted. *n*=2. **(F)** IB of MAP4K4-FLAG and components of STRIPAK complex after anti-FLAG-IP in DAOY cells. **(G)** IB analysis of MAP4K4 and components of STRIPAK complex after anti-STRN4 (top) or anti-STRIP1 (bottom) immunoprecipitation in DAOY cells transfected for 48 h with the indicated siRNA. Normal rabbit IgG was used as control for the IP. **(H)** Top: MAP4K4 domain architecture and MAP4K4 constructs used. Numbers represent the amino acids corresponding to full-length MAP4K4. KD, kinase domain; ID, intermediate domain; CNH, citron homology domain; CC, coiled-coil. Lower: Co-IP and IB analysis of STRN4 and STRIP1 in DAOY cells transduced with lentiviral vectors expressing full-length or truncated MAP4K4-FLAG. Asterisks indicate relevant bands. **(I)** Quantification of co-IP from H (*n*=3, means ± SEM). See also Figure S1.

Figure 2: STRIPAK complex disruption impairs GF-induced collagen I invasion. (A) Invasion and representative images of 3D spheroid invasion assay (SIA) of DAOY cells transfected with 10 nM of the indicated siRNA ± GFs (EGF: 30 ng/ml, bFGF: 100 ng/ml). Invasion is calculated as sum of the distances of each migrating cell from the center of the spheroids (*n*=4, means ± SEM). **(B)** SIA of UW228 cells transfected with 10 nM of the indicated siRNA ± 100 ng/ml bFGF (*n*=4, means ± SEM). **(C)** SIA and representative images of DAOY cells ± 100 ng/ml bFGF and ± increasing concentrations of PP2A inhibitor LB-100 (*n*=3, means ± SEM). * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001 (one-way ANOVA). See also Figure S3.

Figure 3: STRN3 is necessary for invasive growth of SHH and group 3 MB cell model in cerebellar tissue. (A) Maximum intensity projections (MIP) of representative confocal sections of organotypic cerebellum slice culture (OCSCs) implanted with DAOY tumor cell spheroids for 48h. Green: Lifeact-EGFP; blue: calbindin (Purkinje cells); red: GFAP. Middle: Inverted greyscale images of MIP of LA-EGFP channel, lower: Inverted greyscale images of MIP of EdU-Click-IT staining. Scale bar: 200 μm. **(B)** Quantification of LA-EGFP area of DAOY spheroids shown in A (*n*=3, means ± SD). **(C)** OCSCs culture implanted with tumor spheroids derived from HD-MBO3 cells KO for MAP4K4, STRN4, STRN3, or STRIP1. MIP of representative slices five days after implantation and ± treatment with 30 ng/ml EGF or 12.5 ng/ml bFGF. Green: Lifeact-EGFP; blue: calbindin (Purkinje cells); red: GFAP; yellow: Edu-Click-IT. Scale bar: 200 μm. **(D)** Areas of LA-EGFP positive HD-MBO3 spheroids showed in C (*n*=3, means ± SD). **(E)** Number of EdU-positive nuclei normalized by the area in pixels of the tumor spheroids (*n*=3, means ± SD). * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001 (unpaired Student's *t*-test). See also Figure S4 and S5.

Figure 4: STRN3/4 and MAP4K4 display opposing functions towards MB cell proliferation, clonal growth and YAP/TAZ target gene expression. (A, B) Representative images and quantification of colony formation assay of DAOY (A) and HD-MBO3 cells (B) KO for MAP4K4, STRN4, STRN3, or STRIP1 ± 100 ng/ml bFGF (n=3, means ± SEM). **(C)** qRT-PCR analysis of YAP/TAZ target genes in DAOY cells 48 h after transfection with siRNAs as indicated. (n=4, means ± SEM). **(D)** qRT-PCR analysis of YAP/TAZ target genes in HD-MBO3 with CRISPR/Cas9-mediated knockout of MAP4K4, STRN4, STRN3, or STRIP1 (n=3, means ± SEM). **(E)** qRT-PCR analysis of *CTGF* expression in DAOY cells 48 h after transfection with the indicated siRNA ± treatment for 4 h with 5 µM LB-100 (n=3, means ± SEM). **(F, G)** qRT-PCR analysis of *CTGF* expression in DAOY (F) or HD-MBO3 cells (G) transfected with siRNA for 48 h, serum-starved overnight, and treated with 100 ng/ml bFGF for 1 h and/or 5 µM LB-100 for 4 h where indicated (n=3, means ± SEM). * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001 (unpaired Student's *t*-test). See also Figure S6.

Figure 5: Novel protein kinases C are pro-migratory effectors of MAP4K4 and STRN3. (A) Kinase activity prediction analysis in DAOY cells transfected with the indicated siRNAs and ± stimulation with 100 ng/ml bFGF for 15 min. The plots show the predicted top 25 differentially activated serine/threonine kinases (STK). The x-axis indicates differences in the activity of the predicted protein kinase between the two groups, with effect size (values) and direction (>0=activation; <0=inhibition). The color of the bars represents the specificity score (darker to light is higher to lower score). Values >0.9 were considered as statistically relevant. First graph from left: comparison of bFGF stimulated siCTRL cells vs. untreated (UT); all other graphs: comparison of the indicated siTarget vs. siCTRL with bFGF stimulation. Highlighted in red are novel PKC isoforms (δ, ε, η and θ) (n=3). **(B)** IB analysis and quantification of PKCθ T_{538} phosphorylation in DAOY cells 72 h after siRNA transfection ± 100 ng/ml bFGF for 15 min (n=4, means ± SEM). **(C)** Quantification and representative images of SIA with DAOY cells stimulated with 100 ng/ml bFGF and treated with increasing concentrations of PKCθ inhibitor. Invasion was quantified as the sum of invasion distances from the center of the spheroids. DMSO was used as negative control (n=4, means ± SEM). **(D)** Maximum intensity projections (MIP) of representative confocal sections of OCSCs implanted with DAOY tumor cell spheroids and treated with 5 µM PKCθ inhibitor or DMSO for five days. Green: Lifeact-EGFP; blue: calbindin (Purkinje cells); red: GFAP; yellow: Edu-Click-IT. Scale bar: 200 µm. **(E)** Quantification of LA-EGFP area of DAOY spheroids shown in D (n=3, means ± SD). **(F)** Quantification of number of EdU-positive nuclei normalized by the area of DAOY spheroids shown in D (n=3, means ± SD). **(G)** Quantification and representative images of SIA of siCTRL or siPKCθ transfected DAOY cells ± 100 ng/ml bFGF (n=4, means ± SEM). **(H)** MIP of representative confocal sections of OCSC implanted with transfected DAOY tumor cell spheroids and treated with 12.5 ng/ml bFGF for 48h. Green: Lifeact-EGFP; blue: calbindin (Purkinje cells); red: GFAP; yellow: Edu-Click-IT. Scale bar: 200 µm. **(I)** Quantification of LA-EGFP area of DAOY spheroids shown in H (n=3, means ± SD). Statistical analyses in C and G were performed by one-way ANOVA, in B, E, F and I by unpaired Student's t-test. * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001. See also Figure S7 and S8.

Figure 6: MAP4K4 and STRN3 mediate VASP$_{S157}$ phosphorylation. (A) Heat map of increased (red) and decreased (blue) phosphorylation of Ser/Thr (STK, left) or Tyr (PTK, right) consensus peptides associated with lysates from siRNA transfected DAOY cells treated ± 100 ng/ml bFGF for 15 min. Log$_2$ of the fold change between siTarget and siCTRL (n=3). **(B)** Venn diagram displaying the number of Ser/Thr and Tyr peptides with significantly changed phosphorylation (p < 0.1) in each siTarget versus siCTRL with bFGF stimulation (n=3). **(C)** Volcano plots representing the changes in phosphorylation of STK (circles) and PTK (diamonds) peptides in siMAP4K4 (upper) or siSTRN3 (lower) versus siCTRL with bFGF stimulation. Blue: peptides with significantly decreased (fold change < 0.6), red: peptides with significantly increased phosphorylation (fold change > 1.67). The p-values were calculated versus siCTRL by ANOVA and post-hoc Dunnett's test in the BioNavigator software. n=3. See also Table S3. **(D)** Single confocal section of VASP localization in DAOY cells seeded on collagen-coated plates ± 100 ng/ml bFGF for 15 min. 3X magnifications of boxed areas are shown. Red: VASP, Green: Lifeact-EGFP. Blue: DNA Hoechst. Scale bar: 20 µm. **(E)** IB and quantification of VASP$_{S157}$ phosphorylation in DAOY cells 72 h after siRNA transfection ± 100 ng/ml bFGF for 15 min (n=4, means ± SEM). **(F)** IB and quantification of VASP$_{S157}$ phosphorylation in DAOY cells expressing full-length or truncated (KD-ID1-ID2) MAP4K4-FLAG ± 100 ng/ml bFGF for 15 min (n=3, means ± SEM). **(G)** In vitro VASP S157 phosphorylation of FLAG-immunoprecipitated DAOY cells expressing FLAG-tagged MAP4K4 and transfected for 48 h with the indicated siRNA ± 100 ng/ml bFGF for 15 min. The *in vitro* kinase reaction was conducted incubating the IP fraction with a recombinant VASP protein. Parental WT DAOY cells were used as a negative control for the IP. **(H)** Quantification and representative images of SIA of siCTRL or siVASP transfected DAOY cells ± 100 ng/ml bFGF (n=4, means ± SEM). Statistical analysis in E and F was performed by unpaired Student's t-test, in H by one-way ANOVA. * p < 0.05, ** p < 0.01, **** p < 0.0001. See also Figure S9.

Figure 7: Proposed model of MAP4K4 and STRN3 regulation of cell migration and proliferation in MB. MAP4K4 is activated downstream of RTK signaling and exerts both pro-invasive and anti-proliferative activities in MB cells. Invasion regulation: STRN3 cooperates with MAP4K4 to control cell migration, invasion, and actin cytoskeleton remodeling, independently from PP2A activity. This cooperation of MAP4K4 and STRN3 is necessary for the FGFR-induced activation of motility-associated effectors nPKCs and VASP. Proliferation regulation: MAP4K4 causes LATS1/2 phosphorylation, which promotes the activation of the Hippo pathway. Hippo pathway activation results in the phosphorylation and degradation of the transcriptional regulator YAP. The consequent repression of YAP transcriptional activity leads to a proliferation arrest. STRN3 promotes PP2A-mediated inactivation of MAP4K4 function toward Hippo activation, resulting in an increase of YAP transcriptional activity and cell proliferation.

Figure 1

Figure S1

Figure S3