The MAP4K4-STRIPAK complex promotes growth and tissue invasion in medulloblastoma

Jessica Migliavacca[1], Buket Züllig[1], Charles Capdeville[1], Michael Grotzer[2] and Martin Baumgartner[1,*]

[1] Division of Oncology, Children's Research Center, University Children's Hospital Zürich, Zürich, Switzerland

[2] Division of Oncology, University Children's Hospital Zürich, Zürich, Switzerland

*e-mail: Martin.Baumgartner@kispi.uzh.ch

Abstract

Proliferation and motility are mutually exclusive biological processes associated with cancer that depend on precise control of upstream signaling pathways with overlapping functionalities. We find that STRN3 and STRN4 scaffold subunits of the STRIPAK complex interact with MAP4K4 for pathway regulation in medulloblastoma. Disruption of the MAP4K4-STRIPAK complex impairs growth factor-induced migration and tissue invasion and stalls YAP/TAZ target gene expression and oncogenic growth. The migration promoting functions of the MAP4K4-STRIPAK complex involve the activation of novel PKCs and the phosphorylation of the membrane targeting S157 residue of VASP through MAP4K4. The anti-proliferative effect of complex disruption is associated with reduced YAP/TAZ target gene expression and results in repressed tumor growth in the brain tissue. This dichotomous functionality of the STRIPAK complex in migration and proliferation control acts through MAP4K4 regulation in tumor cells and provides relevant mechanistic insights into novel tumorigenic functions of the STRIPAK complex in medulloblastoma.

Introduction

Medulloblastoma (MB) is the most common malignant brain tumor in children[1]. Four distinct molecular subgroups (Wingless (WNT), Sonic Hedgehog (SHH), group 3, and group 4) can be discriminated with a total of 12 subtypes that display differences in genetic alterations, clinical features, and outcomes[2–4]. With multimodal therapies, the survival rates across all MB age groups have reached 70-85%[5–7]. However, the characteristic tendency of MB cells to locally infiltrate the cerebrospinal fluid (CSF) via leptomeningeal dissemination reduces the efficacy of current treatments[8]. One key driver of tissue invasion and growth in MB is the Fibroblast Growth Factor Receptor (FGFR) signaling pathway[9,10] The molecular mechanisms underlying MB tissue infiltration and growth are poorly understood and cannot be specifically targeted[11]. Infiltration, dissemination, and long-distance spreading of tumor cells are mediated by the transient or permanent induction of cell motility[12]. Cell migration relies on the complex orchestration of cytoskeletal remodeling, membrane dynamics, and cell adhesion to the extracellular matrix (ECM)[13–15]. Parallel, potentially redundant signaling pathways control these processes, and anti-dissemination therapies should specifically target the pro-migratory activity of a central component that integrates signals of several pathways.

We have identified the serine/threonine mitogen-activated protein kinase kinase kinase kinase 4 (MAP4K4/HGK/NIK), which is upregulated in MB patient samples, as a critical promoter of receptor tyrosine kinase (RTK)-induced MB cell dissemination[16,17]. MAP4K4 was originally identified as a Ste20 germinal center-like kinase[18] involved in the activation of the c-jun N-terminal kinase (JNK)[19,20]. MAP4K4 controls a wide range of biological processes[21–26], which can be explained by the interaction of MAP4K4 with a range of structurally and functionally different effectors. The pro-migratory function of MAP4K4 is in part due to its capability to directly phosphorylate the ezrin, radixin, moesin (ERM) proteins, which mediate lamellipodium formation in response to growth-factor (GF) stimulation[27]. MAP4K4 directly couples GF signaling to actin polymerization through phosphorylation of the actin-related protein 2 (Arp2) subunit of the Arp2/3 complex, which increases its actin nucleation activity and triggers membrane protrusion in response to EGF[28]. In MB, MAP4K4 promotes turnover and activation of the receptor tyrosine kinase c-MET and of the β-1 integrin adhesion receptor, which is necessary for membrane protrusion at the leading edge of invading cells[17]. However, the mechanisms orchestrating and balancing oncogenic functions of MAP4K4 in MB and other solid tumors remain unknown.

Using proximity-dependent biotin identification (BioID), we identified MAP4K4 as a component of the

recently identified Striatin-interacting phosphatase and kinase (STRIPAK) complex[29–31] in MB cells. We

explored whether MAP4K4-STRIPAK components are implicated in FGFR-driven cell dissemination,

tissue invasion and growth in MB, and we identified underlying changes in relevant signaling pathways

by kinase activity profiling. Our study identified the MAP4K4-STRIPAK complex as a central hub and

bifurcation point, which orchestrates tissue invasion and growth in MB tumors. It provides relevant

mechanistic insights in the regulation of these central processes in MB, which will contribute to the

design of more effective therapies for this devastating disease in the future.

Results

MAP4K4 is a component of the STRIPAK complex in MB cells

We used BioID to assess the MAP4K4 interactome and identified proteins biotinylated by FLAG-

tagged N- or C-terminally biotin ligase-fused (BioID2[32]) full-length MAP4K4 (BioID2-MAP4K4 and

MAP4K4-BioID2) or BioID2-FLAG alone through mass spectrometry (MS)[33,34] (Figure S1a). The

expression of 3xFLAG-tagged MAP4K4-BioID2 was confirmed by anti-FLAG immunoblot (IB) analysis

and of biotinylation activity by horseradish peroxidase (HRP)-coupled streptavidin binding (Figure 1a).

Biotinylated proteins were enriched by affinity-capturing using streptavidin-conjugated beads and

subjected to MS. We identified a total of 156 MAP4K4-specific proteins (Figure 1b, Table S1). Gene

ontology (GO) analysis with these proteins predicted functions such as endosomal transport[17], actin

polymerization[28], or regulation of cell adhesion[26] (Figure 1c), and found enrichment of components of

the STRIPAK complex (Figure 1c,d), a supramolecular scaffold held together by the striatin family

proteins (STRN, STRN3, STRN4). STRNs act as a regulatory subunit of the protein phosphatase PP2A

and are associated with STRN-interacting proteins (STRIP1/2) and with members of the germinal center

kinase (GCK) families (Figure 1e)[29,35]. We confirmed the interaction of 3xFLAG-BioID2-MAP4K4 with

STRN3, STRN4, and STRIP1 in HEK-293T cells (Figure 1f). In agreement with other studies[36–39], these

data confirm MAP4K4 association with the STRIPAK complex in cells, possibly through direct interaction

with STRN4 and STRN3.

We generated MB cell lines stably expressing either the 3XFLAG-BioID2-MAP4K4 fusion proteins or

BioID2 alone to test MAP4K4-STRIPAK complex interaction in these cells (Figure 1g). BioID2-MAP4K4

localized to the cytoplasm, the primary residency of MAP4K4, where it co-localized with biotinylated

proteins (Figure S1b). In contrast, BioID2 distributes throughout the cells, with enrichment in the nucleus.

Streptavidin affinity purification-MS and anti-FLAG immunoprecipitation confirmed the association of

MAP4K4 with the STRIPAK complex in MB cells (Figure 1h,i, Figure S1c, Table S2). Unlike other

germinal center kinases that have been reported to interact with STRIP1 in an activity-dependent

manner[40], inhibition of MAP4K4 by GNE-495 did not impact its constitutive association with STRN4,

STRN3, and STRIP1 (Figure S1d). We generated DAOY cells stably expressing 3XFLAG-tagged single

domains or truncated versions of MAP4K4 (Figure 1J). We found that N-terminally truncated MAP4K4

(ID1-ID2-CNH, 290-1273) pulled-down STRN4 and STRIP1 more effectively than C-terminally truncated

KD-ID1-ID2 (1-954) (Figure 1j,k). The kinase domain (KD, 1-289) or the intermediate domains (ID1, 290-

634 or ID2, 635-954) alone are not sufficient for pulling down STRN4 or STRIP1. Combined, these

results suggest that the citron homology domain (CNH, 954-1273) is required for MAP4K4-STRIPAK

interaction.

STRIPAK complex is required for bFGF-induced collagen I invasion of MB cells

STRIPAK complex members STRN4, STRN3, and STRIP1 are highly expressed in MB, and the

expression of STRNs correlates positively with site-specific phosphorylation of MAP4K4 in ID1 and

negatively with phosphorylation in ID2 *in vivo* ([41] and Figure S2a-d), suggesting a potential contribution

of these proteins to MAP4K4 regulation and MB pathogenesis. MAP4K4 promotes GF-induced MB cell

motility [16,17]. Hence, we first tested the implication of STRIPAK complex proteins in invasive motility in

DAOY (SHH) MB cell lines with a stable knockout (KO) of STRN4, STRN3, or STRIP1 (Figure S3a,b).

Using the spheroid collagen I invasion assay (SIA[42]), we observed that STRN4 KO abrogated GF-

induced cell invasion in DAOY cells, while KO of either STRIP1 or STRN3 had little or no effect (Figure

2A, Figure S3c).

We next determined the effect of simultaneous depletion of various STRIPAK components on GF-

induced migration using siRNA (Figure S3d,e). siSTRN4 confirmed STRN4 implication in bFGF-induced

collagen I invasion (Figure 2b). siSTRN3 also moderately reduced bFGF-induced invasion, whereas

siSTRIP1 caused a marked increase in both EGF- and bFGF-induced invasion (Figure 2b). siSTRIP1-

induced invasion was reverted when STRN4 was depleted simultaneously, indicating that the activity

repressed by STRIP1 involves a STRN3/4-dependent mechanism. Consistent with a functional

redundancy of STRN3 and STRN4, we observed that combined depletion of STRN3 and STRN4 had

an additive effect on EGF- or bFGF-induced collagen I invasion. Thus, STRN3 and STRN4 are, together

with MAP4K4, required for GF-induced collagen I invasion in MB, while STRIP1 has an inhibitory function in this process.

STRIPAK complex can regulate associated kinases by enabling dephosphorylation through the PP2A catalytic subunit[43,44]. Inhibition of PP2A sensitizes tumors to radiation and/or chemotherapy and attenuates cell migration[45,46]. The PP2A inhibitor LB-100 decreased EGF- and bFGF-induced collagen I invasion in a dose-dependent manner (Figure 2c, Figure S3f), without affecting cell viability (Figure S3g). PP2A is thus not acting as a negative regulator of STRIPAK complex-associated MAP4K4 towards GF-induced migration in MB, whereas STRN3/4 enable or facilitate the pro-invasive phenotype in GF-stimulated MB cells.

STRN3 controls GF-induced tissue expansion of tumor cell

We used an *ex vivo* organotypic cerebellum slice culture model (OCSCs) to test STRIPAK implication in MB tissue invasion. OCSCs maintain key features of the normal cerebellum cytoarchitecture and constitute a physiologically relevant microenvironment model[47], where FGFR signaling contributes to MB growth and tissue invasion[9]. We first tested whether the depletion of MAP4K4 or STRIPAK components prevented growth and tissue invasion of DAOY cells (Figure 3a). Treatment of the co-culture with EGF further enhances brain tissue invasion of the tumor cells. Depletion of MAP4K4 caused rounded cell morphology and moderately reduced EGF-induced invasion (Figure 3b,c, Figure S4). Depletion of STRN3 caused compaction of the tumor cell spheroids and prevented tissue invasion in basal and EGF-stimulated conditions. Conversely, depletion of STRIP1 induced a more infiltrative behavior, similar to what was observed *in vitro* (Figure 2b)*.* The combined depletion of MAP4K4 and STRN3 in DAOY cells caused a dramatic reduction of tumor area, which is not solely caused by reduced cell proliferation (Figure 3c,d).

To confirm STRIPAK complex implication in tissue invasion, we implanted HD-MBO3 group 3 MB cell lines KO for either MAP4K4, STRN4, STRN3, or STRIP1 (Figure S3b) in OCSCs. HD-MBO3 cells are invasive and display a high growth rate[47]. We observed a moderate for MAP4K4 KO and a significant reduction in the expansion for STRN3 KO cell-derived tumors, both after five days in basal or GF-stimulated conditions (Figure 4a,b, Figure S5). Moreover, the number of EdU-positive nuclei was considerably reduced in sgSTRN3 and sgSTRN4 but not in sgMAP4K4 cells (Figure 4c).

This data supports a model of STRIPAK-MAP4K4 complex control of growth and invasion of MB tumor cells in the cerebellum tissue environment.

 Opposing functions of STRN3/4 and MAP4K4 in cell proliferation and YAP/TAZ target gene regulation

STRN3 and STRN4 but not MAP4K4 depletion reduce MB cell growth in the cerebellum tissue. We

therefore tested MAP4K4-STRIPAK complex implication in colony-forming capability and clonal growth.

Depletion of MAP4K4 caused a marked increase in colony number independent of growth factor

stimulation in DAOY cells (Figure 5a). Conversely, depletion of STRN3 impaired colony formation and

clonal growth. Consistently depletion of either STRN4 or STRN3 caused a significant decrease in the

number of HD-MB03 colonies (Figure 5b), demonstrating that an intact STRIPAK complex is necessary

for clonal growth and colony expansion in MB cells. The effect of STRN3 depletion is particularly striking

under bFGF stimulation in HD-MB03, which promotes massive colony-forming activity in this group 3

MB model.

MAP4K4 is an activator of the Hippo tumor suppressor pathway, repressing YAP/TAZ target gene

expression and growth[48,49]. By antagonizing these growth-repressive MAP4K4 functions, the STRIPAK

complex halts Hippo pathway activation and promotes YAP/TAZ transcriptional activity[37,38,50,51]. We

found that MAP4K4 depletion significantly increased the expression of the YAP/TAZ target genes CTGF,

CYR61, and ANKRD1, which supports a Hippo-pathway activating function of MAP4K4 in MB cells

(Figure 5c,d). Conversely, suppression of STRN3 or STRN4 decreased transcription of these genes and

reverted the transcriptional activation induced by MAP4K4 depletion (Figure 5c).

Thus, MAP4K4 blocks MB cell proliferation, possibly by promoting the activation of the Hippo

pathway, whereas the STRIPAK complex and, in particular, STRN3, antagonizes Hippo activation and

is necessary for proliferation of MB cells (Figure S6).

Novel protein kinases C are downstream effectors of MAP4K4 and STRN3 in promigratory signaling

To identify effectors of the MAP4K4-STRIPAK complex downstream of FGFR activation, we profiled

changes in global kinase activities using the PamGene array technology. FGFR signaling caused

significantly increased activities of CDKs, PKC, SRC kinases, and FAK (Fig. 6a and Figure S7a). Several

of these and additional bFGF-induced kinase activities require the expression of MAP4K4 or STRIPAK

complex components (Figure 6a, Figure S7a). The most differently regulated Ser/Thr (STK) kinases

cluster in the AGC (cAMP-dependent, cGMP-dependent, and PKC) family of kinases (Figure S7b-e).

The activities of several AGC kinases were reduced when MAP4K4, STRN4, or STRN3 were

downregulated and increased in STRIP1-depleted cells. Consistently, depletion of MAP4K4, STRN3 or

STRN4, but not of STRIP1, caused a striking decrease in bFGF-induced activity of "novel" protein

kinases C (nPKCs, PKCδ, ε, η and θ) (Figure 6a). This suggests that MAP4K4 and STRIPAK

components are necessary for bFGF-induced activation of nPKCs and implies an effector role of nPKCs

downstream of MAP4K4 and STRN3. STRIP1 depletion increased the activity of a MAP4K4 signature

(Figure 6a), indicating that STRIP1 represses MAP4K4 in MB cells. The activated nPKC could constitute

druggable effector kinases to repress MB cell dissemination. We tested this possibility using inhibitors

specific for PKCδ (Rottlerin,[52]) and PKCθ (PKCθ I,[53]), and we observed a dose-dependent reduction of

bFGF-induced cell dissemination in DAOY cells (Figure 6b,c). The combination of both inhibitors further

reduced dissemination, suggesting that PKCδ and PKCθ functions in migration control are partially

overlapping (Figure 6d). Notably, the inhibition of classical PKCα, β, γ kinases by Bisindolylmaleimide I

(BIM) did not significantly repress MB cell dissemination (Figure 6e).

Collectively these data demonstrate that MAP4K4 and STRN3 promote a pro-migratory phenotype

in MB cells through the activation of common downstream effectors, which include novel but not classical

PKC family members.

MAP4K4 and STRN3 enable bFGF-induced VASP Ser157 phosphorylation

Depletion of MAP4K4 in bFGF-stimulated DAOY cells resulted in an overall decrease in the

phosphorylation of multiple STK and PTK consensus phosphorylation sites (Figure 7a, Figure S8, Table

S3). Many of these alterations were phenocopied by the depletion of STRN3 (Figure 7a). Conversely,

depletion of STRIP1 caused a global increase in the phosphorylation of STK substrates (Figure 7a,

Figure S8a). Phosphorylation profiles of PTK substrates were similar under all conditions (Figure 7a,

Figure S8b), with 23 substrates uniquely altered by MAP4K4 or STRN3 depletion (Figure 7b), including

the phospholipase C-γ (PLC-γ)-binding site[54] of FGFR1 (FGFR$_{Y766}$), and the SRC phosphorylation site

of AKT (AKT$_{Y326}$)[55] (Figure 7c). Additionally, we observed the reduction of FAK1$_{Y570/576}$ and FAK2$_{Y573/579}$

phosphorylation in siMAP4K4 and siSTRN3, supporting the previously described role of MAP4K4 in

mediating FAK protein activation and focal adhesion assembly[17] (Figure 7c).

Exploring potential direct substrates of MAP4K4, we found that both MAP4K4 and STRN3 depletion

decreased the phosphorylation of VASP$_{S157}$ (Figure 7c), an F-actin elongation promoting factor[56,57].

Phosphorylation of S157 enables VASP translocation to the plasma membrane at the leading edge of

migrating cells[58,59] and regulates anti-capping activity[57]. In bFGF-stimulated cells, VASP accumulated

in the cortical cytoskeleton and the plasma membrane (Figure 7d), and in cells invading the brain tissue,

we observed filopodia only in control but not in MAP4K4 or STRN3-depleted cells (Figure 7e).

Consistently, depletion of MAP4K4 or STRN3 caused a robust reduction of bFGF-induced $VASP_{S157}$

phosphorylation in cells seeded on collagen I matrix (Figure 7f). Under the same condition, $VASP_{S157}$

phosphorylation is repressed by the MAP4K4 inhibitor GNE-595 in DAOY (Figure 7g) and HD-MBO3

cells (Figure 7h), supporting the notion that MAP4K4 directly phosphorylates this residue both under

basal condition as well as in response to FGFR activation.

Taken together, these results identified $VASP_{S157}$ phosphorylation as a novel effector mechanism of

MAP4K4 and STRN3 downstream of FGFR signaling (Figure 7i).

Discussion

We found that MAP4K4 interacts with the STRIPAK complex in MB cells and revealed a dual function

of the complex in growth and invasion control. An intact complex is necessary to maintain proliferation

and to drive migration and invasion induced by FGFR signaling. Depletion of MAP4K4 shifts complex

function towards increased clonal growth, increased YAP/TAZ target gene expression, increased tumor

cell proliferation in the cerebellar tissue, and decreased invasiveness. Depletion of STRN3 repressed

clonal growth, reduced YAP/TAZ target gene expression, and caused a significant reduction of tumor

growth and invasion in the cerebellar tissue. We furthermore discovered a novel function of the

MAP4K4-STRIPAK complex as a regulator of a set of migration and motility-associated effectors. These

include novel PKCs, FAK, and VASP, which mechanistically link complex functions to cytoskeleton

dynamics and cell motility. Thus, the MAP4K4-STRIPAK complex orchestrates signal transmission from

activated RTKs either towards an invasion or a growth-promoting phenotype in MB.

The interaction of the STRIPAK complex with MAP4K4 we found in MB confirms previous findings

describing direct interaction of STRIPAK with MST1/2, MST3/4, and MAP4Ks in other cell

models[35,39,43,44]. In MB, MAP4K4 represses YAP/TAZ target gene induction and its depletion promotes

colony formation and clonal growth, consistent with a canonical function of the STRIPAK complex[37,38,51],

where STRN3/4 block growth suppressing activities of MAP4K4 and additional kinases [48,49,60]. From

these findings, we conclude that MAP4K4 and STRN3 have opposing functions towards Hippo signaling

in MB and that the STRIPAK complex exerts a growth promoting function by suppressing the activity of

MAP4K4 specific to this context.

Parallel to antagonistic proliferation regulation of MAP4K4 and STRN3/4, we discovered that

MAP4K4 and STRN3/4 display a co-operative function towards bFGF-induced invasion control. We

identified AGC kinases and, in particular, novel PKCs (nPKCs), which are known promoters of tumorigenesis[61], as downstream effectors of MAP4K-STRIPAK in invasion control, complementing previous reports of tumor invasion control by STRIPAK with mechanistic insight[62,63]. nPKC-driven invasion in MB may additionally involve the activation of adhesion and cytoskeleton-associated proteins like ezrin- radixin-moesin (ERM), β1D integrin, and focal adhesion kinase (FAK)[64–68], the activity of latter we also found increased by bFGF in a MAP4K4 and STRN3/4-dependent manner.

Inhibition of PKCδ and PKCθ but not of classical PKCs caused robust blockade of FGFR-driven invasiveness. However, additional mechanisms of MAP4K-STRIPAK invasion regulation are likely. We consider the phosphorylation of $VASP_{S157}$ particularly relevant because it couples STRIPAK function to VASP localization, cytoskeleton regulation, and cell migration[57,58]. We propose that MAP4K4-mediated $VASP_{Ser157}$ phosphorylation contributes to GF-induced MB cell motility and invasion by promoting filopodia formation. In support of this is our observation in previous studies[9] and herein that tissue-invading MB tumor cells display bFGF-induced, filopodia-like protrusions and that loss of these filopodia-like protrusions is associated with reduced invasion. We found that MAP4K4 and STRN3 promote the activity of several AGC kinases that are implicated in $VASP_{Ser157}$ phosphorylation, like PKG and PKD1[69]. VASP contributes to the etiology of prostate cancer, and $VASP_{Ser157}$ phosphorylation was recognized as a marker of metastatic progression[70]. $VASP_{Ser157}$ phosphorylation by STRIPAK-controlled MAP4K4 could have broad implications in controlling invasion and viability of MB cells in the brain tissue by providing signaling inputs from environment-sensing filopodia[71,72]. Growth cone turning and neuronal pathfinding depend on filopodia function[73], whereby in particular low and high amplitude Ca^{2+} signaling is involved in environment sensing and filopodia signaling[71]. Consistent with MB cells possibly exploiting this feature of the developing nervous system, we detected a calcium-dependent kinase signature downstream of bFGF-induced MAP4K4-STRIPAK. Depletion of STRIP1 increased GF-induced migration in MB, perhaps by antagonizing MAP4K4 function. In support of this is an increase in MAP4K4 kinase activity signature after STRIP1 depletion, thus positioning STRIP1 as a negative regulator of MAP4K4-driven mesenchymal invasiveness in MB. We observed a rounded, filopodia-depleted morphology in cells devoid of STRN3 or of MAP4K4. STRIP1 has been associated with a contractile motility mode, supporting migration in confined, stiff spaces[74]. Rounded, blebbing motility may contribute to brain invasion, albeit at lower efficacy, and we previously observed a similar phenomenon in cells treated with the SRC-BCR/ABL inhibitor dasatinib, which represses mesenchymal migration[75]. This

would explain why MAP4K4-depleted cells still display some invasiveness as MAP4K4 is mostly

involved in mesenchymal migration in MB cells. Together, our data indicate that co-operation of MAP4K4

and STRN3/4 favors filopodia formation through the induction of Ca^{2+}-activated kinases and the

phosphorylation of $VASP_{S157}$, which in concert with the previously identified regulation of ERM proteins

and the Arp2/3 complex by MAP4K4[17,27,28], promote invasive motility in environments of variable

complexity and stiffness.

In conclusion, we found that MAP4K4 exerts pro-invasive and anti-proliferative activities in MB cells

and that STRIPAK complex components contribute to and modulate these MAP4K4 functions

downstream of FGFR signaling. This dualism of MAP4K4 function in tumor cells provides an explanation

why MAP4K4 inhibition alone is not sufficient to shrink tumor volume *ex vivo,* despite its anti-invasive

activity. Remarkably however, co-depletion of MAP4K4 and STRN3 caused a near-complete eradication

of the tumor cells and completely abrogated cell dissemination *in vitro*. These findings argue strongly

for the development of STRN3 targeting strategies as therapeutic interventions to block MB cell

proliferation and invasion. Peptide disruption of the STRN3-PP2A interaction has anti-tumor effects by

impairing the phosphatase inactivation of the kinase in the complex and by activating the Hippo

pathway[50]. However, PP2A inhibition repressed invasion in MB cells in a dose-dependent manner,

indicating that STRIPAK-associated PP2A is rather unlikely to repress MAP4K4 pro-invasive activity

and that co-targeting of MAP4K4 and STRN3 or of their common downstream effectors such as nPKCs,

FAK and VASP could represent a successful strategy to impair both MB cell proliferation and spreading.

Phosphorylation status of these proteins in patient samples may inform clinicians in the future about the

potential susceptibility to therapies targeting this pathway.

Material and methods

Reagents

Protein phosphatase PP2A inhibitor LB-100 (S7537), PKC-θ inhibitor (S6577), Rottlerin (S7862), and

Bisindolylmaleimide I (S7208) were purchased from Selleckchem, Houston, TX, USA. MAP4K4 inhibitor

GNE-495 was kindly provided by Genentech Inc.[76]. EGF (100-47) and bFGF (100-18B) were purchased

from PeproTech, London, UK. If not otherwise indicated, the cells were treated with growth factors at

these concentrations*:* EGF: 30 ng/ml, bFGF: 100 ng/ml.

Cells and cell culture

DAOY human MB cells were purchased from the American Type Culture Collection (ATCC, Rockville,

MD, USA) and cultured in Iscove's Modified Dulbecco's Medium (IMEM, Sigma). HD-MBO3 group 3 MB

cells were generously provided by Till Milde (DKFZ, Germany) and cultured in RPMI medium (Sigma).

Human embryonic kidney HEK-293T cells were cultured in Dulbecco's modified Eagle's medium

(DMEM, Sigma). Cell culture media were supplemented with 10% fetal bovine serum (FBS, Sigma), 1%

Penicillin-Streptomycin (ThermoFisher Scientific), and 1% GlutaMAX (Gibco). DAOY LA-EGFP and HD-

MBO3 LA-EGFP cells were produced by lentiviral transduction of DAOY and HD-MBO3 cells with pLenti-

LA-EGFP. The cells were maintained at 37°C in a humidified atmosphere containing 5% CO_2. All cells

were confirmed to be mycoplasma negative. Cell line authentication and cross-contamination testing

were performed by Multiplexion by single nucleotide polymorphism (SNP) profiling.

Plasmids

Lentiviral constructs were ordered from VectorBuilder (Santa Clara, CA, USA). For BioID experiments,

3xFLAG-tagged BioID2 biotin ligase was fused to either the N-terminus (3xFLAG-BioID2-MAP4K4) or

the C-terminus (MAP4K4-BioID2-3xFLAG) of human MAP4K4 cDNA (NM_145686.4). An extended

flexible linker consisting of 13 repeats of GGGGS was inserted between MAP4K4 and the BioID2

ligase[32]. As negative controls, only BioID2-FLAG was used. Details of the vectors can be found

at https://en.vectorbuilder.com/vector/VB180410-1463qqf.html   (N-BioID-MAP4K4),

https://en.vectorbuilder.com/vector/VB180410-1356dmp.html   (C-BioID2-MAP4K4),   and

https://en.vectorbuilder.com/vector/VB180411-1121jky.html (BioID2-CTRL). See also Figure S1. For

the analysis of MAP4K4 domains, we designed different constructs encoding 3xFLAG-tagged truncated

versions of MAP4K4 cDNA (Figure 1j). Details of the vector can be found at:

https://en.vectorbuilder.com/vector/VB190724-1021fsg.html   (full-length   MAP4K4),

https://en.vectorbuilder.com/vector/VB190724-1072aza.html (ID1-ID2-CNH, residues 290-1273),

https://en.vectorbuilder.com/vector/VB190724-1048drb.html   (KD,   residues   1-289),

https://en.vectorbuilder.com/vector/VB190724-1089brg.html   (KD-ID1-ID2,   residues   1-954),

https://en.vectorbuilder.com/vector/VB190724-1058sct.html   (ID1,   residues   290-634),   and

https://en.vectorbuilder.com/vector/VB190724-1062xrk.html (ID2, residues 635-954). All plasmids were

verified by Sanger sequencing.

Mouse maintenance

Mouse protocols for organotypic brain slice culture were approved by the Veterinary Office of the Canton

Zürich (Approvals ZH134/17, ZH116/20). Wild type C57BL/6JRj pregnant females were purchased from

Janvier Labs and were kept in the animal facilities of the University of Zürich Laboratory Animal Center.

Immunoblotting (IB)

To analyze total cell lysates by immunoblot, 1×10^5 DAOY wild type or 2.5×10^5 HD-MBO3 cells were

lysed using RIPA buffer supplemented with protease (Complete Mini, Roche) and phosphatase

inhibitors (PhosSTOP, Roche) and cleared by centrifugation. Protein concentration was assessed using

the Pierce BCA Protein Assay Kit (Thermo Fisher Scientific) according to the manufacturer's

instructions. Protein separation was performed on Mini-Protean TGX (4-15% or 4-20%) SDS-PAGE gel

(Bio-Rad) and transferred to PVDF membranes (Bio-Rad). After 1 h of blocking with 5% non-fat milk,

membranes were probed with primary antibodies listed in Supplementary material (Table 1). GAPDH or

β-tubulin were used as internal loading control. HRP-linked secondary antibodies (1:5000) were used

to detect the primary antibodies. Chemiluminescence detection was carried out using ChemiDoc Touch

Gel and Western Blot imaging system (Bio-Rad). The integrated density of immuno-reactive bands was

quantified using ImageJ (National Institutes of Health, USA).

To assess VASP S157 phosphorylation, the cells were seeded on a collagen-coated plate and starved

for 16 h before treatment with growth factor and inhibitors. Protein extraction was performed by directly

incubating the cells with a lysis buffer containing 2.5% SDS and 125 mM Tris-HCl (pH 6.8) at 95°C for

5 minutes. The lysates were resolved by SDS-PAGE and analyzed by IB as described above.

Biotin-streptavidin affinity purification

The protocol for isolating proteins biotinylated by biotin ligase was adapted from the BioID method [32,34].

For proximity-labeling experiments with HEK-293T, 7×10^6 cells were transiently transfected with N-

BioID2-M4K4, C-BioID2-M4K4, or BioID2-CTRL plasmids using jetPRIME reagent (Polyplus

Transfection) according to manufacturers' instructions. Non-transfected WT HEK-293T cells were used

as a negative control (to exclude biotin-independent binders). Twenty-four hours after the transfection,

the cells were incubated with 50 µM biotin for 16 h. For proximity-labeling experiments with DAOY MB

cells, we generated cells stably transfected with N-BioID2-M4K4, C-BioID2-M4K4, or BioID2-CTRL

lentivirus. 5×10^6 cells were incubated with biotin as above. Parental WT DAOY cells were used as a

negative control. After 16 h of biotin incubation and two PBS washes, the cells were lysed at 4°C with 1

ml of lysis buffer containing 50 mM Tris-HCl (pH 7.5), 150 nM NaCl, 1 mM EDTA, 1 mM EGTA, 1%

Triton X-100, 0.1% SDS and protease inhibitor tablet (Roche). The protein concentrations were

normalized among the samples. Two mg of protein samples were incubated with 75 µl of pre-equilibrated

Dynabeads MyOne Streptavidin T1 magnetic beads (65602, Thermo Fisher Scientific) in low-binding

Eppendorf tubes. The mixtures were incubated for 3 h at 4 °C with end-over-end rotation. The beads

were collected using a magnetic rack and washed twice with lysis buffer and twice with 50 mM

ammonium bicarbonate (pH 8.3). 10% of the streptavidin-conjugated beads were used for

immunoblotting. Proteins were eluted by heating the beads at 95°C for 5 min in Laemmli Sample Buffer

(1610747, Bio-Rad) with 50 mM DTT followed by magnetic separation. The remaining beads with bound

proteins were transferred to a fresh centrifuge tube, resuspend in 150 µl of 50 mM ammonium

bicarbonate, and used for MS analysis. For each condition, three biological replicates were performed

and analyzed by MS.

Mass spectrometry (MS)

Tryptic digestion of proteins isolated by streptavidin pull-down was performed based on the protocol

described in[33]. Data acquisition was performed by the Functional Genomics Center Zurich (FGCZ). In

brief, 50 µl of 8 M urea were added to 150 µl of bead-protein suspensions in 50 mM ammonium

bicarbonate (pH 8.3). Beads were reduced by adding 4 µl of 100 mM Tris (2-carboxyethyl) phosphine

(TCEP), alkylated with 4 µl of 500 mM chloroacetamide (CAA) and incubated at 30°C for 1 h in the dark

using an Eppendorf Thermomixer at 700 rpm. Sample volume was adjusted by adding 200 µl of digest

buffer (10 mM Tris-HCl, 2 mM $CaCl_2$, pH 8.2) to dilute the 8 M urea to 1 M before trypsin digestion. The

mixture was treated with 1 µg/sample trypsin and incubated for overnight digestion in the dark at 30°C

using an Eppendorf Thermomixer at 1000 rpm. The supernatant was collected in a fresh Eppendorf tube

after magnetic separation of the beads. The beads were washed twice with 100 µl 10 mM Tris-HCl pH

8.2/10% acetonitrile. These washes were pooled with the first eluate and acidified with 60 µl 5%

trifluoroacetic acid (TFA). The Samples were desalted on SepPack C18 (Thermo Fisher Scientific),

dried, and dissolved in 20 µl 0.1% formic acid. The samples were diluted 10-fold and transferred to

autosampler vials for liquid chromatography-tandem mass spectrometry (LC-MS/MS). 2 µl of samples

were injected. LC-MS/MS analysis was performed on an Orbitrap Fusion Tribrid Mass Spectrometer

(Thermo Fisher Scientific). Peptides were separated by Easy Nano LC 1000 liquid chromatography

(Thermo Fisher Scientific) at 0.3 ml/min flow rate interfaced to EASY-Spray Source (Thermo Fisher

Scientific). A trap column (Acclaim PepMap 100 C18) and an analytical column (EASY-Spray C18) were

used for separation. Peptides were eluted using an acetonitrile gradient.

Proteomic data analysis

The MS/MS raw data of three biological replicates were combined and searched against a SwissProt

database with *Homo sapiens* taxonomy using Mascot (version 2.6.2, London, UK), with a parent ion

tolerance of 10 ppm and a fragment ion mass tolerance of 0.030 Da. The database was also expanded

with frequently observed contaminants, such as bovine serum albumin, trypsin, and keratins

carbamidomethylation of cysteine residues (C) was specified as a fixed modification and oxidation of

methionine (M) was set as a variable modification. Scaffold (version 4.8.9, Proteome Software Inc.,

Portland, OR) was used to validate MS/MS-based peptide and protein identifications. 1% and 0.1% false

discovery rate (FDR) cutoff on peptide and protein, respectively, was used to control for false

identifications. Protein identifications were accepted if they contained at least 2 identified peptides per

protein. Proteins that contained similar peptides and could not be differentiated based on MS/MS

analysis alone were grouped to satisfy the principles of parsimony. Proteins sharing significant peptide

evidence were grouped into clusters. Proteins detected in the parental control samples (WT cells lacking

BioID2 expression) were subtracted from the results. The significance of enrichment was tested using

a two-tailed student's *t*-test with equal variances. Hits were considered significantly enriched in

MAP4K4-BioID2 cells compared the unspecific background (BioID2-CTRL) if they had > 2-fold

enrichment and p-values < 0.05. Gene ontology (GO) enrichment analysis of MAP4K4-interacting

partners was performed using Metascape (http://metascape.org/) [77].

Co-immunoprecipitation (Co-IP)

2.5×10^6 DAOY or 3.5×10^6 HEK-293T cells were lysed at 4°C with 400 µl of buffer containing 50 mM

Tris-HCl (pH 7.5), 150 nM NaCl, 0.3% NP-40 with protease and phosphatase inhibitor cocktail. Lysates

were cleared by centrifugation, and protein concentrations were normalized among the samples. For

immunoprecipitation, 30 µl of Dynabeads Protein G magnetic beads (10004D, Thermo Fisher Scientific)

were coated with 3 µg of anti-FLAG antibody by resuspending them in 60 µl PBS containing 0.02%

Tween-20 (P1379, Sigma), and rotated for 45 min at 4°C. 1 mg of protein samples was incubated with

the antibody-coated Dynabeads for 3 h at 4 °C with end-over-end rotation. The beads were washed four

times with TBS buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl). The proteins were eluted by incubating

the beads with 35 µl 100 µg/ml 3xFLAG-Peptide (F4799, Sigma Aldrich) at room temperature for 20 min

or by adding Laemmli Sample Buffer (1610747, Bio-Rad) containing 50 mM DTT and denaturation at

95°C for 5 min. The beads were then magnetically separated from the eluates and the samples analyzed

by IB as described above.

Immunofluorescence analysis (IFA)

To visualize MAP4K4-BioID2 proteins, stably transduced DAOY cells were plated on glass coverslips

and incubated with 50 µM biotin for 16 h. The cells were fixed and permeabilized as described in[78]. The

fixed cells were incubated with Streptavidin Alexa Fluor 594 conjugate (Invitrogen) for 1.5 h at RT,

followed by incubation with primary antibodies overnight at 4°C. Secondary antibodies were incubated

for 2 h at RT.

For VASP localization analysis, 2000 LA-EGFP DAOY cells were seeded in a 384 well plate (781090,

Greiner Bio-One) coated overnight with 0.07 µg/µl collagen I (5005-B, Cell systems) diluted in 70%

EtOH. After seeding, the cells were starved for 16 h and treated with 100 ng/ml bFGF for 15 min. Cells

were fixed and incubated with primary and secondary antibodies as described above.

Imaging acquisition was performed using an SP8 confocal microscope (Leica). The list of the primary

and secondary antibodies used is provided in Supplementary material (Table 1).

Lentivirus Production and Transduction

Lentiviral particles were produced by transfection of HEK-293T cells with transfer plasmids, psPAX2

(#12260, Addgene) and pCMV-VSV-G plasmids (#8454, Addgene)[79] in a ratio of 5:3:2 using

polyethylenimine (24765-2, Polysciences). Virus supernatant was harvested 30 h after transfection,

filtered, and concentrated using Amicon Ultra 15 ml Centrifugal Filters (Millipore). For the transduction,

human MB cells were incubated for 24 hours with supernatant containing the viral particles and 8

µg/ml polybrene. Two days after transduction, cells were selected with puromycin (DAOY: 2 µg/ml, HD-

MBO3: 1 µg/ml).

Generation of CRISPR/Cas9-mediated knockout cells

Gene-specific single-guide RNA (sgRNAs) were designed using the online design tool at

https://design.synthego.com. Only highly specific target sites were selected, and the respective

sequences are listed in Supplementary material (Table 2). Oligonucleotides specific for the target sites

were synthesized with BsmBI restriction site overhangs by Microsynth (Balgach, Switzerland) and

436 cloned into the LentiCRISPRv2 transfer plasmid (Addgene plasmid #52961) with a single tube restriction

437 and ligation method as described in[80]. Production of lentiviral vectors and cell transduction was

438 performed as described above. The efficiency of the knockouts was tested by immunoblot.

RNA interference

DAOY cells at approximately 70% confluency were transfected with DharmaFECT Transfection Reagent

4 (T-2004-03, Dharmacon) or Lipofectamine RNAiMAX Transfection Reagent (13778075, Thermo

Fisher Scientific) with 10 nM siRNAs following the manufacturer's instruction. The list of the siRNA used

is provided in Supplementary material (Table 3). Non-Targeting Control siRNA (siCTRL, D-001210-02-

05, Dharmacon) was used as a negative control for siRNA transfection. After 48 or 72 hours, RNA and

proteins were isolated from the cells to determine gene expression by qRT-PCR and to evaluate protein

expression by IB. For SIA, the cells were re-seeded to form spheroids in ultra-low adhesion plates 24 h

after the transfection.

RNA extraction and quantitative Real-Time PCR (qRT-PCR)

Total RNA was isolated from cells with the RNeasy mini kit (74106, Qiagen). cDNA was obtained by

retro-transcription of 1 µg total RNA using High-capacity cDNA Reverse Transcription Kit (4368813,

Applied Biosystems). qRT-PCR was performed using PowerUp Syber Green (A25776, Thermo

Scientific) under conditions optimized for the 7900HT Fast Real-Time PCR System (Applied

Biosystems). The primers used were synthesized by Microsynth AG (Balgach, Switzerland). Primer

sequences are detailed in Supplementary material (Table 4). The relative expression levels of each

gene of interest were calculated using the $2^{-\Delta\Delta Ct}$ method normalized to 18s.

Spheroid invasion assay (SIA)

SIA was performed and quantified according to[42]. In brief: 2500 cells/100 µl per well were seeded in

cell-repellent 96 well microplates (650970, Greiner Bio-One, Kremsmünster, Austria). The cells were

incubated at 37°C for 24 h to form spheroids. 70 µl medium was removed and replaced with 70 µl of a

solution containing 2.5% bovine collagen I (CellSystems, Troisdorf, Germany). Polymerized collagen I

hydrogels were overlaid with 100 µl of serum-free media containing growth factors and/or inhibitors 2x

concentrated. The cells were allowed to invade the collagen matrix for 24 h, after which they were

stained with Hoechst (1:2000, B2883, Sigma Aldrich) for 3-4 h. Images were acquired with an

AxioObserver 2 mot plus fluorescence microscope (Zeiss, Munich, Germany) or with an Operetta high-

content imaging system (PerkinElmer) at 5x magnification. When acquired with the Operetta microscope, the spheroids were localized in the well by acquiring 3 z-planes with 250 μm z-spacing, re-scanned by acquiring 16 z-planes with 25 μm z-spacing and a maximum intensity projection was computed and used for further analysis. Cell invasion was determined as the sum of the distance invaded by the cells from the center of the spheroid as quantified using an automated cell dissemination counter (aCDc)[42] or using Harmony 4.5 software (PerkinElmer).

Colony formation assay

DAOY (400 cells/well) or HD-MBO3 (600 cells/well) cells were seeded in a 6-well plate in complete growth medium containing 10% FBS. The cells were allowed to adhere for 48 h, and the medium was replaced with fresh growth medium containing 1% FBS ± GF. The cells were cultured at 37°C for 10 days (DAOY) or 15 days (HD-MBO3) with medium changes every third or fourth day. The colonies were fixed in methanol for 10 min at -20°C and then stained with 0.5% crystal violet for 15 min. The number and the area of the colonies were quantified by ImageJ using the plugin colony area.

CellTox green cytotoxicity assay

Cytotoxicity was determined using the CellTox™ Green Cytotoxicity Assay (G8742, Promega, Dübendorf, Switzerland), according to manufacturer's instructions. 2500 DAOY cells were seeded in 100 μl per well of complete medium in 96-well Corning black wall microplates with clear round bottom (Corning Incorporated, NY, USA). The cells were incubated at 37°C for 24 h to form spheroids. 50 μl of the medium were then removed from each well and replaced with 50 μL of fresh medium containing 1:500 of CellTox™ Green Dye and different drug concentrations. Wells with medium but without cells were used to measure background fluorescence. The fluorescence representing nonviable cell was measured after 24, 48, and 72 h of incubation using Cytation 3 imaging reader (BioTek, Sursee, Switzerland) at Ex/Em 485/520 nm.

Ex vivo organotypic cerebellum slice culture (OCSC)

Cerebellar slices were prepared from P8-10 C57BL/6JRj mouse pups as described in[47]. Slices were kept in culture for 15 days. Medium was changed daily, and cultures monitored for signs of cell death. Tumor spheroids from DAOY or HD-MBO3 LA-EGFP cells were used. For DAOY cells, the cells were re-seeded to form spheroids in ultra-low adhesion plate 24 h after transfection with siRNA and implanted on the slices 48 h after transfection. The slice-spheroid co-cultures were treated with EGF (30 ng/ml) or

bFGF (12.5 ng/ml) where indicated for 2 days (DAOY) or 5 days (HD-MBO3). Following the treatment,

the co-cultures were fixed and stained for calbindin, GFAP, or anti-human nuclei, as described in[47], and

analyzed using IFA. To evaluate the extent of proliferation both in the tumor cells and in the slice culture,

we performed 5-ethynyl-2′-deoxyuridine (EdU) staining using the Click-iT EdU Alexa Fluor 647 imaging

kit (C10340, Invitrogen). Four-color image acquisition was performed on an SP8 confocal microscope

(Leica). The perimeter and the area of the infiltrating spheroids, and the number of EdU-positive nuclei

were measured using ImageJ.

Patient gene expression and proteomic data

Gene expression and proteomic data of pediatric brain tumor patients were obtained from the ProTrack

data portal (http://pbt.cptac-data-view.org/) from the Clinical Proteomic Tumor Analysis Consortium

(CPTAC) and the Children's Brain Tumor Tissue Consortium (CBTTC). The dataset used includes a

cohort of 218 tumor samples representing seven distinct histological diagnoses, including

medulloblastoma (MB), low-grade glioma (LGG), high-grade glioma (HGG), ependymoma (EP),

craniopharyngioma (CP), ganglioglioma (GG), and atypical teratoid rhabdoid tumor (ATRT)[41].

Protein kinase activity profiling

4.5×10^5 DAOY cells were transfected with siRNA as described above. 72 h after the transfection and

O/N starvation, cells were stimulated with 100 ng/ml bFGF for 15 minutes and lysed in M-PER protein

extraction reagent (Thermo Fischer Scientific) supplemented with Halt Phosphatase Inhibitor Cocktail

and Halt Protease Inhibitor Cocktail EDTA free (Thermo Fischer Scientific). Lysates were cleared by

centrifugation, and protein concentrations were determined by Pierce BCA Protein Assay Kit (Thermo

Fisher Scientific). Aliquots of 1 mg/ml of lysates were prepared and snap-frozen in liquid nitrogen. Kinase

activity profiles were performed using the PamChip® serine/threonine (STK) and protein tyrosine (PTK)

peptide microarray system from PamGene ('s-Hertogenbosch, The Netherlands), according to standard

manufacturer's instructions. The PamChip® consists of 4 identical peptide arrays, each array containing

144 (STK) or 196 (PTK) peptides. The technology measures the phosphorylation of the arrayed STK

and PTK peptides and compares derived peptide phosphorylation signatures to known consensus

signatures of kinases to predict kinase activities. Three PamChip® microarrays were processed at the

same time on the PamStation®12 (PamGene). One and five µg of protein extract were used for the STK

array and the PTK array protocol, respectively, and the enzymatic assay was started by adding ATP

(final concentration 100 μM). For the PTK array, fluorescein isothiocyanate (FITC)–labeled pY20

antibody was incubated on the chip, and the phosphorylation of the individual Tyr peptides was followed

by fluorescence detection in real-time. For the STK array, the primary antibody mixture was incubated

with the chip, and the secondary FITC- conjugated antibody was used to quantify the signal. For each

condition, three biological replicates were performed and analyzed in three different runs.

Software and data analysis

The fluorescent signal intensity for each peptide on the PamChip® was quantitated by BioNavigator

software (PamGene) and represented as linear regression slope as a function of exposure time. A local

background correction was used per spot. Nominal CV (Coefficient of variation) was calculated per

peptide with a 2-component error fit model, using the overall mean as input. Only peptides that showed

nominal CV <0.5 were included in the analysis. Additionally, Combat correction/normalization was

performed to correct for differences in batch effects. After averaging signal intensities across the 3

biological replicates per condition, the signal ratio between siTarget and siCTRL was used to calculate

fold change (FC) values. Peptides with a fold change of at least 35% (i.e., FC > 1.35 or FC < 0.65) were

considered differentially phosphorylated. To generate the peptide phosphorylation heatmap and

compare the global phosphorylation levels, the linear regression slope of each peptide was $\log_2$

transformed. Peptides differently phosphorylated ($p < 0.1$) in siTarget vs. siCTRL were identified by

ANOVA and postHoc Dunnett's tests. Venn diagram analyses were performed in Venny 2.1

(https://bioinfogp.cnb.csic.es/tools/venny/). An upstream kinase prediction tool based on the PamApp

on BioNavigator Analysis software was used to generate a putative list of kinases responsible for

phosphorylating the phosphosites on the PamChip. The prediction of the differentially activated

upstream kinases for each experimental group (siTarget vs. siCTRL, or bFGF vs. UT) is carried out by

a comparison of the phosphorylation sites of the peptides on the array with databases containing

experimental and literature-based protein modifications such as HPRD, PhosphoELM

(http://phospho.elm.eu.org), PhosphoSite Plus (https://www.phosphosite.org), Reactome

(http://reactome.org), UNIPROT (https://www.uniprot.org/) and in silico computational predictions

database such as PhosphoNet (http://www.phosphonet.ca/). The prediction tool functional rank-orders

the top kinases differentially activated between two compared groups. The mean kinase statistic

represents the difference in the predicted protein kinase activity between the two compared groups. A

positive value indicates an activation, while a negative value an inhibition. The specificity score indicates

the specificity of the normalized kinase statistics with respect to the number of peptides used for predicting the corresponding kinase. The higher the specificity score, the higher is the power of the prediction. Values of the specificity score >0.9 were considered as statistically relevant. To group the kinases into sequence families, phylogenetic trees were created using the web-based kinome tool CORAL (http://phanstiel-lab.med.unc.edu/CORAL/).

Statistical analysis

Statistical analyses were performed using GraphPad Prism 8 software (San Diego, California). Statistical significance of differences between groups was determined using Student's *t*-test (unpaired, two-tailed) or one-way ANOVA repeated-measures test with Bonferroni's correction. The experiments were performed in at least three independent biological replicates (n=3) unless differently indicated, and results are shown as mean ± standard deviation (SD) or standard error of the mean (SEM). The results were considered significant when $p < 0.05$ (*$p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$). Where indicated, asterisks show statistical significances between control and test sample.

References

1. Hatten, M. E. & Roussel, M. F. Development and cancer of the cerebellum. *Trends Neurosci.* **34**, 134–142 (2011).

2. Cavalli, F. M. G. *et al.* Intertumoral Heterogeneity within Medulloblastoma Subgroups. *Cancer Cell* **31**, 737--754.e6 (2017).

3. Schwalbe, E. C. *et al.* Novel molecular subgroups for clinical classification and outcome prediction in childhood medulloblastoma: a cohort study. *Lancet Oncol.* **18**, 958–971 (2017).

4. Hovestadt, V. *et al.* Medulloblastomics revisited: biological and clinical insights from thousands of patients. *Nat. Rev. Cancer* **20**, 42–56 (2020).

5. Gajjar, A. *et al.* Risk-adapted craniospinal radiotherapy followed by high-dose chemotherapy and stem-cell rescue in children with newly diagnosed medulloblastoma (St Jude Medulloblastoma-96): long-term results from a prospective, multicentre trial. *Lancet Oncol.* **7**, 813–820 (2006).

6. Gerber, N. U. *et al.* Recent developments and current concepts in Medulloblastoma. *Cancer Treat. Rev.* **40**, 356–365 (2014).

7.   Rieken, S. *et al.* Outcome and prognostic factors of radiation therapy for medulloblastoma. *Int.*
J. Radiat. Oncol. Biol. Phys. **81**, 7–13 (2011).

8.   Wu, X. *et al.* Clonal selection drives genetic divergence of metastatic medulloblastoma. **482**,
529–533 (2012).

9.   Santhana Kumar, K. *et al.* TGF-β Determines the Pro-migratory Potential of bFGF Signaling in
Medulloblastoma. *Cell Rep.* **23**, 3798-3812.e8 (2018).

10. Neve, A. *et al.* Crosstalk between SHH and FGFR signaling pathways controls tissue invasion in
medulloblastoma. *Cancers (Basel).* **11**, (2019).

11. Juraschka, K. & Taylor, M. D. Medulloblastoma in the age of molecular subgroups: A review. *J.*
Neurosurg. Pediatr. **24**, 353–363 (2019).

12. Friedl, P. & Wolf, K. Tumour-cell invasion and migration: diversity and escape mechanisms. *Nat.*
Rev. Cancer **3**, 362–374 (2003).

13. Lauffenburger, D. A. & Horwitz, A. F. Cell migration: A physically integrated molecular process.
Cell **84**, 359–369 (1996).

14. Ridley, A. J. *et al.* Cell Migration: Integrating Signals from Front to Back. *Science (80-.).* **302**,
1704–1709 (2003).

15. Makrilia, N., Kollias, A., Manolopoulos, L. & Syrigos, K. Cell adhesion molecules: Role and
clinical significance in cancer. *Cancer Invest.* **27**, 1023–1037 (2009).

16. Kumar, K. S. *et al.* The Ser/Thr kinase MAP4K4 drives c-Met-induced motility and invasiveness
in a cell-based model of SHH medulloblastoma. *Springerplus* **4**, 19 (2015).

17. MAP4K4 controlled integrin beta1 activation and c-Met endocytosis are associated with invasive
behavior of medulloblastoma cells. *Oncotarget* (2018).

18. Dan, I., Watanabe, N. M. & Kusumi, A. The Ste20 group kinases as regulators of MAP kinase
cascades. *Trends Cell Biol.* **11**, 220–230 (2001).

19. Su, Y. C., Han, J., Xu, S., Cobb, M. & Skolnik, E. Y. NIK is a new Ste20-related kinase that binds
NCK and MEKK1 and activates the SAPK/JNK cascade via a conserved regulatory domain.
EMBO J. **16**, 1279–1290 (1997).

20.   Tesz, G. J. *et al.* Tumor Necrosis Factor α (TNFα) Stimulates Map4k4 Expression through TNFα
Receptor 1 Signaling to c-Jun and Activating Transcription Factor 2. *J. Biol. Chem.* **282**, 19302–
19312 (2007).

21.   Tripolitsioti, D., Grotzer, M. A. & Baumgartner, M. The Ser/Thr Kinase MAP4K4 Controls Pro-
Metastatic Cell Functions. *J. Carcinog. Mutagen.* **08**, 1–8 (2017).

22.   Roth Flach, R. J. *et al.* Endothelial protein kinase MAP4K4 promotes vascular inflammation and
atherosclerosis. *Nat. Commun.* **6**, (2015).

23.   Danai, L. V *et al.* Inducible Deletion of Protein Kinase Map4k4 in Obese Mice Improves Insulin
Sensitivity in Liver and Adipose Tissues. *Mol. Cell. Biol.* **35**, 2356–2365 (2015).

24.   Collins, C. S. *et al.* A small interfering RNA screen for modulators of tumor cell motility identifies
MAP4K4 as a promigratory kinase. *Proc. Natl. Acad. Sci.* **103**, 3775–3780 (2006).

25.   Prolo, L. M. *et al.* Targeted genomic CRISPR-Cas9 screen identifies MAP4K4 as essential for
glioblastoma invasion. *Sci. Rep.* 1–12 (2019) doi:10.1038/s41598-019-50160-w.

26.   Vitorino, P. *et al.* MAP4K4 regulates integrin-FERM binding to control endothelial cell motility.
Nature **519**, 425–430 (2015).

27.   Baumgartner, M. *et al.* The Nck-interacting kinase phosphorylates ERM proteins for formation of
lamellipodium by growth factors. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 13391–13396 (2006).

28.   LeClaire, L. L., Rana, M., Baumgartner, M. & Barber, D. L. The Nck-interacting kinase NIK
increases Arp2/3 complex activity by phosphorylating the Arp2 subunit. *J. Cell Biol.* **208**, 161–
170 (2015).

29.   Goudreault, M. *et al.* A PP2A Phosphatase High Density Interaction Network Identifies a Novel
Striatin-interacting Phosphatase and Kinase Complex Linked to the Cerebral Cavernous
Malformation 3. *Mol. Cell. proteomics* **8**, 157–171 (2009).

30.   Chen, C. *et al.* Striatins Contain a Noncanonical Coiled Coil That Binds Protein Phosphatase 2A
A Subunit to Form a 2 : 2 Heterotetrameric Core of Striatin-interacting Phosphatase and Kinase
(STRIPAK) Complex *. *J. Biol. Chem.* **289**, 9651–9661 (2014).

31.   Kück, U., Radchenko, D. & Teichert, I. STRIPAK , a highly conserved signaling complex , controls
multiple eukaryotic cellular and developmental processes and is linked with human diseases

Discovery of striatin. *Biol. Chem.* **400**, 1–18 (2019).

32.   Kim, D. I. *et al.* An improved smaller biotin ligase for BioID proximity labeling. *Mol. Biol. Cell* **27**,

1188–1196 (2016).

33.   Kim, D. I. *et al.* Probing nuclear pore complex architecture with proximity-dependent biotinylation.

Proc. Natl. Acad. Sci. 2453–2461 (2014) doi:10.1073/pnas.1406459111.

34.   Roux, K. J., Kim, D. I., Raida, M. & Burke, B. A promiscuous biotin ligase fusion protein identifies

proximal and interacting proteins in mammalian cells. *J. Cell Biol.* **196**, 801–810 (2012).

35.   Couzens, A. L. *et al.* Protein interaction network of the mammalian hippo pathway reveals

mechanisms of kinase-phosphatase interactions. *Sci. Signal.* **6**, (2013).

36.   Seo, G. *et al.* MAP4K Interactome Reveals STRN4 as a Key STRIPAK Complex Component in

Hippo Pathway Regulation. *Cell Rep.* **32**, (2020).

37.   Kim, J. W. *et al.* STRIPAK directs PP2A activity toward MAP4K4 to promote oncogenic

transformation of human cells. *Elife* **9**, 1–27 (2020).

38.   Chen, R., Xie, R., Meng, Z., Ma, S. & Guan, K. L. STRIPAK integrates upstream signals to initiate

the Hippo kinase cascade. *Nat. Cell Biol.* **21**, (2019).

39.   Hyodo, T. *et al.* Misshapen-like kinase 1 (MINK1) is a novel component of striatin-interacting

phosphatase and kinase (STRIPAK) and is required for the completion of cytokinesis. *J. Biol.*

Chem. **287**, 25019–25029 (2012).

40.   Tang, Y. *et al.* Architecture, substructures, and dynamic assembly of STRIPAK complexes in

Hippo signaling. *Cell Discov.* **5**, (2019).

41.   Petralia, F. *et al.* Integrated Proteogenomic Characterization across Major Histological Types of

Pediatric Brain Cancer. *Cell* **183**, 1962--1985.e31 (2020).

42.   Kumar, K. S. *et al.* Computer-assisted quantification of motile and invasive capabilities of cancer

cells. *Sci. Rep.* **5**, 15338 (2015).

43.   Gordon, J. *et al.* Protein phosphatase 2a (PP2A) binds within the oligomerization domain of

striatin and regulates the phosphorylation and activation of the mammalian Ste20-Like kinase

Mst3. *BMC Biochem.* **12**, 1–18 (2011).

44.   Zheng, Y. *et al.* Homeostatic Control of Hpo/MST Kinase Activity through Autophosphorylation-

Dependent Recruitment of the STRIPAK PP2A Phosphatase Complex. *Cell Rep.* **21**, 3612–3623

(2017).

45.   Ho, W. S. *et al.* PP2A inhibition with LB100 enhances cisplatin cytotoxicity and overcomes

cisplatin resistance in medulloblastoma cells. *Oncotarget* **7**, 12447–12463 (2016).

46.   Yang, D., Okamura, H., Morimoto, H., Teramachi, J. & Haneji, T. Protein phosphatase 2A C α

regulates proliferation , migration , and metastasis of osteosarcoma cells. *Lab. Investig.* **96**,

1050–1062 (2016).

47.   Neve, A., Kumar, K. S., Tripolitsioti, D., Grotzer, M. A. & Baumgartner, M. Investigation of brain

tissue infiltration by medulloblastoma cells in an ex vivo model. *Sci. Rep.* **7**, (2017).

48.   Meng, Z. *et al.* MAP4K family kinases act in parallel to MST1/2 to activate LATS1/2 in the Hippo

pathway. *Nat. Commun.* **6**, 1–13 (2015).

49.   Zheng, Y. *et al.* Identification of Happyhour/MAP4K as Alternative Hpo/Mst-like Kinases in the

Hippo Kinase Cascade. *Dev. Cell* **34**, 642–655 (2015).

50.   Tang, Y. *et al.* Selective Inhibition of STRN3-Containing PP2A Phosphatase Restores Hippo

Tumor-Suppressor Activity in Gastric Cancer. *Cancer Cell* 1–14 (2020)

doi:10.1016/j.ccell.2020.05.019.

51.   Ribeiro, P. S. *et al.* Combined Functional Genomic and Proteomic Approaches Identify a PP2A

Complex as a Negative Regulator of Hippo Signaling. *Mol. Cell* **39**, 521–534 (2010).

52.   Gschwendt, M. *et al.* Rottlerin, a novel protein kinase inhibitor. *Biochemical and Biophysical*

Research Communications vol. 199 93–98 (1994).

53.   Cywin, C. L. *et al.* Discovery of potent and selective PKC-θ inhibitors. *Bioorganic Med. Chem.*

Lett. **17**, 225–230 (2007).

54.   Mohammadi, M. *et al.* A tyrosine-phosphorylated carboxy-terminal peptide of the fibroblast

growth factor receptor (Flg) is a binding site for the SH2 domain of phospholipase C-gamma 1.

Mol. Cell. Biol. **11**, 5068–5078 (1991).

55.   Chen, R. *et al.* Regulation of Akt/PKB Activation by Tyrosine Phosphorylation. *J. Biol. Chem.*

276, 31858–31862 (2001).

56. Skoble, J., Auerbuch, V., Goley, E. D., Welch, M. D. & Portnoy, D. A. Pivotal role of VASP in
Arp2/3 complex-mediated actin nucleation, actin branch-formation, and Listeria monocytogenes
motility. *J. Cell Biol.* **155**, 89–100 (2001).

57. Barzik, M. *et al.* Ena/VASP proteins enhance actin polymerization in the presence of barbed end
capping proteins. *J. Biol. Chem.* **280**, 28653–28662 (2005).

58. Benz, P. M. *et al.* Differential VASP phosphorylation controls remodeling of the actin
cytoskeleton. *J. Cell Sci.* **122**, 3954–3965 (2009).

59. Harbeck, B., Hüttelmaier, S., Schlüter, K., Jockusch, B. M. & Illenberger, S. Phosphorylation of
the vasodilator-stimulated phosphoprotein regulates its interaction witn actin. *J. Biol. Chem.* **275**,
30817–30825 (2000).

60. Lu, J. *et al.* MEKK2 and MEKK3 orchestrate multiple signals to regulate Hippo pathway. *J. Biol.*
Chem. **95**, 106408 (2021).

61. Garg, R. *et al.* Protein kinase C and cancer: What we know and what we do not. *Oncogene* **33**,
5225–5237 (2014).

62. Du, Q. Y. *et al.* High STRN Expression Promotes HCC Invasion and Migration but Not Cell
Proliferation or Apoptosis through Facilitating Epithelial-Mesenchymal Transition. *Biomed Res.*
Int. **2020**, (2020).

63. Wong, M. *et al.* Silencing of STRN4 suppresses the malignant characteristics of cancer cells.
Cancer Sci. **105**, 1526–1532 (2014).

64. Lee, T. H. *et al.* Rottlerin, a natural polyphenol compound, inhibits upregulation of matrix
metalloproteinase-9 and brain astrocytic migration by reducing PKC-δ-dependent ROS signal.
J. Neuroinflammation **17**, 1–13 (2020).

65. Sarkar, S. & Yong, V. W. Reduction of protein kinase C delta attenuates tenascin-C stimulated
glioma invasion in three-dimensional matrix. *Carcinogenesis* **31**, 311–317 (2010).

66. Brenner, W. *et al.* Migration of renal carcinoma cells is dependent on protein kinase Cδ via β1
integrin and focal adhesion kinase. *Int. J. Oncol.* **32**, 1125–1131 (2008).

67. Cannon, J. L. *et al.* PKCθ regulates T cell motility via ezrin-radixin-moesin localization to the
uropod. *PLoS One* **8**, (2013).

68.   Madaro, L. *et al.* PKCθ signaling is required for myoblast fusion by regulating the expression of

caveolin-3 and β1D integrin upstream focal adhesion kinase. *Mol. Biol. Cell* **22**, 1409–1419

(2011).

69.   Döppler, H. & Storz, P. Regulation of VASP by phosphorylation Consequences for cell migration.

Cell Adhes. Migr. **7**, 482–486 (2013).

70.   Hasegawa, Y., Murph, M., Yu, S., Tigyi, G. & Mills, G. B. Lysophosphatidic acid (LPA)-induced

vasodilator-stimulated phosphoprotein mediates lamellipodia formation to initiate motility in PC-

3 prostate cancer cells. *Mol. Oncol.* **2**, 54–69 (2008).

71.   Heckman, C. A. & Plummer, H. K. Filopodia as sensors. *Cell. Signal.* **25**, 2298–2311 (2013).

72.   Jacquemet, G., Hamidi, H. & Ivaska, J. Filopodia in cell adhesion, 3D migration and cancer cell

invasion. *Curr. Opin. Cell Biol.* **36**, 23–31 (2015).

73.   Steketee, M. B. & Tosney, K. W. Three functionally distinct adhesions in filopodia: Shaft

adhesions control lamellar extension. *J. Neurosci.* **22**, 8071–8083 (2002).

74.   Madsen, C. D. *et al.* STRIPAK components determine mode of cancer cell migration and

metastasis. *Nat. Cell Biol.* **17**, 68–80 (2015).

75.   Schönholzer, M. T. *et al.* Real-time sensing of MAPK signaling in medulloblastoma cells reveals

cellular evasion mechanism counteracting dasatinib blockade of ERK activation during invasion.

Neoplasia **22**, 470–483 (2020).

76.   Ndubaku, C. O. *et al.* Structure-Based Design of GNE-495, a Potent and Selective MAP4K4

Inhibitor with Efficacy in Retinal Angiogenesis. *ACS Med. Chem. Lett.* **6**, 913–918 (2015).

77.   Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-level

datasets. *Nat. Commun.* **10**, (2019).

78.   Ma, M. & Baumgartner, M. Intracellular Theileria annulata Promote Invasive Cell Motility through

Kinase Regulation of the Host Actin Cytoskeleton. *PLoS Pathog.* **10**, e1004003--4 (2014).

79.   Stewart, S. A. *et al.* Lentivirus-delivered stable gene silencing by RNAi in primary cells. *Rna* **9**,

493–501 (2003).

80.   McComb, S. *et al.* Activation of concurrent apoptosis and necroptosis by SMAC mimetics for the

treatment of refractory and relapsed ALL. *Sci. Transl. Med.* **8**, (2016).

81.   Jeong, B. *et al.* Cryo-EM structure of the Hippo signaling integrator human STRIPAK. *Nat. Struct.*

Mol. Biol. 1–10 (2021) doi:10.1038/s41594-021-00564-y.

82.   Varnaitė, R. & MacNeill, S. A. Meet the neighbors: Mapping local protein interactomes by

proximity-dependent labeling with BioID. *Proteomics* **16**, 2503–2518 (2016).

Acknowledgments

The authors thank Dr. Savithri Rangarajan (PamGene International) for scientific support and analysis

of PamChip imaging data and Dr. Jens Sobek (Functional Genomics Center Zurich) for assistance with

the experiments performed on PamStation®12. We would like to thank members of the Functional

Genomics Center Zurich for mass spectrometry support. Imaging was performed with equipment

maintained by the Centre for Microscopy and Image Analysis, University of Zurich. This study was

supported by grants from the Swiss National Science foundation (SNF_31003A_165860/1,

SNF_310030_188793) and the Swiss Cancer League (SCL_KLS-3834-02-2016) to MB, the University

of Zürich (Candoc) to J.M. and from the Childhood Cancer Foundation to MG.

Author contributions

761   J.M. contributed to designing the study, planned and conducted all the experiments, prepared the

762   figures, and wrote the manuscript. B.S. generated DAOY cells stably expressing different MAP4K4

763   domains. C.C. provided experimental support for image acquisition and quantification with the Operetta

764   microscope. M.G. helped to draft the study. M.B. designed the study and wrote the manuscript.

Figures

Figure 1: BioID identifies MAP4K4-STRIPAK complex interaction in MB cells. (a) Immunoblot
analysis (IB) of HEK-293T total cell lysate or after immunoprecipitation (IP) with streptavidin-conjugated
beads and detection with HRP-streptavidin. *n*=2 experiments. **(b)** Venn diagram of affinity captured
proteins from HEK-293T cells. Comparison of BioID2-MAP4K4 with BioID2-CTRL is shown. Values of
enriched proteins (p < 0.05, one-way ANOVA) are shown. *n*=3. **(c)** Gene ontology (GO) classifications
of MAP4K4-interacting proteins. **(d)** Volcano plot of interacting proteins in N-BioID2-MAP4K4 (left) or C-
BioID2-MAP4K4 (right) HEK-293T cells compared to BioID2-CTRL. $\log_{10}$ ratios of cumulated peptide
detection values plotted against negative $\log_{10}$ p-values of Student's *t*-test. Enriched proteins are

highlighted in red (p < 0.05 and > 2-fold enrichment) and members of the STRIPAK complex indicated.
(e) Architecture of the STRIPAK complex, adapted from[81]. **(f)** IB detection of BioID2-MAP4K4-FLAG and
components of STRIPAK complex in anti-FLAG-IP in HEK-293T cells. **(g)** IB of FLAG-tagged BioID2
alone or fused to MAP4K4 in DAOY cell lysate or after pull-down with streptavidin-conjugated beads.
(h) Venn diagram of affinity-captured proteins from DAOY BioID2-MAP4K4 cells compared to BioID2-
CTRL. *n*=2. **(I)** IB of MAP4K4-FLAG and components of STRIPAK complex after anti-FLAG-IP in DAOY
cells. **(j)** Top: MAP4K4 domain architecture and MAP4K4 constructs used. Numbers represent the
amino acids corresponding to full-length MAP4K4. KD, kinase domain; ID, intermediate domain; CNH,
citron homology domain; CC, coiled-coil. Lower: Co-IP and IB analysis of STRN4 and STRIP1 in DAOY
cells transduced with lentiviral vectors expressing full-length or truncated MAP4K4-FLAG. Asterisks
indicate relevant bands. **(k)** Quantification of co-IP from **j** (*n*=3, means ± SD). See also Figure S1.
786

Figure 2: STRIPAK complex disruption impairs GF-induced collagen I invasion. (a) Invasion and
representative images of 3D spheroid invasion assay (SIA) of DAOY cells KO for MAP4K4, STRN4,
STRN3, or STRIP1 ± bFGF (100 ng/ml). Invasion is calculated as sum of the distances of each migrating
cell from the center of the spheroids (*n*=3, means ± SD). **(b)** SIA of DAOY cells transfected with 10 nM
of the indicated siRNA ± GFs (EGF: 30 ng/ml, bFGF: 100 ng/ml). *n*=4, means ± SD. **(c)** SIA and
representative images of DAOY cells ± 100 ng/ml bFGF and ± increasing concentrations of PP2A
inhibitor LB-100 (*n*=3, means ± SD). * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001 (one-way
ANOVA). See also Figure S3.
795

Figure 3: Depletion of MAP4K4 and STRN3 blocks tissue invasion in SHH MB cell model. (a)
Timeline of organotypic cerebellum slice culture (OCSC) and tumor spheroid implantation. **(b)** Maximum
intensity projections (MIP) of representative confocal sections of OCSCs implanted with DAOY tumor
cell spheroids ± 30 ng/ml EGF for 48h. Green: Lifeact-EGFP; blue: calbindin (Purkinje cells); red: GFAP.
Middle: Inverted greyscale images of MIP of LA-EGFP channel, lower: Inverted greyscale images of
MIP of EdU-Click-IT staining. Scale bar: 200 µm. **(c)** Quantification of LA-EGFP area of spheroids shown
in **b** (*n*=3, means ± SD). **(d)** Number of EdU-positive nuclei normalized by the area of the tumor
spheroids (*n*=3, means ± SD). * p < 0.05, ** p < 0.01, *** p < 0.001 (unpaired Student's *t*-test). See also
Figure S4.
Figure 4: STRN3 is necessary for invasive growth of group 3 MB cell model in cerebellar tissue.
(a) Organotypic cerebellum slice culture implanted with tumor spheroids derived from HD-MBO3 cells
KO for MAP4K4, STRN4, STRN3, or STRIP1. MIP of representative slices five days after implantation
and ± treatment with 30 ng/ml EGF or 12.5 ng/ml bFGF. Green: Lifeact-EGFP; blue: calbindin (Purkinje
cells); red: GFAP; yellow: Edu-Click-IT. Scale bar: 200 µm. **(b)** Areas of LA-EGFP positive spheroids
showed in **a**. *n*=3, means ± SD. **(c)** Number of EdU-positive nuclei normalized by the area in pixels of
the tumor spheroids (*n*=3, means ± SD). * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001 (unpaired
Student's *t*-test). See also Figure S5.
813

**Figure 5: STRN4/3 and MAP4K4 display opposing functions towards MB cell proliferation, clonal
growth and YAP/TAZ target gene expression. (a)** Colony formation assay of DAOY cells KO for
MAP4K4, STRN4, STRN3, or STRIP1 ± EGF or bFGF treatment (EGF: 30 ng/ml, bFGF: 100 ng/ml).
Representative images (left) and quantification (right, *n*=3, means ± SD). **(b)** Colony formation assay
using HD-MBO3 cells with treatments as in a. (*n*=3, means ± SD). **(c)** qRT-PCR analysis of YAP/TAZ
target genes in DAOY cells 48 h after transfection with siRNAs as indicated. *n*=4, means ± SD. **(d)** qRT-
PCR analysis of YAP/TAZ target genes in HD-MBO3 with CRISPR/Cas9-mediated knockout of
MAP4K4, STRN4, STRN3, or STRIP1 (*n*=3, means ± SD). Statistical analyses in **a** and **b** were
performed by one-way ANOVA, in **c** and **d** by unpaired Student's *t*-test. * p < 0.05, ** p < 0.01, *** p <
0.001, **** p < 0.0001. See also Figure S6.
824

Figure 6: Novel protein kinases C are pro-migratory effectors of MAP4K4 and STRN3. (a) Kinase
activity prediction analysis in DAOY cells transfected with the indicated siRNAs and ± stimulation with
100 ng/ml bFGF for 15 min. The plots show predicted top 25 differentially activated serine/threonine
kinases (STK). The x-axis indicates differences in the activity of the predicted protein kinase between
the two groups, with effect size (values) and direction (>0=activation; <0=inhibition). The color of the
bars represents the specificity score (darker to light is higher to lower score). Values >0.9 were

considered as statistically relevant. First graph from left: comparison of bFGF stimulated siCTRL cells
vs. untreated (UT); all other graphs: comparison of the indicated siTarget vs. siCTRL with bFGF
stimulation. Highlighted in red are novel PKC isoforms (δ, ε, η and θ) ($n=3$). **(b-e)** Quantification and
representative images of SIA with DAOY cells stimulated with 100 ng/ml bFGF and treated with
increasing concentrations of PKC inhibitors. Invasion was quantified as the sum of invasion distances
from the center of the spheroids. **b:** Rottlerin (PKCδ inhibitor, $n=4$, means ± SD). **c:** PKCθ inhibitor ($n=4$,
means ± SD). **d:** Combination of Rottlerin and PKCθ inhibitor ($n=3$, means ± SD). **e:** Bisindolylmaleimide
I (BIM, PKCα/β/γ inhibitor, $n=4$, means ± SD). * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$ (one-way ANOVA).
See also Figure S7.
840

Figure 7: MAP4K4 and STRN3 mediate VASP$_{S157}$ phosphorylation. (a) Heat map of increased (red)
and decreased (blue) phosphorylation of Ser/Thr (STK, left) or Tyr (PTK, right) consensus peptides
associated with lysates from siRNA transfected DAOY cells treated ± 100 ng/ml bFGF for 15 min. Log$_2$
of the fold change between siTarget and siCTRL ($n=3$). **(b)** Venn diagram displaying the number of
Ser/Thr and Tyr peptides with significantly changed phosphorylation ($p < 0.1$) in each siTarget versus
siCTRL with bFGF stimulation ($n=3$). **(c)** Volcano plots representing the changes in phosphorylation of
STK (circles) and PTK (diamonds) peptides in siMAP4K4 (left) or siSTRN3 (right) versus siCTRL with
bFGF stimulation. Blue: peptides with significantly decreased (fold change < 1.35), red: peptides with
significantly increased phosphorylation (fold change > 1.35). ($n=3$). See also Figure S8 and Table S3.
(d) Single confocal section of VASP localization in DAOY cells seeded on collagen-coated plates ± 100
ng/ml bFGF for 15 min. 3X magnifications of boxed areas are shown. Red: VASP, Green: Lifeact-EGFP.
Blue: DNA (Hoechst). Scale bar: 20 μm. **(e)** Maximum intensity projection of LA-EGFP fluorescence of
tissue-invading HD-MBO3 CTRL or KO MAP4K4 or STRN3 cells. Scale bar: 10 μm. **(f)** IB and
quantification of VASP$_{S157}$ phosphorylation in DAOY cells 72 h after siRNA transfection ± 100 ng/ml
bFGF for 15 min ($n=3$, means ± SD). **(g-h)** IB and quantification of VASP$_{S157}$ phosphorylation in DAOY
(g) and HD-MBO3 cells **(h)** after treatment for 4 h with increasing concentration of GNE-495 ± 100 ng/ml
bFGF for 15 min. Quantification represents the ratio of p-VASP$_{S157}$ over total VASP relative to untreated
control (**g**: $n=3$, means ± SD. **h:** $n=2$, means ± SD). **(i)** Model for MAP4K4 and STRIPAK regulation of
PKC and p-VASP$_{S157}$ and its putative implication in filopodia formation.