Reprogrammed blastoids contain amnion-like cells but not trophectoderm

Cheng Zhao[1], Alvaro Plaza Reyes[1,2], John Paul Schell[1], Jere Weltner[1], Nicolás M. Ortega[1,11], Yi Zheng[3],

Åsa K. Björklund[4], Janet Rossant[5], Jianping Fu[3,6,7], Sophie Petropoulos*[1,8,9,10] & Fredrik Lanner*[1,11]

1. Department of Clinical Science, Intervention and Technology, Karolinska Institutet, and Division
of Obstetrics and Gynecology, Karolinska. Universitetssjukhuset, Stockholm, Sweden.
2. Department of Regeneration and Cell Therapy, Andalusian Molecular Biology and Regenerative
Medicine Centre (CABIMER), Seville, Spain.
3. Department of Mechanical Engineering, University of Michigan, Ann Arbor, MI 48109, USA.
4. Dept of Cell and Molecular Biology, National Bioinformatics Infrastructure Sweden, Science for
Life Laboratory, Uppsala University, Husargatan 3, SE-752 37 Uppsala, Sweden.
5. Program in Developmental and Stem Cell Biology, Hospital for Sick Children, Toronto, Ontario,
Canada.
6. Department of Cell & Developmental Biology, University of Michigan Medical School, Ann
Arbor, MI 48109, USA.
7. Department of Biomedical Engineering, University of Michigan, Ann Arbor, MI 48109, USA.
8. Département de Médecine, Université de Montréal, Montreal Canada
9. Centre de Recherche du Centre Hospitalier de l'Université de Montréal, Axe Immunopathologie,
H2X 19A Montreal Canada
10. Département de Biochimie et Médecine Moléculaire, Université de Montréal, Montréal Canada.
11. Ming Wai Lau Center for Reparative Medicine, Stockholm Node, Karolinska Institutet,
Stockholm, Sweden.
25

Corresponding author

Abstract

With the advancement of human stem cell cultures and interest in understanding human embryogenesis,

human blastocyst-like structures, or human blastoids, are being developed. Thus far, two different strategies

have been taken, generating blastoids from naive human pluripotent stem cells (hPSC)[1–3] or through

reprogramming of adult human somatic cells[4]. All these studies utilize comparative transcriptome analyses

to authenticate blastoid cells and identify their counterparts in the human blastocyst[5,6]. However, the validity

of comparative transcriptome analysis is critically hinged on including relevant reference data, not only

those from targeted cell types but also from potential alternative cell lineages. Thus, we sought to reevaluate

the single-cell transcriptome data from the blastoids based on a more comprehensive cellular reference,

which includes data from *in vitro* cultured human[6,7] and non-human primate (NHP, *Cynomolgus macaque*)

blastocysts[8], a human stem cell-based post-implantation amniotic sac embryoid (PASE) model[9], and an *in

vivo* gastrulation-stage human embryo specimen[10], using four different analysis strategies. Our analyses

unequivocally support that blastoids developed by reprogramming adult human somatic cells largely fail to

generate cells with a transcriptome profile consistent with the human blastocyst trophectoderm. Instead,

cells identified as trophectoderm-like have a transcriptional profile more similar to the amniotic ectoderm

in the gastrulating human and NHP embryos. To facilitate cell lineage identifications in human embryo

models, we further identified a set of human amniotic ectoderm and trophectoderm markers that could be

utilized to distinguish these two lineages. We further built a neural-network based online prediction tool,

which accurately discerns the full cellular composition of blastoids.

Results

Blastoids are promising tools to study early human development, resolving technical, ethical, and legal limitations of natural human embryos[11]. However, the utility of blastoids is critically dependent on their faithful recapitulation of natural human blastocyst development. Human pluripotent stem cells (hPSCs) with primed pluripotency have been suggested to have the capacity to differentiate into the trophectoderm (TE)[12]. More recent studies, however, suggest that the purported TE cells derived from primed hPSCs might instead be mis-annotated extraembryonic mesoderm or amniotic ectoderm[13–15]. Similarly, putative TE cells in mouse blastoids generated from extended pluripotent stem cells (EPSCs) are now believed to instead represent mesoderm-like cells[11]. In light of these conflicting views and a critical need for accurate annotations of cell lineages in embryoids, we sought to reevaluate transcriptome data from all human blastoids[1–4]. To discern cell lineage identities and examine the possible presence of mesoderm- or amniotic ectoderm-like cells in the blastoids, we generated an integrated dataset, combining blastoid transcriptome data with reference data from *in vitro* cultured human blastocysts[3,6,7], a human stem cell-based post-implantation amniotic sac embryoid (PASE) model[9], and a Carnegie Stage (CS) 7 human gastrula[10]. UMAP projection of this integrated data clearly revealed distinct cell clusters, including those annotated as the epiblast, TE, endoderm, amnion, and mesoderm lineages based on known lineage markers (Fig. 1a and Extended Data Fig. 1a-c). Notably, cells annotated as trophoblast-like cells (TLC) in reprogrammed blastoids (Liu-Blastoids) did not contribute to the TE cluster; rather, the majority of TLC in Liu-Blastoids overlaps with the amnion cluster (Fig. 1a-b and Extended Data Fig. 1b). In contrast, all three naive hPSC-derived blastoids contain expected cell types, with most of the annotated epiblast-like cells (ELC), hypoblast-like cells (HLC), and TLC overlapping with their counterparts in the human embryo. Furthermore, inclusion of transcriptome data from naive and primed hPSCs as well as trophoblast stem cells representative of post-implantation human trophoblast lineage[2,9] together with the integrated dataset enabled us to explore developmental stages of blastoid cell lineages (Extended Data Fig. 1a). Among the naive hPSC-derived blastoids, those cultured in the PXGL condition (KAG-Blastoids[2] and YAN-Blastoids[3]) generally comprise cell lineages that align better with those in preimplantation human

blastocysts, whereas blastoids generated from hPSCs cultured in the 5i/L/A condition still contain a notable

portion of cells resembling those in postimplantation human embryos (Yu-Blastoids[1]) (Fig. 1b). For

convenience and further exploration, our transcriptome comparative analyses can be browsed at

(http://petropoulos-lanner-labs.clintec.ki.se/).

A single human dataset containing transcriptome of both amniotic ectoderm and TE is currently lacking.

Thus, we utilized recently published data of cultured non-human primate (NHP, *Cynomolgus macaque*)[8],

which includes transcriptome of both amniotic ectoderm and TE, to perform a second analysis. Integration

of the CS7 human gastrula and NHP data confirmed proper alignments of corresponding cell lineages from

these two species (Extended Data Fig. 1c). Using the NHP data as the reference for comparative

transcriptome analyses of the blastoids clearly shows that the majority of TLC in Liu-Blastoids clusters

with the amnion as opposed to the TE lineage (Fig. 1b), corroborating our finding in Fig. 1b. We should

note that this finding could be reached by common integration methods MNN, Harmony or CCA (Seurat)

(Extended Data Fig. 1d-e).

We next analyzed the proportion of cell types identified in the blastoids using our re-annotations obtained

from the multi-data integration. TE cells should be the dominating cell type in human blastocysts (Extended

Data Fig. 2a-b). In contrast to human blastocysts, all blastoids contain a much greater EPI population

(Extended Data Fig. 2b). It is also notable that in addition to containing a significant portion of amnion-like

cells, Liu-Blastoids also carried substantial numbers of mesoderm-like and unidentified cells (Fig. 1a-b and

Extended Data Fig. 2b-c). Mesoderm-like and unidentified cells can also be found in Yu-Blastoids and to

a lesser degree in KAG-Blastoids (Fig. 1a-b and Extended Data Fig. 2b-c).

Differential gene expression analysis incorporating TE lineages from both pre- and post-implantation

human embryos[6,7] and amnion cells from the CS7 gastrula[10] allowed the identification of 173 and 14 genes

highly enriched in TE lineage and amnion, respectively (Fig. 2a, Extended Data Table1.tsv). We further

validated these gene signatures by examining two additional preimplantation human blastocyst datasets[3,5]

and the NHP data (Fig. 2a and Extended Data Fig. 2d). TLC in Liu-Blastoids showed distinct expression

of the amnion markers (Fig. 2b). Examining normalized expression of selected amnion markers further

showed that TLC in Liu-Blastoids highly expresses amnion markers such as *ISL1*, *GABRP*, and *IGFB5*

(Fig. 2b)[8,10,16], with low expression of TE lineage markers including *GATA2, GCM1,* or *BIN2*. ISL1 and

GABRP are specifically expressed in the amnion but not in the TE lineage of *in vitro* cultured NHP

embryos, at both the transcriptional and protein levels[8]. Furthermore, ISL1 and GABRP are not detectable

in the TE of *in vitro* cultured human blastocysts at both the transcriptional and protein levels[17].

In an effort to compare multiple datasets without the need of integration or single-cell normalization, we

collapsed all the datasets into pseudo-bulk. A PCA plot was generated for the human TE, TSC, and amnion

datasets together with the blastoids based on the 187 differentially expressed genes (DEGs) (Fig. 2c). The

first component, which explains 72.78% of all variations, represents the TE to amnion axis. While TLCs of

the three hPSC-derived blastoids are located on the left side of the plot with the TE datasets, TLC of Liu-

Blastoids are positioned well to the right towards the amnion of the CS7 human gastrula and the amnion-

like cells (AMLC) in the PASE model. To further validate this analysis, we incorporated the NHP data to

generate another PCA plot (Extended Data Fig. 2d-e). This cross-species analysis further supports that TLC

of Liu-Blastoids are more similar to the amnion at the transcriptome level.

Finally, we used neural-network to build an Early Embryogenesis Prediction tool (http://petropoulos-

lanner-labs.clintec.ki.se/) based on data from human blastocysts and the CS7 human gastrula[6,7,10] (Fig. 2d

and Extended Data Fig. 3a-c). The prediction performance of this tool was validated using a large number

of additional datasets[3,5,18], with a specificity of 97.3% and a sensitivity of 88.7% (Extended Data Fig. 3d).

The neural network prediction also supports that Liu-Blastoids lack TE and instead those annotated TLC

in Liu-Blastoids are more similar to the amnion at the transcriptome level (Fig. 2d). In conclusion, to discern

cell lineages in human embryoids including blastoids based on transcriptome data, it is imperative to use

multiple independent analyses together with unbiased, comprehensive reference datasets. To validate cell

identities in human embryoids, expression of key cell fate markers, such as those identified in this work,

should also be clearly demonstrated whenever possible. We envision our analyses, identified amnion and

TE markers and the Early Embryogenesis Prediction tool should be useful for authentication of human

embryoids, thus promoting the healthy development of this emerging field.

Figure. 1 Integration of blastoid cells with an extensive reference dataset, including data from human,

non-human primate and human embryoid references.

a&b, UMAP projection of the eight-dataset integration. Embryonic and blastoid cells are shown in **a** and

b, respectively. The color of each data point represents the cell annotations retrieved or restored for each

publication. The light grey colored data points represent cells used in the multi-data integration which were

not used to infer cell annotations. The shape of data points for the embryonic cells indicates data source.

Cells not assigned to a major lineage from the blastoid datasets were labeled as "Undefined" and colored in

dark grey. **c,** UMAP projection of Mutual Nearest Neighbours (MNN) cross-species integration, including

cells from each blastoid dataset and the *in vitro* non-human primate (NHP) embryo at Day 14, colored by

cell type. The shape of data points indicates species source. Embryonic cells from Yanagida et al., and

naïve, primed and Okae cells from Kawaga et al., were also included in the integration.

Figure. 2 Validation of cell lineage identities in blastoids.

a, Expression of significant differentially expressed genes in embryonic amnion, TE, and trophectoderm-

like cell (TLC) from the four blastoid models. TE including TE cells from Blakeley et al. (TE-1), from

Petropoulos et al. (TE-2), from Yanagida et al. (TE-3), from pre-implantation cells (TE-4), and from post-

implantation cells (TE-5) in Xiang et al. For visualization, cells from the same cell type were randomly

down-sampled to 50. The color shown in the left and middle heatmap represents scaled values of log-

transformed expression in each cell. The heatmap to the right of the main heatmap represents scaled average

log-transformed expression values calculated from TLCs of the blastoid datasets. **b,** Violin plots showing

log-transformed expression of key amnion and TE markers in amnion, TE, amnion-like cells (AMLC), and

TLC from the four blastoids. **c,** Principal components analysis (PCA) of the pseudo-bulk expression profiles

of amnion, TE, AMLC and Tsw-AMLC from the PASE model, Okae TSC cells, and TLCs from the four

blastoids. TE (1-5) in **a** and **b** represent the same source of TE as in **c. d,** Alluvial plot comparing the

original cell-type annotations (epiblast-like cells (ELC), hypoblast-like cells (HLC), and TLC from the four

blastoids) to the predicted identities obtained from the Early Embryogenesis Prediction tool. The

predictions identified as "ambiguous" or "NoSigHits" were removed for visualization.

Acknowledgements

We thank members of the laboratories of F.L. and S.P. for discussions. This work was supported by the

Swedish Research Council (F.L., S.P.) , Ragnar Söderberg Foundation (F.L.), Ming Wai Lau Center for

Reparative Medicine (F.L.), Center for Innovative Medicine (F.L.), Wallenberg Academy Fellow (F.L.),

Swedish Society for Medical Research (S.P.), Emil och Wera Cornells Stiftelse (S.P.), Natural Sciences

and Engineering Research Council of Canada Discovery Grant (S.P), The Canadian Institutes of Health

Research (S.P., J.R.), Päivikki and Sakari Sohlberg Foundation (J.W.), and Sigrid Jusélius Foundation

(J.W.). SP holds the Canada Research Chair Functional Genomics in Reproduction and Development.

Contributions

C.Z. and F.L. conceived the study with advice from A.P-R., J.P.S, and S.P., C.Z., A.P-R., and J.P.S.,

performed the data analyses with help from J.W., S. P., N.M.O. Y.Z., J.R., J.F., Å.B. The manuscript was

written by C.Z., A.P-R, S.P., and F.L., with input from all of the authors.

Corresponding author

Correspondence to fredrik.lanner@ki.se

Ethics declarations

Competing interests

The authors declare no competing interests.

Reference
172
1.   Yu, L. *et al.* Blastocyst-like structures generated from human pluripotent stem cells. *Nature*
(2021) doi:10.1038/s41586-021-03356-y.
2.   Kagawa, H. *et al.* Human blastoids model blastocyst development and implantation. *Nat. 2021* 1–
9 (2021) doi:10.1038/s41586-021-04267-8.
3.   Yanagida, A. *et al.* Naive stem cell blastocyst model captures human embryo lineage segregation.
Cell Stem Cell **28**, 1016-1022.e4 (2021).
4.   Liu, X. *et al.* Modelling human blastocysts by reprogramming fibroblasts into iBlastoids. *Nature*
(2021) doi:10.1038/s41586-021-03372-y.
5.   Blakeley, P. *et al.* Defining the three cell lineages of the human blastocyst by single-cell RNA-
seq. *Development* (2015) doi:10.1242/dev.123547.
6.   Petropoulos, S. *et al.* Single-Cell RNA-Seq Reveals Lineage and X Chromosome Dynamics in
Human Preimplantation Embryos. *Cell* **165**, 1012–1026 (2016).
7.   Xiang, L. *et al.* A developmental landscape of 3D-cultured human pre-gastrulation embryos.
Nature **577**, 537–542 (2020).
8.   Yang, R. *et al.* Amnion signals are essential for mesoderm formation in primates. *Nat. Commun.*
2021 121 **12**, 1–14 (2021).
9.   Zheng, Y. *et al.* Controlled modelling of human epiblast and amnion development using stem
cells. *Nature* **573**, 421–425 (2019).
10.   Tyser, R. C. V. *et al.* Single-cell transcriptomic characterization of a gastrulating human embryo.
Nat. 2021 6007888 **600**, 285–289 (2021).
11.   Posfai, E., Lanner, F., Mulas, C. & Leitch, H. G. All models are wrong, but some are useful:
Establishing standards for stem cell-based embryo models. *Stem Cell Reports* **16**, 1117 (2021).
12.   Xu, R.-H. *et al. BMP4 initiates human embryonic stem cell differentiation to trophoblast. Nature*
biotechnology vol. 20 (2002).
13.   Chhabra, S. & Warmflash, A. BMP-treated human embryonic stem cells transcriptionally
resemble amnion cells in the monkey embryo. *Biol. Open* **10**, (2021).
14.   Bernardo, A. S. *et al.* BRACHYURY and CDX2 mediate BMP-induced differentiation of human
and mouse pluripotent stem cells into embryonic and extraembryonic lineages. *Cell Stem Cell* **9**, 144–155
(2011).
15.   Guo, G. *et al.* Human naive epiblast cells possess unrestricted lineage potential. *Cell Stem Cell*
28, 1040-1056.e6 (2021).
16.   Roost, M. S. *et al.* KeyGenes, a Tool to Probe Tissue Differentiation Using a Human Fetal
Transcriptional Atlas. *Stem Cell Reports* **4**, 1112–1124 (2015).
17.   Zheng, Y. *et al.* Single-Cell Analysis of Embryoids Reveals Lineage Diversification Roadmaps of
Early Human Development. *SSRN Electron. J.* (2021) doi:10.2139/SSRN.3952092.
18.   Zhou, F. *et al.* Reconstituting the transcriptome and DNA methylome landscapes of human
implantation. *Nat. 2019 5727771* **572**, 660–664 (2019).

Integrated analysis including embryonic cells and blastoid models

a

Embryonic dataset

b

Liu et al., 2021(Liu-Blastoids)

Yu et al., 2021(Yu-Blastoids)

Kagawa et al., 2021(KAG-Blastoids)

Yanagida et al., 2021(YAN-Blastoids)

Legend (a):
- Prelineage
- ICM−TE_trans
- ICM
- Epiblast
- Endoderm
- non-embryonic cells
- TE
- Primitive streak
- Amnion
- Mesoderm
- ExE_Mes
- ● Tyser et al., 2021
- ■ Xiang et al., 2020
- ▲ Petropoulos et al., 2016
- ◆ Yanagida et al., 2021

Legend (b):
- ● ELC
- ● HLC
- ● TLC
- ● Undef
- ● additional cells from mutli-data integration

c

Cross-species integration including non-human primate embryos and blastoid models

NHP & Liu-Blastoids

NHP & Yu-Blastoids

NHP & Kagawa et al., 2021(KAG-Blastoids, other cells)

NHP & Yanagida et al., 2021 (YAN-Blastoids, embryonic cells)

Legend (c):
- Amnion(NHP)
- E−Amnion(NHP)
- EPI−AM−trans(NHP)
- Undef
- Epiblast
- ELC
- Naive
- Primed
- Endoderm
- HLC
- ICM−TE_trans
- ICM
- TE
- TLC
- Okae
- PriS
- ExE_Mech
- Mesoderm
- ● Human
- ▲ NHP

Expression of top DEGs between amnion and TE-lineage

a

Expression of key amnion and TE markers

b

PCA of pseudo-bulk data

c

Lineage calling through neural-network prediction

d