A hybrid PDE–ABM model for viral dynamics with application to SARS–CoV–2 and influenza

Sadegh Marzban [1], Renji Han [1], Nóra Juhász [1], and Gergely Röst[1]

[1]Bolyai Institute, University of Szeged, H-6720 Szeged, Hungary

Corresponding Author: Sadegh Marzban
e-mail: sadegh.marzban@math.u-szeged.hu

Abstract

We propose a hybrid partial differential equation – agent-based (PDE–ABM) model to describe the spatio-temporal viral dynamics in a cell population. The virus concentration is considered as a continuous variable and virus movement is modelled by diffusion, while changes in the states of cells (i.e. healthy, infected, dead) are represented by a stochastic agent-based model. The two subsystems are intertwined: the probability of an agent getting infected in the ABM depends on the local viral concentration, and the source term of viral production in the PDE is determined by the cells that are infected.

We develop a computational tool that allows us to study the hybrid system and the generated spatial patterns in detail. We systematically compare the outputs with a classical ODE system of viral dynamics, and find that the ODE model is a good approximation only if the diffusion coefficient is large.

We demonstrate that the model is able to predict SARS–CoV–2 infection dynamics, and replicate the output of in vitro experiments. Applying the model to influenza as well, we can gain insight into why the outcomes of these two infections are different.

Keywords: hybrid model, spatiotemporal dynamics, agent based model, viral kinetics, virus diffusion, infection, SARS–CoV–2, influenza

1. Introduction

A widely used concept to mathematically describe in-host viral dynamics is to model the main quantities by means of an ODE system. While this method proves to be extremely useful in many applications, the richness of information it can provide is naturally limited since it uses time as its single variable – in several research areas it is necessary to extend the cornerstone ideas of this model in order to grasp and investigate important spatial aspects as well. In the case of modelling the in-host spread of COVID-19 and influenza, we can benefit from taking into account the major size difference between viruses and cells of the body: as a result we have developed a hybrid partial differential equations – agent based model (PDE–ABM) spatiotemporal system, where virus concentration is

modeled as a continuous variable, while for the case of epithelial cells we take a discrete spatial approach. Our implementation of the latter model is based on a free and open source software package, HAL (Hybrid Automata Library). The present article has two main focuses: firstly, we explore the similarities and differences between the classical ODE model and the hybrid PDE-ABM model by observing the temporal evolution of cells and viral concentration in the respective systems. Secondly, we investigate the results that are found beyond the limitations of the ODE model, i.e. we explore conclusions that can be drawn exclusively from the more complex spatiotemporal PDE-ABM model. Finally, for completeness we include well-posedness theorems regarding the two main systems.

Mathematical models are powerful tools in the hands of research teams across the globe since the appearance of the COVID-19 disease caused by the novel SARS-CoV-2 coronavirus. In the currently still ongoing and virtually unprecedented pandemic circumstances these mathematical models are invaluable as they are able to optimize high-impact decisions and make successful predictions based on computer simulations alone. This means that their results are obtained at an ideally low cost in a complex situation where real-life and real-time experiments to obtain these same results would be either extremely risky or simply not possible. A further advantage of these models is their wide range of applicability: mathematical models can be used in large-scale problems as for example forecasting the effect of mass vaccination programs or a nation-wise lockdown, but they can equally be applied for small-scale phenomena such as analysing a drug's mechanism of action or the viral transmission between individual cells. The present article focuses on the latter: we use a hybrid mathematical approach to study the in-host spread of SARS-CoV-2 infection on a cellular level.

Throughout the investigation of the above mentioned hybrid model we need a simple and already comprehensively understood model we can use as a reference system – for this purpose we will consider a specific version of an ODE prototype model, the May-Nowak system (for an extensive overview see [5]; while for concrete applications we refer to [3], [4] and [16]). This classical model is given by

$$\begin{cases} H_t = -\beta HV, & t > 0, \\ I_t = \beta HV - \delta I, & t > 0, \\ V_t = pI - cV, & t > 0, \end{cases}$$

where the $H = H(t), I = I(t)$ and $V = V(t)$ functions respectively correspond to the number of healthy cells, infected cells and virus particles, and they depend on time only. This model plays a crucial part in our article in several ways: *a)* in the development process of the more sophisticated hybrid model we use the ODE model to compare and match the respective solutions, *b)* we exploit the fact that there is more information available on the exact parameters of this classical model and use their converted version in the hybrid system, and finally, *c)* we can demonstrate both the advantages and challenges offered by the relatively new hybrid system in comparison to a well-known model.

The above ODE model can not capture the effects of spatial spread, and hence it undoubtedly ignores any occurring spatial phenomenon. In several biological applications

we simply can not afford to leave out such an important factor from the main model: the most obvious example for this is virus diffusion itself, which makes it possible for viruses to physically reach susceptible cells in the first place. One of the most powerful tools that does capture these crucial spatial mechanisms is the rich analytical theory of partial differential equations – PDE models are often used to investigate complex systems in viral infection dynamics, see [1, 2, 3] just to mention a few examples.

Agent-based modeling is another (and in fact quite different) approach to describe complex space-time dynamics in viral spread. The cornerstone idea of this method is to define a discrete heterogeneous state space where the so-called agents or elements have collective interactions with each other and they change their states correspondingly [6, 7]. The ABM concept has several important applications within the field of virus dynamics; in [10] the authors have used agent-based models to evaluate COVID-19 transmission risks, [9] focuses on the spatiotemporal dynamics of influenza, while [8] approaches the topic of viral infections from another viewpoint, here the authors investigate interactions between innate and adaptive immunity.

As seen above, different frameworks – including agent-based models and PDE systems – have been applied to construct viral dynamics models. An agent-based model is a great tool to include randomness and natural variability into the system, while PDEs are faster to evaluate numerically: by forming bridges between these two modeling strategies we arrive to a hybrid system, the so-called hybrid PDE-ABM model, which unites the different advantages of these methods (see [11]). Hybrid PDE-agent-based models have been under intense investigation in several fields of biology. Butner et al. [12] presented a hybrid model to study the role of biological factors (such as cellular differentiation pathways, endocrine levels etc.) during the development of the mammary gland, where PDEs are used to model biologically relevant molecular distributions and agents represent cells. In [13], Garbey et al. studied a hybrid PDE-ABM model to analyze vascular adaptation. However, to our knowledge, this hybrid approach has not yet been applied to explore the specific topic of viral infections' spatiotemporal dynamics. In fact, this is exactly what the main novelty of the present article consists in: while both the PDE-ABM hybrid model as a mathematical construction and the most urging current biological questions are already well-known, we bring these separately existing concepts into one single framework and examine several fundamental viral dynamic features in a new light. For the purpose of investigating the viral spread of SARS-CoV-2 and influenza, our hybrid model is chosen and tailored by carefully considering certain dimension-related aspects of the problem: virus particles are several magnitude smaller than body cells ([25], [26], [27]), and as a consequence, it is natural to capture virus concentration by a continuous function, while cells are modeled in a straightforward way as discrete entities in space. In other words, we obtain our hybrid model by merging a) a PDE representing virus concentration, and b) an agent-based model describing body cells and their three possible states (healthy, infected, and dead). We complete the hybrid PDE-ABM model by formulating a meaningful connection between the separate parts, i.e. by giving definition to all the considerable interactions and feedback processes that take place between the model's discrete and continuous parts, such as for example cells responding to their environment's virus concentration level. Then we

explore the emerging spatio-temporal dynamics in detail, and apply the model to SARS–CoV–2 and influenza infections.

2. Methods

The purpose of this section is to give a detailed description of the two main systems we use to model virus spread. We begin by defining the hybrid PDE–ABM model and introducing its variables, then we briefly outline a classical ODE model – the latter can be viewed as a simpler mean-field model that approximates the more complex hybrid framework by averaging over spatial variables. Once all the fundamental features of both systems are defined, we continue this section by expressing the connection between the corresponding model parameters, and conclude with the numerical implementation of the hybrid system.

2.1 The hybrid PDE–ABM model

As described in the Introduction, the main hybrid system is constructed via forming bridges between two important and rather different modeling techniques: we merge a discrete agent based model and a continuous PDE, and we create a meaningful connection between them by carefully designing their interactions with each other. We highlight the fact the model we obtain this way is defined in both space and time.

We begin by setting the notation for the domain we construct our model upon: let Ω be the mathematical representation of the part of lung tissue (or, in the case of an in vitro investigation, the relevant area of the examination) we are considering – for consistency reasons, we assume that Ω is an open set. Now we are ready to introduce the discrete part of our hybrid system.

One of the most important modeling decisions in the first part of the model construction is approaching epithelial cells as discrete agents. In more detail, we define a two dimensional ABM state space by introducing a lattice of $k_1 \times k_2$ agents representing lung cells: naturally, $k_1, k_2 \in \mathbb{N}$, and cells are identified via the (i,j) indices (corresponding to the respective agent's place in the grid), where $(i,j) \in L = \{(i,j) | 1 \leq i \leq k_1,\ 1 \leq j \leq k_2\}$. By introducing the $\Omega_{i,j}$ notation for the open set occupied by the (i,j)-th cell, we obtain that $\bar{\Omega} = \bigcup_{(i,j) \in L} \bar{\Omega}_{i,j}$.

Regarding the states of the ABM space's agents, each agent can have three possible states in our approach. This is formally grasped by following state function, which represents the concept that a cell is either *healthy*, *infected*, or *dead*:

$$s_{i,j}(t) = \begin{cases} \text{H,} & \text{if the } (i,j)\text{-th cell is healthy at time } t \\ \text{I,} & \text{if the } (i,j)\text{-th cell is infected at time } t \\ \text{D,} & \text{if the } (i,j)\text{-th cell is dead at time } t. \end{cases}$$

In terms of state dynamics, we use the following assumptions:

- for simplicity, we do not account for cell division or cell birth taking place during the time frame of the infection;

- the only reason for cell death is viral infection itself, i.e. death related to any other natural cause is ignored;

- all healthy cells are susceptible target cells;

- once a cell gets infected, it is not possible for it to become healthy again;

- the *healthy* → *infected* state change: a healthy cell may become infected once the virus has reached the given cell; moreover, infection is randomized and it occurs with a probability of P_I, where we highlight that P_I increases linearly with the virus concentration in the given cell (for more details see Section 2.3 on P_I's relationship to other parameters and Section 2.4 on implementation);

- the *infected* → *dead* state change: an infected cell dies with a probability of P_D, but similarly to infection, death too is approached from a stochastic viewpoint.

The stochastic element in the above state changes is of key importance. The inclusion of natural randomness makes the complex PDE-ABM hybrid model not only more realistic than the relatively simple ODE model, but also compared to a pure PDE system.

We note that apart from following the state changes of all cells on an individual level, we also introduce three additional system variables as functions taking discrete (in fact, natural) values: $H(t)$, $I(t)$ and $D(t)$ respectively denote the total number of healthy, infected, and dead cells in the two-dimensional ABM state space.

Now we are ready to consider the second part of our hybrid model. Let us begin by observing that the size of an epithelial cell is relatively significant ([26], [27]) and hence indeed it was a natural idea above to define cells as separate entities and follow their states on an individual level (instead of exclusively considering the total number of healthy, infected and dead cells as continuous functions for example). One of the cornerstone observations in the second part of the model is that viruses, on the other hand, are several magnitude smaller compared to epithelial cells [25] – we incorporate this simple but crucial biological fact into our system by modeling virus concentration as a continuous function. Hence, in the second part of the model we approach virus concentration $V^h = V^h(t, x, y)$ as a variable that is continuous in both space and time, and as such, we capture it by means of a partial differential equation:

$$\frac{\partial V^h(t, x, y)}{\partial t} = D_V \Delta V^h - \mu_V \cdot V^h(t, x, y) + \sum_{(i,j) \in L} g_{i,j}(t, x, y), \quad t > 0, \ (x, y) \in \Omega, \quad (1)$$

where D_V denotes the virus diffusion coefficient, μ_V is a constant ratio representing virus removal, while $g_{i,j}$ stands for the viral source term (the latter is assumed to be continuous) for the (i, j)-th cell. The above equation essentially grasps the basic concept that viruses spread across the domain primarily via diffusion (convective flows are ignored in this

model), the immune system removes viruses in a constant ratio, while new virus particles are generated by infected cells in a process described by the $g_{i,j}$ functions:

$$g_{i,j}(t,x,y) = \begin{cases} 0, & \text{if } s_{i,j}(t) = \text{H and } (x,y) \in \Omega_{i,j} \\ f_{i,j}(t,x,y), & \text{if } s_{i,j}(t) = \text{I and } (x,y) \in \Omega_{i,j} \\ 0, & \text{if } s_{i,j}(t) = \text{D and } (x,y) \in \Omega_{i,j} \\ 0 & \text{if } (x,y) \notin \Omega_{i,j} \end{cases} \quad (2)$$

We point out that in the above formula we do not specify any particular exact $f_{i,j}(t,x,y)$ form for the viral source term $g_{i,j}(t,x,y)$ for the case when $s_{i,j}(t) = \text{I}$ and $(x,y) \in \Omega_{i,j}$ (i.e. when the (i,j)-th cell is infected). As noted in [21], very little is known about the shape, duration and magnitude of viral burst, and hence, we allow any reasonable and smooth $f_{i,j}(t,x,y)$ function here that meets the following two criteria. Firstly, any concrete choice for $g_{i,j}$ needs to represent the general fact the (i,j)-th epithelial cell starts secreting virions at some point after it becomes infected. Thus, naturally, $g_{i,j}$ takes positive values at least in some subset of $\Omega_{i,j}$ at some point after the event of infection (but of course, at formulated above, outside of this given cell $g_{i,j}$ is zero). Secondly, the $g_{i,j}$ function needs to be defined in a way so that the well-posedness of system (1) is guaranteed: specifically, any concrete definition of the viral source term has to be Hölder continuous with Hölder exponent $\alpha \in (0,1)$, i.e. we assume $g_{i,j}(t,x,y) \in C^{\frac{\alpha}{2},\alpha}((0,\infty) \times \bar{\Omega})$ for any $t > 0$, $(x,y) \in \bar{\Omega}$ and any $(i,j) \in L$. The latter ensures the global existence of system (1)'s solution (see Appendix A). As an example, we can consider a definition where we use the mollified version of a constant rate release within the (i,j)-th cell. As for the details and changes related to the construction of $g_{i,j}$ in the implementation process, see Section 2.4.

The PDE-ABM hybrid model can not be complete without explicitly taking into account the meaningful interactions that each part-system has on the other, hence we briefly highlight these connections again. The continuous viral part effects the agent-based subsystem through the *healthy* $\rightarrow$ *infected* state change described in the ABM section above. On the other hand, the discrete ABM part makes a difference within the continuous viral equation thanks to the $g_{i,j}$ source functions (representing the fact that once a cell gets infected, at some point it starts spreading the virus).

2.2 The ODE model

In this subsection we consider a simpler viral dynamics system which will serve as a reference model for the hybrid framework in our epidemiological investigations. We consider the following – only time-dependent – ODE system:

$$\begin{cases} \frac{dH(t)}{dt} = -\beta H(t)V(t), & t > 0, \\ \frac{dI(t)}{dt} = \beta H(t)V(t) - \delta I(t), & t > 0, \\ \frac{dV(t)}{dt} = pI(t) - cV(t), & t > 0, \\ \frac{dD(t)}{dt} = \delta I(t), & t > 0, \\ H(0) = H_0 \geq 0, I(0) = I_0 \geq 0, V(0) = V_0 \geq 0, D(0) = D_0 \geq 0, \end{cases} \quad (3)$$

where the $H(t), I(t), V(t), D(t)$ functions represent the number of healthy cells, the number of productively infected cells, the number of viruses released by infected cells and the dead cells at time t, respectively; β denotes the healthy cells' infection rate, δ stands for the death rate of infected cells, p is the virus production rate, and c is the virus removal rate. We emphasise the fact that unlike the hybrid PDE-ABM model, the above ODE system is not defined in space.

Technically, the $V(t)$ function can be defined to represent either the number of viruses or the virus concentration itself depending on the specific application we are considering. In Section 3's simulation results $V(t)$ stands for concentration.

Simply because of the physical meaning behind these functions, the respective initial values are naturally set in the region $\Gamma = \{(u_1, u_2, u_3, u_4) \in \mathbb{R}^4 : u_1, u_2, u_3, u_4 \geq 0\}$. For the well-posedness and uniform boundedness of model (3) see Appendix B.

Setting the right-hand side of the equations in (3) equal to zero, we obtain that there are at most two kinds of ecologically meaningful equilibria in $\mathbb{R}_+^4$: the extinct equilibrium $e_0(0, 0, 0, 0)$ and the infection-free equilibrium $e_*(H_0, 0, 0, 0)$. The basic reproduction number of system (3) is given by

$$\mathcal{R}_0 = \frac{p\beta H_0}{c\delta}, \quad (4)$$

for the derivation of this number and how it governs the threshold dynamics of the system, see Appendix C.

2.3 The connection between the two main models and their parameters

In this subsection we examine the relationship between the two main models' respective parameters. Expressing these connections is not always a trivial task as the hybrid PDE-ABM framework exists in space, while the ODE model's functions are defined as variables only in time. In order to match the two different systems' corresponding parameters, we need to fix some basic features of the spatial domain: first of all, let A denote the complete area of the state space; moreover, for simplicity let us assume that each cell has the area of a unit space – for the latter we introduce the notation σ^2. Finally, let τ denote the unit step in time.

1. **Parameters related to cell death:** We emphasise that the two main models use different approaches to quantify the chance of an infected cell's death – on the one hand, the (3) ODE model works with a δ death *rate*; on the other hand, the hybrid

system uses a P_D *probability*. We can easily obtain a conversion between probability and rate by following the exact meaning behind these parameters. When infected cells die with a death rate δ, their natural decay can be described by the function $e^{-\delta t}$; hence, the probability of an infected cell's death between any two arbitrary time points t_1 and t_2 is given by

$$\frac{e^{-\delta t_1} - e^{-\delta t_2}}{e^{-\delta t_1}} = 1 - e^{-\delta(t_2 - t_1)}.$$

Specifically, for a time interval of length τ the above formula means that an infected cell dies within that given time frame with a probability of $1 - e^{-\delta\tau}$.

Applying the Taylor expansion of the exponential function and combining it with the fact that τ is small, we arrive to the $1 - e^{-\delta\tau} \approx \delta\tau$ approximation, i.e. we the connection between δ and P_D is given by

$$P_D \approx \delta \cdot \tau. \tag{5}$$

2. **Parameters related to new infections:** In the (3) ODE model the first equation corresponds to the infection of healthy cells: in other words, the $\beta \cdot V \cdot H$ term represents new infections per unit time, where β is the infection rate, V is the total number of viruses, and H is the total number of healthy cells at a given time. Respectively, in the PDE-ABM hybrid model we define the probability of a cell's infection in a way that this probability is directly proportional to the $V_{\Omega_{i,j}}$ virus concentration in the (i,j)-th cell (n.b. virus concentration per unit space), to the β infection rate, to the A area of the domain and to the τ time unit:

$$P_I \approx \beta \cdot A \cdot V_{\Omega_{i,j}} \cdot \tau. \tag{6}$$

3. **Parameters related to virus production:** The (3) ODE model's parameter p corresponds to virus production rate per unit time. Respectively, in our spatial hybrid model's virus dynamical equation (1) the parameter $g_{i,j}$ represents the virus production rate per unit time per unit space^2 – in particular, $g_{i,j} = f_{i,j}$ within infected cells (see 2), hence, clearly, $f_{i,j}$ matches p.

4. **Parameters related to virus removal:** Analogously, the respective pair of the hybrid model's m_V parameter is the ODE system's c virus removal constant.

5. **Parameters related to virus diffusion:** We highlight that the hybrid model's diffusive constant D_V does not have a corresponding parameter in the ODE system, as the latter model is defined only in time and diffusivity is strictly related to spacial dimensions.

We summarize the respective parameters' relationship to each other in Table 1.

Table 1: Description of the hybrid PDE-ABM model's parameters and their relationship to the ODE system. Naturally, the notations [healthy cell], [infected cell] and [dead cell] represent a healthy cell, an infected cell and a dead cell – as a unit – respectively. Analogously, [virion]/σ^2 stands for the concentration unit.

Symbol	Parameter	Unit	Value corresponding to the ODE model
P_I	Probability of infection for the (i,j)-th cell		$\beta \cdot A \cdot V_{\Omega_{i,j}} \cdot \tau$
P_D	Probability of death of an infected cell		$\delta \cdot \tau$
$f_{i,j}$	Virus production rate of an infected cell	$\frac{[\text{virion}]}{\sigma^2 \cdot \tau}$	p
D_V	Virus diffusion	$\sigma^2 \tau^{-1}$	
μ_V	Virus removal rate	τ^{-1}	c
H_0	The initial number of healthy cells	[healthy cell]	H_0
I_0	The initial number of infected cells	[infected cell]	I_0
$V_0^h(x,y)$	The initial value of virus concentration	[virion]/σ^2	V_0
D_0	The initial number of dead cells	[dead cell]	D_0

τ: time unit σ: space unit

Figure 1: An ABM state space where most cells are healthy, while a small number of randomly infected cells start spreading the virus in the domain.

2.4 Implementation

This section describes important technical details related to the hybrid PDE-ABM framework's implementation. Our numerical computations are based on a free and open source java software package, HAL (Hybrid Automata Library) [14]; our source code is publicly accessible in the Github repository [15].

We simulate the PDE-ABM model's behaviour in an $\Omega \subset \mathbb{R}^2$ two-dimensional bounded domain, hence the ΔV^h Laplacian takes the form of $V_{xx}^h + V_{yy}^h$, where $(x, y) \in \Omega$. The resolution of the state space is chosen to be 200×200, which means we work with a total number of 40.000 cells (this choice is mainly due to the hybrid model's computational demands – we found that concerning the number of cells 40.000 is a reasonable value for exploring virus dynamics on an ordinary computer).

In terms of spatial boundary conditions, we assume zero flux on the boundaries of the hybrid model.

As for the initial conditions in our simulations, we consider the case where we have zero virus concentration in the beginning, but we do have a small number of infected cells in the domain to start with – the virus will spread from these infected cells and as a result, more and more originally healthy cells become infected. As an initial setting, we assume that 5% of domain's cells are infected, and these infected cells are distributed randomly in space. Figure 1 shows the initial state of our hybrid PDE-ABM model.

After giving a clear definition of the initial and boundary conditions in our implementation, we highlight another interesting technical detail: the internal realization of the virus production functions. At the abstract definition of the hybrid PDE-ABM model we assumed that the virus source term in (1) was continuous. This was important for certain theoretical reasons (see Appendix A), but as we move on to implementation-related decisions and solutions, we make a simplifying step regarding this source function. We take a discretized approach, namely, throughout the simulations, a given cell's $g_{i,j}$ virus production function (and in particular, $f_{i,j}$) is quantified assuming that infected cells have a constant virus production rate; formally, $g_{i,j}(t, x, y) = $ a non-zero constant when $s_{i,j}(t) = \text{I}$ and $(x, y) \in \Omega_{i,j}$. We highlight that what really happens at the implementation

process is that in each time loop and for each cell (thus, for all $(i, j) \in L$) we solve equation (7) to obtain the virus concentration in a given cell:

$$\frac{\partial V^h(t, x, y)}{\partial t} = D_V \Delta V^h - \mu_V \cdot V^h(t, x, y) + g_{i,j}(t, x, y), \quad t > 0, \; (x, y) \in \Omega_{i,j}. \quad (7)$$

We also emphasise that in order to make the model as realistic as possible, we take some probabilistic considerations into account, and as a result we use a stochastic approach in the implementation of both cell infection and cell death.

- *Stochastic implementation of a healthy cell's infection:* In the $P_I = \beta \cdot A \cdot V_{\Omega_{i,j}}$ expression (obtained in (6)) we assume that the infection rate β and the A area of the domain are constant, but $V_{\Omega_{i,j}}$ is changing in both space and time – once the virus concentration is recalculated for a given cell, we then compare the correspondingly updated P_I value with an $r_{i,j}^I$ random number, which is generated newly for each cell. In case the cell's P_I value is greater than $r_{i,j}^I$, the cell in question gets infected.

- *Stochastic implementation of an infected cell's death:* Similarly, a random $r_{i,j}^D$ number is generated for each infected cell at each time-step, and the given cell's death is determined by comparing this random number with P_D: naturally, the infected cell dies if P_D is greater than $r_{i,j}^D$.

All of the random numbers noted above are real numbers from standard uniform distribution on the interval (0,1) and they are obtained by means of Java's random number generator.

Figure 2 shows the flow diagram of the program we implemented to simulate the virus spread and observe the spatial distributions of infected cells.

3. Results

We present and explain our most significant computational results in this section. In particular, we illustrate the advantages of the spatiotemporal hybrid system over the classical ODE model, we show an example where we successfully recreate the actual results of an in vitro experiment assessing SARS-CoV-2 propagation, moreover, by means of the PDE–ABM system we explore and compare the spatial patterns of influenza and COVID–19.

3.1 Comparing the hybrid PDE–ABM model and the ODE model through their solutions

This section is principally dedicated to illustrating the impact of the PDE–ABM model's built-in extra information regarding spatial factors such as diffusion. In the following steps we compare the numerical solutions of the (only time-dependent) ODE model and the hybrid spatiotemporal PDE–ABM system. We emphasise the fact that in this subsection

Figure 2: The program flow diagram of the PDE-ABM model's numerical simulation based on HAL.

Figure 3: A sample solution of model (3) considering the parameters given in (8) with $H_0 = 40000 - I_0$, $I_0 = 20$, $V_0 = 0$ and $D_0 = 0$ as initial values. **Left** Number of cells over time. **Right** Virus concentration over time.

we consider a generic scenario that does not necessarily correspond to influenza or SARS–CoV–2 specifically. Our main goal here is to outline a comparison itself between the two main models making sure that their parameters represent the same virus infection in identical circumstances – with the very important exception of virus diffusion, which incorporates spatial effects.

As for the ODE model's simulation, we consider the following parameter values:

$$\beta = 10^{-6}, \quad \delta = 8 \times 10^{-3}, \quad p = 4 \times 10^{-2}, \quad c = 2 \times 10^{-2}. \tag{8}$$

Figure 3 depicts the solution of the ODE model (3) using the parameter values defined in (8) and the initial conditions $H_0 = 40000 - I_0$, $I_0 = 20$, $V_0 = 0$ and $D_0 = 0$.

Once the ODE parameters are set, most of the corresponding hybrid parameter values are automatically defined through the connections described in Table 1. We obtain

$$P_I = 0.04 \times V_{\Omega_{i,j}}, \quad P_D = 8 \times 10^{-3}, \quad f_{i,j} = 4 \times 10^{-2}, \quad \mu_V = 2 \times 10^{-2}. \tag{9}$$

Naturally – due to the fact that the ODE model is defined only in time – the hybrid system's diffusion coefficient is an exceptional parameter missing from the list above. To demonstrate the notable difference made by virus diffusion in the hybrid model's numerical solutions (and to compare the latter with the ODE model's solution), we simulate our spatial model with two different diffusion coefficients:

$$D_{V1} = 0.2, \quad D_{V2} = 20 \cdot D_{V1} = 4. \tag{10}$$

Figure 4 and Figure 5 show the different cell and virus dynamics emerging from scenarios that are identical apart from their diffusion coefficients. Both graphs demonstrate cell states on the left and virus spread on the right, captured in four different time points. Infected cells (denoted by red squares) are apparently distributed randomly at the initial state, virus spread in the domain is clearly originating from infected cells; moreover, the two figures illustrate that a higher diffusion value results in a higher number of infected cells.

Similarly, Figures 6 and 7 confirm that different diffusion coefficient values lead to significantly different results regarding the entire system: the inclusion of spatial effects and the diffusion coefficient itself play a crucial role in virus dynamics.

Figure 4: Sample hybrid PDE-ABM results considering (9) and $\boldsymbol{D_{V1} = 0.2}$ at $t = 0$, 240, 480, and 720 (**A** - **D**). Infected, healthy, and dead cells (denoted by red, green, and black squares, respectively) are shown on the left and virus spread is depicted on the right in all four subfigures (**A** - **D**).

Figure 5: Sample hybrid PDE-ABM results considering (9) and $\boldsymbol{D_{V2} = 4}$ at $t = 0$, 240, 480, and 720 (**A** - **D**). Infected, healthy, and dead cells (denoted by red, green, and black squares, respectively) are shown on the left and virus spread is depicted on the right in all four subfigures (**A** - **D**).

Figure 6: A sample solution of the hybrid PDE-ABM model considering parameters specified in (9) and $D_{V1} = 0.2$. **Left** Number of cells over time. **Right** Virus concentration over time.

Figure 7: A sample solution of the hybrid PDE-ABM model considering parameters specified in (9) and $\bar{D}_{V2} = 4$. **Left** Number of cells over time. **Right** Virus concentration over time.

Figure 8: Comparison between the ODE model and the hybrid PDE-ABM system with two different diffusion values (as given in (10)). It is clearly seen that the outputs are very similar for a high diffusion coefficient, as opposed to the case of low diffusion coefficient. **A** Virus concentration over time. **B** Number of healthy cells over time. **C** Number of infected cells over time. **D** Number of dead cells over time.

Finally, in Figure 8 we compare the solution of the ODE system with the respective numerical solutions calculated for our spatial hybrid model considering the previously described two different virus diffusion values. Figure 8 illustrates a special relationship between the solutions of the two main systems: increasing the spatial model's diffusion coefficient apparently impacts the hybrid system's solution in a way that it, roughly speaking, gets closer to the ODE model's numerical solution – this corresponds to the fact that the ODE system represents a scenario where all particles can interact with any other particle, i.e. in some sense the diffusion coefficient is infinite. In other words, the lower the actual virus diffusion value is, the more important it is to consider a more complex spatiotemporal system to model the respective virus spread.

3.2 Applications

In one way or another, all applications of the proposed PDE–ABM hybrid model naturally focus on exploiting or predicting spatial information regarding virus spread. The first application detailed in Section (3.2.1) is dedicated to the latter; here we perform a simulation to obtain as much information as possible on the spatial dynamics of a COVID–19 infection and we compare our computer-generated predictions with actual results of in vitro experiments. On the other hand, Section (3.2.2) is dedicated to observing some important differences between influenza and COVID–19 – these are identified more easily thanks to the additional spatial information that is integrated into the hybrid model.

3.2.1 Predicting the spatiotemporal spread of SARS–CoV–2 in human airway epithelial cells

Perhaps one of the most straightforward applications is the prediction of spatial virus propagation itself. In this first scenario we simulate the spatial spread of a SARS-CoV-2 infection over the course of seven days and we compare our results with real-life observations obtained by scientific experiments. Specifically, we consider Figure(4B) in [29], where the authors examine the spread of fluorescent SARS-CoV-2 in human airway epithelial cells at the indicated days with or without peptide treatment – naturally, when we compare our results, we focus on the graphs listed in the line that corresponds to our scenario (i.e. investigating uninhibited virus propagation excluding any treatment).
Examining Figure(4B) in [29] and Figure 9 of the present article we observe that the key events and features of virus spread match in a reassuring way: the first significant and bigger sign of infection appears at day 4, a clearly visible peak is reached at day 6, while a slight decrease in the infection's severity starts to show between day 6 and day 7. At the same time we highlight that there is a natural limit to how accurately our simulations can recreate the specific events depicted in [29]: firstly, to our best knowledge, the exact values of virus death rate, infection rate and diffusion are not specified in [29]. Secondly, the authors have not disclosed the number of cells they worked with – this means that the resolution of the corresponding ABM fields are different (theirs being unknown). The latter is particularly important as it is most likely one of the reasons why the computer-simulated results are less "sharp" compared to the experimental scenario's

Figure 9: The simulated spatiotemporal dynamics of SARS–CoV–2 virus spread in human airway epithelial cells – the results were obtained by our source code implementing the hybrid PDE–ABM model. This sequence of pictures from our model output shows a striking resemblance to Figure(4B) in [29], where the latter depicts real experimental results assessing viral propagation.

images: roughly speaking, the generated results resemble ink spreading in water to some extent, while the in-vitro results have an apparent grain-like structure. This matter can possibly be the subject of future investigation.

3.2.2 Investigating and comparing the main properties of influenza and COVID-19

While Section 3.1 focused on comparing the ODE model's and the hybrid system's respective solutions in a generic scenario, the present – and in some sense, main – section's goal is to explore these solutions for two specific viruses, namely, influenza and SARS-CoV-2. In order to investigate the propagation of an influenza or COVID-19 infection, we need to set up both of our models with parameters that represent the features of the previously mentioned viruses as accurately as possible. This is a nontrivial task not only because SARS-CoV-2 is relatively new for scientists, but also because several other small, technical, but important issues arise when we want to compare two different kinds of solutions.

The ODE model's parameter setting is summed up in Table 2. The parameters representing influenza and COVID-19 are defined according to [16] and [18], respectively. We highlight two practical considerations we used during the parameterization process. *i)* Firstly, in order to make sure that the respective results of the ODE system and the hybrid model are comparable, we need to initialize them with the same number of healthy cells. Due to the high computational load of the latter system, this number is naturally limited – in our case we use $H_0 = 4 \times 10^4$. However, this number is lower than the number of healthy cells used in [16] and [18], which is problematic as it affects the value of R_0. Hence, to counterbalance the impact of the different H_0 value, we modify the value of β correspondingly. *ii)* Secondly, the viral load for influenza in [16] is given in $(TCID_{50}/ml)$ units, while [18] specifies the viral load index for COVID-19 in $(copies/ml)$. To overcome this, we apply the results of [20], we assume that a $TCID_{50}$ unit corresponds to 2×10^3 viral copies and we change the β and p values in the influenza parameter setting correspondingly.

As for the hybrid PDE-ABM model, the exact parameter values we use during the simulations are given in Table 3. With the exception of the diffusion coefficient, all of the hybrid model's parameter values have been calculated indirectly: we used the ODE parameters as a starting point and we combined these with the results detailed in Table 1 –

knowing the formal connection between the respective ODE and PDE-ABM parameters we can easily obtain the hybrid parameters in question. Finally, we note that although virus diffusion can vary due to several external parameters, we generally assume that the diffusion coefficient for influenza and COVID-19 corresponds to the values given in [21], and [22] respectively.

We highlight an important difference between the units of Table 1 (Section 2) and Table 3 (Section 3 / 3.2). While in the more theoretical Section 2 we conveniently used abstract units such as σ^2/τ, in the present section dedicated to applications all units become concrete (e.g. μm^2/min). This raises two interesting technical questions – the first one being the relationship between the real-life space unit μm and the abstract space unit σ still internally used within the implementation. The reassuring answer is that even though we can set an anatomically realistic choice such as $\sigma = 14\mu$m, the specific constant we use here eventually falls out from the virus dynamic equation where the diffusion coefficient multiplies the Laplacian; and hence, the equation itself remains unchanged. The second question concerns the relationship between the 3-dimensional ml unit and the 2-dimensional, internally used σ^2 (technically, we still solve equation (7) on the latter). This issue can easily be resolved by means of a simple projection: using that $1\text{ml} = 10^{12}\mu m^3$, we obtain that 1 ml corresponds to $10^8\mu m^2$ i.e. $10^8/14^2\sigma^2$ on the 2-dimensional lattice. As the constant appearing at the conversion multiplies all terms in (7), the actual equation we need to numerically solve stays identical.

Now we are ready to compare the two different models' predictive performances. Firstly, we consider the case of influenza. Figure 10 shows the respective numerical solutions obtained by the ODE model and the hybrid system – we note that the solution given by the ODE model is similar to the one described in [19]. We also refer back to an observation we made in Section 3.1 regarding the diffusion coefficients: the lower this diffusion value is, the further the hybrid and ODE models' respective solutions are from each other. Now, according to [21], our hybrid model's diffusion coefficient for influenza is set to 0.19 $\mu m^2 min^{-1}$, which is – in terms of virus diffusion in general – a relatively small value. Therefore, there is an apparent difference between the two solutions we obtained with the two different models. Finally, we highlight another important aspect in connection with Figure 10. The authors of [16] and [17] suggest that in the upper respiratory tract about 30%-50% of the epithelial cells are destroyed at the peak of infection. This corresponds to our simulated results in a reassuring way.

The second subject of our comparative investigations is COVID-19. Similarly to the concept of the previous image, Figure 11 demonstrates the simulated results given by the ODE model and the PDE-ABM hybrid system, but in this case the computations were executed with the parameters representing COVID-19. As noted in Table 3, the D_V parameter for SARS-CoV-2 is set to 0.6 $\mu m^2 min^{-1}$. This value is significantly higher compared to influenza's diffusion coefficient, and hence, as we expect, this time the ODE model and the hybrid system generate solutions that are visibly closer to each other than their respective solutions computed for influenza.

From this point in this section we examine influenza and SARS-CoV-2 propagation exclusively by means of the hybrid PDE-ABM system.

Table 2: Description of the (3) **ODE model's** parameters. Naturally, the notations [healthy cell], [infected cell] and [dead cell] represent a healthy cell, an infected cell and a dead cell – as a unit – respectively.

Symbol	Parameter	Unit	Value for influenza	Value for COVID-19	Ref
β	Infection rate	[virus copy] $^{-1}\cdot$ min^{-1}	0.94×10^{-7}	1.01×10^{-7}	
δ	The infected cell death rate	min^{-1}	2.66×10^{-3}	7.2×10^{-4}	[16] - [18]
p	The virus production rate	[virus copy] $\cdot$ min$^{-1}\cdot$ [infected cell] $^{-1}$	1.666×10^{-2}	3.72×10^{-3}	[16] - [18]
c	The virus removal rate	min^{-1}	2.08×10^{-3}	1.67×10^{-3}	[16] - [18]
H_0	The initial number of healthy cells	[healthy cell]	$40000 - 20$	$40000 - 20$	
I_0	The initial number of infected cells	[infected cell]	20	20	
V_0	The initial value of virus concentration	[virus copy] / ml	0	0	
D_0	The initial number of dead cells	[dead cell]	0	0	

Table 3: Description of the **hybrid PDE-ABM model's** parameters. Naturally, the notations [healthy cell], [infected cell] and [dead cell] represent a healthy cell, an infected cell and a dead cell – as a unit – respectively.

Symbol	Parameter	Unit	Value for influenza	Value for COVID-19	Ref
P_I	Probability of infection		3.76×10^{-3} $\times V_{\Omega_{i,j}}$	4.04×10^{-3} $\times V_{\Omega_{i,j}}$	
P_D	Probability of the infected cell death		2.66×10^{-3}	7.2×10^{-4}	
$f_{i,j}$	Virus production from an infected cell	[virus copy] $\cdot$ ml $^{-1} \cdot$ min^{-1}	1.666×10^{-2}	3.72×10^{-3}	
μ_V	The virus removal rate	min^{-1}	2.08×10^{-3}	1.67×10^{-3}	
D_V	Virus diffusion	μm^2 min^{-1}	0.19	0.6	[21] - [22]
H_0	The initial number of healthy cells	[healthy cell]	$40000 - 20$	$40000 - 20$	
I_0	The initial number of infected cells	[infected cell]	20	20	
$V_0^h(x,y)$	The initial value of virus concentration	[virus copy] / ml	0	0	
D_0	The initial number of dead cells	[dead cell]	0	0	

Figure 10: Comparison of the ODE model (solid lines) and the hybrid PDE-ABM model (dashed lines) for *influenza* using the parameters in Tables 2 and 3. All figures depict a change taking place over the course of seven days; specifically, **A** Viral load (*copies/ml*), **B** Number of healthy cells, **C** Number of infected cells, **D** Number of dead cells.

Figure 11: Comparison of the ODE model (solid lines) and the hybrid PDE-ABM model (dashed lines) for *SARS–CoV–2* using the parameters in Tables 2 and 3. All figures depict a change taking place over the course of seven days; specifically, **A** Viral load (*copies/ml*), **B** Number of healthy cells, **C** Number of infected cells, **D** Number of dead cells.

Figure 12: The difference between influenza and SARS–CoV–2 infections. The viral load **(Left)** and the number of infected cells **(Right)** simulated for influenza (dashed line) and COVID-19 (solid line) using the hybrid PDE-ABM model.

In order to compare the dynamical features of SARS-CoV-2 and influenza, we firstly simulate the viral load and the number of infected cells for both of these two viruses in Figure 12 (the respective solutions for the two different viruses are shown together). We highlight that in terms of the number of infected cells Figure 12 demonstrates that even though the peak itself occurs sooner in influenza, the value taken at this peak is much higher for COVID-19.

As a final step, we exploit the great advantage of having established a model that provides information not only about the temporal behaviour of the total number of healthy, infected and dead cells, but is also able to monitor an infection on an individual cellular level. Figures 13 and 14 demonstrate the distributional patterns of influenza and SARS-CoV-2 propagation. Comparing Figure 13/(**A**) and (**B**) with Figure 14 it is particularly visible that while influenza propagation shows spatial patterns that, roughly speaking, resemble flocks or well-outlined explosions, there is no analogous phenomenon occurring in the case of COVID-19: the latter is distributed among cells in a much more homogeneous manner through all stages of infection.

4.   Discussion

In this paper we have applied two different models for investigating the dynamical aspects of virus spread. The first model we considered was a hybrid PDE-ABM system, which is essentially a result of merging a discrete state space representing epithelial cells with a continuous reaction-diffusion equation grasping virus concentration. At the same time, we have used the so-called May-Nowak system – a well-known version of the classical ODE model – as a reference system. As for theoretical completeness, we provide a rigorous analysis of both models in the Appendix, including a well-posedness result related to the hybrid model and the study of the ODE model's temporal dynamics.

The hybrid model's computational implementation and the careful exploration of its results is in some sense the heart of our work – we highlight that our program code is based on a free and open source Java library, HAL (Hybrid Automata Library), commonly used for oncology modeling.

Compared to the ODE system, both the decisive advantages and the main difficulties

Figure 13: Simulated spatiotemporal numerical solutions captured **A** 20 hours, **B** 33 hours, **C** 53 hours, and **D** 67 hours after *influenza* infection. The computations were performed by means of the hybrid PDE-ABM model. Infected, healthy, and dead cells (denoted by red, green, and black squares, respectively) are shown on the left, while virus spread is depicted on the right in all four subfigures (**A** - **D**).

Figure 14: Simulated spatiotemporal numerical solutions captured **A** 20 hours, **B** 33 hours, **C** 53 hours, and **D** 67 hours after *SARS-CoV-2* infection. The computations were performed by means of the hybrid PDE-ABM model. Infected, healthy, and dead cells (denoted by red, green, and black squares, respectively) are shown on the left, while virus spread is depicted on the right in all four subfigures (**A** - **D**).

of the hybrid model are naturally related to the PDE-ABM system's added (and quite high-level) complexity: the inclusion of spatial effects. On the one hand, the limiting factors of this model include an increased computational demand and the fact that it is virtually impossible to consider a really large number of cells on an ordinary computer (we worked with a slice of tissue consisting of $4 \cdot 10^4$ cells). On the other hand however, this hybrid model provides us with the invaluable spatial distribution of infection spread: by running simulations for influenza and SARS-CoV-2 propagation, our spatiotemporal PDE-ABM system demonstrated that COVID-19 spreads more homogeneously among epithelial cells than influenza does. This simply would not have been possible using the ODE model as the latter is defined only in time and it represents a scenario where all cells can interact with all virus particles, or in other words, it implicitly assumes an infinite diffusion coefficient in some sense. Real-life viruses however clearly have a finite diffusion rate. This also means that if the specific virus in question has a relatively low diffusion rate, then the ODE model's predictions regarding infection dynamics will be less accurate: the lower the diffusion coefficient, the more important it is to apply the more complex and more suitable spatial hybrid model. This phenomenon can be observed in Figure 8: a small diffusion value results in great differences between the respective solutions of the PDE-ABM system and the (3) ODE model, while the solutions are indeed close to each other for a larger diffusion value.

In terms of verifying the accuracy and correctness of our proposed model, we highlight the results of Section (3.2.1) – we have relatively successfully recreated the real results of a scientific in vitro experiment: our computer-simulated results matched the actual events and features of infection spread on a satisfactory level. Regarding correctness, we refer to Figure 10 as well: about 30 to 50% of the epithelial cells are destroyed in the upper respiratory system at the peak of infection, which corresponds to the observations of [16] and [17].

As for possible further improvements and applications of our hybrid spatial model, we mention two main points. Firstly; fine-tuning features such as immune response processes, time delay between infection and virus production, and the phenomenon of cell regeneration are ignored in our current study. These can be the subject of possible future work, although we note that the present model itself can also be considered to be highly realistic in specific cases where some of the above mentioned elements are naturally negligible (e.g. at the short early phase of an infection the immune system has typically not responded yet, while the time frame is too short for cell regeneration to be relevant). Secondly, we plan to apply the hybrid system for parameter fitting analogously as [16] used the ODE model for a similar task for the case of influenza A. In more detail, the authors of [16] calculated a best fit of the ODE model using experimental data on viral load – they extracted viral kinetic parameters such as infection rate, virus production rate, viral clearance rate, and the half-life of free infectious viruses. We simulated the corresponding scenario with both systems and – as expected, considering the relatively low diffusion value of influenza – there was an apparent difference between the respective numerical solutions of the ODE model and the hybrid system. According to our results, [16] somewhat underestimates the R_0 value: as Figure 10 shows, in order to obtain a

solution with the PDE-ABM model that corresponds to the ODE solution (and hence, to the real curve), it seems that the R_0 value of the hybrid model needs to be higher than the value estimated by [16] to fit the experimental data. This is another example of how the assumption of homogeneous virus spread can be misleading – the kinetic values obtained by Baccam et al. [16] could be adjusted towards their real biological value by means of the hybrid PDE-ABM model. Thus, parameter estimation and fitting the stochastic hybrid model to various virological data is something we also consider as valuable future work.

Finally, we conclude by emphasising that the complex spatiotemporal information the hybrid model can provide on virus propagation is vital from the aspect of disease treatment and our proposed model can be of great use when assessing potential COVID-19 treatment strategies.

The hybrid PDE-ABM model is better suited for thorough and detailed virus spread assessment than the classical ODE system. Following virus propagation on an individual cellular level and taking important spatial effects into account results in a much more accurate and complete picture regarding the infection's outcome. Even though the additional integrated details clearly come at a price in terms of computational demand, this pays off very well in the form of information on spatial virus distribution and more accurate predictions.

Acknowledgments

RH acknowledges the financial support from Youth Foundation of Zhejiang University of Science and Technology (Grant No. 2021QN001, 2021QN046). SM and RH were supported by Hungarian grant NKFIH KKP 129877. NJ was supported by EFOP-3.6.1-16-2016-00008 and TUDFO/47138-1/2019-ITM. GR was supported by NKFIH FK 124016 and 2020-2.1.1-ED-2020-0000.

Appendix A. The well-posedness and uniform boundedness of the PDE model

In this subsection, we shall establish the global existence and uniform boundedness of the virus concentration function described by equation (1). We make the following suitable assumption:

$$(\mathbf{H_1}): g_{i,j}(t,x,y) \in C^{\frac{\alpha}{2},\alpha}([0,\infty) \times \bar{\Omega}) \cap L^{\infty}((0,\infty) \times \Omega)$$
$$\text{is nonnegative for each } i \text{ and } \alpha \in (0,1).$$

Theorem A.1 (Well-posedness and uniform boundedness) Let's suppose that the parameters D_V, μ_V are positive. Then for a non-negative initial value function $V_0^h(x,y) \in C^0(\bar{\Omega})$ system (1) has a unique non-negative global solution $V^h(t,x,y)$ defined on $[0,\infty) \times \bar{\Omega}$. Moreover, the solution $V^h(t,x,y)$ is uniformly bounded and satisfies $\limsup_{t\to\infty} \max_{\bar{\Omega}} V^h(t,x,y) \leq \max\{\frac{B}{\mu_V}, \sup_{\bar{\Omega}} V_0^h(x,y)\}$, where B is defined in (11).

Proof. We use the lower-supper method for our proof. According to $(\mathbf{H_1})$, we define

$$\Big\| \sum_{(i,j)\in L} g_{i,j}(t,x,y) \Big\|_{L^\infty((0,\infty)\times\Omega)} := B. \tag{11}$$

Let $\overline{V}(t,x,y) = \big(\sup_{\bar{\Omega}} V_0^h(x,y) - \frac{B}{\mu_V}\big)e^{-\mu_V t} + \frac{B}{\mu_V} := V^\star(t)$ and $\underline{V}(t,x,y) = 0$. Since

$$\frac{\partial \overline{V}(t,x,y)}{\partial t} - D_V\Delta\overline{V}(t,x,y) + \mu_V\overline{V}(t,x,y) - \sum_{(i,j)\in L} g_{i,j}(t,x,y)$$

$$\geq \frac{dV^\star(t)}{dt} + \mu_V V^\star(t) - B = 0 \tag{12}$$

and

$$\frac{\partial \underline{V}(t,x,y)}{\partial t} - D_V\Delta\underline{V}(t,x,y) + \mu_V\underline{V}(t,x,y) - \sum_{(i,j)\in L} g_{i,j}(t,x,y)$$

$$= -\sum_{(i,j)\in L} g_{i,j}(t,x,y) \leq 0. \tag{13}$$

Clearly the boundary condition satisfies $\frac{\partial \overline{V}}{\partial \nu} = 0 \geq 0 = \frac{\partial \underline{V}}{\partial \nu}$ and the initial condition satisfies $\overline{V}(0,x,y) = \sup_{\bar{\Omega}} V_0^h(x,y) \geq 0 = \underline{V}(0,x,y)$. Thus $(\overline{V}(t,x,y), \underline{V}(t,x,y))$ is a pair of coupled supper solution and lower solution of system (1). From $(\mathbf{H_1})$, we get that the function $f(t,x,y,V^h) = -\mu_V V^h + \sum_{(i,j)\in L} g_{i,j}(t,x,y)$ is Hölder continuous with exponent α with respect to t and x,y, and $f_{V^h}(t,x,y,V^h) = -\mu_V \in C(Q_T\times[m,M])$, where $Q_T = (0,T]\times\Omega$, $m = \min_{\bar{Q}_T}\underline{V}(t,x,y)$ and $M = \max_{\bar{Q}_T}\overline{V}(t,x,y)$. In view of [24, Theorem 2.4.6], system (1) has a unique solution defined on $(0,T]\times\Omega$ satisfying

$$0 \leq V^h(t,x,y) \leq V^\star(t). \tag{14}$$

By the arbitrariness of T, the solution of system (1) exists globally in time. Clearly, from the definition of $V^\star(t)$, we know that $V^\star(t) \leq \max\{\frac{B}{\mu_V}, \sup_{\bar{\Omega}} V_0^h(x,y)\}$. Then it follows from (14) that $\limsup_{t\to\infty} \max_{\bar{\Omega}} V^h(t,x,y) \leq \max\{\frac{B}{\mu_V}, \sup_{\bar{\Omega}} V_0^h(x,y)\}$, which ends the proof of the theorem. $\qquad\square$

In fact, since $\sum_{(i,j)\in L} g_{i,j}(t,x,y) \in C^{\frac{\alpha}{2},\alpha}([0,\infty)\times\bar{\Omega})$, it follows from the regularity theory of parabolic equations that system (1) has a unique classical solution $V^h(t,x,y) \in C^{1+\frac{\alpha}{2},2+\alpha}([0,\infty)\times\bar{\Omega})$. Here the sub-super solutions method is mainly employed to estimate the upper bound of virus particles in the whole spatial domain Ω as time evolves.

Appendix B. The well-posedness and uniform boundedness of the ODE model

Theorem A.2 Let's suppose $(H_0, I_0, V_0, D_0) \in \Gamma$ for our initial data. Then system (3) has a unique solution $(H, I, V, D) \in \Gamma$, i.e. Γ is positively invariant. Furthermore, the solution of system (3) satisfies

$$0 \leq H(t) \leq H_0, \quad 0 \leq \limsup_{t \to \infty} V(t) \leq \frac{p(N - V_0)}{c},$$

$$0 \leq \limsup_{t \to \infty} I(t) \leq (N - V_0) \min\{1, \frac{p\beta H_0}{c\delta}\}, \ 0 \leq D(t) \leq N - V_0. \tag{15}$$

where $N = H_0 + I_0 + V_0 + D_0$.

Proof. **Existence and Uniqueness.** By the continuity of right side of Eq. (3), the solutions of system (3) exist globally in $(0, \infty)$. Since the terms on the right-hand side of system (3) satisfy the local Lipschitz property respectively, the uniqueness of solutions follows from the standard theory of differential equations. This completes the proof.
Positive Invariance and Uniform Boundedness. For $H \geq 0, I \geq 0, V \geq 0, D \geq 0$, since

$$\dot{H}(t)\big|_{H=0} = 0 \geq 0, \quad \dot{I}(t)\big|_{I=0} = \beta HV \geq 0,$$

$$\dot{V}(t)\big|_{V=0} = pI \geq 0, \quad \dot{D}(t)\big|_{D=0} = \delta I \geq 0, \tag{16}$$

it follows from [23, Theorem 5.2.1] that the solution $(H(t), I(t), V(t), D(t))$ of system (3) is nonnegative for all $t \geq 0$ whenever $H_0 \geq 0, I_0 \geq 0, V_0 \geq 0, D_0 \geq 0$. Therefore Γ is positively invariant with respect to system (3).
Since $\dot{H}(t) = -\beta H(t)V(t) \leq 0$, we get $0 \leq H(t) \leq H_0$. Also from system (3), we obtain

$$\frac{d(H(t) + I(t) + D(t))}{dt} = 0 \quad \Rightarrow H(t) + I(t) + D(t) = N - V_0. \tag{17}$$

As $H(t) \geq 0, D(t) \geq 0$, we get $I(t) \leq N - V_0$. From the third equation of system (3), we have

$$\frac{dV(t)}{dt} \leq p(N - V_0) - cV,$$

which implies that

$$\limsup_{t \to \infty} V(t) \leq \frac{p(N - V_0)}{c}. \tag{18}$$

From (18), we know that for any $\varepsilon > 0$ there exists $T(\varepsilon) > 0$, s.t. when $t \geq T(\varepsilon)$, $V(t) \leq \frac{p(N - V_0)}{c} + \varepsilon$. Thus

$$\frac{dI(t)}{dt} \leq \beta H_0 (\frac{p(N - V_0)}{c} + \varepsilon) - \delta I.$$

By the comparison principle of ordinary differential equations, we get

$$\limsup_{t\to\infty} I(t) \leq \frac{p\beta H_0(N - V_0)}{c\delta}. \tag{19}$$

Since $I(t) \leq N - V_0$, we get $\limsup_{t\to\infty} I(t) \leq (N - V_0)\min\{1, \frac{p\beta H_0}{c\delta}\}$. This completes the proof. $\square$

Appendix C. The threshold dynamics of the ODE model

The basic reproduction number $\mathcal{R}_0$ can be deduced by simple reasoning: a single infected cell produces virus with rate p during its expected lifetime $1/\delta$, summing up to p/δ viruses. Virus particles generate infected cells (in a healthy cell population) with rate βH_0, in an expected time period $1/c$. Overall we find the number of new infected cells originated from the initial cell as given in (4). For a more formal derivation, one can also use the next generation approach [28].

Theorem A.3 If $\mathcal{R}_0 > 1$, then the infection-free equilibrium e_* is unstable, which means an infection can be established. Furthermore, system (3) cannot undergo a Hopf bifurcation around e_*.

Proof. The Jacobian matrix J_* around the infection-free equilibrium e_* is given by

$$J_* = \begin{bmatrix} 0 & 0 & -\beta H_0 & 0 \\ 0 & -\delta & \beta H_0 & 0 \\ 0 & p & -c & 0 \\ 0 & \delta & 0 & 0 \end{bmatrix}. \tag{20}$$

Thus the characteristic equation can be calculated as

$$\lambda^2[\lambda^2 + (c + \delta)\lambda + c\delta - p\beta H_0] = 0, \tag{21}$$

which has eigenvalues $\lambda_{1,2} = 0, \lambda_{3,4} = \frac{-(c+\delta)\pm\sqrt{(c+\delta)^2-4(c\delta-p\beta H_0)}}{2}$. Clearly, if $\mathcal{R}_0 > 1$, then the characteristic Eq. (21) has a positive real eigenvalue, which indicates that e_* is unstable. In addition, it is easy to see that there are no pairs of purely imaginary roots in characteristic Eq.(21) for any variational parameter, which implies that a Hopf bifurcation can not occur. This completes the proof. $\square$

References

[1] O. Stancevic, C.N. Angstmann, J.M. Murray, B.I.Henry, Turing patterns from dynamics of early HIV infection, Bull. Math. Biol. 75 (2013) 774-795.

[2] X. Lai, X. Zou, A reaction diffusion system modeling virus dynamics and CTLs response with chemotaxis, Disc. Const. Dyn. Sys. 21 (2016) 2567-2585.

[3] N. Bellomo, K.J. Painter, Y. Tao, M. Winkler, Occurence vs. absence of taxi-driven instabilities in a May-Nowak model for virus infection, SIAM. J. Appl. Math. 79 (2019) 1990-2010.

[4] J. Yang, X. Wang, F. Zhang, A Differential Equation Model of HIV Infection of CD T-Cells with Delay, Discrete Dynamics in Nature and Society (2008).

[5] M. Nowak, R. M. May, Virus dynamics: mathematical principles of immunology and virology, Oxford University Press, 2000.

[6] C. Macall, M. North, Tutorial on agent-based modelling and simulation, J. Simulat. 4 (2010) 151-162.

[7] Z. Wang, J.D. Butner, R. Kerketta et al., Simulating cancer growth with multiscale agent-based modelling, Semin. Cancer Biol. 30 (2015) 70-78.

[8] V.A. Folcik, G.C. An, C.G. Orosz, The basic immune simulator: an agent-based model to study the interactions between innate and adaptive immunity, Theor. Biol. Med. Model 4 (39) (2007) 1-18.

[9] O.M. Cliff, N. Harding, M. Piraveenan et al., Investigating spatiotemporal dynamics and synchrony of influenza epidemics in Australia: An agent-based modelling approach, Simula. Modelling Pract. Theor. 87 (2018) 412-431.

[10] E. Cuevas, An agent-based model to evaluate the COVID-19 transmission risks, Computers in Biology and Medicine 121 (2020) 103827.

[11] T.S. Deisboeck, Z. Wang, P. Macklin et al., Multiscale cancer modelling, Annu. Rev. Biomed. Eng. 13 (2011) 127-155.

[12] J.D. Butner, Y.L. Chuang, E. Simbawa et al., A hybrid agent-based model of the developing mammary terminal end bud, J. Theor. Biol. 407 (2016) 259-270.

[13] M. Garbey, S. Casarin, S.A. Berceli, A versatile hybrid agent-based, particle and partial differential equations method to analyze vascular adaptation, Biomechanics and Modeling in Mechanobiology 18 (2019) 29-44.

[14] R.R. Bravo, E. Baratchart, J. West, et al., Hybrid Automata Library: A flexible platform for hybrid modeling with real-time visualization. PLoS Comput. Biol. 16(3) (2020) e1007635.

[15] S. Marzban, R. Han, N. Juhász, G. Röst, Hybrid PDE-ABM model for viral dynamics, https://github.com/sadeghmarzban/Hybrid-PDE-ABM-for-viral-dynamics, 2021.

[16] P. Baccam, C. Beauchemin, C.A. Macken et al., Kinetics of Influenza A Virus Infection in Humans, Journal of Virology, 80 (2006) 7590-7599.

[17] G.A. Bocharov, A. A. Romanyukha, Mathematical model of antiviral immune response III. Influenza A virus infection, J. Theor. Biol. 167 (1994) 323-360.

[18] E.A. Hernandez-Vargas, J.X. Velasco-Hernandez, In-host Mathematical Modelling of COVID-19 in Humans, Annu. Rev. Control 50 (2020) 448-456.

[19] E.A. Hernandez-Vargas, Modeling and Control of Infectious Diseases in the Host, Academic Press, 2019.

[20] W. Yang, S. Elankumaran, L.C. Marr, Concentrations and size distributions of airborne influenza A viruses measured indoors at a health centre, a day-care centre and on aeroplanes, J. Roy. Soc. Interface, 8(61) (2011) 1176-1184.

[21] C. Beauchemin, S. Forrest, F.T. Koster, Modeling Influenza Viral Dynamics in Tissue, Springer, Berlin, Heidelberg, 2006

[22] T.J. Sego, J.O. Aponte-Serrano, G.J. Ferrari, et al., A modular framework for multiscale, multicellular, spatiotemporal modeling of acute primary viral infection and immune response in epithelial tissues and its application to drug therapy timing and effectiveness, PLoS Comput. Biol. 16 (2020) e1008451.

[23] H.L. Smith, Monotone dynamical systems: an introduction to the theory of competitive and cooperative systems, American Mathematical Society, providence, Rhode Island, 1995

[24] Q.Y. Ye, Z.Y. Li, M.X. Wang et al., Introduction to reaction-diffusion equations, Science Press, Beijing, 2011 (in Chinese).

[25] Y.M. Bar-On, A. Flamholz, R. Phillips et al., SARS-CoV-2 (COVID-19) by the numbers, eLife 9 (2020) e57309.

[26] M. Carcaterra, C. Cristina, Alveolar epithelial cell type II as main target of SARS-CoV-2 virus and COVID-19 development via NF-Kb pathway deregulation: A physiopathological theory, Medical Hypotheses 146 (2021) 110412.

[27] R.J. Mason, Biology of alveolar type II cells, Respirology 11 (2006) S12-S15.

[28] P. van den Driessche, J. Watmough, Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission, Math. Biol. 180 (2002) 29-48.

[29] V.K. Outlaw, F.T. Bovier, M.C. Mears, et al., Inhibition of Coronavirus Entry In Vitro and Ex Vivo by a Lipid-Conjugated Peptide Derived from the SARS-CoV-2 Spike Glycoprotein HRC Domain, mBio. 11 (2020) e01935-20.