The integrated role of resource memory and scent-based territoriality in the emergence of home-ranges

Meryl Theng[1*]
meryltheng@gmail.com

Thomas A. A. Prowse[1]
thomas.prowse@adelaide.edu.au

Phillip Cassey[1]
phill.cassey@adelaide.edu.au

Steven Delean[1]
steven.delean@adelaide.edu.au

Chloe Bracis[2]
chloe.bracis@gmail.com

[1]Ecology & Environmental Science, School of Biological Sciences, University of Adelaide, Australia.
[2]Université Grenoble Alpes, TIMC-IMAG / BCM, Grenoble, France
* Corresponding author: meryltheng@gmail.com; +61 432347080; Floor/Room G24A, Benham, North Terrace, University of Adelaide, Adelaide, Australia

Keywords: movement ecology, animal space use, individual-based models, mechanistic models, mechanistic

Statement of authorship: MT performed all simulations and analysed output data and wrote the original draft with support from all authors. Both MT and CB conceptualised and designed the study. CB designed and provided the model.

Abstract

Despite decades of animal movement research, we still lack an integrated understanding behind the emergence of home-ranges. We explored the effects of integrating two key processes of home-ranging arising from individual movement decisions: (i) memory-based resource use; and (ii) scent-based territoriality. Both mechanisms led to the formation of exclusive, negative resource- and density-dependent home-ranges, which were responsive to perturbations in the conspecific environment (i.e., removing individuals). Home-range patterns (size, size skewness and overlap) were primarily influenced by memory and scent decay rate parameters, demonstrating that home-ranging is ultimately a balance between an animal's inherent exploratory tendency and its desire to avoid conspecifics. Model application to a population of feral cats demonstrated that general space use patterns could be approximated through simulation, and replication of finer-scale space use patterns is plausible with further model development. Our modelling framework can provide a foundation for sophisticated theoretical models of space use in interacting animals.

1   Introduction

Many animal species constrain their movement to specific home-ranges (HR), which emerge from behaviours ultimately to survive and reproduce (Burt 1943). The emergent HRs observed result from movement decisions of individual animals, which are driven by dynamic processes including their need to access resources while avoiding costly interactions with conspecifics and predators (Borger *et al.* 2008; Nathan *et al.* 2008). Despite decades of extensive research, it is only recently that models of HR emergence have been developed (Ranc *et al.* 2020). Two key processes are proposed to play important roles in HR formation in territorial animals: (1) optimising resource acquisition by referencing a cognitive memory (i.e., resource memory); and (2) minimising resource competition through defensive cues (i.e., territoriality) (Borger *et al.* 2008; Powell & Mitchell 2012; Spencer 2012; Fagan *et al.* 2013). While there have been successes in modelling these underlying mechanisms separately, integration of the two to form a general predictive theory of HR emergence has remained a key challenge (Potts & Lewis 2014).

A fundamental characteristic of animal HRs is the regular revisitation to locations such as foraging areas, dens, watering holes and movement corridors (a.k.a. 'site fidelity'). Animal memory provides a plausible biological explanation of this phenomenon and recent empirical evidence supports this hypothesis (Bracis & Mueller 2017; Merkle *et al.* 2017; Ranc *et al.* 2021). While quantifying memory is particularly challenging, theoretical analyses have demonstrated that memory-based foraging processes can produce emergent home ranges and more efficient resource use (Van Moorter *et al.* 2009; Bracis *et al.* 2015; Riotte-Lambert *et al.* 2015). Modelling memory mechanisms essentially captures the underlying process behind formation of HR boundaries, and spatio-temporal patterns of site use and fidelity within a HR. It allows replication of the dynamic nature of HRs as a response to a changing environment (e.g., Potts *et al.* 2013; Bateman *et al.* 2015), which is a key advance from non-

mechanistic models that have commonly assigned localising centres or HR boundaries to achieve stable, but unrealistically static HRs (Borger *et al.* 2008).

Competitive interactions can drive spatial segregation of HRs, particularly in territorial animals that maintain and defend exclusive territories against conspecifics. In classical mechanistic models of animal movement, territoriality is modelled as scent-mediated conspecific avoidance (Giuggioli *et al.* 2013; Potts & Lewis 2014), which has been demonstrated as a significant underlying driver of observed variation in individual HRs and changes in HR patterns following population change in territorial carnivores (Lewis & Murray 1993; Moorcroft *et al.* 2006; Bateman *et al.* 2015). While these models have led to realistic patterns of HR formation, most have imposed a redirect-to-centre response to scent to stabilise otherwise unconstrained expansion of HRs caused by diffusive movement (Borger *et al.* 2008; Potts & Lewis 2014). This non-mechanistic component is not suitable for modelling HRs of animals that are not central place foragers or denning animals, nor does it allow the emergence of dynamic localising behaviours as a response to changing environments (e.g., HR shifts following resource depletion or conspecific removal). Moreover, the redirect-to-centre response does not explain the underlying localising movement behaviours in the absence of conspecifics (e.g., in sparsely populated habitats), which can result from memory processes.

Though critical insight has been gained from modelling resource memory and territoriality separately, the integration of these two important aspects of HR formation is unexplored. Each component essentially provides a mechanistic explanation for what the other lacks: resource memory is an attractive driver for individuals to preferentially acquire resources from previously visited sites, while territoriality is a repulsive driver which causes individuals to establish exclusive HRs in a multi-individual context (Potts & Lewis 2014). Here, we advance toward a more holistic understanding of HR formation, by developing what

is, to our knowledge, the first theoretical model integrating resource memory and territorial

processes to simulate realistic patterns of space use by territorial animals. We extend a two-

state memory-based model for non-territorial animals (Bracis *et al.* 2015) to include multiple

individuals that interact through scent-mediated conspecific avoidance behaviour. To explore

and illustrate the effects of integrating these two components, we investigate how the

interplay of resource memory and territoriality influences: (1) the emergence of individual

HRs; (2) the relationship between HR size, population density and resource availability; and

(3) the response of animal HR to changes in the conspecific environment (i.e., removing

individuals). Next, we conduct a sensitivity analysis on the movement model to assess

parameter importance for HR patterns (size, size skewness and overlap) and other metrics of

movement (step-length, step-length distribution skewness, daily distance) to guide the

calibration of model parameters using telemetry data.

2 Methods

We first describe the movement model's concept (**section 3.1**; summarised in **Fig. 1**), then

detail the parameterisation process used for simulations in the following **sections 2.2** and **2.3**.

All parameter values used in simulations are collated in **Table 1**.

2.1 Movement model description

In our model, simulated individuals move around a dynamic resource landscape, learning

about the intrinsic quality of the landscape as they consume resources (based on Bracis *et al.*

2015; **Fig. 1a**).

2.1.1 Movement process

86   An animal's movements through the landscape are described by a continuous trajectory with

87   a current position of $Z(t) = \int_0^t V(t') \, dt' + Z_0$, with a velocity of $V(t)$ and initial position

88   of Z_0. The autocorrelated, directed, continuous movement process is given by

$$dV = \frac{1}{\tau}(\mu(t) - V)dt. \tag{1}$$

This is similar to the Ornstein-Uhlenbeck process where τ is the time scale of autocorrelation

and instead of the white noise component, stochasticity is introduced through the bias vector

$\mu(t) = (v, \angle\theta)$ of magnitude $v = ||\mu(t)||$ (controlling average speed) and angle θ

(direction). A Poisson process with rate parameter λ determines when angle θ is updated,

which is then selected from an angular probability distribution derived from resource

memory or scent processes, depending on the behavioural state. Finally, individuals switch

between feeding and searching states, characterised by different values for τ and v, based on

the current resource consumption $C(t)$ (**Fig. 1b**).

2.1.2   Resource memory

The resource Q is modelled as continuously varying in space across the landscape. Resources

deplete as they are consumed by individuals and regenerate logistically at a rate of β_R, but do

not shift in space. Thus, it is advantageous for the individual to leave recently depleted

patches but return to high quality areas over the long term. Animals consume resources

according to a spatial kernel (a bivariate normal distribution with length scale γ_C) and

consumption rate β_C. Animals have a resource memory with two different streams of

information: a short-term stream S that pushes the animal away from recently visited

locations even if they are attractive, and a long-term stream L that attracts the animal back to

high quality areas (Van Moorter *et al.* 2009). The latter can either be initiated fully informed,

with the intrinsic resource quality Q_0, or naively, with a homogenous map of value $M *$

indicating the animal's expectation for unvisited areas which can be more optimistic or

pessimistic and thus affect exploratory tendency (see Bracis & Wirsing, In review). $M *$ is

also the value to which long-term memory L decays. The two streams are combined into a

single memory map M, which is used to inform the movement process.

The resource memory contribution to this angular probability distribution is computed

by integrating transects of the resource memory map radiating out from the individual's

location with the resource memory value at each point weighted by distance

116
$$g(\theta) = \frac{\int_0^r M(r,\theta) f_z(r) dr}{\int_0^{2\pi} \int_0^r M(r,\theta') f_z(r) dr d\theta'}, \tag{2}$$

117   where θ represents the angle, and $f_z(r)$ is the kernel function that weights according to

118   distance (here exponential with length scale γ_z). The foraging memory movement model is

119   described in further detail in Bracis *et al.* (2015).

2.1.3   Scent-marking (territoriality)

As individuals move about the landscape, they also deposit scent, which decays over time,

thereby marking their territory. The amount of scent on the landscape, D, is governed by the

deposition rate, β_D, (i.e., how much scent is deposited), the deposition spatial scale, γ_D, (i.e,

how broadly the scent is deposited in the vicinity of the forager) and the exponential decay

rate ϕ_D. D is limited to a maximum value of D_0 and we take D_0 to be one. Thus, the change

in scent for each forager at location $z = (x, y)$ is given by the equation

127
$$\frac{\partial D}{\partial t} = \beta_D f_D(|z - Z|)(D_0 - D) - \phi_D D, \tag{3}$$

128   where f_D is the spatial kernel (here exponential with scale γ_D). Scent deposition is tracked per

129   individual, and foragers are indifferent to their own scent, but repulsed by the scents of all

130   other conspecifics.

131   The repulsion of individuals by the scent of conspecifics is represented with an angular

132   conspecific safety metric that scales between 0–1 but is not constrained to sum to 1 (like

predation risk in (Bracis *et al.* 2018b)). It represents the relative 'safety' in each direction in terms of avoiding conspecifics, with 0 meaning not safe (high levels of conspecific scent) and 1 meaning safe (no conspecific scent). It is calculated by integrating the summed values of all other foragers' deposited scent according to

$$d_i(\theta) = 1 - max\left(1, \int_0^r \Psi_D \sum_{n \neq i} D_n(r, \theta) f_w(r) dr \right), \tag{4}$$

where Ψ_D is the response strength and $f_w(r)$ is a spatial kernel (here the exponential kernel with length scale γ_w) that represents that decay of scent perception with increasing distance.

2.1.4 Decision rules

To create the final angular probability distribution from which the angle θ is drawn to inform the movement process, the angular probability distribution based on the resource memory $g(\theta)$ is multiplied by the conspecific safety metric $d_i(\theta)$ for individual i, then normalised, giving

$$h_i(\theta) = \frac{g(\theta)d_i(\theta)}{\int_0^{2\pi} g(\theta')d_i(\theta')d\theta'}, \tag{5}$$

and the orientation angle of the bias term $\theta(t) = \angle\mu(t)$, in the movement process is then drawn from $h_i(\theta)$, which is specific for each individual.

2.2 Theoretical examples

For this section, the movement model was simulated on a 200×200 pixel heterogenous resource landscape, generated with the function *nlm_fbm* in R package *NLMR* (*fract_dim* = 0.75). Initial locations of animals were random, and the landscape boundaries were reflective.

2.2.1 Impacts of resource memory and territoriality

To demonstrate the effect of resource memory and territoriality on emergent HRs, we

simulated animal movement trajectories for three scenarios: (1) territoriality only, (2)

memory only; and (3) territoriality and memory. In scenario 1, the movement process was a

resource-dependent two-state random walk, with $g(\theta)$ in Eq. 5 being uniform while

scenarios 2 and 3 use the memory process of Eq. 2. All three scenarios were run across three

different densities of individuals ($n = 5, 10, 15$), creating 100 replicate simulations for each

combination of scenario and density. Individual HR size was estimated from the simulation

outputs using a kernel density estimator function *kernelUD* in the R package *adehabitatHR*.

The 95% HR polygons were used to extract resource values within the landscape using the R

packages *sp* and *raster*, after which the mean resource value within each HR was computed.

Variance in each individual's HR size was also visualised with respect to density and

resource availability. A two-way ANOVA was used to evaluate the effect of density and

scenario on $\log_{10}$-scaled HR size.

2.2.2   Impacts of population manipulation

To demonstrate that HRs are both an emergent property and dynamic when conditions

change in our model, we tested the effect of individual removal on remaining individuals'

HRs. We first simulated 100 datasets with a density of 10 individuals (i.e., a control setting).

We simulated a further 100 datasets with the same parameters and starting conditions as the

control setting, and then used a three-phase approach to remove individuals and quantify the

effects of their removals on the HRs of the remaining individuals. The three phases were:

(Phase 1) establishment of HR ($t = 1$–20,000); (Phase 2) remove four out of ten individuals (t

$= 20,001$), followed by a transition period to allow the scent of removed individuals to nearly

disappear ($t = 20,001$–30,000); and (Phase 3) exploration and establishment by remaining

individuals ($t = 30,001$–50,000). To quantify the effects of removal, we compared the

aggregate overlap in area (95% HR) between initial area covered by removed individuals

(calculated from Phase 1 with 5,000 timesteps burn-in) and the area covered by remaining

individuals before and after removal (calculated from Phase 1 and 3 respectively with 5,000

timesteps burn-in. A paired t-test was used to compare differences between the dependent

before-after samples. For comparisons in the control setting, four of the 10 individuals were

randomly "marked for removal" (but not actually removed). Aggregate overlap in area

covered between "marked for removal" and "unmarked" individuals was calculated using the

function *over* from R package *rgeos*.

2.3   Model calibration to empirical data using sensitivity analysis

To facilitate a better mechanistic understanding of emergent properties of the model and to

explore the potential for model applications, we (1) conducted a sensitivity analysis (SA) to

evaluate input parameter influence; and (2) used the SA results to guide calibration of

parameters to match general patterns in an empirical case study that consisted GPS-locations

from 11 adult female and male feral cats (*Felis catus*) from a 2019 eradication operation

conducted in Kangaroo Island (South Australia) by the Department for Environment and

Water South Australia (see **Text S1** for details on the empirical study and simulation setup).

2.3.1   Sensitivity analysis

We assessed the relative influence of the following nine input parameters on emergent

movement patterns: resource regeneration rate, searching speed, feeding speed, short-term

memory decay rate, long-term memory decay rate, memory expectation, scent deposition

rate, scent decay rate and scent response spatial scale. Emergent patterns were summarised

with six summary statistics for 1,000 sampled input parameter combinations (one replicate

simulation per combination): step-length, step-length distribution skewness, daily distance,

HR size, HR size distribution skewness and HR overlap. The influence of input parameters on each respective summary statistic was estimated with boosted regression trees (BRT) to account for complex, non-linear relationships and parameter interactions (Elith *et al.* 2008; Prowse *et al.* 2016). We calculated the relative influence of each parameter (scaled to sum to 100), with higher numbers indicating stronger influence on the response. In models with interactions, we measured the interaction strength for top pairwise interactions between input parameters, with values near zero represent negligible interactions. Details on summary statistic calculation and model fitting are in **Text S2** and **S3** respectively.

2.3.2   Parameter calibration

We proceeded with parameter calibration via a multi-step process: (1) optimising to a parameter set with which to achieve the target values for the six summary statistics simultaneously (using respective BRT models); and (2) manual calibration to five target criteria: HR size and HR size skew (***quantitatively evaluated***), HR overlap, cumulative HR size and spatial distribution of locations (***visually evaluated***). The second step was iterated until no further improvement was found (i.e., improving one metric made another worse). We then ran 10 simulations using our final set of parameters and compared the outcomes with the empirical data. Details for this section are in **Text S4**.

All simulations (section 2.2 and 2.3) were performed using Java v. 13.0.2 on a High-Performance Computing cluster. Analyses of model outputs were performed in R v. 3.6.1 (R Core Team 2020).

3　Results

3.1　Resource memory and territoriality lead to exclusive, negative resource- and density-dependent HRs

Individuals in the territoriality-only scenario established distinct non-overlapping territories and space use was independent of resource (**Fig. 2a**). In the memory-only scenario, individual space use favoured high resource areas but overlapped substantially (**Fig. 2b**). With both territorial and resource memory processes operating, individuals established distinct territories that utilised high resource areas with relatively little overlap (**Fig. 2c**).

Across all three scenarios, individuals in low resource areas tended to have larger HRs than in high resource areas (**Fig. 2d**). This relationship was strongly log-linear in the memory-only and territoriality + memory scenarios (Pearson's $r = -0.75$; $p < 0.001$), though variation in both HR size and mean resource were larger for the latter. The resource and HR size relationship in the territoriality-only scenario was consistent with the other scenarios at mid to high resource areas but starts to trend towards a positive HR-resource relationship in low resource areas, with a much lower lower-limit of mean resource compared to the other two scenarios.

An increase in density caused a decrease in mean HR size in the territoriality-only (mean(HR size)$_{density=5/10/15}$ = 11,860/6,670/4,600 unit^2) and territoriality + memory (mean(HR size)$_{density=5/10/15}$ = 6,730/4,200/2,840 unit^2) scenarios, while it caused a slight increase in mean HR size in the memory-only scenario (mean(HR size)$_{density=5/10/15}$ = 11,230, 12,080, 12,730 unit^2; two-way ANOVA, $F_{2,8995} = 450$, $p < 0.001$; **Fig. 2e**). Overall, the smallest HRs were observed in the territoriality + memory scenario (two-way ANOVA, $F_{2,8995} = 2,580$, $p < 0.001$).

3.2　Individuals move into recently unoccupied areas

Aggregate HR overlap between remaining individuals and removed individuals was

markedly higher in Phase 3 compared to Phase 1 (mean(HR size)$_{\text{Phase 1/3}}$ = 7,000/14,430 unit^2;

paired t-test: t_{99} = -19.9, $p < 0.001$) of the removal simulations, while there was a smaller

increase in the overlap between individuals (mean(HR size)$_{\text{Phase 1/3}}$ = 6,580/8,980 unit^2; paired

t-test, t_{99} = -8.2, $p < 0.001$) in the control simulations (**Fig. 3**). One example removal

simulation showed space use shifts in remaining individuals into areas previously occupied

by removed individuals. Individuals in the control example demonstrated smaller shifts in

space use on average, explaining the small increase in aggregate HR overlap from Phase 1 to

3.

253   3.3   Sensitivity analysis

HR size was strongly influenced by memory decay rates and scent decay rate (**Fig. 4a**).

Short-term memory decay rate had a strong negative effect on HR size, whereas long-term

memory and scent decay rates had a generally positive effect varying in strength at different

points in their parameter ranges (**Fig. 4b**). Interactions led to a combined effect that if scent

decay rate was high, and either short-or long-term memory decay rate was low and high

respectively, a 4–6 times increase in HR size was observed (**Fig. S7**). Longer right-tailed

skews in HR size distribution were primarily caused by slower rates of scent decay, faster

rates of short-term memory decay and larger scent response spatial scales (**Fig. 4c**). Smaller

overlaps in HR were predominantly associated with slow scent decay rates and fast short-

term memory decay rates (**Fig. 4d**).

Distance metrics (step-length, step-length skewness, daily distance) were mainly

influenced by resource regeneration rate and search speed (**Fig. 4a**). Step-length and daily

distance increased, while step-length skewness decreased, with lower resource regeneration

rates and higher search speeds (**Fig. S3**–**5**).

3.4   Empirical application

The simulated data generally captured the long right-tailed distribution of HR sizes that reflected populations of predominantly resident individuals (with smaller and distinct HRs) and a small number of transient individuals (with larger and less distinct HRs, and sparser location density) evident in the empirical cat movement data (**Fig. 5a**). Median HR was the same in the simulated and empirical data (1.9 km^2). However, simulated HR size range tended to be smaller than the empirical data – approximately a tenth of simulated individuals had HRs smaller than one km^2 and few ranged wider than nine km^2 across the replicates. Furthermore, the cumulative HR in the simulations reflected a less stable system compared to the empirical data (i.e., curves did not approach a plateau for most individuals; **Fig. 5b; Table S2** for all). Spatial patterns of simulated locations resembled empirical movement patterns in that locations tended to cluster linearly along high resources, but to a markedly lesser extent. The intense clustering of locations in specific sites observed for cats was not reproduced in the simulated data (**Fig. 5c**).

4   Discussion

We showed that integrating two key drivers of HR formation, resource memory and territoriality, gives rise to spatially distinct and dynamic HRs that vary in size according to resource distribution. We also demonstrated that stable HR patterns are ultimately a balance between an animal's inherent exploratory tendency and its desire to avoid conspecifics. Additionally, we approximated the general spatial measures and patterns of space use of feral cats, including the phenomenon of transience, through simulations characterised by specific assumptions about resource memory and avoidance of conspecifics.

In our multi-individual simulation environment, two mechanisms were required to replicate patterns of HR formation in territorial animals: one that tended towards fidelity to

previously visited, high quality areas (resource memory); and another that drove spatial

avoidance between individuals (territoriality). Further, our model captured realistic responses

to density with the emergence of transient behaviour by some individuals as density increased

and all higher quality areas were occupied. The absence of territoriality meant that

individuals were acting independently (of conspecific presence), which caused large overlaps

in emergent HRs (**Fig. 2b**). Whereas the absence of resource memory resulted in unrealistic

HR formation that was independent of resource quality; movement was only constrained by

the scent of surrounding conspecifics (**Fig. 2a**). The combination of both mechanisms led to

the emergence of distinct individual HRs distributed across high resource areas of the

landscape (**Fig. 2c**). In addition to being spatially realistic, biologically meaningful

relationships were also captured. Simulated individuals' HR sizes correlated negatively with

both resource availability and population density, which is congruent with ecological

evidence from both field and experimental studies (Baker *et al.* 2000; Santangeli *et al.* 2012;

Šálek *et al.* 2014; Schoepf *et al.* 2015). This is biologically intuitive: when resources are

abundant, an individual needs less space to meet its metabolic needs (McLoughlin &

Ferguson 2016); when population density is high, the amount of available, unoccupied space

is smaller, which limits individual HR sizes (Schradin *et al.* 2010).

Home-ranging is a dynamic property emerging from our model, continuously adapting

to the dynamic resource landscape and presence of surrounding conspecifics. This was

emphasised in our removal simulations, where remaining individuals migrated into areas

previously occupied by removed individuals once their scent decayed. The phenomenon of

expanding into newly vacated habitat can be found in numerous species, such as mice

(Schoepf *et al.* 2015), chipmunks, (Mares *et al.* 1976), coyotes (Moorcroft *et al.* 2006), and

red foxes (Potts *et al.* 2013). Non-mechanistic models (e.g., with imposed HR centres and/or

boundaries) fail to respond to changing environments (e.g., individuals leaving or dying,

fluctuating resource availability), whereas in our model dynamic behaviour arises directly from underlying mechanisms. Notably, the memory mechanism in our model provides the basis for HR stabilisation in the absence of conspecific avoidance, which has largely remained an imposed property in existing mechanistic territorial models (Potts & Lewis 2014).

Our sensitivity analysis revealed that emergent HR sizes were most sensitive to memory (short- and long-term) and scent decay rates. These rate parameters essentially control the decay in effects of modelled mechanisms that act on space use. Short-term memory is a repulsive force pushing the individual to move beyond areas it has visited, where a slower decay results in more exploratory (or diffusive) movement leading to larger HR sizes. On the other hand, long-term memory is an attractive force that attracts the individual to areas, based on either an *informed* memory of previously visited high resource areas, or an *uninformed* expectation that areas beyond the local area of movement (that are unexplored or forgotten) are better. If the animal's expectation for unexplored area matches the mean resource value, faster long-term memory decay rates led to larger HR sizes because the memory expectation was generally better than most individuals' local area of movement (i.e., resource distribution was skewed so the mean was greater than the median value). If unexplored area expectations were low, we expect long-term memory decay to have the opposite negative effect on HR size. Unlike resource memory, the effect of scent is externally conferred by conspecifics. When conspecific scent lingers at slower decay rates, it acts as an external negative pressure limiting HR sizes. While most individuals have smaller HR sizes with slow scent decay rates, a few individuals ranged widely in low resource areas, causing the longer right-tailed skew in overall HR size distribution.

The effects of resource memory and scent on emergent HRs from our model raises an interesting point: HR formation in territorial animals can be attributed to both memory-

mediated philopatric behaviour operating at the individual-level (i.e., inherent exploratory vs.

recursive tendency) and spatial competition at the population-level. Existing theoretical

models have independently demonstrated that distinct and stable HRs can form from either

resource memory-mediated movement (Van Moorter *et al.* 2009; Bracis *et al.* 2015) *or* scent-

or memory-mediated conspecific avoidance (Potts *et al.* 2012; Giuggioli *et al.* 2013; Ellison

et al. 2020) alone. In territorial species, both mechanisms are likely crucial to HR formation.

While site revisits owing to memory processes are commonly observed, it is also common for

animals to 'probe' the boundaries of their HR and expand into newly unoccupied habitat.

Thus, it is important to consider the degree to which each component drives HR formation

within the same modelling framework to understand how territorial animals integrate the

opposing motivators and the relative strengths of each response. Mechanistic home-range

analyses (MHRAs) on carnivores have integrated and disentangled the effects of scent-

mediated conspecific avoidance from other drivers of HR formation (e.g., resource selection

and steep terrain avoidance) in their modelling frameworks (Moorcroft *et al.* 2006; Bateman

et al. 2015). However, the use of memory as mechanistic basis for philopatry (e.g., site

fidelity) within analytical models is still in its' infancy. The analytical framework used in

recent studies to incorporate memory-mediated behaviours (e.g., step selection and

conspecific avoidance; Oliveira-Santos *et al.* 2016; Ellison *et al.* 2020) can plausibly be

extended to memory-mediated philopatry. Integrating both territoriality and memory-

mediated philopatry into MHRAs to uncover key drivers of HR formation is an important

area of future work (Fagan *et al.* 2013).

Despite challenges in in simultaneously fitting multiple metrics to a real system with

our movement model, we successfully approximated the general patterns. The BRT models

of the sensitivity analysis output were able to approximate simulation outcomes and

facilitated parameter calibration to real data using multiple summary metrics. Following

manual calibration, our simulations reproduced the target median and positively skewed variation in HR size of the empirical cat GPS dataset. The underestimate of the upper limit of HR size may be due to the spatial limitation of the truncated simulation landscape (to prevent long simulation times) from the total landscape covered by the tagged feral cats. Nonetheless, we were able to replicate the emergence of wide-ranging individuals (or transients) amidst resident individuals, as seen by the positively skewed variation in individual HR size, reflecting a real phenomenon observed in our data and many other territorial species (Holmes *et al.* 1996; Mitchell *et al.* 2015; Hinton *et al.* 2016). While transience itself is a complex movement phenomenon, its emergence can occur with the core mechanisms of resource memory and territoriality, without modelling further complexities such as individual differences in motivation (e.g., varying exploratory tendencies).

We were only able to partially replicate finer scale spatio-temporal patterns estimated for feral cats (i.e., cumulative HR) for several reasons. First, the less stable simulated cumulative HR curves could indicate that our burn-in period was not long enough for individuals to establish stable HRs and that burn-in time should be considered in parameter calibration. However, the simulated patterns were not unrealistic and have been observed in real systems (e.g., Eurasian lynx, Breitenmoser-Würsten *et al.* 2007; Brazilian mesocarnivores, Bianchi *et al.* 2016). Secondly, we were not able to quantitatively describe the cumulative HR curves. Quantitative characterisation of such curves (including similar metrics such as net squared displacement) is an active area of research that is challenging and remains only partially solved (Spitz *et al.* 2017).

Ultimately, improving resemblance to our real data likely requires additional mechanisms for movement behaviours common in carnivores. For example, the lack of location point clusters and linear location concentrations along linear resources in our simulated data suggests that we did not account for intermittent resting behaviour and

preferential use of corridors respectively (Noss *et al.* 1996). Integrating rest site and corridor

layers into the resource map could potentially improve emergent spatial patterns. However,

the underlying utilisation mechanism is different from modelled behaviours (of continuous

resource consumption), which may warrant building additional mechanisms and landscape

layers into the model (e.g., a resting state dependent on the animal's internal bioenergetic

state; landscape permeability). Additionally, methods that attempt to identify site fidelity and

quantify recursions can potentially be used to parameterise such a model (Richardson *et al.*

2017; Bracis *et al.* 2018a).

Our flexible modelling framework simulates continuous-time movement trajectories

that can be applied to a variety of populations exhibiting cue-based territorial behaviour.

Models can be calibrated and validated against empirical movement data, which is becoming

increasingly available (Kays *et al.* 2015). While further development may be necessary to

replicate some fine-scale movement patterns, our model can be used to simulate generic

territorial space use, in terms of broadly matching HR characteristics. Potential applications

include simulation-based evaluations of behavioural structure in animal movement (e.g.,

Gurarie *et al.* 2016), and population estimation of mobile animals (e.g., Theng *et al.*, In

Review). Another potential application is in animal-mediated seed dispersal research, which

has identified the need to integrate frugivory and disperser movement and to consider more

process-based movement approaches (Nield *et al.* 2019).

In conclusion, we developed a broadly applicable modelling framework that integrates

two key underlying mechanisms of home-range formation, resource memory and

territoriality, to simulate complex patterns of territorial animal space use. Our model

application to a target population of feral cats demonstrated that general space use patterns

evident in the empirical data could be approximated through simulation, and while replication

of finer-scale space use patterns is plausible, we have identified key considerations for future

417 investigations. Our understanding of how animals integrate multiple aspects of spatial

418 localisation is only beginning to emerge, and our modelling approach can provide a

419 foundation for sophisticated theoretical models of space use in interacting animals.

Acknowledgements

M.T. was supported by an Adelaide Postgraduate Scholarship Award (International). This work was supported with supercomputing resources provided by the Phoenix HPC service at the University of Adelaide. Empirical data was from a 2019 feral cat control study conducted by the Department for Environment and Water South Australia.

References

Baker, P.J., Funk, S.M., Harris, S. & White, P.C. (2000). Flexible spatial organization of urban foxes, Vulpes vulpes, before and during an outbreak of sarcoptic mange. *Anim Behav*, 59, 127-146.

Bateman, A.W., Lewis, M.A., Gall, G., Manser, M.B. & Clutton-Brock, T.H. (2015). Territoriality and home-range dynamics in meerkats, Suricata suricatta: a mechanistic modelling approach. *J Anim Ecol*, 84, 260-271.

Bianchi, R.C., Olifiers, N., Gompper, M.E. & Mourao, G. (2016). Niche Partitioning among Mesocarnivores in a Brazilian Wetland. *PLoS One*, 11, e0162893.

Bracis, C. & Wirsing, A. J. (In review). Prey foraging behavior after predator reintroduction is driven by resource knowledge and exploratory tendency. *Frontiers in Ecology and Evolution*.

Borger, L., Dalziel, B.D. & Fryxell, J.M. (2008). Are there general mechanisms of animal home range behaviour? A review and prospects for future research. *Ecol Lett*, 11, 637-650.

Bracis, C., Bildstein, K.L. & Mueller, T. (2018a). Revisitation analysis uncovers spatio-temporal patterns in animal movement data. *Ecography*, 41, 1801-1811.

Bracis, C., Gurarie, E., Rutter, J.D. & Goodwin, R.A. (2018b). Remembering the good and the bad: memory-based mediation of the food–safety trade-off in dynamic landscapes. *Theoretical Ecology*, 11, 305-319.

Bracis, C., Gurarie, E., Van Moorter, B. & Goodwin, R.A. (2015). Memory Effects on Movement Behavior in Animal Foraging. *PLoS One*, 10, e0136057.

Bracis, C. & Mueller, T. (2017). Memory, not just perception, plays an important role in terrestrial mammalian migration. *Proc Biol Sci*, 284.

Breitenmoser-Würsten, C., Zimmermann, F., Molinari-Jobin, A., Molinari, P., Capt, S.,

Vandel, J.-M. *et al.* (2007). Spatial and Social stability of a Eurasian lynx Lynx lynx

population: an assessment of 10 years of observation in the Jura Mountains. *Wildlife*

Biology, 13, 365-380.

Burt, W.H. (1943). Territoriality and Home Range Concepts as Applied to Mammals.

Journal of Mammalogy, 24.

Elith, J., Leathwick, J.R. & Hastie, T. (2008). A working guide to boosted regression trees. *J*

Anim Ecol, 77, 802-813.

Ellison, N., Hatchwell, B.J., Biddiscombe, S.J., Napper, C.J. & Potts, J.R. (2020).

Mechanistic home range analysis reveals drivers of space use patterns for a non-

territorial passerine. *J Anim Ecol*.

Fagan, W.F., Lewis, M.A., Auger-Methe, M., Avgar, T., Benhamou, S., Breed, G. *et al.*

(2013). Spatial memory and animal movement. *Ecol Lett*, 16, 1316-1329.

Giuggioli, L., Potts, J.R., Rubenstein, D.I. & Levin, S.A. (2013). Stigmergy, collective

actions, and animal social spacing. *Proceedings of the National Academy of Sciences*,

110, 16904-16909.

Gurarie, E., Bracis, C., Delgado, M., Meckley, T.D., Kojola, I. & Wagner, C.M. (2016).

What is the animal doing? Tools for exploring behavioural structure in animal

movements. *J Anim Ecol*, 85, 69-84.

Hinton, J.W., Proctor, C., Kelly, M.J., van Manen, F.T., Vaughan, M.R. & Chamberlain, M.J.

(2016). Space Use and Habitat Selection by Resident and Transient Red Wolves

(Canis rufus). *PLoS One*, 11, e0167603.

Holmes, R.T., Marra, P.P. & Sherry, T.W. (1996). Habitat-Specific Demography of Breeding

Black-Throated Blue Warblers (Dendroica caerulescens): Implications for Population

Dynamics. *The Journal of Animal Ecology*, 65.

Kays, R., Crofoot, M.C., Jetz, W. & Wikelski, M. (2015). ECOLOGY. Terrestrial animal

tracking as an eye on life and planet. *Science*, 348, aaa2478.

Lewis, M.A. & Murray, J.D. (1993). Modelling territoriality and wolf–deer interactions.

Nature, 366, 738-740.

Mares, M.A., Watson, M.D. & Lacher, T.E., Jr. (1976). Home range perturbations in Tamias

striatus : Food supply as a determinant of home range and density. *Oecologia*, 25, 1-

12.

McLoughlin, P.D. & Ferguson, S.H. (2016). A hierarchical pattern of limiting factors helps

explain variation in home range size. *Écoscience*, 7, 123-130.

Merkle, J.A., Potts, J.R. & Fortin, D. (2017). Energy benefits and emergent space use

patterns of an empirically parameterized model of memory-based patch selection.

Oikos, 126, 185-195.

Mitchell, N., Strohbach, M.W., Pratt, R., Finn, W.C. & Strauss, E.G. (2015). Space use by

resident and transient coyotes in an urban–rural landscape mosaic. *Wildlife Research*,

42.

Moorcroft, P.R., Lewis, M.A. & Crabtree, R.L. (2006). Mechanistic home range models

capture spatial patterns and dynamics of coyote territories in Yellowstone. *Proc Biol*

Sci, 273, 1651-1659.

Nathan, R., Getz, W.M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D. *et al.* (2008). A

movement ecology paradigm for unifying organismal movement research. *Proc Natl*

Acad Sci U S A, 105, 19052-19059.

Nield, A.P., Nathan, R., Enright, N.J., Ladd, P.G., Perry, G.L.W. & Buckley, Y. (2019). The

spatial complexity of seed movement: Animal-generated seed dispersal patterns in

fragmented landscapes revealed by animal movement models. *Journal of Ecology*,

108, 687-701.

Noss, R.F., Quigley, H.B., Hornocker, M.G., Merrill, T. & Paquet, P.C. (1996). Conservation

Biology and Carnivore Conservation in the Rocky Mountains. *Conservation Biology*,

10, 949-963.

Oliveira-Santos, L.G., Forester, J.D., Piovezan, U., Tomas, W.M. & Fernandez, F.A. (2016).

Incorporating animal spatial memory in step selection functions. *J Anim Ecol*, 85,

516-524.

Potts, J.R., Harris, S. & Giuggioli, L. (2012). Territorial dynamics and stable home range

formation for central place foragers. *PLoS One*, 7, e34033.

Potts, J.R., Harris, S. & Giuggioli, L. (2013). Quantifying behavioral changes in territorial

animals caused by sudden population declines. *Am Nat*, 182, E73-82.

Potts, J.R. & Lewis, M.A. (2014). How do animal territories form and change? Lessons from

20 years of mechanistic modelling. *Proc Biol Sci*, 281, 20140231.

Powell, R.A. & Mitchell, M.S. (2012). What is a home range? *Journal of Mammalogy*, 93,

948-958.

Prowse, T.A.A., Bradshaw, C.J.A., Delean, S., Cassey, P., Lacy, R.C., Wells, K. *et al.*

(2016). An efficient protocol for the global sensitivity analysis of stochastic

ecological models. *Ecosphere*, 7.

R Core Team (2020). R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing Vienna, Austria.

Ranc, N., Cagnacci, F., Moorcroft, P.R. & Gill, J. (2020). Knowing your neighbours: How

memory-mediated conspecific avoidance influences home ranges. *Journal of Animal*

Ecology, 89, 2746-2749.

Ranc, N., Moorcroft, P.R., Ossi, F. & Cagnacci, F. (2021). Experimental evidence of

memory-based foraging decisions in a large wild mammal. *Proc Natl Acad Sci U S A*,

118.

Richardson, T.O., Giuggioli, L., Franks, N.R. & Sendova-Franks, A.B. (2017). Measuring

site fidelity and spatial segregation within animal societies. *Methods Ecol Evol*, 8,

965-975.

Riotte-Lambert, L., Benhamou, S. & Chamaille-Jammes, S. (2015). How memory-based

movement leads to nonterritorial spatial segregation. *Am Nat*, 185, E103-116.

Šálek, M., Drahníková, L. & Tkadlec, E. (2014). Changes in home range sizes and population

densities of carnivore species along the natural to urban habitat gradient. *Mammal*

Review, n/a-n/a.

Santangeli, A., Hakkarainen, H., Laaksonen, T. & Korpimäki, E. (2012). Home range size is

determined by habitat composition but feeding rate by food availability in male

Tengmalm's owls. *Animal Behaviour*, 83, 1115-1123.

Schoepf, I., Schmohl, G., König, B., Pillay, N. & Schradin, C. (2015). Manipulation of

population density and food availability affects home range sizes of African striped

mouse females. *Animal Behaviour*, 99, 53-60.

Schradin, C., Schmohl, G., Rödel, H.G., Schoepf, I., Treffler, S.M., Brenner, J. *et al.* (2010).

Female home range size is regulated by resource distribution and intraspecific

competition: a long-term field study. *Animal Behaviour*, 79, 195-203.

Spencer, W.D. (2012). Home ranges and the value of spatial information. *Journal of*

Mammalogy, 93, 929-947.

Spitz, D.B., Hebblewhite, M. & Stephenson, T.R. (2017). 'MigrateR': extending model-

driven methods for classifying and quantifying animal movement behavior.

Ecography, 40, 788-799.

Theng, M., Milleret, C., Bracis, C., Cassey, P. & Delean, S. (2021). Confronting spatial

capture-recapture models with realistic animal movement simulations. *Ecology*.

Van Moorter, B., Visscher, D., Benhamou, S., Börger, L., Boyce, M.S. & Gaillard, J.-M.

(2009). Memory keeps you at home: a mechanistic model for home range emergence.

Oikos, 118, 641-652.

Table 1. Overview of simulation parameters used in all four sets of simulations in the study. In theoretical example 1 (TE1), we tested the impacts of resource memory and territoriality on emergent home-ranges across three scenarios (territoriality only, resource memory only, territoriality and memory) and densities ($n = 5, 10, 15$). In theoretical example 2 (TE2), we test the impact of population manipulation by removing four out of ten individuals mid-simulation on emergent space use, and compare it with a control scenario without individual removal. In the sensitivity analysis (SA), we assessed the relative influenced of nine parameters across a range of values (grey cells). Finally, we ran simulations with a set of parameters calibrated to the empirical feral cat data (Cat). Units are arbitrary in the simulations, L is used for generic length units and T is used for generic time units.

Parameter	Definition	Units	TE1	TE2	SA	Cat
Simulations						
Δt	Model time step	T	1	1	1	1
T	Simulation length (time steps)	T	25,000	50,000*	21,600	21,600
	Burn-in (time steps)	T			5,040	5,040
n	Number of individuals		5, 10, 15	10	30	30
	Number of individuals removed		-	4 (removal),	-	-
				0 (control)		
Landscapes						

–	Dimensions	L	200 × 200	200 × 200	400 × 600	400 × 600
Consumption						
β_R	Resource regeneration rate	1/T	0.01	0.01	10^{-3}–10^{-1}	0.01
β_C	Consumption rate	1/T	0.5	1	1	1
γ_C	Consumption spatial scale	L	1	1	1	1
Resource memory†						
ϕ_L	Long-term memory decay rate	1/T	10^{-5}	10^{-6}	10^{-7}–10^{-3}	10^{-7}
ϕ_S	Short-term memory decay rate	1/T	0.01	0.01	10^{-4}–10^{-1}	0.005
Ψ_M	Short-term memory factor		2	2	2	2
β_L, β_S	Learning rates	1/T	1	1	1	1
γ_L, γ_S	Learning spatial scale	L	1	1	1	1
γ_Z	Memory spatial scale foraging	L	10	10	10	10
$M*$	Memory expectation (value uninformed)		5e-5	5e-4	0–2e^{-4}	5e-5
	Memory full informed?	Boolean	True	True	True	True
Scent behaviour						

	Switched on?	Boolean	True, False, True	True	True	True
β_D	Deposition rate	1/T	3	3	0–5	3
γ_D	Deposition spatial scale	L	1	1	1	1
ϕ_D	Decay rate	1/T	10^{-5}	10^{-4}	10^{-6}–10^{-2}	10^{-6}
Ψ_D	Response strength		2	2	2	2
γ_w	Response spatial scale	L	2	2	0.5–10	3
Movement‡						
	Type	Categorical	Kinesis, Memory, Memory	Memory	Memory	Memory
τ_S, τ_F	Autocorrelation time scale	T	4, 2	4, 2	4, 2	4, 2
v_S	Search speed	L/T	2	2	1–6	5
v_F	Feed speed	L/T	0.2	0.2	0–0.5	0.05
λ	Mean time to update θ	T	1	1	1	1

†L = long-term resource memory, S = short-term resource memory
‡S = searching, F = feeding.

Figure 1. A graphical representation of the movement model. a) An overview of all the

processes involved. The individual interacts with the external environment through a series of

actions, which serves to update its' internal state and ultimately influences movement

decisions. b) The movement process is formulated on continuous time, where the updating of the individual's direction and behavioural state are independent processes. Behavioural state ultimately determines the processes used in selecting the next direction (represented by an angular probability density function of memory and scent) and the speed/correlation values. The processes in (a) represented in (b) are highlighted in colours.

Figure 2. Simulation results from three scenarios (territoriality, memory, territoriality and memory): (a–c) sample realisations of animal movement trajectories from each scenario ($n =$ 10), where individual trajectories start at random (colours represent individuals); relationship

between HR size and (d) mean resource value within HR (T: territoriality, M: memory, T +

M: territoriality + memory) and (e) density (n = 5, 10, 15) from 100 replicates.

Figure 3. Aggregate home-range (HR) overlap between removed and remaining individuals

before (Phase 1; $t = 5{,}001$–$20{,}000$) and after removal (Phase 3; $t = 35{,}001$–$50{,}000$) in the

control and removal scenarios (100 replicates each). Beneath each scenario is a sample realisation of movement trajectories in Phase 1 (top panel) and 3 (bottom panel) of the control (left panel) and removal (right panel) simulation. Individual trajectories decrease in opacity according to time. Cool colours represent remaining individuals, warm colours represent removed individuals (in the control scenario, individuals were randomly grouped).

Figure 4. Results from the sensitivity analysis. (a) Relative influence (%) of nine model input parameters on six emergent movement summary

statistic from our boosted regression tree (BRT) analysis. Partial dependency plots showing effects of the top three contributing parameters on

(b) home-range size, (c) home-range size skew and (d) home-range overlap. Partial dependency plots show the effect of a given explanatory

variable on each home-range metric, while holding the effects of other explanatory variables at their average. We used 500 bootstrap replicates to calculate the 95% confidence intervals, depicted in grey. Note that several of the x-axes are on a $\log_{10}$ scale. Full plots showing partial effects of all parameters on each summary statistic can be found in **Fig. S3–S9**.

Figure 5. A comparison of movement patterns from parameterised simulations and the

empirical data: a) frequency distribution of HR size of simulated individuals obtained from

10 replicates bootstrap sampled 10 times each (samples represented as histograms in translucent black fill) to match the empirical sample of 11 tagged feral cats in Kangaroo Island, South Australia (histogram with empty fill and red border); b) one realisation of cumulative HR size over time of 11 randomly selected individuals (left) against that of the feral cats (right) and their; c) corresponding movement locations (left) vs. movement locations from feral cats (right; note that some cat locations go beyond the landscape shown here).