**Title**

Adapting to extremes: reconstructing niche evolution in Australian *Acacia* over time and

space.

Authors: Xia Hua[1,2*], Marcel Cardillo[1], Lindell Bromham[1]

[1]Macroevolution and Macroecology, Research School of Biology, Australian National

University, Canberra ACT 0200 Australia

[2]Mathematical Sciences Institute, Australian National University, Canberra ACT 0200

Australia

**Abstract**

Macroevolutionary analysis is increasingly being used to study biodiversity responses to

climate change, for example by using phylogenetic node ages to infer periods of

diversification, or phylogenetic reconstruction of traits to infer adaptation to particular

stresses. Here we apply a new macroevolutionary method to investigate the responses of a

diverse plant genus, *Acacia*, to increasing aridity and salinity in Australia from the Miocene

to the present. The Niche Evolution Model (NEMo) combines environmental niche

modelling with phylogenetic comparative methods in a single statistical framework, to

estimate current environmental tolerances, reconstruct the history of niche evolution and infer

rates of change in key aspects of environmental tolerance. Using a large database of *Acacia*

occurrence records and presence-absence survey sites, we find that both spatial and temporal

patterns in niche evolution of *Acacia* are consistent with the aridification history of Australia

and suggest high niche lability along both axes, which has allowed *Acacia* to quickly exploit

new niches created during the aridification of the continent, and resulting in their current

dominance of many habitats across Australia. This study demonstrates that phylogenetic

studies of niche evolution can move beyond application of simple trait-based models,

allowing the underlying processes of speciation, adaptation and dispersal to be explicitly

modelled in a macroecological and macroevolutionary context.

Keywords: aridity tolerance, salt tolerance, aridification, niche lability, Environmental Niche

Modelling, Phylogenetic Comparative Methods.

**Introduction**

The climatic and environmental history of Australia during the Neogene and Quaternary

Periods has been a story of increasing aridification. Dry climates may have existed from the

Eocene (Carpenter et al 2014), but evidence from charcoal and pollen records reveals an

increase in fire frequency and prevalence of fire-adapted, scleromorphic plant taxa, pointing

to an expansion of the arid zone from the early Miocene onwards (Byrne et al 2008). As the

arid zone expanded, previously widespread mesic forest habitats contracted to the coastal

margins, and plant groups responded in different ways (Byrne & Murphy 2020, Crisp et al

2004, Weston & Jordan 2017). Some groups (such as *Nothofagus*) remained confined to

humid forest conditions and contracted with these habitats to small coastal refuges. Other

groups responded to new ecological opportunities provided by the expanding arid zone, and

diversified into numerous xeromorphic lineages. Some plant families, especially Proteaceae,

Myrtaceae and Fabaceae, contain both mesic and xeromorphic lineages. Even some large

genera, such as *Acacia, Grevillea* and *Hakea,* have both mesic and xeromorphic

representatives. Plant groups such as these, in which close relatives occupy a range of

ecoclimatic conditions, provide good case studies for studying the evolution of tolerance to

extreme environmental conditions.

The expansion of the arid zone in Australia led to increasingly challenging

environments characterized by stressful conditions for plant growth, including extreme

drought, low soil fertility, frequent fires and high soil salinity. At the same time, areas that

are extreme for most plants represent an opportunity for lineages to specialize, escape

competition, and diversify. Arid-adapted lineages evolved a range of strategies to cope with

water stress, broadly classified into escape, avoidance, and tolerance (Delzon 2015). We see

all three strategies in *Acacia*, such as phyllode shedding (escape), scleromorphic structures

like sunken stomata or needle-shaped phyllodes (avoidance), and physiological adaptations

such as low leaf water potential (tolerance: Moore 2013). The wide range of mechanisms

adopted by *Acacia* species to cope with water stress are reflected in their geographic

distributions and varied climatic niches.

To understand how opportunities provided by the drying environments shaped

present-day distribution and diversity, we need to examine the dynamics of lineage

adaptation and diversification in response to extreme conditions. Macroevolutionary and

macroecological analyses based on phylogenies have provided a variety of ways of

investigating adaptation to harsh environments (Bromham et al. 2020). For example, the ages

of nodes in molecular phylogenies have been used to infer bursts of diversification associated

with periods of drying in *Acacia* (Renner et al. 2020), and the movement of rainforest-

adapted lineages into dry environments (Crayn et al. 2006). In addition to relative timing of

diversification, phylogenies can provide information on evolution of specific adaptations to

extreme conditions. For example, Crisp et al. (2011) inferred the phylogenetic history of

traits such as post-fire epicormic resprouting to investigate the timing of adaptation of

*Eucalyptus* to increasing fire frequency. Jordan et al. (2008) reconstructed the evolution of

stomatal protection structures (which are adaptations to dry climates) in Proteaceae to show

that such structures evolved relatively few times in the ancestors of dry-climate clades. As

well as physiological and morphological traits, phylogenetic approaches have been applied to

reconstructing the evolution of environmental niches. For example, by comparing alternative

evolutionary models of climatic niche evolution for clades from different regions, Skeels and

Cardillo (2017) showed that within four large genera (*Protea, Moraea, Banksia*, and *Hakea*),

different lineages are evolving towards different climatic optima in different bioclimatic

regions. Onstein et al. (2016) used the same approach to show that Proteaceae lineages in

open vegetation and closed forest also have different inferred climatic optima.

Phylogenetic analyses such as these are useful, but have a number of limitations. To

permit the application of standard stochastic models of trait evolution, such as Brownian

Motion (BM) or Ornstein-Uhlenbeck (OU), in most macroevolutionary studies, the

environmental niche is simplified to a point-estimate for a species, for example by taking the

mean value across a species distribution (Evans et al 2009; Kozak & Wiens 2010;

Münkemüller et al 2015; Renner et al 2020). But this is problematic for two main reasons.

First, it averages away most of the potentially informative variation in environmental

conditions across the range of a species. Second, models of trait evolution such as BM may

not be appropriate for modelling the evolution of environmental tolerance. In particular,

applying trait-evolution models to current environmental data to infer ancestral niches (e.g.,

Renner et al 2020, Skeels & Cardillo 2017) assumes that environmental traits such as aridity

or temperature evolve randomly and independently along branches for each lineage. This

assumption may often be violated because environmental niche evolution is likely to be

driven by climatic changes, which most likely apply simultaneously and directionally across

all lineages within a given region.

More fundamentally, phylogenetic analyses of niche evolution with changing climate

compartmentalise the inference of contemporary niche (inferred from occurrence data) and

the history of niche evolution (inferred from phylogenetic models). These are treated as two

separate problems, and subject to independent analyses with separate statistical frameworks.

But contemporary niches are the product of macroevolutionary processes and inferring these

processes requires accurate inference of the niche from contemporary distributions, so the

two are fundamentally linked. The logical step, then, is to link their analyses too.

Here, we infer the history of adaptation to extreme environments by applying a novel

method, the Niche Evolution Model (NEMo), that combines Environmental Niche Modelling

(ENM) and reconstruction of niche evolution in a single statistical framework. This method

uses occurrence records to characterise the distribution of each tip species in a phylogeny,

and reconstructs niche evolution by explicitly modelling key driving processes: (1) shift in

areas of tolerable conditions, (2) adaptation of lineages to changing conditions, (3) dispersal

away from areas that are no longer suitable and into accessible areas of suitable habitat, (4)

speciation, and (5) divergence along lineages' environmental tolerances. A strength of this

method is that it allows us to infer the placement of these key processes on the phylogeny and

to reconstruct ancestral niches at any time point, without making the unrealistic assumption

that niches evolve under a simple stochastic trait-based model. Unlike previous approaches,

NEMo can infer the occurrence density of these processes of niche evolution on the

phylogeny, so it allows us to directly compare the timing of niche shift to periods of climatic

change, or compare the timing of niche evolution along different environmental axes, using

standard statistical tests, without pre-defining the covariance structure to account for

phylogenetic nonindependence. Information on paleoclimate can either be incorporated into

the model as prior information, or can be held aside as independent information against

which to test the robustness of the inference.

We use NEMo to investigate macroevolutionary responses of the Australian plant

genus *Acacia* to aridification. *Acacia* is one of the largest and most widely-distributed plant

genera in Australia, with over 900 described species occupying nearly all major habitat types,

often as dominant taxa. Many *Acacia* species are known to be tolerant of a range of extreme

environmental conditions, and several previous studies have used *Acacia* species as case

studies for understanding tolerance to extremes, including salinity and other aspects of soil

geochemistry (Bui et al. 2014a, 2014b), and water stress (Atkin et al. 2002). There has been

rapid progress in understanding the evolutionary history of *Acacia* in recent years (Maslin et

al. 2003; Murphy et al. 2010; Mishler et al. 2014). Nearly all *Acacia* species are conspicuous,

woody plants, so spatial occurrence records for *Acacia* are abundant and widespread. We

combine a curated database of *Acacia* occurrence records with extensive survey data, along

with predictors of sampling bias, to characterise the current distribution of species in the

phylogeny. We use NEMo to reconstruct the evolution of adaptation to increasing

aridification in Australian *Acacia,* focusing on niche evolution in response to increased

aridity and changes in the distribution and severity of soil salinity (frequently associated with

aridification (Gilkes et al. 2003). Specifically, we ask two questions. First, have periods of

rapid niche evolution along the aridity and salinity axes corresponded with major periods of

aridification in Australia? Second, on which lineages did adaptation to increased aridity and

salinity occur? Answering these two questions may help us understand whether *Acacia*

responded to continental aridification by a burst of adaptation, allowing the genus to radiate

through the arid zone.

**Methods**

*1. Phylogenetic and Spatial Data*

We based our analysis on a published phylogeny of 508 *Acacia* species, representing half of

species currently recognized (Mishler et al. 2014). We timescaled branch lengths under a

relaxed clock model using the R package ape (Paradis and Schliep 2018), with the root date

constrained between 21 and 23.9 million years ago (the estimated range of crown divergence

dates (Miller et al. 2013).

To describe the current environmental conditions under which each species is found,

we used a published dataset of 132,295 carefully checked presence locations of these 508

*Acacia* species, filtered from a total of 750,000 *Acacia* records from the Australian Virtual

Herbarium (Misher et al. 2014). We also assembled species presence-absence data from

floristic survey databases for New South Wales (BioNet Systematic Flora Survey;

environment.nsw.gov.au/research/VISplot.htm), Victoria (Victorian Biodiversity Atlas flora

records; environment.vic.gov.au/biodiversity/victorian-biodiversity-atlas), and Western

Australia (Flora Surveys of the Yilgarn; naturemap.dbca.wa.gov.au). We only included data

from complete floristic surveys conducted later than 2005 (to ensure consistency in

taxonomy) under government custodianship (to ensure consistent standards of survey

quality). As a result, our presence-absence survey data included 27,850 survey sites across

three states of Australia. Survey data are particularly valuable because they represent

presence/absence data, not just opportunistic presence records, and are not subject to the

same sampling biases as atlas or herbarium occurrence records. These presence-absence data

were analysed together with the presence locations. To represent degree of aridity and

salinity at each presence location for each species and each survey site, we used information

from the geographic layer of maximum monthly aridity index (Williams et al. 2012) and the

geographic layer of root zone soil electrical conductivity (Bui et al. 2017).

The majority of occurrence records for most species are presence-only records,

typically collected opportunistically, so their spatial distributions tend to be biased towards

regions that are more easily accessible to researchers, especially along road networks, and

areas of particular interest, such as protected areas or hotspots of diversity (Fithian et al.

2015; Haque et al. 2017). To correct for this sampling bias, we included two additional

factors: distance to the nearest road (using data from www.ga.gov.au/geoscience/national)

and whether it is in a protected area, such as a conservation reserve or national park (from

www.environment.gov.au/land/nrs/science/capad).

*2. Niche evolution model*

The Niche Evolution Model (NEMo) uses a Bayesian framework that includes three major

components: model of niche evolution, phylogenetic comparative method (PCM), and

ecological niche model (ENM). NEMo models species distributions in niche space, not

geographic space. The aim is to infer not only species' niches at the contemporary tips of the

phylogeny, but also to estimate the amount of change in niches along every branch of the

phylogeny. This method has been shown to be effective at reconstructing past periods of

niche evolution and better than standard ENMs at predicting aspects of contemporary niches

(see Hua et al. submitted for details, available on bioRxiv).

The niche evolution component calculates how three aspects of niche (fundamental,

available, realized) change under three basic processes (speciation, adaptation, dispersal).

"Fundamental niche" represents the range of conditions that each species can tolerate at a

given point in time, expressed as a range of values along a relevant environmental axis. We

expect species to evolve along this environmental axis over evolutionary time, through the

processes of adaptation to new conditions (by selecting for individuals at the extreme values

of the distribution) and through speciation along environmental gradients (so that a species

splits along its tolerance range leaving one daughter species that is more skewed toward the

extreme conditions). "Available niche" describes the range of conditions in areas that are

accessible to the species, regardless of whether the species can persist in those conditions.

The potential of the species to occupy the available niche is governed by its fundamental

niche and dispersal ability. We also include barriers to dispersal as nuisance parameters,

which could represent physical or biotic barriers. "Realized niche" consists of the locations

where a species is found, which is shaped by the suitability of conditions and the opportunity

to disperse to and occupy those areas.

The PCM part of the model infers the locations on the phylogeny of adaptation and

speciation events, assuming these events are distributed along the branches of the phylogeny

according to a Poisson process. Dispersal in niche space is modelled as a continuous process

between any two events (adaptation or speciation), or nodes of the phylogeny. The

distribution of these events on the phylogeny gives a possible history of niche evolution.

Given this history, we use the model of niche evolution to calculate the three aspects of niche

of each tip species. The ENM component of the method then assesses the fit of the inferred

niches to the presence and/or absence locations of the tip species. The fit of the inferred

niches to the data then feeds back to our inferences of the history of niche evolution and the

niches of tip species. Details of the method, and description of tests of its performance and

reliability, are given in Hua et al. (submitted), and the code with step-by-step instructions for

implementing the model can be found at

https://github.com/huaxia1985/NEMo. Below we briefly describe the parameters of the

model applied to this case study.

*3. Model Parameters*

NEMo has two universal parameters which are assumed to be constant across the phylogeny:

adaptation rate and dispersal rate. Additional parameters that are specific to each adaptation,

speciation, and dispersal events are: 1) amount of environmental change in an adaptation

event, 2) proportion of a tolerance range that is inherited by one descendent species (its sister

species inherits the other part) from the ancestral species in a speciation event, 3) where a

dispersal barrier locates along the environmental axis during a dispersal process.

We apply separate NEMo analyses to the evolution of environmental niches in *Acacia*

along aridity and along salinity. For each of these niche axes, the values for the available

niche and the fundamental niche at the root of the phylogeny were assumed to follow normal

distributions. The starting values for their means and standard deviations were set to their

maximum likelihood estimates given the likelihood of each tip species' presence-only data

and presence-absence data, assuming that no events of niche evolution occurred on the

phylogeny. The starting values for adaptation rate and dispersal rate were arbitrarily set to

0.03 and 1, respectively. Dispersal rate had rather flat prior, an exponential distribution with

rate of 0.1. The parameters for the niches at the root and the adaptation rate had relatively

narrow priors, a lognormal distribution with logarithmic mean of their initial values and

logarithmic standard deviation of 0.5. The narrow priors for the root niches were used to

ensure that root niches do not exceed observed maximum and minimum values (Williams et

al. 2012; Bui et al. 2017) too often during MCMCs. The narrow prior for adaptation rate was

used to ensure a finite likelihood value, as parameters of the fundamental niche change

proportionally to adaptation rate on a log scale, so there is a narrow range of adaptation rate

that gives finite likelihood values. The starting value for the occurrence rate of adaptation

events and the occurrence rate of speciation events was arbitrarily set to 0.1. A narrow prior

was applied on both occurrence rates, an exponential distribution with rate of 10 to reduce the

degree of overestimation on the number of events (see Hua et al. submitted). For both aridity

and salinity, we ran 20 independent MCMCs for $1 \times 10^6$ generations, with a thinning interval

of 1000 and the first 600,000 generations discarded as burnin.

***4. Accounting for sampling bias***

We used an inhomogeneous Poisson point process (Renner et al. 2015) as the ENM

component of NEMo. Because opportunistic sampling bias can influence the niche estimation

250 (Fithian et al. 2015), we modified the likelihood function to account for sampling bias, using

251 the method of Fithian et al. (2015). The log-likelihood of the presence-only data of each tip

252 species is modified as,

253
$$\ell_P = \sum_{i \in I_{PO}} \left[ \log(\lambda_{r_i}) + \delta z_i \right] - \frac{\mathcal{D}}{N} \sum_{i \in I_{BG}} \lambda_{f_i} \lambda_{a_i} e^{\delta z_i}$$

254 , where $z_i$ are values of the factors that influence sampling effort at the $i^{\text{th}}$ location and $\delta$ are

255 the coefficients of the two sampling factors we included - distance of the location to the

256 nearest road and whether the location is in a protected area - which are assumed to have the

257 same value for all tip species. The starting values of coefficients were set to their maximum

258 likelihood estimates given the likelihood function, and a rather flat prior was used: an

259 exponential distribution with rate of 0.1. In the second term of the equation, $\lambda_f$ is the inferred

260 fundamental niche of a tip species, giving the proportion of individuals of the species that can

261 tolerate a certain level of aridity and salinity, and $\lambda_a$ is the inferred available niche of the tip

262 species, giving the proportion of areas accessible to the species that have certain level of

263 aridity and salinity. If we were not accounting for sampling bias, the second term of the

264 equation would be a direct integral of $\lambda_f \lambda_a$ (Hua et al. submitted), but because each location

265 now has a specific value for sampling bias, we can only approximate the integral by

266 randomly sampling $N$ locations from the surroundings of the species presence locations to

267 construct a set of background locations $I_{BG}$ and averaging the term to integrate over all the

268 background locations. Note that these background locations are only used to numerically

269 approximate the second term of the equation, so they are not pseudo-absence locations

270 (Fithian et al. 2015).

271  For each *Acacia* species, background locations were sampled uniformly from areas

272 within an arbitrary large radius (20 *km*) of each presence location of the species, so the

273 weight of each background location is $\frac{\mathcal{D}}{N}$, where $\mathcal{D}$ is the total area that falls within the 20 *km*

radius of each species presence location, and $N$ is the total number of background locations,

which was ten times the number of the species presence locations, with a minimum of 1000

for species with few presence locations. Based on the environmental conditions over all the

background locations, we used Sturges' formula (Sturges 1926) to decide the bin size to

discretise the aridity index and the electrical conductivity.

*5. Estimating rate and amount of niche evolution*

The amount of niche evolution per branch is calculated as the sum of the differences in the

mean of the realized niche before and after each event (both adaptation and speciation events)

inferred along each branch of the phylogeny over the posterior samples, calculating both the

net amount (positive or negative) and the absolute amount. The net amount of niche evolution

divided by branch length gives the direction of niche evolution, and the absolute amount of

niche evolution divided by branch length gives the rate of niche evolution. Last, we

calculated the intensity of the inferred environmental change during adaptation events at each

time point over the whole phylogeny and compared the distribution of this intensity over time

to the aridification history of Australia (see details in Hua et al. submitted).

**Results**

*1. Spatial Patterns of Niche Evolution*

The *Acacia* phylogeny is characterized by a mixture of large, young clades (recent, rapid

divergences), and smaller, older clades (longer-persisting lineages: Figure 1). Three recent

radiations, each around 1 Ma old, include most present-day species in Northern Australia,

Southern Australia, and Eastern Australia. Of these, the Northern Australian clade has the

highest average rate of niche evolution on both aridity and salinity axes (Figure 1). The

smaller, older clades are mainly in Western Australia, with many extant species originating

earlier than the three large radiations. These smaller older clades have, on average, much

lower rate of niche evolution on both aridity and salinity axes than the recent radiations

(Figure 1). Present-day species that occupy regions that are arid and prone to high salinity

(deserts and Mediterranean ecosystems) are found throughout the phylogeny, forming small

clusters of related species. In contrast, present-day species that occupy the mesic zone are

found mainly in young clades as large clusters of closely-related species.

The net amount of niche evolution inferred by NEMo along the phylogeny is shown in Figure

1. Most of the evolutionary change on both aridity and salinity axes has been pushing species

towards tolerance of the extreme conditions. We can see this in the patterns of niche

evolution along the aridity axis in different habitat types (Figure 2A). The largest changes in

niche are in lineages currently found in the most arid environments, while little or no niche

shift is in lineages currently distributed in forests. The same pattern is repeated over the

whole phylogeny, and also within each of the recent radiations (Figure 2A). Similarly, on the

salinity axis, NEMo inferred the greatest amount of niche evolution towards higher salinity

leading to species currently distributed in the Mediterranean ecosystems (Figure 2B). The

same pattern is inferred in the three recent radiations, where there is little change along the

salinity axis in lineages currently distributed in forests (Figure 2B).

*2. Temporal Patterns of Niche Evolution*

On the aridity axis, the inferred intensity of environmental change during each adaptation

event peaks at 13 Ma and near the present (after 2.5 Ma), corresponding to major drying

periods of Australia (Figure 3). On the salinity axis, the inferred intensity of environmental

change during each adaptation event peaks near the present (after 2.5 Ma), with two

additional lower peaks around 10 Ma and 17 Ma (Figure 3).

Across the phylogeny, the net amount of inferred niche evolution due to both

adaptation and speciation events along the salinity and aridity axes showed a significant

positive correlation (Figure 4; Kendall's rank correlation coefficient=0.25, $p<0.001$).

Although significant, the correlation is mostly driven by niche evolution along the terminal

branches, and there is a great deal of scatter, with relatively more aridity evolution on tip

branches leading to present-day species found in the arid zone (Deserts & Xeric Shrublands),

and relatively more salinity evolution on tip branches leading to present-day species found in

Mediterranean ecosystems. The greater amounts of aridity and salinity evolution along the tip

branches are not due to longer branch length, as NEMo also infers large amount of aridity

and salinity evolution along very short tip branches (Figure 4).

**Discussion**

Aridification has been the prevailing environmental trend across most of the Australian

continent through the Neogene and Quaternary. In order to understand the history underlying

Australia's current plant diversity, we need effective methods for reconstructing the

evolutionary responses of plant clades to the environmental changes brought about by

aridification. In this study, we use the large, widespread genus *Acacia* as a case study to

reconstruct its history of niche evolution along aridity and salinity axes, using the newly

developed method NEMo. We reconstruct both spatial and temporal patterns in niche

evolution of *Acacia* and show (1) a preponderance of evolutionary change in niches towards

more extreme values of aridity and salinity, (2) a concentration of niche evolution in the

ancestors of species currently found in arid and Mediterranean regions in both the old clades

and the recent radiations, (3) peaks in niche evolution throughout the aridification history of

Australia, (4) correlated evolution of aridity and salinity tolerance, particularly along tip

branches. As we explain below, these results are consistent with the adaptation of *Acacia*

lineages to drier climates, allowing them to exploit new niches opening up through the

aridification of the continent. Our results point to an evolutionary flexibility in *Acacia* that is

likely to have played a key role in their current near-ubiquitous distribution and dominance of

many habitats across Australia.

The preponderance of niche evolution towards more extreme values of both aridity

and salinity suggests that aridification has been a dominant driver of *Acacia* evolution and

diversification across Australia during the Neogene & Quaternary. The concentration of

niche evolution on branches leading to species currently found in more arid and salt-affected

regions, in both old and recent clades, suggests that the recent radiations of *Acacia* were not

triggered by adaptation to aridity and salinity. Instead, extant species occupying arid and salt-

affected regions are likely to have evolved from ancestral lineages in mesic areas which do

not show high rates of niche evolution along these axes. This interpretation indicates

evolutionary flexibility in *Acacia*, which is supported by the high rate of niche evolution

along both aridity and salinity axes in the three recent radiations. This enhanced rate of niche

evolution was also found by a recent study of diversification in *Acacia* using traditional

PCMs, which hypothesized that Pleistocene glacial-interglacial climate cycling ($< 2.5$Ma)

drove rapid climatic adaptation, which promoted diversification and led to large radiations

(Renner et al. 2020).

Unlike traditional PCMs, NEMo is able to identify key events of niche evolution

along the *Acacia* phylogeny. We found that most changes in niche along the aridity axis were

in the recent radiations, predominantly after 1Ma and along tip branches. But niche evolution

in aridity was also inferred along two branches ancestral to all these radiations (marked with

asterisk in Figure 1), between 12 to 15 Ma and around 5Ma (Figure 1). It is likely that the

niche evolution events that occurred along the two ancestral branches left descendant

lineages of *Acacia* able to tolerate higher aridity than the other lineages in Tropical and

Temperate forests and grasslands during the Pleistocene, so that *Acacia* was able to replace

many other groups during the drying period of the Pleistocene. These two pulses of niche

evolution along the aridity axis are also consistent with the aridification history of Australia.

*Acacia* emerged during a period of aridification (28-23Ma), and we found peaks of

adaptation to increasing aridity in *Acacia* during the two subsequent drying periods,14-5Ma

and after 5Ma (Byrne et al. 2008).

Comparing salt tolerance to aridity tolerance in *Acacia*, we found that the amount of

niche evolution along aridity and salinity axes was highly correlated across branches of the

*Acacia* phylogeny. This result is consistent with the observation that saline soils are

frequently associated with aridity, so many species are adapted to both aridity and salinity

(Gong et al. 2017). Correlated evolution was also suggested between salinity and alkalinity

tolerance in *Acacia* using phylogenetic independent contrasts (Bui et al. 2014b). The NEMo

method offers a way to account for phylogenetic nonindependence in testing correlated

evolution, without making *a priori* assumptions about the phylogenetic covariance matrix. In

addition, the NEMo method allows us to detect branches that have large amounts of niche

evolution along both aridity and salinity axes. These branches are mostly terminal branches

scattered across the *Acacia* phylogeny and are mostly ancestral to species currently

distributed in areas with both high aridity and high salinity. This result is consistent with high

niche lability along both aridity and salinity in Australian *Acacia*, as suggested by the high

rate of niche evolution in the three radiations.

Inferred peaks of increased salinity are on recent lineages, which is consistent with

previous findings of a phylogenetically "tippy" distribution of salt tolerant species in grasses

(Bennett et al. 2013; Moray et al.  2015). Consequently, even if salt tolerance has evolved

many times through old adaptation events, these events rarely gave rise to persistently salt

tolerant lineages of large clades of related salt tolerant taxa, leaving less phylogenetic signal

for us to reconstruct adaptation events in the deep part of the phylogeny than adaptation

events near the tip of the phylogeny (Bromham 2016). The high niche lability in Australian

*Acacia* further agrees with previous hypothesis that the tippy distribution of salt tolerant

species is due to high niche lability (Bromham et al. 2016).

It is important to note that for the NEMo analysis, as for most macroevolutionary

analyses, we have assumed that the phylogeny has accurate topology and provides an

approximate scaling of evolutionary events in time. Clearly, phylogenetic estimates of either

topology or the timing of evolutionary events are not error-free (Bromham et al. 2018;

Guindon 2020; Nie et al. 2019). In particular, uncertainty or bias in branch length estimates

arising from imperfect characterisation of molecular rates over time may have a non-trivial

impact on macroevolutionary inference (Duchêne et al. 2017). Phylogenies such as this one

where the only calibration is a root date may be particularly vulnerable to errors in position of

node heights in some parts of the tree (Duchêne et al. 2014). One possible solution is to apply

the NEMo analysis on posterior samples of phylogenies, but perhaps a better advance will be

to embed the NEMo method into phylogenetic reconstruction, which allows us to directly

link niche evolution events to diversification events (Hua et al. submitted). Another issue

with our current inference of niche evolution in *Aacia* is incomplete sampling, as the *Aacia*

phylogeny only includes half of the known extant species. One obvious solution is to sample

more species, but another way to increase the amount of information in the analysis of niche

evolution is to incorporate historical data, such as paleoclimate, as priors for the occurrence

density of niche evolution events, to supplement current species distribution data (Hua et al.

submitted). Here we have used only part of the available information in the inference process

in order to demonstrate the utility of the method (so we can compare our inferred

evolutionary history to known paleoclimatic history), but the method allows for all available

information to contribute to stronger inference.

**References**

Atkin O., Schortemeyer M., McFarlane N., Evans J.R. 2002. Variation in the components of

relative growth rate in 10 *Acacia* species from contrasting environments. Plant Cell

Environ. 21:1007–1017.

Bennett T.H., Flowers T.J., Bromham L. 2013. Repeated evolution of salt-tolerance in

grasses. Biol. Lett. 9:20130029.

Bromham L. 2016. Testing hypotheses in macroevolution. Stud. Hist. Philos. Sci. A 55:47–

59.

Bromham L., Hua X., Cardillo M. 2016. Detecting macroevolutionary self-destruction from

phylogenies. Syst. Biol. 65:109–127.

Bromham L., Duchêne S., Hua X., Ritchie A., Duchêne D., Ho S. 2018. Bayesian molecular

dating: opening up the black box. Biol. Rev. 93:1165–1191.

Bromham L., Hua X., Cardillo M. 2020. Macroevolutionary and macroecological approaches

to understanding the evolution of stress tolerance in plants. Plant Cell Environ. 43:1–

15.

Byrne M., Yeates D.K., Joseph L., Kearney M., Bowler J., Williams M.A., Cooper S.,

Donnellan S.C., Keogh J.S., Leys R., Melville J., Murphy D.J., Porch N., Wyrwoll

K.H. 2008. Birth of a biome: insights into the assembly and maintenance of the

Australian arid zone biota. Mol. Ecol. 17:4398–4417.

Byrne M., Murphy D.J. 2020. The origins and evolutionary history of xerophytic vegetation

in Australia. Aust. J. Bot. 68:195–207.

Bui E.N., González-Orozco C.E., Miller J.T. 2014a. *Acacia*, climate, and geochemistry in

Australia. Plant and Soil 281:161–175.

Bui E.N., Thornhill A., Miller J.T. 2014b. Salt- and alkaline-tolerance are linked in *Acacia*.

Biol. Lett. 10:20140278.

Bui E.N., Wilford J., de Caritat P. 2017. Predictive spatial modelling for mapping soil salinity

at continental scale, EGU General Assembly, Vienna, Austria. doi:

10.13140/RG.2.2.25552.20483.

Carpenter R.J., McLoughlin S., Hill R.S., McNamara K.J., Jordan G.J. 2014. Early evidence

of xeromorphy in angiosperms: stomatal encryption in a new eocene species of

*Banksia* (Proteaceae) from Western Australia. Am. J. Bot. 101:1486–1497.

Crayn D.M., Rossetto M., Maynard D.J. 2006. Molecular phylogeny and dating reveals an

Oligo-Miocene radiation of dry-adapted shrubs (former Tremandraceae) from

rainforest tree progenitors (Elaeocarpaceae) in Australia. Am. J. Bot. 93:1328–1342.

Crisp M., Cook L., Steane D.A. 2004. Radiation of the Australian flora: what can

comparisons of molecular phylogenies across multiple taxa tell us about the evolution

of diversity in present day communities? Philos. Trans. R. Soc. B 359:1551–1571.

Crisp M.D., Burrows G.E., Cook L.G., Thornhill A.H., Bowman D.M.J.S. 2011. Flammable

biomes dominated by eucalypts originated at the Cretaceous–Palaeogene boundary.

Nat. Commun. 2:193.

Delzon S. 2015. New insight into leaf drought tolerance. Funct. Ecol. 29:1247–1249.

Duchêne S., Lanfear R., Ho S.Y.W. 2014. The impact of calibration and clock-model choice

on molecular estimates of divergence times. Mol. Phylogenetics Evol. 78:277–289.

Duchêne D.A., Hua X., Bromham L. 2017. Phylogenetic estimates of diversification rate are

affected by molecular rate variation. Journal of Evolutionary Biology 30:1884–1897.

Evans M.E.K., Smith S.A., Flynn R.S., Donoghue M.J. 2009. Climate, niche evolution, and

diversification of the "bird-cage" evening primroses (Oenothera, sections Anogra and

Kleinia). Am. Nat. 173:225–240.

Fithian W., Elith J., Hastie T., Keith D.A. 2015. Bias correction in species distribution

models: pooling survey and collection data for multiple species. Methods Ecol. Evol.

6:424–438.

Gilkes B., Lee S., Singh B. 2003. The imprinting of aridity upon a lateritic landscape: an

illustration from southwestern Australia. C. R. Geosci. 335:1207–1218.

Gong Y., Lv G., Guo Z., Chen Y., Cao J. 2017. Influence of aridity and salinity on plant

nutrients scales up from species to community level in a desert ecosystem. Sci. Rep.

7:6811.

Guindon S. 2020. Rates and Rocks: Strengths and Weaknesses of Molecular Dating Methods.

Front. Genet. 11:526–526.

Haque M.M., Nipperess D.A., Gallagher R.V., Beaumont L.J. 2017. How well documented is

Australia's flora? Understanding spatial bias in vouchered plant specimens. Austral

Ecol. 42:690–699.

Hua X., Cardillo M., Bromham L. 2020. Tracking niche change through time: simultaneous

inference of ecological niche evolution and estimation of contemporary niches.

bioRxiv XXXXXXXXXX

Jordan G. J., Weston P.H., Carpenter R.J., Dillon R.A., Brodribb T.J. 2008. The evolutionary

relations of sunken, covered, and encrypted stomata to dry habitats in Proteaceae. Am.

J. Bot. 95:521–530.

Kozak K., Wiens J. 2010. Accelerated rates of climatic-niche evolution underlie rapid species

diversification. Ecol. Lett. 13:1378–1389.

Maslin B.R., Miller J.T., Seigler D.S. 2003. Overview of the generic status of *Acacia*

(Leguminosae: Mimosoideae). Aust. Syst. Bot. 16:1–18.

Miller J.T., Murphy, D.J., Brown G.K., Richardson D.M., González-Orozco C.E. 2011. The

evolution and phylogenetic placement of invasive Australian *Acacia* species.

Diversity Distrib. 17:848–860.

Miller J.T., Murphy D.J., Ho S.Y.W., Cantrill D.J., Seigler D. 2013. Comparative dating of

*Acacia*: combining fossils and multiple phylogenies to infer ages of clades with poor

fossil records. Aust. J. Bot. 61:436−445.

Mishler B.D., Knerr N., González-Orozco C.E., Thornhill A.H., Laffan S.W., Miller J.T.

2014. Phylogenetic measures of biodiversity and neo- and paleo-endemism in

Australian *Acacia*. Nat. Comm. 5:4473.

Moore G.M. 2013. Adaptations of Australian tree species relevant to water scarcity in the

urban forest. Arboric. Urban. For. 39:109−115.

Moray C., Hua X., Bromham L. 2015. Salt tolerance is evolutionarily labile in a diverse set

of angiosperm families. BMC Evol. Biol. 15:90.

Münkemüller, T., Boucher, F.C., Thuiller, W., & Lavergne, S. (2015). Phylogenetic niche

conservatism – common pitfalls and ways forward. Funct. Ecol. 29:627−639.

Murphy D. J., Brown G. K., Miller J. T., Ladiges P. Y. 2010. Molecular phylogeny of *Acacia*

Mill. (Mimosoideae: Leguminosae): evidence for major clades and informal

classification. Taxon 59:7–19.

Nie Y., Foster C.S.P., Zhu T., Yao R., Duchêne D.A., Ho S.Y.W., Zhong B. 2019.

Accounting for uncertainty in the evolutionary timescale of green plants through

clock-partitioning and fossil calibration strategies. Syst. Biol. 69:1−16.

Onstein R.E., Jordan G.J., Sauquet H., Weston P.H., Bouchenak-Khelladi Y., Carpenter R.J.,

Linder H.P. 2016. Evolutionary radiations of Proteaceae are triggered by the

interaction between traits and climates in open habitats. Glob. Ecol. Biogeogr.

25:1239–1251.

Owen C.L., Marshall D.C., Hill K.B.R., Simon C. 2017. How the aridification of Australia

structured the biogeography and influenced the diversification of a large lineage of

Australian cicadas. Syst. Biol. 66:569−589.

Paradis E., Schliep K. 2018. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics, bty633.

Renner I.W., Elith J., Baddeley A., Fithian W., Hastie T., Phillips S.J., Popovic G., Warton

D.I. 2015. Point process models for presence-only analysis. Methods Ecol. Evol.

6:366–379.

Renner M.A.M., Foster C.S.P., Miller J.T., Murphy D.J. 2020. Increased diversification rates

are coupled with higher rates of climate space exploration in Australian *Acacia*

(Mimosoideae). New Phytol. 226:609–622.

Skeels A., Cardillo M. 2017. Environmental niche conservatism explains the accumulation of

species richness in Mediterranean-hotspot plant genera. Evolution 71: 582–594.

Sturges H.A. 1926. The choice of a class interval. J. Am. Stat. Assoc. 21:65–66.

Weston P.H., Jordan G.J. 2017. Evolutionary biogeography of the Australian flora in the

Cenozoic era. In 'Australian vegetation,' 3rd edn. (Ed. DA Keith) pp. 40–62.

(Cambridge University Press: Cambridge, UK)

Williams K.J., Belbin L., Austin M.P., Stein J.L., Ferrier S. 2012. Which environmental

variables should I use in my biodiversity model? Int. J. Geogr. Inf. Sci. 26:2009–

2047.

Figure 1. Phylogenetic distribution of the inferred amount of niche evolution along axes of

aridity (left) and salinity (right). The blue to red color scale gives the sum of the amount of

change in the mean of realized niche values before and after each niche evolution event

inferred along each branch, averaged over posterior samples. Shifts towards less extreme

conditions are in blue and shifts towards more extreme conditions are in red. The ecoregion

map shows the current distribution of eight ecoregions in Australia. Each tip species has a

stacked bar showing the relative proportion of its current presence locations in each

ecoregion, in the same color scheme as the ecoregion map. We also include two additional

maps. Left one shows the spatial variation in the aridity index in Australia, with lower values

in more arid areas. Right one shows the variation in root zone soil electrical conductivity,

with higher values in areas with higher salinity. The mean and the 95% range of estimated

rate of niche evolution over all posterior samples are marked for each major clade in the

phylogeny.

Figure 1 cont.

Figure 2. Boxplots of the net amount of niche evolution along aridity axis A) and salinity axis

B). "All species" plots the sum of the amount of niche evolution along the branches that

connect the root to each tip of the phylogeny. "Northern clade", "Southern clade", "Eastern

clade" plot the sum of the amount of niche evolution along the branches that connect the

MCRA of the three clades of recent radiation to each tip in these clades in Figure 1. Tips are

grouped by the ecoregion that the majority of current presence location of the tip species fall

in, using the same color scheme as the ecoregion map in Figure 1. T-test is used to first

compare within vegetation types: desert, Mediterranean, grasslands, forests. If regions of the

same vegetation type do not differ, then these regions are grouped and compared to other

vegetation types. The statistical significance of a t-test between two bioregions linked by

lines is shown above the corresponding horizontal line.

A) The amount of niche evolution along aridity axis

B) The amount of niche evolution along salinity axis

NS: *p*>0.05    *: 0.01<*p*<0.05    **: 0.001<*p*<0.01    ***: *p*<0.001

Deserts & Xeric Shrublands

Montane Grasslands & Shrublands
Temperate Grasslands, Savannas & Shrublands
Tropical & Subtropical Grasslands, Savannas & Shrublands

Mediterranean Forests, Woodlands & Scrub_SE
Mediterranean Forests, Woodlands & Scrub_SW

Temperate Broadleaf & Mixed Forests
Tropical & Subtropical Moist Broadleaf Forests

Figure 3. Histogram of the inferred dates of adaptation events in *Acacia* drought tolerance

(solid) and salt tolerance (dashed).

Figure 4. Correlation in the amount of niche evolution per branch along aridity and salinity.

Each datapoint is a branch, with internal branches in black. The size of each datapoint is

proportional to the tenth root of the branch length. Terminal branches are colored by the

ecoregion that the majority of current presence location of the corresponding tip species fall

in, using the same color scheme as the ecoregion map in Figure 1.

582