Strapline: Research article

Title: Adapting to extremes: reconstructing evolution in response to changing climate

over time and space in the diverse Australian plant genus *Acacia*.

Running title: Evolution of *Acacia* in a changing climate.

Authors: Xia Hua[1,2], Marcel Cardillo[1], Lindell Bromham[1*]

Affiliations:

[1]Macroevolution and Macroecology, Research School of Biology, Australian National

University, Canberra ACT 0200 Australia

[2]Mathematical Sciences Institute, Australian National University, Canberra ACT 0200

Australia

[*]Corresponding author:

Lindell Bromham

Research School of Biology

Australian National University

Canberra ACT 0200

Australia

Tel: +61 2 6125 5945

E-mail: lindell.bromham@anu.edu.au

Acknowledgements

We thank Joe Miller for providing *Acacia* distribution data, and we are grateful to Joe Miller,

Elisabeth Bui, John La Salle, Rebecca Pirzl, and Lee Belbin for sharing their insight and

expertise with us.

Statement of significance

Acacia species are found throughout Australia, from rainforests to deserts, and are striking in

their environmental adaptability, so they are a perfect case study for understanding evolution

of tolerance to environmental extremes in a changing climate. We use the largest database of

spatial distribution records yet assembled, using both surveys and atlas data, and a new

analytical method that combines the strengths of environmental niche modelling with

phylogenetic comparative methods, to demonstrate rapid evolution in aridity and salinity

tolerance in response to aridification of the Australian continent during the Neogene and

Quaternary.

Abstract

Aim: Macroevolutionary analysis is increasingly being used to study biodiversity responses

to climate change, for example by using phylogenetic node ages to infer periods of

diversification, or phylogenetic reconstruction of traits to infer adaptation to particular

stresses. Here we apply a recently developed macroevolutionary method to investigate the

responses of a diverse plant genus, *Acacia*, to increasing aridity and salinity in Australia from

the Miocene to the present. We ask whether increase in tolerance of aridity and salinity

coincided with periods of aridification, and if it allowed the radiation of *Acacia* into a wide

range of niches.

Taxon: *Acacia*

Location: Australia

Methods: We applied the Niche Evolution Model (NEMo), which combines Environmental

(or Ecological) Niche Modelling (ENM) with phylogenetic comparative methods (PCM) in a

single statistical framework, to a large database of *Acacia* presence-only records and

presence-absence survey sites in order to infer current environmental tolerances of Australia

Acacia species and reconstruct the evolution of environmental tolerance to increasing aridity

and salinity.

Results: We find that patterns in evolution of *Acacia,* over time and across different habitat

types, are consistent with the aridification history of Australia and suggests substantial ability

to adapt to high aridity and salinity.

Main conclusions: Our results suggest that many Acacia lineages have been able to exploit

new environments created during the aridification of Australia through evolution of

environmental tolerance, resulting in their current dominance of many habitats across the

continent. This study demonstrates that phylogenetic studies of the evolution of responses to

changing environment can move beyond application of simple trait-based models, allowing

the underlying processes of speciation, adaptation and dispersal to be explicitly modelled in a

macroecological and macroevolutionary context.

Keywords: aridification, Environmental Niche Modelling (ENM), macroecology,

macroevolution, Phylogenetic Comparative Methods (PCM), stress tolerance

Introduction

The climatic and environmental history of Australia during the Neogene and Quaternary

Periods has been a story of increasing aridification. Dry climates may have existed from the

Eocene (Carpenter et al 2014), but evidence from charcoal and pollen records reveals an

increase in fire frequency and prevalence of fire-adapted, scleromorphic plant taxa, pointing

to an expansion of the arid zone from the early Miocene onwards (Byrne et al 2008). As the

arid zone expanded, previously widespread mesic forest habitats contracted to the coastal

margins. Plant groups responded in different ways (Byrne & Murphy 2020, Crisp et al 2004,

Weston & Jordan 2017). Some (such as *Nothofagus*) remained confined to humid forest

conditions and contracted with these habitats to small mesic refuges. Others responded to

new ecological opportunities provided by the expanding arid zone, and diversified into

numerous xeromorphic lineages. Some plant families, especially Proteaceae, Myrtaceae and

Fabaceae, contain both mesic and xeromorphic lineages. Even some large genera, such as

Acacia, Grevillea and *Hakea,* have both mesic and xeromorphic representatives. Plant groups

such as these, in which close relatives occupy a range of ecoclimatic conditions, provide

good case studies for studying the evolution of tolerance to extreme environmental

conditions.

The expansion of the arid zone in Australia led to increasingly challenging

environments characterized by stressful conditions for plant growth, including drought, low

soil fertility, frequent fires and high soil salinity. At the same time, areas that are extreme for

most plants represent an opportunity for lineages to specialize, escape competition, and

diversify (e.g. Thornhill et al 2016). Arid-adapted lineages have evolved a range of strategies

to cope with water stress, broadly classified into escape, avoidance, and tolerance (Delzon

2015). All three strategies occur in *Acacia* (Moore 2013), including phyllode shedding

(escape), scleromorphic structures like sunken stomata or needle-shaped phyllodes

(avoidance), and physiological adaptations such as low leaf water potential (tolerance). The

wide range of mechanisms adopted by *Acacia* species to cope with water stress are reflected

in their distribution over extremely varied climatic conditions.

To understand how opportunities provided by drying environments shaped present-

day distribution and diversity, we need to examine the dynamics of lineage adaptation in

response to extreme conditions. Macroevolutionary and macroecological analyses based on

phylogenies have provided a variety of ways of investigating adaptation to harsh

environments (Bromham et al. 2020). For example, the ages of nodes in molecular

phylogenies have been used to infer bursts of diversification associated with periods of

drying in *Acacia* (Renner et al. 2020), and the movement of rainforest-adapted lineages into

dry environments (Crayn et al. 2006). In addition to relative timing of diversification,

phylogenies can provide information on evolution of specific adaptations to extreme

conditions. For example, Crisp et al. (2011) inferred the phylogenetic history of traits such as

post-fire epicormic resprouting to investigate the timing of adaptation of the eucalypts to

increasing fire frequency. Jordan et al. (2008) reconstructed the evolution of stomatal

protection structures (which are adaptations to dry climates) in Proteaceae to show that such

structures evolved relatively few times in the ancestors of dry-climate clades. As well as

physiological and morphological traits, phylogenetic approaches have been applied to

reconstructing the evolution of adaptation to environmental conditions. For example, by

comparing alternative evolutionary models of evolution for clades from different regions,

Skeels and Cardillo (2017) showed that within four large genera (*Protea, Moraea, Banksia*,

and *Hakea*), different lineages are evolving towards different climatic optima in different

bioclimatic regions. Onstein et al. (2016) used the same approach to show that Proteaceae

lineages in open vegetation and closed forest also have different inferred climatic optima.

Phylogenetic analyses such as these are useful, but have a number of limitations. To

permit the application of standard stochastic models of trait evolution (e.g Brownian Motion

(BM) or Ornstein-Uhlenbeck (OU)) in most macroevolutionary studies, the "environmental

niche" is simplified to a point-estimate for a species, which is then treated as a species trait in

the analysis. Usually, the trait is the mean value of an environmental variable across species'

presence locations (Evans et al 2009; Kozak & Wiens 2010; Münkemüller et al 2015; Renner

et al 2020). But this approach is problematic for three main reasons.

First, the distribution of the value of environmental variables across a species'

presence locations reflects the species' realised environmental niche (that is, the conditions

under which it is found rather than the fundamental niche which describes the conditions it is

able to thrive in). While the mean of the distribution may be a fair representation for narrow-

range endemics found under uniform conditions, for most species an average value is not an

ideal representation of the range of conditions under which the species can persist. Often the

questions we want to ask are not just about the inferred midpoint of the range, but also

concern the expansion into more extreme values. For example, a species with sparse presence

under extreme conditions (reflected in a "long tail" of its realised niche distribution) is likely

to have lower average fitness under extreme conditions than a species with frequent presence

under extreme conditions (reflected in a "fat tail" of its realised niche distribution). This

information can only be captured by considering the whole niche distribution, and is lost

when environmental variables are summarised as average values. So we need an evolutionary

model to allow changes in the whole niche distribution rather than just changes in a point-

estimate of the niche distribution.

Second, even if we have an evolutionary model to allow changes in the realised

environmental niche of a species over time, changes in the realised niche do not necessarily

indicate evolution in the species' fundamental niche. For a species with wide tolerance range,

changes in its realised niche may be due to changes to its distribution, for example through

imposition of physical barriers. We need an evolutionary model to use contemporary species

distribution data to infer changes in the fundamental niche over time.

Third, standard phylogenetic methods rely on relatively simple models of trait

evolution such as BM, which may not be appropriate for modelling changes in either realised

or fundamental environmental niche. In particular, applying trait-evolution models to current

environmental data to infer "ancestral niches" (e.g., Renner et al 2020, Skeels & Cardillo

2017) assumes that environmental traits such as aridity or temperature evolve randomly and

independently along branches for each lineage. This assumption will be violated by change in

response to climatic changes which apply simultaneously and directionally across all lineages

within a given region. For example, Renner et al. (2020) reconstruct ancestral niches for

Acacia by using a BM model to infer ancestral values of climatic variables such as

precipitation, temperature and soil moisture. Under BM, the ancestor of a group of species

found in both warm and cool environments is considered most likely to have occupied an

area with a climate intermediate between the descendants, which is unlikely to be a fair

model for the evolution of the Australian flora as it responds to strong directional climate

change.

More fundamentally, phylogenetic analyses of evolution with changing environments

compartmentalise the inference of contemporary realised niche (inferred from species

distribution data) and the evolution in the fundamental niche (inferred from phylogenetic

models). These are treated as two separate problems, and subject to independent analyses

with separate statistical frameworks. But contemporary realised niches are the product of

macroevolutionary processes: inferring these processes requires inference of the fundamental

niche from contemporary distributions, so the two are fundamentally linked. The logical step,

then, is to link their analyses too.

Here, we infer the history of adaptation to extreme environments by applying a

recently developed method, the Niche Evolution Model (NEMo), that combines

Environmental Niche Modelling (ENM) and reconstruction of the evolutionary history of

fundamental niche in a single statistical framework (Hua et al. 2021). This method uses

species distribution data to characterise the realised niches of each contemporary species at

the tips of a phylogeny, and reconstructs evolution of the fundamental niche and the

associated changes in the realised niche by modelling key driving processes: (1) shift in areas

of tolerable conditions and adaptation of lineages to changing conditions (hereafter referred

to as an "adaptation event"); (2) dispersal away from areas that are no longer suitable and

into accessible areas of suitable habitat; (3) speciation and divergence along lineages'

environmental tolerances (hereafter "speciation event"). NEMo solves the three limitations of

our current phylogenetic analyses of niche evolution listed above by modelling changes in the

whole niche distribution along a phylogeny, accounting for fundamental niche and realised

niche separately, and simultaneously estimating contemporary fundamental and realised

niches and the evolution of fundamental niches along phylogeny.

Another strength of this method is that it allows us to infer the placement of

adaptation and speciation events on the phylogeny and to reconstruct ancestral fundamental

and realised niche at any time point, without making unrealistic assumptions that niches

evolve under a simple stochastic model. Unlike previous approaches, NEMo can infer the

occurrence density of evolutionary events on the phylogeny, allowing us to compare the

timing of shifts in environmental tolerance to periods of climatic change, or compare the

relative timing of evolution of tolerance along different environmental axes, using standard

statistical tests, without pre-defining the covariance structure to account for phylogenetic

non-independence. Information on paleoclimate can either be incorporated into the model as

prior information on the occurrence rate of these events, or can be held aside as independent

information against which to test the robustness of the inference (Hua et al 2021). By

explicitly accounting for the effect of niche evolution history on species current niche, NEMo

also improves the performance of ENMs at predicting contemporary realised niche and it can

predict contemporary fundamental niche (Hua et al. 2021).

We use NEMo to investigate macroevolutionary responses of the Australian plant

genus *Acacia* to aridification. *Acacia* is one of the largest and most widely-distributed plant

genera in Australia, with over 900 described species occupying nearly all major habitat types,

often as the dominant taxa. Many *Acacia* species are known to be tolerant of a range of

extreme environmental conditions, and several previous studies have used *Acacia* species as

case studies for understanding tolerance to extremes, including salinity and other aspects of

soil geochemistry (Bui et al. 2014a, 2014b), and water stress (Atkin et al. 2002). There has

been rapid progress in understanding the evolutionary history of *Acacia* in recent years

(Maslin et al. 2003; Murphy et al. 2010; Mishler et al. 2014; Renner et al. 2020; Dale et al.

2020). Nearly all *Acacia* species are conspicuous, woody plants, so spatial presence records

for *Acacia* are abundant and widespread. We combine a curated database of *Acacia* presence

records with extensive presence-absence survey data, along with predictors of sampling bias,

to characterise the current distribution of species in the phylogeny. We use NEMo to

reconstruct the evolutionary response to increasing aridification in Australian *Acacia,*

focusing on tolerance of aridity and soil salinity (Gilkes et al. 2003). Specifically, we ask two

questions. First, did evolution of tolerance to increased aridity and salinity allow present-day

Acacia species to distribute across many different habitats? Second, did increased tolerance

to aridity and salinity coincide with major periods of aridification in Australia? Answering

these two questions may help us understand whether *Acacia* responded to continental

aridification by a burst of adaptation, allowing the genus to radiate through the arid zone.

Methods

1. Phylogenetic and Spatial Data

We based our analysis on the maximum clade credibility tree of published dated

phylogenies of 505 *Acacia* species, representing half of species currently recognized (Renner

et al. 2020, Figure S1). To describe the current environmental conditions under which each

species is found, we used a published dataset by Misher et al. (2014), which includes 132,295

carefully checked presence locations of these 505 *Acacia* species, filtered from a total of

750,000 *Acacia* presence-only records from the Australian Virtual Herbarium. To represent

degree of aridity and salinity at each presence location for each species, we extracted the

value of two environmental factors at each location, one from the geographic layer of

maximum monthly aridity index (Williams et al. 2012) that takes into account both

precipitation and evaporation and the other from the geographic layer of root zone soil

electrical conductivity (Bui et al. 2017) that measures soil salinity at root zone.

Presence-only records are typically collected opportunistically, so their spatial

distributions tend to be biased towards regions that are more easily accessible to researchers,

especially along road networks, and areas of particular interest, such as protected areas or

hotspots of diversity (Fithian et al. 2015; Haque et al. 2017), and this pattern is clearly

illustrated in *Acacia* occurrence records (Figure 1). To correct for this sampling bias, we

included bias factors: distance to the nearest road (using data from Coarse Cultural

Topographic Datasets available at https://dev.ecat.ga.gov.au/geonetwork) and whether it is in

a protected area, such as a conservation reserve or national park (from

www.environment.gov.au/land/nrs/science/capad). If the parts of the species range that are

accessible from roads or included in protected areas are more likely to be sampled, then we

need to account for sampling bias in our model, because otherwise our sample locations

represent a biased characterization of the possible conditions under which the species is

found. If distance from roads or protected area are uninformative for sample location – that

is, if a species range is evenly sampled and does not show more records near a road – then

this parameter will have negligible influence on our results.

To improve model identifiability between the effect of environmental factors and the

effect of bias factors on the presence probability of a species at a location, we also assembled

species presence-absence data from government-managed floristic survey databases. Survey

data is a valuable addition to herbarium records because, unlike presence-only data, it

provides information on both presence and absence (i.e. places where the flora was surveyed

and the species was not found). Although the location of survey sites are biased, just like the

presence-only locations, the same bias applies to both the number of presences and absences

at each survey site, so the number of presences is unbiased compared to the number of

absences (Fithian et al. 2015). Therefore, survey data is very useful for characterising the

effect of sampling bias on estimating species environmental niche (Fithian et al. 2015).

There are three government-managed online databases that allow easy extraction of

the survey data and include information of each survey, including the survey sites, survey

time, survey type, plot size, GPS accuracy, and custodianship. These databases are from New

South Wales (BioNet Systematic Flora Survey;

environment.nsw.gov.au/research/VISplot.htm), Victoria (Victorian Biodiversity Atlas flora

records; environment.vic.gov.au/biodiversity/victorian-biodiversity-atlas), and Western

Australia (Flora Surveys of the Yilgarn; naturemap.dbca.wa.gov.au). While these surveys

include only three states (which cover nearly half of the land area of Australia), here they are

only used to increase the accuracy in estimating the coefficients of the two sampling bias

factors. With the information of each survey, we filtered survey data that are from complete

floristic surveys conducted in the previous 15 years under government custodianship to

ensure consistent standards. As a result, our presence-absence survey data included 27,850

survey sites across three states of Australia. These presence-absence data were analysed

together with the presence-only locations.

2. Niche Evolution Model (NEMo)

NEMo uses a Bayesian framework that includes three major components: model of niche

evolution, phylogenetic comparative method (PCM), and environmental niche model (ENM).

NEMo models species distributions in "niche space", not geographic space, by making

assumptions about the dispersal process in the model of niche evolution (see below). We

model niche space in this way because we rarely have reconstructed paleoclimate with high

resolution over a large region, without which it is impossible to model species distribution

over geographic space on geological time scale. So NEMo provides a widely applicable

approach to allow us model the evolution of the whole niche distribution under more

biologically informed processes, at the cost of making simplified assumptions on the

dispersal process. In comparison, traditional PCMs make simplified assumptions on the

process of niche evolution and do not model dispersal process.

The niche evolution model calculates how three classically recognized aspects of

niche (fundamental, available, realized) change over time under three basic processes

(adaptation, dispersal, speciation). "Fundamental niche" represents the range of conditions

that each species can tolerate at a given point in time, expressed as the proportion of

individuals of a species that can tolerate a given condition, or a value along a relevant

environmental axis. We expect species to evolve along this environmental axis over

evolutionary time, through the processes of adaptation to new conditions (by selecting for

individuals at the extreme values of the distribution) and through speciation along

environmental gradients (so that a species splits along its tolerance range leaving one

daughter species that is more skewed toward the extreme conditions). "Available niche"

describes the range of conditions in areas that are accessible to the species, regardless of

whether the species can persist in those conditions. The potential of the species to occupy the

available niche is governed by its fundamental niche and dispersal ability. We also include

barriers to dispersal as nuisance parameters, which could represent physical or biotic barriers.

"Realized niche" describes the locations where a species is actually found, which is shaped

by the suitability of conditions and the opportunity to disperse to and occupy those areas.

The PCM part of the model infers the locations on the phylogeny of adaptation and

speciation events, assuming these events are distributed along the branches of the phylogeny

according to a Poisson process with constant occurrence rate over the phylogeny. The

distribution of these events on the phylogeny gives a possible history of evolution in the

fundamental environmental niche. Dispersal in niche space is modelled as a continuous

process between any two events (adaptation or speciation), or nodes of the phylogeny. It is

assumed to be a diffusion process to reach the realized niche at equilibrium given the

potential of the species and its available niche that is truncated at dispersal barriers. So

dispersal updates the realised niche as a response to the evolution in the fundamental niche

and changes in the environmental conditions in the available niche. Given this history, we use

the niche evolution model to calculate the three classic aspects of the niche of each

contemporary (tip) species. The ENM component of the method then assess the fit of the

inferred niches to the presence and/or absence locations of the tip species. The fit of the

inferred niches to the data then feeds back to our inferences of the history of the evolutionary

events underlying shifting environmental tolerances along the phylogeny, and the niches of

tip species. Details of the method, and description of tests of its performance and reliability,

are given in Hua et al. (2021). Code with step-by-step instructions for implementing the

model can be found at

https://github.com/huaxia1985/NEMo. Below we briefly describe the parameters of the

model applied to this case study.

3. Accounting for sampling bias

We used an inhomogeneous Poisson point process (Renner et al. 2015) as the ENM

component of NEMo. Because opportunistic sampling bias can influence the niche estimation

(Fithian et al. 2015), we modified the likelihood function to account for sampling bias, using

the method of Fithian et al. (2015): details are given in the Supplementary Information.

4. Model Parameters

NEMo has four universal parameters which are assumed to be constant across the phylogeny:

heritability, dispersal rate, the occurrence rate of adaptation events, and the occurrence rate of

speciation events. We apply separate NEMo analyses to the evolution of environmental

tolerances in *Acacia* along aridity and salinity axes. For each of these axes, the values for the

"available niche" at the root of the phylogeny were assumed to follow a normal distribution

that is truncated at the observed maximum and minimum values of the axis (Williams et al.

2012; Bui et al. 2017); the values for the "fundamental niche" were assumed to follow

$f(y) = e^{-(y/\alpha)^{\beta}}$, where y is a value along the axis (see Hua et al. 2021 for details of the

function). The starting values for these parameters and their priors are described in the

Supplementary Information.

5. MCMC approximation

For both aridity and salinity, we ran 20 independent reversible-jump Markov Chain Monte

Carlo (rjMCMCs) for 3×10^6 generations, with a thinning interval of 1000: details provided in

the Supplementary Information (Figure S2). For both analyses, the 20 rjMCMCs agree on a

general pattern of the amount of change in the fundamental niche along *Acacia* phylogeny

(Figure S3 and S4), so we use the niche evolution history with the highest posterior

probability to illustrate this general pattern for both analyses (Figure S1).

Results

The *Acacia* phylogeny is characterized by a mixture of large, young clades (recent, rapid

divergences), and smaller, older clades (longer-persisting lineages: Figure S1). The young

clades, each around 10 Ma old, include most present-day species in tropical forests and

grasslands and the surrounding desert areas (in clades 10 and 11; Figure S1), as well as most

present-day species in subtropical forests and grasslands (in clades 7 and 8; Figure S1). The

smaller, older clades, each around 20 Ma old, mainly include present-day species in habitats

classified as "Mediterranean forests, woodlands & scrub" in World Wildlife Fund's

categories of terrestrial ecoregions (https://www.worldwildlife.org/biome-

categories/terrestrial-ecoregions; Figure S1) (i.e., ecosystems occurring in Mediterranean

type climates) and the surrounding desert areas (in clades 1, 3-6, 8; Figure S1). Present-day

species in Mediterranean-type ecosystems also form smaller clades that are sisters to the

recent radiations in tropical and subtropical forests and grasslands (in clades 10-11 and 7-8;

Figure S1). In short, present-day species that occupy regions that are arid and prone to high

salinity in today's Australia (Mediterranean ecosystems and surrounding deserts) are

scattered throughout the phylogeny, forming generally smaller clades than present-day

species that occupy the relatively mesic zone with low salinity (tropical and subtropical

forests and grasslands), which are found mainly in large and young clades.

1. The amount of niche evolution along **Acacia** *lineages*

The amount of evolution in the fundamental niche along the environmental tolerance axes

inferred by NEMo along the phylogeny is shown as branch color in Figure S1, with red

indicating increased tolerance of the extreme conditions over the branch and blue indicating

decreased tolerance. Both aridity and salinity analyses infer more red than blue branches, but

the result is much stronger in the salinity analysis (Figure S3 and S4), which suggests that

evolutionary change, particularly on salinity axis, has been pushing species towards tolerance

of more extreme conditions. We can also see this pattern in the inferred realised niche at the

root of the phylogeny. The root had realised niche on aridity axis under conditions equivalent

to the medium aridity level of today's Australia, but its realised niche on salinity axis is under

conditions equivalent to the lowest salinity level of today's Australia (Figure S1). So our

analyses suggest that the common ancestor of *Acacia* was likely to have been able to tolerate

the aridity level of most non-desert areas in today's Australia, but was possibly sensitive to

soil salinity.

We can also compare the inferred environmental niches of present-day species in

different habitats (using World Wildlife Fund's categories of terrestrial ecoregions), as well

as comparing them to the common ancestor (Figure 2). On the aridity axis, present-day

species mainly found in tropical forests have the most similar environmental niches (in all

three aspects: fundamental, available, realised) to the common ancestor (Figure 2A-2C), a

pattern which is also reflected in the absence of niche evolution along the branches

connecting the root and the most recent common ancestor (MRCA) of the radiation in

tropical forests and grasslands (Figure S1). The largest changes in environmental niches (in

all three aspects) are in lineages currently found in deserts, the most arid environments

(Figure 2A-2C). Along the branches leading to these desert species, there are often multiple

rounds of evolution to tolerate higher aridity on the phylogeny, both in the old clades and the

recent radiations (Figure S1). Lineages in all the other habitats have similar inferred

fundamental niche (Figure 2A), although their inferred realised niches tend to differ (Figure

2C).

In contrast, on the salinity axis, our analysis inferred large shifts in both fundamental

niche and realised niche towards higher salinity since the common ancestor of *Acacia* in all

president-day lineages (Figure 2D-2F). The amount of shift is largest in species currently

distributed in the Mediterranean ecosystems, and is smallest in extant tropical and montane

species (Figure 2D-2F). Lineages in all the other habitats have similar inferred fundamental

niche and realised niche (Figure 2D,2F). Shifts in most lineages are due to the major niche

evolution events at the very basal branches of the phylogeny. Along the branches leading to

species currently distributed in the Mediterranean ecosystems, there are multiple rounds of

evolution to tolerate higher salinity, both in the old clades and the recent radiations (Figure

S1).

2. The timing of niche evolution events

Figure S1 shows the inferred occurrences of adaptation and speciation events along the

phylogeny, with red indicating events that increase the tolerance of the extreme conditions

and blue indicating events that decrease the tolerance. Since there have been three significant

drying periods in Australia during the Cenozoic Era: around 28-23 Million years ago (Ma),

14-5 Ma, and throughout the Pleistocene since 2.6 Ma (Owen et al. 2017), we count the

number of inferred adaptation and speciation events between the time intervals of the root

age, 23 Ma, 14 Ma, 5 Ma, and 2.6 Ma. A common pattern of these niche evolution events

along aridity and salinity axes is that speciation events have similar occurrence density per

lineage over all time intervals (Figure 3A,3C), while adaptation events have increasing

occurrence density per lineage towards the present, with significantly high occurrence density

during the Pleistocene (Figure 3B,3D). This burst of adaptation events along aridity and

salinity axes in the Pleistocene mainly occurs in clades 7-11 that include most of the present-

day species in the tropical and subtropical forests and grasslands (Figure S5).

On the aridity axis, no adaptation or speciation events are inferred along the basal

branches between the root of the phylogeny and 23 Ma (Figure S1). There tends to be higher

occurrence density of adaptation events with increasing aridity than adaptation events with

decreasing aridity during the drying periods in the Pleistocene and between 14-5 Ma, and

vice versa between 5-2.6 Ma, the short period when Australia became wet and warm again

(Figure 3B). The differences between the occurrence densities of adaptation events with

increasing aridity versus decreasing aridity are statistically significant during the Pleistocene

and between 14-5 Ma. Both have 95% confidence intervals of the difference in the

occurrence densities above zero (the Pleistocene: 0.036~0.096; 14-5 Ma: 0.001~0.020). In

contrast, our analysis infers continuously increased tolerance to salinity over the *Acacia*

phylogeny from the very basal branches of the phylogeny to the present (Figure S1). The

occurrence density of adaptation events with increasing salinity is consistently higher than

the occurrence density of adaptation events with decreasing salinity over all time intervals

(Figure 3D). This result is consistent with the idea that soil salt might have been continuously

accumulated in Australia over geological time as a result of the antiquity and flatness of the

continent, as well as the complex relationship between salinisation and aridification which is

not a simple linear relationship (George et al. 2008).

3. Uncorrelated evolution on aridity and salinity axes

To test if evolution to tolerate aridity occurs in concert with evolution to tolerate salinity in

Acacia, we plot the inferred amount of evolution in the fundamental niche along each branch

of the *Acacia* phylogeny (i.e., the branch color in Figure S1) on the aridity axis versus that on

the salinity axis in Figure 4A. The Pearson's correlation test suggests no correlation in the

amount of evolution on the aridity and salinity axes (t_{1004}=0.47, p=0.64; Figure 4A). Yet,

present-day species that tolerate high level of salinity tend to tolerate high level of aridity

(t_{503}=2.83, p=0.005; Figure 4B), which is mostly driven by species in the deserts (t_{97}=3.72, p

<0.001), the tropical grasslands (t_{95}=4.23, p <0.001), and the Mediterranean forests in South

Australia (t_{31}=2.20, p=0.04; Figure S6).

Discussion

We reconstructed the evolution of the widespread plant genus *Acacia* over time and across

different habitat types and infer a preponderance of evolutionary change towards more

extreme values of aridity and salinity, inferring that the *Acacia* common ancestor could

tolerate medium aridity level and very low salinity level, compared to contemporary levels in

Australia. Our results suggest that there are multiple rounds of evolution of environmental

tolerance in the ancestors of species currently found in arid and Mediterranean regions in

both the old clades and the recent radiations. In contrast, we infer relatively little increase in

aridity in recent radiations that lead to present-day species in tropical and subtropical forests

and grasslands. Our results suggest different tempo and mode of evolution of aridity and

salinity tolerance: while adaptation events increasing aridity tolerance are more frequent

during the major periods of aridification in Australia, adaptation to higher salinity occurs

over all periods. However, both salinity and aridity tolerance are inferred to have undergone a

burst of adaptation events on both axes in the Pleistocene. These differing patterns suggest a

decoupling of evolution of aridity and salinity tolerance, even though the tolerance level of

aridity and salinity correlates in present-day species. As we explain below, these results are

consistent with the adaptation of *Acacia* lineages to drier climates, allowing them to exploit

new environmental opportunities opening up through the aridification of the continent. Our

results are consistent with evolutionary flexibility in *Acacia* that may have played a key role

in their current near-ubiquitous distribution across Australia and their dominance of many

habitats.

The preponderance of evolution towards more extreme values of both aridity and

salinity suggests that aridification has been a dominant driver of *Acacia* evolution and

diversification across Australia during the Neogene and Quaternary. The multiple events of

niche evolution to tolerate higher aridity and salinity on branches leading to species currently

found in more arid and salt-affected regions suggest strong adaptation ability of *Acacia* to

tolerate extreme environments. MRCAs of recent radiations in tropical and subtropical

forests and grasslands can tolerate the same level of aridity as the common ancestor of all

Acacia species included in the phylogeny. Since the common ancestor of *Acacia* already had

the ability to tolerate the aridity level typical of non-desert areas in today's Australia, our

results support the hypothesis proposed by Renner et al. (2020) that *Acacia* was able to

replace many other species groups in these mesic areas during the drying period of the

Pleistocene, because these areas had increased aridity during the drying period and *Acacia*

tolerated higher aridity. We also find a burst of adaptation events on both aridity and salinity

axes in the Pleistocene, particularly in these recent radiations. This enhanced rate of niche

evolution was also found by Renner et al. (2020) using traditional PCMs, which hypothesized

that Pleistocene glacial-interglacial climate cycling (< 2.5Ma) drove rapid climatic

adaptation. Yet, we find no evidence that niche evolution along aridity and salinity axes

promoted diversification and led to the radiations, as Renner et al. (2020) hypothesized,

because we find that the occurrence density of speciation events that are associated with

niche evolution is not elevated during the Pleistocene. This result does not rule out the

possibility that diversification is promoted by niche evolution along environmental axes other

than aridity or salinity, but our analyses show that a clade with both high diversification rate

and high rate of niche evolution does not necessarily suggest that niche evolution promotes

diversification.

Comparing salt tolerance to aridity tolerance in *Acacia*, we find that the amount of

evolution along aridity and salinity axes was not correlated across branches of the *Acacia*

phylogeny. However, our inferred fundamental niches of present-day species are correlated

between their tolerance abilities to aridity and salinity. This result is consistent with the

observation that saline soils are frequently associated with aridity, so species need to adapt to

both aridity and salinity to survive (Gong et al. 2017). Traditional PCMs based on the

distribution data of tip species have been applied to test correlated evolution between

different types of tolerance in *Acacia*. For example, correlated evolution was suggested

between salinity and alkalinity tolerance in *Acacia* using phylogenetically independent

contrasts (Bui et al. 2014b). Since traditional PCMs often assume Brownian motion

evolution, they often infer correlated evolution if aspects of tip realised niches are correlated.

Our analysis show that even when tip niches on different axes are correlated, we cannot

assume that niche evolution along these axes occurs in concert, simply because there could be

different histories to generate similar tip niches.

It is important to note that for the NEMo analysis, as for most macroevolutionary

analyses, we have assumed that the phylogeny has accurate topology and provides an

approximate scaling of evolutionary events in time. Clearly, phylogenetic estimates of either

topology or the timing of evolutionary events are not error-free (Bromham et al. 2018;

Guindon 2020; Nie et al. 2019). In particular, uncertainty or bias in branch length estimates

arising from imperfect characterisation of molecular rates over time may have a non-trivial

impact on macroevolutionary inference (Duchêne et al. 2017). Phylogenies such as this one

where the only calibration is a root date may be particularly vulnerable to errors in position of

node heights in some parts of the tree (Duchêne et al. 2014). One possible solution is to apply

the NEMo analysis on posterior samples of phylogenies, but perhaps a better advance will be

to embed the NEMo method into phylogenetic reconstruction, which allows us to directly

link evolution on environmental tolerance to diversification events.

Another issue with our current inference of evolution in *Acacia* is incomplete

sampling, as the *Acacia* phylogeny only includes half of the known extant species, and

sampling is not random either with respect to area or lineage (Renner et al 2020). One

obvious solution is to sample more species, but another way to increase the amount of

information in the analysis is to incorporate historical data, such as paleoclimate or fossil

distribution data, to supplement current species distribution data. For example, we could

adjust priors for the occurrence rate of adaptation events with increasing aridity to allow

higher rate under the known drying periods in Australia. Or we could include fossil locations

and reconstructed paleoclimate by using the ENM equations to calculate the probability of

sampling fossils at the locations given its inferred niches on the phylogeny and incorporate

the probability into the posterior probability of NEMo. Here we have deliberately used only

part of the available information in the inference process, so we can compare our inferred

evolutionary history to known paleoclimatic history in order to demonstrate the utility of the

method, but the method allows for all available information to contribute to stronger

inference.

Conclusion

To understand how plants in today's Australia have responded to continuing aridification

through geological time, we use a newly developed approach to reconstruct the evolution of

tolerance to increasing aridity and soil salinity in a major Australian plant genus *Acacia.* By

explicitly modelling changes in the fundamental niche, realised niche, and available niche of

Acacia along its phylogeny, we find evidence that is consistent with *Acacia* having great

adaptability to increasing aridity and salinity and that *Acacia*'s ability to tolerate aridity

allows it to dominate not only the arid areas, but also the mesic areas by surviving the drying

periods during Pleistocene in Australia. But unlike previous studies, we find no evidence that

Acacia's adaptability to increasing aridity and salinity promoted its radiation in today's mesic

areas. Neither do we find evidence for correlated evolution between tolerance to aridity and

salinity, even when present-day *Acacia* species often need to adapt to both high aridity and

high soil salinity. These results demonstrate the value of modelling biologically-informed

processes of niche evolution, because we cannot make reliable inferences on processes from

patterns without modelling the processes.

References

Atkin O., Schortemeyer M., McFarlane N., Evans J.R. 2002. Variation in the components of

relative growth rate in 10 *Acacia* species from contrasting environments. Plant Cell

Environ. 21:1007–1017.

Bennett T.H., Flowers T.J., Bromham L. 2013. Repeated evolution of salt-tolerance in

grasses. Biol. Lett. 9:20130029.

Bromham L., Hua X., Cardillo M. 2016. Detecting macroevolutionary self-destruction from

phylogenies. Syst. Biol. 65:109–127.

Bromham L., Duchêne S., Hua X., Ritchie A., Duchêne D., Ho S. 2018. Bayesian molecular

dating: opening up the black box. Biol. Rev. 93:1165–1191.

Bromham L., Hua X., Cardillo M. 2020. Macroevolutionary and macroecological approaches

to understanding the evolution of stress tolerance in plants. Plant Cell Environ. 43:1–

15.

Byrne M., Yeates D.K., Joseph L., Kearney M., Bowler J., Williams M.A., Cooper S.,

Donnellan S.C., Keogh J.S., Leys R., Melville J., Murphy D.J., Porch N., Wyrwoll

K.H. 2008. Birth of a biome: insights into the assembly and maintenance of the

Australian arid zone biota. Mol. Ecol. 17:4398–4417.

Byrne M., Murphy D.J. 2020. The origins and evolutionary history of xerophytic vegetation

in Australia. Aust. J. Bot. 68:195–207.

Bui E.N., González-Orozco C.E., Miller J.T. 2014a. *Acacia*, climate, and geochemistry in

Australia. Plant and Soil 281:161–175.

Bui E.N., Thornhill A., Miller J.T. 2014b. Salt- and alkaline-tolerance are linked in *Acacia*.

Biol. Lett. 10:20140278.

Bui E.N., Wilford J., de Caritat P. 2017. Predictive spatial modelling for mapping soil salinity

at continental scale, EGU General Assembly, Vienna, Austria. doi:

10.13140/RG.2.2.25552.20483.

Carpenter R.J., McLoughlin S., Hill R.S., McNamara K.J., Jordan G.J. 2014. Early evidence

of xeromorphy in angiosperms: stomatal encryption in a new eocene species of

Banksia (Proteaceae) from Western Australia. Am. J. Bot. 101:1486–1497.

Crayn D.M., Rossetto M., Maynard D.J. 2006. Molecular phylogeny and dating reveals an

Oligo-Miocene radiation of dry-adapted shrubs (former Tremandraceae) from

rainforest tree progenitors (Elaeocarpaceae) in Australia. Am. J. Bot. 93:1328–1342.

Crisp M., Cook L., Steane D.A. 2004. Radiation of the Australian flora: what can

comparisons of molecular phylogenies across multiple taxa tell us about the evolution

of diversity in present day communities? Philos. Trans. R. Soc. B 359:1551–1571.

Crisp M.D., Burrows G.E., Cook L.G., Thornhill A.H., Bowman D.M.J.S. 2011. Flammable

biomes dominated by eucalypts originated at the Cretaceous–Palaeogene boundary.

Nat. Commun. 2:193.

Dale, E.E., Larcombe, M.J., Lee, W.G., Higgins, S.I. 2020. Diversification is decoupled from
biome fidelity: *Acacia* – a case study. J. Biogeogr. 47:538–552.
doi:10.1111/jbi.13768.

Delzon S. 2015. New insight into leaf drought tolerance. Funct. Ecol. 29:1247–1249.

Duchêne S., Lanfear R., Ho S.Y.W. 2014. The impact of calibration and clock-model choice
on molecular estimates of divergence times. Mol. Phylogenetics Evol. 78:277–289.

Duchêne D.A., Hua X., Bromham L. 2017. Phylogenetic estimates of diversification rate are
affected by molecular rate variation. Journal of Evolutionary Biology 30:1884–1897.

Evans M.E.K., Smith S.A., Flynn R.S., Donoghue M.J. 2009. Climate, niche evolution, and
diversification of the "bird-cage" evening primroses (Oenothera, sections Anogra and
Kleinia). Am. Nat. 173:225–240.

Fithian W., Elith J., Hastie T., Keith D.A. 2015. Bias correction in species distribution
models: pooling survey and collection data for multiple species. Methods Ecol. Evol.
6:424–438.

George R., Clarke J., English P. 2008. Modern and palaeogeographic trends in the
salinisation of the Western Australian wheatbelt: a review. Aust. J. Soil Res. 46:751–
767.

Gilkes B., Lee S., Singh B. 2003. The imprinting of aridity upon a lateritic landscape: an
illustration from southwestern Australia. C. R. Geosci. 335:1207–1218.

Gong Y., Lv G., Guo Z., Chen Y., Cao J. 2017. Influence of aridity and salinity on plant
nutrients scales up from species to community level in a desert ecosystem. Sci. Rep.
7:6811.

Guindon S. 2020. Rates and Rocks: Strengths and Weaknesses of Molecular Dating Methods.
Front. Genet. 11:526–526.

Haque M.M., Nipperess D.A., Gallagher R.V., Beaumont L.J. 2017. How well documented is

Australia's flora? Understanding spatial bias in vouchered plant specimens. Austral

Ecol. 42:690–699.

Hua X., Cardillo M., Bromham L. 2021. Tracking niche change through time: simultaneous

inference of ecological niche evolution and estimation of contemporary niches.

bioRxiv. https://doi.org/10.1101/2019.12.29.890608

Jordan G. J., Weston P.H., Carpenter R.J., Dillon R.A., Brodribb T.J. 2008. The evolutionary

relations of sunken, covered, and encrypted stomata to dry habitats in Proteaceae. Am.

J. Bot. 95:521–530.

Kozak K., Wiens J. 2010. Accelerated rates of climatic-niche evolution underlie rapid species

diversification. Ecol. Lett. 13:1378–1389.

Maslin B.R., Miller J.T., Seigler D.S. 2003. Overview of the generic status of *Acacia*

(Leguminosae: Mimosoideae). Aust. Syst. Bot. 16:1–18.

Mishler B.D., Knerr N., González-Orozco C.E., Thornhill A.H., Laffan S.W., Miller J.T.

2014. Phylogenetic measures of biodiversity and neo- and paleo-endemism in

Australian *Acacia*. Nat. Comm. 5:4473.

Moore G.M. 2013. Adaptations of Australian tree species relevant to water scarcity in the

urban forest. Arboric. Urban. For. 39:109−115.

Moray C., Hua X., Bromham L. 2015. Salt tolerance is evolutionarily labile in a diverse set

of angiosperm families. BMC Evol. Biol. 15:90.

Münkemüller, T., Boucher, F.C., Thuiller, W., & Lavergne, S. (2015). Phylogenetic niche

conservatism – common pitfalls and ways forward. Funct. Ecol. 29:627−639.

Murphy D. J., Brown G. K., Miller J. T., Ladiges P. Y. 2010. Molecular phylogeny of *Acacia*

Mill. (Mimosoideae: Leguminosae): evidence for major clades and informal

classification. Taxon 59:7–19.

Nie Y., Foster C.S.P., Zhu T., Yao R., Duchêne D.A., Ho S.Y.W., Zhong B. 2019.

Accounting for uncertainty in the evolutionary timescale of green plants through

clock-partitioning and fossil calibration strategies. Syst. Biol. 69:1−16.

Onstein R.E., Jordan G.J., Sauquet H., Weston P.H., Bouchenak-Khelladi Y., Carpenter R.J.,

Linder H.P. 2016. Evolutionary radiations of Proteaceae are triggered by the

interaction between traits and climates in open habitats. Glob. Ecol. Biogeogr.

25:1239–1251.

Owen C.L., Marshall D.C., Hill K.B.R., Simon C. 2017. How the aridification of Australia

structured the biogeography and influenced the diversification of a large lineage of

Australian cicadas. Syst. Biol. 66:569−589.

Renner I.W., Elith J., Baddeley A., Fithian W., Hastie T., Phillips S.J., Popovic G., Warton

D.I. 2015. Point process models for presence-only analysis. Methods Ecol. Evol.

6:366–379.

Renner M.A.M., Foster C.S.P., Miller J.T., Murphy D.J. 2020. Increased diversification rates

are coupled with higher rates of climate space exploration in Australian *Acacia*

(Mimosoideae). New Phytol. 226:609–622.

Skeels A., Cardillo M. 2017. Environmental niche conservatism explains the accumulation of

species richness in Mediterranean-hotspot plant genera. Evolution 71: 582–594.

Sturges H.A. 1926. The choice of a class interval. J. Am. Stat. Assoc. 21:65–66.

Thornhill, A.H., Mishler, B.D., Knerr, N.J., González-Orozco, C.E., Costion, C.M., Crayn,

D.M., Laffan, S.W. and Miller, J.T., 2016. Continental-scale spatial phylogenetics of

Australian angiosperms provides insights into ecology, evolution and

conservation. *Journal of Biogeography*, *43*(11), pp.2085-2098.

Weston P.H., Jordan G.J. 2017. Evolutionary biogeography of the Australian flora in the

Cenozoic era. In 'Australian vegetation,' 3rd edn. (Ed. DA Keith) pp. 40–62.

(Cambridge University Press: Cambridge, UK)

Williams K.J., Belbin L., Austin M.P., Stein J.L., Ferrier S. 2012. Which environmental

variables should I use in my biodiversity model? Int. J. Geogr. Inf. Sci. 26:2009–

2047.

Biosketch: Xia Hua is a mathematical biologist who specialises in finding analytical

solutions to a wide range of evolutionary problems, including research in molecular

evolution, macroevolution and macroecology and language evolution. Marcel Cardillo's

research aims to understand large-scale patterns of biodiversity, with a current focus on

Australian plants. Lindell Bromham designs comparative analyses for investigating

mechanisms for the generation of diversity, from molecular evolution to macroevolution

to languages and cultures. Together, they are the three principal investigators from the

Macroevolution and Macroecology Group at the Australian National University.

Editor: Margaret Byrne

Data availability: This study uses publicly available data on *Acacia* phylogeny and

distribution, from sources cited in the text. Code with step-by-step instructions for

implementing the model can be found at https://github.com/huaxia1985/NEMo.

Figure 1. *Acacia* is a diverse genus of trees and shrubs occupying many habitats across

Australia. For example, silver wattle (a, *A. dealbata*) is found in mesic habitats in southern

Australia, dead finish (b, *A. tetragonophylla*) grows in arid and semi-arid areas, and yellow

wattle (c, *A. flavescens*) grows in humid tropical areas. While often occurring as understory

shrubs and trees, in some places *Acacia* is the dominant species, such as mulga (d, *A.*

aneura). (e) Map of occurrence records illustrating the patterns of sampling bias: each red dot

is a species presence location; black lines are roads in Australia; grey polygons are protected

areas. Presence locations are clearly distributed along roads. For areas that are distant to

roads, presence locations tend to fall in protected areas. Credit for images of *Acacia,* via

Wikimedia commons under Creative Commons licenses: (a) SABENCIA Bertu Ordiales; (b)

Melburnian; (c) and (d) Mark Marathon.

Figure 2. Boxplots of the total amount of change in the three aspects of environmental niche

(fundamental, available, realized) from root to each tip along aridity axis (A-C) and salinity

axis (D-F). Tips are grouped by the habitat that the majority of current presence location of

the tip species fall in, using the same color scheme as the habitat map in Figure S1.

Figure 3. Occurrence density of the inferred speciation and adaptation events per branch that

fall within each time interval. The mean of the occurrence density is calculated as the count

of speciation or adaptation events occurred within a time interval divided by the sum of

branch lengths within the time interval, so that the occurrence density is corrected for the

number of branches. The bars show the 95% confidence interval of the occurrence density,

assuming that occurrences follow Poisson process. A similar plot for each clade in the

phylogeny is in Figure S5.

Figure 4. Correlation in the evolution of environmental tolerance along aridity and salinity.

A) plots the amount of change in parameter value α of the fundamental niche long each

branch. So each datapoint is a branch, with internal branches in black and terminal branches

colored by the habitat that the majority of current presence location of the corresponding tip

species fall in. B) plots the total amount of change in parameter value α of the fundamental

niche from root to each tip species. So each datapoint is a tip species, colored by the habitat

that the majority of the species' presence location fall in. The plot for species in each habitat

is in Figure S6. The same color scheme is used for habitats as Figure S1.

719