SARS-CoV-2 Infection Causes Strong mTORC1 Inhibition and Massive Polysome

Collapse

Haripriya Parthasarathy*[,1], Divya Gupta*[,1], Dixit Tandel[1,2], Vishal Sah[1,2], Abdul Hamid

Siddiqui[1], Abhirami P Suresh[1], and Krishnan Harinivas Harshan[€,1,2]

*These authors contributed equally

[1]CSIR-Centre for Cellular and Molecular Biology, Hyderabad, India 500007

[2]Academy for Scientific and Innovative Research (AcSIR), Ghaziabad-201002, India

[€]Correspondence: hkrishnan@ccmb.res.in

Keywords

SARS-CoV-2, COVID-19, Delta variant, Global translation, mTORC1, Polysome profiling,

5'-TOP

ABSTRACT

Viruses employ distinct strategies to ensure efficient translation of their mRNAs over the host transcripts. SARS-CoV-2 targets host mRNAs and ribosomes to favor its own protein synthesis. However, the modulation of the key signal pathways that control the host protein translation machinery during SARS-CoV-2 infection has not been sufficiently addressed. Here, by employing an early variant and a Delta variant isolate that evolved later in the pandemic demonstrates that SARS-CoV-2 infection results in massive polysome collapse starting from 24 hpi, a hallmark of global translation inhibition. Unexpectedly, eIF2α phosphorylation, commonly targeted by viruses to induce translation arrest, was not involved in the translation arrest, suggesting that SARS-CoV-2 countermeasures by the virus to suppress ISR. eIF4E phosphorylation remained unaltered during the infection, ruling out its involvement in the preferential translation of SARS-CoV-2 transcripts. We find that SARS-CoV-2 infection consistently causes mTORC1 inhibition in a comparable manner across both variants indicating that the virus likely targets mTORC1 pathway to suppress host translation. Interestingly, mTORC1 inhibition by SARS-CoV-2 did not impact the polysomal loading of ribosomal protein transcripts rpS3 and rpL26, suggesting that 5'TOP mRNAs are spared from the translation suppression and that ribosomal protein synthesis remains active during the infection. Pharmacological activation of mTORC1 did not significantly impact viral replication, suggesting that mTORC1 inhibition might be selectively restricting the host mRNAs from accessing the translation machinery, facilitating a more robust translation of viral transcripts. This study provides new insights into the molecular interactions by which

41 SARS-CoV-2 variants, despite their different clinical outcomes, converge on a conserved

42 mechanism to manipulate host translation regulatory pathways.

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has emerged as the

deadliest of all known coronaviruses up to date, causing more than 7 million deaths

worldwide(1,2). Inside a host cell, the 5'-capped and polyadenylated viral genomic RNA

undergoes multiple rounds of translation to generate two long polypeptides that are

cleaved by proteases to generate non-structural proteins that assemble to form the

replicase complex. Translation of sub-genomic mRNAs generate structural proteins that

are temporally regulated, indicating the importance of tightly regulated translation for

successful infection by SARS-CoV-2(3) . Given that translation control is central to viral

life cycle, perturbation of host translation emerged as a common strategy observed in

SARS-CoV-2(4)

Viruses establish distinct relationships with the host translation machinery, ranging from

subverting it to coaxing, or partly manipulating it. Poliovirus completely shuts down host

translation to utilize the now available machinery for cap-independent translation of viral

transcripts (5). Several other viruses inhibit host translation to varying degrees while

allowing a set of mRNAs to translate (6,7). Yet, some other viruses such as hepatitis C

virus (HCV) do not cause an apparent suppression of host translation, but still use a cap-

independent mechanism for their translation. Coronaviruses that gained significance in

the recent past such as SARS-CoV and MERS-CoV are known to inhibit host translation

through a variety of mechanisms such as inhibition of 40S subunit and 80S formation,

mRNA degradation, and disrupting protein trafficking (7–12) Extensive studies have

established a potent involvement of several SARS-CoV-2 non-structural proteins in the

arrest of host translation. Most notably, Nsp1 interferes with host translation through its

interaction with 40S ribosomes(7,13–15). Reports also indicate SARS-CoV-2 mediated

preferential destruction of host mRNAs lead to their reduced translation events (7,16).

While a number of different mechanisms by which SARS-CoV-2 interferes with host

translation activities have been proposed, newer reports suggest significant differences

between the variants that have emerged over time (17,18)

Global translation activities in higher eukaryotes are regulated by three major pathways.

Mechanistic target of Rapamycin (mTOR) pathway is the most studied of these and is

known to regulate translation of a sub-set of mRNAs with a 5' terminal oligo pyrimidine

(5' TOP) stretch (19–21) . mTOR complex 1 (mTORC1) is active in metabolically active

cells and promotes translation by facilitating the free availability of the cap-binding

initiation factor, eIF4E (22). One of the substrates of mTORC1, eIF4E binding protein 1

(4EBP1), inhibits translation activities by sequestering eIF4E (23). mTORC1 mediated

phosphorylation of 4EBP1 lowers its affinity towards eIF4E thereby making it available for

cap-binding. mTORC1 also facilitates translation by phosphorylating the small ribosomal

protein rpS6(24), eIF4B and helicase eIF4A through p70S6K (25). Several viruses are

reported to target mTORC1 in order to suppress host translation activities(11). Inhibition

of mTORC1 is known to cause a major drop in active polysomes and translation activities

(20,21).

MAPKs p38 and ERK1/2 are known to regulate the phosphorylation of eIF4E through their substrate Mnk1/2(26,27). Even though eIF4E phosphorylation does not alter its affinity for the 5' cap of the mRNAs, phosphorylated eIF4E is commonly detected in several cancers leading several researchers to hypothesize that this phosphorylation results in preferential translation of a set of mRNAs (28,29). A third mechanism of regulation of global translation is the phosphorylation of eIF2α at S52, a key event leading to reduced recycling rate of eIF2 ternary complexes that is critical for new events of translation initiations (30). The four kinases GCN2, HRI, PKR and PERK, known as integrated stress response (ISR) kinases, coordinate this phosphorylation relaying various upstream signals (31) PKR, phosphorylates eIF2α after the detection of dsRNA replication intermediates in the cytosol during viral infections. This results in severe translational suppression in the virus infected cells as demonstrated in several cases (32,33).

While the majority of studies aimed at understanding the impact of SARS-CoV-2 infection on host translation have demonstrated a major translation inhibition by the virus, our study focused on delineating the impact of the virus on host signaling pathways that have a major impact on translation. We analysed this by employing two distinct SARS-CoV-2 variants with one of the representing an early lineage isolated during the early months of the pandemic, while the other one representing Delta variant that emerged during the later phase in the pandemic. Our findings demonstrate that both variants trigger a comparable and sustained polysome dissociation beginning at 24 hrs post infection and persisting upto 96 hrs. Further assessment of pathways regulating host translation revealed that p38 MAPK was phosphorylated by both variants throughout the course of

infection and its inhibition also resulted in lower viral replication. Interestingly, despite

activated p38 MAPK, eIF4E phosphorylation was limited. However, both variants

consistently, and strongly inhibited the mTORC1 pathway. Nevertheless, the translation

of 5' TOP transcripts continued unhindered in cells infected with both variants, indicating

escape of these mRNAs from virus-mediated mTORC1 inhibition as well as translation

arrest. Collectively, our studies demonstrate that SARS-CoV-2 infection causes severe

arrest of host translation machinery, and this is potentially supported by a strong inhibition

of mTORC1 in a selective manner to spare the key host transcripts. Importantly, these

effects were conserved between the two variants despite their distinct stage of

emergence and different clinical manifestations, indicating that SARS-CoV-2 employs a

shared molecular strategy to modulate host translation.

RESULTS

SARS-CoV-2 causes major polysome collapse

Previous reports have characterized a strong suppression of translation of host transcripts

upon SARS-CoV-2 infection through Nsp1-mediated host mRNA degradation. However,

few have explored the influence of the viral infection on host translation apparatus and

signal pathways that maintain it. We studied the effect of SARS-CoV-2 infection on

polysome formation as an indicator of translation activity. We used an earlier variant

isolate B.1.1.8 and a later variant isolate of Delta in this study to infect Caco2 cells(34).

Both variants established robust infection as evident from nucleocapsid (N) expression

(Figures 1A and D). A significant collapse of actively translating polysome pool was

observed in B.1.1.8 infection as early as 24 hours post infection (hpi), (Figure 1B and C),

indicating a rapid suppression of translation activities. Quantification of the polysome area

demonstrated a polysome depletion by about 40% in the virus infected samples by 24 hpi

against the uninfected control cells, indicating that the virus causes a marked reduction

in translation activity. Notably, this reduction in the translating pool of polysomes was

maintained throughout the infection course. A moderate increase in the 80S peak was

observed during the infection, suggesting that the fresh rounds of translation initiation

were not impacted. Delta infection also showed a comparable pattern of polysome

dissociation (Figures 1E and F). Curiously, no proportionate increase in the size of the

mRNPs fraction was noticed, suggesting that the loss in the polysomes did not result in

large-scale degradation of the mRNA associated with them (Supplementary Figures S1A

and B). Under these conditions, N protein maintained its levels, indicating that the viral

translation is unaffected by the loss in polysome fractions. The possibility that the

polysome collapse is an outcome of cell death due to viral infection was ruled out by the

cytotoxicity assay (Figures 1G for B.1.1.8 and H for Delta).

To further characterize the molecular means to this polysome loss, we studied mTOR,

MAPK, and eIF2α pathways upon infection. Strong signals of N protein and viral RNA

were detected by 12 hpi in both B.1.1.8 and Delta infections (Supplementary Figures S2

A-C; S3 A-C, respectively). Interestingly, none of the three pathways indicated any signs

of regulation suggesting that SARS-CoV-2 infection does not result in an early inhibition

of protein translation initiation (Supplementary Figures S2 D-L; Figures S3 D-L,

respectively). We subsequently extended our studies during 24-48 hpi by when the virus

had successfully established infection.

SARS-CoV-2 does not induce ISR

eIF2α phosphorylation-mediated translational arrest is often used by viruses to achieve preferential translation of viral proteins (32,33). We first asked if SARS-CoV-2 infection potentially leads to eIF2α phosphorylation resulting in polysome dissociation and translational arrest. Unexpectedly, eIF2α phosphorylation remained unchanged at 24 hpi, followed by a sharp decrease from 48 hpi onwards in B.1.1.8 infection, Although much delayed, Delta infection also caused a severe eIF2α dephosphorylation. Collectively, these findings suggest that SARS-CoV-2 mediated translational suppression does not primarily involve engagement of the eIF2α pathway (Figures 2A and C).

Inhibition of eIF2α phosphorylation by SARS-CoV-2 is an indication that the virus prefers to suppress ISR or does not want to limit the availability of ternary complexes for the interest of its own translation. Next, we asked if activating ISR would be deleterious to the virus. Caco2 cells infected with SARS-CoV-2 were treated with sodium arsenite, a general inducer of eIF2α phosphorylation and its impact on the viral titer was studied. Sodium arsenite induced strong eIF2α phosphorylation in the infected cells, but failed to reduce the levels of viral N protein (Figure 2E). However, a moderate drop in viral RNA as well as infectious viral titers were observed in the supernatant of arsenite treated cells as compared with the control (Figures 2F and G), suggesting that the viral translation is partially impacted by elevated eIF2α phosphorylation, and the virus may have evolved ways to translate under conditions of depleted canonical ternary complex.

MAPK-eIF4E axis does not mediate polysome collapse during SARS-CoV-2 infection

Since eIF2α did not mediate polysome collapse in SARS-CoV-2 infection, we asked if the two MAPKs, ERK1/2 or p38 MAPK mediate this outcome. We first investigated the

phosphorylation of ERK/2 in SARS-CoV-2 infected cells. Interestingly, as in the case of

eIF2α, ERK1/2 phosphorylation remained unchanged until 24 hpi, but started to decrease

from 48 hpi onwards (Figures 3 A-B and D-E). However, eIF4E phosphorylation remained

unchanged throughout the infection cycle (Figures 3C and F). These results, consistent

for both the variant infections, indicated that SARS-CoV-2-induced polysome collapse is

not mediated through ERK1/2-eIF4E.

Next, we analyzed the status of p38 MAPK during the infection. Unlike ERK1/2, p38

MAPK phosphorylation was induced during SARS-CoV-2 infection from 48 hpi in both

variant infections (Figures 4 A-D). We tested if p38 MAPK could be an important player

in SARS-CoV-2 replication by inhibiting p38 MAPK activity in SARS-CoV-2-infected

Caco2 cells. Inhibition was confirmed by the reduced eIF4E phosphorylation (Figure 4E).

Despite a substantial and significant drop in SARS-CoV-2 RNA in the supernatants of

inhibitor-treated cultures (Figure 4F), the infectious titers remained unchanged (Figure 4

G). Unlike other reports (35,36), our results ruled out any significant impact of p38 on

SARS-CoV-2 translation. Collectively, these results suggested that while p38 MAPK may

be required for viral RNA replication, it does not play an active role in virus production.

SARS-CoV-2 strongly inhibits mTORC1, but is unperturbed by the inhibition

As eIF2α and MAPK pathways were ruled out for their potential role in the polysome

collapse during SARS-CoV-2 infection, we extended our studies to mTORC1. 4EBP1, a

key substrate of mTORC1 and an important mediator of translation initiation, was used

as the indicator of mTORC1 activity. The kinetics of 4EBP1 phosphorylation during

SARS-CoV-2 infection in Caco2 cells was analyzed. We noticed a marked

dephosphorylation of 4EBP1 in SARS-CoV-2 infected cells beyond 48 hpi (Figures 5 A-

D), suggesting an inhibition of mTORC1 during infection. Both B.1.1.8 and Delta

infections led to a comparable and sustained inhibition of mTORC1 activity, which

coincided temporally with the observed polysome collapse. Since mTORC1 inhibition and

polysome collapse shared similar time-frames, SARS-CoV-2 appears to inhibit mTORC1

in order to subvert the host translation machinery. The significance of this inhibition on

the viral replication was further studied by inhibiting mTORC1 during the 24-hrs of

infection by Rapamycin (Figure 5E). Though a noticeable drop in 4EBP1 phosphorylation

was observed in Rapamycin treated cells, no significant change in N levels, or viral RNA

or infectious titers were observed (Figures 5F and G). To further test the role of mTORC1

on SARS-CoV-2, we activated mTORC1 using MHY1485, and analyzed its impact on

SARS-CoV-2 replication. Infected cells were treated with MHY1485 (10 µM) at 24 hpi and

further continued up to 48 hpi. Despite a strong activation of mTORC1 and 4EBP1

phosphorylation with MHY1485, no significant change in viral RNA, N, or viral titers was

observed (Figure 5H-J). These results confirmed that the inhibition of mTORC1 seen in

SARS-CoV-2 infection has no consequence to the viral translation. Together, these

results demonstrate that mTORC1 inhibition represents a conserved host response

during infection and does not significantly alter viral replication efficiency.

5' TOP mRNAs escape translation repression and mTORC1 suppression

Polysome profiles of SARS-CoV-2 infected cultures showed no loss in overall ribosome

abundance (Figures 1C and F). Translation of the mRNAs of ribosomal proteins (RPs) is

regulated through the 5' TOP element. Since the 5' TOP mRNAs are regulated by

mTORC1, we asked if the inhibition of mTORC1 pathway and translational repression

negatively impacts ribosomal biogenesis by restricting the translation of the 5' TOP

transcripts.

We assessed cell lysates from mock- and SARS-CoV-2-infected cells for two ribosomal

proteins, rpS3 (ribosomal protein S3) and rpL26 (ribosomal protein L26), as

representatives for both subunits and 5' TOP transcripts. Sustained levels of rpS3 were

observed in the infected samples of both variants throughout the infection compared to

the mock (B.1.1.8- Figures 6 A-B, Delta- Figures 6H-I). Notably, the levels of rpL26

increased with the progress of infection in the B.1.1.8 variant (B.1.1.8- Figures 6A-C,

Delta- Figures 6H-J). Thus, despite a strong polysome dissociation and inhibition of

mTORC1, ribosomal protein synthesis goes on unperturbed indicating that inhibition of

mTORC1 activity is not affecting the translation of 5' TOP mRNAs.

We next measured the cellular abundance and their ribosomal loading (association with

translating ribosomes) of these two RP transcripts by quantitative RT-PCR. Cellular RNA

and RNA from polysome fractions from across all time points after infection with both

B.1.1.8 and Delta variants were analyzed. Steady-state levels of *RPS3* transcripts were

observed in both B.1.1.8 and Delta variant infected cells (Figures 6D and K), and the

transcripts were also associated with the polysomes throughout the infection (Figures 6E

and L). Higher *RPL26* transcript levels were observed in the cells infected with both

variants from 48 hpi (Figures 6 F and M). These transcripts were loaded onto polysome

at a higher efficiency in cells infected with either of the variants, (Figures 6G and N).

Subsequently, we studied the levels of viral RNA in cellular and polysome pools from

B.1.1.8 and Delta infected Caco2 cultures in order to monitor the viral replication and the

association of viral transcripts with the translating ribosomes. The abundance of viral RNA

levels peaked at 48 hpi for both B.1.1.8 and Delta variants (Supplementary Figures S4A

and C). A continuous association of viral RNAs with the polysomes was also observed in

both variants (Supplementary Figures S4B and D), indicating unperturbed translation of

viral RNA despite a significant depletion of the translating polysome pool. Overall, these

results indicate that mTORC1 inhibition by SARS-CoV-2 infection does not affect the

polysome loading of the viral RNA as well as ribosomal protein transcripts, and their

translation, despite an mTORC1-suppressed environment.

DISCUSSION

Several studies have demonstrated that SARS-CoV-2 infection suppresses host protein

translation (3,5,7,14,37), with Nsp1 now being a well-documented driver of this effect by

associating with 40S ribosomes to block host mRNA entry (7,16). Nsp2 also contributes

to host immune suppression by stimulating viral translation even under hypoxic

conditions, while specifically inhibiting interferon translation through interaction with

specific translation initiation factors (38,39). Other viral proteins such as Orf9b

significantly impact the innate immune system to regulate host responses, collectively

inhibiting translation (7,34,40). However, the detailed picture of the complex, multilayered,

regulatory networks that are frequently modulated by various viruses in order to establish

their authority over the process of protein translation has not been established yet. Our

study was aimed at understanding the overall picture of translation machinery and to gain

insight into the signal pathways that normally regulate the machinery. We monitored

polysome profiling to capture the polysome association in SARS-CoV-2 infected colon

epithelial cell line Caco2 at multiple time intervals. These results were subsequently

correlated with the modulation of the key regulatory pathways that control translation

initiation. Using two variants of SARS-CoV-2, We identified that SARS-CoV-2 controls the

translation machinery through the inhibition of mTORC1 pathway, while eIF2α and MAPK

pathways are not affected in ways that could inhibit translation. Our results indicate that

SARS-CoV-2 inhibits mTORC1, which is likely behind a massive collapse of polysomes

thereby limiting translation of host transcripts. Interestingly, mTORC1 inhibition does not

impact the polysome association of viral transcripts, and transcripts encoding ribosomal

proteins. Our results find that the virus uses mechanisms to prevent polysome association

that primarily targets host transcript while making sure that its own translation and those

of ribosomal proteins are protected through mechanisms yet to be understood. The

results were consistent between the two variants, highlighting the conserved mechanism

targeted by SARS-CoV-2 in its pursuit to control host translation apparatus.

Similar to a previously published study (16), we did not observe any change in any of the

three pathways that regulate translation during the early hours of infection, indicating any

significant change on host translation signaling onsets much later. The only pathway that

was impacted by SARS-CoV-2 was mTORC1. Unlike other reports that depict activated

mTORC1 (41,42) we consistently observed an inhibition of this pathway with both the

variants. Pharmacological activation of this pathway however did not have an impact on

viral titers despite a curious drop in RNA levels, which indicates that the virus is able to

sustain its translation despite an impact on its replication. Similar responses have been

reported for other viruses such as Dengue (DENV). DENV translation, despite being cap-

dependent, is unaffected by mTORC1 inhibition (43) Notwithstanding a strong inhibition

of mTORC1 by the virus, translation of 5' TOP transcripts, which are chiefly regulated by

mTOR pathways, continued unhindered. This indicates a concerted effort by the virus to

target specific transcripts regulated by mTORC1 without compromising on the

requirements of the virus. It would be interesting to identify the host transcripts that are

specifically targeted by the virus, which might lead to understanding the precise

mechanisms by which this discrimination is established.

p38 and ERK1/2 MAPK signaling converge at cap-binding eIF4E to regulate translation

initiation. Both kinases have a wide array of substrates that regulate a large number of

processes including cell survival, stress, apoptosis, and interferon production. In our

study, we noted that only ERK1/2, but not p38 MAPK, was dephosphorylated by SARS-

CoV-2-mediated signaling activities. p38 activation remained sustained during both

variant infections. The decreased viral replication upon p38 inhibition indicates SARS-

CoV-2's dependence on p38 activity for continued viral activities. This has been observed

by other groups as well (44,45). Activated p38 promotes viral replication, influences innate

immune signalling, and therefore viral infectivity (44). The impact of p38 inhibition on

SARS-CoV-2 replication and inflammatory cytokine levels that has been observed by us

and other groups (45) is possibly why its activation status is unaffected throughout

evolution of the virus. Between the activated p38 MAPK and dephosphorylated state of

ERK1/2, the lack of significant impact on eIF4E phosphorylation is expected. Since eIF4E

phosphorylation is understood to affect only a select set of mRNAs translationally (46),

we believe that its contribution to the global suppression of translation activities caused

by SARS-CoV-2 infection could be limited and more studies are necessary to determine

its impact.

It is fascinating to note that SARS-CoV-2 infection manages to avoid eIF2α

phosphorylation. eIF2α is phosphorylated by one of its four kinases most of which are

activated upon various stress exerted on the cell. RNA viruses often impart intense stress

on ER that is relayed to PERK (47,48). PKR, one of the dsRNA sensors is often activated

by RNA viral infections. These observations indicate that SARS-CoV-2 depends on the

canonical mechanism of translation initiation that requires the availability of active ternary

complexes, which eIF2α is a part of. Intriguingly, one report found that PKR inhibition

could work as a potent antiviral against SARS-CoV-2(49), suggesting that despite any

purported effect on eIF2α phosphorylation in our study, its kinases may still be involved

in curbing SARS-CoV-2 infection. Since eIF2α phosphorylation results in the inhibition of

new initiation events that would adversely affect the translation of viral transcripts as well,

SARS-CoV-2 might have evolved strategies to bypass this modification. One such report

suggests that Nsp3 and Nsp4 of SARS-CoV-2 suppress PERK, which is also involved in

the unfolded protein response pathway (50), which could also impact global translation

through eIF2α. Another report indicates that SARS-CoV-2 inhibits stress granule

assembly, a process largely regulated by eIF2α, and this may be a mechanism to allow

continued viral replication (51). Together, we believe that the inhibited status of eIF2α

found in both variants during active infection allows viral translation to continue

unimpeded.

To summarize our findings, we observe the complex relation between host translation

regulation pathways in response to viral infection. The most striking impact was observed

on mTOR pathway although this did not deter translation of certain key mTOR substrates

such as 5'TOP transcripts. Surprisingly, unlike other RNA viruses, eIF2α phosphorylation

does not seem to be sustained or increased along the course of infection. Our study

focuses on the interplay between host and viral systems after the establishment of active

infection, as opposed to other studies that have focused on its immediate impact. Our

observations clarify that there is a dynamic rewiring in cellular signalling in response to

virus-mediated effects on translation. Findings from our lab also indicate at novel means

by which SARS-CoV-2 evades host innate immune responses as well that may also

explain how some of our observations differ from other RNA viruses (Tandel et al).

LIMITATIONS

While this study establishes a conserved mechanism of host translational suppression

by distinct SARS-CoV-2 variants, it is limited by the use of a single epithelial cell model,

which may not fully capture the heterogeneity of infection across different tissue

environments. The experiments were performed with pre-Omicron lineages, and it

remains to be determined whether newer variants exhibiting altered pathogenicity retain

similar translational control or not. Furthermore, although our findings clearly links

mTORC1 inhibition with polysome collapse , the specific viral or host determinants

driving this effect are not dissected here and needs further investigation. Nonetheless,

by integrating polysome profiling with signalling analyses, this study provides a

comprehensive framework that advances understanding of how SARS-CoV-2

modulates host translation.

Experimental procedures

Antibodies and inhibitors

All primary antibodies were purchased from Cell Signaling Technologies except the

GAPDH and anti-SARS-CoV-2 Nucleocapsid were procured from Thermo Fisher

Scientific. HRP-conjugated anti-rabbit and anti-mouse secondary antibodies were

purchased from Jackson ImmunoResearch. Rapamycin and MHY1485 were from Sigma,

whereas the p38 VIII inhibitor and Sodium arsenite were from Merck Millipore.

Cell culture

Caco2, human colorectal adenocarcinoma epithelial cells were procured from ATCC and

were cultured in Dulbecco's Modified Eagle's Medium (DMEM; from Gibco) with 20%

Fetal Bovine Serum (FBS; Hyclone), 1× penicillin-streptomycin cocktail (Pen-strep,

Gibco), and 1× non-essential amino acids (NEAA, Gibco), at 37°C and 5% CO_2. Cells

were continuously passaged at 70-80% confluency and mycoplasma contamination was

monitored periodically.

SARS-CoV-2 Infection and quantification

Two Indian isolates of SARS-CoV-2 strains were used in this study (GSSAID id:

EPI_ISL_458046, (B.1.1.8), and EPI_ISL_2775201 (Delta) (52,53). All the viral cultures

were propagated in Vero (CCL-81) cells in serum and antibiotics-free conditions. Caco2

cells were infected at 1 MOI for 3 hrs in serum-free conditions after which the media was

replaced with complete media and further incubated until the time of harvesting as per

experiment requirement. At the time of harvesting, the cells were first trypsinized and

collected separately for protein and RNA study. The extracellular RNA from the

supernatants was isolated (MACHEREY-NAGEL GmbH & Co. KG) and the SARS-CoV-

2 RNA was quantified using a commercial kit (LabGun™ COVID-19 RT-PCR Kit)

following manufacturers protocol in Roche LightCycler 480. The viral E gene was

normalized with internal control provided in the kit, and fold changes were calculated

using $2^{(-\Delta\Delta Ct)}$ method. The infectious viral particle numbers in the cell culture supernatant

were quantified using the plaque-forming unit (PFU/mL) assay. Briefly, the supernatant

was log diluted (10^{-1}-10^{-7}) in 1× serum-free DMEM and used for infecting Vero monolayer-

grown twelve-well plates. 3 hpi, the cells were briefly washed and were overlaid with

agarose: DMEM mix (in 1:1 ratio; 2 x DMEM with 5% FBS and 1% penicillin-streptomycin

mixed with equal volumes of 2% LMA), after which the plates were incubated undisturbed

for 6 days at 37°C. Later, the cells were fixed with 4% formaldehyde and stained with

crystal violet. The clear zones were counted and PFU was calculated as PFU/mL.

Inhibitions and infection

For the inhibition experiments, 0.45×10^6 Caco2 cells were seeded in a six-well format

and 24 hrs later the cells were infected with SARS-CoV-2 variants, at 1 MOI for 3 hrs in

serum-free media. For rapamycin inhibition, the infection media was replaced with serum-

sufficient media containing 25 nM Rapamycin or DMSO, and incubated for 22 hrs. At the

end of the treatment, the cells were harvested and protein and RNA were prepared. For

the activation of mTORC1, cells were treated with 10 μM MHY1485 or DMSO at 24 hpi

and harvested at 48 hpi. The p38 inhibition was carried out in a similar fashion to the

MHY1485 experiment, at a final concentration of 10 μM. For ISR activation, SARS-CoV-

2 variant infected cells were treated with 10 μM of sodium arsenite or water for 8 hrs at

16 hpi. The intracellular and extracellular RNA were subjected to qRT-PCR, and the

protein lysates were subjected to western blotting for confirmation of inhibition or

activation.

Polysome preparation and RNA precipitation

Polysomes were fractionated as explained elsewhere (54). Caco2 cells were grown in

175cm^2 flasks till 70% confluency and subsequently infected with SARS-CoV-2 variants

at 1 MOI. Media was changed after 2 hrs, and cells were harvested at 24, 48, 72, and 96

hpi, along with mock-infected cells grown alongside for each time point. The cells were

incubated for 5-10 minutes, harvested, and washed twice with a solution of ice-cold

1×PBS containing 100 µg/mL cycloheximide, to freeze the polysomes on the mRNAs.

They were subsequently lysed in polysome lysis buffer containing 20 mM Tris-Cl pH 8.0,

140 mM KCl, 1.5 mM MgCl$_2$, 0.5 mM DTT, 1% Triton X-100, 1× protease inhibitor, 0.5

mg/L heparin,100 µg/mL cycloheximide, and RNase inhibitor. Crude RNA was quantified

using a spectrophotometer, and 90 µg was layered onto 11 mL of 10- 50% linear sucrose

gradient (20 mM Tris-Cl pH 8.0, 140 mM KCl, 1.5 mM MgCl$_2$, 0.5 mM DTT, 100 µg/mL

cycloheximide, 1mM PMSF, 10-50% sucrose). The resulting gradients were centrifuged

in an SW41 Ti rotor (Beckman Coulter) at 35,000 r.p.m. at 4°C for 3.5 hrs. The polysome

samples were fractionated using the Teledyne ISCO fraction collector system and

absorbance was measured and graphically noted at 254 nM. Polysome profiles of mock

and infected cells for each time point were digitized and overlaid using Inkscape. Later,

total polysome RNA was precipitated by adding 1/10th volume of 3M sodium acetate and

an equal volume of absolute ethanol, incubated at -20°C and centrifuged at 16000g, and

the pellet obtained was airdried and resuspended in DEPC water. To this, 4M LiCl was

added to a final concentration of 2.5M, incubated at -20°C overnight. After centrifugation,

the pellet was washed twice with 70% ethanol, airdried and resuspended in DEPC water.

Subsequently, the RNA samples were purified using MN NucleoSpin RNA kit following

manufacturer's protocol. The abundance of polysome-associated mRNAs were quantified

through qRT-PCR.

Quantitative RT-PCR

Cellular and polysomal RNA were extracted using the Nucleospin RNA kit (MACHEREY-

NAGEL) according to the manufacturer's instructions. Equal quantities of RNA were

reverse-transcribed into cDNA using the Primescript Reverse transcriptase kit (Takara)

as per the manufacturer's protocol. Subsequently, equal amounts of cDNA were used for

specific gene expression quantification using specific primers (Table 1) and the SYBR

Green Master Mix (Takara) on a LightCycler 480 system (Roche). Host-specific

transcripts were normalized to GAPDH, while viral transcripts were normalized using the

internal control gene included in the LabGun™ COVID-19 RT-PCR Kit. Relative fold

changes between control and experimental samples were calculated using the $2^{(-\Delta\Delta Ct)}$

method and presented as graphs.

Immunoblotting

Protein pellets were lysed in 1 × Nonidet P-40 lysis buffer (1% Nonidet P-40, 50 mM Tris-

HCl, 150 mM NaCl (pH 7.5), EGTA, 1 mM sodium orthovanadate, 10 mM sodium

pyrophosphate, 100 mM NaF, and 1 mM PMSF) incubated on ice for 20 minutes with

intermittent vortexing and centrifuged at 13,000 rpm for 15 minutes at 4°C. The

supernatants containing the proteins were collected and quantified using BCA reagents

(G Biosciences). Lysates were mixed with 6× denaturing dye and the proteins were

resolved using SDS-PAGE and transferred to PVDF membranes. The membranes were

blocked in 5% BSA dissolved in 1× TBST before the addition of primary antibodies.

Primary antibodies against the proteins of interest were diluted in the blocking buffer,

added to the membrane, and incubated overnight at 4°C. Later, the membranes were

washed in 1× TBST, secondary antibodies conjugated with HRP were added and the

blots were developed on a Bio-Rad Chemidoc MP system using SuperSignal West Pico

PLUS (Thermo Fisher) and SuperSignal West Femto Maximum Sensitivity (Thermo

Fisher) chemiluminescent substrate kits.

Statistical analysis

For each experiment, at least three independent replicates (until specified) were used to

calculate mean ± SEM and plotted graphically wherever indicated. Statistical significance

was measured using a two-tailed, unpaired Student t-test, and the resultant p values were

represented as *, **, *** indicating p values ≤ 0.05, 0.005, and 0.0005, respectively.

Data availability

All data pertaining to this manuscript are included within the manuscript.

Institutional biosafety

Institutional biosafety clearance was obtained by K.H.H., for the experiments pertaining

to SARS-CoV-2.

Supporting information

This article contains supporting information.

Acknowledgements

We thank Mohan Singh Moodu and Amit Kumar for assisting with logistics and Karthika

S Nair for her assistance with some experiments.

Funding

This study was supported by a grant from Council of Scientific and Industrial Research

(CSIR), Govt. of India (6/1/FIRST/2020-RPPBDD-TMD-SeMI) to K.H.H. H.P, D.G., and

D.K received research fellowship from CSIR. V.S received research fellowship from the

Department of Biotechnology (DBT), Govt. of India.

Contributions

The experiments were conceived by H.P., D.G., and K.H.H. H.P., and D.G., performed

polysome profiling, and qRT-PCRs. D.G., D.K., and V.S prepared infectious SARS-CoV-

2, performed infections, quantified them and analyzed data. H.P., A.P.S., D.G., D.K., and

A.H.S. performed immunoblotting. H.P., and D.G., assisted K.H.H in writing the

manuscript.

Conflicts of interest

The authors declare that they have no conflicts of interest with the contents of this article.

Table 1

Primer name	Primer sequence 5'-3'
GAPDH RT For	ATGGGGAAGGTGAAGGTCG
GAPDH RT Rev	GGGGTCATTGATGGCAACAATA
rpL26 For	GAAGGGAGCCTGTGGAAATAGA
rpL 26 Rev	GTGGTCTGGAAATTCCGTGAAGAC
rpS3 For	GCAAGATGGCAGTGCAAATATCC
rpS3 Rev	CTCGCACCTCAACTCCAGAGTAGC

Figure 1

Figure 2

A

B

C

D

E

F

G

Figure 3

A

B

C

D

E

F

Figure 4

A

B.1.1.8	24h	48h	72h	96h	
SARS-CoV-2	- +	- +	- +	- +	kDa

T180/ Y182 p38 — 43, 33

p38

β-Tubulin — 54

B

C

Delta	24h	48h	72h	96h	
SARS-CoV-2	- +	- +	- +	- +	kDa

T180/ Y182 p38 — 43, 33

p38 — 43, 33

GAPDH — 43, 33

D

E

p38 VIII	- +	- +	- +	
B.1.1.8	- -	+ +	- -	
Delta	- -	- -	+ +	kDa

S209 eIF4E — 29

eIF4E

Nucleocapsid — 54, 43

GAPDH — 33

F

G

Figure 5

Figure 6

FIGURE LEGENDS

Figure 1: SARS-CoV-2 causes major polysomal collapse. (A, D) Immunoblot analysis of mock and B.1.1.8 **(A)** or Delta **(D)** infected Caco2 cells over 24, 48, 72, and 96 hrs of infection. Cell lysates were separated by SDS-PAGE and probed for SARS-CoV-2 nucleocapsid protein. **(B, E)** Polysome profiles of Caco2 cells infected with B.1.1.8 and Delta for 24, 48, 72, and 96 hrs. The cells were treated with 100 µg/mL CHX before harvesting and lysed in polysome lysis buffer. Equal quantities of lysates were layered onto continuous sucrose gradients ranging from 10-50%, subjected to ultracentrifugation, and fractionated along with measuring absorbance at 254 nm. The digitized profiles for infected and uninfected samples for each time point were overlaid to assess any change in global translation levels. The profiles were digitalized using Inkscape software. Arrows indicate inferred positions of 40S, 60S, and 80S peaks, based on known sedimentation standards (George et al., J Biol Chem, 2012) **(C, F)** Graphs showing % area under the curve (AUC) of mock and SARS-CoV-2 infected polysome profiles over 24, 48, 72, and 96 hrs of infection for B.1.1.8 and Delta respectively. The area under the curve was calculated using ImageJ software. The polysome area, excluding mRNPs fractions, was normalized to the profile's total area. Graphs represent data from three independent sets for B.1.1.8 **(B)** and Delta **(E)**, respectively. **(G)** and **(H)** Graphs showing percentage cell viability of infected cells compared to mock at different time intervals from B.1.1.8 and Delta infection respectively. All values are plotted as mean ± SEM. p-values were calculated using Student's t-test and represented as * and **, indicating p-values ≤ 0.05 and 0.005, respectively.

Figure 2: SARS-CoV-2 does not induce ISR-eIF2α axis. (A, C) Immunoblots showing changes in phosphorylation of eIF2α at S52, and its expression in Caco2 cells over different time points of infection with B.1.1.8 **(A)** and Delta **(C)**, respectively. **(B, D)** Densitometric analysis from the panel for B.1.1.8 **(B)** and Delta **(D)**, respectively. **(E)** SARS-CoV-2 variants infected Caco2 cells were treated with sodium arsenite at 10 μM concentration for 8 hrs starting at 16 hpi. The effect of treatment was scored by the phosphorylation status of eIF2α at S52, and the abundance of viral protein under arsenite treatment was also studied. **(F)** Relative fold change in SARS-CoV-2 viral Envelope levels in infected cells treated with either vehicle or 10 µM arsenite. Value from vehicle-treated cells were used for normalization, and the relative fold change is represented graphically. **(G)** Graph showing changes in viral titers with vehicle or arsenite treatment quantified as PFU/mL. Graphs represent data from at least three sets and are plotted as mean ± SEM. p-values are represented as *, ** and ***, indicating p-values ≤ 0.05, 0.005 and 0.0005, respectively, n.s. represents non-significant.

Figure 3: MAPK-eIF4E axis does not mediate polysome collapse during 5vfr34bCoV-2 infection. (A) Immunoblots depicting phosphorylation and expression status of ERK1/2 and eIF4E from Caco2 cells infected with B.1.1.8 for 24, 48, 72, and 96 hrs along with mock-infected controls. **(B)** Densitometric quantitation of T202/Y204 ERK1/2 phosphorylation from the panel in A. **(C)** Densitometric quantitation of S209 eIF4E phosphorylation from the panel in A. **(D)** Immunoblot showing changes in phosphorylation and expression status of ERK1/2 and eIF4E with Delta infection. **(E)** Densitometric quantitation of T202/Y204 ERK1/2 phosphorylation from the panel in D. **(F)** Densitometric quantitation of S209 eIF4E phosphorylation from the panel in D.

Figure 4: SARS-CoV-2 activates the p38 MAPK pathway independent of variants and this activation benefits viral replication. (A, C) Immunoblots showing changes in phosphorylation of p38-MAPK and its expression in Caco2 cells over different time points when infected with B.1.1.8 (A) and Delta (C), respectively. **(B, D)** Densitometric quantification of T180/Y182 phosphorylation from the panel in A (B) and panel in C (D), respectively. **(E)** SARS-CoV-2 variants infected Caco2 cells were treated with p38 VIII inhibitor (p38i) at 10 µM concentration for 24 hrs beginning at 24 hpi until harvesting. The inhibition was scored by the dephosphorylation status of eIF4E and viral protein abundance under p38 inhibited environment was also studied. **(F)** Relative fold change in SARS-CoV-2 extracellular E gene RNA, in DMSO and p38-inhibited cells. **(G)** Infectious viral titer measure in DMSO and p38-inhibited supernatants, quantified as PFU/mL. Graphs represent data from at least three sets and are plotted as mean ± SEM. p-values are represented as *, ** and ***, indicating p-values ≤ 0.05, 0.005 and 0.0005, respectively, n.s. represents non-significant.

Figure 5: SARS-CoV-2 strongly inhibits mTORC1, but is unperturbed by the inhibition (A, C) Immunoblots showing changes in phosphorylation of 4EBP1 and its expression in Caco2 cells over different time points when infected with B.1.1.8 (A) and Delta (C), respectively. **(B, D)** Densitometric quantification of T37/46 phosphorylation from the panel in A (B) and panel in C (D), respectively. **(E)** Caco2 cells were infected with SARS-CoV-2 variants for two hrs, treated with either DMSO or 25nM of Rapamycin up to 24 hpi, and harvested. A similar setup in mock cells was used. Inhibition was assessed by the drop in 4EBP1 phosphorylation in Rapamycin-treated mock and infected background. **(F)** Graph depicting relative fold changes in intracellular viral Envelope gene

levels over rapamycin treatment in SARS-CoV-2 variant infected Caco2 cells. **(G)** Graph

showing levels of viral titer with rapamycin treatment quantified as PFU/mL. **(H)** Mock and

SARS-CoV-2 variants infected Caco2 cells were treated with either DMSO or 10 µM

MHY1485 at 24 hpi and harvested at 48 hpi, and successful activation was assessed by

increased phosphorylation of 4EBP1. **(I)** Graph showing relative fold changes in

intracellular viral Envelope gene levels over MHY1485 treatment. **(J)** Graph showing

levels of viral titer with MHY1485 treatment quantified as PFU/mL. Graphs represent data

from at least three sets and are plotted as mean ± SEM. p-values are represented as *,

** and ***, indicating p-values ≤ 0.05, 0.005 and 0.0005, respectively, n.s. represents non-

significant.

Figure 6: 5'TOP mRNAs escape translation repression despite subdued mTORC1

activity during infection. (A) Immunoblot analysis of ribosomal protein rpS3 and rpL26

throughout infection with B.1.1.8. **(B, C)** Graphs showing densitometric quantification of

rpS3 (B), and rpL26 (C) levels from the panel in A. **(D, E)** Cellular abundance of *RPS3*

(D) and changes in its polysome association (E) with B.1.1.8 infection. **(F, G)** Cellular

abundance of *RPL26* (F) and changes in its polysome association (G) with B.1.1.8

infection **(H)** Immunoblot showing the expression level of rpS3 and rpL26 under Delta

infection in Caco2 cells **(I, J)** Graphs showing densitometric quantification of rpS3 and

rpL26 levels from the panel in H, n=2. **(K, L)** Changes in cellular abundance of *RPS3* (K),

and changes in its polysome association (L) with Delta infection. **(M, N)** Changes in

cellular levels of *RPL26* (M) and changes in its polysome association with Delta infection.

Graphs represent data from at least three sets until specified and are plotted as mean ±

SEM. p-values are represented as *, ** and ***, indicating p-values ≤ 0.05, 0.005 and

0.0005, respectively, n.s. represents non-significant.

References

1.   Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with
a new coronavirus of probable bat origin. Nature. 2020 Mar;579(7798):270–3.

2.   https://www.worldometers.info/coronavirus/. 2024.

3.   V'kovski P, Kratzel A, Steiner S, Stalder H, Thiel V. Coronavirus biology and replication:
implications for SARS-CoV-2. Nat Rev Microbiol. 2021 Mar;19(3):155–70.

4.   Thoms M, Buschauer R, Ameismeier M, Koepke L, Denk T, Hirschenberger M, et al. Structural
basis for translational shutdown and immune evasion by the Nsp1 protein of SARS-CoV-2.
Science. 2020 Sep 4;369(6508):1249–55.

5.   Walsh D, Mohr I. Viral subversion of the host protein synthesis machinery. Nat Rev Microbiol.
2011 Oct 17;9(12):860–75.

6.   Jan E, Mohr I, Walsh D. A Cap-to-Tail Guide to mRNA Translation Strategies in Virus-Infected
Cells. Annu Rev Virol. 2016 Sep 29;3(1):283–307.

7.   Finkel Y, Gluck A, Nachshon A, Winkler R, Fisher T, Rozman B, et al. SARS-CoV-2 uses a
multipronged strategy to impede host protein synthesis. Nature. 2021 Jun;594(7862):240–5.

8.   Kamitani W, Huang C, Narayanan K, Lokugamage KG, Makino S. A two-pronged strategy to
suppress host protein synthesis by SARS coronavirus Nsp1 protein. Nat Struct Mol Biol. 2009 Nov
18;16(11):1134–40.

9.   Nakagawa K, Lokugamage KG, Makino S. Viral and Cellular mRNA Translation in Coronavirus-
Infected Cells. Adv Virus Res. 2016;96:165–92.

10.   Banerjee AK, Blanco MR, Bruce EA, Honson DD, Chen LM, Chow A, et al. SARS-CoV-2 Disrupts
Splicing, Translation, and Protein Trafficking to Suppress Host Defenses. Cell. 2020 Nov
25;183(5):1325-1339.e21.

11.   Stern-Ginossar N, Thompson SR, Mathews MB, Mohr I. Translational Control in Virus-Infected
Cells. Cold Spring Harb Perspect Biol. 2019 Mar 1;11(3).

12.   de Breyne S, Vindry C, Guillin O, Condé L, Mure F, Gruffat H, et al. Translational control of
coronaviruses. Nucleic Acids Res. 2020 Dec 16;48(22):12502–22.

13.   Lapointe CP, Grosely R, Johnson AG, Wang J, Fernández IS, Puglisi JD. Dynamic competition
between SARS-CoV-2 NSP1 and mRNA on the human ribosome inhibits translation initiation. Proc
Natl Acad Sci U S A. 2021 Feb 9;118(6).

14.   Schubert K, Karousis ED, Jomaa A, Scaiola A, Echeverria B, Gurzeler LA, et al. SARS-CoV-2 Nsp1
binds the ribosomal mRNA channel to inhibit translation. Nat Struct Mol Biol. 2020
Oct;27(10):959–66.

15.   Thoms M, Buschauer R, Ameismeier M, Koepke L, Denk T, Hirschenberger M, et al. Structural
basis for translational shutdown and immune evasion by the Nsp1 protein of SARS-CoV-2.
Science. 2020 Sep 4;369(6508):1249–55.

16.   Puray-Chavez M, Lee N, Tenneti K, Wang Y, Vuong HR, Liu Y, et al. The Translational Landscape of
SARS-CoV-2-infected Cells Reveals Suppression of Innate Immune Genes. mBio. 2022 Jun
28;13(3):e0081522.

17.   Metzler M, Tharyan RG, Klann K, Grikscheit K, Bojkova D, Cinatl J, et al. SARS-CoV-2 Variants Show
Different Host Cell Proteome Profiles With Delayed Immune Response Activation in Omicron-
Infected Cells. Mol Cell Proteomics. 2023 May;22(5):100537.

18.   Carabelli AM, Peacock TP, Thorne LG, Harvey WT, Hughes J, COVID-19 Genomics UK Consortium,
et al. SARS-CoV-2 variant biology: immune escape, transmission and fitness. Nat Rev Microbiol.
2023 Mar;21(3):162–77.

19.   Jefferies HB, Reinhard C, Kozma SC, Thomas G. Rapamycin selectively represses translation of the
"polypyrimidine tract" mRNA family. Proc Natl Acad Sci U S A. 1994 May 10;91(10):4441–5.

20.   Thoreen CC, Chantranupong L, Keys HR, Wang T, Gray NS, Sabatini DM. A unifying model for
mTORC1-mediated regulation of mRNA translation. Nature. 2012 May 2;485(7396):109–13.

21.   Hsieh AC, Liu Y, Edlind MP, Ingolia NT, Janes MR, Sher A, et al. The translational landscape of
mTOR signalling steers cancer initiation and metastasis. Nature. 2012 Feb 22;485(7396):55–61.

22.   Hay N, Sonenberg N. Upstream and downstream of mTOR. Genes Dev. 2004 Aug 15;18(16):1926–
45.

23.   Ma XM, Blenis J. Molecular mechanisms of mTOR-mediated translational control. Nat Rev Mol
Cell Biol. 2009 May;10(5):307–18.

24.   Hara K, Yonezawa K, Weng QP, Kozlowski MT, Belham C, Avruch J. Amino acid sufficiency and
mTOR regulate p70 kinase and eIF-4E BP1 through a common effector mechanism. J Biol Chem.
1998 Jun 5;273(23):14484–94.

25.   Zoncu R, Efeyan A, Sabatini DM. mTOR: from growth signal integration to cancer, diabetes and
ageing. Nat Rev Mol Cell Biol. 2011 Jan;12(1):21–35.

26.   Pyronnet S, Imataka H, Gingras AC, Fukunaga R, Hunter T, Sonenberg N. Human eukaryotic
translation initiation factor 4G (eIF4G) recruits mnk1 to phosphorylate eIF4E. EMBO J. 1999 Jan
4;18(1):270–9.

27.   Joshi B, Cai AL, Keiper BD, Minich WB, Mendez R, Beach CM, et al. Phosphorylation of eukaryotic
protein synthesis initiation factor 4E at Ser-209. J Biol Chem. 1995 Jun 16;270(24):14597–603.

28.   Scheper GC, Proud CG. Does phosphorylation of the cap-binding protein eIF4E play a role in
translation initiation? Eur J Biochem. 2002 Nov;269(22):5350–9.

29.   Lazaris-Karatzas A, Montine KS, Sonenberg N. Malignant transformation by a eukaryotic initiation
648   factor subunit that binds to mRNA 5' cap. Nature. 1990 Jun 7;345(6275):544–7.

649   30.   Krishnamoorthy T, Pavitt GD, Zhang F, Dever TE, Hinnebusch AG. Tight binding of the
650   phosphorylated alpha subunit of initiation factor 2 (eIF2alpha) to the regulatory subunits of
651   guanine nucleotide exchange factor eIF2B is required for inhibition of translation initiation. Mol
652   Cell Biol. 2001 Aug;21(15):5018–30.

653   31.   Pakos-Zebrucka K, Koryga I, Mnich K, Ljujic M, Samali A, Gorman AM. The integrated stress
654   response. EMBO Rep. 2016 Oct;17(10):1374–95.

655   32.   Rojas M, Arias CF, López S. Protein kinase R is responsible for the phosphorylation of eIF2alpha in
656   rotavirus infection. J Virol. 2010 Oct;84(20):10457–66.

657   33.   Bechill J, Chen Z, Brewer JW, Baker SC. Coronavirus infection modulates the unfolded protein
658   response and mediates sustained translational repression. J Virol. 2008 May;82(9):4492–501.

659   34.   Tandel D, Sah V, Singh NK, Potharaju PS, Gupta D, Shrivastava S, et al. SARS-CoV-2 Variant Delta
660   Potently Suppresses Innate Immune Response and Evades Interferon-Activated Antiviral
661   Responses in Human Colon Epithelial Cells. Microbiol Spectr. 2022 Oct 26;10(5):e0160422.

662   35.   Higgins CA, Nilsson-Payant BE, Bonaventure B, Kurland AP, Ye C, Yaron TM, et al. SARS-CoV-2
663   hijacks p38β/MAPK11 to promote virus replication. mBio. 2023 Jun 22;

664   36.   Faist A, Schloer S, Mecate-Zambrano A, Janowski J, Schreiber A, Boergeling Y, et al. Inhibition of
665   p38 signaling curtails the SARS-CoV-2 induced inflammatory response but retains the IFN-
666   dependent antiviral defense of the lung epithelial barrier. Antiviral Res. 2023 Jan;209:105475.

667   37.   Shang J, Wan Y, Luo C, Ye G, Geng Q, Auerbach A, et al. Cell entry mechanisms of SARS-CoV-2.
668   Proc Natl Acad Sci U S A. 2020 May 26;117(21):11727–34.

669   38.   Xu Z, Choi JH, Dai DL, Luo J, Ladak RJ, Li Q, et al. SARS-CoV-2 impairs interferon production via
670   NSP2-induced repression of mRNA translation. Proc Natl Acad Sci U S A. 2022 Aug
671   9;119(32):e2204539119.

672   39.   Korneeva N, Khalil MI, Ghosh I, Fan R, Arnold T, De Benedetti A. SARS-CoV-2 viral protein Nsp2
673   stimulates translation under normal and hypoxic conditions. Virol J. 2023 Mar 30;20(1):55.

674   40.   Alexander MR, Brice AM, Jansen van Vuren P, Rootes CL, Tribolet L, Cowled C, et al. Ribosome-
675   Profiling Reveals Restricted Post Transcriptional Expression of Antiviral Cytokines and
676   Transcription Factors during SARS-CoV-2 Infection. Int J Mol Sci. 2021 Mar 25;22(7).

677   41.   Appelberg S, Gupta S, Svensson Akusjärvi S, Ambikan AT, Mikaeloff F, Saccon E, et al.
678   Dysregulation in Akt/mTOR/HIF-1 signaling identified by proteo-transcriptomics of SARS-CoV-2
679   infected cells. Emerg Microbes Infect. 2020 Dec;9(1):1748–60.

680   42.   Zambalde ÉP, Dias TL, Maktura GC, Amorim MR, Brenha B, Santos LN, et al. Increased mTOR
681   Signaling and Impaired Autophagic Flux Are Hallmarks of SARS-CoV-2 Infection. Curr Issues Mol
682   Biol. 2022 Dec 31;45(1):327–36.

43. Edgil D, Polacek C, Harris E. Dengue Virus Utilizes a Novel Strategy for Translation Initiation When
Cap-Dependent Translation Is Inhibited. J Virol. 2006 Mar 15;80(6):2976–86.

44. Higgins CA, Nilsson-Payant BE, Bonaventure B, Kurland AP, Ye C, Yaron TM, et al. SARS-CoV-2
hijacks p38β/MAPK11 to promote virus replication. mBio. 2023 Aug 31;14(4):e0100723.

45. Faist A, Schloer S, Mecate-Zambrano A, Janowski J, Schreiber A, Boergeling Y, et al. Inhibition of
p38 signaling curtails the SARS-CoV-2 induced inflammatory response but retains the IFN-
dependent antiviral defense of the lung epithelial barrier. Antiviral Res. 2023 Jan;209:105475.

46. Furic L, Rong L, Larsson O, Koumakpayi IH, Yoshida K, Brueschke A, et al. eIF4E phosphorylation
promotes tumorigenesis and is associated with prostate cancer progression. Proc Natl Acad Sci U
S A. 2010 Aug 10;107(32):14134–9.

47. Pavio N, Romano PR, Graczyk TM, Feinstone SM, Taylor DR. Protein synthesis and endoplasmic
reticulum stress can be modulated by the hepatitis C virus envelope protein E2 through the
eukaryotic initiation factor 2alpha kinase PERK. J Virol. 2003 Mar;77(6):3578–85.

48. Su HL, Liao CL, Lin YL. Correction for Su et al., "Japanese Encephalitis Virus Infection Initiates
Endoplasmic Reticulum Stress and an Unfolded Protein Response". J Virol. 2024 Jul
23;98(7):e0040724.

49. Jain S, Rego S, Park S, Liu Y, Parn S, Savsani K, et al. RNASeq profiling of COVID19-infected
patients identified an EIF2AK2 inhibitor as a potent SARS-CoV-2 antiviral. Clin Transl Med. 2022
Nov;12(11):e1098.

50. Davies JP, Sivadas A, Keller KR, Wojcikiewicz RJH, Plate L. SARS-CoV-2 Nonstructural Proteins 3
and 4 tune the Unfolded Protein Response. bioRxiv. 2023 Jun 12;

51. Dolliver SM, Kleer M, Bui-Marinos MP, Ying S, Corcoran JA, Khaperskyy DA. Nsp1 proteins of
human coronaviruses HCoV-OC43 and SARS-CoV2 inhibit stress granule formation. PLoS Pathog.
2022 Dec;18(12):e1011041.

52. Gupta D, Parthasarathy H, Sah V, Tandel D, Vedagiri D, Reddy S, et al. Inactivation of SARS-CoV-2
by β-propiolactone causes aggregation of viral particles and loss of antigenic potential. Virus Res.
2021 Nov;305:198555.

53. Gupta D, Ahmed F, Tandel D, Parthasarathy H, Vedagiri D, Sah V, et al. Equine immunoglobulin
fragment F(ab')2 displays high neutralizing capability against multiple SARS-CoV-2 variants. Clin
Immunol. 2022 Apr;237:108981.

54. George A, Panda S, Kudmulwar D, Chhatbar SP, Nayak SC, Krishnan HH. Hepatitis C virus NS5A
binds to the mRNA cap-binding eukaryotic translation initiation 4F (eIF4F) complex and up-
regulates host translation initiation machinery through eIF4E-binding protein 1 inactivation. J Biol
Chem. 2012 Feb 10;287(7):5042–58.

717
718