Live imaging of delamination in *Drosophila* shows epithelial cell motility and invasiveness

are independently regulated

Mikiko Inaki* and Kenji Matsuno

Department of Biological Sciences, Graduate School of Science, Osaka University, 1-1

Machikaneyama-cho, Toyonaka, Osaka 560-0043, Japan

*Corresponding author: minaki@bio.sci.osaka-u.ac.jp

Abstract

Delaminating cells undergo complex, precisely regulated changes in cell-cell adhesion, motility,

polarity, invasiveness, and other cellular properties. Delamination occurs during development

and in pathogenic conditions such as cancer metastasis. We analyzed the requirements for

epithelial delamination in *Drosophila* ovary border cells, which detach from the structured

epithelial layer and begin to migrate collectively. We used live imaging to examine cellular

dynamics, particularly epithelial cells' acquisition of motility and invasiveness, in

delamination-defective mutants during the time period in which delamination occurs in the

wild-type ovary. We found that border cells in *slow border cells* (*slbo*), a delamination-defective

mutant, lacked invasive cellular protrusions but acquired motility, while JAK/STAT-inhibited

border cells lost both invasiveness and motility. Our results indicate that invasiveness and

motility, which are cooperatively required for delamination, are regulated independently. Our

reconstruction experiments also showed that motility is not a prerequisite for acquiring

invasiveness.

Key words: delamination, cell migration, locomotion, invasion, motility

Summary statement

Live imaging of epithelial delamination reveals that cell motility and invasiveness, both of which

are required for delamination, are regulated independently and that motility is not a prerequisite

for invasiveness.

Introduction

The delamination of epithelial cells, which normally form a layered structure, occurs when the cells become motile and detach from the organized cell layer. In the developing embryo, delamination allows cells to leave the epithelial layer and move where they are needed. Disrupting this precisely regulated process results in morphological abnormalities, cancer metastasis, and other pathological conditions (Stuelten et al., 2018; Thiery et al., 2009). Delamination occurs in *Drosophila* gastrulation when the endodermal layer invaginates; epithelial cells detach from the layer and convert into mesenchymal cells to form the mesoderm (Ko and Martin, 2020; Thiery et al., 2009). In vertebrates, neural crest cells delaminate from the dorsal neural tube and migrate to form bone, neurons, glia, and other mesodermal cells (Gouignard et al., 2018; Szabó and Mayor, 2018). Delamination also occurs during cancer metastasis, in which tumor cells lose adhesion, digest epithelial-tissue basal membranes, and invade the body cavity (Stuelten et al., 2018; Thiery et al., 2009).

Although delamination is also called epithelial-to-mesenchymal transition (EMT), a term that reflects the dynamic changes in cell polarity, adhesion, and cytoskeletal and membrane structures (Campbell and Casanova, 2016; Nieto et al., 2016; Yang et al., 2020), the delaminated cells often retain apicobasal polarity and other epithelial cell characteristics. The requirements for epithelial delamination remain elusive. To define these requirements, we analyzed the

delamination of *Drosophila* border cells, which derive from the follicle-cell layer that surrounds

germline cells—that is, the oocyte and supporting nurse cells in the developing egg chambers of

the *Drosophila* ovary (Montell, 2006). The cells at the ends of the follicular layer, called pole

cells, have distinct properties. Anterior pole cells secrete signals that activate JAK/STAT

signaling in neighboring follicle cells (Silver and Montell, 2001). Follicle cells that receive

JAK/STAT signaling start expressing *slow border cells* (*slbo*), which encodes a C/EBP

transcription factor, and become motile border cells (Montell et al., 1992). After delaminating

from the follicular layer, border cells migrate collectively toward the oocyte, where they are

essential for fertilization. Although this collective migratory process has been studied extensively

as a model for cancer metastasis (Bianco et al., 2007; Cai et al., 2014; Cliffe et al., 2017; Dai et

al., 2020; Inaki et al., 2012; Wang et al., 2010; Yang et al., 2012), little is known about the

process by which border cells delaminate. Border cells retain apicobasal polarity and cell-cell

adhesion during delamination (De Graeve et al., 2012; Pinheiro and Montell, 2004; Szafranski

and Goode, 2004; Wang et al., 2018). Actin regulators such as Profilin and Fascin have been

implicated in delamination (Ghiglione et al., 2018; Lamb et al., 2020). However, the changes in

cellular properties prerequisite to delamination need further clarification.

We used live imaging to analyze the dynamic cellular changes involved in delamination.

Live imaging studies of delamination in various systems, including EMT during neural crest cell

development and at gastrulation, have described morphological features in delaminating cells

and the involvement of actomyosin contractility (Clay and Halloran, 2013; Liu et al., 2017;

Ramkumar et al., 2016; Saykali et al., 2019). However, the hierarchy of gene regulation has not

been analyzed. Here, we analyzed border cell delamination at single-cell resolution through a

live imaging system that allowed us to genetically dissect and reconstruct the complex cellular

processes of delamination. We found that *slbo*-mutant border cells, which have been considered

non-motile (Montell et al., 1992), acquire basic motility. This motility is lost in

JAK/STAT-defective border cells. We also found that forced Slbo expression in

JAK/STAT-defective border cells rescued the cells' ability to form invasive protrusions, but did

not rescue motility or delamination. These results indicate that epithelial cells acquire motility

and invasiveness separately, and that motility is not a prerequisite for invasiveness.

Results

Locomotion is impaired upon JAK/STAT inhibition, but not in *slbo* mutants

Delaminating border cells in the *Drosophila* ovary undergo a series of complex cellular changes

(Fig. 1A). After the cells are specified and distinguished from other follicle cells, they start to

send out small extensions (Fig. 1A). Their affinity to other follicle cells changes, and the shape

of the cell cluster becomes more rounded (Fig. 1A2-6). The cluster sends out front extensions,

mostly single, in the direction the cluster will migrate (Fig. 1A3-6). The extensions progressively

elongate and contract, eventually forming a large protrusion, and the body of the cluster moves

toward the protrusion and finally detaches from the epithelial layer (Fig. 2A).

We investigated the cellular properties required for epithelial cell delamination by live

imaging of border cells in delamination-defective mutants. To compare the behavior of wild-type

and delamination-defective cells during the period in which wild-type border cells become

migratory, it was necessary to define developmental egg chamber stages. Border cells are

thought to delaminate around egg-chamber stage 8 to early stage 9, but there are no objective

markers for these stages. Follicle-cell retraction has been used to define stage 9, but it is hard to

judge exactly when retraction starts. To obtain internal markers for egg chamber development,

we measured several parameters, including the length and width of the oocyte and egg chamber,

from movies of delamination in wild-type cells. We chose oocyte length as a suitable marker for

egg-chamber stages because it consistently increases over time and appears to be a major

contributor to egg chamber growth (Fig. 1B1-3). Based on our analysis of 19 movies of

wild-type delamination, we defined the delamination time as the point when the entire cluster,

including trailing cells, has detached from the follicular layer. In movies of wild-type cells, 85%

of delamination events occurred when the oocyte was 50–70 microns long (Fig. 1C). Thus, we

used oocyte length as a marker of the wild-type delamination period, corresponding to an oocyte

length of 50–70 microns (Fig. 1C).

Movies of wild-type cells were analyzed for the 30-min period prior to the delamination

point. Movies of delamination-defective mutants were analyzed for the 30-min period starting

with the first frame corresponding to the wild-type delamination period, assessed by oocyte

length. During this period, wild-type border cells form a round cluster that jiggles and forms an

extension in the direction of future migration (Fig. 2A1-3, Movie 1). In delamination-defective

mutants, we first analyzed movies of egg chambers with JAK/STAT inhibition, which causes

border cell differentiation to fail (Silver and Montell, 2001). Follicle cells next to anterior pole

cells use a receptor named Domeless (Dome) to receive ligands from the anterior pole cell and

activate JAK/STAT signaling. To inhibit JAK/STAT signaling, we expressed a dominant-negative

(DN) form of Dome receptor lacking the cytoplasmic domain (Dome$^{\Delta CYT}$) specifically in follicle

cells, where border cells are differentiated, using *slbo*-gal4 along with UAS-10×GFP as a cellular

marker (Poukkula et al., 2011; Rørth et al., 1998; Silver and Montell, 2001). The

JAK/STAT-inhibited border cells appeared to retain an epithelial morphology with apicobasal

polarity, as do other follicle cells, and were not motile (Fig. 2B1-3, Movie 2). Next, we examined

border cells in a mutant for *slbo*, a downstream target of JAK/STAT signaling in border cells

(Silver and Montell, 2001). We found that border cells in homozygotes for the hypomorphic

allele *slbo*^{1310}, in which border cell migration is almost completely abolished (Montell et al.,

1992; Rørth et al., 1998), failed to delaminate during the wild-type delamination period (Fig. 2C).

However, their behavior differed from JAK/STAT-inhibited border cells (Fig. 2B,C) in that they

formed round clusters in which individual border cells moved back and forth, similar to

locomotion (Fig. 2C1-3, Movie 3). We quantified this movement as angular velocity by

measuring the movement of border cells relative to the center of the cluster (Fig. 2F). Angular

velocity in *slbo*-mutant border cells was similar to wild-type, but was much lower in

JAK/STAT-inhibited border cells—similar, in fact, to that of non-motile follicle cells in the

posterior end (Fig. 2D,G). We confirmed the loss of cell motility in another allelic combination

of *slbo*, *slbo*^{1310}/*slbo*^{e7b}, in which border cell migration is completely blocked (Fig. S1, (Montell

et al., 1992). These results indicate that *slbo*-mutant border cells acquired motility, allowing them

to move about, even though they could not invade the egg chamber. To see whether this

locomotion in individual cells was an active process involving cytoskeletal forces, we used the

inhibitor CytochalasinD to block actin polymerization. CytochalasinD treatment completely

blocked border cell delamination (n=13) and cell movement (Fig. 2E,G, Movie 4), indicating that

the observed locomotive behavior is an active process and is required for delamination.

Invasiveness is impaired in *slbo* mutants

Although the locomotive behavior of *slbo*-mutant border cells was similar to that of wild-type

border cells, *slbo*-mutant border cell clusters failed to delaminate and had no front extensions

(Fig. 3A,B). Wild-type border-cell clusters send extensions toward the direction of future

migration (Fig. 3A). These protrusions extend and retract over time, and although a cell with

extensions is sometimes replaced by another, a cell with extensions is almost always present

during the delamination period (Movie 1). We quantified the front extensions as previously

described (Poukkula et al., 2011) with slight modifications. Briefly, we binarized Z-stack images,

separated the cluster body and extension by scanning a circle with a diameter slightly larger than

that of the cell nucleus, and measured the size (length) and persistence (duration) of the front

extensions. We found that extensions formed by wild-type border-cell clusters were up to 18

microns long and persisted through nearly the entire 30-min analysis period; in contrast,

slbo-mutant border cells had no front extensions (Fig. 3B,E,G,H, Movie 3). We concluded that

slbo-mutant border cells acquired locomotion (motility) but not invasiveness, which might

explain why they failed to delaminate. Inhibiting actin polymerization with CytochalasinD

completely blocked the formation (n=9) and maintenance (n=17) of front extensions in wild-type

border cells, showing that the extension formation and maintenance are active processes of the

actin cytoskeleton. These results indicate that wild-type delaminating border cells acquire

motility and invasiveness separately, and that invasiveness is regulated by *slbo* while motility is

regulated by other factors downstream of JAK/STAT signaling.

Invasiveness induced by *slbo* is mediated partly by *shg*

Slbo upregulates gene expression of *shotgun* (*shg*), which encodes E-cadherin (E-cad) in border

cells (Mathieu et al., 2007; Tepass et al., 1996). We investigated the *shg* mutant *shg^{PB4354}*, in

which *shg* upregulation in border cells is specifically suppressed (Mathieu et al., 2007). During

the wild-type delamination period, *shg^{PB4354}*-mutant border cells did not delaminate but had

normal locomotion, as did *slbo*-mutant border cells (Fig. 3C, Fig. S2, Movie 5). Unlike

slbo-mutant border-cell clusters, *shg^{PB4354}*-mutant border-cell clusters had front extensions (Fig.

3C, Movie 5); however, these extensions were much shorter than wild-type in both length and

duration (Fig. 3E-H), suggesting that *shg* is required both to form and to maintain invasive

extensions. These results explain why *shg^{PB4354}*-mutant border cells could not delaminate during

the wild-type delamination period. Next, we tested whether the overexpression of E-cad in

border cells, which rescues *shg^{PB4354}* mutants, could rescue *slbo*-mutant phenotypes (Mathieu et

al., 2007). The *slbo*-mutant border cells with forced E-cad expression had front extensions,

suggesting that E-cad expression induced by Slbo is responsible for border cell invasiveness (Fig.

3D, Movie 6). However, the front extensions were still significantly shorter than in wild-type

border cells (Fig. 3G,H) and E-cad expression did not rescue the impaired delamination

phenotype in *slbo* mutants, suggesting that other components downstream of *slbo* are also

involved in delamination.

Invasive protrusions must have sufficient reach to attach across distances

To determine how front extensions contribute to delamination, we examined delamination in

guidance-deficient mutants since the extensions must be guided toward the path of future

migration. Border cell migration is guided by signaling through two receptor tyrosine kinases,

PVR and EGFR, and blocking both receptors severely delays delamination (Duchek and Rørth,

2001; Duchek et al., 2001). To produce a guidance-deficient condition, we expressed DN forms

of PVR and EGFR specifically in border cells. These cells' front extensions during the wild-type

delamination period were shorter and less persistent than those of wild-type cells (Fig. 4A,D,E,

Movie 7). The guidance-deficient cells had normal locomotive behavior (Fig. 4C). However,

although the guidance-deficient cells eventually moved sideways and delaminated, in most cases

their route was less direct than that of wild-type cells (Fig. 4B, Movie 8). As the

guidance-deficient cells approached the delamination point, their front extensions and

movements were much like those observed in earlier stages of the period of analysis (Fig. 4C-E).

We further found that the distance from the anterior tip of the egg chamber to the tip of the

border-cell extension (defined as reach) was similar in wild-type and guidance-deficient mutants

at the point of delamination (Fig. 4F). Wild-type and guidance-deficient border-cell clusters

reached to approximately the same point before detaching completely, which suggests that the

extension tip reached substrates it could grab hold of to aid in detaching the cluster, and that it is

the cluster's ability to reach such a substrate, rather than the actual length of the front extension,

that is important for delamination. Consistent with this idea, the maximum length of the front

extension did not correlate with the point of delamination, and the length of the front extension at

the time of delamination varied more than the distance from the tip of the extension to the

anterior tip of the egg chamber (Fig. S3). Locomotion and front extensions were similar in

shg-mutant and guidance-deficient border cells (Fig. 3C,G,H, Fig. S2). However, *shg*-mutant

border cells could not delaminate in the same time frame as guidance-deficient mutants, which

might indicate that *shg*-mutant extensions were less able to grab onto of nurse cell substrates. Dai

et al. recently revealed that a border cell's preferential path is a juncture with multiple nurse cells

(Dai et al., 2020). We found a hot spot close to the first multi-cell juncture with more than 3

nurse cells, suggesting that the preference for multi-cell junctures comes from not just more

room, but also more membrane to grab hold of.

Motility is not a prerequisite for invasiveness

E-cad induces invasive extensions in *slbo*-mutant border cells, which normally have no

detectable extensions (Fig. 3B,G,H). We examined whether E-cad could induce invasive

extensions in JAK/STAT-inhibited border cells, which lack locomotion. Under normal conditions,

border cells appear to send out extensions after acquiring motility (Fig. 1A). To our surprise,

forced E-cad expression induced JAK/STAT-inhibited follicle-like cells to send out front

extensions, without changing the morphology of other follicle cells in most cases (Fig. 5A,

Movie 9), and the length and duration of these front extensions were very like wild-type

extensions (Fig. 5D,E). Thus, non-motile cells were able to form extensions, confirming that

motility and invasiveness are regulated independently. We investigated this possibility further by

overexpressing Slbo in JAK/STAT-inhibited border cells. JAK/STAT-inhibited follicle-like cells

with forced Slbo expression produced front extensions, as did those with E-cad (Fig. 5B, Movie

10), but other border cells were unaffected in shape and were less motile than wild-type (Fig. 5C),

supporting the idea that invasive extension does not require cell motility. These results also ruled

out the possibility that residual Slbo activity conferred motility to the $slbo^{1310}$ mutant. Both E-cad

and Slbo overexpression induced extensions that were not sufficiently invasive for successful

delamination.. Extensions induced by E-cad were very thin and grew without ever retracting,

suggesting that they lacked machineries to produce traction forces (Fig. 5A4). In contrast,

extensions induced by Slbo were thicker and retractable, suggesting that other downstream

components of Slbo serve to make the extensions functional. This conclusion is consistent with

the results of rescue experiments in *slbo* mutants (Fig. 5B4,3D).

Discussion

Epithelial cells undergo a series of cellular events as they approach delamination: Cells

de-adhere from neighboring cells, begin moving actively as individual cells, send protrusions

toward the tissue they will move into, and then detach from the original cell layer. We showed

that motility and invasiveness are acquired independently in delaminating *Drosophila* border

cells. Border cells are differentiated by JAK/STAT signaling. Downstream of JAK/STAT

signaling, the transcription factor Slbo regulates border cell invasiveness, while motility is

regulated by other downstream factors. In models of cancer metastasis, cell proliferation and

migratory behavior are thought to be genetically distinct. However, cell motility and

invasiveness are not often considered separately (Cagan et al., 2019; Stuelten et al., 2018). Here,

we revealed that motility and invasiveness are regulated independently in delamination. In

wild-type border cells, individual cells begin to jiggle slightly, develop small spikey extensions,

and eventually form a round cluster that sends out a single large extension (Fig. 1A). Thus, it

might seem that cells first acquire motility and then invade the path of migration. However, our

results showed that cell motility and invasiveness are acquired independently; thus, cell motility

is not a prerequisite for invasiveness. This was confirmed when forced Slbo expression in

JAK/STAT-inhibited border cells rescued invasive extensions without affecting cell motility (Fig.

5B-E). Although guidance-defective border cell clusters produced extensions that were smaller

and less persistent than wild-type, these clusters eventually delaminated (Fig. 4D,E) by sideways

movements (Fig. 4B). Thus, the locomotive and invasive behaviors of border cells cooperatively

contribute to delamination.

Acquiring motility involves actin cytoskeleton regulators, since inhibiting actin

polymerization diminished border cell motility (Fig. 2E,G). Border cells retain apicobasal

polarity during delamination (Pinheiro and Montell, 2004; Wang et al., 2018). Thus, border cells

may become motile and form clusters without any major changes in the distribution of proteins

involved in apicobasal polarity. Changes in adhesion between border cells and neighboring

follicle cells may affect the shape and formation of border-cell clusters. Some of these properties

might be regulated downstream of JAK/STAT signaling by transcription factors other than Slbo.

The few downstream targets of JAK/STAT signaling identified in border cells include Apontic

and Socs36, both of which modulate JAK/STAT signaling itself (Monahan and Starz-Gaiano,

2013; Starz-Gaiano et al., 2008).

The acquisition of invasiveness involves E-cad downstream of Slbo. E-cad is required in

both border cells and nurse cells for border cell migration (Mathieu et al., 2007). E-cad is

required for border cell protrusions, which are formed by homophilic interaction with E-cad on

the nurse cells. Although it might appear that E-cad upregulation in border cells is involved in

cluster integrity, E-cad upregulation is transcriptionally regulated by Slbo (Mathieu et al., 2007),

and a lack of Slbo only affects the invasiveness of the border-cell cluster (Fig. 2C). Furthermore,

forced expression of E-cad in the JAK/STAT-inhibited border-cell region did not restore

border-cell clusters to a round shape (Fig. 5A). Thus, E-cad upregulation in border cells

contributes solely to invasiveness. Cluster extensions induced by forced Slbo expression were

more functional than those induced by E-cad (Fig. 5B), suggesting that other factors are involved

in the formation, maintenance, and retraction of invasive extensions. These factors likely include

actin cytoskeleton regulators. Extensive searches for Slbo targets (Borghese et al., 2006; Wang et

al., 2006) have identified some actin-associated proteins, including Myosin VI, encoded by

jargur, and Fascin, encoded by *singed* (Borghese et al., 2006; Geisbrecht and Montell, 2002).

Myosin VI stabilizes E-cad in border cells, and Fascin regulates stiffness in nearby nurse cells

(Geisbrecht and Montell, 2002; Lamb et al., 2020; Lamb et al., 2021). To generate the necessary

traction forces for delamination, it is likely that more actin-associated proteins are involved in

border-cell extension, or that proteins that have already been identified play additional roles. To

clarify their contributions, reconstruction experiments should be conducted using multiple

downstream genes.

Materials and methods

Fly strains

All flies were raised at 25 °C with standard media. We used the following fly strains: *slbo*-gal4

(Rørth et al., 1998), *slbo^{1310}* (Montell et al., 1992), *slbo^{e7b}* (null, Rørth, 1994), *shg^{PB4354}* (Mathieu

et al., 2007), UAS-*Dome^{ΔCYT}* (Silver and Montell, 2001), UAS-*shg* (Mathieu et al., 2007),

UAS-*slbo* (Rørth et al., 2000), UAS-DN-PVR, UAS-DN-EGFR (Duchek and Rørth, 2001), and

UAS-10×GFP (Poukkula et al., 2011). We used the following genotypes for reconstruction

experiments: *slbo^{1310}/ slbo^{1310}*; UAS-*shg* / *slbo*-gal4, UAS-10×GFP, UAS-*shg*, UAS-*Dome^{ΔCYT}* /

slbo-gal4, UAS-10×GFP, and UAS-*slbo*, UAS-*Dome^{ΔCYT}* / *slbo*–gal4, UAS-10×GFP.

Live imaging and drug treatment

Live imaging was as previously described (Bianco et al., 2007). Newly emerged females of the

appropriate genotypes were collected and maintained one day on dry yeast followed by one day

on fresh yeast. Ovaries were taken in Schneider's medium (Gibco, 21720) containing 5 μg/ml of

Insulin (Sigma-Aldrich, 19278). Egg chambers were dissected out in the media and moved into

cover-glass chambers (Lab-Tek,155411) coated with 0.1 mg/ml poly-L-Lysine (Sigma-Aldrich,

P7405) using truncated pipette tips. We added 250μm of imaging media, consisting of

Schneider's medium containing 2.5% Fetal Bovine Serum (heat-inactivated, Biowest), 5 μg/ml

Insulin, 2 mg/ml Trehalose (Sigma-Aldrich, 90208), 5 μM Methoprene (Sigma-Aldrich, 33375),

1 μg/ml 20-hydroxyecdysone (Sigma-Aldrich, H5142), and 9 μM FM4-64 (Invitrogen, T13320).

Images were acquired by confocal microscopy (SP5, Leica) with a 63× objective; sections were

2.5 μm apart and covered the cluster every 2 min for 2 h. For quality control of the movies, we

used transmission images to see whether the growth and cellular integrity of the egg chamber

were maintained for 2 h. To inhibit actin polymerization, CytochalasinD (Sigma, C8273) was

added to the imaging media at a final concentration of 25 μM.

Image analysis and quantification

For egg chamber staging, we selected transmission images of single Z slices showing the

maximum egg chamber area, and measured the width and height of the egg chamber and the

oocyte. We selected oocyte length as a marker for the wild-type delamination period,

corresponding to an oocyte length of 50–70 microns. We analyzed movies of wild-type

Drosophila for 30 min prior to the delamination point; for mutants, we analyzed the 30 min

period starting with the first frame in the wild-type delamination period (assessed according to

oocyte width). For guidance-receptor mutants, we analyzed both time periods. To analyze

single-cell movement, we tracked individual cells using nuclear position, marked by absence of

GFP expression. We tracked the center of the anterior pole cells as the cluster center. We used an

Image J macro to measure changes in the angle of border cells relative to the cluster center, and

calculated the angular velocity. We analyzed front extensions using custom macros slightly

modified from Poukkula et al. (2011). After making binarized images, we manually deleted

stretched cells that linked border-cell clusters, then analyzed extensions as described previously

(Poukkula et al., 2011). We could not use the same method for rescue experiments for JAK/STAT

inhibition because the border cells did not form a round cluster, so we separated extensions by

drawing a line where the extension base touched other border cells and measured the length and

duration of the extensions. Statistical analyses were performed using Welch's or Student's t tests,

depending on the results of verification of standard deviation by F test. We calculated two-sided

p values.

Acknowledgements

We would like to thank Pernille Rørth and the Institute of Molecular and Cell Biology in

Singapore for their support. This work was supported by Sumitomo, Futaba, and Takeda Science

Foundations.

References

References

Bianco, A., Poukkula, M., Cliffe, A., Mathieu, J., Luque, C.M., Fulga, T.A., and Rørth, P. (2007).

Two distinct modes of guidance signalling during collective migration of border cells. Nature *448*,

362-365. 10.1038/nature05965.

Borghese, L., Fletcher, G., Mathieu, J., Atzberger, A., Eades, W.C., Cagan, R.L., and Rørth, P.

(2006). Systematic analysis of the transcriptional switch inducing migration of border cells. Dev

Cell *10*, 497-508. 10.1016/j.devcel.2006.02.004.

Cagan, R.L., Zon, L.I., and White, R.M. (2019). Modeling Cancer with Flies and Fish. Dev Cell *49*,

317-324. 10.1016/j.devcel.2019.04.013.

Cai, D., Chen, S.C., Prasad, M., He, L., Wang, X., Choesmel-Cadamuro, V., Sawyer, J.K., Danuser,

G., and Montell, D.J. (2014). Mechanical feedback through E-cadherin promotes direction sensing

during collective cell migration. Cell *157*, 1146-1159. 10.1016/j.cell.2014.03.045.

Campbell, K., and Casanova, J. (2016). A common framework for EMT and collective cell migration.

Development *143*, 4291-4300. 10.1242/dev.139071.

Clay, M.R., and Halloran, M.C. (2013). Rho activation is apically restricted by Arhgap1 in neural

crest cells and drives epithelial-to-mesenchymal transition. Development *140*, 3198-3209.

10.1242/dev.095448.

Cliffe, A., Doupé, D.P., Sung, H., Lim, I.K.H., Ong, K.H., Cheng, L., and Yu, W. (2017). Quantitative

3D analysis of complex single border cell behaviors in coordinated collective cell migration. Nature

Communications *8*.

Dai, W., Guo, X., Cao, Y., Mondo, J.A., Campanale, J.P., Montell, B.J., Burrous, H., Streichan,

S., Gov, N., Rappel, W.J., and Montell, D.J. (2020). Tissue topography steers migrating. Science

370, 987-990. 10.1126/science.aaz4741.

De Graeve, F.M., Van de Bor, V., Ghiglione, C., Cerezo, D., Jouandin, P., Ueda, R., Shashidhara,

L.S., and Noselli, S. (2012). Drosophila apc regulates delamination of invasive epithelial clusters.

Dev Biol *368*, 76-85. 10.1016/j.ydbio.2012.05.017.

Duchek, P., and Rørth, P. (2001). Guidance of cell migration by EGF receptor signaling during

Drosophila oogenesis. Science *291*, 131-133. 10.1126/science.291.5501.131.

Duchek, P., Somogyi, K., Jékely, G., Beccari, S., and Rørth, P. (2001). Guidance of cell migration

by the Drosophila PDGF/VEGF receptor. Cell *107*, 17-26. 10.1016/s0092-8674(01)00502-5.

Geisbrecht, E.R., and Montell, D.J. (2002). Myosin VI is required for E-cadherin-mediated border

cell migration. Nat Cell Biol *4*, 616-620. 10.1038/ncb830.

Ghiglione, C., Jouandin, P., Cérézo, D., and Noselli, S. (2018). The *Drosophila* insulin pathway

controls *Profilin* expression and dynamic actin-rich protrusions during collective cell migration.

Development *145*. 10.1242/dev.161117.

Gouignard, N., Andrieu, C., and Theveneau, E. (2018). Neural crest delamination and migration:
Looking forward to the next 150 years. Genesis *56*, e23107. 10.1002/dvg.23107.
Inaki, M., Vishnu, S., Cliffe, A., and Rørth, P. (2012). Effective guidance of collective migration
based on differences in cell states. Proc Natl Acad Sci U S A *109*, 2027-2032.
10.1073/pnas.1115260109.
Ko, C.S., and Martin, A.C. (2020). The cellular and molecular mechanisms that establish the
mechanics of Drosophila gastrulation. Curr Top Dev Biol *136*, 141-165. 10.1016/bs.ctdb.2019.08.003.
Lamb, M.C., Anliker, K.K., and Tootle, T.L. (2020). Fascin regulates protrusions and delamination
to mediate invasive, collective cell migration in vivo. Dev Dyn *249*, 961-982. 10.1002/dvdy.186.
Lamb, M.C., Kaluarachchi, C.P., Lansakara, T.I., Mellentine, S.Q., Lan, Y., Tivanski, A.V., and
Tootle, T.L. (2021). Fascin limits Myosin activity within *Drosophila* border cells to control
substrate stiffness and promote migration. Elife *10*. 10.7554/eLife.69836.
Liu, J.A., Rao, Y., Cheung, M.P.L., Hui, M.N., Wu, M.H., Chan, L.K., Ng, I.O., Niu, B., Cheah,
K.S.E., Sharma, R., et al. (2017). Asymmetric localization of DLC1 defines avian trunk neural crest
polarity for directional delamination and migration. Nat Commun *8*, 1185.
10.1038/s41467-017-01107-0.
Mathieu, J., Sung, H.H., Pugieux, C., Soetaert, J., and Rorth, P. (2007). A sensitized
PiggyBac-based screen for regulators of border cell migration in Drosophila. Genetics *176*,
1579-1590. 10.1534/genetics.107.071282.
Monahan, A.J., and Starz-Gaiano, M. (2013). Socs36E attenuates STAT signaling to optimize motile
cell specification in the Drosophila ovary. Dev Biol *379*, 152-166. 10.1016/j.ydbio.2013.03.022.
Montell, D.J. (2006). The social lives of migrating cells in Drosophila. Curr Opin Genet Dev *16*,
374-383. 10.1016/j.gde.2006.06.010.
Montell, D.J., Rorth, P., and Spradling, A.C. (1992). slow border cells, a locus required for a
developmentally regulated cell migration during oogenesis, encodes Drosophila C/EBP. Cell *71*, 51-62.
10.1016/0092-8674(92)90265-e.
Nieto, M.A., Huang, R.Y., Jackson, R.A., and Thiery, J.P. (2016). EMT: 2016. Cell *166*, 21-45.
10.1016/j.cell.2016.06.028.
Pinheiro, E.M., and Montell, D.J. (2004). Requirement for Par-6 and Bazooka in Drosophila border
cell migration. Development *131*, 5243-5251. 10.1242/dev.01412.
Poukkula, M., Cliffe, A., Changede, R., and Rørth, P. (2011). Cell behaviors regulated by guidance
cues in collective migration of border cells. J Cell Biol *192*, 513-524. 10.1083/jcb.201010003.
Ramkumar, N., Omelchenko, T., Silva-Gagliardi, N.F., McGlade, C.J., Wijnholds, J., and Anderson,
K.V. (2016). Crumbs2 promotes cell ingression during the epithelial-to-mesenchymal transition at
gastrulation. Nat Cell Biol *18*, 1281-1291. 10.1038/ncb3442.
Rørth, P. (1994). Specification of C/EBP function during Drosophila development by the bZIP basic

region. Science *266*, 1878-1881. 10.1126/science.7997882.

Rørth, P., Szabo, K., Bailey, A., Laverty, T., Rehm, J., Rubin, G.M., Weigmann, K., Milán, M.,

Benes, V., Ansorge, W., and Cohen, S.M. (1998). Systematic gain-of-function genetics in Drosophila.

Development *125*, 1049-1057.

Rørth, P., Szabo, K., and Texido, G. (2000). The level of C/EBP protein is critical for cell migration

during Drosophila oogenesis and is tightly controlled by regulated degradation. Mol Cell *6*, 23-30.

10.1016/s1097-2765(05)00008-0.

Saykali, B., Mathiah, N., Nahaboo, W., Racu, M.L., Hammou, L., Defrance, M., and Migeotte, I. (2019).

Distinct mesoderm migration phenotypes in extra-embryonic and embryonic regions of the early mouse

embryo. Elife *8*. 10.7554/eLife.42434.

Silver, D.L., and Montell, D.J. (2001). Paracrine signaling through the JAK/STAT pathway activates

invasive behavior of ovarian epithelial cells in Drosophila. Cell *107*, 831-841.

10.1016/s0092-8674(01)00607-9.

Starz-Gaiano, M., Melani, M., Wang, X., Meinhardt, H., and Montell, D.J. (2008). Feedback inhibition

of Jak/STAT signaling by apontic is required to limit an invasive cell population. Dev Cell *14*,

726-738. 10.1016/j.devcel.2008.03.005.

Stuelten, C.H., Parent, C.A., and Montell, D.J. (2018). Cell motility in cancer invasion and

metastasis: insights from simple model organisms. Nat Rev Cancer *18*, 296-312. 10.1038/nrc.2018.15.

Szabó, A., and Mayor, R. (2018). Mechanisms of Neural Crest Migration. Annu Rev Genet *52*, 43-63.

10.1146/annurev-genet-120417-031559.

Szafranski, P., and Goode, S. (2004). A Fasciclin 2 morphogenetic switch organizes epithelial cell

cluster polarity and motility. Development *131*, 2023-2036. 10.1242/dev.01097.

Tepass, U., Gruszynski-DeFeo, E., Haag, T.A., Omatyar, L., Török, T., and Hartenstein, V. (1996).

shotgun encodes Drosophila E-cadherin and is preferentially required during cell rearrangement

in the neurectoderm and other morphogenetically active epithelia. Genes Dev *10*, 672-685.

10.1101/gad.10.6.672.

Thiery, J.P., Acloque, H., Huang, R.Y., and Nieto, M.A. (2009). Epithelial-mesenchymal transitions

in development and disease. Cell *139*, 871-890. 10.1016/j.cell.2009.11.007.

Wang, H., Qiu, Z., Xu, Z., Chen, S.J., Luo, J., Wang, X., and Chen, J. (2018). aPKC is a key polarity

determinant in coordinating the function of three distinct cell polarities during collective

migration. Development *145*. 10.1242/dev.158444.

Wang, X., Bo, J., Bridges, T., Dugan, K.D., Pan, T.C., Chodosh, L.A., and Montell, D.J. (2006).

Analysis of cell migration using whole-genome expression profiling of migratory cells in the

Drosophila ovary. Dev Cell *10*, 483-495. 10.1016/j.devcel.2006.02.003.

Wang, X., He, L., Wu, Y.I., Hahn, K.M., and Montell, D.J. (2010). Light-mediated activation reveals

a key role for Rac in collective guidance of cell movement in vivo. Nat Cell Biol *12*, 591-597.

10.1038/ncb2061.

Yang, J., Antin, P., Berx, G., Blanpain, C., Brabletz, T., Bronner, M., Campbell, K., Cano, A.,

Casanova, J., Christofori, G., et al. (2020). Guidelines and definitions for research on

epithelial-mesenchymal transition. Nat Rev Mol Cell Biol *21*, 341-352. 10.1038/s41580-020-0237-9.

Yang, N., Inaki, M., Cliffe, A., and Rørth, P. (2012). Microtubules and Lis-1/NudE/dynein regulate

invasive cell-on-cell migration in Drosophila. PLoS One *7*, e40632. 10.1371/journal.pone.0040632.

Figures

Fig. 1.

Delamination is a complex cellular process. (A) A time series showing the delamination process

in wild-type *Drosophila* border cells. (B) A time series of transmission images of the egg

chamber during delamination. Blue arrows indicate oocyte length, which increased with time.

(C) Oocyte length at the delamination point was determined from 17 movies of wild-type border

cells. We defined an oocyte length of 50-70 µm as a marker of the wild-type delamination period.

Elapsed time from the start of the movie is shown at lower left. For all images, anterior is left.

Fig. 2.

Locomotion is impaired upon JAK/STAT inhibition in wild-type but not in *slbo*-mutant border

cells. (A-E) Single Z slices from movies show the delamination of wild-type (A), DN-Dome

expressing (B), *slbo*-mutant (C), and CytochalasinD-treated (D) border cells and a movie of

posterior follicle cells (E). Red and blue circles indicate the cluster center and a border cell

nucleus, respectively. (F) A schematic showing how cell movement was quantified by measuring

the change in the angle of the border-cell nucleus (blue circle) relative to the cluster center (red

circle). (G) The angular velocity of wild-type (WT), DN-Dome-expressing (*Dome*), *slbo*-mutant

(*slbo*), and CytochalasinD-treated (CytoD) border cells relative to the cluster center, and the

angular velocity of posterior follicle cells relative to the posterior pole cell center (FC). $12 \leqq$

number of cells (n) $\leqq 24$, $6 \leqq$ number of egg chambers (N) $\leqq 12$. Error bars indicate SEM. ***,

$p < 0.001$. Time shown at the lower left indicates elapsed time from the start of the movie (white)

and time prior to the delamination point (yellow). For all images, anterior is left.

Fig. 3

Invasiveness is regulated by *slbo*, partly though *shg*. (A-D) Z stacks from delamination movies

of border cells, as follows: wild-type (A), *slbo*-mutant (B), *shg*-mutant (C), and *slbo*-mutant with

E-cad expression (D). Light blue arrows indicate the distance (reach) from the anterior end of the

egg chamber to the tip of the border-cell extension at delamination (A3). (E,F) The cluster body

(blue) and extensions (red) separated by image analysis from the images shown in (A2,C2).

Light blue circles: the centroid of the cluster body. (G,H) The maximum length (G) and duration

(H) of the front extensions of border-cell clusters, as follows: wild-type (WT), *slbo*-mutant (*slbo*),

shg-mutant (*shg*), and *slbo*-mutant with E-cad expression (*slbo*+E-cad). $6 \leqq N \leqq 14$. Error bars

indicate SEM. ***, $p<0.001$, **, $p<0.01$. Time shown at the lower left indicates time elapsed

from the start of the movie (white) and the time prior to the delamination point (yellow). For all

images, anterior is left.

Fig. 4

Invasive protrusion is required to reach a distance. (A,B) Z stacks from movies of

guidance-deficient mutants in the wild-type (A), and the actual (B) delamination period. Light

blue arrow indicates from the anterior end of the egg chamber to the reach of the border-cell

cluster at delamination (B3). (C-E) The angular velocity (C), maximum length (D), and

persistence (E) of front extensions of border-cell clusters in wild-type (WT), and

guidance-deficient mutants in the wild-type (GM_e) and actual (GM_d) delamination periods.

(C) $16 \leqq n \leqq 24$, $8 \leqq N \leqq 12$. (D,E) $8 \leqq N \leqq 14$ (F) The distance from the anterior end of the egg

chamber to the tip of the border-cell extension at the delamination point for wild-type (WT) and

the guidance-deficient mutant (GM_d). $8 \leqq N \leqq 13$. Error bars indicate SEM. ***, $p<0.001$, **,

$p<0.01$. Time shown at the lower left indicates time elapsed from the start of the movie (white)

and time prior to delamination (yellow). For all images, anterior is left.

Fig. 5

Motility is not a prerequisite for invasiveness. (A,B) Z stacks from movies of border cells

expressing DN-Dome with E-cad (A) and Slbo (B). E-cad and Slbo induced front extensions in

JAK/STAT-inhibited border cells. (C-E) Angular velocity and (C) the maximum length (D) and

duration of front extensions in wild-type (WT) or DN-Dome expressing (*Dome*) border-cell

clusters with either E-cad (*Dome*+E-cad) or Slbo (*Dome*+Slbo) expression. (C) 16≦n≦24, 8≦

N≦12. (D,E) 6≦N≦14. Error bars indicate SEM. ***, p<0.001, **, p<0.01. Time elapsed from

the start of the movie is indicated at the bottom right. For all images, anterior is left.

Supplemental figures

Fig. S1

The angular velocity of wild-type (WT) and $slbo^{1310}/slbo^{e7b}$ border cells. Angular velocities were

comparable in these conditions. n=10, N=5. Error bars indicate SEM.

Fig. S2

Angular velocity for the following border cells: wild-type (WT), *shg*-mutant (*shg*), and

slbo-mutant with E-cad expression (*slbo*+E-cad). Angular velocities were comparable in these

conditions. $10 \leqq n \leqq 26$, $5 \leqq N \leqq 13$. Error bars indicate SEM.

Fig. S3

(A,B) Examples of changes in front extension length (A) and in cluster reach (the distance

between the anterior end of the egg-chamber and the extension tip) (B) during delamination. X

axis: time frame. Red lines: delamination point. Front extension length varied quite a bit, while

cluster reach tended to increase over time. The maximum extension length did not correlate with

the delamination point. (C,D) The maximum extension length (C) and the cluster reach (D) just

before delamination. Length varied more than reach.

Movies

Movie 1

Delaminating wild-type *Drosophila* border cells form a cluster of jiggling cells. The cluster

extends a protrusion in the direction of future migration, and the tip finds an attachment point.

The cell cluster detaches from the epithelial layer and follows the protrusion as it contracts.

Elapsed time (minutes) shows at lower right.

Movie 2

JAK/STAT-inhibited follicle cells failed to differentiate into border cells, move into clusters, or

extend protrusions in the period corresponding to wild-type delamination. JAK/STAT signaling

was inhibited with DOME lacking a cytoplasmic domain. Elapsed time shows at the lower

right.

Movie 3

slbo-mutant border cells formed clusters and moved within the clusters, but did not form

protrusions or delaminate during the period when wild-type cells would delaminate. Elapsed

time shows at the lower right.

Movie 4

slbo-mutant border cells formed clusters but lost motility when treated with CytochalasinD,

which inhibits actin polymerization, and failed to delaminate during the period of wild-type

delamination period. Elapsed time shows at the lower right.

Movie 5

shg-mutant border cells formed clusters of motile cells and sent out short extensions, but did not

delaminate during the wild-type delamination period. Elapsed time shows at the lower right.

Movie 6

slbo-mutant border cells with E-cad expression formed front extensions but did not delaminate

during the wild-type delamination period. Elapsed time shows at the lower right.

Movie 7

Guidance-deficient border cells had normal motion but formed front extensions that were shorter

and less persistent than wild-type extensions.

Movie 8

Guidance-deficient border cells delaminated successfully. Elapsed time shows at the lower right.

Movie 9

E-cad expression did rescued invasive extensions in JAK/STAT-inhibited border cells did not

rescue cell movement. Elapsed time shows at the lower right.

Movie 10

Slbo overexpression induced front extensions in JAK/STAT-inhibited follicle-like cells, but the

cells failed to delaminate. Elapsed time shows at the lower right.