1 **Elucidating the diversity of microeukaryotes and epi-endophytes in the brown**
2 **algal holobiome**
3
4

5 Marit F. Markussen Bjorbækmo[1,2*], Juliet Brodie[2], Ramiro Logares[3], Stephanie
6 Attwood[2], Stein Fredriksen[4], Janina Fuss[1], Kamran Shalchian-Tabrizi[1], Anders Wold-
7 Dobbe[1], Anders K. Krabberød[1], David Bass[2,5*]
8
9

[1.] University of Oslo, Department of Biosciences, Section for Genetics and
Evolutionary Biology (EVOGENE), N-0316, Oslo, Norway

[2.] The Natural History Museum (NHM), Department of Life Sciences, London SW7
5BD, United Kingdom

[3.] Institut de Ciències del Mar (CSIC), Department of Marine Biology and
Oceanography, ES-08003, Barcelona, Catalonia, Spain

[4.] University of Oslo, Department of Biosciences, Section for Aquatic biology and
Toxicology (AQUA), N-0316, Oslo, Norway
18
19 [5.] Centre for Environment Fisheries and Aquaculture Science (CEFAS), Weymouth,
20 Dorset DT4 8UB, United Kingdom
21
22
23 [*] Correponding authors:
24 David Bass
25 e-mail: david.bass@cefas.co.uk
26 Phone:
27
Marit F. Markussen Bjorbækmo
e-mail: m.m.bjorbakmo@gmail.com
Phone:

Abstract

Background

Brown algae (Phaeophyceae) are essential species in coastal ecosystems where they

form kelp forests and seaweed beds that support a wide diversity of marine life. Host-

associated microbial communities are an integral part of phaeophyte biology. The

bacterial microbial partners of phaeophytes have received far more attention than

microbial eukaryotes. The pre-requisite to understand the ecology of phaeophytes and

their host-associated microbes is to know their diversity, distribution and community

dynamics. To our knowledge, this is the first study to investigate phaeophyte-associated

eukaryotes (the eukaryome) using broadly targeting 'pan-eukaryotic' primers and high

throughput sequencing (HTS). Using this approach, we aimed to unveil the eukaryome

of seven large common phaeophyte species. We also aimed to assess whether these

macroalgae harbour novel eukaryotic diversity and to ascribe putative functional roles

to the host-associated eukaryome, based on taxonomic affiliation and phylogenetic

placement.

Results

Our sequence dataset was dominated by phaeophyte reads, from the host species and

potential symbionts. We also detected a broad taxonomic diversity of eukaryotes in the

phaeophyte holobiomes, with OTUs taxonomically assigned to ten of the eukaryotic

major Kingdoms or supergroups. A total of 265 microeukaryotic and epi-endophytic

operational taxonomic units (OTUs) were defined, using 97% similarity cut off during

clustering, and were dominated by OTUs assigned to Stramenopiles, Alveolata and

Fungi. C Almost one third of the OTUs we detected have not been found in previous

molecular environmental surveys, and represented potential novel eukaryotic diversity.

This potential novel diversity was particularly diverse in phylogenetic groups

comprising heterotrophic and parasitic organisms, such as labyrinthulids and

Oomycetes, Cercozoa, and Amoebozoa.

Conclusions

Our findings provide important baseline data for future studies of seaweed-associated

microorganisms, and demonstrate that microeukaryotes and epi-endophytic eukaryotes

should be considered as an integral part of phaeophyte holobionts. The potential novel

eukaryotic diversity we found and the fact that the vast majority of macroalgae in

marine habitats remain unexplored, demonstrates that brown algae and other seaweeds

are potentially rich sources for a large and hidden diversity of novel microeukaryotes

and epi-endophytes.

Keywords [3-10]

Symbiome, epiphyte, endophyte, protist, microeukaryote, eukaryome, Phaeophyceae,

seaweed, holobiont, fungi

Background

Seaweeds are essential to the health of our planet, providing core ecosystem services,

and food and shelter to a wide diversity of marine life, from microbes to mammals, [1,

2 and references therein]. Brown algal seaweeds (Phaeophyceae) are Stramenopiles, a

radiation of eukaryotes which is phylogenetically distant from red- and green algal

seaweeds (Archaeplastida; Chlorophyta and Rhodophyta). The phaeophytes are one of

the most diversified groups of benthic algae and comprise approximately 2000 species,

encompassing complex multicellular species which have diversified and evolved since

the Mesozoic Era ~250 MYA [2, 3]. Large phaeophytes, especially those in the orders

Laminariales, Fucales and Tilopteridales are dominant members of intertidal and

shallow subtidal ecosystems worldwide, where they function as ecosystem engineers

forming complex underwater forests. These coastal ecosystems are some of the most

productive habitats on the planet [4], where seaweed beds and kelp forests are

indispensable biodiversity hotspots, which also provide habitat and nursery for a wide

range of taxa, including many commercially important species [5]. Phaeophytes have

been shown to accommodate up to 100.000 individuals of different invertebrate species

per square metre [6].

Phaeophytes and other seaweeds also harbour a wide diversity of microbial

epibionts and endobionts, comprising eukaryotes [7-9], prokaryotes [10-12] and viruses

[13-16]. These, together with the host, are collectively referred to as the seaweed

holobiont and can be considered as a localized ecosystem living on and in a host [10,

17-19]. It is increasingly recognized that host-associated microbial communities

(microbiomes) are an integral part of the host biology, exerting diverse and strong

influences on their hosts [19, 20]. While the term microbiome usually refers mostly or

exclusively to bacteria, it is important to consider the whole microbial symbiome,

including microbial eukaryotes (microeukaryotes) and larger epiphytic, epizootic, and

endophytic symbionts, to enable a comprehensive understanding of holobiont

functioning [19, 21].

Symbiotic interactions in its broadest sense refers to all intimate ecological

interactions between two species [22]. According to the fitness effect on the members

involved in the symbiotic relationship, the associations cover a wide spectrum of

interactions along the gradual continuum between positive mutualism and negative

parasitism. Some symbiotic interactions are obligate while other are more temporary

and fluctuating, and the nature of the interaction can vary depending on the symbiont's

interactions with its hosts, other symbionts and environmental conditions [20]. To date,

bacterial symbionts of phaeophytes have received most study [e.g., 10, 11, 12, 23], and

have been shown to represent complex and highly dynamic relationships that can have

everything from fundamental to detrimental effects on their hosts [10, 24-28].

Microeukaryotes in the phaeophyte symbiome are far less well known. Only a

small number of studies have investigated these associations, mostly using traditional

culturing/cell isolation methods or targeted molecular approaches. These studies

demonstrate that a broad taxonomic diversity of microeukaryotes are associated with

phaeophytes, including surface dwelling heterotrophic diatoms, dinoflagellates and

ciliates [8], 'naked amoeba' [29], epiphytic and endophytic diatoms [30-32], and green

algal endophytes [33], in addition to parasitic or saprotrophic labyrinthulids [34, 35],

oomycetes [7, 36, 37], phytomyxids [38-40] and fungi [9, 41-44]. The nature of these

microeukaryote-host relationships is mostly unknown, although some symbionts can

have detrimental effects on their macroalgal hosts, for example phytomyxids [38-40],

oomycetes [7, 36, 37] and chytridiomycete fungi [41]. Other microeukaryotes are

suspected to have a beneficial effect on their hosts, for example fungi that are mutualists

[9, 45, 46] and fungal endophytes that might protect seaweeds against pathogenic

protists [44].

Adding to the complexity of the microeukaryotic biodiversity associated with

phaeophytes, their thalli are often overgrown with a wide variety of smaller seaweeds

which represents a continuum between epiphytism and endophytism. These epi-

endophytes can have negative effects such as imposing physical and physiological

stress on their host, or positive ecosystem effects such as increased available habitats

and food for both macroscopic and microscopic life [47]. As macroalgal symbiomes

comprise multi-cellular algae as epiphytes [48, 49] and potentially pathogenic

endophytes [50-53] in addition to microeukaryotes, the almost complete lack of

knowledge of these eukaryotes in a holobiont context/perspective is a fundamental

knowledge gap which needs to be filled in order to improve our understanding of

phaeophyte holobionts.

The overall aim of this study was to unveil the eukaryome of seven large

phaeophytes which are key components in coastal ecosystems: *Fucus vesiculosus*, *F.*

serratus, *Himanthalia elongata* and *Ascophyllum nodosum* (order Fucales), *Laminaria*

digitata and *Saccharina latissima* (order Laminariales) and *Saccorhiza polyschides*

(order Tilopteridales). We achieved this aim by using using broadly-targeted pan-

eukaryotic primers and High Throughput Sequencing (HTS). This application in host

environments has so far been limited due to the technical limitation of co-amplifying

host DNA when using 'universal' eukaryotic primers [20]. However, with sufficient

sequencing depth now tractable to amplify host-associated eukaryotes much more

comprehensively than was previously possible, this will become increasingly

established as a standard tool for investigating host-associated eukaryomes [54, 55].

We also aimed to assess whether phaeophytes harbour a novel or differentially

structured diversity of (micro)eukaryotes, and to ascribe putative functional roles to the

microeukaryotes associated with these phaeophytes (e.g., putative parasites).

Results

Composition of the brown algal eukaryome

A total of 256 eukaryotic Operational Taxonomic Units (OTUs) were defined by
UPARSE and post-UPARSE trimming. Their read abundance in each library are shown
in Supplementary Table S1. Each library represented five phaeophyte samples pooled
according to species and geographic location, and is hereafter referred to as samples.
The richness recovered from each sample ranged from 26 to 101 OTUs, and the
taxonomic profile of these are shown as proportional OTU abundance in Fig. 1,
Supplementary Table S1. The sequence dataset was dominated by phaeophyte reads
(97.3% of all sequences, Supplementary Table S2). Phaeophytes also displayed the
highest OTU richness (18.9% of the total OTUs; Fig. 2, and accountable for 27.3-
70.4% of the OTUs per sample; Supplementary Table S2). Nevertheless, there was a
broad taxonomic diversity of microeukaryotes in the phaeophyte samples with OTUs
taxonomically assigned to ten of the eukaryotic major Kingdoms or supergroups (Fig.
1, Fig. 2).

Phylogenetic diversity of eukaryotic OTUs

Phylogenetic analyses were conducted for taxonomic groups displaying either high
taxonomic diversity and/or taxonomic groups previously documented to be in a
symbiotic relationship (*sensu lato*) with phaeophytes. The resulting phylogenies and
heat maps of proportional read abundance (log_{10}) of each OTU in the different samples
are shown in Figs. 3-7 and Supplementary Figs 1-4, which include OTUs and the closest
sequence matches in NCBI GenBank as of 05.05.2020 (re-blasted 11.04.2021).

For some taxonomic groups, the heatmaps indicated that the majority of OTUs

were only detected associated with one phaeophyte sample, such as Cercozoa (Fig. 4),

Fungi (Fig. 5), Ciliophora (Supplementary Fig. 1), Apicomplexa and Perkinsea

(Supplementary Fig. 2), Dinoflagellata (Supplementary Fig. 4) and Amoebozoa and

Centroheliozoa (Supplementary Fig. 3). However, most of these taxonomic groups also

comprised OTUs that were found in multiple samples, across host orders and

geographic locations (e.g., the fungal ascomycete OTU 226; Fig. 5, the ciliate OTU 89;

Supplementary Fig. 1, and the centroheliozoan OTU 13; Supplementary Fig. 3 were

found in samples belonging to Fucales, Tilopteridales and Laminariales from both

Norway and the UK). For other taxonomic groups such as Bacillariophyta (Fig. 6) and

Labyrinthulomycetes & Oomycetes (Fig. 3), the heatmaps show that a large proportion

of the OTUs were detected in multiple samples.

Several OTUs clustered with reference sequences that have been found

associated with various hosts in previous studies, such as the labyrinthulid and

oomycete OTUs clustering with *Aplanochytrium* (Labyrinthlida) and Olpidiopsidales

(Fig. 3), and the cercozoan vampyrellid OTUs clustering with a reference sequence

previously detected in samples of brown alga (Fig. 4).

Novel diversity in the brown algal eukaryome

We found that ~30% of the OTUs had low percentage identity (<95 %) with any known

close relatives in reference sequence databases and represent potential novel taxonomic

diversity (Supplementary Table 1). The majority of OTUs taxonomically and

phylogenetically placed within Fungi, Bacillariophyta and ciliates displayed high

percentage identity (>95 %) to known reference sequences (Fig. 5; Fig. 6; and

Supplementary Fig. 1, respectively). Conversely, within Labyrinthulomycetes and

Oomycetes (Fig. 3), Cercozoa (Fig. 4) and Amoebozoa and Centroheliozoa

(Supplementary Fig. 3), about 50% of the OTUs had low similarity to known reference

sequences (<95 %) which indicated that these OTUs represented potential novel

diversity. Some of this potential novel diversity was represented by specific clades in

the phylogenetic trees, such as a well-supported labyrinthulid clade within

Thraustochytrida that encompassed four OTUs with very low percentage identity

(ranging between 84.1% to 89.6%). Similarly, within Cercozoa, eight of the nine

Vampyrellida, Phytomyxea, and Novel Clade 9 OTUs had low similarity to reference

sequences (from 85.2% to 93%; Fig 4).

Green (Chlorophyta), red (Rhodophyta) and Brown (Phaeophyceae) algae:

In addition to protist taxa, we also detected OTUs from green, red, and (non-host-

derived) brown algae (Fig. 7). The majority of the nine green algal OTUs (Fig. 7) were

very similar (>99% identity) to characterized and sequenced Ulvales taxa such as

Ulvella (formerly *Acrochaete*) leptochaete and *Umbraulva japonica*. Only a single

green alga (OTU 264) displayed low similarity (90.4%) to reference sequences, and

clustered with *Lithotrichon pulchrum*, Ulvales (Fig. 7). Similarly, the five red algal

OTUs (Fig. 7) displayed >98% similarity to taxa including *Ceramium* sp. and

Cryptopleura ramosa.

There were in total 50 phaeophyte OTUs. Significant levels of microdiversity

in the V4 OTUs were seen in the phaeophyte clade. OTUs corresponding to the host

species were inferred on the basis of their phylogenetic proximity to, and high

proportions of reads associating with, reference sequences from that host, either from

GenBank or sequences generated in this study. Host OTUs are indicated by labeled

brackets on Fig. 7. Many of the OTUs derived from the host species were present in

samples from multiple host species, albeit generally represented with lower read

abundance in the 'non-host' samples.

However, other OTUs did not cluster with host-derived sequences, or from the

longer 18S sequences generated by cloning amplicons from the host. The 26 OTUs in

the "uncharacterized clade" on Fig. 7 were clearly distinct from any characterized or

environmental sequences in reference sequence databases. This clade encompassed

OTUs that were mainly detected in the *Fucus* sp. samples, but six of them were also

detected in lower abundance in other phaeophyte samples (Fig. 7). Additionally, seven

OTUs in the *A. nodosum* clade may be host sequences but are obviously different in

sequence from the cloned sequences generated from that host, and OTUs 2 and 326,

presumably deriving from that host (Fig. 7).

Some OTUs grouped strongly with known epiphytic lineages, such as the three

OTUs clustering with genera *Pylaiella* and *Halothrix*. These OTUs were found in all,

or in the majority, of the different pheaophyte hosts (Fig. 7). In the same clades were

the brown algal endophytes isolated and sequenced by Attwood (2005), published here

for the first time, and represented on Fig. 7, all in the 'brown endophytes' clade. These

endophytes were identified as *Myrionema strangulans*, isolated from *Fucus serratus*,

Mastocarpus stellatus, *Chorda filum*, *Osmundea pinnatifida*, and *Chondrus crispus*;

Microspongium tenuisimum, isolated from *Osmundea osmunda* and *C. crispus*;

Streblonema sp. isolated from *M. stellatus*; *Mytriotrichia clavaeformis* and *Ectocarpus*

sp. from *C. crispus*; and *Chorda filum* from a larger *C. filum* individual. Light

microscopy images of the isolated endophytes are shown in Fig. 8.

Three of our OTU sequences occurred in the same clade as these endophytes:

OTU 9 and some 18S clone sequences from *S. polyschides* clustering with *Pylaiella*,

and OTUs 70 and 146, clustering most closely with *Myriotrichia* and *Myrionema* (Fig.

7). All of these were strongly associated with *F. vesiculosus* and *S. polyschides*

individuals, but they were present in almost all host samples.

Analyses of community composition

The clustering of samples was consistent irrespective of whether host OTUs were

included in the analyses (Fig 9: A-B). The three *F. vesiculosus* samples clustered with

F. serratus and *S. polyschides* in both plots (Fig 9: A-B), which indicated that these

have a similar OTU composition. The *A. nodosum* sample showed less similarity with

Fucus and *S. polyschides* when the phaeophyte OTUs were excluded from the analysis

(Fig. 9, A-B). *Himanthalia elongata* did not cluster with any other phaeophyte hosts

and appeared highly dissimilar from the others both including and excluding

phaeophyte OTUs (Fig. 9; A-B). Analyses of similarity (ANOSIM) was highly

significant both with and without phaeophyte OTUs included (p-value 7e-04 and 0.03

respectively), which supported the dissimilarity observed between the phaeophyte host

genera in the NMDS plots (Fig. 9: A-B).

Dissimilarity between host genera (SIMPER values) ranged from 31.4%

(*Laminaria* spp. vs *Saccharina* spp.) to 99.9% (*Ascophyllum* spp. vs *Himanthalia* spp.)

in the dataset including phaeophyte OTUs (Table 1, Supplementary Table S3), where

certain host derived OTUs (OTU 1, 2, 3 and 5) were the main contributors to the

observed dissimilarity. These results also supported some of the OTUs inferred to

derive from the host species based on their phylogenetic placement and read abundance

(Table 1, Fig. 7). When phaeophyte OTUs were excluded the host genera were all

highly dissimilar, with SIMPER values ranging from 83.3% (*Ascophyllum* spp. vs

Saccorhiza spp.) to 99.4% (*Ascophyllum* spp. vs *Himanthalia* spp.). The main

contributors to the observed dissimilarity between the phaeophyte host genera were

green algal, red algal, fungal and oomycete OTUs in addition to an unassigned OTU

(Table 2). There were also several OTUs assigned to a wide range of taxonomic groups

that had some contribution to the dissimilarity (Supplementary Table S3).

IndVal analyses of the dataset including phaeophyte OTUs, indicated that six

OTUs were preferentially associated with *Ascophyllum* (Table 3), of which five were

taxonomically assigned to Phaeophyceae. Four of these OTUs clustered within the

clade labelled "*A. nodosum*" in the phylogenetic tree (Fig. 7, Table 3), while the fifth

OTU was placed in the "Uncharacterised" clade, represented with reads in all

phaeophyte libraries (Fig. 7). The final OTU (25) was a fungal OTU found in all the

libraries except *L. digitata* (UK), but with higher read abundance in *Ascophyllum* than

in the other host genera (Fig. 5). OTU 25 was also the only one found to be significant

by the IndVal analysis of the dataset excluding phaeophytes (p-value 0.0446; Table 2).

Shared OTUs

Although our analyses suggested a large degree of dissimilarity between the different

phaeophyte hosts (Table 1, 2), 70 OTUs were found in multiple samples across the host

orders Fucales, Laminariales and Tilopteridales (Fig. 10, Table 4). Of these, 25 OTUs

were detected in samples belonging to all of the three host orders, of which 15 belonged

to Phaeophyceae, including the three brown algal endophyte OTUs (Fig. 7), three OTUs

from the "uncharacterized" clade (Fig. 7) and several host OTUs that were found in

high abundances in the respective host samples, but also present in lower abundance in

several of the other phaeophyte samples. The shared OTUs also included five diatoms

which clustered with known epiphytes and endophytes of macroalgae in the

phylogenetic tree (Fig. 6, Table 4).

Twenty OTUs were found in samples belonging to Fucales and Laminariales

(Table 4), including Fungi, Oomycetes and Labyrithulomycetes (Table 4, Fig. 5, 3).

The 22 OTUs that overlapped between Fucales and Tilopteridales comprised several

diatom and green algal OTUs (Table 4, Fig 6, 7), and two Oomycete OTUs that

clustered within Olpidiopsidales (Table 4, Fig. 3). The OTUs that were unique to

Fucales, Tilopteridales and Laminariales are shown in Supplementary Table S4.

Discussion

Broad diversity of microeukaryotes associated with brown algae

To our knowledge, this is the first study that has investigated the diversity of eukaryotes

associated with brown algae using high throughput sequencing. Our findings show that,

although the sequence data was dominated by brown algal reads, we were able to

identify a broad diversity of microeukaryotes representing most of the main branches

in the eukaryotic tree of life (Fig 1, 2).

Some of the brown algae we investigated in this study displayed a higher

diversity of microeukaryotes compared to others. Particularly, we detected a higher

diversity associated with *Fucus vesiculosus*, *F. serratus* (Fucales) and *Saccorhiza*

polyschides (Tilopteridales) than *Laminaria digitata* and *Saccharina latissima*

(Laminariales). There are several factors/variables which may have affected the

observed diversity and community composition. Density and composition of

microorganisms associated with brown algae have been shown to vary at several scales:

e.g., among host species, between individuals of the same host species, different thallus

areas of individual hosts, the kind of surface tissue available for microepiphyte

colonization, different habitats and location in the intertidal zone, seasons, different

stages of the life cycle, and different longevity of the host [e.g., 19, 25, 32, 56, 57-59].

In addition, the interaction between the settlers affects the formation of specific

communities [47, 56], including chemical interactions and signalling between the

microbial symbionts [44, 60, 61]. Seaweeds themselves also, to various degrees have

physical and chemical antifouling defence mechanisms which prevents settlement of

microorganisms and epiphytes [31, 62, 63]. For this study, we may also have randomly

sampled different microhabitats of the brown algal thalli, resulting in a highly variable

diversity of microeukaryotes.

The community composition of microeukaryotes associated with *Himanthalia*

elongata stood out as very dissimilar to the other phaeophytes (Fig. 9 B). This seaweed

differs from the other phaeophytes in the type of life cycle and the tissue type sampled

for this study. *H. elongata* is biannual, but the elongated thongs which accounts for

~98% of the thallus is reproductive tissue that degrades at the end of the season

(reproductive season from late July to ~January), leaving only the vegetative basal disc

[64]. The tissue type sampled from *H. elongata* were therefore different and most likely

younger than the tissues sampled from the other phaeophytes, which could explain the

distinctive and low diversity of microeukaryotes detected in this host. A recent study

also reported low diversity of bacteria associated with *H. elongata* [65].

Eukaryotic epiphytes and endophytes in the seaweed holobiont:

Seaweed surfaces provide nutrient-rich and important habitats for epiphytic and

endophytic organisms, from unicellular microbes such as bacteria and protists to

multicellular, usually filamentous seaweeds [e.g., 10, 12, 47, 56, 66]. These

associations can be either facultative (i.e. the epi-endophytes can grow on a variety of

biotic and/or abiotic substrata) or obligate (dependent on one or more specific hosts).

Like many biological categorizations, the boundaries between epiphytes and

endophytes in real-life can be blurred; the nature of the association can be affected by

both biotic and abiotic factors, and can range from mutualism to parasitism [47, 67,

68].

Microeukaryotic epi-endophytes:

Diatoms are regularly found in the biofilm on seaweed surfaces, where they live as

epiphytes, and often occur in large numbers [32, 69-71]. Although many diatoms

associated with phaeophytes are epiphytes, there are also studies which have

demonstrated that some diatom species, such as the ones belonging to the genera

Navicula and *Cocconeis* can live as endophytes, intracellularly in tissues of both brown

algae and red algae [30, 72-74]. Mayombo *et al*. [75, 76] recently suggested that some

diatom taxa might have adapted to an endophytic life to avoid the antifouling

mechanisms of their hosts. Several of the OTUs from our study which were

taxonomically and phylogenetically assigned to different species of Naviculales,

Cocconeidales and Fragilariales (Fig. 6), were present in multiple samples/hosts

independent of geographic location. Our molecular data in combination with previous

observational studies [30, 32, 72-76] supports the hypothesis that some diatom species

such as *Navicula* sp. and *Cocconeis* sp., might have a more intimate, potentially

endophytic, association with seaweeds. This, however, remains to be tested in future

studies.

Ciliates represent another ubiquitous group of microeukaryotes living as

epiphytes in the biofilm on seaweeds [8, 77], and was also the dominating group of

alveolates in our study (Fig. 2). Many ciliates are predators/grazers feeding on bacteria

and microeukaryotes [8], whereas other ciliates can be involved in symbiotic

interactions [78]. Most of the ciliates detected in our study were only found in single

samples (Supplementary Fig. 1), which might indicate that these ciliates were "random

visitors". A few of the OTUs were, however found in several samples from both

Norway and the UK, and were highly similar to reference sequences from different

marine environmental samples, which suggest that these ciliates are generalist biofilm

formers on different organisms and substrates. However, one of these OTUs (OTU 256,

Spirotrichea, Supplementary Fig. 1) was almost identical to a reference sequence found

in samples of *Ascophyllum nodosum* used as live bait wrapping [79], which could point

towards that some of these ciliates have a closer association with brown algae. As

emphasised by del Campo et al. [20], "it is likely that many microeukaryotes routinely

detected or observed/cultured from environmental samples may also be host associated,

even if transiently".

Host-associated phaeophytes, chlorophytes and rhodophytes

In addition to the phaeophyte OTUs clearly deriving from the host species, we detected

a diversity of other sequence types from putative brown algal epi-endophytes (Fig. 7).

The diversity of 'OTUs' apparently deriving from the host organisms was much higher

than expected. This may partly be caused by artefactual sequence variants amplified to

a discernible extent by the depth of Illumina sequencing. But the distinctiveness of

some of these OTUs from obviously host-derived sequences was so great in some cases

(e.g., the strongly *Fucus*-associated 'uncharacterised' clade in Fig. 7) that we propose

that these represented hitherto unknown epi-endobionts. In other cases we have

conservatively ascribed OTU diversity clustering with host sequences (e.g., the clades

bracketed as *A. nodosum* and *H. elongata* in Fig. 7) as being host-derived, but there

remains the possibility that they represented different organisms.

Some of the phaeophyte OTUs detected in our study, such as the ones clustering

with epiphytic *Pylaiella* and *Halothrix* and the other phaeophyte epi-endophytes (Fig.

7), clearly also represented epi-/endophytic lineages [e.g., 48, 58, 80]. These OTUs

were detected in all the phaeophyte hosts included in this study, indicating a broad

distribution and host range.

By analysing sequences from cultured phaeophyte symbionts of brown algae

and red algae (Attwood 2005) we show that the epi-endophyte genera *Microspongium*,

Myriotrichia, *Myrionema*, and *Streblonema* form a clade with *Chorda* and *Ectocarpus*

(growing as epi-endophytes), and previously sequenced phaeophyte epi-endophytes

Pylaiella, *Halothrix*, and *Scytosiphon*, which all together form a maximally supported

clade with *Saccorhiza*. While this significantly expands the 18S sequence

representation of phaeophyte epi-endophytes, it does not account for the vast majority

of host-associated phaeophyte sequences in our data that are not very obviously host-

derived. Further culturing and is situ experiments are required to explain these

suggestive findings, which might represent microscopic life-stages (zoospores,

gametophytes, gametes and juvenile sporophytes) of brown algae growing epi-

/endophytically on substrata, including on macroscopic sporophytes of other seaweeds

[81-84]. This can also explain why the host-derived OTUs were detected in lower

abundance in the other brown algal samples (Fig. 7).

Another possible explanation is that small filamentous brown algae were

growing epi-endophytically in the larger brown algal hosts [48, 85]. Such endophytic

infections are described as common diseases of brown algae [e.g., 52, 53, 67, 68, 86].

Hosts containing brown algal endophytes can either be asymptomatic, show weak

symptom such as dark spots or warts, or present severe symptoms such as

morphological changes to the brown algal host tissues [52, 67]. Substrata and brown

algal thalli often contain many more brown algal endophytes than expected based on

visual inspections prior to culturing [47, 53, 85].

The green algal OTUs were all similar to reference sequences of taxa that are

known to be endophytes in red (*Chondrus crispus Mastocarpus* stellatus, *Osmundea*

spp. and *Dumontia contorta*) and brown algae (*Fucus serratus* and *Chorda filum*) [33,

87, 88] .

All red algal OTUs were similar to reference sequences of taxa that are common

epiphytes on seaweeds and a variety of other substrata. Several of the taxonomic names

for the reference sequences included in the phylogeny are out of date (Fig. 7):

Ceramium spp. is an aggregate of species (i.e. *Ceramium rubrum* in Fig. 7) which are

members of the order Ceramiales together with *Calithamnion* spp., *Polysiphonia* sensu

lato (*Polysiphonia fucoides* is now *Vertebrata fucoides*) and *Cryptoleura* spp. Red algal

endophytes are common in other rhodophytes and chlorophytes (REF!). We did not,

however find any endophytic rhodophytes in our study, which might indicate that

phaeophytes are not prone to endophytic infections by rhodophytes, but instead are

more susceptible to endophyte infestations by chlorophytes and endophytic

phaeophytes.

Novel diversity and putative parasites

We demonstrate that brown algal holobionts encompass potential novel eukaryotic

diversity, as almost one third of the OTUs we detected have not been found in molecular

environmental surveys before. This was more pronounced in some phylogenetic groups

than in others, in particular the Labyrinthulids and Oomycetes (Fig. 3), Cercozoa (Fig.

4), Amoebozoa and Centroheliozoa (Supplementary Fig. 3). Some of the OTUs were

taxonomically and phylogenetically placed with known and putative parasites of

seaweeds. Our findings and the fact that the vast majority of macroalgae in marine

habitats remain unexplored, demonstrates that brown algae and other seaweeds are

potentially rich sources for a large and hidden diversity of novel microeukaryotes.

Labyrinthulomycetes are heterotrophic Stramenopiles which are abundant and

diverse in a wide range of marine and freshwater habitats where they play important

roles as saprotrophs/decomposers [89, 90]. Some Labyrinthulomycetes are also known

as important symbionts (parasites, mutualists or commensals) of marine organisms,

including brown algal seaweeds [91-93]. Although most thraustochytrids are free

living, a few species have been associated with disease in marine metazoans, including

the quahog parasite QPX which parasitizes clams. Several of the OTUs in our study

clustered in three highly supported clades within Labyrinthulida (Fig 3), and had high

similarity to *Aplanochytrium* labyrinthulids associated with various hosts such as corals

[94], the pseudoparenchymatous brown alga *Elachista* sp. [93], seagrass [95] and sea

stars [96].

Certain Oomycetes are common parasites of brown and red algae [37, 97-99],

and *Olpidiopsis* species have been shown to have cosmopolitan occurrence and broad

host ranges [98, 100]. In this study, we found OTUs clustering within Olpidiopsidales.

Some of these were highly similar to reference sequences of *Anisolpidium rosenvingei*

and *A*. *ectocarpii* (Fig 3), which are parasites of filamentous brown algae [36]. Others

displayed high similarity to *Pontisma lagenidioides* and "*Ectrogella*" (lnfLicSC1,

SC2-a, SC3 and SC4) which parasitizes the diatom *Licmophora* sp. and the red algae

Ceramium virgatum [101, 102]. Further, we detected two novel OTUs with no close

relatives among available reference sequences (OTU 214; 90.9 % identity to JN635125

and OTU 290; 92.4 % identity to KT273921, Fig. 3), of which the closest known

relative is an Oomycete parasitizing diatoms [101, 103].

Cercozoan diversity was dominated by vampyrellid amoebae, with four OTUs

forming a distinct clade with GU385680, which was previously detected on

Ascophyllum nodosum used as live bait wrapping [79]. Vampyrellids exhibit a wide

diversity of feeding strategies and often feed omnivorously; it is striking that this clade

comprises only lineages associated with brown algae, suggesting a specific association.

Phytomyxids, biotrophic parasites of angiosperms, oomycetes, and stramenopile algae

[39], were represented by two OTUs in a clade of marine sediment-derived sequences,

so there is no direct evidence for a specific interaction with macroalgae. OTU 242

groups with endomyxan Novel Clade 9 [104], members of which are found in a wide

range of environment types and have been hypothesized to be parasites of algae.

Fungi associated with brown algae can have everything from detrimental to

beneficial effects on their hosts [105, and references therein]. Two of the Ascomycota

OTUs clustered with taxa known to be parasites of brown algae such as *Sarocladium*

and *Acremonium* [106] (Fig. 5). Several of the basidiomycete OTUs clustered with

reference sequences of *Cryptococcus* and *Cystofilobasidium* that have previously been

found associated to various marine invertebrates, algae and seaweeds (Fig. 5) [107-

110]. In addition, within Chytridiomycota there were several OTUs that clustered with

Rhizophydium sp., which comprise parasites with broad host ranges known to infect

both macroalgae and protists [111, 112]. One OTU (OTU 76; Fig. 5) also displayed

high similarity to *Chytridium polysiphoniae*, a parasite of brown algae [113, 114]. The

final chytrid OTU clustered with *Mesochytrium penetrans*, a parasite of the unicellular

green algae *Chlorococcum minutum* [115], indicating that this could represent a parasite

of one of the green algae found in our study, or that relatives of *M. penetrans* have a

broader host range than currently known.

Conclusions

Brown algal holobionts are major marine habitat formers that provide food and shelter for diverse communities of both macro- and microorganisms. Our findings demonstrate that microeukaryotes should be considered as an integral part of brown algal holobionts, and provides important baseline data for future studies of seaweed-associated microorganisms. The prerequisite to understand the ecology of brown algae and their host-associated microbes, is to know their diversity, distribution and community dynamics. Consequently, in order to obtain an understanding of the interdependence of all the different players in brown algal holobionts it is critical to include the so far largely neglected eukaryotic and microeukaryotic partners in future holobiont investigation. It is however also important to remember that microeukaryotes interact not only with the host but also with the prokaryotes and viruses that share the same host environment, and future holobiont surveys should ideally include all of these components.

Methods

Sample collection

Five individuals each of *Ascophyllum nodosum*, *Fucus vesiculosus*, *Laminaria digitata* and *Saccharina latissima* were sampled from the Oslofjord (59°40′26.9″N, 10°35′13.199″E), while five individuals each of *F. vesiculosus*, *F. serratus* *Himanthalia elongata*, *Saccorhiza polyschides* and *L. digitata* were collected at Newton's Cove (Dorset, UK: 50°36′18″N, 2°26′58″W) in October 2015. Samples were collected by free diving and shore-based collection, kept cool in separate containers of seawater while transported to molecular laboratories where they were immediately subjected to subsampling of tissues for molecular analyses.

All samples were handled under laminar flow hoods to limit their exposure to

airborne contaminants. The brown algae were rinsed in sterile artificial seawater (ASW)

to remove loosely attached organisms (but not true epibionts) and debris from the

surface. This was done in three consecutive steps by vortexing the samples in sterile 50

mL tubes with ASW for 5-10 seconds. The rinsed individuals were placed in sterile

petri dishes, and one subsample of the algal thallus (approximately 1 cm^2 squares) per

individual was excised and placed in separate 2 mL tubes. All algae looked healthy; we

did not specifically target potentially infected tissues.

The samples were freeze-dried under sterile conditions by perforating the

sample tube lids and placing the sample tubes inside sterile culture flasks with 0.2 um

filter caps (Thermo Scientific[TM] Nunclon[TM]). After freeze-drying, samples were

weighted, placed in new 2 mL sample tubes, and stored at -80 °C until DNA extraction.

Endophyte sampling

As part of a previous student project (Attwood 2005), brown algal endophytes were

selected from the B. Rinkel collection at The Natural History Museum, London. The

samples were isolated from hosts as shown in Table 5. Partial 18S-ITS1 rDNA

sequences were amplified using the phaeophyte-specific primer DICI [116] and the

general primer TW7 [117]. The 18S region of this amplicon did not overlap with the

V4 region sequenced by the metabarcoding approach, so longer 18S amplicons were

generated from the brown macroalgal samples used for this study as described below,

to enable as much direct comparison of our newly generated sequences with those from

the endophytes. Full details of the DNA extraction, PCR, and sequencing methods used

for the endophytes are given in Attwood 2005.

Molecular methods

The freeze-dried samples were mechanically disrupted by adding two sterile tungsten

carbide beads to each sample tube and tissue-lyzed at 20Hz for 2 minutes, or longer,

until the tissues were completely pulverised. Thereafter, a modified lysis buffer [118]

was added to the tubes proportional to sample weight. This lysis buffer contains

antioxidant compounds such as PVP (Polyvinylpyrrolidone) and BSA (Bovine Serum

Albumin) that bind polyphenols, and has high salt concentration which decrease the

levels of coextracted polysaccharides. The samples were homogenised and divided in

two or more tubes so that the sample used for DNA isolation did not exceed 20 mg dry

weight tissue. DNA was extracted following the protocol by Snirc *et al*. [118].

PCR amplification of the V4 region of the 18S rDNA gene was performed with

'pan eukaryotic' primers; the forward primer used was a slightly modified version of

the TAReuk454FWD1 primer (5'-CCAGCASCYGCGGTAAT**W**CC-3') and reverse

primer TAReukREV3 (5'-ACTTTCGTTCTTGATYRA-3') [119]. PCR reactions (25

μl) contained 1x KAPA HiFi HotStart ReadyMix (Kapa Biosystems), 0.4 μM of each

primer, 0.5 mg/mL Bovine serum albumin (BSA; Promega) and 1 μl genomic DNA.

The PCR program had an initial denaturation step at 98°C for 2 min, 15 cycles of 30 s

at 98°C, 30 s at 53°C and 45 s at 72°C, then 20 similar cycles except that the annealing

temperature was 48°C, and a final elongation step at 72°C for 10 min. DNA was titrated

into several dilutions to take into account potential inhibiting substances from the

brown algae. Each brown algal individual sample was amplified separately. PCR

products were purified and eluted (20 μl) with ChargeSwitch PCR Clean-Up kit

(ThermoFisher Scientific), and quantified with the dsDNA BR Assay Kit and Qubit 2.0

Fluorometer (ThermoFisher Scientific), before they were pooled equimolarly,

according to species and geographic location; e.g., the purified PCR products of five

Fucus vesiculosus individuals from Norway were pooled into one sample before

sequencing library preparation.

Sequencing libraries were prepared using the TruSeq Nano DNA Library

Preparation Kit at The Natural History Museum, London UK, following the

manufacturer's protocol. High throughput sequencing was conducted on a MiSeq v3

flow cell using 2x300 bp paired-end reads. Base calling was performed by Illumina

Real Time Analysis v1.18.54 and was demultiplexed and converted to FastQ files with

Illumina Bcl2fastq v1.8.4.

The complete sequencing dataset is available at the European Nucleotide

Archive under the study accession number XXXXXXXX (http://www.ebi.ac.uk/ena/

data/view/ XXXXXXXX).

To obtain full length 18S sequences of the brown algal host species, three DNA

isolates from each brown algal species were subjected to PCR amplification using the

general eukaryotic 18S primers NSF83 (5'-GAAACTGCGAATGGCTCATT-3' [120])

and 1528R (5'-TCCTTCTGCAGGTTCACCTAC-3' [121]) utilising illustra™

PuReTaq Ready-To-Go™ PCR beads (GE Healthcare). PCR reactions contained 1

Illustra PuReTaq Ready-To-Go-bead, 1μl of DNA template, 0.5 μM of each primer,

filled to 25μl with water. PCR settings were: initial denaturation at 94°C for 2 min; 35

cycles of denaturation at 94°C for 45 s, annealing at 55°C for 30 s, elongation at 68°C

for 1.5 min; final elongation step at 68°C for 15 min. The PCR products were cloned

using the TOPO TA Cloning Kit for Sequencing (Invitrogen), according to the

manufacturer's protocol. The clones were grown overnight in Luberia-Bertoni (LB)

media amended with 50 μg/mL ampicillin. From each brown algal species, 20-30

bacterial colonies/cloned fragments, were subjected to PCR reactions with the vector

primers T7 and M13R and using ca 0.5 μl of the bacterial suspension as template. PCR

settings were: initial denaturation at 94°C for 10 min; 30 cycles of denaturation at 94°C

for 30 s, annealing at 53°C for 1 min, elongation at 72°C for 2 min; final elongation

step at 72°C for 10 min. The PCR products were cleaned using illustra™ ExoProStar

1-Step (GE Healthcare) and then Sanger sequenced (GATC Eurofins Genomics,

Germany).

Bioinformatics

The sequence data was processed using a workflow for MiSeq amplicon data

(https://github.com/ramalok/amplicon_processing). In short, prior to downstream

analyses error correction was done with BayesHammer [122], before merging the

paired-end reads with PEAR [123]. Thereafter, quality filtering, length check and fasta

conversion (from fastq to fasta) was done with USEARCH [124] before reads were put

into the correct 5' – 3' direction using profile hidden Markov models (hmmer-3.0rc2).

Dereplication, sorting by abundance and discarding of singletons was done using

USEARCH 64bit version. Clustering of reads into Operational Taxonomic Units

(OTUs) was done using the UPARSE algorithm as implemented in USEARCH,

applying a 97% sequence similarity threshold. *De-novo* and reference-based chimera

checking was performed and chimeric sequences discarded. OTUs were taxonomically

assigned using BLASTn against the PR2 v. 4.12.0 [125], GenBank [126] and SILVA

SSU [127] databases. We used a relaxed BLAST search strategy (BLASTn parameters:

-evalue 0.0001 -perc_identity 75) to retrieve distant sequences, and only retrieved the

best hit per query sequence. The PR2 taxonomy was used for taxonomy assignment for

the majority of the following analyses.

"Post-UPARSE trimming" & Diversity analyses

Species (OTU) composition analyses and further trimming of the dataset was done

using R v. 3.6.3 (R Core Team 2013). As we expected the sequence dataset was heavily

dominated by brown algal host reads, and we did therefore not normalise the dataset

because this would have led to removal of true diversity since most of the other OTUs

had very low read numbers compared to the brown algal OTUs. Further, all OTUs

assigned to Metazoa were removed from the dataset in addition to OTUs assigned to

land plants. The 57 OTUs (16.3% of the total number of OTUs) that matched those of

Metazoa are shown in Supplementary Figure S5, Supplementary Table S5.The trimmed

dataset, containing only non-metazoan eukaryotes, was used for further analyses.

The taxonomic composition of OTUs was visualized using the ggplot2 package

v. 3.3.2. [128]. To investigate how the composition of (micro)eukaryotes associated

with the different brown algal hosts, Bray-Curtis pairwise distances were calculated

using vegan v. 2.5-6 [129] for all brown algal samples (including/excluding brown algal

OTUs) and visualised using non-metric multidimensional scaling (NMDS) cluster

analyses (ggplot2 package v. 3.3.2 [128]). Analysis of similarities [ANOSIM; 130] was

calculated using vegan v. 2.5-6 [129] to test if there was a statistical significant

differences between clusters of libraries corresponding to different brown algal host

genera. The similarity percentage [SIMPER; 130] was calculated using the

simper.pretty.R script [131] to identify OTUs that contributed the most to the

dissimilarity observed. Indicator species analyses (IndVal) were carried out using the

indicspecies package v. 1.7.9. [132], to identify OTUs most preferentially associated

with specific brown algal host genera. Venn diagrams intersects were computed using

the function overLapper and plotted using the function vennPlot in the R-package

systemPipeR v. 1.24.3 [133].

Phylogenetic analysis

All OTUs were places into a global eukaryotic alignment of near full-length 18S reference sequence using MAFFT v7.300b with the 'einsi' option for global homology and long gaps for the reference sequences [134]. The shorter OTUs from the V4 region (300-450 bp) were added to the reference alignment using the –addfragment option in MAFFT and ambiguously aligned sites were removed with TrimAl [135] with –gt 0.3 and –st 0.001. Phylogenetic trees were built with RAxML v8.0.26 under the GTRGAMMA model and the autoMRE bootsopping criteria [136]. Based on a combination of the phylogenetic affinities of the OTUs and the taxonomic assignment from PR2, individual phylogenies of the most abundant/diverse groups of microeukaryotes – and those with known associations with brown algae - were constructed: i.e Fungi, Oomycetes & labyrinthulids, Cercozoa, diatoms, red, brown, and green algae, ciliates, other alveolates, Amoebozoa, and centrohelid heliozoans.

Individual sequence datasets were constructed including OTUs from each group whose taxonomic annotation and phylogenetic position were concordant, and their closest Blastn matches from GenBank. Sequence alignments were carried out in MAFFT as described above. Phylogenetic trees were built with MrBayes v.3.2.6 [137]. Two separate MC3 runs with randomly generated starting trees were carried out for 4 million generations each with 1 cold and 3 heated chains. The evolutionary model applied a GTR substitution matrix, with a 4-category autocorrelated gamma correction. All parameters were estimated from the data. The trees were sampled every 1000 generations and the first 1 million generations discarded as burn-in. All phylogenetic analyses were carried out on the Cipres server [138]. Heatmaps were made for all phylogenetic trees to display the proportional read abundance ($\log_{10}$) for each OTU in the different samples.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The dataset generated are available in XX repository, under the accession XXXX-XXXX (link to dataset).

Competing interest

The authors declare that they have no competing interests

Funding

This study has been supported financially by research funds "Sunniva og Egil Baardseths legat, til støtte for forskning på makroalger" and from the University of Oslo and the Natural History Museum, London to MFMB.

Authors' contributions

Acknowledgements

We thank Jonas Thormar and Jens Ådne Haga for help with field sampling of brown algae in Oslofjord. Drøbak Biological Station and Hans Erik Karlsen are thanked for providing sampling equipment and laboratory facilities for sample preparations.

Figure titles and figure legends

Figure 1. Diversity of eukaryotes in brown algal holobionts.

Proportional abundance of OTUs taxonomically assigned to eukaryotic major

Kingdoms or 'supergroup' in the brown algal samples of *Fucus serratus*, *F.*

vesiculosus, *Ascophyllum nodosum*, *Himanthalia elongata*, *Saccorhiza polyschides*,

Laminaria digitata and *Saccharina latissima*. The sampling location for each brown

alga is shown in parenthesis; NO = Norway and UK = The United Kingdom, together

with the total number of OTUs per library. The asterisk (NO*) represent *F. vesiculosus*

sampled in Norway, May 2013. All other samples were collected in October 2015.

Figure 2. Abundance of eukaryotic taxonomic groups.

Percentage representation of OTUs assigned to the different taxonomic groups at (A)

Kingdom or 'supergroup' level and at lower taxonomic level (B) in all of the brown

algal samples combined. To see the percentage of reads and percentage of OTUs for all

the taxonomic groups per brown algal sample, see Supplementary table S2.

Figure 3. Oomycetes and Labyrinthulids phylogeny with heatmap representing

proportional read abundance (log$_{10}$) of OTUs per sample. Bayesian phylogeny of

Oomycetes and Labyrinthulomycetes with bootstrap values from maximum likelihood

analysis added. Black circles represent maximal support (posterior probability = 1, and

bootstrap support = 100%). OTUs from this study are shown in bold. The percentage

identity to the most similar reference sequence is shown in parenthesis after each OTU.

Host or type of environment the reference sequences were retrieved from in previous

studies are listed in parenthesis after each GenBank accession number. The scale bar

represents 0.2 substitutions per site. Abbreviations used for descriptions of

environment: env.S = environmental sample, marine sediment; env.W = environmental

sample, marine water; env.FW = environmental sample, fresh water; env.So =

environmental sample, soil. The heatmap illustrates the $\log_{10}$ read abundance for each

OTU in the brown algal samples of *Fucus serratus*, *F. vesiculosus*, *Ascophyllum*

nodosum, *Himanthalia elongata*, *Saccorhiza polyschides*, *Laminaria digitata* and

Saccharina latissima. The sampling location for each brown alga is shown in

parenthesis; NO = Norway and UK = The United Kingdom. All samples were collected

in October 2015, except *F. vesiculosus* (NO*) which was sampled in Norway, May

2013.

Figure 4. Cercozoa: phylogeny with heatmap representing proportional read

abundance ($\log_{10}$) of OTUs per sample. Bayesian phylogeny of Cercozoa with

bootstrap values from maximum likelihood analysis added. Black circles represent

maximal support (posterior probability = 1, and bootstrap support = 100%). OTUs from

this study are shown in bold. The percentage identity to the most similar reference

sequence is shown in parenthesis after each OTU. Host or type of environment the

reference sequences were retrieved from in previous studies are listed in parenthesis

after each GenBank accession number. The scale bar represents 0.1 substitutions per

site. Abbreviations used for descriptions of environment: env.S = environmental

sample, marine sediment; env.W = environmental sample, marine water; env.FW =

environmental sample, fresh water; env.BW = environmental sample, brackish water;

env.S/W = environmental sample, marine sediment or water; env.So = environmental

sample, soil. The heatmap illustrates the $\log_{10}$ read abundance for each OTU in the

brown algal samples of *Fucus serratus*, *F. vesiculosus*, *Ascophyllum nodosum*,

Himanthalia elongata, *Saccorhiza polyschides*, *Laminaria digitata* and *Saccharina*

latissima. The sampling location for each brown alga is shown in parenthesis; NO =

Norway and UK = The United Kingdom. All samples were collected in October 2015,

except *F. vesiculosus* (NO*) which was sampled in Norway, May 2013.

Figure 5. Fungi: phylogeny with heatmap representing proportional read

abundance ($\log_{10}$) of OTUs per sample. Bayesian phylogeny of Fungi with bootstrap

values from maximum likelihood analysis added. Black circles represent maximal

support (posterior probability = 1, and bootstrap support = 100%). OTUs from this

study are shown in bold. The percentage identity to the most similar reference sequence

is shown in parenthesis after each OTU. Host or type of environment the reference

sequences were retrieved from in previous studies are listed in parenthesis after each

GenBank accession number. The scale bar represents 0.07 substitutions per site.

Abbreviations used for descriptions of environment: env.S = environmental sample,

marine sediment; env.W = environmental sample, marine water; env.FW =

environmental sample, fresh water; env.So = environmental sample, soil; env.A =

environmental sample, air. The heatmap illustrates the $\log_{10}$ read abundance for each

OTU in the brown algal samples of *Fucus serratus*, *F. vesiculosus*, *Ascophyllum*

nodosum, *Himanthalia elongata*, *Saccorhiza polyschides*, *Laminaria digitata* and

Saccharina latissima. The sampling location for each brown alga is shown in

parenthesis; NO = Norway and UK = The United Kingdom. All samples were collected

in October 2015, except *F. vesiculosus* (NO*) which was sampled in Norway, May

2013.

Figure 6. Bacillariophyceae: phylogeny with heatmap representing proportional

read abundance (log$_{10}$) of OTUs per sample. Bayesian phylogeny of diatoms with

bootstrap values from maximum likelihood analysis added. Black circles represent

maximal support (posterior probability = 1, and bootstrap support = 100%). OTUs from

this study are shown in bold. The percentage identity to the most similar reference

sequence is shown in parenthesis after each OTU. Host or type of environment the

reference sequences were retrieved from in previous studies are listed in parenthesis

after each GenBank accession number. The scale bar represents 0.06 substitutions per

site. Abbreviations used for descriptions of environment: env.S = environmental

sample, marine sediment; env.W = environmental sample, marine water; env.Si =

environmental sample, sea ice; GC = gut content. The heatmap illustrates the log$_{10}$ read

abundance for each OTU in the brown algal samples of *Fucus serratus*, *F. vesiculosus*,

Ascophyllum nodosum, *Himanthalia elongata*, *Saccorhiza polyschides*, *Laminaria*

digitata and *Saccharina latissima*. The sampling location for each brown alga is shown

in parenthesis; NO = Norway and UK = The United Kingdom. All samples were

collected in October 2015, except *F. vesiculosus* (NO*) which was sampled in Norway,

May 2013.

Figure 7. Brown-Red-Green: phylogeny with heatmap representing proportional

read abundance (log$_{10}$) of OTUs per sample. Bayesian phylogeny of Phaeophyceae,

Rhodophyta and Chlorophyta with bootstrap values from maximum likelihood analysis

added. OTUs from this study are shown in bold. The percentage identity to the most

similar reference sequence is shown in parenthesis after each OTU. Host or type of

environment the reference sequences were retrieved from in previous studies are listed

in parenthesis after each GenBank accession number. The longer 18S sequences

generated by cloning and sanger sequencing in this study are shown in bold and

includes which brown algal host species they were amplified from. The brown algal

endophytes isolated and sequenced by Attwood (2005) are all in the 'brown

endophytes' clade and are shown in capital letters. The scale bar represents 0.7

substitutions per site. The heatmap illustrates the $\log_{10}$ read abundance for each OTU

in the brown algal samples of *Fucus serratus*, *F. vesiculosus*, *Ascophyllum nodosum*,

Himanthalia elongata, *Saccorhiza polyschides*, *Laminaria digitata* and *Saccharina*

latissima. The sampling location for each brown alga is shown in parenthesis; NO =

Norway and UK = The United Kingdom. All samples were collected in October 2015,

except *F. vesiculosus* (NO*) which was sampled in Norway, May 2013.

Figure 8. Brown algal endophytes

Light microscopy images of phaeophyte endophytes isolated from *Fucus serratus*,

Mastocarpus stellatus, *Chorda filum*, *Osmundea pinnatifida*, and *Chondrus crispus*. (I)

Streblonema sp.: A – terminal sporangium, showing apical spore release; B – branching

thallus from spherical cells; C – growth pattern of new filament; D – discoid parietal

plastid. (II) *Myrionema strangulans*: A – Phaeophyceae hair; B – plurilocular

sporangia; C- two different cell shapes; D – unilocular lateral sporangium. (III)

Microspongium tenuisimum: A - plurilocular sporangia releasing zoospores; B -

Phaeophyceae hair; C – bifurcating branch; D – ovate cell, showing plate-like plastid

with prominent pyrenoids; E – waisted cross cell wall. (IV) *Mytriotrichia clavaeformis*:

A - Phaeophyceae hair; B – clubbed terminal cell; C - unilocular clustering sporangia;

D – empty clubbed terminal sporangium. (V) *Ectocarpus* sp.: A – ribbon-shaped

chloroplast; B – pyrenoid; C – pseudohairs; D – pseudohairs shoing growth from

filament; E – branching growth form. (VI) *Chorda filum*: A – densely pigmented cells;

B – prostrate clumps of branched filaments.

Figure 9. Community composition. Non metric multidimensional scaling (NMDS)

analysis of the composition of eukaryotic OTUs in the brown algal samples based on

Bray-Curtis pairwise distances. (A) including brown algal/Phaeophyceae OTUs and

(B) excluding brown algal/Phaeophyceae OTUs.

Figure 10. Shared and unique OTUs in the brown algal orders.

Venn diagrams displaying the number of shared and unique OTUs between the brown

algal host orders Fucales, Laminariales and Tilopteridales. To see the taxonomic

assignment for the shared OTUs see Table 4, and for the unique OTUs see

Supplementary Table S4.

Supplementary figures:

Supplementary figure 1. Ciliates: phylogeny with heatmap representing

proportional read abundance (log$_{10}$) of OTUs per sample. Bayesian phylogeny of

Ciliates with bootstrap values from maximum likelihood analysis added. Black circles

represent maximal support (posterior probability = 1, and bootstrap support = 100%).

OTUs from this study are shown in bold. The percentage identity to the most similar

reference sequence is shown in parenthesis after each OTU. Host or type of

environment the reference sequences were retrieved from in previous studies are listed

in parenthesis after each GenBank accession number. The scale bar represents 0.2

substitutions per site. Abbreviations used for descriptions of environment: env.S =

environmental sample, marine sediment; env.W = environmental sample, marine water;

env.FW = environmental sample, fresh water; env.WW = environmental sample, waste

water; env.BW = environmental sample, brackish water; env.S/W = environmental

sample, marine sediment or water; env.MBF = environmental sample, marine biofilm;

env.FBF = environmental sample, freshwater biofilm; env.So = environmental sample,

soil. The heatmap illustrates the $\log_{10}$ read abundance for each OTU in the brown algal

samples of *Fucus serratus*, *F. vesiculosus*, *Ascophyllum nodosum*, *Himanthalia*

elongata, *Saccorhiza polyschides*, *Laminaria digitata* and *Saccharina latissima*. The

sampling location for each brown alga is shown in parenthesis; NO = Norway and UK

= The United Kingdom. All samples were collected in October 2015, except *F.*

vesiculosus (NO*) which was sampled in Norway, May 2013.

Supplementary figure 2. Apicomplexa: phylogeny with heatmap representing

proportional read abundance ($\log_{10}$) of OTUs per sample. Bayesian phylogeny of

Apicomplexa and perkinsids with bootstrap values from maximum likelihood analysis

added. Black circles represent maximal support (posterior probability = 1, and bootstrap

support = 100%). OTUs from this study are shown in bold. The percentage identity to

the most similar reference sequence is shown in parenthesis after each OTU. Host or

type of environment the reference sequences were retrieved from in previous studies

are listed in parenthesis after each GenBank accession number. The scale bar represents

0.3 substitutions per site. Abbreviations used for descriptions of environment: env.S =

environmental sample, marine sediment; env.W = environmental sample, marine water;

env.FW = environmental sample, fresh water; env.I = environmental sample, ice; GC

= gut content. The heatmap illustrates the $\log_{10}$ read abundance for each OTU in the

brown algal samples of *Fucus serratus*, *F. vesiculosus*, *Ascophyllum nodosum*,

Himanthalia elongata, *Saccorhiza polyschides*, *Laminaria digitata* and *Saccharina*

latissima. The sampling location for each brown alga is shown in parenthesis; NO =

Norway and UK = The United Kingdom. All samples were collected in October 2015,

except *F. vesiculosus* (NO*) which was sampled in Norway, May 2013.

Supplementary figure 3. Amoebozoa and Centroheliozoa phylogeny with heatmap

representing proportional read abundance (log$_{10}$) of OTUs per sample. Bayesian

phylogeny of Amoebozoa and Centroheliozoa with bootstrap values from maximum

likelihood analysis added. Black circles represent maximal support (posterior

probability = 1, and bootstrap support = 100%). OTUs from this study are shown in

bold. The percentage identity to the most similar reference sequence is shown in

parenthesis after each OTU. Host or type of environment the reference sequences were

retrieved from in previous studies are listed in parenthesis after each GenBank

accession number. The scale bar represents 0.2 and 0.08 substitutions per site for

Amoebozoa and Centroheliozoa respectively. Abbreviations used for descriptions of

environment: env.FWS = environmental sample, freshwater sediment; env.W =

environmental sample, marine water; env.So = environmental sample, soil. The

heatmap illustrates the log$_{10}$ read abundance for each OTU in the brown algal samples

of *Fucus serratus*, *F. vesiculosus*, *Ascophyllum nodosum*, *Himanthalia elongata*,

Saccorhiza polyschides, *Laminaria digitata* and *Saccharina latissima*. The sampling

location for each brown alga is shown in parenthesis; NO = Norway and UK = The

United Kingdom. All samples were collected in October 2015, except *F. vesiculosus*

(NO*) which was sampled in Norway, May 2013.

Supplementary figure 4. Dinoflagellates: phylogeny with heatmap representing proportional read abundance (log$_{10}$) of OTUs per sample. Bayesian phylogeny of Dinoflagellates with bootstrap values from maximum likelihood analysis added. Black circles represent maximal support (posterior probability = 1, and bootstrap support = 100%). OTUs from this study are shown in bold. The percentage identity to the most similar reference sequence is shown in parenthesis after each OTU. Host or type of environment the reference sequences were retrieved from in previous studies are listed in parenthesis after each GenBank accession number. The scale bar represents 0.08 substitutions per site. Abbreviations used for descriptions of environment: env.S = environmental sample, marine sediment; env.W = environmental sample, marine water; env.BW = environmental sample, brackish water. The heatmap illustrates the log$_{10}$ read abundance for each OTU in the brown algal samples of *Fucus serratus*, *F. vesiculosus*, *Ascophyllum nodosum*, *Himanthalia elongata*, *Saccorhiza polyschides*, *Laminaria digitata* and *Saccharina latissima*. The sampling location for each brown alga is shown in parenthesis; NO = Norway and UK = The United Kingdom. All samples were collected in October 2015, except *F. vesiculosus* (NO*) which was sampled in Norway, May 2013.

Supplementary figure 5. Diversity of metazoan OTUs detected in brown algal holobionts.

Proportional abundance of OTUs taxonomically assigned to Metazoa in the brown algal samples of *Fucus serratus*, *F. vesiculosus*, *Ascophyllum nodosum*, *Himanthalia elongata*, *Saccorhiza polyschides*, *Laminaria digitata* and *Saccharina latissima*. The sampling location for each brown alga is shown in parenthesis; NO = Norway and UK

= The United Kingdom. The asterisk (NO*) represent *F. vesiculosus* sampled in

Norway, May 2013. All other samples were collected in October 2015.

References:

1.   Teagle H, Hawkins SJ, Moore PJ, Smale DA: **The role of kelp species as**
biogenic habitat formers in coastal marine ecosystems. *Journal of*
Experimental Marine Biology and Ecology 2017, **492**:81-98.
2.   Bringloe TT, Starko S, Wade RM, Vieira C, Kawai H, De Clerck O, Cock JM,
Coelho SM, Destombe C, Valero M *et al*: **Phylogeny and Evolution of the**
Brown Algae. *Critical Reviews in Plant Sciences* 2020, **39**(4):281-321.
3.   Silberfeld T, Leigh JW, Verbruggen H, Cruaud C, de Reviers B, Rousseau F: **A**
multi-locus time-calibrated phylogeny of the brown algae (Heterokonta,
Ochrophyta, Phaeophyceae): Investigating the evolutionary nature of the
"brown algal crown radiation". *Mol Phylogenet Evol* 2010, **56**(2):659-674.
4.   Hyndes GA, Nagelkerken I, McLeod RJ, Connolly RM, Lavery PS, Vanderklift
MA: **Mechanisms and ecological role of carbon transfer within coastal**
seascapes. *Biol Rev Camb Philos Soc* 2014, **89**(1):232-254.
5.   Smale DA: **Impacts of ocean warming on kelp forest ecosystems**. *New*
Phytologist 2020, **225**(4):1447-1454.
6.   Christie H, Norderhaug KM, Fredriksen S: **Macrophytes as habitat for fauna**.
Marine Ecology Progress Series 2009, **396**:221-233.
7.   Gachon CM, Strittmatter M, Muller DG, Kleinteich J, Kupper FC: **Detection of**
differential host susceptibility to the marine oomycete pathogen
Eurychasma dicksonii by real-time PCR: not all algae are equal. *Appl*
Environ Microbiol 2009, **75**(2):322-328.
8.   Armstrong E, Rogerson A, Leftley JW: **The Abundance of Heterotrophic**
Protists Associated with Intertidal Seaweeds. *Estuarine, Coastal and Shelf*
Science 2000, **50**(3):415-424.
9.   Zuccaro A, Schoch CL, Spatafora JW, Kohlmeyer J, Draeger S, Mitchell JI:
Detection and identification of fungi intimately associated with the brown
seaweed Fucus serratus. *Appl Environ Microbiol* 2008, **74**(4):931-941.
10.   Egan S, Harder T, Burke C, Steinberg P, Kjelleberg S, Thomas T: **The seaweed**
holobiont: understanding seaweed-bacteria interactions. *FEMS Microbiol*
Rev 2013, **37**(3):462-476.
11.   Singh RP, Reddy CR: **Unraveling the Functions of the Macroalgal**
Microbiome. *Front Microbiol* 2015, **6**:1488.
12.   Hollants J, Leliaert F, De Clerck O, Willems A: **What we can learn from**
sushi: a review on seaweed-bacterial associations. *FEMS Microbiol Ecol*
2013, **83**(1):1-16.
13.   McKeown DA, Schroeder JL, Stevens K, Peters AF, Saez CA, Park J, Rothman
MD, Bolton JJ, Brown MT, Schroeder DC: **Phaeoviral Infections Are Present**
in Macrocystis, Ecklonia and Undaria (Laminariales) and Are Influenced
by Wave Exposure in Ectocarpales. *Viruses* 2018, **10**(8).

14. McKeown DA, Stevens K, Peters AF, Bond P, Harper GM, Brownlee C, Brown
MT, Schroeder DC: **Phaeoviruses discovered in kelp (Laminariales)**. *ISME*
J 2017, **11**(12):2869-2873.
15. Lachnit T, Thomas T, Steinberg P: **Expanding our Understanding of the**
Seaweed Holobiont: RNA Viruses of the Red Alga Delisea pulchra.
Frontiers in Microbiology 2016, **6**.
16. Muller DG, Kapp M, Knippers R: **Viruses in marine brown algae**. *Adv Virus*
Res 1998, **50**:49-67.
17. Skillings D: **Holobionts and the ecology of organisms: Multi-species**
communities or integrated individuals? *Biology & Philosophy* 2016,
31(6):875-892.
18. Margulis L, Fester R: **Symbiosis as a Source of Evolutionary Innovation:**
Speciation and Morphogenesis. *Symbiosis Philadelphia Pa* 1991, **11**.
19. van der Loos LM, Eriksson BK, Falcão Salles J: **The Macroalgal Holobiont**
in a Changing Sea. *Trends in Microbiology* 2019, **27**(7):635-650.
20. del Campo J, Bass D, Keeling PJ: **The eukaryome: Diversity and role of**
microeukaryotic organisms associated with animal hosts. *Functional*
Ecology 2019.
21. Bass D, Stentiford GD, Wang HC, Koskella B, Tyler CR: **The Pathobiome in**
Animal and Plant Diseases. *Trends Ecol Evol* 2019, **34**(11):996-1008.
22. de Bary A: **Die Erscheinung der Symbiose**. 1879:30 s.
23. Florez JZ, Camus C, Hengst MB, Buschmann AH: **A Functional Perspective**
Analysis of Macroalgae and Epiphytic Bacterial Community Interaction.
Frontiers in Microbiology 2017, **8**.
24. Lage OM, Graça AP: **Biofilms: an extra coat on macroalgae**. In: *Algae-*
Organisms for Imminent Biotechnology. Edited by Thajuddin N, Dhanasekaran
D: IntechOpen; 2016: 183-210.
25. Wahl M, Goecke F, Labes A, Dobretsov S, Weinberger F: **The second skin:**
ecological role of epibiotic biofilms on marine organisms. *Front Microbiol*
2012, **3**:292.
26. de Mesquita MMF, Crapez MAC, Teixeira VL, Cavalcanti DN: **Potential**
interactions bacteria-brown algae. *Journal of Applied Phycology* 2018,
31(2):867-883.
27. Abdul Malik SA, Bedoux G, Garcia Maldonado JQ, Freile-Pelegrín Y, Robledo
D, Bourgougnon N: **Defence on surface: macroalgae and their surface-**
associated microbiome. In: *Seaweeds Around the World: State of Art and*
Perspectives. 2020: 327-368.
28. Zhang R, Chang L, Xiao L, Zhang X, Han Q, Li N, Egan S, Wang G: **Diversity**
of the epiphytic bacterial communities associated with commercially
cultivated healthy and diseased Saccharina japonica during the harvest
season. *Journal of Applied Phycology* 2020, **32**(3):2071-2080.
29. Rogerson A: **On the abundance of marine naked amoeba on the surface of**
five species of macroalgae. *FEMS Microbiology Letters* 1991, **85**(4):301-312.
30. Baardseth E: **Some aspects of the native intercellular substance in Fucaceae**.
Proc Intl Seaweed Symp 1969, **6**:53-60.
31. Lam C, Grage A, Schulz D, Schulte A, Harder T: **Extracts of North Sea**
macroalgae reveal specific activity patterns against attachment and
proliferation of benthic diatoms: a laboratory study. *Biofouling* 2008,
24(1):59-66.

32.   Totti C, Poulin M, Romagnoli T, Perrone C, Pennesi C, De Stefano M:
Epiphytic diatom communities on intertidal seaweeds from Iceland. *Polar*
Biology 2009, **32**(11):1681-1691.
33.   Rinkel BE, Hayes P, Gueidan C, Brodie J: **A Molecular Phylogeny of**
Acrochaete and Other Endophytic Green Algae (Ulvales, Chlorophyta)1.
Journal of Phycology 2012, **48**(4):1020-1027.
34.   Raghukumar S: **Ecology of the marine protists, the Labyrinthulomycetes**
(Thraustochytrids and Labyrinthulids). *European journal of protistology*
2002, **145**:127-145.
35.   Raghukumar S, Damare VS: **Increasing evidence for the important role of**
Labyrinthulomycetes in marine ecosystems. *Botanica Marina* 2011, **54**:3-11.
36.   Gachon CMM, Strittmatter M, Badis Y, Fletcher KI, West PV, Müller DG:
Pathogens of brown algae: culture studies of Anisolpidium ectocarpii and
A. rosenvingei reveal that the Anisolpidiales are uniflagellated oomycetes.
European Journal of Phycology 2017, **52**(2):133-148.
37.   Strittmatter M, Gachon CMM, Müller DG, Kleinteich J, Heesch S, Tsirigoti A,
Katsaros C, Kostopoulou M, Küpper FC: **Intracellular eukaryotic pathogens**
in brown macroalgae in the Eastern Mediterranean, including LSU rRNA
data for the oomycete Eurychasma dicksonii *Diseases of Aquatic Organisms*
2013, **104**(1):1-11.
38.   Goecke F, Wiese J, Nunez A, Labes A, Imhoff JF, Neuhauser S: **A novel**
phytomyxean parasite associated with galls on the bull-kelp Durvillaea
antarctica (Chamisso) Hariot. *PLoS One* 2012, **7**(9):e45358.
39.   Neuhauser S, Kirchmair M, Bulman S, Bass D: **Cross-kingdom host shifts of**
phytomyxid parasites. *BMC Evolutionary Biology* 2014, **14**(33).
40.   Murúa P, Goecke F, Westermeier R, van West P, Küpper FC, Neuhauser S:
Maullinia braseltonii sp. nov. (Rhizaria, Phytomyxea, Phagomyxida): A
Cyst-forming Parasite of the Bull Kelp Durvillaea spp. (Stramenopila,
Phaeophyceae, Fucales). *Protist* 2017, **168**(4):468-480.
41.   Küpper FC, Müller DG: **Massive occurrence of the heterokont and fungal**
parasites Anisolpidium, Eurychasma and Chytridium in Pylaiella littoralis
(Ectocarpales, Phaeophyceae). *Nova Hedwigia* 1999, **69**:381-389.
42.   Tourneroche A, Lami R, Burgaud G, Domart-Coulon I, Li W, Gachon C, Gèze
M, Boeuf D, Prado S: **The Bacterial and Fungal Microbiota of Saccharina**
latissima (Laminariales, Phaeophyceae). *Frontiers in Marine Science* 2020,
7.
43.   Zuccaro A, Schulz B, Mitchell JI: **Molecular detection of ascomycetes**
associated with Fucus serratus. *Mycological Research* 2003, **107**(12):1451-
1466.
44.   Vallet M, Strittmatter M, Murúa P, Lacoste S, Dupont J, Hubas C, Genta-Jouve
G, Gachon CMM, Kim GH, Prado S: **Chemically-Mediated Interactions**
Between Macroalgae, Their Fungal Endophytes, and Protistan Pathogens.
Frontiers in Microbiology 2018, **9**.
45.   Garbary DJ, MacDonald KA: **The Ascophyllum / Polysiphonia /**
Mycosphaerella Symbiosis. IV. Mutualism in the Ascophyllum /
Mycosphaerella Interaction. *Botanica Marina* 1995, **38**(1-6).
46.   Toxopeus J, Kozera CJ, O'Leary SJB, Garbary DJ: **A reclassification of**
Mycophycias ascophylli (Ascomycota) based on nuclear large ribosomal
subunit DNA sequences. *Botanica Marina* 2011, **54**(3).

47. Potin P: **Intimate Associations Between Epiphytes, Endophytes, and**
Parasites of Seaweeds. In: *Seaweed Biology.* 2012: 203-234.
48. Fredriksen S, Christie H, Andre Sæthre B: **Species richness in macroalgae and**
macrofauna assemblages on Fucus serratus L. (Phaeophyceae) and Zostera
marina L. (Angiospermae) in Skagerrak, Norway. *Marine Biology Research*
2005, **1**(1):2-19.
49. Bjærke MR, Fredriksen S: **Epiphytic macroalgae on the introduced brown**
seaweed Sargassum muticum (Yendo) Fensholt (Phaeophyceae) in
Norway. *Sarsia* 2010, **88**(5):353-364.
50. Heesch S, Peters AF, Broom JE, Hurd CL: **Affiliation of the parasite**
Herpodiscus durvillaeae (Phaeophyceae) with the Sphacelariales based on
DNA sequence comparisons and morphological observations. *European*
Journal of Phycology 2008, **43**(3):283-295.
51. Murúa P, Küpper FC, Muñoz LA, Bernard M, Peters AF: **Microspongium**
alariae in Alaria esculenta: a widely-distributed non-parasitic brown algal
endophyte that shows cell modifications within its host. *Botanica Marina*
2018, **61**(4):343-354.
52. Ellertsdóttir E, Peters AF: **High prevalence of infection by endophytic brown**
algae in populations of Laminaria spp. (Phaeophyceae). *Marine Ecology*
Progress Series 1997, **146**:135-143.
53. Amsler CD, Amsler MO, McClintock JB, Baker BJ: **Filamentous algal**
endophytes in macrophytic Antarctic algae: prevalence in hosts and
palatability to mesoherbivores. *Phycologia* 2009, **48**(5):324-334.
54. Del Campo J, Pons MJ, Herranz M, Wakeman KC, Del Valle J, Vermeij MJA,
Leander BS, Keeling PJ: **Validation of a universal set of primers to study**
animal-associated microeukaryotic communities. *Environ Microbiol* 2019,
21(10):3855-3861.
55. Bass D, del Campo J: **Micro-eukaryotes in animal and plant microbiomes:**
Ecologies of disease? *European Journal of Protistology* 2020.
56. Goecke F, Imhoff JF: **Microbial Biodiversity Associated with Marine**
Macroalgae and Seagrasses. In: *Marine macrophytes as foundation species 1.*
Edited by Olafsson E: CRC Press
Taylor & Francis Group; 2016: 3-25.
57. Quigley CTC, Capistrant-Fossa KA, Morrison HG, Johnson LE, Morozov A,
Hertzberg VS, Brawley SH: **Bacterial Communities Show Algal Host (Fucus**
spp.)/Zone Differentiation Across the Stress Gradient of the Intertidal
Zone. *Frontiers in Microbiology* 2020, **11**.
58. Longtin CM, Scrosati RA, Whalen GB, Garbary DJ: **Distribution of Algal**
Epiphytes across Environmental Gradients at Different Scales: Intertidal
Elevation, Host Canopies, and Host Fronds1. *Journal of Phycology* 2009,
45(4):820-827.
59. Ihua MW, FitzGerald JA, Guihéneuf F, Jackson SA, Claesson MJ, Stengel DB,
Dobson ADW: **Diversity of bacteria populations associated with different**
thallus regions of the brown alga Laminaria digitata. *Plos One* 2020, **15**(11).
60. Tourneroche A, Lami R, Hubas C, Blanchet E, Vallet M, Escoubeyrou K, Paris
A, Prado S: **Bacterial–Fungal Interactions in the Kelp Endomicrobiota**
Drive Autoinducer-2 Quorum Sensing. *Frontiers in Microbiology* 2019, **10**.
61. Vallet M, Chong Y-M, Tourneroche A, Genta-Jouve G, Hubas C, Lami R,
Gachon CMM, Klochkova T, Chan K-G, Prado S: **Novel α-Hydroxy γ-**

Butenolides of Kelp Endophytes Disrupt Bacterial Cell-to-Cell Signaling.
Frontiers in Marine Science 2020, **7**.

62. da Gama BAP, Plouguerné E, Pereira RC: **The Antifouling Defence**
Mechanisms of Marine Macroalgae. In: *Sea Plants.* 2014: 413-440.

63. Goecke F, Labes A, Wiese J, Imhoff JF: **Chemical interactions between**
marine macroalgae and bacteria. *Marine Ecology Progress Series* 2010,
409:267-299.

64. Moss B: **Apical Meristems and Growth Control in Himanthalia Elongata**
(S. F. Gray). *New Phytologist* 1969, **68**(2):387-397.

65. del Olmo A, Picon A, Nuñez M: **The microbiota of eight species of**
dehydrated edible seaweeds from North West Spain. *Food Microbiology*
2018, **70**:224-231.

66. Goecke F, Thiel V, Wiese J, Labes A, Imhoff JF: **Algae as an important**
environment for bacteria – phylogenetic relationships among new bacterial
species isolated from algae. *Phycologia* 2013, **52**(1):14-24.

67. Eggert A, Peters AF, Küpper FC: **The Potential Impact of Climate Change**
on Endophyte Infections in Kelp Sporophytes. In: *Seaweeds and their Role*
in Globally Changing Environments. 2010: 139-154.

68. Correa J: **Infections by pigmented algal endophytes: misuse of concepts and**
terminology. *Revista Chilena de Historia Natural* 1994, **67**:4-8.

69. Lage OM, Graça AP: **Biofilms: An Extra Coat on Macroalgae**. In: *Algae -*
Organisms for Imminent Biotechnology. 2016.

70. Tiffany MA: **Epizoic and Epiphytic Diatoms**. In: *The Diatom World.* 2011:
195-209.

71. Costa MMdS, Pereira SMB, Silva-Cunha MdGGd, Arruda PCd, Eskinazi-Leça
E: **Community structure of epiphytic diatoms on seaweeds in Northeastern**
Brazil. *Botanica Marina* 2016, **0**(0).

72. Baardseth E, Taasen JP: **Navicula dumontiae sp. nov., an Endophytic Diatom**
Inhabiting the Mucilage of Dumontia incrassata (Rhodophycea.pdf.
Norwegian Journal of Botany 1973, **20**:79-87.

73. Hasle GR: **Navicula endophytica Sp.Nov., A pennate diatom with an**
unusual mode of existence. *British Phycological Bulletin* 1968, **3**(3):475-480.

74. Klochkova TA, Pisareva NA, Park JS, Lee JH, Han JW, Klochkova NG, Kim
GH: **An endophytic diatom, Pseudogomphonema sp. (Naviculaceae,**
Bacillariophyceae), lives inside the red alga Neoabbottiella
(Halymeniaceae, Rhodophyta). *Phycologia* 2014, **53**(3):205-214.

75. Mayombo NAS, Majewska R, Smit AJ: **Diatoms associated with two South**
African kelp species:Ecklonia maximaandLaminaria pallida. *African*
Journal of Marine Science 2019, **41**(2):221-229.

76. Mayombo NAS, Majewska R, Smit AJ: **An Assessment of the Influence of**
Host Species, Age, and Thallus Part on Kelp-Associated Diatoms. *Diversity*
2020, **12**(10).

77. Gismervik I: **Podite carrying ciliates dominate the benthic ciliate**
community in the kelp forest. *Aquatic Microbial Ecology* 2004, **36**:305-310.

78. Bjorbækmo MFM, Evenstad A, Røsæg LL, Krabberød AK, Logares R: **The**
planktonic protist interactome: where do we stand after a century of
research? *The ISME Journal* 2019, **14**(2):544-559.

79. Haska CL, Yarish C, Kraemer G, Blaschik N, Whitlatch R, Zhang H, Lin S:
Bait worm packaging as a potential vector of invasive species. *Biological*
Invasions 2011, **14**(2):481-493.

80.   Lee Y: **The Circumscription of the Genus Halothrix Reinke (Chordariales,**
1166   **Phaeophyceae)**. *Algae* 2001, **16**(1):35-43.

1167   81.   Lane CE, Saunders GW: **Molecular investigation reveals epi/endophytic**
1168   **extrageneric kelp (Laminariales, Phaeophyceae) gametophytes colonizing**
1169   **Lessoniopsis littoralis thalli**. *Botanica Marina* 2005, **48**(5-6).

1170   82.   Fox CH, Swanson AK: **Nested PCR detection of microscopic life-stages of**
1171   **laminarian macroalgae and comparison with adult forms along intertidal**
1172   **height gradients**. *Marine Ecology Progress Series* 2007, **332**:1-10.

1173   83.   Hubbard CB, Garbary DJ, Kim KY, Chiasson DM: **Host specificity and**
1174   **growth of kelp gametophytes symbiotic with filamentous red algae**
1175   **(Ceramiales, Rhodophyta)**. *Helgoland Marine Research* 2004, **58**(1):18-25.

1176   84.   Schoenrock KM, McHugh TA, Krueger‐Hadfield SA, Edwards M: **Revisiting**
1177   **the 'bank of microscopic forms' in macroalgal‐dominated ecosystems**.
1178   *Journal of Phycology* 2020, **57**(1):14-29.

1179   85.   Robuchon M, Couceiro L, Peters AF, Destombe C, Valero M: **Examining the**
1180   **bank of microscopic stages in kelps using culturing and barcoding**.
1181   *European Journal of Phycology* 2014, **49**(1):128-133.

1182   86.   Bernard M, Rousvoal S, Jacquemin B, Ballenghien M, Peters AF, Leblanc C:
1183   **qPCR-based relative quantification of the brown algal endophyte**
1184   **Laminarionema elsbetiae in Saccharina latissima: variation and dynamics**
1185   **of host—endophyte interactions**. *Journal of Applied Phycology* 2017,
1186   **30**(5):2901-2911.

1187   87.   Gunnarsson K, Nielsen R: **Culture and field studies of Ulvellaceae and other**
1188   **microfilamentous green seaweeds in subarctic and arctic waters around**
1189   **Iceland**. *Nova Hedwigia* 2016, **103**(1-2):17-46.

1190   88.   Nielsen R, Gunnarsson K, Daugbjerg N, Petersen G: **Description ofUlvella**
1191   **eleganssp. nov. andU. islandicasp. nov. (Ulvellaceae, Ulvophyceae) from**
1192   **Iceland – a study based on morphology of species in culture andtufA gene**
1193   **sequences**. *European Journal of Phycology* 2014, **49**(1):60-67.

1194   89.   Nakai R, Naganuma T: **Diversity and Ecology of Thraustochytrid Protists in**
1195   **the Marine Environment**. In: *Marine Protists.* 2015: 331-346.

1196   90.   Pan J, del Campo J, Keeling PJ: **Reference Tree and Environmental**
1197   **Sequence Diversity of Labyrinthulomycetes**. *Journal of Eukaryotic*
1198   *Microbiology* 2017, **64**(1):88-96.

1199   91.   Raghukumar S: **Ecology of the marine protists, the Labyrinthulomycetes**
1200   **(Thraustochytrids and Labyrinthulids)**. *European Journal of Protistology*
1201   2002, **38**(2):127-145.

1202   92.   Gleason FH, van Ogtrop F, Lilje O, Larkum AWD: **Ecological roles of**
1203   **zoosporic parasites in blue carbon ecosystems**. *Fungal Ecology* 2013,
1204   **6**(5):319-327.

1205   93.   Mystikou A, Peters AF, Asensi AO, Fletcher KI, Brickle P, van West P, Convey
1206   P, Küpper FC: **Seaweed biodiversity in the south-western Antarctic**
1207   **Peninsula: surveying macroalgal community composition in the Adelaide**
1208   **Island/Marguerite Bay region over a 35-year time span**. *Polar Biology* 2014,
1209   **37**(11):1607-1619.

1210   94.   Siboni N, Rasoulouniriana D, Ben-Dov E, Kramarsky-Winter E, Sivan A, Loya
1211   Y, Hoegh-Guldberg OVE, Kushmaro A: **Stramenopile Microorganisms**
1212   **Associated with the Massive CoralFaviasp**. *Journal of Eukaryotic*
1213   *Microbiology* 2010.

95.   Medina-Pons FJ, Terrados J, López-López A, Yarza P, Rosselló-Móra R:
Evaluation of the 18S rRNA clone library approach to study the diversity
of the macroeukaryotic leaf-epiphytic community of the seagrass Posidonia
oceanica (L.) Delile. *Marine Biology* 2009, **156**(9):1963-1976.
96.   FioRito R, Leander C, Leander B: **Characterization of three novel species of**
Labyrinthulomycota isolated from ochre sea stars (Pisaster ochraceus).
Marine Biology 2016, **163**(8).
97.   Gachon CM, Sime-Ngando T, Strittmatter M, Chambouvet A, Kim GH: **Algal**
diseases: spotlight on a black box. *Trends Plant Sci* 2010, **15**(11):633-640.
98.   Badis Y, Klochkova TA, Brakel J, Arce P, Ostrowski M, Tringe SG, Kim GH,
Gachon CMM: **Hidden diversity in the oomycete genus Olpidiopsis is a**
potential hazard to red algal cultivation and conservation worldwide.
European Journal of Phycology 2020, **55**(2):162-171.
99.   Badis Y, Klochkova TA, Strittmatter M, Garvetto A, Murúa P, Sanderson JC,
Kim GH, Gachon CMM: **Novel species of the oomycete Olpidiopsis**
potentially threaten European red algal cultivation. *Journal of Applied*
Phycology 2018, **31**(2):1239-1250.
100. Sekimoto S, Klochkova TA, West JA, Beakes GW, Honda D: **Olpidiopsis**
bostrychiae sp. nov.: an endoparasitic oomycete that infects Bostrychia and
other red algae (Rhodophyta). *Phycologia* 2009, **48**(6):460-472.
101. Garvetto A, Perrineau MM, Dressler‐Allame M, Bresnan E, Gachon CMM:
"Ectrogella" **Parasitoids of the Diatom Licmophora sp. are Polyphyletic**.
Journal of Eukaryotic Microbiology 2019, **67**(1):18-27.
102. Buaya AT, Ploch S, Inaba S, Thines M: **Holocarpic oomycete parasitoids of**
red algae are not Olpidiopsis. *Fungal Systematics and Evolution* 2019.
103. Thines M, Nam B, Nigrelli L, Beakes G, Kraberg A: **The diatom parasite**
Lagenisma coscinodisci (Lagenismatales, Oomycota) is an early diverging
lineage of the Saprolegniomycetes. *Mycological Progress* 2015, **14**(9).
104. Bass D, Chao EE-Y, Nikolaev S, Yabuki A, Ishida K-I, Berney C, Pakzad U,
Wylezich C, Cavalier-Smith T: **Phylogeny of novel naked Filose and**
Reticulose Cercozoa: Granofilosea cl. n. and Proteomyxidea revised. *Protist*
2009, **160**:75-109.
105. Suryanarayanan TS: **Fungal Endosymbionts of Seaweeds**. In: *Biology of*
Marine Fungi. 2012: 53-69.
106. Vicente TFL, Gonçalves MFM, Brandão C, Fidalgo C, Alves A: **Diversity of**
fungi associated with macroalgae from an estuarine environment and
description of Cladosporium rubrum sp. nov. and Hypoxylon aveirense sp.
nov. *International Journal of Systematic and Evolutionary Microbiology* 2021.
107. Duarte AWF, Dayo-Owoyemi I, Nobre FS, Pagnocca FC, Chaud LCS, Pessoa
A, Felipe MGA, Sette LD: **Taxonomic assessment and enzymes production**
by yeasts isolated from marine and terrestrial Antarctic samples.
Extremophiles 2013, **17**(6):1023-1035.
108. Vaca I, Faúndez C, Maza F, Paillavil B, Hernández V, Acosta F, Levicán G,
Martínez C, Chávez R: **Cultivable psychrotolerant yeasts associated with**
Antarctic marine sponges. *World Journal of Microbiology and Biotechnology*
2012, **29**(1):183-189.
109. Lo Giudice A, Azzaro M, Schiaparelli S: **Microbial Symbionts of Antarctic**
Marine Benthic Invertebrates. In: *The Ecological Role of Micro-organisms*
in the Antarctic Environment. 2019: 277-296.

110. Ogaki MB, de Paula MT, Ruas D, Pellizzari FM, García-Laviña CX, Rosa LH:
Marine Fungi Associated with Antarctic Macroalgae. In: *The Ecological*
Role of Micro-organisms in the Antarctic Environment. 2019: 239-255.
111. Gromov BV, Plujusch AV, Mamkaeva KA: **Morphology and possible host**
range of Rhizophydium algavorum sp. nov. (Chytridiales) - an obligate
parasite of algae. *Protistology* 1999, **1**(2):62-65.
112. Frenken T, Alacid E, Berger SA, Bourne EC, Gerphagnon M, Grossart H-P,
Gsell AS, Ibelings BW, Kagami M, Küpper FC *et al*: **Integrating chytrid**
fungal parasites into plankton ecology: research gaps and needs.
Environmental Microbiology 2017, **19**(10):3802-3822.
113. Küpper FC, Maier I, Müller DG, Loiseaux - De Goer S, Guillou L:
Phylogenetic affinities of two eukaryotic pathogens of marine macroalgae,
Eurychasma dicksonii (Wright) Magnus and Chytridium polysiphoniae
Cohn. . *Cryptogamie, Algologie* 2006, **27**(2):165-184.
114. Muller DG, Kupper FC, Kupper H: **Infection experiments reveal broad host**
ranges of Eurychasma dicksonii (Oomycota) and Chytridium
polysiphoniae (Chytridiomycota), two eukaryotic parasites in marine
brown algae (Phaeophyceae). *Phycological Research* 1999, **47**(3):217-223.
115. Karpov SA, Letcher PM, Mamkaeva MA, Mamkaeva KA: **Phylogenetic**
position of the genus Mesochytrium (Chytridiomycota) based on zoospore
ultrastructure and sequences from the 18S and 28S rRNA gene. *Nova*
Hedwigia 2010, **90**(1-2):81-94.
116. Peters AF, Burkhardt E: **Systematic position of the kelp endophyte**
Laminarionema elsbetiae (Ectocarpales sensu lato, Phaeophyceae)
inferred from nuclear ribosomal DNA sequences. *Phycologia* 2019,
37(2):114-120.
117. White TJ: **Analysis of phylogenetic relationships by amplification and direct**
sequencing of ribosomal genes. *PCR protocols: a guide to methods and*
applications 1990:315-322.
118. Snirc A, Silberfeld T, Bonnet J, Tillier A, Tuffet S, Sun J-S: **Optimization of**
DNA Extraction from Brown Algae (Phaeophyceae) Based on a
Commercial Kit1. *Journal of Phycology* 2010, **46**(3):616-621.
119. Stoeck T, Bass D, Nebel M, Christen R, Jones MD, Breiner HW, Richards TA:
Multiple marker parallel tag environmental DNA sequencing reveals a
highly complex eukaryotic community in marine anoxic water. *Mol Ecol*
2010, **19 Suppl 1**:21-31.
120. Hendriks L, Goris A, Neefs J-M, Van De Peer Y, Hennebert G, De Wachter R:
The Nucleotide Sequence of the Small Ribosomal Subunit RNA of the Yeast
Candida albicans and the Evolutionary Position of the Fungi among the
Eukaryotes. *Systematic and Applied Microbiology* 1989, **12**(3):223-229.
121. Medlin L, Elwood HJ, Stickel S, Sogin ML: **The characterization of**
enzymatically amplified eukaryotic 16S-like rRNA-coding regions. *Gene*
1988, **71**(2):491-499.
122. Schirmer M, Ijaz UZ, D'Amore R, Hall N, Sloan WT, Quince C: **Insight into**
biases and sequencing errors for amplicon sequencing with the Illumina
MiSeq platform. *Nucleic Acids Res* 2015, **43**(6):e37.
123. Zhang J, Kobert K, Flouri T, Stamatakis A: **PEAR: A fast and accurate**
Illumina Paired-End reAd mergeR. *Bioinformatics* 2014, **30**:614-620.
124. Edgar RC: **UPARSE: highly accurate OTU sequences from microbial**
amplicon reads. *Nature methods* 2013, **10**.

125. Guillou L, Bachar D, Audic S, Bass D, Berney C, Bittner L, Boutte C, Burgaud
G, de Vargas C, Decelle J *et al*: **The Protist Ribosomal Reference database**
(PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences
with curated taxonomy. *Nucleic Acids Research* 2012, **41**(D1):D597-D604.
126. Sayers EW, Cavanaugh M, Clark K, Ostell J, Pruitt KD, Karsch-Mizrachi I:
GenBank. *Nucleic Acids Research* 2019.
127. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glockner FO:
SILVA: a comprehensive online resource for quality checked and aligned
ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res*
2007, **35**(21):7188-7196.
128. Wickham H: **ggplot2: elegant graphics for data analysis**: Springer 2016.
129. Oksanen J., Blanchet F. G., Friendly M, Kindt R, Legendre P, McGlinn D,
Minchin P. R., O'Hara R. B., Simpson G. L., Solymos P. *et al*: **vegan:**
Community Ecology Package. R package version 2.5-6. In.; 2019.
130. Clarke KR: **Non‐parametric multivariate analyses of changes in**
community structure. *Australian journal of ecology* 1993, **18**(1):117-143.
131. Steinberger A: **asteinberger9/seq_scripts v1.1 (Version v1.1). In**. Zenodo.
http://doi.org/10.5281/zenodo.4270481; 2020.
132. Cáceres MD, Legendre P: **Associations between species and groups of sites:**
indices and statistical inference. *Ecology* 2009, **90**(12):3566-3574.
133. Backman T, Girke T: **systemPipeR: NGS workflow and report generation**
environment. *BMC Bioinformatics* 2016, **17**:388.
134. Katoh K, Standley DM: **MAFFT multiple sequence alignment software**
version 7: Improvements in performance and usability. *Molecular Biology*
and Evolution 2013, **30**:772-780.
135. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T: **trimAl: A tool for**
automated alignment trimming in large-scale phylogenetic analyses.
Bioinformatics 2009, **25**:1972-1973.
136. Stamatakis A: **RAxML version 8: A tool for phylogenetic analysis and post-**
analysis of large phylogenies. *Bioinformatics* 2014, **30**:1312-1313.
137. Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S,
Larget B, Liu L, Suchard MA, Huelsenbeck JP: **MrBayes 3.2: Efficient**
Bayesian Phylogenetic Inference and Model Choice Across a Large Model
Space. *Systematic Biology* 2012, **61**(3):539-542.
138. Miller MA, Pfeiffer W, Schwartz T: **Creating the CIPRES Science Gateway**
for Inference of Large Phylogenetic Trees. In: *In Proceedings of the Gateway*
Computing Environments Workshop (GCE), New Orleans LA. 2010.
1350

Table 1. Summary of pairwise SIMPER analyses between brown algal host samples grouped at genus level. Showing the OTUs that contributed to more than 10% of the observed dissimilarity between host genera according to SIMPER, when Phaeophyceae OTUs were included in the analyses.

		OTUs contributing >10 % to the dissimilarity between brown algal hosts (Including Phaeophyceae OTUs)					
	Phylum	Stramenopiles					
	Class	Phaeophyceae					
	PR2 taxonomy	*Silvetia siliquosa*	*Fucus gardneri*	*Fucus gardneri*	*Silvetia siliquosa*	*Saccorhiza polyschides*	*Agarum clathratum*
	Inferred host taxonomy	*Ascophyllum nodosum*	*Fucus spp.*	*Fucus vesiculosus*	*Himanthalia elongata*	*Saccorhiza polyschides*	*Laminariales (L. digitata + S. latissima)*
SIMPER comparison	Dissimilarity (%)	OTU 2	OTU 49	OTU 41	OTU 5	OTU 3	OTU 1
Fucus – Ascophyllum	74.4	41.3	8.4	4.8			
Fucus – Himanthalia	99.8	20.0	5.1	9.6	36.6		
Fucus – Saccorhiza	97.9	21.1	5.4	10.1		35.8	
Fucus – Laminaria	96.7	21.0	5.4	10.1			38.3
Fucus – Saccharina	97.3	15.2	3.9	7.4			54.3
Ascophyllum – Himanthalia	99.9	54.9	11.7		19.7		
Ascophyllum – Saccorhiza	99.8	56.6	12.1			18.7	
Ascophyllum – Laminaria	99.9	56.4	12.1				20.8
Ascophyllum – Saccharina	99.9	47.3	10.1				33.3
Himanthalia – Saccorhiza	99.7				45.0	41.6	
Himanthalia – Laminaria	99.7				44.7		45.8
Himanthalia – Saccharina	99.8				31.3		61.6
Saccorhiza – Laminaria	98.3					44.2	49.1
Saccorhiza – Saccharina	98.5					30.4	64.6
Laminaria – Saccharina	31.4						

Table 2. Summary of pairwise SIMPER analyses between brown algal host samples grouped at genus level. Showing the OTUs that contributed to more than 10% of the observed dissimilarity between host genera according to SIMPER, when Phaeophyceae OTUs were excluded from the analyses.

		OTUs contributing >10 % to the dissimilarity between brown algal hosts (Excluding Phaeophyceae OTUs)						
	Phylum	Chlorophyta	Rhodophyta	Fungi			Stramenopiles	Unassigned
	Class	Ulvophyceae	Florideo-phycaea	Basidiomycota		Ascomycota	Oomycota	NA
	PR2 tax	Ulvophyceae sp.	*Ceramium rubrum*	Agarico-mycetes sp.	Exobasidio-mycetes sp.	*Torulaspora franciscae*	*Pythium chamaihyphon*	NA
SIMPER comparison	Dissi-milarity (%)	OTU 6	OTU 21	OTU 57	OTU 25	OTU 50	OTU 42	OTU 61
Fucus – Ascophyllum	95.9	33.2	15.7		11.9	4.2		
Fucus – Himanthalia	98.5	44.9						4.8
Fucus – Saccorhiza	94.0	32.6	5.1				5.8	
Fucus – Laminaria	98.5	43.0	3.2					
Fucus – Saccharina	98.7	41.3		12.2				
Ascophyllum – Himanthalia	99.4		41.3		31.1	10.5		4.4
Ascophyllum – Saccorhiza	83.3		13.8		15.9	5.4	7.7	
Ascophyllum – Laminaria	92.9		35.9		28.6	9.8		
Ascophyllum – Saccharina	98.6		36.4	12.8	27.0	9.2		
Himanthalia – Saccorhiza	98.7		13.7				14.2	4.2
Himanthalia – Laminaria	96.5		11.1		3.0			28.6
Himanthalia – Saccharina	98.1			58.2	1.2			17.5
Saccorhiza – Laminaria	91.9		10.4				13.3	
Saccorhiza – Saccharina	98.6		12.2	12.2			12.5	
Laminaria – Saccharina	97.7		7.4	47.2				

Table 3. Indicator species analysis (IndVal) showing OTUs most preferentially associated with the brown algal hosts.

OTU#	Associated with	p-value	Taxonomic assignment
OTU 149	*Ascophyllum*	0.0185 *	Stramenopiles; Ochrophyta; Phaeophyceae; *Silvetia*; *Silvetia siliquosa*
OTU 381	*Ascophyllum*	0.0175 *	Stramenopiles; Ochrophyta; Phaeophyceae; *Silvetia*; *Silvetia siliquosa*
OTU 244	*Ascophyllum*	0.0005 ***	Stramenopiles; Ochrophyta; Phaeophyceae; *Silvetia*; *Silvetia siliquosa*
OTU 2	*Ascophyllum*	0.0020 **	Stramenopiles; Ochrophyta; Phaeophyceae; *Silvetia*; *Silvetia siliquosa*
OTU 49	*Ascophyllum*	0.0127 *	Stramenopiles; Ochrophyta; Phaeophyceae; *Fucus*; *Fucus gardneri*
OTU 25	*Ascophyllum*	0.0446 *	Fungi; Basidiomycota; Ustilaginomycotina; Exobasidiomycetes; Exobasidiomycetes_X sp.
OTU 1	*Laminaria + Saccharina*	0.0033 **	Stramenopiles; Ochrophyta; Phaeophyceae; *Agarum*; *Agarum clathratum*

The Associated with column show which library/host species the OTUs were preferentially associated with. Abbreviations used; Ascophyllum = Ascophyllum nodosum, Laminaria = Laminaria digitata, Saccharina = Saccharina latissima. The asterisk associated with IndVal p-values indicat significance at ≤ 0.001 (***), ≤ 0.01 (**), ≤ 0.05 (*).

Table 4. Taxonomy of the shared OTUs between the three brown algal orders: Fucales, Laminariales and Tilopteridales, as shown in Venn plot (Fig. 10).

Overlap	OTU#	Taxonomic assignment		
Fucales + Laminariales + Tilopteridales (25 OTUs)	OTU 256	Alveolata	Ciliophora	Spirotrichea
	OTU 10, OTU 6	Archaeplastida	Chlorophyta	Ulvophyceae
	OTU 21	Archaeplastida	Rhodophyta	Florideophyceae
	OTU 25	Opisthokonta	Fungi	Basidiomycota
	OTU 94, OTU 233, OTU 234, OTU 68, OTU 112	Stramenopiles	Ochrophyta	Bacillariophyta
	OTU 1, OTU 236, OTU 28, OTU 292, OTU 41, OTU 49, OTU 146, OTU 70, OTU 4, OTU 7, OTU 9, OTU 3, OTU 2, OTU 348, OTU 5	Stramenopiles	Ochrophyta	Phaeophyceae
Fucales + Tilopteridales (22 OTUs)	OTU 111, OTU 148, OTU 137, OTU 188, OTU 151	Alveolata	Ciliophora	Oligohymenophorea, Phyllopharyngea, Spirotrichea
	OTU 298	Amoebozoa	Lobosa	*Vermistella* sp.
	OTU 180, OTU 87	Archaeplastida	Chlorophyta	Ulvophyceae
	OTU 110	Rhizaria	Cercozoa	Endomyxa
	OTU 265, OTU 197, OTU 220, OTU 72, OTU 63, OTU 237, OTU 17	Stramenopiles	Ochrophyta	Bacillariophyta
	OTU 300	Stramenopiles	Ochrophyta	Chrysophyceae
	OTU 105	Stramenopiles	Ochrophyta	Phaeophyceae
	OTU 38, OTU 42	Stramenopiles	Pseudofungi	Oomycota
	OTU 166, OTU 173	Unassigned	NA	NA
Fucales + Laminariales (20 OTUs)	OTU 89, OTU 97	Alveolata	Ciliophora	Oligohymenophorea, Phyllopharyngea
	OTU 107, OTU 191	Alveolata	Dinoflagellata	Dinophyceae
	OTU 69	Archaeplastida	Chlorophyta	Ulvophyceae
	OTU 226, OTU 118, OTU 257	Opisthokonta	Fungi	Ascomycota, Basidiomycota
	OTU 93, OTU 172	Stramenopiles	Ochrophyta	Bacillariophyta
	OTU 23, OTU 26, OTU 389, OTU 149, OTU 244, OTU 326, OTU 381	Stramenopiles	Ochrophyta	Phaeophyceae
	OTU 238	Stramenopiles	Opalozoa	MAST-3
	OTU 295	Stramenopiles	Pseudofungi	Oomycota
	OTU 249	Stramenopiles	Sagenista	Labyrinthulomycetes
Laminariales + Tilopteridales (3 OTUs)	OTU 193, OTU 74	Alveolata	Dinoflagellata	Dinophyceae
	OTU 168	Stramenopiles	Sagenista	Labyrinthulomycetes

Table 5. Hosts with brown algal endophytes selected for isolation from the B. Rinkel collection at the Natural History Museum, London.

Sample ID	Host	Site name, county	Date	Collector
BRE1sn1	*Chondrus crispus*	Sheerness, Kent	22.10.03	J. Brodie
BRE9ls2	*Chondrus crispus*	Lilstock, Somerset	27.10.03	B. Rinkel
BRE38sm11	*Mastocarpus stellatus*	Sidmouth, Devon	28.10.01	B. Rinkel
BRE74ls1	*Fucus serratus*	Lilstock, Somerset	27.10.03	B. Rinkel
BRE98sm12	*Mastocarpus stellatus*	Sidmouth, Devon	28.10.01	B. Rinkel
BRE117sm22	*Osmundea osmunda*	Sidmouth, Devon	28.10.01	B. Rinkel
BRE118sm22	*Osmundea osmunda*	Sidmouth, Devon	28.10.01	B. Rinkel
BRE128hi1	*Osmundea pinnatifida*	Hayling Island, Hampshire	05.02.04	B. Rinkel
BRE152cb2	*Chondrus crispus*	Castle Beach, Pembrokeshire	09.03.04	B. Rinkel
BRE157br1	*Mastocarpus stellatus*	Black Rock, Pembrokeshire	10.03.04	B. Rinkel
E026pz	*Chondrus crispus*	Polzeath, Cornwall	19.06.04	M. Holmes
E030Nb	*Mastocarpus stellatus*	Norway	23.06.04	J. Brodie
E326wr	*Fucus serratus*	West Runton, Norfolk	28.09.04	B. Rinkel
E360sm2	*Chorda filum*	Sidmouth, Devon	28.10.03	B. Rinkel
E361ac	*Chorda filum*	A'Chleit, Bute	10.07.04	B. Rinkel
E365sg	*Mastocarpus stellatus*	Seagreens, Aberdeenshire	16.07.04	B. Rinkel
E368	*Chondrus crispus*	Bouley Bay, Jersey	13.03.05	B. Rinkel
E370	*Chondrus crispus*	Bouley Bay, Jersey	13.03.05	B. Rinkel
E377	*Chondrus crispus*	Greve de Lacq, Jersey	14.03.05	B. Rinkel
LTE19/1	*Osmundea pinnatifida*	Black Rock, Pembrokeshire	10.03.04	B. Rinkel

A. (With Phaeophyceae OTUs)

B. (Without Phaeophyceae OTUs)

- Fucales
- Laminariales
- Tilopteridales