A novel approach to the functional classification of retinal ganglion cells

Abbreviated title: DREADD for RGC classification

Gerrit Hilgen[1, 2 *], Evgenia Kartsaki[1, 3], Viktoriia Kartysh[1,4,5], Bruno Cessac[3], Evelyne Sernagor[1*]

[1] Biosciences Institute, Newcastle University, Newcastle upon Tyne, UK

[2] Health & Life Sciences, Applied Sciences, Northumbria University, Newcastle upon Tyne UK

[3] Université Côte d'Azur, Inria, Biovision team and Neuromod Institute, Sophia Antipolis Cedex, F

[4] Ludwig Boltzmann Institute for Rare and Undiagnosed Diseases (LBI-RUD), 1090 Vienna, AT

[5] Research Centre for Molecular Medicine (CeMM) of the Austrian Academy of Sciences, 1090 Vienna, AT

* Corresponding author: gerrit.hilgen@ncl.ac.uk; evelyne.sernagor@ncl.ac.uk

Financial interests or conflicts of interest: The authors declare no competing financial interests

Abstract

Retinal neurones come in remarkable diversity based on structure, function and genetic identity. Classifying these cells is a challenging task, requiring multimodal methodology. Here, we introduce a novel approach for retinal ganglion cell (RGC) classification, based on pharmacogenetics combined with immunohistochemistry and large-scale retinal electrophysiology. Our novel strategy allows grouping of cells sharing gene expression and understanding how these cell classes respond to basic and complex visual scenes. Our approach consists of increasing the firing level of RGCs co-expressing a certain gene (*Scnn1a* or *Grik4*) using excitatory DREADDs (Designer Receptors Exclusively Activated by Designer Drugs) and then correlate the location of these cells with *post hoc* immunostaining, to unequivocally characterise anatomical and functional features of these two groups. We grouped these isolated RGC responses into multiple clusters based on the similarity of their spike trains. With our approach, combined with immunohistochemistry, we were able to extend the pre-existing list of Grik4 expressing RGC types to a total of 8 and, for the first time, we provide a phenotypical description of 14 Scnn1a-expressing RGCs. The insights and methods gained here can guide not only RGC classification but neuronal classification challenges in other brain regions as well.

Keywords

DREADD, Grik4, Scnn1a, retinal ganglion cells, multielectrode array, classification

Introduction

The retina contains two types of photoreceptors, rods for dim light and cones for daylight and colour vision. Furthermore, cone-contacting bipolar cells can be divided into ON and OFF types and further subdivided into more than a dozen different subpopulations (1). These parallel processed channels are further divided into a variety of functional output channels, so-called retinal ganglion cells (RGCs), which encode different features of the visual

environment. There are ~1 million RGCs in the human retina and ~45,000 in the mouse retina

(2,3), integrating visual information processed from photoreceptors down the retinal neural

network. Different types of RGCs extract very specific features from the visual scenery (4).

This code is transmitted to postsynaptic targets in the brain, leading to visual perception. At

present, more than 40 RGC types have been identified in the mouse retina (5,6). RGC

classification is typically based on common anatomical features (7,8), responses to light (5,9–

11) or on shared gene expression (6,12–14). Classification based on gene expression is

relatively recent, and the majority of RGC groups sharing specific genes have not been

phenotyped yet.

Current approaches for functional characterization of RGC subpopulations at pan-retinal scale

are limited. Multielectrode arrays (MEAs) allow high throughput simultaneous electrical

recording from the RGC population at high spatiotemporal resolution (15). Here we use a

CMOS (complementary metal-oxide-semiconductor) MEA system consisting of 4,096

electrodes (2.67 x 2.67 mm), allowing us to record light responses from hundreds to thousands

of RGCs simultaneously at pan-retinal level and near cellular resolution (16,17). We selected

two genes based on Cre lines availability, on the sparse expression distribution across the

RGC layer and on their novelty for phenotypic characterization (Allen Mouse Brain

Connectivity Atlas (2011)). *Grik4* (glutamate receptor, ionotropic, kainate subunit 4, HGNC:

4582) expressing RGCs have already been partially described using a *Grik4* Cre mouse line

(18). The other gene we investigated is *Scnn1a* (non-voltage gated sodium channel, epithelial

1 subunit alpha, HGNC:10599). Scnn1a Cre-induced recombination (*Scnn1a*-Tg3-Cre line) is

present in sparse Layer 4 neurones, mostly in the somatosensory cortex (19). Current

knowledge of Scnn1a expressing RGCs in the retina is limited to the fact that their dendritic

arbour stratifies in sublaminas S1 and S2 (OFF layers) and in sublamina S4 (ON layer) of the

inner plexiform layer (13). Here, we crossbred Grik4 and Scnn1a Cre mice with DREADD flox

mice to further investigate these RGC types.

Designer Receptors Exclusively Activated by Designer Drugs (DREADDs) (20) technology is

a powerful new approach to pharmacologically dissect out the role of specific neuronal cell

classes in network activity (21,22). DREADDs are an engineered version of G-proteins that

allow precise control of G-protein signalling pathways and can be activated with clozapine-N-

oxide (CNO), a "designer drug" with no endogenous receptors in the organism. Most

commonly used DREADDs are excitatory (hM3Dq, triggering release of calcium from

organelles leading to increase of intracellular concentration of free calcium and to membrane

depolarization). In this study, we have generated Cre recombinase-mediated restricted

expression of cell-specific DREADD (23) expression in either *Grik4* or *Scnn1a* reporter lines.

We have successfully isolated light-evoked responses in RGCs sharing either Scnn1a or Grik4

gene expression by combining excitatory DREADD activation, large-scale retinal CMOS MEA

recording and *post hoc* labelling of DREADD-expressing RGCs. We grouped the RGC

responses into multiple clusters based on the similarity of the spike trains they generate in

response to a series of stationary stimuli. Our method was validated by successfully isolating

existing *Grik4*-RGC types, but it also added several new candidates to the group. It provided

novel classification for 14 subgroups of *Scnn1a*-RGCs.

72 Results

To functionally validate RGC subgroups according to shared gene expression, we first

established an immunohistochemical atlas of these cells. Building such a resource for Grik4

and Scnn1a expressing cells in the mouse retina is important for estimation of the RGC

numbers and types in these two genetic pools. We used the intrinsic fluorescence signal of

Grik4-DREADD (hereafter named Grik4) and Scnn1a-DREADD (hereafter named Scnn1a)

cells to provide a detailed IHC map of Grik4 and Scnn1a expressing cells in retinal whole

mounts and vertical sections. Each DREADD is tagged with hemagglutinin (HA) as well as

mCitrine, allowing to visualize DREADD-expressing cells by immunofluorescence. We first

investigated the distribution of Grik4 and Scnn1a cells in the ganglion cell layer (GCL) in retinal

Figure 1: Grik4 and Scnn1a cells in the GCL are not homogeneously distributed. Whole mount antibody staining against Grik4 (A) and Scnn1a (B) DREADD GFP were imaged at the level of the GCL. All stained GFP cells were counted and the densities were calculated and presented in pseudocolors for 3 Grik4 (C) and Scnn1a (D) retinas. V= ventral, T = temporal, D = dorsal, N = nasal. Scale bar A, B = 1 mm; Scale bar A, B insets II = 100 μm.

whole mounts using an antibody against Green Fluorescent Protein (GFP) to amplify the

intrinsic mCitrine signal (Fig 1). Both lines exhibit sparse cellular distribution in the GCL, with

Scnn1a cells significantly more abundant than Grik4 (Fig 1 A, B insets I and II). We calculated

the cell densities in the GCL of three representative Grik4 (Fig 1 C) and Scnn1a (Fig 1 D)

retinas. Grik4 and Scnn1a cell densities respectively vary between 10-25 and 40-100

cells/mm^2. Both pools exhibit non-even distributions. Grik4 cells are more prominent in the

dorsal-temporal periphery (Fig 1 C, yellow areas), while Scnn1a cells are more prominent in

the ventral-nasal periphery (Fig 1 D, yellow areas).

The GCL consists mainly of RGCs but it also contains displaced amacrine cells (dACs) and

glial cells. We investigated whether dACs contribute to the pool of Grik4 and Scnn1a cells in

the GCL. We used known specific RGC and amacrine cell (AC) markers to estimate the

contribution of different RGC types in both gene pools. First, we double-labelled Grik4 and

Scnn1a retinal vertical sections against GFP (Fig 2 A, B cyan) and RBPMS (Fig 2 A, B

magenta), a selective marker for RGCs in the mammalian retina (24). More than 50% of the

GFP labelled Grik4 cells in the GCL are RBPMS positive (Fig 2 A, arrows) whereas the other

Figure 2: Grik4 and Scnn1a DREADD are expressed in RGCs and ACs. Vertical sections were stained for Grik4 (A, C, cyan) and Scnn1a (B, D, cyan) GFP. Sections were further co-labelled with RBPMS (A, B, magenta), a marker for RGCs, and GABA (C, D, magenta). INL = inner nuclear layer, IPL = inner plexiform layer, GCL = ganglion cell layer. Scale bar in D = 20 μm.

cells are likely dACs. In support, the somata of putative dACs in the GCL are relatively small,

which is a key feature of these cells. A similar pattern was found for Scnn1a cells in the GCL

but here, the majority of Scnn1a cells are RGCs (Fig 2 B, arrows). Sparsely distributed cells

expressing either Grik4 (Fig 2 C, cyan) or Scnn1a (Fig 2 D, cyan) were present in the proximal

inner nuclear layer (INL). These cells did not stain for RBPMS (Fig 2 A, B), confirming they

are amacrine cells (ACs). Most AC types are GABAergic, hence we initially stained for GABA

(Fig 2 C, D, magenta) and GFP in Grik4 and Scnn1a vertical sections. Interestingly, we did

not find any GABA-positive Grik4 ACs (Fig 2 C). On the other hand, we occasionally found

GABAergic Scnn1a cells in the INL or GCL (Fig 2 D, arrow). In conclusion, a major proportion

of Grik4 and Scnn1a cells in the GCL are RGCs but both genes are also sparsely expressed

in ACs in the INL and in dACs in the GCL. Further, GABA is not present in Grik4 cells, but it

is in a very small population of Scnn1a expressing cells.

To further characterize the expression patterns of Scnn1a and Grik4 in RGCs and ACs, retinal

sections were stained for GFP, RBPMS and for the well-described calcium-binding protein

markers Parvalbumin and Calretinin (Fig 3) (25,26). Briefly, Parvalbumin is found in ACs and

Figure 3: Grik4 and Scnn1a DREADD are expressed in multiple RGC types. Vertical sections of Grik4 and Scnn1a retinas were triple stained A, B) for GFP (cyan), RBPMS (magenta) and Calretinin (yellow) or C, D) GFP (cyan), Parvalbumin (magenta) and Calretinin (yellow). INL = inner nuclear layer, IPL = inner plexiform layer, GCL = ganglion cell layer. Scale bar in D = 20 µm.

in 8-14 different RGC types. Calretinin is also present in ACs and in 10 RGC types. In a first

set of experiments (Fig 3 A, B) we labelled respectively against GFP (cyan), RBPMS

(magenta) and Calretinin (yellow) in vertical sections of Grik4 (Fig 3 A) and Scnn1a (Fig 3 B)

retinas. For both retinas, we found a subset of Grik4 and Scnn1a RGCs that were

GFP/RBPMS/Calretinin positive (Fig 3 A, B arrows) and GFP/RBPMS positive cells, which

clearly demonstrates that there are at least two RGC types for each pool. We did not find any

Calretinin positive Grik4 ACs, but we did find some Calretinin positive Scnn1a ACs. For

technical reasons we were not able to combine RBPMS and Parvalbumin staining (both

antibodies were raised in the same species). To extend on these findings and to further

investigate if the genetic pools of Grik4 and Scnn1a consist of more than two RGC types, we

triple-labelled vertical sections (Fig 3 C, D) for GFP (cyan), Calretinin (yellow) and Parvalbumin

(magenta). Our initial findings indicate that the majority of GFP cells in the GCL are RGCs.

These experiments described below thus aimed to find out whether these GFP cells can be

subdivided into smaller groups. We found that the population of Grik4 (Fig 3 C) expressing

cells (cyan) in the GCL consists of at least three different types of cells: only GFP (Fig 3 C,

cyan arrow), GFP/Calretinin (yellow arrow) and GFP/Calretinin/Parvalbumin (white arrows)

cells. It is likely that these three different cell types reflect RGCs and not dACs because we

did not find any putative Grik4 ACs in the INL expressing Calretinin and/or Parvalbumin.

For the pool of Scnn1a expressing cells in the GCL (Fig 3 D), we found four different types:

those expressing only GFP (Fig 3 D, cyan arrow), GFP/Calretinin (yellow arrow),

GFP/Parvalbumin (magenta arrow) and GFP/Calretinin/Parvalbumin (white arrow) cells. A

fraction of the GFP positive ACs in the INL were positive for Calretinin, suggesting that some

P/Calretinin and GFP/Calretinin/Parvalbumin cells in the GCL are potentially dACs. In

summary, the pool of Grik4 and Scnn1a cells in the GCL respectively consists of at least three

and four different RGC types. That information is important for the validation of our novel

approach to the functional classification of these same RGCs.

We next identified Grik4 and Scnn1a RGCs according to the change of their spiking pattern

following DREADD activation with CNO, and we grouped them into functional clusters based

on response similarity. The IHC experiments revealed that DREADDs are expressed in

several RGC types but also in ACs (albeit more sparsely). Hence, the first step to isolate Grik4

and Scnn1a expressing RGCs is to pre-identify cells that showed either a 50% (arbitrarily

chosen) increase in spontaneous firing rate (Fig 4 A) or a 50% increase of the Burst Index (Fig

4 B) in the presence of CNO. That step is followed by correlating the physical position of these

identified RGCs with micrographs of DREADD-GFP expressing cells in the GCL. This step

was necessary because CNO also evoked activity in ACs which in turn altered spiking

activities in non-DREADD expressing RGCs (see also Fig 6). Broadband blocking of these

unknown DREADD ACs with pharmacological agents resulted in up- and down regulated

spiking

Figure 4: DREADD activation leads to spike pattern changes in RGCs. DREADDs in RGCs can be activated with Clozapine N-oxide (CNO) and lead to an increase in firing frequency (A, right half) and sometimes also in bursting activity (B, right half). Similar activities can be evoked with an "allround" AC neurotransmitter blocker (C) without CNO. The cocktail contains: 2 µM Strychnine (glycine receptor antagonist), 20 µM Mecamylamine (nicotinic acetylcholine receptor antagonist), 20 µM Bicuculline (GABA$_A$ antagonist), 5 µM CGP 55845 ((2S)-3-[[(1S) -1- (3,4-Dichlorophenyl)ethyl] amino -2- hydroxypropyl] (phenylmethyl) phosphinic acid hydrochloride, GABA$_B$ antagonist), 50 µM TPMPA (1,2,5,6-Tetrahydropyridin-4-yl methylphosphinic acid, GABA$_C$ antagonist). Plotted in A, B are 20 Grik4 RGCs that showed a minimum of 50% change in firing frequency or bursting activity and in C 100 randomly selected

activities in almost all RGCS (Fig 4 C), hence it is not a suitable approach to isolate the

DREADD RGCs. The process of correlating physical positions with structural imaging allowed

us to unequivocally correlate anatomical and functional features of Grik4 and Scnn1a RGCs.

Here, we provide a brief exemplary description, using an identified Grik4 RGC (Fig 5), to show

how we register GFP labelled cells with their spiking activity for unambiguous identification of

Grik4 RGCs. Further technical details are available in the Method section. The spike

localization cluster centre (magenta circle) is calculated for a pre-identified (see above) Grik4

RGC (Fig 5 B) using the spike interpolation algorithm in Herdingspikes2. In addition, electrical

imaging confirmed the maximal current sink position (Fig 5 C, orange diamond) and visualized

Figure 5: Registering GFP RGCs with nearby isolated spike centre clusters. The spike cluster centre (B, circle) and the electrical imaging (C, diamond) define the potential electrical source location of the isolated RGCs. Spike centres in close proximity to GFP labelled RGCs (D) were registered as potential DREADD-expressing RGCs (E, F, Match). Scale bars A, C, F = 40 µm; B = 80 µm; E = 20 µm.

putative axonal trajectories (Fig 5 C, arrows), thus helping to align the GFP cell structure with

the electrical activity more precisely. Image segmentation techniques are used to threshold

the GFP foreground from the background and a blob detection is applied to detect the outlines

and centre of Grik4 and Scnn1a GFP positive cells in the GCL (Fig 5 D, green outline). Lastly,

the x and y positions are converted to match with MEA images dimensions (Fig 5 E). Spikes

originate from the axon initial segment (AIS) rather than from the cell body. Therefore, we

expect the spike cluster centre to be slightly eccentric with respect to the soma itself. The

exact location of the AIS in RGCs can be very close to the soma, or sometimes rather distant

along the axon (27,28). All the information from panels B-D is needed to estimate if a GFP-

expressing cell corresponds to a nearby spike cluster centre. Two virtual circles with a radius

of 30 and 60 µM (arbitrary values) are drawn around a detected GFP blob centre (Fig 5 E),

and spike cluster centres falling within one of these two zones are registered. From

experience, we reached the conclusion that using the 60 µm zone was sufficient to reliably

determine whether an isolated spike cluster corresponds to a specific GFP-expressing cell.

Such cases are referred to as "Match", and those falling outside these boundaries are

classified as "No Match" (Fig 5 F, Fig 6 A, D). All other recorded spike units, that did not exhibit

changes in spike behaviour in the presence of CNO, were labelled "RGC unit" (Fig 6 A, D) but

not considered for further classification.

We successfully analysed 3 Grik4 and 5 Scnn1a retinas to register "Match" (Fig 6 A, D,

magenta) and "No Match'' (Fig 6 A, D, white) cells. Figure 6 A, D are representative examples

from the cohort of all experiments. Note that the recording area is generally not completely

covering the entire 64 x 64 array. In total, we identified 117 "Match" and 863 "No Match" RGC

in the 3 Grik4 retinas (Fig 6 C) and 307 "Match" and 1066 "No Match" in the 5 Scnn1a retinas

(Fig 6 F). We divided these RGCs according to their response preference for stationary (static,

full field) and non-stationary images (moving bars for direction of orientation selectivity)).

Briefly, cells that were exceeding a threshold value (see Methods) for their direction or

orientation selectivity index (DSI and OSI, respectively) were classified as non-stationary (Fig

6 B, E). Figure 6 is showing an example for DSI grouping but we also grouped cells according

to their OSI. In the following, we focus on "Match" Grik4 and Scnn1a RGCs only. A total of 78

Grik4 stationary and 41 non-stationary RGCs were collected (Fig 6 C). For Scnn1a, 180

stationary and 127 non-stationary RGCs were collected (Fig 6 F). However, most RGCs did

exhibit changes in firing rate (increase or decrease) without revealing any GFP signals, hence

they are unlikely to be Grik4 or Scnn1a RGCs (here called "No Match"). These cells are most

likely other RGC types affected by either Grik4 or Scnn1a expressing ACs, and they

outnumber the "Match" RGCs. In summary, we successfully combined near pan-retinal

Figure 6: Dividing registered Grik4 and Scnn1a RGCs into stationary and non-stationary responders. Moving stimuli were used to estimate the direction selectivity index (DSI) of registered "Match" RGCs (Grik4 A, Scnn1a D, magenta). The DSI was plotted against the probability density (histogram) and fitted with two gaussians (B, E, red line). RGCs exceeding a set threshold (B, E, dotted red line) were labelled "non-Stationary". The registered RGCs from all retinas were grouped into "non-Stationary" and "stationary" (C, F).

recordings with *post hoc* anatomical characterization and isolated GFP-positive Grik4 and

Scnn1a RGCs with increased spiking or bursting activity in the presence of CNO.

After the successful registration of spikes with Grik4/Scnn1a GFP RGCs, the final step is to

classify these cells into functional groups according to the nature of their responses to light.

We recently described a non-parametric approach for RGC classification by using the SPIKE

distance (29,30) as a clustering metric (10). In order to use that approach, it is necessary to

have a stimulus that elicits responses simultaneously over the entire recording area. Here we

used a chirp stimulus inspired from Baden et al (2016) (5) that elicits responses from all RGCs

at the same time to pre-sort stationary and non-stationary RGCs (chirp, Fig 7 B, F, top) for

Figure 7: Clustering of Grik4 stationary and non-stationary RGC responses. RGC responses that showed a high spike train similarity for a chirp stimulus (B, F) were grouped together using gap statistics (A). For each RGC of the groups, the PSTH was calculated and the mean PSTH was plotted (B, F, colored lines). The means for Bias Index (C, D, G), Response Duration (C), Baseline Spiking (D), Burst change (E), Spike Rate change (E) and orientation selectivity index (OSI, G) index were scatter plotted and their standard deviations were used as the circle diameters.

SPIKE distance measure and hierarchical agglomerative clustering. We manually validated

our detected clusters by grouping several response parameters, e.g. bias index (ON, ON-OFF

or OFF) or response duration (transient or sustained), from the chirp and moving bars (for

example Fig 7 C-E, G). Manual grouping was only feasible because the expected cluster

numbers for Grik4 and Scnn1a were small (respectively 3 and 4 according to our IHC

analyses). The pairwise SPIKE distances were determined from all trials of the chirp stimulus

and the resulting distance matrix of stationary and non-stationary RGCs was clustered with a

hierarchical clustering algorithm followed by the construction of a dendrogram as shown in

Jouty et al., 2018 (10). To find the optimal number of clusters for stationary and non-stationary

SPIKE distances we used gap statistics (10,31). For stationary and non-stationary RGCs, gap

statistics estimated four response clusters of Grik4 RGCs (Fig 7 A). Additional response

parameters were extracted from the chirp peri-stimulus time histogram (PSTH) e.g., bias index

or response duration of these 4 stationary RGC types. We found potentially ON transient (#1,

n = 30), ON sustained (#2, n = 13), ON-OFF (#3, n = 16) and OFF sustained (#4, n = 10)

response types. Some RGC units were discarded during the process of SPIKE distance

grouping as they showed a low signal-to-noise ratio i.e., high response variability (10). To

validate these four detected clusters, we plotted the mean bias index against the mean

response duration from all RGCs in a cluster (Fig 7 C), revealing four distinct groups that

corresponded to the confirmed stationary RGC types. Moreover, spontaneous activity follows

very specific patterns for certain RGC types and can therefore help classification. Plotting the

mean Bias Index against mean spontaneous firing rate (baseline firing) revealed that the ON

sustained cluster #2 (Fig 7 B, C, cluster 2) exhibited a very strong baseline firing rate (Fig 7

D). The same plot also shows that the three other RGC types have a moderate baseline firing

rate and further confirmed the 4 well separated RGC types. Whether the spontaneous firing

rate becomes burstier or simply increases monotonously in the presence of CNO can be used

to further group RGCs into clusters (Fig 7 E). Interestingly, the effect of CNO on cluster #2 is

minimal (Fig 7 E) whereas it is maximal for cluster #3 (Fig 7 E). The non-stationary RGCs

barely responded to the chirp stimulus (Fig 7 F, scale bar) but that response was, albeit

modest, reliable and unique for the different clusters. We used the orientation (OS) and

direction selectivity (DS) index in addition to the chirp PSTH parameters to define the four non-

stationary Grik4 clusters (Fig 7 F, G). We found ON - OS (#5, n = 13), ON-OFF - DS (#6, n =

13; #8, n = 7) and OFF – OS (#7, n = 6) response groups in the pool of non-stationary Grik4

RGCs. The mean DS index was almost identical for these 4 groups; hence we used the OS

index for manual validation. While the ON – OS (#5) and the OFF – OS (#7) are clearly

distinguishable from each other, the difference is less pronounced between the two ON-OFF

– DS clusters (#6 & #8). They have a similar mean BI, but Cluster 6 tends to be marginally

more OS (Fig 7 G). Clusters 6 and 8 would therefore need further evaluation to determine

whether they are truly different RGC types. In summary (Table 1), we found 4 stationary RGCs

types and most likely 3, non-stationary RGC types that share the Grik4 gene pool.

Grik 4 Cluster	Grik 4 stationary	Grik 4 Cluster	Grik4 non-stationary
#1	G20 ON high freq after Baden et al., 2016	#5	G17 ON local trans OS after Baden et al., 2016
#2	G22 ON sustained after Baden et al., 2016; Putative PixON after Johnson et al., 2018	#6	G12 ON-OFF DS1 after Baden et al., 2016; Vaney, Sivyer & Taylor, 2012; Temporal, dorsal cluster 10, 16, 24 after Tran et al., 2019
#3	G11 ON-OFF local after Baden et al., 2016; Putative HD1/2 after Jacoby & Schwartz 2017	#7	G4 OFF slow after Baden et al., 2016; Putative F-midi OFF after Rousso et al., 2017; Cluster 28 after Tran et al., 2019
#4	G5 OFF alpha sustained after Baden et al., 2016; Krieger et al., 2017 Cluster 42 after Tran et al., 2019	#8 (#5?)	G12 ON-OFF DS1 after Baden et al., 2016; Vaney, Sivyer & Taylor, 2012

Table 1: Summary of identified Grik4 clusters and potential matches with existing literature. NB these are suggestions.

We proceeded in a similar way with stationary and non-stationary registered Scnn1a RGCs

(Fig 8). Gap statistics suggested 7 clusters for both groups (Fig 8 A). The dendrogram of the

chirp PSTH consists of ON transient (#1, n = 30; #2, n = 27; #3, n = 47), ON sustained (#6, n

= 16; #7, n = 12), ON-OFF (#5, n = 15) and OFF (#4, n = 28) like response for Scnn1a RGC

types (Fig 8 B). Further analysis revealed that the ON sustained (#7), OFF (#4) and ON-OFF

(#5) response parameters (Fig 8 C - Bias Index vs Response Duration; Fig 8 D – Bias Index

vs Baseline) were clearly distinct from each other. At the same time, the distinction is less

Figure 8: Clustering of Scnn1a stationary and non-stationary RGC responses. RGC responses that showed a high spike train similarity for a chirp stimulus (B, F) were grouped together using gap statistics (A). For each RGC of the groups, the PSTH was calculated and the mean PSTH was plotted (B, F, colored lines). The means for Bias Index (C, D, G), Response Duration (C), Baseline Spiking (D), Burst change (E), Spike Rate change (E) and orientation selectivity index (OSI, G) index were scatter plotted and their standard deviations were used as the circle diameters.

non-stationary Scnn1a RGCs (Fig 8 F) clustered into OFF-DS (#11, n = 13), ON-OFF– DS

(#12, n = 28; #14, n = 2), ON-OFF – OS (#13, n = 13), ON – DS (#9, n = 8; #10, n = 12) and

ON - OS (#8, n = 29) like responses. Although there was some overlap between certain

clusters (#9, #10, #11, #14) when plotting the Bias Index and OS index means (Fig 8 G), their

chirp PSTH plots were substantially different, suggesting that these cells do belong to distinct

functional groups. Therefore, it is likely that the Scnn1a RGC pool consists of several DS and

OS cells which will necessitate further investigation to establish their basic functional

differences. In summary (Table 2), we found a minimum of four stationary and a maximum of

seven non-stationary Scnn1a RGC types.

Scnn1a Cluster	Scnn1a stationary	Scnn1a Cluster	Scnn1a non-stationary
#1	G18 ON trans after Baden et al., 2016	#8	G17 ON trans (OS) after Baden et al., 2016
#2	G18 ON trans after Baden et al., 2016	#9	G26 ON DS sustained 2 after Baden et al., 2016
#3	G18 ON trans after Baden et al., 2016	#10	?
#4	G8 OFF alpha trans after Baden et al., 2016; Krieger et al., 2017; Cluster 45 after Tran et al., 2019	#11	G2 OFF DS after Baden et al., 2016; Putative F-mini OFF after Rousso et al., 2017; Cluster 4 after Tran et al., 2019
#5	G10 W3 local edge detector after Baden et al., 2016; Zhang et al., 2012; Cluster 6 after Tran et al., 2019	#12	G12 ON-OFF DS1 after Baden et al., 2016; Nasal, ventral cluster 12, 16 after Tran et al., 2019
#6	G18 ON trans after Baden et al., 2016	#13	G14 ON-OFF local OS after Baden et al., 2016
#7	G22 ON sustained after Baden et al., 2016; Putative PixON after Johnson et al., 2018	#14	?

Table 2: Summary of identified Scnn1a clusters and potential matches with existing literature. NB these are suggestions.

Discussion

We established a novel approach for RGC classification, based on grouping of cells sharing

gene expression and similarities in their light response patterns. A previous study used

pharmacogenetics in combination with MEA recordings for Parvalbumin RGC classification

(33). PV expressing cells in the retina are manifold with at least 8 RGC types (26) and a distinct

number of amacrine cells. Other approaches combined MEA recordings with anatomical

imaging (34,35). We successfully used a combination of all three methods, excitatory

DREADD activation in retinal cells, immunohistochemistry and large-scale retinal

electrophysiology to provide a significantly more comprehensive pan-retinal phenotypical

description of Grik4- and Scnn1a-expressing RGCs in the mouse retina. Indeed, we extended

the pre-existing list of Grik4 expressing RGC types, which validates our approach, and

indicates that it has more powerful detection potential than previous ones. For the first time,

we also provide a functional description of Scnn1a-expressing RGCs. We also showed that

certain types of ACs express Grik4 and Scnn1a and that exciting these cells with DREADDs

activation leads to spike firing change in many undefined RGCs. Our approach is not restricted

to retinal cells but is widely applicable to other neurons from other brain regions. It is a scalable

multimodal approach and can provide fast grouping of large cohorts of neurons with similar

gene expression.

Grik4 expression in the retina has been described (13,35–38) but there is functional

characterization only for two RGC types (37,38). Johnson et al., (38) reported that the Grik4-

expressing PixON RGC type is an ON sustained type and has strong spontaneous activity.

Our Grik4 cluster #2 from the stationary RGCs (Tab 1) has the same key phenotypic features

- sustained responses and strong spontaneous activity. Cluster #2 also matches with the ON

sustained group (G) 22 from Baden et al. 2016, (5). Grik4 clusters #1, #3 and #4 potentially

resemble G20 (ON high freq), G11 (ON-OFF local) and G5 (OFF alpha sustained) from that

study. Cluster #3 is potentially also resembling HD1/2 after Jacoby & Schwartz 2017 (39) and

the #4 OFF alpha sustained is further described in Krieger et al. 2017, (40) The latter one is

parvalbumin positive (5,14,40) and in line with our IHC results that revealed Grik4 and

parvalbumin expressing RGCs. Baden et al. (2016) found many more parvalbumin-expressing

RGC types rather than the traditionally known 8 types (26). We suspect that there is some

overlap between Grik4 and yet undescribed parvalbumin, potentially also calretinin, RGC

types. We currently investigate the functional features of parvalbumin and calretinin co-

expressing RGCs using a similar methodology (41). Further, the existence of Grik4-expressing

ON-OFF – DS RGCs has been demonstrated in the mouse retina [22]. Cluster #6 of our non-

stationary RGCs also show ON-OFF DS RGCs (42) and potentially resemble G12 (ON-OFF

DS1) from Baden et al. (2016) and the dorsal-temporal clusters 10, 16 and 24 after Tran et

al., 2019 (6). Grik4 cluster #8 is also G12 ON-OFF DS though the responses are less

pronounced than in cluster #6. Our non-stationary Grik4 cluster #5 has similarities with G17

(ON local trans - OS) and cluster #7 with G4 (OFF slow) from Baden et al. (2016). The latter

one also shows similarities to the F-midi OFF cell responses described in Roussou et al. 2017,

(43) and Tran et al. 2019, (6). It would also explain the findings of parvalbumin and calretinin

positive Grik4 RGCs in our experiments (Fig 3 C) since F-midi OFF cells are supposed to

express both (43).

For Scnn1a, we found seven stationary and seven non-stationary RGC response groups (Tab

2). Interestingly, the Scnn1a ON sustained Cluster #7 showed remarkable similarity with the

PixON characteristics described earlier - sustained responses and high baseline firing. There

are no reports about Grik4 and Scnn1a co-expressing cells but it is not uncommon that certain

RGC types share the same genes e.g. *Foxp* and *Brn3 (43)* or *Pvalb* and *Calb2 (41,43,44)*.

Our Scnn1a stationary clusters #1, #2, #3 and #6 were grouped by gap statistics into 4 different

groups but our post-hoc analysis (Fig 8 C, D) suggests one group. Therefore, these four

clusters may reflect just one, or maybe two functional groups. They all exhibit ON transient

responses but are not able to follow high frequencies in the chirp (like cluster #1 in Grik4),

thus they resemble G18 (ON trans) in Baden et al. (2016). Cluster #4 has OFF transient

response characteristics and could resemble G8 (OFF alpha trans, see also (40)) after Baden

et al., 2016 and cluster 45 after Tran et al. 2019, (6). The Scnn1a stationary cluster #5 shows

weak ON and OFF responses to chirp and could potentially be the W3 local edge detector

(G10 in Baden et al. 2016) (45). The grouping of the non-stationary Scnn1a clusters #8 - #14

shows a variety of response types. Scnn1a cluster #8 shows strong ON transient and slight

OS responses and may relate to G17 (ON trans OS) in Baden et al. (2016). Scnn1a cluster

#9 shows very weak and sluggish ON DS responses and is not really following the oscillating

components in the chirp. It shows similarities to G26 (ON DS sust 2) from Baden et al. (2016).

Our Scnn1a OFF DS cluster #11 is most likely G2 (OFF DS) in Baden et al. (2016) and/or

potentially the F-mini OFF described in Roussou et al. (2016) and Tran et al., 2019. The F-

mini-OFF is calretinin (occasionally parvalbumin as well) positive which would fit with our IHC

results. The non-stationary Scnn1a ON-OFF DS cluster #12 may relate to G12 (ON-OFF DS

1) and nasal-ventral clusters 12 &16 after Tran et al., 2019. Cluster #13 is potentially G14

(ON-OFF local OS) from Baden et al. 2016. Clusters #10 and #14 show very weak chirp

responses and it is difficult to assign them to any existing response type. In summary, our

approach is able to group individually MEA recorded RGC response types into established

response clusters (Baden et al. (2016 and Tran et al., 2019)) obtained with Ca^{2+} imaging or

genetically, respectively.

DREADDs were also expressed in one or several yet undescribed AC types. For Scnn1a, we

found few GABAergic ACs but most GFP-positive ACs were not characterized further. For

Grik4, we did not find any GABAergic ACs, nor were we able to define the AC type with our

experimental means. Excitatory activation of Grik4 and Scnn1a ACs does have a profound

effect on postsynaptic RGC activity. A single AC affects nearly every RGC within reach of its

dendrites (46). Briefly, assuming that CNO acts simultaneously on some ACs and RGCs, we

are confronted with the problem of untangling the underlying circuits. For example, let us

consider a simple network with one population of DREADD ACs which modulate one

population of bipolar cells and one population of DREADD-RGCs. Adding CNO will have a

direct effect on RGCs, increasing their activity (firing and/or bursting rate). At the same time,

ACs activity increases, thereby reinforcing lateral inhibition. This situation is already complex

enough if we consider non linearities in the cell's responses. But as those cells are embedded

in a network, the net effect of CNO can have many forms leading to a priori undetermined

situations. For example, it is known that cascade of inhibition can result in excitation (e.g.

push-pull effect) (47). A possible solution to this problem relies on a quantitative analysis that

requires considering the factors constraining individual cell responses, without and with CNO,

and the network connectivity. It is possible to propose a map of CNO induced scenarios in

simple situations, with a suitable space of relevant biophysical parameters. These questions

will be addressed on modelling and mathematical grounds in a forthcoming paper.

Finding DREADD expression in ACs was initially challenging. We managed to circumvent the

interference of DREADD activated ACs by combining MEA recordings with IHC, which is not

a straightforward solution and renders experiments much more complex. Our experimental

design was based on Cre-Lox recombination, resulting in DREADD expression in all cells with

the same promoter gene in the organism. In future work, DREADD expression should be

targeted only to RGCs via intraocular viral transfection in order to avoid such strong side

effects (48). Blocking AC input onto postsynaptic targets (bipolar cells or RGCs) would be

another solution but has a big caveat: such disinhibition leads to massive increase in activity

levels in all RGCs, hence masking RGC specific DREADD activation in specific subgroups

where DREADDs are directly expressed on RGCs.

Methods

Animals and retina preparation:

All experimental procedures were approved by the ethics committee at Newcastle University

and carried out in accordance with the guidelines of the UK Home Office, under control of the

Animals (Scientific Procedures) Act 1986. Grik4 (C57BL/6-Tg(Grik4-cre)G32-4Stl/J, the

Jackson Laboratory, MA, JAX Stock No: 006474) and Scnn1a (B6;C3-Tg(Scnn1a-cre)3Aibs/J,

JAX Stock No: 009613) mice were cross-bred with Gq-DREADD mice (B6N;129-Tg(CAG-

CHRM3*,-mCitrine)1Ute/J, JAX Stock No: 026220) to generate a strain of mice with the

excitatory Gq-DREADD expressed in Grik4 and Scnn1a expressing cells (from here on

named: Grik4-DREADD and Scnn1a-DREADD, respectively). In addition, we crossbred the

Grik4 and Scnn1a lines with an inhibitory DREADD (B6.129-*Gt(ROSA)26Sortm1(CAG-*

CHRM4,-mCitrine)Ute*/J, JAX Stock No: 026219) but the effect on RGC firing rate was

negligible and the litters were used only for immunofluorescence studies. Male and female

wild-type, Grik4-DREADD and Scnn1a-DREADD mice, housed under a 12-hour light-dark

cycle and aged between postnatal days (P) 53-148 were used for the experiments. Mice were

dark-adapted overnight and killed by cervical dislocation. Eyes were enucleated, and following

removal of the cornea, lens, and vitreous body, they were placed in artificial cerebrospinal fluid

(aCSF) containing the following (in mM): 118 NaCl, 25 $NaHCO_3$, 1 NaH_2PO_4, 3 KCl, 1 $MgCl_2$,

2 $CaCl_2$, 10 glucose, and 0.5 l-Glutamine, equilibrated with 95% O_2 and 5% CO_2. The ventral

and dorsal orientation was marked after enucleation. The retina was isolated from the eye cup

and flattened for MEA recordings. For vertical cryosections, mouse eyecups were fixed in 4%

paraformaldehyde (PFA; Alfa Aesar, MA) in 0.1 M phosphate buffer solution (PBS) for 2 x 20

minutes at room temperature and washed with PBS several times. For whole mounts, retinas

were isolated from the eye cup and mounted on nitrocellulose paper (Sartorius, Germany) and

transferred to 4% PFA in PBS (2 x 20 min), rinsed in PBS and prepared for further procedures.

All procedures involving live animals and retinas were performed in dim red light and the room

was maintained in darkness throughout the experiment.

Immunohistochemistry and image acquisition:

After enucleation and fixation, the retinal tissue was processed in different ways for vertical

and whole mount immunohistochemistry (IHC). For vertical sections, eyecups were

cryoprotected in 30% sucrose in PBS overnight at 4°C and embedded in OCT Tissue TeK

(Sakura, NL) at -20°C on the following day. Vertical sections (15-20 μm) were cut on a

OTF5000 cryostat (Bright Instruments, UK) and collected on Superfrost microscope slides

(Thermo Fisher). Vertical sections and whole mounts were blocked with 10% normal goat

serum (NGS) and/or 10% normal donkey serum (NDS) in PBS for at least 30 minutes at room

temperature.

After the blocking procedure and a short rinse in PBS, Vertical sections were incubated with

primary antibodies in 5% NGS (and/or NDS) + 1% Triton X-100 + PBS overnight at 4°C. Whole

mounts were incubated free-floating with primary antibodies in 5% NGS (and/or NDS) + 1%

Triton X-100 + PBS for 4-5 days at 4°C. Incubation with secondary antibodies in 1% Triton X-

100 in PBS was carried out for 2 hours at room temperature for vertical sections or overnight

at 4°C for whole mounts. Details of the primary antibodies are as follows: anti-GFP (chicken,

Abcam 13970, 1:500-1000), anti-RBPMS (rabbit, Phosphosolutions 1830, 1:1000), anti-GABA

(mouse, Sigma Aldrich A0310, 1:1000), anti-Calretinin (mouse, Swant 6B3, 1:1000) and anti-

Parvalbumin (rabbit, Swant PV27, 1:1000). Secondary antibodies are as follows (all

concentrations 1:500): goat anti-chicken CF488, goat anti-rabbit Alexa568/Alexa647 and

donkey anti-mouse Alexa647. After washing several times with PBS, sections and whole

mounts were mounted in Vectashield (Vector Laboratories, UK). All incubations and washing

procedures were performed in the dark.

Images were captured using either a Zeiss Axio Imager upright microscope with Apotome

structured illumination fluorescence (using 20x/40x air objectives) or a Zeiss LSM800 confocal

microscope with 40x oil objective (Zeiss, Germany) operated with Zen software. Whole mount

images were stitched together using Zen software, all other image post processing was done

with Fiji (https://fiji.sc), Adobe Photoshop (Adobe, CA) and MATLAB (Mathworks, MA). The

steps between single sections of confocal stacks were not exceeding 1 µm and 3 - 5 sections

were superimposed with Fiji for presentation. For the cell density maps, cells were manually

counted using the "Cell Counter" plugin in Fiji and a bivariate histogram (MATLAB *hist3*, bin

size 60 x 60 µm) was calculated for the cell densities. For visualizing purposes, a 2-D Gaussian

filtering was applied (MATLAB *imgaussfilt*, sigma 3).

 Large-scale, high-density multielectrode array recordings and light stimulation:

Recordings were performed on the BioCamX platform with high-density-multielectrode array

(HD-MEA) Arena chips (3Brain GmbH, Lanquart, Switzerland), integrating 4096 square

microelectrodes in a 2.67 x 2.67 mm area and aligned in a square grid with 42 µM spacing.

The isolated retina was placed, RGC layer facing down, onto the MEA chip and flattened by

placing a small piece of translucent polyester membrane filter (Sterlitech Corp., Kent, WA,

USA) on the retina followed by a home-made anchor. Retinas were maintained at 33°C using

an in-line heater (Warner Instruments LLC, Hamden, CT, USA) and continuously perfused

using a peristaltic pump (~1 ml min−1). Retinas were left to settle on the MEA for at least 2

hours before recording. The platform records at a sampling rate of ~18 kHz/electrode when

using the full 64×64 array. Recordings are filtered at 50Hz high-pass filter using BrainWaveX

software (3Brain) and stored in hdf5-based data format. Spikes were detected and sorted

using Herdingspikes2 (https://github.com/mhhennig/HS2) as in (16). Briefly, spikes were first

detected as threshold crossings individually on each channel, and then merged into unique

events based on spatial and temporal proximity. For each detected spike, a location was

estimated based on the signal centre of mass. Spike sorting was performed by clustering all

events using a feature vector consisting of the locations and the first two principal components

of the largest waveform.

Light stimuli were projected onto the retina as described elsewhere (17). Briefly, the projector

irradiance was attenuated using neutral density filters to mesopic light levels (white 4 µW/cm^2).

For stimuli we used a full field 'chirp' stimulus consisting of various 1-sec contrast steps,

increasing frequency (1-15Hz) and contrast modulations (1-93 Michelson contrast) which was

repeated 5 times. We also used random black and white moving bars (width 100 µM, 12

directions (30° separation)), 800 mm/s, and the whole sequence repeated 5 times. For the

chirp and motion stimuli, we estimated each unit's instantaneous firing rate for the different

stimuli by convolving its spike train with a Gaussian kernel smoothing function (standard

deviation (SD) = 25 ms). We then averaged the trials and extracted several features including

the Bias Index and the response duration (see (16)). On average, recording from one retina

yields light responses from hundreds to thousands of individual RGCs.

Figure 9: Registering spike location with cell labelling using blood vessel landmarks. Control points in the fixed pre-labelled (A) and moving post-labelled (B) blood vessel image were aligned (C, insets a & b from A and B, respectively) and the resulting transformation matrix (D) was fixed aligned with the spike locations (E) and additional immunostainings (F). D = dorsal; V = ventral; N = nasal; T = temporal; Grid electrode size 2.65 x 2.65 mm

Registering RGC activity with IHC – blood vessels as reference marker:

To register RGC activity with IHC, we used pre- and post-labelled blood vessels as reference

markers (Fig. 1). For pre-labelled blood vessel staining, the eye cup was incubated for 1 hr in

aCSF + 5 µM Sulforhodamine 101 (SR101, Sigma Aldrich, MO) and afterwards transferred to

aCSF + 0.06 µM SR101 for MEA recording procedures. After recording, the weight and the

polyester membrane were carefully removed to expose the retina, and the MEA well rinsed

with aCSF. If the blood vessels were not visible, a 50% Optiprep (Sigma Aldrich, MO) /50%

aCSF solution was used for rinsing and further clearing. The MEA chip was mounted on a

microscope stage (Olympus AX70, CoolLED pE fluorescence) while continuing rinsing every

2-3 min with oxygenated aCSF to ensure the tissue remains healthy throughout the procedure.

The best stained blood vessel layer (either superficial or deep vascular plexus) was meander-

like imaged (Fig. 9 A). For every captured blood vessel, images were acquired in the same

location all the way down to the electrodes as well. Neighbouring images were manually

stitched together in Adobe Photoshop (Adobe, CA) later. The retina was then carefully

removed from the MEA, flattened on a nitrocellulose membrane for IHC procedure. Blood

vessel staining was enhanced by using an anti-mouse secondary antibody in the same

excitation spectrum (normally Alexa568) and blood vessels were imaged again (same plexus

as while the retina was imaged on the MEA (Fig. 9 B). Both blood vessel staining images, pre-

and post-tissue fixation were correlated with each other using the Control Point Selection Tool

in Matlab (Fig. 9 C). Briefly, the images taken in the live retina on the MEA were used as the

fixed (reference) image and images acquired post fixation as the moving image. Minimum 50

reference points were picked for the fixed and moving images to create a geometric

transformation (Matlab *fitgeotrans, lwm, 50*). That geometric transformation was applied to the

GFP image. A stack (hereafter called MEA images) with the MEA electrode image, blood

vessels pre- and post-fixation and GFP image was created in Adobe Photoshop (Fig. 9 F).

Registering RGC activity with IHC – Grik4/Scnn1a DREADD RGC identification:

The main procedure to identify and register RGC activity (spike clusters) with the MEA images

is described in the main results (Figures 5, 9). Only units that did show a significant change in

spontaneous firing rate (sampled during at least 5 min) before and after adding CNO to the

chamber were selected for Grik4/Scnn1a RGC identification and response clustering. Units

with either >50% or <50% change in firing rate in CNO were considered as potential

Grik4/Scnn1a candidates. We also used the Burst Index (32) to look at potential changes in

activity levels induced by CNO, with potential Grik4/Scnn1a cells having changes in their Burst

Index exceeding +/- 50%. Only these selected RGCs were further identified in the next steps.

During the spike sorting process with Herdingspikes2, the physical x and y spike localization

for each detected spike from a single RGC unit is calculated (Figure 5 B). The spike location

coordinates are given for a 64 x64 grid (the electrode layout of our MEA). Occasionally, we

confirmed the cluster centre (e.g., the centre of mass of negative spike amplitude peaks)

calculated by Herdingspikes2 by overlaying the x and y localization for the negative voltage

peak (Figure 5 C magenta diamond) obtained through electrical imaging. For electrical

imaging, the voltage fluctuations from the spike origin electrode and the neighbouring 7

electrode rows and columns were read out. For a minimum of 1000 spikes per RGC unit,

voltage fluctuations before and after (here 5 ms) each spike were normalized, and maximally

projected with absolute values (Figure 5 C) to visualize the current sink and the axonal

trajectories. The spike localization and current sink coordinates were multiplied using a scaling

factor to match the MEA images stack. GFP images were transformed to 8-bit images, a

threshold (Matlab *max entropy*) was applied, potential gaps filled (Matlab *imfill 'holes'*) and

binarized (Matlab *bwareaopen 50*). Binary images were analyzed (Matlab *regionprops*),

centroids and diameters for every potential GFP blob were calculated and the pairwise

distance between centroids and spike locations was measured for all combinations.

Registering RGC activity with IHC – Grik4/Scnn1a response clustering:

All potential Grik4/Scnn1a RGC units linked to a GFP blob (see Results) were further analyzed

and grouped according to their response to a chirp stimulus. But before that grouping, cells

were pre-classified into motion sensitive (responding preferably to either direction or

orientation of motion) or cells that respond best to static stimuli (full field, chirp). We calculated

the direction (DSI) and orientation (OSI) selectivity index as described elsewhere (16) but used

a cut-off value for every experiment independently. Briefly, we fitted the tabulated data with a

mixture of two gaussians (Figure 6 B, E). The fit parameters were computed using maximum

likelihood parameter estimates (Matlab *mle*) and mu2 (the second peak, Figure 6 B, E) was

used as cut-off value. The trail variability and signal to noise ratio was calculated (16) for each

stationary/non-stationary RGC, and units with a value below the 25th percentile were

discarded. The chirp responses of the remaining Grik4 and Scnn1a stationary/non-stationary

RGC units were used to establish response groups. We used a similar approach to calculate

the SPIKE distance as described in our earlier work (10). Briefly, the SPIKE distances were

computed using the open-source package PYSPIKE (30). The pairwise distances between

two units were determined by computing the pairwise distances of all trials of the chirp

stimulus. The resulting distance matrix was then clustered with a hierarchical clustering

algorithm (Python, *Scipy library dendrogram*). Matlab and Python scripts and other extended

material will be made available via https://github.com/GerritHilgen/DREADD_RGCclass.

525   Acknowledgments

This project was funded by the Leverhulme Trust (RPG-2016-315 to ES and BC), by

Newcastle University (Faculty of Medical Sciences Graduate School and Pro-Vice Chancellor

Discretionary Fund). We thank Matthias Hennig for help with the SPIKE distance calculations

and Chris Williams who worked on related, unpublished aspects of the project. We also thank

the Bioimaging Unit at Newcastle University for providing excellent service and help for this

project.

533   Authors' contributions

534   Conceptualization: G.H., E.S., B.C.; Software: G.H. Methodology: G.H., E.S.; Formal Analysis:

535   GH., Investigation, G.H., V.K., E.K.; Writing – Original Draft: G.H., E.S., B.C., V.K., J.K.;

536   Project administration: E.S., B.C.; Funding Acquisition: E.S., B.C.

538   Legends

Figure 1: Grik4 and Scnn1a cells in the GCL are not homogeneously distributed. Whole mount

antibody staining against Grik4 (A) and Scnn1a (B) DREADD GFP were imaged at the level

of the GCL. All stained GFP cells were counted and the densities were calculated and

presented in pseudocolours for 3 Grik4 (C) and Scnn1a (D) retinas. V= ventral, T = temporal,

D = dorsal, N = nasal. Scale bar A, B = 1 mm; Scale bar A, B insets II = 100 µm.

Figure 2: Grik4 and Scnn1a DREADD are expressed in RGCs and ACs. Vertical sections were

stained for Grik4 (A, C, cyan) and Scnn1a (B, D, cyan) GFP. Sections were further co-labelled

with RBPMS (A, B, magenta), a marker for RGCs, and GABA (C, D, magenta). INL = inner

nuclear layer, IPL = inner plexiform layer, GCL = ganglion cell layer. Scale bar in D = 20 µm.

Figure 3: Grik4 and Scnn1a DREADD are expressed in multiple RGC types. Vertical sections

of Grik4 and Scnn1a retinas were triple stained A, B) for GFP (cyan), RBPMS (magenta) and

Calretinin (yellow) or C, D) GFP (cyan), Parvalbumin (magenta) and Calretinin (yellow). INL =

inner nuclear layer, IPL = inner plexiform layer, GCL = ganglion cell layer. Scale bar in D = 20

µm.

Figure 4: DREADD activation leads to spike pattern changes in RGCs. DREADDs in RGCs

can be activated with Clozapine N-oxide (CNO) and lead to an increase in firing frequency (A,

right half) and sometimes also in bursting activity (B, right half). Similar activities can be evoked

with an "allround" AC neurotransmitter blocker (C) without CNO. The cocktail contains: 2 µM

Strychnine (glycine receptor antagonist), 20 µM Mecamylamine (nicotinic acetylcholine

receptor antagonist), 20 µM Bicuculline ($GABA_A$ antagonist), 5 µM CGP 55845 ((2S)-3-[[(1S)

-1- (3,4-Dichlorophenyl)ethyl] amino -2- hydroxypropyl] (phenylmethyl) phosphinic acid

hydrochloride, $GABA_B$ antagonist), 50 µM TPMPA (1,2,5,6-Tetrahydropyridin-4-yl

methylphosphinic acid, $GABA_C$ antagonist). Plotted in A, B are 20 Grik4 RGCs that showed a

minimum of 50% change in firing frequency or bursting activity and in C 100 randomly selected

Scnn1a RGC units. Control in A & B means without CNO, in C without blockers. Vertical lines

represent spike times plotted as a function of time.

Figure 5: Registering GFP RGCs with nearby isolated spike centre clusters. The spike cluster

centre (B, circle) and electrical imaging (C, diamond) define the potential electrical source

location of the isolated RGCs. Spike centres in close proximity to GFP labelled RGCs (D) were

registered as potential DREADD-expressing RGCs (E, F, Match). Scale bars A, C, F = 40 µm;

B = 80 µm; E = 20 µm.

Figure 6: Dividing registered Grik4 and Scnn1a RGCs into stationary and non-stationary

responders. Moving stimuli were used to estimate the direction selectivity index (DSI) of

registered "Match" RGCs (Grik4 A, Scnn1a D, magenta). The DSI was plotted against the

probability density (histogram) and fitted with two gaussians (B, E, red line). RGCs exceeding

a set threshold (B, E, dotted red line) were labelled "non-Stationary". The registered RGCs

from all retinas were grouped into "non-Stationary" and "stationary" (C, F).

Figure 7: Clustering of Grik4 stationary and non-stationary RGC responses. RGC responses

that showed a high spike train similarity for a chirp stimulus (B, F) were grouped together using

gap statistics (A). For each RGC of the groups, the PSTH was calculated and the mean PSTH

was plotted (B, F, coloured lines). The means for Bias Index (C, D, G), Response Duration

(C), Baseline Spiking (D), Burst change (E), Spike Rate change (E) and orientation selectivity

index (OSI, G) index were scatter plotted and their standard deviations were used as the circle

diameters.

Figure 8: Clustering of Scnn1a stationary and non-stationary RGC responses. RGC responses

that showed a high spike train similarity for a chirp stimulus (B, F) were grouped together using

gap statistics (A). For each RGC of the groups, the PSTH was calculated and the mean PSTH

was plotted (B, F, coloured lines). The means for Bias Index (C, D, G), Response Duration

(C), Baseline Spiking (D), Burst change (E), Spike Rate change (E) and orientation selectivity

index (OSI, G) index were scatter plotted and their standard deviations were used as the circle

diameters.

Figure 9: Registering spike location with cell labelling using blood vessel landmarks. Control

points in the fixed pre-labelled (A) and moving post-labelled (B) blood vessel image were

aligned (C, insets a & b from A and B, respectively) and the resulting transformation matrix

(D) was fixed aligned with the spike locations (E) and additional immunostainings (F). D =

dorsal; V = ventral; N = nasal; T = temporal; Grid electrode size 2.65 x 2.65 mm

Table 1: Summary of identified Grik4 clusters and potential matches with existing literature.

NB these are suggestions.

Table 2: Summary of identified Scnn1a clusters and potential matches with existing literature.

NB these are suggestions.

References

1. Euler T, Haverkamp S, Schubert T, Baden T. Retinal bipolar cells: elementary building blocks of vision. Nat Rev Neurosci. 2014 Aug;15(8):507–19.

2. Curcio CA, Allen KA. Topography of ganglion cells in human retina [Internet]. Vol. 300, The Journal of Comparative Neurology. 1990. p. 5–25. Available from: http://dx.doi.org/10.1002/cne.903000103

3. Jeon C-J, Strettoi E, Masland RH. The Major Cell Populations of the Mouse Retina [Internet]. Vol. 18, The Journal of Neuroscience. 1998. p. 8936–46. Available from: http://dx.doi.org/10.1523/jneurosci.18-21-08936.1998

4. Masland RH. The fundamental plan of the retina. Nat Neurosci. 2001 Sep;4(9):877–86.

5. Baden T, Berens P, Franke K, Román Rosón M, Bethge M, Euler T. The functional diversity of retinal ganglion cells in the mouse. Nature. 2016 Jan 21;529(7586):345–50.

6. Tran NM, Shekhar K, Whitney IE, Jacobi A, Benhar I, Hong G, et al. Single-Cell Profiles of Retinal Ganglion Cells Differing in Resilience to Injury Reveal Neuroprotective Genes. Neuron. 2019 Dec 18;104(6):1039–55.e12.

7. Coombs J, van der List D, Wang G-Y, Chalupa LM. Morphological properties of mouse retinal ganglion cells. Neuroscience. 2006 Jun 19;140(1):123–36.

8. Völgyi B, Chheda S, Bloomfield SA. Tracer coupling patterns of the ganglion cell subtypes in the mouse retina. J Comp Neurol. 2009 Feb 10;512(5):664–87.

9. Zeck GM, Masland RH. Spike train signatures of retinal ganglion cell types. Eur J Neurosci. 2007 Jul;26(2):367–80.

10. Jouty J, Hilgen G, Sernagor E, Hennig MH. Non-parametric Physiological Classification of Retinal Ganglion Cells in the Mouse Retina. Front Cell Neurosci. 2018 Dec 7;12:481.

11. Sanes JR, Masland RH. The types of retinal ganglion cells: current status and implications for neuronal classification. Annu Rev Neurosci. 2015 Jul 8;38:221–46.

12. Rheaume BA, Jereen A, Bolisetty M, Sajid MS, Yang Y, Renna K, et al. Single cell transcriptome profiling of retinal ganglion cells identifies cellular subtypes. Nat Commun. 2018 Jul 17;9(1):2759.

13. Martersteck EM, Hirokawa KE, Evarts M, Bernard A, Duan X, Li Y, et al. Diverse Central Projection Patterns of Retinal Ganglion Cells. Cell Rep. 2017 Feb 21;18(8):2058–72.

14. Laboissonniere LA, Goetz JJ, Martin GM, Bi R, Lund TJS, Ellson L, et al. Molecular signatures of retinal ganglion cells revealed through single cell profiling. Sci Rep. 2019 Oct 31;9(1):15778.

15. Marblestone* AH, Zamft* BM, Maguire YG, Shapiro MG, Cybulski TR, Glaser JI, et al. Physical principles for scalable neural recording. Front Comput Neurosci [Internet]. 2013 [cited 2021 Apr 5];7. Available from: https://www.frontiersin.org/articles/10.3389/fncom.2013.00137/pdf

16. Hilgen G, Pirmoradian S, Pamplona D, Kornprobst P, Cessac B, Hennig MH, et al. Pan-retinal characterisation of Light Responses from Ganglion Cells in the Developing Mouse Retina. Sci Rep. 2017 Feb 10;7:42330.

17. Portelli G, Barrett JM, Hilgen G, Masquelier T, Maccione A, Di Marco S, et al. Rank Order Coding: a Retinal Information Decoding Strategy Revealed by Large-Scale Multielectrode Array Retinal Recordings. eNeuro [Internet]. 2016 May;3(3). Available from: http://dx.doi.org/10.1523/ENEURO.0134-15.2016

18. Nakazawa K, Quirk MC, Chitwood RA, Watanabe M, Yeckel MF, Sun LD, et al. Requirement for hippocampal CA3 NMDA receptors in associative memory recall. Science. 2002 Jul 12;297(5579):211–8.

19. Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, et al. A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. Nat Neurosci. 2010 Jan;13(1):133–40.

20. Armbruster BN, Li X, Pausch MH, Herlitze S, Roth BL. Evolving the lock to fit the key to create a family of G protein-coupled receptors potently activated by an inert ligand. Proc Natl Acad Sci U S A. 2007 Mar 20;104(12):5163–8.

21. Roth BL. DREADDs for Neuroscientists. Neuron. 2016 Feb 17;89(4):683–94.

22. Urban DJ, Roth BL. DREADDs (designer receptors exclusively activated by designer drugs): chemogenetic tools with therapeutic utility. Annu Rev Pharmacol Toxicol. 2015;55:399–417.

23. Zhu H, Aryal DK, Olsen RHJ, Urban DJ, Swearingen A, Forbes S, et al. Cre-dependent DREADD (Designer Receptors Exclusively Activated by Designer Drugs) mice. Genesis. 2016 Aug;54(8):439–46.

24. Rodriguez AR, de Sevilla Müller LP, Brecha NC. The RNA binding protein RBPMS is a selective marker of ganglion cells in the mammalian retina [Internet]. Vol. 522, Journal of Comparative Neurology. 2014. p. 1411–43. Available from: http://dx.doi.org/10.1002/cne.23521

25. Lee E-S, Lee J-Y, Jeon C-J. Types and density of calretinin-containing retinal ganglion cells in mouse. Neurosci Res. 2010 Feb;66(2):141–50.

26. Yi C-W, Yu S-H, Lee E-S, Lee J-G, Jeon C-J. Types of parvalbumin-containing retinotectal ganglion cells in mouse. Acta Histochem Cytochem. 2012 Jun 28;45(3):201–10.

27. Verbist C, Müller MG, Mansvelder HD, Legenstein R, Giugliano M. The location of the axon initial segment affects the bandwidth of spike initiation dynamics. PLoS Comput Biol. 2020 Jul;16(7):e1008087.

28. Colbert CM, Johnston D. Axonal action-potential initiation and Na+ channel densities in the soma and axon initial segment of subicular pyramidal neurons. J Neurosci. 1996 Nov 1;16(21):6676–86.

29. Kreuz T, Chicharro D, Houghton C, Andrzejak RG, Mormann F. Monitoring spike train synchrony. J Neurophysiol. 2013 Mar;109(5):1457–72.

30. PySpike—A Python library for analyzing spike train synchrony. SoftwareX. 2016 Jan 1;5:183–9.

31. Tibshirani R, Walther G, Hastie T. Estimating the number of clusters in a data set via the gap statistic. J R Stat Soc Series B Stat Methodol. 2001;63(2):411–23.

32. Jones TA, Leake PA, Snyder RL, Stakhovskaya O, Bonham B. Spontaneous discharge patterns in cochlear spiral ganglion cells before the onset of hearing in cats. J Neurophysiol. 2007 Oct;98(4):1898–908.

33. Drinnenberg A, Franke F, Morikawa RK, Jüttner J, Hillier D, Hantz P, et al. How Diverse Retinal Functions Arise from Feedback at the First Visual Synapse. Neuron. 2018 Jul 11;99(1):117–34.e11.

34. Li PH, Gauthier JL, Schiff M, Sher A, Ahn D, Field GD, et al. Anatomical identification of extracellularly recorded cells in large-scale multielectrode recordings. J Neurosci. 2015 Mar 18;35(11):4663–75.

35. Pisano F, Zampaglione E, McAlinden N, Roebber J, Dawson MD, Mathieson K, et al. Large scale matching of function to the genetic identity of retinal ganglion cells. Sci Rep. 2017 Nov 13;7(1):15395.

36. Ivanova E, Hwang G-S, Pan Z-H. Characterization of transgenic mouse lines expressing Cre recombinase in the retina. Neuroscience. 2010 Jan 13;165(1):233–43.

37. Rivlin-Etzion M, Zhou K, Wei W, Elstrott J, Nguyen PL, Barres BA, et al. Transgenic mice reveal unexpected diversity of on-off direction-selective retinal ganglion cell subtypes and brain structures involved in motion processing. J Neurosci. 2011 Jun 15;31(24):8760–9.

38. Johnson KP, Zhao L, Kerschensteiner D. A Pixel-Encoder Retinal Ganglion Cell with Spatially Offset Excitatory and Inhibitory Receptive Fields. Cell Rep. 2018 Feb 6;22(6):1462–72.

39. Jacoby J, Schwartz GW. Three Small-Receptive-Field Ganglion Cells in the Mouse Retina Are Distinctly Tuned to Size, Speed, and Object Motion. J Neurosci. 2017 Jan 18;37(3):610–25.

40. Krieger B, Qiao M, Rousso DL, Sanes JR, Meister M. Four alpha ganglion cell types in mouse retina: Function, structure, and molecular signatures. PLoS One. 2017 Jul 28;12(7):e0180091.

41. Hilgen G, Lockwood E, Ratcliff JEJ, Sernagor E. Pan-retinal characterization of parvalbumin-calretinin co-expressing cells in the mouse retina. F1000Res [Internet]. 2019 Sep 9 [cited 2021 Apr 28];8. Available from: http://dx.doi.org/10.7490/f1000research.1117479.1

42. Vaney DI, Sivyer B, Taylor WR. Direction selectivity in the retina: symmetry and asymmetry in structure and function. Nat Rev Neurosci. 2012 Feb 8;13(3):194–208.

43. Rousso DL, Qiao M, Kagan RD, Yamagata M, Palmiter RD, Sanes JR. Two Pairs of ON and OFF Retinal Ganglion Cells Are Defined by Intersectional Patterns of Transcription Factor Expression. Cell Rep. 2016 May 31;15(9):1930–44.

44. Kovács-Öller T, Szarka G, Tengölics ÁJ, Ganczer A, Balogh B, Szabó-Meleg E, et al. Spatial Expression Pattern of the Major Ca-Buffer Proteins in Mouse Retinal Ganglion Cells. Cells [Internet]. 2020 Mar 25;9(4). Available from: http://dx.doi.org/10.3390/cells9040792

45. Zhang Y, Kim I-J, Sanes JR, Meister M. The most numerous ganglion cell type of the mouse retina is a selective feature detector. Proc Natl Acad Sci U S A. 2012 Sep 4;109(36):E2391–8.

46. de Vries SEJ, Baccus SA, Meister M. The projective field of a retinal amacrine cell. J Neurosci. 2011 Jun 8;31(23):8595–604.

47. Deny S, Ferrari U, Macé E, Yger P, Caplette R, Picaud S, et al. Multiplexed computations in retinal ganglion cells of a single type. Nat Commun. 2017 Dec 6;8(1):1964.

48. Storchi R, Milosavljevic N, Eleftheriou CG, Martial FP, Orlowska-Feuer P, Bedford RA, et al. Melanopsin-driven increases in maintained activity enhance thalamic visual response reliability across a simulated dawn. Proc Natl Acad Sci U S A. 2015 Oct 20;112(42):E5734–43.