1  **Establishment of CRISPR/Cas9-based knock-in in a hemimetabolous insect: targeted gene**
2  **tagging in the cricket *Gryllus bimaculatus***
3

4  Yuji Matsuoka[1,3*], Taro Nakamura[2,3*], Takahito Watanabe[1,4], Austen A. Barnett[2,5], Sayuri
5  Tomonari[1], Guillem Ylla[6], Carrie A. Whittle[7], Sumihare Noji[8], Taro Mito[1*], Cassandra G.
6  Extavour[2,7,9*]
7
1. Department of Life Systems, Institute of Technology and Science, the University of
Tokushima Graduate School, 201 Minami-Jyosanjima-cho, Tokushima City, 770-8506, Japan
2. Department of Organismic and Evolutionary Biology, 16 Divinity Avenue, Cambridge MA
02138, USA
3. Current address: National Institute for Basic Biology, Nishigonaka 38, Myodaiji, Okazaki 444-
8585, Aichi, Japan
4. Bio-Innovation Research Center, Tokushima University, 2272-2 Ishii, Ishii-cho, Myozai-gun,
Tokushima 779-3233, Japan
5. Current address: DeSales University, 2755 Station Avenue, Center Valley, PA 18034, USA
6. Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Krakow, 30-
387, Poland
7. Howard Hughes Medical Institute, Chevy Chase MD, USA
8. Tokushima University, 2-14 Shinkura-cho, Tokushima City, 770-8501, Japan
9. Department of Molecular and Cellular Biology, 16 Divinity Avenue, Cambridge MA 02138,
USA
23
24
25 **\* Correspondence to:**     Yuji Matsuoka matsuoka@nibb.ac.jp
26                             Taro Nakamura taro@nibb.ac.jp
27                             Taro Mito mito.taro@tokushima-u.ac.jp
28                             Cassandra G. Extavour extavour@oeb.harvard.edu
29

**Running Title:** CRISPR/Cas9 knock-ins in a cricket

**Abstract**

Studies of traditional model organisms like the fruit fly *Drosophila melanogaster* have

contributed immensely to our understanding of the genetic basis of developmental processes.

However, the generalizability of these findings cannot be confirmed without functional genetic

analyses in additional organisms. Direct genome editing using targeted nucleases has the

potential to transform hitherto poorly-understood organisms into viable laboratory organisms for

functional genetic study. To this end, here we present a method to induce targeted genome

knock-out and knock-in of desired sequences in an insect that serves as an informative contrast

to *Drosophila*, the cricket *Gryllus bimaculatus*. The efficiency of germ line transmission of

induced mutations is comparable to that reported for other well-studied laboratory organisms,

and knock-ins targeting introns yield viable, fertile animals in which knock-in events are directly

detectable by visualization of a fluorescent marker in the expression pattern of the targeted gene.

Combined with the recently assembled and annotated genome of this cricket, this knock-

in/knock-out method increases the viability of *G. bimaculatus* as a tractable system for functional

genetics in a basally branching insect.

**Keywords**: CRISPR/Cas9, Orthoptera, genome editing, Hox genes, *Ultrabithorax*, *abdominal-A*

Matsuoka, Nakamura et al.

**Introduction**

In what is often called the "post-genomic era," (Wainberg et al., 2021), massive

advances in nucleic acid sequencing chemistry over the last two decades have given scientists

access to greater volumes of gene sequence data than ever before (Kulski, 2016; Papageorgiou et

al., 2018). However, this wealth of genomic information has highlighted two major gaps in our

understanding of gene function and evolution. First, comparative genomic data and increased

taxon sampling in functional genetic, developmental and cellular biology have revealed that the

biology of many traditional laboratory model organisms is not representative of the broader

clades to which they belong (Goldstein and King, 2016). Second, our ability to accurately deduce

gene function from sequence data is limited to those genes that display high sequence and

structural conservation (Ashburner et al., 2000; Consortium et al., 2020), and tools for

manipulating gene function have been developed for only a small fraction of organisms (Russell

et al., 2017). Addressing these problems calls for both increased taxon sampling and

development of techniques to enable targeted alteration of gene function in understudied

organisms. Here we address both of these issues by developing a method for targeted genome

editing, including both knock-out and knock-in editing, in a basally branching insect model

organism, the cricket *Gryllus bimaculatus*.

*G. bimaculatus* is an emerging model organism in a variety of fields of biology (Horch

et al., 2017a; Kulkarni and Extavour, 2019). Ease of husbandry (Horch et al., 2017b), detailed

developmental staging tables (Donoughe and Extavour, 2016), established gene expression

analysis methods (Horch et al., 2017b), and an assembled and annotated genome (Ylla et al.,

2021) make this cricket a highly amenable hemimetabolous laboratory model system (Kulkarni

and Extavour, 2019). In contrast to the ontogenetically derived model *Drosophila melanogaster*,

many aspects of cricket embryogenesis are thought to resemble putative ancestral developmental

modes of insects (Davis and Patel, 2002). For example, the function of several axial patterning

genes has been analyzed and compared to that of their *D. melanogaster* homologues, revealing

that the gene regulatory networks governing axial patterning have undergone considerable

evolutionary change across insects (Mito et al., 2007; Mito et al., 2008; Matsuoka et al., 2015).

*G. bimaculatus* is also used for the analysis of gene function in tissue regeneration (Mito et al.,

2002; Nakamura et al., 2007; Bando et al., 2013), as this cricket can regenerate amputated

organs, including legs and antennae, after several rounds of juvenile molts. In addition, *G.*

*bimaculatus* is used for the analysis of gene function and neuronal circuits in neuronal activity,

including learning, memory and circadian clocks (Hedwig and Sarmiento-Ponce, 2017;

Matsumoto et al., 2018; Mizunami and Matsumoto, 2017; Tomiyama et al., 2020). Moreover,

several species of cricket are being farmed as a new food source for humans because of their

high protein and nutrient content (Huis et al., 2013).

Genome editing techniques using artificial nucleases were previously established in *G.*

*bimaculatus* (Watanabe et al., 2012). However, construction of artificial nucleases is laborious.

More recently, use of the clustered regulatory interspaced short palindromic repeat

(CRISPR)/associated Cas9 nuclease (CRISPR/Cas9) has emerged and been verified as an

efficient tool for genome editing in several arthropod species, including the fruit fly *Drosophila*

*melanogaster* (Gratz et al., 2013), the beetle *Tribolium castaneum* (Gilles et al., 2015), the

mosquito *Aedes aegypti* (Kistler et al., 2015), multiple butterfly species (Li et al., 2015;

Matsuoka and Monteiro, 2018; Zhang et al., 2017), the amphipod *Parhyale hawaiensis*, (Martin

et al., 2016), the clonal raider ant *Ooceraea biroi* (Trible et al., 2017), the European honeybee

*Apis mellifera* (Kohno and Kubo, 2018), and the milkweed bug *Oncopeltus fasciatus* (Reding

Matsuoka, Nakamura et al.

and Pick, 2020). Among crickets, we and others have reported CRISPR/Cas9-mediated

knockouts in *Acheta domesticus* (Dossey e tal. 2023) and in *G. bimaculatus* (Bai et al. 2023,

Nakamura et al. 2022). Knock0in has also been reported for *A. domesticus* (Dossey et al. 2023),

but this approach has not been used to generate the types of enhancer trap or protein trap lines

that have proven so beneficial to advancing developmental biology research in other insects

(Morin et al. 2001). Here we briefly review this technique, investigate and demonstrate its utility

for targeted gene trap knock-in genome modification in *G. bimaculatus*, the most widely used

orthopteran model for developmental biology.

In the CRISPR/Cas9 system, short guide RNAs (sgRNAs) recruit Cas9 nuclease to the

target sequence, and Cas9 then introduces a double strand break (DSB) at the target sequence.

The presence of DSBs triggers the activity of the cell's DNA repair machinery, either <u>n</u>on-

<u>h</u>omologous <u>e</u>nd <u>j</u>oining (NHEJ) or <u>h</u>omology <u>d</u>irected <u>r</u>epair (HDR). NHEJ is an error-prone

machinery, such that insertions or deletions can be generated at the break point (Branzei and

Foiani, 2008). By using artificial nucleases to trigger NHEJ, we previously succeeded in

generating *G. bimaculatus* mutant lines (Watanabe et al., 2012). HDR, however, would offer

more precise repair machinery, as the break is repaired through use of a homologous template.

By supplying a donor template containing sequence homologous to the target, in principle a

desired donor sequence can be integrated into the genome though HDR. Such gene knock-ins,

while highly desirable for detailed analysis of the function of genomic regions, are more difficult

to achieve than gene knock-outs because of the low efficiency of HDR in eukaryotes (Hagmann

et al., 1998). Although success with HDR has been reported in some insects including the silk

moth *Bombyx mori* (Ma et al., 2014; Zhu et al., 2015), multiple mosquito species (Hammond et

al., 2016; Kistler et al., 2015; Purusothaman et al., 2021) and mosquito cell lines (Rozen-Gagnon

et al., 2021), the beetle *Tribolium castaneum* (Gilles et al., 2015), the medfly *Ceratitis capitata*

(Aumann et al., 2018), and the squinting bush brown butterfly *Bicyclus anynana* (Connahs et al.,

2022), our attempts at HDR-based knock-in techniques have never succeeded in *G. bimaculatus*

(unpublished observations). Recently, an efficient gene knock-in method through NHEJ was

developed in the zebrafish *Danio rerio* (Auer et al., 2014). In this method, both the genome and

the donor vector are cleaved *in vivo*, then the terminal genomic and donor sequences are

combined through NHEJ. The method is efficient and can integrate longer constructs into the

genome than knock-ins achieved through HDR (Auer et al., 2014, Yoshimi et al., 2016). Bosch

and colleagues (Bosch et al., 2019) subsequently reported that this knock-in strategy also works

in *D. melanogaster*.

Here, we present evidence that the CRISPR/Cas9 system functions efficiently in *G.*

*bimaculatus*. We demonstrate the utility of this technique for developmental biology by

performing functional analysis of the *G. bimaculatus* orthologues of the Hox genes

*Ultrabithorax* (*Gb-Ubx*) and *abdominal-A* (*Gb-abd-A*). Furthermore, using a donor vector

containing an autonomous expression cassette, we demonstrate that gene knock-in by a

homology-independent method works efficiently in *G. bimaculatus*. We show that this

homology-independent gene knock-in method can be applied to identify mutant individuals

simply by detecting marker gene expression in this cricket. Efficient targeted genome editing,

now including both knock-out and gene-tagging knock-in techniques, will pave the way for

making this cricket a much more sophisticated model animal for functional genetic laboratory

studies.

**Results**

Matsuoka, Nakamura et al.

*Targeted mutagenesis of the Gb-lac2 locus*

To determine whether the CRISPR/Cas9 system was functional in the cricket, we first tried to

perform a targeted gene knock-out of the *laccase 2* (*Gb-lac2*) gene (Table 1), which regulates

tanning of the arthropod cuticle following molting (Arakane et al., 2005). We chose this gene

because of its easily detectable loss-of-function phenotype, and because we had previously

successfully generated stable mutant lines for this gene by using artificial nucleases (Watanabe et

al., 2012). sgRNA target sites were first chosen using the CasOT tool (Xiao et al., 2014), and

then we chose the final target sequence from among these candidate sequences based on the

number of mismatches relative to the other sequences in the genome (> 3 mismatches in the

whole sgRNA sequence) and GC content (70 ± 10% for the whole sgRNA sequence). Based on

these criteria, we designed sgRNAs against the fifth exon of *Gb-lac2*, which is close to the target

regions of the previously reported artificial nuclease experiment (Fig. 1D; Table 2) (Watanabe et

al., 2012).

We co-injected 0.5 μg/μl sgRNA and 0.5 μg/μl of Cas9 mRNA into 232 fertilized

cricket eggs within 1-3 hours after egg laying (h AEL) (Table 3). Five days after injection, we

evaluated the frequency of mutant alleles in individual eggs using the Surveyor™ nuclease assay

(Qiu et al., 2004) (see Materials and Methods for detailed mechanism and procedure). We

detected cleaved fragments in 13 out of 29 eggs examined (Table 3, Fig. 1C).

We observed mosaic pigmentation of the cuticle in 92% of $G_0$ hatchlings that emerged

from the individual eggs injected with the *Gb-lac2* sgRNA, consistent with Cas9-mediated

interruption of the *Gb-lac2* gene in some, but not all, somatic cells of the $G_0$ hatchlings. We

raised these hatchlings to adulthood (Fig. 1A'), and crossed these $G_0$ adults with wild type

Matsuoka, Nakamura et al.

crickets of the opposite sex, to determine the efficiency of germ line transmission of the Cas9-

induced *Gb-lac2* mutations to the $G_1$ generation. We found that 47.4% of those injected with the

*Gb-lac2* gRNA transmitted the mutation to their offspring (Fig. 1B). To determine the nature of

the *Gb-lac2* Cas9-induced alleles, we isolated genomic DNA from each line and analyzed the

sequence of the *Gb-lac2* locus. We found that several different types of indel mutations were

introduced at the target locus (Fig. 1D). These results indicate that this CRISPR/Cas9-mediated

genome editing system is functional in the cricket.

*Targeted mutagenesis of the Gb-Ubx locus via knock-out*

To compare phenotypes obtained with targeted gene disruption to those obtained with the RNA

interference (RNAi) method that has hitherto been the most common method of performing

functional genetics in this cricket (Mito and Noji, 2008), we used the CRISPR/Cas9 system to

perform functional analyses of the *G. bimaculatus* ortholog of the Hox gene *Ultrabithorax* (*Gb-*

*Ubx*) (Table 1). We previously examined RNAi-induced phenotypes for *Gb-Ubx* in developing

abdominal segments (Barnett et al., 2019), providing a basis for comparison with CRISPR-

induced mutants. We designed sgRNA for a sequence within an exon upstream of the

homeodomain (Fig. 2A), and co-injected 0.5 µg/µl of this sgRNA and 1 µg/µl Cas9 mRNA into

167 fertilized cricket eggs within 1-3 h AEL. Seven days following injection, we extracted

genomic DNA from a small subset of injected eggs and performed the Surveyor$^{TM}$ assay to

determine the efficiency of gene targeting. We found that *Gb-Ubx* mutations had been induced in

all examined eggs (n=16) (Fig. 2B). The remaining 151 injected $G_0$ embryos gave rise to ten

adults, which we backcrossed individually to wild type adults of the opposite sex. We randomly

chose approximately 30 $G_1$ eggs from each of the ten $G_0$ crosses, extracted genomic DNA from

Matsuoka, Nakamura et al.

the pooled embryos, and performed the Surveyor$^{TM}$ assay. We found that six out of ten $G_0$

crickets transmitted *Gb-Ubx* mutations to the next generation. We selected one of the six $G_1$ *Gb-*

*Ubx$^{CRISPR}$* lines, which had a frame-shift mutation in the *Gb-Ubx* locus, for further phenotypic

analysis. These *Gb-Ubx$^{CRISPR}$* mutants displayed two different classes of phenotype: (1)

*Contraction of the T3 leg*. Wild type *G. bimaculatus* adults have large, conspicuous T3 jumping

legs. However, heterozygous *Gb-Ubx$^{CRISPR}$* mutants had smaller T3 legs than wild type animals

(Fig. 2C). Homozygous *Gb-Ubx$^{CRISPR}$* mutants obtained in the $G_2$ generation had T3 legs that

were even smaller than those of heterozygotes, almost the same size as T1/T2 legs (Fig. 2C).

This specific phenotype is not directly comparable with previously reported *Gb-Ubx* RNAi

experiments (Barnett et al., 2019), because embryos in those experiments were not reared to

hatching. However, the *Gb-Ubx$^{CRISPR}$* phenotypes were in good correspondence with those

previously observed for *Ubx* RNAi in the cricket *Acheta domesticus* (Mahfooz et al., 2007). (2)

*Transformation of the A1 appendage*. Wild type *G. bimaculatus* germ band stage embryos

possess two appendage-like organs on the A1 segment called pleuropodia (Fig. 2D; Rathke,

1844; Wheeler, 1892). Instead of the pleuropodia present in wild type adults, the appendage

outgrowths on the A1 segment of homozygous *Gb-Ubx$^{CRISPR}$* mutants were transformed towards

leg-like structures, which is consistent with detection of expression of *Gb-Dll* and other leg

patterning genes in the leg-like structures. (Fig. 2D and Supplemental Figure 1). This phenotype

matches that previously observed in *Gb-Ubx* RNAi embryos (Fig. 2E; Barnett et al., 2019). *Gb-*

*Ubx$^{CRISPR}$* heterozygous mutants were fertile but homozygous mutants were lethal. Therefore, to

maintain this line, *Gb-Ubx$^{CRISPR}$* heterozygous mutant animals of separate sexes were crossed to

each other, and we performed the Surveyor$^{TM}$ assay to isolate heterozygous mutants from among

their offspring.

Matsuoka, Nakamura et al.

To confirm whether the production of Gb-Ubx protein was indeed disrupted by these

CRISPR-induced mutations, we performed immunostaining with the "UbdA" monoclonal

antibody FP6.87 (Kelsh et al., 1994), which recognizes both Ubx and Abd-A proteins, and was

previously reported to cross-react in multiple *Gryllus* species including *G. bimaculatus* (Barnett

et al., 2019; Mahfooz et al., 2004). In wild type embryos, the UbdA antibody revealed the

expected combination of the Gb-Ubx and Gb-Abd-A expression patterns (Fig. 2E and 2F;

Barnett et al., 2019; Mahfooz et al., 2004). In *Gb-Ubx$^{CRISPR}$* embryos, however, while the UbdA

expression domain was retained in the abdomen where *Gb-abd-A* is expressed, expression was

undetectable in the T3 and A1 segment where *Gb-Ubx* is expressed (Fig. 2F), suggesting that the

CRISPR-induced *Gb-Ubx* mutations interfered with Gb-Ubx protein production. We note that in

previous reports, UbdA expression was not affected in *Gb-Ubx$^{RNAi}$* embryos, which might be

because RNAi does not disrupt target gene expression completely (REF).

We further characterized the phenotype of *Gb-Ubx$^{CRISPR}$* embryos by observing the *Gb-*

*Dll* expression pattern (Fig. 2D). In wild type embryos, *Gb-Dll* is expressed strongly in the

presumptive tarsus of all three developing T1, T2 and T3 legs (Fig. 2D). In the tibia, however,

*Gb-Dll* is expressed differently in T1/T2 and T3 legs. Specifically, in T3 there is a sharply

defined border between a distal domain of high expression and a proximal domain of lower

expression, but T1 and T2 lack such a border and display moderate proximal expression as well

as strong distal expression in (Niwa et al., 1997) (Fig. 2D). In *Gb-Ubx$^{CRISPR}$* embryos, *Gb-Dll*

expression in the presumptive tibia of the T3 leg lacked a strong boundary between high and low

tibial expression levels, and instead resembled the expression pattern of wild type T1 or T2 legs

(Fig. 2D). In addition, the T3 leg-specific pattern of multiple leg-patterning genes was absent in

*Gb-Ubx$^{CRISPR}$* embryos, which is consistent with disruption of the *Gb-Ubx* locus (Supplementary

Matsuoka, Nakamura et al.

234    Fig.1). Taken together, these results suggest that the CRISPR/Cas9 system induced mutations

235    specifically at the *Gb-Ubx* locus, which disrupted *Gb-Ubx* function.

*In-depth analysis of mutagenesis profile for the CRISPR/Cas9 system in G. bimaculatus*

238    To optimize the genome editing procedure, we wished to evaluate whether and how the timing of

239    injection affected NHEJ mutagenesis. For detailed assessment of this mutagenesis, we therefore

240    performed in-depth analysis of the CRISPR mutants using next generation sequencing.

241            Our previous study had revealed early cellular dynamics during cricket embryogenesis

(Nakamura et al., 2010), allowing us to assess whether specific mutagenesis events were

correlated with cellular behaviors during early development. As in *D. melanogaster* (Foe and

Alberts, 1983) early mitotic divisions in *G. bimaculatus* embryos are syncytial, meaning that

mitosis takes place without cytokinesis, resulting in multiple energids (nuclei surrounded by

aqueous cytoplasm but lacking a unique lipid bilayer) within a single cell membrane (Donoughe

et al. 2021; Donoughe and Extavour, 2016; Nakamura et al., 2010; Sarashina et al., 2003). To

evaluate whether and how the timing of injection affected mutagenesis outcomes, we chose four

early embryonic time points following the one-hour (1h) embryo collection period, as follows

(Supplementary Fig. 2A): (1) At the 1h injection time point, energids start to migrate from the

center of the egg to the cortex. (2) At the 3h injection time point, energids continue to become

distributed throughout the yolk, accompanied by mitotic cycles. (3) At the 5h injection time

point, energids have become nearly uniformly distributed throughout the egg cortex and begin

tangentially oriented nuclear division. (4) The 9h injection time point is one to eight hours before

cellularization (Donoughe and Extavour, 2016). We co-injected 0.5 μg/μl of the *Gb-Ubx* sgRNA,

the *Gb-lac2* sgRNA or an sgRNA targeting *abdominal-A* (*Gb-abd-A*); see "*Knock-in of donor*

Matsuoka, Nakamura et al.

*vector sequence at the Gb-abd-A locus*" below) described above, and 1 μg/μl of Cas9 mRNA into

the eggs at each of these time points. At 5d AEL, we isolated genomic DNA from three

individual embryos for each injection time point and used it for amplicon sequencing. We

examined the sequences of the on-target site and of the single highest predicted potential off-

target site for each of the *Gb-Ubx* and *Gb-abd-A* genes. For each sample we performed amplicon

sequencing with three replicates.

For *Gb-Ubx* on-target gene disruptions, we found that the rate of NHEJ-induced

mutations decreased with the age of the embryo at injection (Supplementary Fig. 2B). The rate of

NHEJ-induced mutations at the studied off-target site was less than 1.3% for all injection time

points (Supplementary Fig. 2C), suggesting that off-target effects may be minimal in this system.

The same trend was also observed for *Gb-abd-A* on-target mutations (Supplementary Fig. 2D-E;

see section "*Targeted mutagenesis of the Gb-abd-A locus via knock-in*" below). This result is

well correlated with the phenotypic severity observed in the $G_0$ hatchlings emerging from the

*Gb-lac2^CRISPR* embryos. *Gb-lac2^CRISPR* embryos injected at the two earlier time points (1h and 3h)

gave rise to hatchlings with broad patches of white cuticle (Supplementary Fig. 2H). In contrast,

the embryos injected at 5h showed milder phenotypes (Supplementary Fig. 2H) and the embryos

injected at 9h showed only little detectable phenotype (Supplementary Fig. 2H).

*Knock-in of donor vector sequence at the Gb-Ubx locus*

In addition to targeted sequence deletions, targeted sequence knock-in is a highly desirable

technique that would expand our ability to understand the functions of genomic regions of

interest. We had previously attempted to achieve targeted gene knock-ins through homology-

dependent repair, but this method has not worked in *G. bimaculatus* in our hands to date (data

Matsuoka, Nakamura et al.

not shown). In a homology-independent knock-in method reported for *D. rerio* and *D.*

*melanogaster* (Auer et al., 2014; Bosch et al., 2019), both genome and donor vector are cleaved

*in vivo*, then the cut ends of genome and donor vector are combined through NHEJ. This method

is more efficient than the homology-dependent method, potentially because NHEJ is highly

active throughout the cell cycle in eukaryotes (Hagmann et al., 1998). However, due to the nature

of NHEJ, the orientation of integration of the donor vector sequence cannot be controlled. In

addition, indel mutations are generated at the junction point. To try to circumvent these issues,

which might otherwise prevent functional knock-in, we generated a donor vector containing an

autonomous expression cassette comprising the *G. bimaculatus actin* (*Gb-act*) promoter

followed by the *eGFP* coding sequence (Nakamura et al., 2010). As a sgRNA recognition site,

we included a partial *DsRed* gene sequence (Auer et al., 2014), which is native to the coral

*Discosoma sp.* (Baird  et al., 2000) and not present in the cricket genome. We predicted that

successful knock-in of this donor sequence into the genome would result in eGFP expression

being driven by the *Gb-act* promoter regardless of the insert's orientation or any potential

induced indel mutations. To try to further increase the utility of this tool to facilitate

identification of targeted gene disruptions, we targeted knock-in of the donor sequence to an

exon of the target gene, which we anticipated would result in disruption of target gene function.

Our goal was to be able to identify such successfully knocked-in individuals by detectable eGFP

expression in the known expression domains of the target gene.

We chose the *Gb-Ubx* locus for this targeted knock-in strategy and used the same

sgRNA as that used for the knock-out experiment described above (see section "*Targeted*

*mutagenesis of the Gb-Ubx locus via knock-out*", Fig. 2A). We co-injected 50 ng/µl of sgRNA

for the *Ubx* locus, 50 ng/µl of sgRNA for the donor vector, 100 ng/µl of Cas9 mRNA, and 100

Matsuoka, Nakamura et al.

ng/µl of donor vector into fertile cricket eggs. By seven days after injection, four out of 85

injected embryos (4.7%) showed mosaic eGFP expression in the T3 trunk and leg (Fig. 3B'). Of

the 85 injected embryos, 18 individuals (21.2%) grew to adulthood. We crossed them

individually with wild type counterparts of the opposite sex and evaluated eGFP expression in

their offspring. One out of the 18 $G_0$ crickets (5.6%) produced $G_1$ embryos with eGFP expression

in a pattern identical to that of *Gb-Ubx* (Fig. 3C'; Barnett et al., 2019; Matsuoka et al., 2015;

Zhang et al., 2005). The eGFP expression was detectable through the eggshell even at late

embryonic stages (Fig. 3C'). At adult stages, $G_1$ knock-in crickets showed detectable eGFP

expression in the hind wing and T3 legs (Supplementary Fig. 3A' and B').

To confirm the integration of donor sequence into the genome, we performed PCR and

sequence analysis. We designed specific primers for each 5' and 3' junction point (a and b in Fig.

3A). All three examined embryos showed the expected amplicon size for both junctions (Fig.

3G), suggesting that at least two copies of the donor vector fragment were integrated into the

genome. Sequence analysis further confirmed the integration of the donor plasmid into the

genome, and that indel mutations were generated at each junction (Fig. 3H). To determine how

many copies of donor sequence were integrated into the genome, we performed copy number

estimation by quantitative RT-PCR. The copy number of the donor plasmid was estimated as the

copy ratio of the *eGFP* gene to that of the endogenous *orthodenticle* gene (*Gb-otd*), which is

known to have only one copy in the genome (Nakamura et al., 2010; Ylla et al., 2021). The

results of this analysis indicated that three copies of the donor plasmid were likely integrated into

the genome (Supplementary Fig. 4A).

To determine whether the function of the target gene was indeed disrupted by this

knock-in/knock-out strategy, we examined eGFP expression and morphology in $G_2$ *Gb-*

*Ubx$^{CRISPR-KI}$* embryos. Among the G$_2$ *Gb-Ubx$^{CRISPR-KI}$* embryos, we found they displayed one of

two different intensities of eGFP expression (Fig. 3E' and 3F'). We determined their genotype by

quantitative RT-PCR, and found that the individuals with strong eGFP expression were

homozygous mutant, and the individuals with weak eGFP expression were heterozygotes

(Supplementary Fig. 4B). Consistent with the genotyping result, crickets with weak eGFP

expression displayed no detectable morphological abnormalities (Fig. 3E and 3E'). Crickets with

strong eGFP expression, however, had smaller T3 legs and formed leg-like structures rather than

pleuropodia on the A1 segment (Fig. 3F and 3F') These phenotypes, which were the same as

those observed in the *Gb-Ubx$^{CRISPR}$* homozygous mutant (Fig. 2C and 2D), further suggested that

the strong eGFP-expressing crickets were homozygous mutants.

*Knock-in of donor vector sequence at the Gb-abd-A locus*

To confirm the efficiency and utility of this method, we next chose the Hox gene *Gb-abdominal-*

*A* (*abd-A*) as a target (Table 1). We designed sgRNAs for the sequence within the exon just

upstream of the homeodomain (Fig. 4A). We co-injected 50 ng/µl of sgRNA for the *Gb-abd-A*

locus, 50 ng/µl of sgRNA for the donor vector, 100 ng/µl of Cas9 mRNA, and 100 ng/µl of donor

vector into fertilized cricket eggs.

Of 38 injected G$_0$ embryos, five showed mosaic eGFP expression in the abdomen (Fig.

4B). Four G$_0$ adults were individually backcrossed with wild type counterparts of the opposite

sex to obtain multiple G$_1$ crickets. We obtained one stable transgenic line, in which eGFP

expression in G$_2$ embryos was similar to the previously documented expression pattern of *Gb-*

*abd-A* transcript (Barnett et al., 2019; Matsuoka et al., 2015; Zhang et al., 2005) (compare Fig.

4B with Supplementary Fig. 1K). In a replicate injection experiment, we obtained a second such

Matsuoka, Nakamura et al.

transgenic line (Table 3). PCR and sequence analysis confirmed that one of the two lines

contained the plasmid fragment in the sense orientation, and the second line contained the

plasmid fragment in the antisense orientation (Fig.4C). Copy number estimation analysis results

suggested that a single plasmid fragment was integrated into the genome in each line

(Supplementary Fig.4B).

In the $Gb$-$abd$-$A^{KI\text{-}exon}$ lines, eGFP expression was detectable in nymphs even through

the cuticle (Supplementary Fig. 3C', D'). We further detected eGFP expression in adult male and

female internal organs (Supplementary Fig. 5). In wild type females, a pair of ovaries, each

comprising hundreds of ovarioles, is located in the anterior abdomen (Nandchahal, 1972).

Mature eggs are stored in an egg chamber at the posterior of each ovariole, and eggs are

subsequently moved posteriorly through the oviduct. The posterior end of the oviduct is

connected to the uterus, where fertilization takes place, located at the base of the ovipositor

(Supplementary Fig. 5C''). In $Gb$-$abd$-$A^{KI\text{-}exon}$ mutant females, eGFP expression was detected in

the posterior portion of the oviduct (compare Supplementary Fig. 5B' and B'' to Supplementary

Fig.5A'). We found, however, that the oviduct and uterus were not connected in $Gb$-$abd$-$A^{KI\text{-}exon}$

mutant females, and these females were not able to lay eggs.  $Gb$-$abd$-$A^{KI\text{-}exon}$ mutant females

also generated ectopic ovipositors (Supplementary Fig. 6U). In $Gb$-$abd$-$A^{KI\text{-}exon}$ males, ubiquitous

eGFP expression was detected throughout the testis (compare Supplementary Fig. 5E' and E'' to

Supplementary Fig. 5D'). The observed eGFP expression in females is reminiscent of the

expression pattern of $D.$ $melanogaster$ $abd$-$A$ in the developing female genital disc (which gives

rise to the somatic reproductive structures including the oviduct in this fruit fly (Epper, 1983;

Sánchez and Guerrero, 2001), and in the adult oviducts (Foronda et al., 2006).

We also detected eGFP expression at the anterior tip of the testis in $Gb$-$Ubx^{KI\text{-}exon}$ adult

Matsuoka, Nakamura et al.

males (Supplementary Fig. 5G). *abd-A* expression has not, to our knowledge, been previously

detected in the *D. melanogaster* male genital disc (Freeland and Kuhn, 1996), which gives rise to

male somatic reproductive structures. However, high-throughput sequencing data from the

modENCODE project do report *abd-A* expression in the adult *D. melanogaster* testis (Brown et

al., 2014).

To confirm whether the eGFP expression observed in *Gb-abd-A$^{KI-exon}$* and *Gb-Ubx$^{KI-}$*

*$^{exon}$* animals reflects the endogenous expression of target genes in a given tissue, we performed

quantitative RT-PCR on eGFP-positive and eGFP-negative tissues. In a previous study, we

performed an RNA-seq analysis using *G. bimaculatus* adult brain and gonad tissue from both

sexes (Whittle et al. 2021). In this dataset, we did not detect expression of *Gb-Ubx* nor *Gb-adb-A*

in the brain (Fig. S7A, B). We therefore used brain as a negative control tissue for the expression

of *Gb-Ubx* and *Gb-abd-A*. We found a clear correlation between *eGFP* and *Gb-abd-A* expression

levels in the oviduct (eGFP-positive) and the uterus (eGFP-negative) in *Gb-abd-A$^{KI-exon}$* females

(Supplementary Fig. 7C and D). In the adult testis of *Gb-abd-A$^{KI-exon}$* males, where we observed

strong *eGFP* expression, we confirmed that the expression of *eGFP* and *Gb-abd-A* was also well

correlated (Supplementary Fig. 7E). These results indicate that *eGFP* expression correlates well

with *Gb-abd-A* expression, and are consistent with the interpretation that *eGFP* expression in

these knock-in crickets reflects target gene expression.

In contrast, we observed no correlation between *Gb-Ubx* and *eGFP* expression levels

in either eGFP-positive (anterior tip of testis) or eGFP-negative (rest of the testis) tissues, even

though *eGFP* expression levels showed a clear correlation between those tissues (Supplementary

Fig. 7F and G). It might be possible that *Gb-Ubx* is expressed at low levels throughout the ovary

and that we were therefore unable to detect *eGFP* expression in this organ except for at the tips

Matsuoka, Nakamura et al.

of ovaries. We found that the *Gb-Ubx* expression level in the wild type ovary and ovary tip was

comparable to the level detected in the brain; in other words, it was effectively undetectable

(Supplementary Fig. 7H). These results suggest that *Gb-Ubx* is not expressed, or is expressed

only at levels undetectable in ou transcriptome (Whittle et al. 2021), at the tip of wild type

ovaries, even though eGFP expression was detected there in *Gb-Ubx*$^{KI-exon}$ adults (Supplementary

Fig. 5G). Taking into account the fact that three copies of the donor sequence were integrated in

the *Gb-Ubx*$^{KI-exon}$ line (Supplementary Fig. 4), it might be possible that the observed eGFP

expression in the tip of the ovary from *Gb-Ubx*$^{KI-exon}$ was an artifact due to the multiple copies of

the expression cassette integrated at the *Gb-Ubx* locus, rather than being reflective of a true wild

type *Gb-Ubx* expression domain. It is also possible that the *Gb-act* promoter used in the

expression cassette might be more or differently active compared to the endogenous *Gb-Ubx*

promoter. The fact that the *Gb-Ubx*$^{KI-exon}$ line did not show any abnormality in ovary

development is consistent with this notion. Taken together, we conclude that it is possible to

visualize target gene expression by knocking an expression cassette into a target gene locus, but

careful verification should be done before making a conclusion as to all domains and levels of

target gene expression.

*Targeted insertion of an expression cassette into an intron of the Gb-abd-A locus*

Kimura and colleagues (Kimura et al., 2014) demonstrated that in *D. rerio* the homology

independent method could be applied for trapping endogenous enhancer activity by inserting a

donor sequence containing an expression cassette into the 5'UTR of genes of interest. We aimed

to apply this technique to *G. bimaculatus* by attempting to knock-in a donor vector into the

intronic region of *Gb-abd-A* (Fig. 5A).

Matsuoka, Nakamura et al.

We co-injected an sgRNA against an intron of *Gb-abd-A*, together with all other

relevant reagents as described above (sgRNA against the donor vector, donor vector, and Cas9

mRNA) (Fig. 5A). Of 100 injected eggs, two eggs showed mosaic expression of eGFP in the

abdomen (Fig. 5B). When the donor sequence was inserted into an exon in the previous

experiment (see "*Knock-in of donor vector sequence at the Gb-abd-A locus*" above), eGFP

expression was accompanied by a phenotype of ectopic leg-like structure development on

abdominal segments (Fig.4B; Supplementary Fig. 6), as previously observed in *Gb-abd-A* RNAi

experiments (Barnett et al., 2019). However, when the plasmid fragment was inserted into an

intron, the region expressing eGFP did not generate ectopic leg-like structures in $G_0$ embryos

(Fig. 5B). This apparent absent or minimal loss-of-function phenotype in the intron knock-in

embryos might explain the relatively high survival rate of the intron-targeted $G_0$ embryos (28%

of injected $G_0$ embryos survived to adulthood) compared to that of the exon-targeted $G_0$ embryos

(4.9% to 8.5% of 41 and 47 injected embryos survived to adulthood; Table 3).

We obtained one *Gb-abd-A*$^{KI-intron}$ line, in which we confirmed that one donor vector

sequence was likely integrated into the target region in a forward orientation (Fig. 5C). We

carefully inspected the morphology of the *Gb-abd-A*$^{KI-intron}$ adult crickets to assess whether

potential post-embryonic functions of the target gene were affected by insertion of the donor

sequence into an intron. $G_1$ heterozygous *Gb-abd-A*$^{KI-intron}$ females did not show the

supernumerary ovipositors observed in $G_1$ heterozygous *Gb-abd-A*$^{KI-exon}$ female adults

(Supplementary Fig .6W compared with 6U), and they laid eggs normally (data not shown). For

further confirmation, we examined the morphology of $G_2$ homozygous *Gb-abd-A*$^{KI-intron}$ mutants.

Approximately 25% of examined $G_2$ eggs showed strong eGFP expression, which we interpret as

likely indicative of a homozygous mutant. All $G_2$ embryos with strong eGFP expression

Matsuoka, Nakamura et al.

generated leg-like structures on the abdomen (Fig. 5B), suggesting that the function of the target

gene was somewhat affected in the homozygous condition, unlike in the mosaic condition

exhibited by $G_0$ embryos (Fig. 5B). Thus, even though the knock-in into an intron somewhat

affected target gene expression. $G_0$ embryos with intronic insertions did not show abnormalities,

unlike the embryos with exonic insertions. This increases the developmental success of intronic

knock-in embryos, and accordingly, the possibility of establishing a transgenic line.

**Discussion**

In the present study, we demonstrated that targeted knockout and knock-in by using the

CRISPR/Cas9 system works efficiently in the cricket *G. bimaculatus*. We performed functional

analysis of CRISPR/Cas9-induced mutations in the Hox genes *Gb-Ubx* and *Gb-abd-A* during

embryogenesis and at post-embryonic stages. We found that the cleavage efficiency of the

CRISPR/Cas9 system was much higher than that previously reported for artificial nucleases in

this cricket (Watanabe et al., 2012; Watanabe et al., 2014). We demonstrated that gene knock-in

via a homology-independent method is effective in this cricket, and successfully applied it to

functional analysis of Hox genes by knocking a donor sequence into an exon of the target gene to

disrupt the function of the target gene (knock-in/knock-out). In addition, we succeeded in

trapping endogenous gene activity using this method and revealed a number of new expression

domains that had not been previously observed with traditional methods (Barnett et al., 2019;

Matsuoka et al., 2015; Zhang et al., 2005). This homology-independent method is technically

simpler than the homology-dependent methods, as the donor plasmid does not need to be newly

made for each target region.

Matsuoka, Nakamura et al.

*CRISPR/Cas9 system vs RNA interference*

Delivering the proper amount of genome editing constructs at the proper time is important for

efficient outcomes. For example, gene editing in embryos injected around 3h AEL, at which

stage energids are distributing throughout the yolk, was highly efficient, but both detectable

mosaic phenotypes (of *Gb-lac2* crispants) and the rate of NHEJ decreased with later injections at

blastoderm stages (Supplementary Fig. 2B, D, and H). Thus, we suggest that optimization of

delivery conditions can be achieved by using pigmentation genes as an index.

In our study, reproducibility and severity of phenotypes generated with CRISPR/Cas9

were greater than those obtained with RNAi (Fig. 2D and 2F). We speculate that the efficiency of

RNAi-mediated knockdown may be influenced by when and at what levels the target gene is

expressed. In the case of *Gb-abdA* and *Gb-Ubx*, these genes are expressed much later in

development (embryonic stage 5, about 1.5 days after egg laying) than the stage at which we

performed injections (within 3 hours after egg laying) (Matsuoka et al., 2015, Donoughe et al.,

2016. Correspondingly, much higher concentrations of dsRNA proved necessary to produce even

mild phenotypes (5-6 µg/µl; Fig. 2E), than those typically used for RNAi against most genes in

this cricket (1-2 ug/ul; e.g. Donoughe et al., 2014). In contrast, indel mutations generated at

earlier stages by genome editing techniques resulted in clearly detectable, severe phenotypes

(Supplementary Fig. 2). Several studies have demonstrated that genome editing techniques can

sometimes be adequate for functional analysis of target genes in mosaic $G_0$ individuals (e.g.

Daimon et al., 2015; Martin et al., 2016; Matsuoka and Monteiro, 2018). However, it is often

difficult or impossible to unambiguously identify mutant cells in such mosaics. In this regard,

Matsuoka, Nakamura et al.

establishment and maintenance of stable mutant lines as performed herein, allows for less

ambiguous phenotypic analysis in this cricket.

Although the CRISPR/Cas9 system is efficient, it offers little to no conditionality,

which can complicate study of the many genes that act pleiotropically during development

(Minelli, 2016). For example, in the case of *Gb-abd-A*, the gene acts to repress leg formation in

the abdomen at embryonic stages (Supplementary Fig. 6), while at adult stages, it regulates

proper development of female genitalia (Supplementary Fig. 7). Likely because of this latter

phenotype, we were unable to obtain homozygous *Gb-abdA^{CRISPR}* animals. To overcome this

problem, sophisticated genetic methods, like balancer chromosomes (Miller et al., 2019), will

need to be developed in the future. In this regard, RNAi offers more options for conditional

control of gene function. By controlling the timing of injection of dsRNA, target gene activity

can be knocked down at any desired developmental stage in *G. bimaculatus* (Dabour et al., 2011;

Nakamura et al., 2008; Takahashi et al., 2009). Thus, while the CRISPR/Cas9 system is a

powerful new tool for gene function analyses, RNAi remains a useful technique for this system.

*Application of homology-independent knock-in method for functional analysis of endogenous*

*genes*

Homology-independent knock-in methods will expand our ability to analyze the function of

target genes in this hemimetabolous insect model. Here, we demonstrated one such application,

the KI/KO method, which allows for isolation of mutants without PCR-based genotyping. When

analyzing mutant phenotypes, affected individuals must typically be distinguished either by their

morphology or by molecular methods to detect changes in target gene product levels or

functions. In the case of the *Gb-Ubx* mutant, we would have needed to distinguish subtle

Matsuoka, Nakamura et al.

differences in the T3 and A1 embryonic segments (Fig. 2C), requiring destructive sampling.

Moreover, antibodies against target genes may not be routinely available in many cases. The

KI/KO method allows us to distinguish mutant individuals based on marker gene expression.

Even heterozygous and homozygous mutants can sometimes be distinguished based on the

intensity of marker gene expression. A similar strategy was employed in mosquitos via HDR

(McMeniman et al., 2014). In the present study, we could easily identify the eGFP expression

resulting from the KI/KO event because it matched the previously characterized expression

pattern for *Gb-Ubx* (Barnett et al., 2019; Matsuoka et al., 2015; Zhang et al., 2005). However, for

target genes with previously uncharacterized expression domains, analysis may be more

complex.

The promoter used in all expression cassettes herein is the same one used in a previous

study to drive ubiquitous constitutive expression (Nakamura et al., 2010). Nevertheless, all

knocked-in lines showed an eGFP expression pattern that was spatially and temporally restricted

like that of the target gene. We speculate that the promoter in the expression cassette acts as a

minimal promoter, and that the observed eGFP expression results from trapping endogenous

enhancer activity. The eGFP expression was not caused by the fusion to the endogenous gene

product, since both the line containing an inverted orientation of the donor sequence, and the

knock-in line targeting an intronic region, showed similar eGFP expression patterns. To enhance

the usefulness of this method, identification and use of a ubiquitous and strong promoter could in

principle drive exogenous marker gene expression in the whole embryo without being subject to

positional effects.

A remarkable feature of the homology-independent knock-in method is the length of

sequence that can be integrated. In case of knock-in through HDR, a few kb of sequence can be

integrated into the genome in arthropods (Gilles et al., 2015; McMeniman et al., 2014). In this

study, through NHEJ, at least six kb of plasmid sequence was integrated into the genome.

Furthermore, in some cases, three copies of plasmid sequence were integrated in tandem into the

genome (Supplementary Fig. 4). In this case, we speculate that first the donor plasmids were

digested and combined via NHEJ, and then the combined fragment was knocked in into the

genome via NHEJ, suggesting that homology-independent knock-in might be able to integrate

several tens of kb of sequence into the genome. For example, one study showed that a 200 kb

BAC vector could be integrated into a rodent genome through a similar strategy (Yoshimi et al.,

2016). This method might therefore be used for direct functional comparison of genomic regions

by exchanging homologous regions between related species of interest.

542           The efficiency of knock-in through NHEJ is high, but to improve its feasibility as a

technique for functional genetic analysis in this cricket, future studies may be able to further

enhance efficiency by optimizing at least one of two parameters, as follows: (1) *Enhance*

*expression level of a fluorescent cassette*. Empirically, $G_0$ crickets showing mosaic eGFP

expression tend to transmit their knocked-in transgene to their offspring. To increase the

efficiency of obtaining knock-in lines, future efforts should therefore focus on increasing the

number of mosaic marker gene expression cassettes in $G_0$ embryos. We also observed that some

expression cassettes seemed to show higher expression levels than others and were therefore

easier to detect in mosaic $G_0$s. Inclusion of inducible expression elements such as a heat shock

promoter or a modified Gal4/UAS system might help to enhance the activity of the expression

cassette. However, enhancement of fluorescent cassette expression may cause artificial or ectopic

expression as we observed in the ovaries of *Gb-Ubx$^{KI-exon}$* (Supplementary Fig. 5G''). (2)

*Introduce insulator sequences into the donor cassette*. Positional effects might also in principle

Matsuoka, Nakamura et al.

prevent the full potential activity of the expression cassette. In this study, our vector plasmid

contained insulators of the sea urchin *Hemicentrotus pulcherrimus* arylsulfatase gene (Takagi et

al., 2011) on either side of the expression cassette (see Materials and Methods), but we

nonetheless detected eGFP expression in a pattern matching that of the target gene. To achieve

more effective insulation, future studies might evaluate several different combinations of

insulator orientations, which can affect insulator activity (Tchurikov et al., 2009). Alternatively,

other insulators such as that of the *gypsy* retrotransposon (Modolell et al., 1983) might be

additional options for future optimization (Carballar-Lejarazú et al., 2013).

In conclusion, we provide evidence that the CRISPR-Cas9 system works well for both

knock-in and knock-out in the cricket *G. bimaculatus*. In depth analysis of CRISPR-Cas9-

induced mutations revealed optimized injection timing. In addition, we succeeded in the targeted

functional knock-in of exogenous sequences into the genome through NHEJ, resulting in

expression-tag reporter lines for target genes.

**Materials and Methods**

*Cricket husbandry*

All adult and juvenile *Gryllus bimaculatus* were reared in plastic cages at 26–30 °C and 50%

humidity under a 10-h light, 14-h dark photoperiod. They were fed on artificial fish food (Tetra)

or Purina cat food (item #178046). For microinjections, fertilized eggs were collected on a wet

kitchen towel in a plastic dish and incubated at 28 °C as previously described (Barry et al., 2019;

Watanabe et al., 2017).

Matsuoka, Nakamura et al.

*Construction of sgRNA vectors*

sgRNA target sequences were designed and their off-target sites were predicted with the CasOT

program (Xiao et al., 2014). From the suggested candidates, we selected target sequences having

high GC content (around 70%), beginning with guanidine for efficient *in vitro* transcription by

the T7 promoter, and with off-target sites containing at least 3 mismatches as per Ren and

colleagues (Ren et al., 2014). Two synthetic oligonucleotides (5'-ATAG-$N_{19}$-3' and 5'-AAA-$N_{20}$)

were annealed and inserted into the Bsa□ site of the modified pDR274 vector (Addgene plasmid

#42250) to expand its utility (the $GGN_{18}NGG$ sequence was present in the original pDR274,

while in the modified vector, a $GN_{19}NGG$ sequence was used). We confirmed insertion by

Sanger sequence analysis.

*Synthesis of sgRNA and mRNA*

For sgRNA synthesis, the template for *in vitro* transcription was digested from the vectors

generated as described above with Dra□. For Cas9 mRNA synthesis, the template for *in vitro*

transcription was digested from pMLM3613 (Addgene catalogue #42251) with Pme□. Both

sgRNA and Cas9 mRNA were *in vitro* transcribed using mMESSAGE mMACHINE T7 Kit (Life

Technologies catalogue #AM1344), and purified by ethanol precipitation. For the Cas9 mRNA,

we attached a poly-A tail by using a poly-A tailing Kit (Life Technologies catalogue #AM1350).

The concentration of synthesized RNAs was estimated by NanoDrop and gel electrophoresis.

*Construction of donor plasmids*

The *eGFP*bait-2A-RFP donor plasmid was generated in a pUC57 vector by commercial artificial

Matsuoka, Nakamura et al.

composition (GeneScript). The *DsRed*bait-*G'act-eGFP* donor plasmid was generated based on

the *eGFP*bait-2A-RFP donor plasmid. First, 2A-RFP was digested using BglII and NotI. *Gb-act-*

*eGFP* was also digested from a pXL-BacII- G'act-*eGFP* vector and ligated to generate the

*eGFP*bait-G'act-*eGFP* vector. Then, we digested this *eGFP*bait vector using BglII and SacI. We

amplified          *DsRed*bait          with          primers          (5'          to          3')          DsRed_fwd:

GCTCAGATCTCTTGGAGCCGTACTGGAAC,                          and                          DsRed_rev:

GTACGAGCTCCATCACCGAGTTCATGCG. The amplicon was ligated to generate the

*DsRed*bait-G'act-*eGFP* donor plasmid. The *DsRed*bait-2×Ars_rev-G'act-*eGFP*-2×Ars_fwd

donor plasmid was generated based on the *DsRed*bait-G'act-*eGFP* donor plasmid. The Ars

insulator sequence ArsInsC from *H. pulcherrimus* (Takagi et al., 2011) was amplified from an

ArsInsC-containing plasmid (kind gift of Assoc. Prof. Naoaki Sakamoto, Hiroshima University,

Japan) and integrated on either side of the expression cassette in the donor plasmid.

*Microinjection*

Cas9 mRNA, sgRNA, and donor vectors were injected into 2-5h AEL cricket eggs. Cricket eggs

were aligned in a groove 0.7 mm deep and 0.7 mm wide made with 2% agarose in 1x phosphate-

buffered saline (PBS) using a custom mold as previously described (Barry et al., 2019; Watanabe

et al., 2017) and filled with 1xPBS. Needles for injection were made by pulling glass capillaries

with filament (Narishige catalogue # GD-1) with a pipette puller (Sutter Instrument catalogue #

P-1000IVF), using the following pulling program: (1) x3 Heat; 858, Pull; 0, Velocity; 15, Time;

250, Pressure; 500, and (2) x1 Heat; 858, Pull; 80, Velocity; 15, Time; 200. To minimize the

invasiveness of the injection, the tips of the pulled needles were sharpened and ground to a 20°

Matsuoka, Nakamura et al.

angle by using a Micro Grinder (Narishige catalogue # EG-400). Approximately 5 nl of solution

was injected into eggs with a Micro Injector (Narishige catalog # IM300). After injection, eggs

were moved to a fresh Petri dish and submerged in fresh 1xPBS containing 50 U/ml penicillin

and 50 μg/ml streptomycin (15070-063, Thermo Fisher), and incubated at 28°C. During the

incubation period, the 1xPBS with penicillin and streptomycin was replaced every day. We

observed fluorescent protein expression at the stages when the target gene was known to be

expressed. Genomic DNA was extracted from 7 d AEL eggs and adult T3 legs and used for

insertion mapping and sequence analyses. After two days of incubation, injected cricket eggs

were moved to wet filter paper in a fresh Petri dish for hatching.

*Detection of indel mutations*

After Cas9 nuclease digests a target sequence, the disrupted sequence is repaired by either the

NHEJ or the HDR cell machinery. To confirm a KO mutation, we searched for errors repaired by

the NHEJ pathway, which sometimes introduces or deletes nucleotides at the digested site during

the repair process. Since disruption or repair are unlikely to take place identically in all cells of

an injected $G_0$ embryo, $G_0$ animals are expected to contain heterogeneous sequences at the

CRISPR targeted site, and thus to be heterozygous for a putative Cas9-induced indel. To confirm

the activity of the sgRNAs, the Surveyor[TM] Mutation Detection Kit (Transgenomic,) was used.

This assay relies on a "surveyor" nuclease that can recognize and digest a heteroduplex DNA

structure. First, genomic DNA was extracted from whole eggs or part of the T3 leg by a phenol

chloroform method as previously described (Barry et al., 2019; Watanabe et al., 2017).

Subsequently, approximately 200 bp of the targeted region was amplified by PCR from genomic

DNA (Table 4). PCR conditions were optimized to reduce non-specific amplification or

Matsuoka, Nakamura et al.

smearing. To create the putative heterogeneous DNA structure for the nuclease assay, the PCR

product was heated to 98°C for five minutes, and then re-annealed by gradually cooling down to

30°C. Half of the PCR product was digested with the Surveyor[TM] nuclease, and the other half

was used as a negative control and incubated without the nuclease. Digestion was confirmed by

agarose gel electrophoresis. For sgRNAs that yielded indels in the target sequence, digest of the

PCR product by the Surveyor[TM] nuclease is expected to produce split fragments around the

CRISPR targeted site relative to the negative control; the latter should not be digested by the

nuclease and thus should remain intact and run at the same size as the original amplicon. Positive

PCR products were extracted from the gel, purified with the QIAquick Gel Extraction Kit

(Qiagen catalogue #28506), and sub-cloned into the pGEM-Teasy vector (Promega catalogue

#A1360) using TA-cloning. The vectors were used for Sanger sequence analysis.

*Amplicon sequence analysis*

After a 1h egg collection, eggs were incubated for the desired length of time at 28°C. We co-

injected 0.5 µg/µl sgRNA and 1 µg/µl Cas9 mRNA into fertilized cricket eggs after each of these

incubation periods. Five days after injection, genomic DNA was extracted individually from

three eggs from each of the four tested injection times; the latter analysis was performed in

biological triplicate. Amplicon sequence analysis was performed by using MiSeq (Illumina), and

the preparation of DNA libraries and sequencing reactions were performed according to the

manufacturer's instructions. We read ~10,000 reads for on-target regions and ~50,000 reads for

off-target regions. We chose off-target sites that were most similar to the target site relative to

other sequences in the genome. As we mentioned above about the criteria for selecting target site,

the off-target site has three mismatch sequences compared to the target site. The assembly of

Matsuoka, Nakamura et al.

output paired end reads was performed by using CLC Genomic Workbench (CLC Bio, QIAGEN

Digital Insights). The relative proportions of reads containing indels and substitutions in the

individual eggs were calculated with the online-tool CRISPResso (Pinello et al., 2016). We used

the Integrative Genomic Viewer (Broad Institute) for investigation of the distribution of indels

and substitutions (Thorvaldsdóttir et al., 2013).

*Insertion mapping*

Genomic DNA was extracted from eGFP-positive eggs of each line. Due to the specifics of this

knock-in method, two types of insertion of vector fragment (sense and antisense orientations)

would be expected to occur; we therefore performed PCR using primers designed against either

side of the putative junction. PCR was performed using target region-specific (upstream or

downstream of sgRNA recognition site) and donor vector-specific primers (sequence within

*eGFP* for forward integration and M13Fw for reverse integration). Primer sequences are listed in

Table 4. Positive PCR products were extracted from the gel, purified by using the QIAquick Gel

Extraction Kit (Qiagen catalogue #28506), and sub-cloned into the pGEM-Teasy vector

(Promega catalogue #A1360) using TA-cloning. The vectors were used for Sanger sequence

analysis.

*Embryo fixation, whole mount* in situ *hybridization, and immunohistochemistry*

Embryos were dissected in 1xPBS and fixed with 4% paraformaldehyde in 1xPBS + 0.1% Tween

(PBT) for 1h at 4℃. The fixed embryos were dehydrated stepwise in 25%, 50%, 75%, and 100%

methanol in 1xPBT with five minutes per wash. The dehydrated embryos were stored in 100%

methanol at -30℃. Whole-mount *in situ* hybridization with digoxigenin (DIG)-labeled antisense

Matsuoka, Nakamura et al.

RNA probes was performed as previously described (Niwa et al., 2000; Zhang et al., 2005).

Immunohistochemistry was performed as follows: Fixed embryos were rehydrated stepwise in

75%, 50%, and 25% solutions of methanol/ PBT and finally in 100% PBT for five minutes in

each solution. After blocking with 1% bovine serum albumin (BSA) (Thermo Fisher) in PBT for

one hour at room temperature, embryos were incubated with an anti-UbdA antibody FP6.87

(Kelsh et al., 1994) (Developmental Studies Hybridoma Bank) diluted 1:200 in 1% BSA/PBT

overnight at 4℃. After washing with PBT three times, embryos were incubated in 1% BSA/PBT

for one hour at room temperature, and then incubated with Alexa Fluor 488-conjugated Goat

Anti-mouse IgG(H+L) (Invitrogen catalogue #A32723) diluted 1:400 in 1% BSA/PBT for one

hour at 4℃. After washing the embryos with PBT once for 10-60 minutes, embryos were

counter-stained with DAPI (Sigma catalogue #10236276001) stock solution 1mg/mL diluted

1:1000 in PBT for ten minutes, and then washed with PBT two times for 10-60 minutes per

wash. PBT was then substituted with 25% and 50% glycerol/PBT to clear embryos for

microscopy.

*Copy number estimation and detection of endogenous Hox gene expression in eGFP-positive*

*tissues by using quantitative RT-PCR*

To estimate the number of plasmid fragments integrated into the genome via NHEJ events, we

performed quantitative RT-PCR using genomic DNA from individual 5 day-old embryos of wild

type, *Gb-Ubx$^{KI-exon}$*, and *Gb-abd-A$^{KI-exon}$* lines, and compared relative quantity values of the

inserted *eGFP* gene to an endogenous gene *Gb-otd*, which is present in a single copy in the *G.*

*bimaculatus* genome (Ylla et al., 2021). Genomic DNA was extracted from embryos using Cica

Matsuoka, Nakamura et al.

713 geneus® Total DNA Prep kit (for Tissue) (Kanto Chemical), according to the manufacture's

714 protocol. Real-time quantitative PCR was performed using the power SYBR Green PCR Master

715 Kit (Applied Biosystems catalogue #4368577) and an ABI 7900 Real Time PCR System

716 (Applied Biosystems) as described previously (Nakamura et al., 2008). Primer sequences are

717 listed in Table 4.

718   To examine whether the eGFP expression observed in gonads of *Gb-Ubx*$^{KI-exon}$ and *Gb-*

719 *abd-A*$^{KI-exon}$ reflects the endogenous expression of *Gb-Ubx* or *Gb-abd-A* in those tissues, we

720 performed quantitative RT-PCR using total cDNA from tissues with or without eGFP expression.

721 We dissected out gonads and brain in 1xPBS, and then transferred the dissected tissues to RNA

722 later (Thermo Fisher). Total mRNA was extracted with RNeasy MinElute Cleanup Kit (Qiagen),

723 according to the manufacture's protocol. Equal amounts of total RNA were used for reverse

724 transcription reaction with SuperScript III Reverse Transcriptase (ThermoFisher). Quantitative

725 RT-PCR was performed with THUNDERBIRD® SYBR® qPCR Mix (TOYOBO) and

726 LightCycler 96 (Roche). The expression levels of *eGFP, Gb-Ubx* and *Gb-abd-A* were normalized

727 to the level of *Gb-beta-tubulin.* Primer sequences used were as previously described (Barnett et

728 al., 2019).

Matsuoka, Nakamura et al.

**Acknowledgements**

We thank N. Sakamoto for the Ars plasmid, and the members of the Extavour and Mito labs for

discussion.

**Competing Interests**

No competing interests declared.

**Funding**

This work was supported by National Science Foundation award number IOS-1257217 and

funds from the Howard Hughes Medical Institute and Harvard University to CGE, an Overseas

Research Fellowship (grant # 693) from the Japan Society for the Promotion of Science (JSPS)

to TN, a Grant-in-Aid for Young Scientists B (grant # JP16K21199, JP26870415) from the JSPS

to TW, and a Grant-in-Aid for Scientific Research (B) (grant # JP26292176) from the JSPS to

TM.

**Data availability**

Not applicable.

Matsuoka, Nakamura et al.

**Tables**

**Table 1.** *G. bimaculatus* **genes in this study disrupted by targeted genome modification**

| Gene | Functions | Tissue distribution | Phenotype of genome-modified knockout/knockdown cricket | Refs |
|------|-----------|---------------------|--------------------------------------------------------|------|
| *Laccase 2* | Phenol oxidase, cuticle tanning (sclerotization and pigmentation) | No data | *Lac2* knock-out nymphs show defect in pigmentation. | (Watanabe et al., 2014); this study |
| *Ultrabithorax* | Enlargement of T3 leg Indentity of A1 pleuropodia | T3 and A1 segment | - *Ubx* knock-out embryos show partial transformation of A1 pleuropodia into T3 thoracic leg, and of T3 thoracic leg into T2 thoracic leg. <br> - Embryonic lethal. | This study |
| *abdominal-A* | Repression of leg formation in abdomen | Abdomen | - *abd-A* knock-out embryos show generation of leg-like structures on the abdomen. <br> - *abd-A* knock-out nymphs show fusion of abdominal segments. <br> - *abd-A* knock-out female adults generated ectopic ovipositors and had defects in oviducts and uterus attachment. | This study |

Matsuoka, Nakamura et al.

**Table 2. sgRNA sequences used in this study**

| sgRNA name | sgRNA sequence (5'->3') (**_Bold underline_** _indicates PAM sequence_) |
|---|---|
| _Gb-lac2_ exon | GGGGTCCTGGCCCGGGTTGA**CGG** |
| _Gb-Ubx_ exon | GGGTAGAAGGTGTGGTTGGC**GGG** |
| _Gb-Ubx_ intron | GGACTGGCCACGCTCCAAGG**AGG** |
| _Gb-abd-A_ exon | GGGGCAAGGCTCACCCGTGA**TGG** |
| _Gb-abd-A_ intron | GCTCGCGGTGTTTTACGGCT**GGG** |

Matsuoka, Nakamura et al.

**Table 3. Efficiency of CRISPR/Cas9-mediated genome editing in *G. bimaculatus*.** *nd = no data*

| | *Shown in Figure* | *# eggs injected* | *# injected embryos with eGFP expression or phenotype* | *# embryos developed by 7d AEL (% of injected embryos)* | *# hatched nymphs (% of embryos developed by 7d AEL)* | *# fertile adults (% of nymphs hatched)* | *% injected embryos yielding fertile adults* | *# fertile adults showing germ line transmission* | *% fertile adults showing germ line transmission* | *% injected embryos yielding fertile adults showing germline transmission* |
|---|---|---|---|---|---|---|---|---|---|---|
| *Gb-Lac2*[CRISPR] KO | 1 | 232 | nd | nd | nd | 38 (nd) | 16.4% | 18 | 47.4% | 7.8% |
| *Gb-Ubx*[CRISPR] KO | 2, S1, S2 | 167 | nd | nd | nd | 10 (nd) | 59.9% | 6 | 60.0% | 3.6% |
| *Gb-Ubx*[KI-exon] | 3, S3, S4 | 85 | 4 (4.7%) | 58 (68.2%) | 30 (51.7%) | 25 (83.3%) | 29.4% | 1 | 4.0% | 1.2% |
| *Gb-abd-A*[KI-exon] 1st trial | 4, S2, S3, S4, S5, S6 | 47 | 5 (10.6%) | 38 (80.8%) | 9 (23.7%) | 4 (44.4%) | 8.5% | 1 | 25.0% | 2.1% |
| *Gb-abd-A*[KI] exon 2nd trial | n/a | 41 | 0 (0.0%) | 36 (87.8%) | 22 (61.1%) | 2 (9.1%) | 4.9% | 1 | 50.0% | 2.4% |
| *Gb-abd-A*[KI-intron] | 5, S5, S6 | 100 | 2 (2.0%) | 77 (77.0%) | 47 (61.0%) | 28 (59.6%) | 28.0% | 1 | 3.6% | 1.0% |
| *Gb-Ubx*[KI-intron] | S7 | 73 | 5 (6.8%) | 62 (84.9%) | 59 (95.2%) | 22 (37.3%) | 30.1% | 2 | 9.1% | 2.7% |
| Control (Donoughe and Extavour, 2016) | n/a | 42 | n/a | 40 (95.2%) | 28 (70.0%) | nd | nd | n/a | n/a | n/a |
| Control (Ewen-Campen et al., 2012) | n/a | 78 | n/a | nd | 64 (82.0%) | nd | nd | n/a | n/a | n/a |

Matsuoka, Nakamura et al.

**Table 4. Primers used in this study.**

| Primer name | Primer seq (5'->3') |
| --- | --- |
| *Gb-lac2* genotype Fw | GCGCGACCCCAACAGCAACC |
| *Gb-lac2* genotype Rv | TGGCCCTTGCCGTTGATGAGC |
| *Gb-Ubx* exon genotype KO Fw | CGTTTGTGAAACGTATGGCCCGTTA |
| *Gb-Ubx* exon genotype KO Rv | GTCCCTGGGCTCCTGGAACACG |
| *Gb-Ubx* exon genotype KI genome Fw | AACACGTGCTCCCTCAACTC |
| *Gb-Ubx* exon genotype KI genome Rv | TGAAACGTATGGCCCGTTAT |
| *Gb-Ubx* exon genotype KI vector 5' Rv | GTCGCATGCTCCTCTAGACTCG |
| *Gb-Ubx* intron genotype Fw | GCAGAACCGTTTCATGAATGT |
| *Gb-Ubx* intron genotype Rv | ATTCTCGCCCTTATGCAGAG |
| *Gb-abd-A* exon genotype Fw | CCGATTCCATGGTGAACTA |
| *Gb-abd-A* exon genotype Rv | AGAACGGAACGCAGTGAGTTAG |
| *Gb-abd-A* exon genotype vector 3' | GAACTTCAGGGTCAGCTTGC |
| *Gb-abd-A* exon genotype vector 5' | CACAAGGCACAAATGCTCGT |
| *Gb-abd-A* intron Fw | CGGATCTATTCGGCCATTT |
| *Gb-abd-A* intron Rv | TCAAACGGATCTTCCTCTCG |
| *Gb-otd* qPCR Fw | CATTCACGTCTCCGCCATAC |
| *Gb-otd* qPCR Rv | GCTCCATCAACAGGCAAACA |
| *eGFP* qPCR Fw | CAGAAGAACGGCATCAAGGT |
| *eGFP* qPCR Rv | GGGTGCTCAGGTAGTGGTTG |
| *Gb-beta-actin* qPCR Fw | TTGACAATGGATCCGGAATGT |
| *Gb-beta-actin* qPCR Fw | AAAACTGCCCTGGGTGCAT |

Matsuoka, Nakamura et al.

## References

**Arakane, Y., Muthukrishnan, S., Beeman, R. W., Kanost, M. R. and Kramer, K. J.** (2005). *Laccase 2* is the phenoloxidase gene required for beetle cuticle tanning. *Proc National Acad Sci* **102**, 11337–11342.

**Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M., Davis, A. P., Dolinski, K., Dwight, S. S., Eppig, J. T., et al.** (2000). Gene Ontology: tool for the unification of biology. *Nat Genet* **25**, 25–29.

**Auer, T. O., Duroure, K., Cian, A. D., Concordet, J.-P. and Bene, F. D.** (2014). Highly efficient CRISPR/Cas9-mediated knock-in in zebrafish by homology-independent DNA repair. *Genome Res* **24**, 142–153.

**Aumann, R. A., Schetelig, M. F., and Häcker, I.** (2018) Highly efficient genome editing by homology-directed repair using Cas9 protein in *Ceratitis capitata*. *Insect Biochem Mol Biol* **101**, 85-93.

**Bai, Y., He, Y., Shen, C.-Z., Li, K., Li, D.-H. and He, Z.-Q.** (2023). CRISPR/Cas9-Mediated genomic knock out of tyrosine hydroxylase and yellow genes in cricket *Gryllus bimaculatus*. *PLoS ONE* **18**, e0284124.

**Baird, G. S., Zacharias, D. A. and Tsien, R. Y.** (2000). Biochemistry, mutagenesis, and oligomerization of DsRed, a red fluorescent protein from coral. *Proc National Acad Sci* **97**, 11984–11989.

**Bando, T., Ishimaru, Y., Kida, T., Hamada, Y., Matsuoka, Y., Nakamura, T., Ohuchi, H., Noji, S. and Mito, T.** (2013). Analysis of RNA-Seq data reveals involvement of JAK/STAT signalling during leg regeneration in the cricket *Gryllus bimaculatus*. *Development* **140**, 959–964.

**Barnett, A. A., Nakamura, T. and Extavour, C. G.** (2019). Hox genes limit germ cell formation in the short germ insect *Gryllus bimaculatus*. *Proc National Acad Sci* **116**, 16430–16435.

**Barry, S. K., Nakamura, T., Matsuoka, Y., Straub, C., Horch, H. W. and Extavour, C. G.** (2019). Injecting *Gryllus bimaculatus* Eggs. *J Vis Exp*.

**Bosch, J. A., Colbeth, R., Zirin, J. and Perrimon, N.** (2019). Gene Knock-Ins in Drosophila Using Homology-Independent Insertion of Universal Donor Plasmids. *Genetics* **214**, genetics.302819.2019.

**Branzei, D. and Foiani, M.** (2008). Regulation of DNA repair throughout the cell cycle. *Nat Rev Mol Cell Bio* **9**, 297–308.

**Brown, J. B., Boley, N., Eisman, R., May, G. E., Stoiber, M. H., Duff, M. O., Booth, B. W., Wen, J., Park, S., Suzuki, A. M., et al.** (2014). Diversity and dynamics of the *Drosophila* transcriptome. *Nature* **512**, 393–399.

**Carballar-Lejarazú, R., Jasinskiene, N. and James, A. A.** (2013). Exogenous *gypsy* insulator sequences modulate transgene expression in the malaria vector mosquito, *Anopheles stephensi*. *Proc National Acad Sci* **110**, 7176–7181.

Matsuoka, Nakamura et al.

**Connahs, H., Tan, E. J., Ter, E. T., Dion, E., Matsuoka, Y., Bear, A., and Monteiro, A.** (2022) The *yellow* gene regulates behavioural plasticity by repressing male courtship in *Bicyclus anynana* butterflies. *Proc Biol Sci* **289**, 20212665.

**Consortium, T. G. O., Carbon, S., Douglass, E., Good, B. M., Unni, D. R., Harris, N. L., Mungall, C. J., Basu, S., Chisholm, R. L., Dodson, R. J., et al.** (2020). The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids Res* **49**, D325–D334.

**Dabour, N., Bando, T., Nakamura, T., Miyawaki, K., Mito, T., Ohuchi, H. and Noji, S.** (2011). Cricket body size is altered by systemic RNAi against insulin signaling components and epidermal growth factor receptor. *Dev Growth Differ* **53**, 857–869.

**Daimon, T., Uchibori, M., Nakao, H., Sezutsu, H. and Shinoda, T.** (2015). Knockout silkworms reveal a dispensable role for juvenile hormones in holometabolous life cycle. *Proc National Acad Sci* **112**, E4226–E4235.

**Davis, G. K. and Patel, N. H.** (2002). Short, long, and beyond: Molecular and Embryological Approaches to Insect Segmentation. *Annual Review of Entomology* **47**, 669–699.

**Donoughe, S. and Extavour, C. G.** (2016). Embryonic development of the cricket *Gryllus bimaculatus*. *Dev Biol* **411**, 140–156.

**Donoughe, S., Nakamura, T., Ewen-Campen, B., Green, D. A., Henderson, L. and Extavour, C. G.** (2014). BMP signaling is required for the generation of primordial germ cells in an insect. *Proc National Acad Sci* **111**, 4133–4138.

**Donoughe, S., Hoffmann, J., Nakamura, T., Rycroft, C. and Extavour, C.G.** (2022). Nuclear speed and cycle length co-vary with local density during syncytial blastoderm formation in a cricket. *Nature Communications* **13**, 3889.

**Dossey, A. T., Oppert, B., Chu, F.-C., Lorenzen, M. D., Scheffler, B., Simpson, S., Koren, S., Johnston, J. S., Katoka, K. and Ide, K.** (2022). Genome and Genetic Engineering of the House Cricket (*Acheta domesticus*): A Resource for Sustainable Agriculture. *Biomolecules* **13**, 589.

**Epper, F.** (1983). The evagination of the genital imaginal discs of *Drosophila melanogaster*. *Wilhelm Roux's Archives Dev Biology* **192**, 275–279.

**Ewen-Campen, B., Srouji, J. R., Schwager, E. E. and Extavour, C. G.** (2012). *oskar* Predates the Evolution of Germ Plasm in Insects. *Curr Biol* **22**, 2278–2283.

**Foe, V. E. and Alberts, B. M.** (1983). Studies of nuclear and cytoplasmic behaviour during the five mitotic cycles that precede gastrulation in *Drosophila* embryogenesis. *Journal of Cell Science* **61**, 31–70.

**Foronda, D., Estrada, B., Navas, L. de and Sánchez-Herrero, E.** (2006). Requirement of *abdominal-A* and *Abdominal-B* in the developing genitalia of *Drosophila* breaks the posterior downregulation rule. *Development* **133**, 117–127.

**Freeland, D. E. and Kuhn, D. T.** (1996). Expression patterns of developmental genes reveal segment and parasegment organization of *D. melanogaster* genital discs. *Mech Develop* **56**, 61–72.

**Gilles, A. F., Schinko, J. B. and Averof, M.** (2015). Efficient CRISPR-mediated gene targeting and transgene replacement in the beetle *Tribolium castaneum*. *Development* **142**, 2832–2839.

Matsuoka, Nakamura et al.

**Goldstein, B. and King, N.** (2016). The Future of Cell Biology: Emerging Model Organisms. *Trends Cell Biol* **26**, 818–824.

**Gratz, S. J., Cummings, A. M., Nguyen, J. N., Hamm, D. C., Donohue, L. K., Harrison, M. M., Wildonger, J. and O'Connor-Giles, K. M.** (2013). Genome Engineering of *Drosophila* with the CRISPR RNA-Guided Cas9 Nuclease. *Genetics* **194**, 1029–1035.

**Hagmann, M., Bruggmann, R., Xue, L., Georgiev, O., Schaffner, W., Rungger, D., Spaniol, P. and Gerster, T.** (1998). Homologous Recombination and DNA-End Joining Reactions in Zygotes and Early Embryos of Zebrafish (*Danio rerio*) and *Drosophila melanogaster. Biol Chem* **379**, 673–682.

**Hammond, A., Galizi, R., Kyrou, K., Simoni, A., Siniscalchi, C., Katsanos, D., Gribble, M., Baker, D., Marois, E., Russell, S., Burt, A., Windbichler, N., Crisanti, A. and Nolan, T.** (2016). A CRISPR-Cas9 gene drive system targeting female reproduction in the malaria mosquito vector *Anopheles gambiae*. Nat Biotechnol **34**, 78-83.

**Hedwig, B. and Sarmiento-Ponce, E. J.** (2017). Song pattern recognition in crickets based on a delay-line and coincidence-detector mechanism. *Proc Royal Soc B* **284**, 20170745.

**Horch, H. Wi., Mito, T., Popadic, A., Ohuchi, H. and Noji, S.** (2017a). *The Cricket as a Model Organism, Development, Regeneration, and Behavior*.

**Horch, H., Liu, J., Mito, T., Popadić, A. and Watanabe, T.** (2017b). Protocols in the Cricket. In *The Cricket as a Model Organism* (ed. Horch, H.), Mito, T.), Popadic, A.), Hideyo), Ohuchi), and Noji, S.), pp. 327–370.

**Huis, A. van, Itterbeeck, J. V., Klunder, H., Mertens, E., Halloran, A., Muir, G. and Vantomme, P.** (2013). *Edible insects: Future prospects for food and feed security*. Rome: Food and Agriculture Organization of the United Nations.

**Kelsh, R., Weinzierl, R. O. J., White, R. A. H. and Akam, M.** (1994). Homeotic gene expression in the locust *Schistocerca*: An antibody that detects conserved epitopes in ultrabithorax and abdominal-A proteins. *Dev Genet* **15**, 19–31.

**Kimura, Y., Hisano, Y., Kawahara, A. and Higashijima, S.** (2014). Efficient generation of knock-in transgenic zebrafish carrying reporter/driver genes by CRISPR/Cas9-mediated genome engineering. *Sci Rep* **4**, 6545.

**Kistler, K. E., Vosshall, L. B. and Matthews, B. J.** (2015). Genome Engineering with CRISPR-Cas9 in the Mosquito *Aedes aegypti. Cell Reports* **11**, 51–60.

**Kohno, H., Kubo, T.** (2018). *mKast* is dispensable for normal development and sexual maturation of the male European honeybee. *Sci Rep*. **8**:11877.

**Kulkarni, A. and Extavour, C. G.** (2019). Chapter 8: The Cricket *Gryllus bimaculatus*: Techniques for Quantitative and Functional Genetic Analyses of Cricket Biology. In *Evo-Devo: Non-model Species in Cell and Developmental Biology* (ed. Twozydlo, W.) and Bilinski, S. M.), pp. 183–216.

**Kulski, J. K.** (2016). Next Generation Sequencing - Advances, Applications and Challenges.

**Li, X., Fan, D., Zhang, W., Liu, G., Zhang, L., Zhao, L., Fang, X., Chen, L., Dong, Y., Chen, Y., et al.** (2015). Outbred genome sequencing and CRISPR/Cas9 gene editing in butterflies. *Nat Commun* **6**, 8212.

Matsuoka, Nakamura et al.

**Ma, S., Chang, J., Wang, X., Liu, Y., Zhang, J., Lu, W., Gao, J., Shi, R., Zhao, P. and Xia, Q.** (2014). CRISPR/Cas9 mediated multiplex genome editing and heritable mutagenesis of *BmKu70* in Bombyx mori. *Sci Rep* **4**, 4489.

**Mahfooz, N. S., Li, H. and Popadić, A.** (2004). Differential expression patterns of the hox gene are associated with differential growth of insect hind legs. *Proc National Acad Sci* **101**, 4877–4882.

**Mahfooz, N., Turchyn, N., Mihajlovic, M., Hrycaj, S. and Popadić, A.** (2007). *Ubx* Regulates Differential Enlargement and Diversification of Insect Hind Legs. *Plos One* **2**, e866.

**Martin, A., Serano, J. M., Jarvis, E., Bruce, H. S., Wang, J., Ray, S., Barker, C. A., O'Connell, L. C. and Patel, N. H.** (2016). CRISPR/Cas9 Mutagenesis Reveals Versatile Roles of Hox Genes in Crustacean Limb Specification and Evolution. *Curr Biol* **26**, 14–26.

**Matsumoto, Y., Matsumoto, C. S. and Mizunami, M.** (2018). Signaling Pathways for Long-Term Memory Formation in the Cricket. *Front Psychol* **9**, 1014.

**Matsuoka, Y. and Monteiro, A.** (2018). Melanin Pathway Genes Regulate Color and Morphology of Butterfly Wing Scales. *Cell Reports* **24**, 56–65.

**Matsuoka, Y., Bando, T., Watanabe, T., Ishimaru, Y., Noji, S., Popadić, A. and Mito, T.** (2015). Short germ insects utilize both the ancestral and derived mode of Polycomb group-mediated epigenetic silencing of Hox genes. *Biol Open* **4**, 702–709.

**McMeniman, C. J., Corfas, R. A., Matthews, B. J., Ritchie, S. A. and Vosshall, L. B.** (2014). Multimodal Integration of Carbon Dioxide and Other Sensory Cues Drives Mosquito Attraction to Humans. *Cell* **156**, 1060–1071.

**Miller, D. E., Cook, K. R. and Hawley, R. S.** (2019). The joy of balancers. *Plos Genet* **15**, e1008421.

**Minelli, A.** (2016). Evo-Devo and Phylogenetics. In *Evolutionary Developmental Biology*, pp. 1–12.

**Mito, T. and Noji, S.** (2008). The Two-Spotted Cricket *Gryllus bimaculatus*: An Emerging Model for Developmental and Regeneration Studies. *Cold Spring Harb Protoc* **2008**, pdb.emo110-pdb.emo110.

**Mito, T., Inoue, Y., Kimura, S., Miyawaki, K., Niwa, N., Shinmyo, Y., Ohuchi, H. and Noji, S.** (2002). Involvement of *hedgehog*, *wingless*, and *dpp* in the initiation of proximodistal axis formation during the regeneration of insect legs, a verification of the modified boundary model. *Mech Develop* **114**, 27–35.

**Mito, T., Kobayashi, C., Sarashina, I., Zhang, H., Shinahara, W., Miyawaki, K., Shinmyo, Y., Ohuchi, H. and Noji, S.** (2007). *even-skipped* has gap-like, pair-rule-like, and segmental functions in the cricket *Gryllus bimaculatus*, a basal, intermediate germ insect (Orthoptera). *Dev Biol* **303**, 202–213.

**Mito, T., Ronco, M., Uda, T., Nakamura, T., Ohuchi, H. and Noji, S.** (2008). Divergent and conserved roles of *extradenticle* in body segmentation and appendage formation, respectively, in the cricket *Gryllus bimaculatus*. *Dev Biol* **313**, 67–79.

**Mizunami, M. and Matsumoto, Y.** (2017). Roles of Octopamine and Dopamine Neurons for Mediating Appetitive and Aversive Signals in Pavlovian Conditioning in Crickets. *Front*

Matsuoka, Nakamura et al.

*Physiol* **8**, 1027.

**Modolell, J., Bender, W. and Meselson, M.** (1983). *Drosophila melanogaster* mutations suppressible by the suppressor of Hairy-wing are insertions of a 7.3-kilobase mobile element. *Proc National Acad Sci* **80**, 1678–1682.

**Morin, X., Daneman, R., Zavortink, M. and Chia, W.** (2001). A protein trap strategy to detect GFP-tagged proteins expressed from their endogenous loci in *Drosophila*. *Proc National Acad Sci* **98**, 15050–15055.

**Nakamura, T., Mito, T., Bando, T., Ohuchi, H. and Noji, S.** (2007). Molecular and Cellular Basis of Regeneration and Tissue Repair. *Cell Mol Life Sci* **65**, 64.

**Nakamura, T., Mito, T., Miyawaki, K., Ohuchi, H. and Noji, S.** (2008). EGFR signaling is required for re-establishing the proximodistal axis during distal leg regeneration in the cricket *Gryllus bimaculatus* nymph. *Dev Biol* **319**, 46–55.

**Nakamura, T., Yoshizaki, M., Ogawa, S., Okamoto, H., Shinmyo, Y., Bando, T., Ohuchi, H., Noji, S. and Mito, T.** (2010). Imaging of Transgenic Cricket Embryos Reveals Cell Movements Consistent with a Syncytial Patterning Mechanism. *Curr Biol* **20**, 1641–1647.

**Nakamura, Y., Tomonari, S., Kawamoto, K., Yamashita, T., Watanabe, T., Ishimaru, Y., Noji, S. and Mito, T.** (2022). Evolutionarily conserved function of the even-skipped ortholog in insects revealed by gene knock-out analyses in *Gryllus bimaculatus*. *Dev Biol* **485**, 1–8.

**Nandchahal, N.** (1972). Reproductive organs of *Gryllodes sigillatus* (Walker) (Orthoptera: Gryllidae). *J Nat Hist* **6**, 125–131.

**Niwa, N., Saitoh, M., Ohuchi, H., Yoshioka, H., and Noji, S.** (1997) Correlation between *Distal-less* expression patterns and structures of appendages in development of the two-spotted cricket, *Gryllus bimaculatus*. *Zoological Science* **14**, 115-125.

**Niwa, N., Inoue, Y., Nozawa, A., Saito, M., Misumi, Y., Ohuchi, H., Yoshioka, H. and Noji, S.** (2000). Correlation of diversity of leg morphology in *Gryllus bimaculatus* (cricket) with divergence in dpp expression pattern during leg development. *Development* **127**, 4373–4381.

**Papageorgiou, L., Eleni, P., Raftopoulou, S., Mantaiou, M., Megalooikonomou, V. and Vlachakis, D.** (2018). Genomic big data hitting the storage bottleneck. *Embnet J* **24**, e910.

**Pinello, L., Canver, M. C., Hoban, M. D., Orkin, S. H., Kohn, D. B., Bauer, D. E. and Yuan, G. C.** (2016). Analyzing CRISPR genome-editing experiments with CRISPResso. *Nat Biotechnol* **34**, 695–697.

**Purusothaman, D. K., Shackleford, L., Anderson, M. A. E., Harvey-Samuel, T. and Alphey, L.** (2021). CRISPR/Cas-9 mediated knock-in by homology dependent repair in the West Nile Virus vector *Culex quinquefasciatus* Say. *Sci Rep* **11**, 14964.

**Qiu, P., Shandilya, H., D'Alessio, J. M., O'Connor, K., Durocher, J. and Gerard, G. F.** (2004). Mutation detection using Surveyor nuclease. *Biotechniques* **36**, 702–707.

**Rathke, H.** (1844). Entwickelungsgeschichte der Maulwurfsgrille (Gryllotalpa vulgaris). *Archiv für Anatomie, Physiologie und Wissenschaftliche Medicin* 27–37.

**Reding, K., Pick, L.** (2020). High-Efficiency CRISPR/Cas9 Mutagenesis of the white Gene in the Milkweed Bug *Oncopeltus fasciatus. Genetics* **215**:1027-1037.

**Ren, X., Yang, Z., Xu, J., Sun, J., Mao, D., Hu, Y., Yang, S.-J., Qiao, H.-H., Wang, X., Hu,**

**Q., et al.** (2014). Enhanced Specificity and Efficiency of the CRISPR/Cas9 System with Optimized sgRNA Parameters in *Drosophila*. *Cell Reports* **9**, 1151–1162.

**Rozen-Gagnon, K., Gu, M., Luna, J. M., Luo, J. D., Yi, S., Novack, S., Jacobson, E., Wang, W., Paul, M. R., Scheel, T. K. H., Carroll, T. and Rice, C. M.** (2021). Argonaute-CLIP delineates versatile, functional RNAi networks in *Aedes aegypti*, a major vector of human viruses. *Cell Host Microbe* **29**, 834-848.

**Russell, J. J., Theriot, J. A., Sood, P., Marshall, W. F., Landweber, L. F., Fritz-Laylin, L., Polka, J. K., Oliferenko, S., Gerbich, T., Gladfelter, A., et al.** (2017). Non-model model organisms. *Bmc Biol* **15**, 55.

**Sánchez, L. and Guerrero, I.** (2001). The development of the *Drosophila* genital disc. *Bioessays* **23**, 698–707.

**Sarashina, I., Shinmyo, Y., Hirose, A., Miyawaki, K., Mito, T., Ohuchi, H., Horio, T. and Noji, S.** (2003). Hypotonic buffer induces meiosis and formation of anucleate cytoplasmic islands in the egg of the two-spotted cricket *Gryllus bimaculatus*. *Dev Growth Differ* **45**, 103–112.

**Takagi, H., Inai, Y., Watanabe, S. -i, Tatemoto, S., Yajima, M., Akasaka, K., Yamamoto, T. and Sakamoto, N.** (2011). Nucleosome exclusion from the interspecies-conserved central AT-rich region of the Ars insulator. *J Biochem* **151**, 75–87.

**Takahashi, T., Hamada, A., Miyawaki, K., Matsumoto, Y., Mito, T., Noji, S. and Mizunami, M.** (2009). Systemic RNA interference for the study of learning and memory in an insect. *J Neurosci Meth* **179**, 9–15.

**Tchurikov, N. A., Kretova, O. V., Moiseeva, E. D. and Sosin, D. V.** (2009). Evidence for RNA synthesis in the intergenic region between enhancer and promoter and its inhibition by insulators in *Drosophila melanogaster*. *Nucleic Acids Res* **37**, 111–122.

**Thorvaldsdóttir, H., Robinson, J. T. and Mesirov, J. P.** (2013). Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* **14**, 178–192.

**Tomiyama, Y., Shinohara, T., Matsuka, M., Bando, T., Mito, T. and Tomioka, K.** (2020). The role of clockwork orange in the circadian clock of the cricket *Gryllus bimaculatus*. *Zoological Lett* **6**, 12.

**Trible, W., Olivos-Cisneros, L., McKenzie, S. K., Saragosti, J., Chang, Ni-Chen, Matthews, B. J., Oxley, P. R., Kronauer, D. J. C.** (2017). *orco* Mutagenesis Causes Loss of Antennal Lobe Glomeruli and Impaired Social Behavior in Ants. *Cell*. **170**:727-735.

**Wainberg, M., Kamber, R. A., Balsubramani, A., Meyers, R. M., Sinnott-Armstrong, N., Hornburg, D., Jiang, L., Chan, J., Jian, R., Gu, M.** (2021). A genome-wide atlas of co-essential modules assigns function to uncharacterized genes. *Nat Genet* 1–12.

**Watanabe, T., Ochiai, H., Sakuma, T., Horch, H. W., Hamaguchi, N., Nakamura, T., Bando, T., Ohuchi, H., Yamamoto, T., Noji, S., et al.** (2012). Non-transgenic genome modifications in a hemimetabolous insect using zinc-finger and TAL effector nucleases. *Nat Commun* **3**, 1017.

**Watanabe, T., Noji, S. and Mito, T.** (2014). Gene knockout by targeted mutagenesis in a

hemimetabolous insect, the two-spotted cricket *Gryllus bimaculatus*, using TALENs. *Methods* **69**, 17–21.

**Watanabe, T., Noji, S. and Mito, T.** (2017). Genome Editing in Animals, Methods and Protocols. *Methods Mol Biology* **1630**, 219–233.

**Wheeler, W. M.** (1892). On the appendages of the first abdominal segment of embryo insects. *Transactions of the Wisconsin Academy of Sciences, Arts and Letters* **8**, 87–140.

**Whittle, C. A., Kulkarni, A., Chung, N. and Extavour, C.G.** (2021). Adaptation of codon and amino acid use for translational functions in highly expressed cricket genes. *BMC Genomics* **22**, 234.

**Xiao, A., Cheng, Z., Kong, L., Zhu, Z., Lin, S., Gao, G. and Zhang, B.** (2014). CasOT: a genome-wide Cas9/gRNA off-target searching tool. *Bioinformatics* **30**, 1180–1182.

**Ylla, G., Nakamura, T., Itoh, T., Kajitani, R., Toyoda, A., Tomonari, S., Bando, T., Ishimaru, Y., Watanabe, T., Fuketa, M., et al.** (2021). Insights into the genomic evolution of insects from cricket genomes. *Communications Biology* **4,** 733.

**Yoshimi, K., Kunihiro, Y., Kaneko, T., Nagahora, H., Voigt, B. and Mashimo, T.** (2016). ssODN-mediated knock-in with CRISPR-Cas for large genomic regions in zygotes. *Nat Commun* **7**, 10431.

**Zhang, H., Shinmyo, Y., Mito, T., Miyawaki, K., Sarashina, I., Ohuchi, H. and Noji, S.** (2005). Expression patterns of the homeotic genes Scr, Antp, Ubx, and abd-A during embryogenesis of the cricket *Gryllus bimaculatus*. *Gene Expr Patterns* **5**, 491–502.

**Zhang, L., Martin, A., Perry, M. W., van der Burg, K. R. L., Matsuoka, Y., Monteiro, A. and Reed, R. D.** (2017). Genetic Basis of Melanin Pigmentation in Butterfly Wings. *Genetics* **205**, 1537–1550.

**Zhu, L., Mon, H., Xu, J., Lee, J. M. and Kusakabe, T.** (2015). CRISPR/Cas9-mediated knockout of factors in non-homologous end joining pathway enhances gene targeting in silkworm cells. *Sci Rep* **5**, 18103.

Matsuoka, Nakamura et al.

**Figure 1. *Gb-laccase2* knock-out G$_0$ and G$_1$ phenotypes.** (A) The cuticle of adult wild type (WT) *G. bimaculatus* is uniformly dark brown or black. Head and anterior thorax shown in dorsal view. (A') *Gb-lac2* gene somatic mutagenesis in G$_0$ animals can be detected by the presence of white or light brown spots of cuticle (asterisks). (B) Representative cuticle phenotypes of G$_1$ *Gb-lac2* mutant nymphs at one day after hatching. Homozygous mutants (-/-) showed homogeneous pale brown cuticle. Heterozyous mutants (-/+) showed slight reduction in black pigmentation. Control first instar nymphs have dark melanized cuticle by one day after hatching. Scale bar:1 mm. (C) Surveyor$^{TM}$ assay result. The control experiment, in which the PCR product was amplified from the genome of a wild type control individual, did not produce any band of the expected size after Surveyor$^{TM}$ endonuclease treatment. In contrast, the PCR product amplified from the genome of CRISPR reagent-injected animals included small fragments of the expected size after Surveyor$^{TM}$ endonuclease treatment (black arrowhead). M: electrophoresis marker. (D) Sequence analysis of *Gb-lac2* mutant alleles #6-2 and #1-10 induced by the CRISPR/Cas9 system. Top row: wild-type sequences; green: Protospacer Adjacent Motif (PAM) sequence; red: target sequence; arrowheads: predicted double strand break site. Asterisks at right show induced frame-shift mutations.

Matsuoka, Nakamura et al.

**Figure 2. Knock-out vs knock-down phenotype of *Gb-Ubx*.** (A) Schematic diagram of *Gb-Ubx* locus. White boxes: exons; red box: homeodomain; black arrowhead: sgRNA target site. (B) Surveyor$^{TM}$ assay results with G$_0$ eggs. Plus (+) indicates the PCR products digested by Surveyor$^{TM}$ nuclease; minus (-) indicates the PCR products with no digestion (no nuclease added). L = ladder. (C) Phenotype of heterozygous and homozygous *Gb-Ubx*$^{CRISPR}$ mutant stage 11 embryos. Anterior is to the left here and in all other figures. The size of the T3 leg (region between white arrowheads) was decreased mildly and severely in heterozygous and homozygous mutants, respectively. T3 legs are delineated with black dotted line in higher magnification images at right figures. Anterior (leftmost) white arrowhead marks the posterior end of T3 segment. Posterior (rightmost) white arrowhead marks the junction of femur and tibia. (D) Expression pattern of *Gb-Dll* in thoracic and abdominal appendages of wild type and homozygous *Gb-Ubx*$^{CRISPR}$ mutant embryos. Tarsus is to the left (distal); tibia is to the right (proximal). In wild type embryos, *Gb-Dll* is expressed strongly in the presumptive tarsus of developing T1, T2 and T3 legs (L1, L2 and L3 respectively). In the tibia of T3, different from T1 and T2, a sharply defined border between a distal domain of high expression is detectable. In homozygous *Gb-Ubx*$^{CRISPR}$ mutant embryos, this T3-specific expression pattern was not detected. In wild type embryos, *Gb-Dll* is ubiquitously expressed in the pleuropodia, but it showed the leg-like expression pattern in homozygous *Gb-Ubx*$^{CRISPR}$ mutant embryos. (E) Ubx/Abd-A (UbdA) protein expression pattern in *Gb-Ubx*$^{RNAi}$ stage 8 embryos. In stage 8 *Gb-Ubx*$^{RNAi}$ embryos, UbdA protein expression was undetectable in the T3 leg (L3) but was still detected in the A1 segment. (F) Ubx/Abd-A (UbdA) protein expression pattern in homozygous *Gb-Ubx*$^{CRISPR}$ stage 9 embryos. In *Gb-Ubx*$^{CRISPR}$ embryos, only the T3/A1 Gb-Ubx expression domain was undetectable. Scale bar: 100 μm. Pp: pleuropodium. Embryonic staging as per (Donoughe and Extavour, 2016).

Matsuoka, Nakamura et al.

**Figure 3. Exonic knock-in/Knock-out against *Gb-Ubx*.** (A) Scheme of knock-in experiment targeting a *Gb-Ubx* exon. White boxes: exons; red box: homeodomain; black arrowhead: sgRNA target site. Donor vector contains bait sequence (orange box) and expression cassette with the following elements: *Gryllus actin* promoter (blue arrow) followed by *eGFP* coding sequence (green box), and flanking *Ars* insulators (gray arrows). After co-injection of the donor vector with sgRNA for the donor vector, sgRNA for the genomic target site, and Cas9 mRNA, two patterns of insertion are predicted to occur due to NHEJ. (B) eGFP expression in *Gb-Ubx*$^{KI-exon}$ $G_0$ stage 11 embryos. 4.7% of $G_0$ embryos showed mosaic eGFP expression in the T3 legs (Table 3). (C) In $G_1$ stage 14 embryos, the eGFP expression pattern was identical to that of the previously reported expression pattern of *Gb-Ubx* (Barnett et al., 2019; Matsuoka et al., 2015; Zhang et al., 2005). Asterisks in (B) and (C) mark position of embryonic head. (D) Wild type embryo does not show any eGFP expression. (E) $G_2$ heterozygous mutants show an eGFP expression pattern identical to the pattern of *Gb-Ubx*. (F) $G_2$ homozygous mutants showed strong eGFP expression, and also showed phenotypes characteristic of *Gb-Ubx*$^{CRISPR}$ mutants, including shortened T3 legs and formation of leg-like structures on the A1 segment (white arrowhead). (G) Assessment of knock-in event by using PCR and Sanger sequencing. We designed PCR primers specific for each putative junction (black arrows flanking areas **a** and **b** of panel (A)). All three homozygous $G_2$ individual mutant animals assayed showed bands of the expected size for each junction. (H) Sequence analysis using the same primers indicated in panel A for genotyping confirmed that several deletions were generated due to NHEJ events at each junction. Black: genomic sequence; Blue: CRISPR target sequence; red: PAM sequence; pink: deleted nucleotides. Scale bar: 200 μm in (B)-(F). Embryonic staging as per (Donoughe and Extavour, 2016).

Matsuoka, Nakamura et al.

**Figure 4. Exonic knock-in/Knock-out against *Gb-abd-A*.** (A) Scheme of knock-in experiment against a *Gb-abd-A* exon. White box: exons; red box: homeodomain; black arrowhead: sgRNA target site. We used the same donor vector construct as that used in the experiment against *Gb-Ubx* (Fig. 3), substituting a *Gb-abd-A* exon-specific sgRNA. Two patterns of insertion are predicted to occur due to NHEJ. (B) Expression of eGFP in $G_0$ and $G_2$ *Gb-abd-A$^{KI-exon}$* embryos. 10.6% of $G_0$ *Gb-abd-A$^{KI-exon}$* embryos showed mosaic eGFP expression in the abdomen of stage 11 embryos (Table 3). We found that eGFP expression was accompanied by ectopic phenotypic leg-like structures (white arrowheads), consistent with the loss of *Gb-abd-A* activity. In $G_2$ *Gb-abd-A$^{KI-exon}$* stage 11 embryos, the expression pattern of eGFP was identical to the previously reported expression pattern of *Gb-abd-A* (Barnett et al., 2019; Matsuoka et al., 2015; Zhang et al., 2005). (C) Assessment of knock-in events by using PCR and Sanger sequencing. Genomic DNA was extracted from homozygous $G_2$ *Gb-abd-A$^{KI-exon}$* stage 11 embryos and used as a PCR template. We designed PCR primers specific for each putative junction (black arrows and regions **a**, **b** and **c**, **d** in panel (A)). The expected amplicon size was detected for each junction. L = ladder. (D) Sequence analysis using the primers for genotyping confirmed that multiple deletions or insertions were generated, likely due to the NHEJ events at each junction. Black: genomic sequence; Blue: CRISPR target sequence; red: PAM sequence; pink: deleted or inserted nucleotides. Scale bar: 200 μm in (B). Embryonic staging as per (Donoughe and Extavour, 2016).

Matsuoka, Nakamura et al.

**Figure 5. Intronic knock-in against *Gb-abd-A*.** (A) Scheme of knock-in experiment targeted to a *Gb-abd-A* intron. White boxes: exons; red box: homeodomain; black arrowhead: sgRNA target site. We used the same donor vector construct as that used in the experiment against *Gb-Ubx* (Fig. 3), substituting a *Gb-abd-A* intron-specific sgRNA. Two patterns of insertion are predicted to occur due to NHEJ. (B) Expression pattern of eGFP in $G_0$ and $G_2$ stage 11 *Gb-abd-A^{KI-intron}* embryonic abdomens. In *Gb-abd-A^{KI-intron}* embryos, patchy eGFP expression was observed but embryos did not show the ectopic abdominal appendage phenotype that was observed in *Gb-abd-A^{KI-exon}* embryos (Fig. 4B). *Gb-abd-A^{KI-intron}* $G_2$ heterozygous embryos show abdominal eGFP expression corresponding to the known pattern of embryonic *Gb-abd-A* transcripts (Barnett et al., 2019; Matsuoka et al., 2015; Zhang et al., 2005)*,* and the embryos did not show any morphological abnormality. *Gb-abd-A^{KI-intron}* $G_2$ homozygous embryos generated ectopic leg-like structures on the abdomen (white arrowheads) as observed in $G_0$ *Gb-abd-A^{CRISPR}* embryos (compare with (B')). (C) Assessment of knock-in event by using PCR with *Gb-abd-A^{KI-intron}* $G_2$ homozygous embryos. We designed PCR primers specific for each putative junction (black arrows and regions **a** and **b** in panel (A)). Scale bar: 500 μm in (B). Embryonic staging as per (Donoughe and Extavour, 2016).