Extreme longevity of highly fecund termite queens achieved by mitochondrial and insulin upregulation without harmful lipid signatures or accumulation

Sarah Séité[1, 2#], Mark C. Harrison[3#], David Sillam-Dussès[4], Roland Lupoli[1, 2], Tom J. M. Van Dooren[5, 6], Alain Robert[4], Laure-Anne Poissonnier[7], Arnaud Lemainque[8], David Renault[9, 10], Sébastien Acket[11], Muriel Andrieu[12], José Viscarra[13], Hei Sook Sul[13], Z. Wilhelm de Beer[7], Erich Bornberg-Bauer[3] and Mireille Vasseur-Cognet[1, 2, 14*]

1- UMR IRD 242, UPEC, CNRS 7618, UPMC 113, INRAe 1392, Paris 7 113, Institute of Ecology and Environmental Sciences of Paris, Bondy, France.

2- University of Paris-Est, Créteil, France.

3- Institute for Evolution and Biodiversity, University of Münster, Münster, Germany.

4- Sorbonne University Paris Nord, Laboratory of Experimental and Comparative Ethology UR4443, Villetaneuse, France.

5- UMR UPMC 113, IRD 242, UPEC, CNRS 7618, INRA 1392, PARIS 7 113, Institute of Ecology and Environmental Sciences of Paris, Paris, France.

6- Naturalis Biodiversity Center, Leiden The Netherlands.

7- Department of Biochemistry, Genetics and Microbiology, Forestry and Agriculture Biotechnology Institute, University of Pretoria, Pretoria, South Africa.

8- Genoscope, François-Jacob institute of biology, Atomic energies Agency, University of Paris-Saclay, Evry, France.

9- University of Rennes, CNRS, ECOBIO (Ecosystems, biodiversity, evolution) - UMR 6553, Rennes, France.

10- University Institute of France, Paris, France

11- University of Technology of Compiègne, UPJV, UMR CNRS 7025, Enzyme and Cell Engineering, Royallieu research Center, Compiègne, France.

12- Cochin Institute, UMR INSERM U1016, CNRS 8104, University of Paris Descartes,

CYBIO Platform, Paris, France.

13- Department of Nutritional Sciences and Toxicology, University of California, Berkeley,

USA

14- INSERM, Paris, France

* Correspondence and requests for materials should be addressed to MVC

email: mireille.vasseur@inserm.fr

https://orcid.org/0000-0001-6963-4114

#These authors contributed equally

ABSTRACT

Eusocial termite queens achieve nearly maximal fertility throughout their extremely long life

without apparent signs of aging. Termites represent, therefore, an ideal model for aging

research. To investigate the molecular mechanisms underlying their long reproductive life, we

carried out transcriptomic, lipidomic and metabolomic analyses on fat bodies of sterile short-

lived workers, long-lived kings and five stages spanning twenty years of adult queen

maturation. In mature reproductives, genes supporting a robust mitochondrial functioning or

associated with genome stability were upregulated. In most organisms, insulin signaling

increases fertility but decreases lifespan, often accompanied by harmful lipid signatures. Our

findings suggest that an upregulation of insulin-like peptide (Ilp9) in the fat body of termite

queens is accompanied by a specific lipid metabolism, limiting fat storage, thus sustaining both

high fertility and maintaining extreme lifespan. Our results highlight potential molecular targets

for research into aging-related metabolic diseases linked to the accumulation of excess fat.

INTRODUCTION

Aging affects almost all living organisms. It is characterized by the failure of several cellular and physiological functions. This often leads to a deleterious accumulation of lipids[1], while a reduction in mitochondrial functioning exacerbates oxidative stress[2]. In most multicellular organisms, surgical or genetic interventions which reduce fecundity can increase lifespan[3], suggesting that fecundity and longevity are negatively correlated[4,5]. This pattern is usually explained by a trade-off, where resources allocated to fecundity are no longer available for somatic maintenance and thus survival[4,6,7]. However, the regulatory mechanisms and signaling pathways controlling the allocation of resources in this trade-off remain insufficiently understood[8].

Our understanding of the causes of aging stems to a large extent from studies on short-lived model organisms[9,10]. Eusocial insects such as the fungus-growing termite *Macrotermes natalensis* (Termitidae, Blattodea) have reproductive castes which outlive their sterile siblings by orders of magnitude despite having the same genome[11–13]. Queens lay thousands of fertile eggs per day[14], and both queens and kings can live for several decades, while workers live for only a few weeks. Long-lived termite queens seem close to their maximum possible fertility for a prolonged time without apparent signs of aging and thus with a negligible cost of reproduction[15]. Among social insects, termites are an excellent new natural model system for aging research. Several molecular mechanisms involved in bypassing the fecundity/longevity trade-off have been identified in social insects, but their integrated physiological effects remain poorly understood[15–19].

Recently, transcriptomic studies in different taxa of social insects have pointed out the importance of downstream components of the Insulin/insulin-like growth factor (IGF-1) signaling (IIS) and target of rapamycin (TOR) network for aging [20–22]. IIS and TOR signaling have been broadly identified as key actors in the allocation of resources and aging of

organisms[23]. IIS is an evolutionarily conserved pathway which controls food intake, energy

stores, energy expenditure, growth and fecundity via the IRS–PI3K–AKT signaling cascade

which targets, in particular, TOR complex 1 (TORC1), inducing protein translation and *de novo*

lipogenesis[24]. In *Drosophila*, a reduction of the activity of IIS or TOR pathways can extend

lifespan substantially while reducing female fecundity[25–27]. Beyond the influence of these two

signaling pathways on fecundity in *Drosophila*, dysregulation of the IIS or TOR pathways, for

instance by the impairment of lipid metabolism, leads to the development of metabolic, age-

related pathologies from *Drosophila* to humans, such as type 2 diabetes and insulin

resistance[28,29].

Physiological consequences of differences between social insect castes in IIS and TOR

pathways and targets have thus far not been assessed using metabolomics or lipidomics. The

fat body, which acts like the mammalian liver and white adipose tissue, is central for

intermediary metabolism and energy balance in insects[30,31]. In *Macrotermes* termites, two

castes with different reproductive roles (queens and kings) live up to high age and can be

compared with short-lived castes. This permits a separation of shared lifespan prolonging

mechanisms from those that differ between the sexes. Some differences can be attributed to

sex-specific reproductive roles and could help to understand the absence of a cost of

reproduction in females or in both sexes. Whereas a gene expression module has been proposed

to characterize "queen-ness" (QCM) in termites, the analysis did not account for properties

shared with kings[21]. Neither have the fat bodies of kings and queens of highly eusocial termites

been investigated.

We studied *M. natalensis* abdominal fat bodies of five queen reproductive stages (adult

virgin to twenty-year-old queen), long-lived reproductive kings and short-lived,

nonreproductive female workers. We identified multiple biological pathways affecting aging

through OMICS analyses. In queens and kings, an anticipated downregulation of the TORC1

signaling occurs but also an unexpected 800-fold upregulation of an insulin-like peptide gene

which we called *Ilp9*. The upregulation of the *Ilp9* gene coincides with high glucose levels in

the hemolymph of mature queens. This apparent insulin increase is associated with an

upregulation of gene programs involved in the synthesis of proteins and specific lipids with low

oxidation potential, destined for oogenesis rather than fat storage. We were able to link this

phenomenon to a non-canonical downregulation of a gene coding a diacylglycerol

acyltransferase (*midway*; *mdy*) in mature reproductives. Consistent with transcriptomic data,

lipidomic analyses in the fat body of queens demonstrated a decrease in the concentration of

preferentially stored lipids (triglycerides) and an increase in the concentration of lipids destined

for oogenesis (diglycerides). Furthermore, we observed an upregulation of genes supporting a

robust mitochondrial functioning and of aging-related genes associated with genome stability

which may drive their long-lifespan.

RESULTS

Analysis of gene expression patterns in fat bodies

Macrotermes natalensis queens (QT4) and kings (KT4), which were over 20 years old, and

short-lived female (minor) workers (FW) were sampled in field colonies in 2016 and in 2018.

Incipient colonies were each founded from one male and one female imago (T0) collected in

the field and raised for 31 months following a protocol based on the natural life history of

Macrotermes species[32,33] (Fig 1a and Methods). Queens of these laboratory colonies were

sampled at 3 months of age (QT1), 9 months (QT2) and 31 months (QT3). To investigate how

caste and sexual maturity affect gene expression, we analysed a total of 25 transcriptomes of

abdominal fat bodies. Sterile female workers FW, queens QT0-QT4 and mature kings KT4

were each sampled from at least 3 independent colonies (Fig. 1a). RNA-sequencing data

showed that variation in expression between castes was greater than between replicates

(Principal Component Analysis, PCA of top 500 genes in terms of variance, Fig. 1b). This PCA

showed that FW expression was very distinct from that of KT4 and all queen stages, with a

strong separation along the first axis (52% variance explained). Removing FW expression from

the PCA allowed a clearer separation of the reproductive individuals within the first two axes

(58% variance explained). Four distinct groups were found: young queens (QT0 and QT1),

QT2, KT4 and mature queens (QT3 and QT4) (Fig. 1b). We then carried out a signed, weighted

gene co-expression network analysis[34]. The resulting gene co-expression network (GCN)

allowed us to identify nine modules of particularly strongly co-expressed genes (Fig. 1c). We

identified regions of the GCN that were up- or downregulated between castes or with queen

age. In accordance with the PCA (Fig. 1b), we found modules strongly and uniquely correlated

with FW (blue module, 1103 genes) and KT4 (plum1 module, 116 genes). One yellow module

was upregulated in mature queens, QT3 and QT4 (yellow module, 537 genes). Finally, the light-

cyan module (3455 genes) may be related to long-lived reproductives QT4 and KT4, though

the latter only as a trend (KT4 0.39 p = 0.05). This largest of the modules was enriched for

functions related to transcription and general protein synthesis. We subsequently tested for

expression differences between specific pairs of castes or ages in all or a subset of individual

genes.

Long-lived reproductives share expression patterns underlying remarkable lifespan

Despite their pronounced sexual dimorphism, the king and the queen in *M. natalensis* have

similar longevity. A comparison of transcriptomic profiles in fat bodies of KT4 and QT4 with

FW allows us to look at genes potentially underlying lifespan (Supplementary Fig.1),

irrespective of reproductive efforts distinguishing kings from queens. A total of 1454 genes

were upregulated in fat bodies of KT4 and QT4 including 350 genes that differed between KT4

and QT4. In addition, 2208 genes were downregulated in KT4 and QT4, 468 of which were

significantly different between KT4 and QT4. We observed significantly upregulated

expression of components of IIS signaling in the fat bodies of KT4 and QT4 with *Ilp9* (800-

fold increase), *pi3k59f* and two insulin targets *eIF6* and *crc*[35] (Fig. 2 and Supplementary Tables

1 & 2). These results were accompanied by the upregulation of gene expression involved in

several processes related to protein and macromolecule synthesis (Supplementary Fig. 1). In

contrast, the expression of *InR2* and several actors involved in the TORC1 signaling pathway

(*tor*, *raptor* and its *S6K* substrate) were downregulated in QT4 and KT4, as well as the *mdy*

lipid metabolism gene (Fig. 2 and Supplementary Tables 1 & 2). *Mdy* is a diacylglycerol

acyltransferase gene coding an enzyme catalysing the final step of triglyceride (TG) synthesis

from diglycerides (DG)[36]. This downregulation of the *mdy* gene suggests a decrease of fat

storage in reproductives. Moreover, the expression of many genes important for genome

stability, including genes involved in DNA damage response and telomere maintenance, was

upregulated in KT4 and QT4 (Fig. 2 and Supplementary Tables 1 & 2). In addition, several

genes coding for the OXPHOS systems and the mitochondrial ribosomal proteins,

mitochondrial transport and mitochondrial fission were upregulated in QT4 and KT4 (Fig. 2

and Supplementary Tables 1 & 2). These results suggest that mitochondrial function was

maintained in the fat body of long-lived reproductives with likely beneficial effects on cell

integrity and oxidative status. Accordingly, we observed a downregulation of the expression of

antioxidant genes in QT4 and KT4 compared to FW (Fig. 2 and Supplementary Table 1 & 2).

Together, these results suggest that long-lived reproductives share several expression

differences from workers in agreement with a reduction or absence of aging, but also an

unexpected upregulated expression of several components of the IIS pathway and a

downregulated expression of *mdy* and antioxidant genes.

The fat body of highly fecund and long-lived queens indicates a specific energetic gene program

To better understand differences in metabolic gene programs which might be explained by mature queen reproduction and relate to the absence of costs of reproduction, we mapped the differentially expressed genes (DEGs) in the fat bodies of FW and QT4 to the GCN and focused on patterns where QT4 differ from FW and KT4, which oogenesis might explain. These DEGs occupied very distinct regions of the network, with genes upregulated in FW mainly contained within the blue, light-yellow and light-green modules and DEGs in reproductives mainly contained within light-cyan (Fig. 1c). Although reproductives have several DEGs in common, differences between KT4 and QT4 occur: genes contained in the plum module were specifically up-regulated in KT4 while genes in the yellow module were specifically up-regulated in QT4 (Fig. 1c). Upregulated gene expression in FW was centered around cell signaling and responses to stimuli, whereas gene expression in QT4 and KT4 was mainly related to macromolecule synthesis with higher protein production in the KT4 fat body (Fig. 1c and Supplementary Fig. 2). Several genes involved in IIS pathway were differentially regulated in QT4 relative to FW and KT4. *InR3 and pdk1* expression were upregulated in QT4 compared to FW, and downregulated in KT4 (Supplementary Table 3). *pi3k59f* and *crc* were more strongly upregulated in QT4 than in KT4 (Supplementary Table 2). In QT4, we also observed an upregulation relative to FW of *pi3k21b* genes encoding the kinases involved in the IIS pathway (Supplementary Table 2). In contrast, *pi3k92e* gene expression was downregulated in the fat body of QT4 relative to FW (Supplementary Table 2). In response to carbohydrates, insulin concentration increases and activates, via the IIS pathway, glycolysis and *de novo* lipogenesis and increase mRNA translation[35,37]. We found major differences in gene expression within carbohydrate and lipid metabolism pathways between QT4 and both FW and KT4. Several genes, located in the blue and light-green GCN modules and involved in the glycogenesis and

trehalose energetic storage pathways, were upregulated in FW (Supplementary Tables 1-3).

Trehalose-6-phosphate synthase was upregulated relative to QT4, but not to KT4

(Supplementary Table 3). In QT4, on the other hand, several genes of the yellow and light-

cyan GCN modules which are key players in carbohydrate catabolism were upregulated relative

to KT4 and FW, particularly genes encoding enzymes involved in glycolysis (e.g. *cg6650* also

known as *adpgk*, *pfk* and *pkm;* Supplementary Table 3), as well as hexosamine biosynthetic

pathway (HBP) and pentose phosphate pathway (PPP) (Supplementary Tables 2 & 3). These

were found next to those promoting fatty acid (FA) synthesis from carbohydrates (*acc* and

fasn1; Supplementary Table 3). Interestingly, the expression of genes promoting FA activation,

esterification and elongation was also upregulated in QT4 relative to KT4 and FW

(Supplementary Table 3). Heightened DG transport in QT4 is suggested by an upregulation

relative to FW and KT4 of expression of lipoprotein genes essential for oogenesis, such as the

female-specific vitellogenin (*vg*) and the diacylglycerol-carrying lipoprotein (*hdlbp*)

(Supplementary Tables 2 & 3).

215   Overall, the results of this comparison suggest that the metabolism of FW tends to

facilitate energetic storage while the QT4 metabolism seems to be oriented towards the

utilization of energy resources for reproduction (oogenesis), more strongly so than KT4.

Transcriptomic analysis of queens at different stages highlights adult developmental

remodeling and expression changes within the highly fecund mature stage.

Concomitantly with the development of colonies, the imaginal fat body of virgin females (QT0)

with fat storage tissue becomes replaced by a royal fat body (QT4) which is highly oriented

towards specific protein synthesis and secretion[38]. Since endoreduplication (the replication of

DNA during the S-phase of the cell cycle without subsequent cell division) was specifically

reported for the fat body of queen termites[39], we recorded polyploidization in the fat body

during queen maturation using flow cytometry. Moreover, to investigate the molecular

mechanisms involved in the establishment of the dedicated energetic metabolism in mature

QT4 and in the change of fat body identity with maturation, we compared gene expression

between adult females in five life stages (Fig. 1a). Different regions of the gene co-expression

network were upregulated in each stage (Fig. 3), and we focus on comparing expression

differences between adjacent stages.

First of all, the level of expression of *Ilp9*, *InR3* and *vg* genes was not different between

QT3 and QT4 (Supplementary Table 4 and Fig. 4). However, *crc* gene expression

(Supplementary Table 4) and genes involved in glycolysis and OXPHOS, mitochondrial

membrane transport proteins and mitochondrial ribosomes, were all strongly upregulated in

QT4 compared to QT3 (Supplementary Table 4). Accordingly, GO terms suggest an increase

of several processes linked to protein synthesis and secretion (Fig. 3). We can conclude from

this that among mature queen stages, substantial further upregulation of important processes

can occur with reproductive age and that these reinforce the energy metabolism discussed

above. In particular, mitochondrial functions seem to be upregulated substantially with age.

Other main aspects of mature queen metabolism might be present already in young queens,

before massive reproduction occurs.

Queens live several weeks without food after mating, due to a lack of workers which

feed all colony members[38]. The imaginal fat body of virgin females (QT0) at the start of this

starvation period displayed high expression of genes associated with intracellular signal

transduction, such as the superfamily Ras which notably controls cell growth. We observed that

Egfr gene expression and downstream signaling cascades, such as *Ras/Mapk* signaling were

upregulated in comparison to QT1 (Supplementary Table. 4). GO analysis revealed that several

signaling pathways involved in development and apoptosis were upregulated as well (Fig. 3).

Furthermore, within the IIS pathway, expression of *InR2*, *InR3*, *pi3k21b*, and *Akt1* were

upregulated (Supplementary Table 4).

At the end of the starvation period (stage QT1), we observed an enrichment of GO terms

related to an increase of catabolic processes (Fig. 3), an upregulation of genes involved in

pathways linked to an increased use of reserves. More specifically we observed an upregulation

of *Ampkα* gene expression (AMP-activated Protein Kinase α; Supplementary Table 4) known

to be activated under conditions of low energy to redirect the metabolism and restore energy

balance by inhibiting anabolic processes and promoting catabolic processes and mitochondrial

functioning[40]. Consistent with the increase of *Ampkα* gene expression, we observed increased

expression of genes involved in several catabolic processes such as β-oxidation and proteolysis

(Fig. 3 & 4), as evidenced by the upregulation of several genes coding for subunits of the

proteasome and autophagy and a decrease of anabolic processes such as lipogenesis

(Supplementary Table 4 and Fig. 4). We observed further responses which can be explained as

stress responses to starvation. AMPK is known to influence mitochondrial homeostasis by

initiating both the degradation of damaged mitochondria and mitochondrial biogenesis[40]. We

observed an upregulation of two genes involved in mitochondrial fission (*drp1* and *fis1*) known

to facilitate mitochondrial degradation by autophagy and two genes involved in a specific

degradation of mitochondria via autophagy called mitophagy, *cg5059* also known as *BCL2*

interacting protein 3 and *pink1* (Supplementary Table 4). Moreover, we observed an

upregulation of the expression of genes involved in the OXPHOS system, mitochondrial

ribosome subunits and mitochondrial transport proteins, suggesting an increase in mitochondria

biogenesis (Supplementary Table 4). Finally, we observed an increased expression of

antioxidant genes (Supplementary Table 4) which are known to be upregulated by AMPK in

situations of oxidative stress[41].

At QT2, antioxidant defense processes, as well as *Ampkα* gene expression, were

downregulated again (Supplementary Table 4). At this stage, GO terms showed that several

processes were linked to protein metabolism (Fig. 3). We believe that QT2 are fed by workers.

This is consistent with the increase in the number of workers per colony (Supplementary Fig.

3b) and of fungus development (Supplementary Fig. 3c). *Ilp9*, *pi3k21b* gene expression, and *de*

novo lipogenic genes were upregulated in QT2 compared to QT1, suggesting a first activation

of IIS pathway components in the fat body of QT2 (Supplementary Table 4 and Fig. 4). At the

same time, we observed a downregulation of *mdy* gene expression, even relative to the mature

queen stages (Fig. 4). In addition, several genes involved in the cell cycle, including *mcm2, 5,*

6, 7 and 10, several cyclins and cyclin-dependent kinases, such as *cycE*/*cdk2* and *orc1*, and the

proto-oncogene *myc* were up-regulated (Supplementary Table 4). Associated to these genes,

we observed an increase in QT2 fat body endoreduplication levels, with the proportion of 4C

nuclei now exceeding 70 % (Fig. 5) in comparison to 35% in QT0 and QT1. Taken together,

these data suggest that QT2 is a transitory stage where maturation is prepared or initiated in

many processes.

In QT3, the abdomen has become enlarged. During this period, expression of *Ilp9* and

InR3 genes was upregulated in comparison to QT2 whereas *InR2* gene expression decreased

(Supplementary Table 4). We also observed an upregulation of genes involved in glycolysis,

HBP and PPP (Supplementary Table 4). Similarly, the expression of genes involved in *de novo*

lipogenesis, as well as FA activation, elongation, esterification and transport were upregulated

(Fig. 4). This is in accordance with increased fertility and associated genes, such as *vg* (Fig. 4)

and these expression patterns do not change further in QT4. We found that over 90% of all cells

in the fat bodies of QT3 and QT4 have 4C nuclei (Fig. 5).

Overall, these data suggest that the energy metabolism of young queens shows many

characteristics in accordance with their starvation period. With social nutrition (increase in the

number of workers and food availability), colony sizes increase. The QT2 stage acts as a

transitional period where the energy metabolism of mature queens becomes established. This

is associated with an upregulation of several specific insulin-signaling genes and a

downregulation of genes involved in lipid storage (*mdy*). Other lifespan affecting processes

seem to be initiated in QT2 (e.g., upregulation of IIS pathway components), or in QT4, long

after maturation (e.g. upregulation of mitochondrial functions).

Major changes in metabolites and lipid composition in mature queens reflect their

transcriptomes

We investigated hemolymph metabolomes in FW and queen maturation stages (QT0-QT4).

Metabolomes differed between FW and QT4, and between each pair of successive queen

maturation stages (pairwise perMANOVA each $p < 0.05$, Fig. 6). FW and QT0 seemed little

different overall, while starved QT1 were well separated from the QT2 individuals in transition

towards massive reproduction (Fig. 6). Physogastric QT3 were globally intermediate between

QT0, QT2 and QT4 (Fig. 6). We compared individual metabolite concentrations between FW

and QT4 and between successive queen maturation stages (Supplementary Figure 4 and Table

5). Glucose and galactose are the metabolites present in the largest concentrations in QT4. This

high concentration of glucose[42] occurred similarly in FW but not in any of the younger queen

stages. Surprisingly, the concentration of trehalose was lower in QT4 than in FW or in any other

queen stage. Alanine, glycerol, aspartic acid, phenylalanine and glutamic acid occur in larger

concentrations in QT4 than FW, corresponding with ongoing increased protein and lipid

synthesis. Relative to QT0, starved queens QT1 have increased levels of leucine, threonine,

isoleucine and citric acid, which decrease again in QT2 and are indicative of proteolysis during

starvation.

The lipidomic analysis in hemolymph revealed eighty-one esterified lipid species of

which 34 differed in quantity and composition between FW and QT4 (Fig. 7 and Supplementary

Table 6). Strikingly, DG were significantly elevated in QT4 relative to FW while TG were

significantly diminished (Fig. 7 and Supplementary Table 6). Furthermore, we compared lipid

compositions between the fat bodies of young QT0, the transitional stage QT2 and QT4 mature

queens (Fig. 8a and Supplementary Table. 7). Thirteen TG concentrations were significantly

lower in QT4 relative to QT0 while other changes were often in the same direction as in

hemolymph. We investigated changes in PUFA/MUFA/SFA proportions in hemolymph of FW

and QT4 and in fat bodies of QT0, QT2 and QT4. In hemolymph, PUFA/MUFA/SFA

proportions were not significantly different between FW and QT4 (Fig. 7b). In contrast,

PUFA/MUFA/SFA proportions in fat bodies of QT2 and QT4 were significantly different from

QT0 (PERMANOVA, p-value < 0.05) (Fig. 8b). This difference seemed to be related to a larger

proportion of PUFA in QT0 compared to QT2 and QT4 (Fig. 8b). The determination of fatty

acid composition in fat bodies of queens allowed us to evaluate the levels of oxidative cell

damage. We calculated the relative peroxidability of membranes (peroxidation index, PI),

which decreased drastically from 29.5 at QT0, 7.9 at QT2 to 1.9 at QT4 (Fig. 8c).

Several hemolymph metabolites discriminate young queens, old queens and workers, in

agreement with results from fat body transcriptomics and highlight the importance of simple

sugars. Moreover, we observed that lipid concentrations in QT4 are consistent with the

upregulation of genes involved in the IIS pathway, *de novo* lipogenesis and with a

downregulation of *mdy* gene expression. Indeed, QT0 metabolism, as in FW, tends to facilitate

TG storage while the QT4 metabolism seems to be oriented towards the utilization of resources

for oogenesis without harmful lipid signatures.

DISCUSSION

Our results demonstrate that termite queens mobilize several mechanisms to sustain fertility and enable their remarkable lifespan without apparent costs of long-term massive reproduction. Against expectations, a gene coding an insulin-like peptide and components of the IIS pathway are upregulated in the fat body of highly fertile long-lived queens. The composition of lipids, the scarcity of stored fat observed in fat bodies of long-lived queens and the upregulated mitochondrial functions indicate that the energy metabolism required for reproduction remains efficient up to high age and does not require reserves to be stored in the queen's fat body.

Long lifespan on a carbohydrate diet

The intermediary metabolism in QT4 is centered on a high use of carbohydrates with an increase of glycolysis and the OXPHOS system for energy generation and the synthesis of specific lipids and proteins. The unexpected low trehalose concentration in long-lived queens, coupled with their high glucose levels also occurring in workers suggest that either trehalose is promptly hydrolyzed in physogastric queens or that glucose is an important component of what workers feed to physogastric queens in colonies with a well-developed fungus, without much trehalose synthesis in these queens. Poulsen *et al.*[43] have additionally shown that the diversity of decomposition enzymes encoded by the queen's microbiota is low and geared towards hydrolyzing simple sugars. Overall, all results confirm that long-lived queens receive a highly energetic feed enriched in simple sugars. In many organisms from *Drosophila* to humans, a prolonged carbohydrate-rich diet is associated with chronic metabolic diseases reducing lifespan[29,44]. Increased insulin secretion caused by such a diet leads to the accumulation of triglycerides, and disrupts IIS signaling over time leading to the development of insulin resistance[37,44,45]. In *M. natalensis* QT4, we were able to link the limited production of triglycerides from diglycerides to the non-canonical downregulation of a gene coding a

diacylglycerol acyltransferase (*mdy*), thus limiting a harmful accumulation of triglycerides in

the fat body.

Antioxidant defense is strongest in young queens

In young queens before mating (QT0), we observed an activation of the EGFR signaling

pathway and downstream signaling cascades that are known to coordinate multiple responses

such as the activation of apoptosis and remodeling (as described by Larsen[46], see also caste

determination in the honey bee[47]). After mating, three months pass before the appearance of

workers which feed all colony members. We observed that during the starvation period ending

at the QT1 stage, the absence of nutrition is associated with an upregulation of genes coding

for mitochondria biogenesis (OXPHOS system, mitochondrial ribosome subunits and transport)

and mitochondrial fission and degradation by autophagy. We also observed an upregulation of

the expression of genes coding for antioxidant enzymes which may suggest an increase of ROS

production. Several studies demonstrated that during calorie restriction, mitochondrial stress

induced by a transitory increase of ROS leads to a cellular adaptive response named

mitohormesis which allows an increase in stress resistance and promotes health and vitality[48].

We propose that a mitohermesis response occurs at the QT1 stage. Interestingly, gene

expression coding mitochondrial synthesis becomes upregulated again in mature queens, with

increased production in QT4 and at KT4 suggesting a maintenance of mitochondrial

biosynthesis which could be beneficial for health span. Associated with these results, we

observed a decrease of genes coding for antioxidant enzymes in reproductives. Similar results

were observed in honey bee[49] and ant[50] queens. These findings suggest that the mitochondria

of long-lived reproductives remain efficient and produce a comparatively low amount of ROS,

thus limiting oxidative damage, one of the main drivers of aging.

 Surprising upregulation of IIS pathway components

We observed several modifications, potentially triggered by social nutrition at the QT2 stage.

We observed a first upregulation of the *vg* gene. As recently suggested for other termite

species[39], polyploidy in the fat body of *M. natalensis* queens can boost their egg production by

increasing protein synthesis, notably VG. In *Drosophila*, seven out of eight insulin-like peptides

(DILPs) are mainly produced in the brain insulin-producing cells except for DILP6, which is

produced in the fat body. *dIlp6* gene expression is up-regulated during starvation[51,52]. In the

social bee, *Apis mellifera*, two *Ilp* genes coding AmILP1 and AmILP2 are expressed in both

tissues[53]. In *M. natalensis,* we identified only two genes from separate loci coding an ortholog

to DILP7 (not expressed in *M. natalensis* fat bodies) and a further paralog found across holo-

and hemimetabolous insects, without a clear ortholog in *Drosophila melanogaster*, which we

called *Ilp9* (Supplementary Fig. 5). During queen maturation, we observed potential increases

in insulin suggested by upregulated expression of *Ilp9.* If this occurs in long-lived queens in

response to increases in some carbohydrates, then this would be reminiscent of processes in

mammalian pancreatic beta-cells[54]. Not only does the expression of *Ilp9* increase concomitantly

with the formation of the royal fat body, it is also approximately 800-fold more expressed in

fat bodies of reproductives compared to workers. The increase of *Ilp9* gene expression was not

associated with an activation of TORC1, although the expression of genes involved in protein

synthesis were upregulated in the fat body of reproductives. However, we observed an increase

of *elF6* gene expression which is involved in insulin-stimulated translation, most notably

controlling adipogenic transcription factors like *crc* (also known as *ATF4*), a member of the

mTOR-independent pathway[35,55]. We propose that the IIS pathway, including the *Ilp9* gene,

activates the eIF6-*crc* gene program in the fat body of mature queens and kings. This in turn

increases the synthesis of proteins involved in lipid synthesis and essential for fecundity, despite

a downregulation of the TORC1 signaling pathway elsewhere described as the main pathway

for protein synthesis[56] and also downregulated in whole bodies of reproductives of lower

termite species[21,22].

No damaging fat signatures and no fat storage in long-lived queens

Simultaneously with the gradual establishment of the pattern of IIS signaling, we observed a

gradual upregulation of *de novo* lipogenic genes. Calculations based on lipid compositions in

fat bodies of queens of different ages highlighted significant changes in the proportion of

PUFA/MUFA/SFA with a decrease of PUFA concentrations. PUFA are a thousand times more

likely to oxidize than MUFA[57]. The oxidation of PUFA can set off an oxidative cascade with

the formation of radicals capable of damaging surrounding macromolecules and tissues[57].

Therefore, this decrease in PUFA concentrations can lead to a reduction in oxidative damage

and thus increase lifespan of queens. In agreement with this, we found a strong, progressive

decline of the peroxidation index (PI) indicating resistance of membranes to peroxidative

damage. Several studies in mammals, birds and invertebrates[57,58] have demonstrated that PI

values inversely correlate with longevity. In honey bees, queens also have a lower PI than

female workers[57]. Moreover, gradually downregulated expression of the *mdy* gene, coding a

diacylglycerol acyltransferase enzyme catalysing the final step of TG synthesis from DG[59],

corroborates the decrease of TG concentrations we observed during adult queen maturation

suggesting very limited storage of lipids in favor of immediate utilization. Also, we observed a

trend for DG to increase, which are known to be preferably used for lipid transport (not free

fatty acid as in vertebrates)[60]. Han & Bordereau already observed almost forty years ago by

electronic microscopy that lipid droplet number decreases during *Macrotermes* termite queen

maturation[38]. The decrease of *mdy* gene expression was also observed in the fat body of mature

kings, suggesting that the low level of stored TG can be generally beneficial for long-lifespan

in reproductives and is not just explaining the absence of a cost of reproduction in queens. This

suggests that the gene network involving *mdy* would be a target for research linked to the

accumulation of excess fat.

449   Although further investigations of, for example, kings of different ages and also

functional studies are necessary, our unique analysis reveals a remodeling of the fat body during

adult queen maturation concomitantly with changes in nutritional and social conditions. The

results suggest that the IIS signaling network may have been adapted in queen fat bodies to

maintain high fertility while also enabling their remarkable lifespan. Together with the

composition of lipids and the scarcity of stored fat observed, this seems to optimize both

sustained fecundity and health span, avoiding hyperfunction[61] and defying the trade-off

between reproduction and survival. Finally, we observed several upregulated cellular functions

in mature reproductives that are otherwise known to decline with age, notably mitochondrial

functions and DNA damage repair. Altogether, our findings on the aging process of *M.*

natalensis highlight potential molecular targets which could be involved in age-related

metabolic diseases in humans.

METHODS

Species studied

The termite *Macrotermes natalensis* (Haviland, 1898) belongs to the Macrotermitinae

(Termitidae, Blattodea), the only subfamily that has established an exo-symbiotic obligatory

relationship with basidiomycete fungi of the genus *Termitomyces*[62]. It lives in large colonies in

Southern Africa where it builds massive mounds[63]. When the colony is mature and in

appropriate environmental conditions, winged male and female imagoes leave the mound

during spring and disperse in synchronous swarms across colonies[64]. Male imagoes locate

female imagoes[65] and after pairing, couples perform dealation and establish new colonies as

queen and king[64]. They mate throughout their life and fertilize all eggs. Few months after the

establishment of the colony, workers explore the environment to collect spores and to inoculate

the fungus comb they build in the nest[66]. During the first years larvae emerge which become

sterile workers or soldiers[67]. Minor workers and soldiers are female and major workers are

male[68]. Mature colonies contain around one million short-lived sterile individuals. Workers

feed all colony members through trophallaxis (transfer of food from mouth to mouth), after a

complex digestion of lignocellulose by the fungus and intestinal microbiota[43]. The royal couple

lives for more than 20 years in a royal cell[11,69]. The old queen lays thousands of eggs per day[14]

while the median lifespan of a worker is 56 days[12,33]. After five to seven years, some larvae

develop into nymphs which become imagoes after several instars.

Sampling

Field colonies opened to collect animals had been followed for over 20 years by Jannette

Mitchell in a field belonging to the University of Pretoria (coordinates in Supplementary Table

8)[64]. Old minor workers (FW) and 20-year old queens (QT4) and kings (KT4) were sampled

from at least twenty-year-old colonies. Less than an hour was taken to reach the royal cells

containing QT4 and KT4. We observed that habitacle volumes were comparable between

colonies (data not shown), suggesting that the colonies were of the same age. QT4 and KT4

also showed limited variability in weight and length between colonies.

Establishment and maintenance of incipient termite colonies

Imagoes were collected in Pretoria (South Africa) in 2016 and 2018 during the spring swarming

flights (coordinates in supplementary Table 8). Mounds were covered with nets to retrieve

imagoes. These were placed in large boxes preserving humidity and immediately transferred to

the laboratory. Individuals were sexed by visual observation of their abdominal sternites,

weight and length were recorded and wings were manually removed. Establishment of

laboratory incipient colonies occurred for both field trips, following a protocol adapted from

Lepage[32] and Han & Bordereau[33] (and Supplementary Fig. 6). In October 2016 and 2018,

imagoes were collected from five colonies each time (males from a single colony, females twice

from four different ones). Per colony providing females (eight in total), 200 incipient colonies

were established. Each paired couple was introduced in a closed plastic box (6 x 5 x 4.5 cm)

filled with sieved soil collected near the mounds. The incipient colonies were kept in a breeding

room with controlled conditions: 28°C, 85% relative humidity and 12:12 photoperiod. Water

was used to keep the soil slightly moistened. Development of the colonies was visually

monitored. At three months, when workers started to explore, small pieces of dry wild oats

were supplied on the surface of the soil and wood was additionally supplied after 4.5 months.

A *Termitomyces sp.* fungus comb with nodules was collected from one mature field colony and

a small part of this comb was introduced in each box. After 3.5 months, mortality was 56% for

the 2016 incipient colonies (± 15% across field colonies of origin) and 30 ± 7% for the 2018

incipient colonies. When the termite populations outgrew their boxes, they were opened on one

side and placed inside bigger ones (18 x 12 x 7.5 cm after 3 months, 36 x 24 x 14 cm after 14

months, and 1000 x 70 x 40 cm after 21 months) filled with sieved moistened soil. Colonies

were checked every two days to supply water and food if needed and to remove moldy food.

Queens were sampled after 3 months (QT1) and 9 months (QT2) in colonies established from

both trips, and after 31 months (QT3) for the 2016 mission. At QT1, the number of minor and

major workers, soldiers, presoldiers, and larvae were counted in colonies where queens were

sampled. At QT2, these were also counted and weight of the fungus measured.

Species identity

Total DNA was isolated from the head and the legs of one imago of each of the 9 colonies. PCR

was performed using the cytochrome oxidase I gene primers: LCO 5'- GGT CAA CAA ATC

ATA AAG ATA TTG G -3' and HCO 5'- TAA ACT TCA GGG TGA CCA AAA AAT CA -

3'[70] and the 650-bp amplified fragment was sequenced and analyzed using the Barcode of Life

Database identification system (www.barcodinglife.org). Species identity of each colony was

confirmed to be *M. natalensis*[43].

Hemolymph and fat body collection

Hemolymph and fat body were collected from cold-anesthetized individuals (FW, QT0, QT1,

QT2, QT3 and QT4). A total of 85 FW were pooled in one hemolymph and fat body sample

while ten QT1, QT2 or QT3 were pooled. We obtained eight samples except for QT3 (three

samples). Hemolymph was collected under a binocular microscope with tapered glass Pasteur

pipettes inserted in the membranous part just behind the head. The mean volume of hemolymph

collected per individual was 0.5 µL for FW, 1.5 µL for QT0, QT1, QT2, 50 µL for QT3, and 1

mL for QT4. Hemolymph samples were collected in cryotubes, quickly frozen in liquid nitrogen

and kept at -80°C until use.

Subsequently, termites were killed by decapitation and abdominal fat body was collected. For

RNA and DNA extraction, the fat body was stored in a tube containing RNAlater buffer

(Invitrogen) and kept at -80°C until use. For lipid and metabolite analyses, nitrogen-frozen fat

bodies were crushed in a tube which was immediately frozen in liquid nitrogen and kept at -

80°C until use. For ploidy analyses, fat body was collected from one individual, stored in a tube

containing 200 µL of Cycletest PLUS DNA Reagent Kit buffer (Becton Dickinson) and kept at

-80°C until use.

RNA profiling

In the 2016 cohort (4 FW, 4 QT0, 4 QT1, 1 QT2, 3 QT3, 4 QT4 and 3 KT4) and 2018 cohort (2 QT2), total RNA was isolated from fat body samples using miRNeasy Micro kit (Qiagen) and RNAse-free DNAse according to the manufacturer's instructions (Qiagen). RNA-Seq library preparations were carried out from 500 ng total RNA using the TruSeq Stranded mRNA kit (Illumina, San Diego, CA, USA) which allows mRNA strand orientation (sequence reads occur in the same orientation as anti-sense RNA). Briefly, poly(A)$^+$ RNA was selected with oligo(dT) beads, chemically fragmented and converted into single-stranded cDNA using random hexamer priming. Then, the second strand was generated to create double-stranded cDNA. cDNA were 3'-adenylated and Illumina adapters added. Ligation products were PCR-amplified. All libraries were subjected to size profile analysis conducted by Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA) and qPCR quantification (MxPro, Agilent Technologies, Santa Clara, CA, USA) using the KAPA Library Quantification Kit for Illumina Libraries (KapaBiosystems, Wilmington, MA, USA), then sequenced using 150 bp paired end reads chemistry on a HiSeq 4000 Illumina sequencer (Illumina, San Diego, CA, USA). An Illumina filter was applied to remove the least reliable data from the analysis. The raw data were filtered to remove any clusters with too much intensity corresponding to bases other than the called base. Adapters and primers were removed on the whole read and low-quality nucleotides were trimmed from both ends (when quality value was lower than 20). Sequences between the second unknown nucleotide (N) and the end of the read were also removed.

RNA-seq analyses

RNAseq reads were mapped against the *M. natalensis* genome[43] using *hisat2* (version 2.1.0[71]) at default settings. Gene expression levels were then generated by counting reads mapping to

each gene of the *M. natalensis* genome (annotation version 2.3) using *htseq-count*[72].

Differential expression analyses were carried out in R (3.5.1) with the DESeq2 package[73],

comparing between all pairs of castes and queen stages, as well as comparing each caste and

queen stage against all others. Genes were considered significantly differentially expressed if

the adjusted *p*-value was less than 0.05. Principal component analyses (PCA) were also carried

out within the DESeq2 package[73]. Counts were transformed using the

varianceStabilizingTransformation function, and the PCA was calculated and plotted using the

plotPCA function. This function carries out a PCA on the top 500 genes, based on variance. A

weighted gene co-expression network (WGCN) was generated with these gene expression

counts, using the R package WGCNA[34]. Normalized counts were extracted from the DESeq2

data set with the *counts* function. These data were filtered for genes with zero variance or with

missing values with the WGCNA function *goodSamplesGenes*. With the remaining 9631 genes,

a signed WGCN was created using a soft power of 14, implementing the biweight

midcorrelation calculation and setting the minimum module size to 30. Modules with a

dissimilarity less than 0.5 were merged using the *mergeCloseModules* function. We related the

expression profiles of the resulting nine modules to castes and queen stages by correlating

(Pearson's *r*) the module eigengenes (first principal component of the expression matrix of each

module) with a binary vector, containing 0s and 1s depending on the membership of each

sample (FW, QT0, QT1, QT2, QT3, QT4 or KT4). A significant positive correlation signifies

an overall upregulation while a negative correlation signifies a downregulation of expression

within the module for a given caste or queen stage.

To visualize the WGCN, we first reduced the WGCN to include only the most highly

connected nodes. We did this by retaining genes with a topological overlap of at least 0.2 with

at least another gene, and by including the top 15 most connected genes within each module.

This reduced WGCN (5823 genes) was exported to Cytoscape (version 3.8.0[74]) with the

exportNetworkToCytoscape function in WGCNA (threshold 0.15). In Cytoscape, the network

was rendered using the Edge-weighted Spring Embedded Layout and nodes were colored by

module membership or expression fold change. GO term enrichment analyses were carried out

with topGO (version 2.34.0[75]), using the classic algorithm. Node size was set to 5, Fisher exact

tests were applied, and we only kept GO terms that matched with 2 genes at least and with a *p*-

value < 0.05.

Analysis of ILPs

Two ILP genes (Mnat_00258 and Mnat_03820) were found in the *M. natalensis* proteome

based on orthology to ILPs in *D. melanogaster*. Orthology groups had been determined in a

previous study [76] using orthoMCL [77]. We checked for further ILP genes within the genome by

mapping the protein sequences of these two *M. natalensis* genes and eight known *D.*

melanogaster ILPs (downloaded from NCBI; accessed February 2021) against the *M.*

natalensis genome. This was carried out with exonerate (v 2.2.0 [78]) using the protein2genome

model at default settings. No further ILP copies were found but the annotations of the two *M.*

natalensis genes were improved based on these exonerate alignments. ILP protein sequences

from 16 further insects (*Aedes aegypti, Apis mellifera, Atta cephalotes, Acromyrmex echinatior,*

Blattella germanica, Camponotus floridanus, Cryptotermes secundus, Ephemera Danica,

Harpegnathos saltator Linepithema humile, Nasonia vitripennis, Pogonomyrmex barbatus,

Rhodnius prolixus, Solenopsis invicta, Tribolium castaneum, Zootermopsis nevadensis and

Strigamia maritima) were extracted from the same orthology database and aligned with the *M.*

natalensis and *D. melanogaster* sequences using MAFFT (v7.397 [79]) with the E-INS-I strategy.

A gene tree was constructed with FastTree (v. 2.1.10 [80]) and visualized with iTOL[81].

Metabolomic analysis

A volume of 20 µL of hemolymph (from 2016 and 2018 experiments; 6 FW, 8 QT0, 7 QT1, 7 QT2, 3 QT3 and 4 QT4 samples) was used to determine metabolic profiles obtained by gas chromatography coupled with mass spectrometry (GC-MS). We used the experimental procedure described in Khodayari *et al.*[82], and adapted in Genitoni *et al.*[83]. Samples were homogenized in 450 µL of ice-cold methanol/chloroform (2:1, v/v) before the addition of 300 µL of ultra-pure water. After they have been vigorously vortexed, the samples were centrifuged for 10 min at 4,000 g (4°C). Then, 100 µL of the upper phase, which contains metabolites, was transferred to new glass vials (Thermofisher), speedvac dried at RT, and vials sealed with PTFE caps. The derivatization of the samples was conducted with a CTC CombiPAL autosampler (CTC Analytics AG, Zwingen, Switzerland), as described in Khodayari *et al.*[82]. The GC-MS platform consisted of an Agilent 7890B gas chromatograph coupled to a 5977B mass spectrometer. The injector was held at 250°C, and the temperature of the oven ranged from 70 to 170°C at 5°C/min, from 170 to 280°C at 7°C/min, and from 280 to 320°C at 15°C/min; at the end of the temperature ramps, the oven remained at 320°C for 4 min. A 30 m fused silica column (HP5 MS 30 m, I.D. 0.25 mm, thickness 0.25 µm, 5% Diphenyl / 95% dimethylpolysiloxane, Agilent Technologies) was used with helium as the gas carrier at 1 mL per min. The temperatures of the transfer line and ion source were 280 and 230°C, respectively. The split mode (split ratio: 2:1) was used for the injection of 1 µL of each sample, and detection was realized by electronic impact (electron energy: 70 eV) in full scan mode. The peaks list was annotated based on their MS fragmentation patterns with MassHunter. Detected metabolites were identified, and calibration curves were used to calculate the concentration of each metabolite.

Statistical analysis

Permutational MANOVA[84] was used to test for differences in multivariate metabolite

concentrations between stages and castes. We subsequently tested for differences in individual

metabolites between pairs of subsequent stages or between FW vs. QT0 and FW vs. QT4. The

concentrations of all metabolites were log transformed and compared between groups by using

Welch tests. Tail probabilities were corrected for multiple testing using the Benjamini-

Hochberg method. Tests were considered significant for a p-value < 0.05 and carried out using

R software (v 3.6.3).

Lipidomic analysis

Lipids were extracted for fatty acid profile analysis gas chromatography with flame ionization

detection (GC-FID) and gas chromatography coupled to mass spectrometry (GC-MS).

Lipidomic analyses were done by liquid chromatography coupled to mass spectrometry (LC-

HRMS/MS). Lipids were extracted from 20 µL of hemolymph (from 4 FW and 4 QT4) or from

fat body (from 4 QT0, 3 QT2 and 3 QT4) using a biphasic solvent system of cold methanol,

methyl tert-butyl ether (MTBE), and water, adapted from Cajka *et al.*[85]. Briefly, the samples

were transferred in 750 µL of MTBE and 150 µL of methanol into a 2 mL screw cap tube. For

lipidomic analysis, 1µL of internal standard (PC 31: 1|PC17:0-PC14:1) at 3,775 µg/mL was

added to each sample. After homogenization with the "Precellys tissue homogenizer" at 5,000

rpm for 5 minutes, 400 µL of H_2O was added to each sample. The samples were then

centrifuged at 13,000 rpm for 5 min. The upper phase containing the lipids (MTBE) was

transferred into a new tube and dried under a stream of nitrogen at 20°C. For fatty acid profile

analysis, extracted lipids were transferred in 100 µL of MTBE, methylated into fatty acids of

methyl esters (FAMEs) after the addition of 10 µL of tetramethylammonium hydroxide

(TMAH). After a centrifugation at 4,000 rpm during 5 min, supernatants were collected and

diluted 3 times into heptane prior to injection into GC-FID and GC-MS. For lipidomic analysis,

lipids extracted were taken up into 100 µL of isopropanol before injection into LC-HRMS/MS.

FAMEs analysis

Fatty acid profiles were separated and analyzed by a gas chromatography with flame ionization

detection (GC-FID 2010 Plus Shimadzu) equipped with a BPX 70 capillary column (SGE,

$30 \, \text{m} \times 0.25 \, \text{mm}$, $0.25 \, \mu\text{m}$) as described in Merlier *et al*.[86]. The fatty acids were identified by

comparison of the retention times of a standard solution of 37 fatty acid methyl esters (Sigma;

47885-U Supelco) in GC-FID and confirmed by high accuracy mass of molecular ions and their

fragments after injection into a GC-MS (Q-Exactive™, Thermo)[86]. The composition of fatty

acids was expressed as a relative percentage of their peak areas with respect to the total peak

area of all the fatty acids. The fat body samples were normalized by dividing peak areas with

total DNA concentration (ng/mL) measured with a Qubit Fluorometer (Thermofisher) and

Qubit dsDNA Assay kit (Invitrogen). The membrane peroxidation index (PI) of lipid extracts

in fat bodies of queens was calculated as the sum of bis-allylic methylene groups per 100 fatty

acids according to the equation[57] :

PI=(percentage of dienoics x 1) + (percentage of trienoics x 2) + (percentage of hexaenoics x 5)

Untargeted lipidomics analysis

The untargeted lipidomics analysis was conducted using a liquid chromatography-high

resolution tandem mass spectrometry (LC-HRMS/MS) analysis used as described and modified

from Ulmer *et al*.[87]. An HPLC 1290 (Agilent Technologies) coupled to a hybrid quadrupole

time-of-flight high definition (QtoF) mass spectrometer Agilent 6538 (Agilent Technologies)

equipped with an ESI dual source was used. Lipids were separated on a C18 Hypersil Gold

(100 x 2.1 mm, 1.9 µm, Thermofisher) at 50°C, using an elution gradient composed of a solution

of 20 mM of ammonium acetate and 0.1% formic acid (ACN: H_2O, 60:40, v/v) (solvent A) and

a solution of 20 mM of ammonium acetate and 0.1% formic acid (IPA: ACN:H_2O, 90:8:2, v/v)

(solvent B). Separation was conducted under the following gradient: 0–2 min from 32% (B),

2–3 min from 32% to 40% (B), 3–8 min from 40% to 45% (B), 8–10 min from 45% to 50%

(B), 10–16 min from 50% to 60% (B), 16–22 min from 60% to 70% (B), 22–28 min from 70%

to 80% (B), 28–30 min from 80% to 99% (B), 30–31 min from 99% to 32% (B), 31–36 min

from 32% to 32% (B). The flow rate was set at 250 µL/min. Two µL of samples were injected.

MS / MS spectra were acquired in positive mode and in negative mode in data dependent and

MS^2 scans were performed on the sixth most intense ions. The source temperature, fragmentor

and the skimmer were set up at 350°C, 150 V and 65 V, respectively. The acquisition was made

in full scan mode between 100 m/z and 1700 m/z, with a scan of two spectra per second. Two

internal references were used for in-run calibration of the mass spectrometer (121.0509,

922.0098 in positive ion mode and 112.9856, 1033.9881 in negative ion mode). MassHunter

B.07 software allowed us to control the parameters of the machine acquired.

Data processing and annotation

MsDial v4.0 software[88] was used for data processing and annotation of lipids. The data

collection was performed by fixing the MS1 and MS2 tolerance, at 0.01 Da and 0.025 Da,

respectively. The peak detection was fixed at 1000 amplitude and a mass slice width at 0.1 Da.

The deconvolution parameters correspond to a sigma window value at 0.5 and a MS/MS

abundance cut off at 10 amplitude. Isotopic ions were kept until 0.5 Da. The peaks list was

annotated based on their unique MS/MS fragmentation patterns using the in-built LipidBlast

mass spectral library in MS-DIAL software. The accurate mass tolerance MS1 and MS2 were

fixed -at 0.01 Da and 0.05 Da respectively. The identification score cut off was fixed at 80%.

Lipids were normalized by the intensity of the internal standard (PC 31: 1|PC17:0-PC14:1).

 Statistical analysis for lipidomic data

Permutational MANOVA[84] on ilr-transformed compositional data[89] was used to test for

differences in SFA, MUFA and PUFA lipid content in hemolymph of FW and QT4. Similarly,

to compare the percentages of SFA, MUFA, PUFA lipid content in fat bodies of different queen

stages we used a permutational MANOVA followed by pairwise post-hoc perMANOVA

comparisons of each stage with QT4 (Holm-Bonferroni correction for multiple comparisons).

To compare log-transformed individual lipid values between FW and QT4, or between QT0,

QT2 and QT4, Welch tests were used, corrected for multiple testing using the Benjamini-

Hochberg method. To compare the peroxidation index (PI) in fat bodies of different queen

stages we used Kruskal-Wallis test followed by Dunn's post hoc comparisons. All tests were

considered significant for a p-value < 0.05 and were carried out using R software (v 4.0.2).

Heatmaps were made using the heatmap function in the Metaboanalyst v 4.0. with Euclidean

distance and clustering using Ward's method.

Ploidy analysis

Fat body from 2016 and 2018 cohort samples (4 QT0, 4 QT1, 3 QT2, 3 QT3 and 6 QT4) was

processed by Flow Cytometric Analysis with a Cycletest PLUS DNA Reagent Kit (BD

Biosciences, Le pont de Claix). All procedures were adapted from Nozaki & Matsuura[39].

Stained nuclei were analyzed for DNA-PI fluorescence using an Accuri C6 Flow Cytometer

(BD Biosciences) at an excitation wavelength of 488 nm and a detector equipped with an 585/45

bandpass filter. Approximately 1,000 cells were acquired for each measurement. Flow

cytometric analyses were performed with the Accuri C6 software v1.0.264.21 (BD

Biosciences). Debris were removed on an FSC-A/SSC-A dotplot and doublet were eliminated

with and PI-FL2-H/ FL2-A dot plot. The nuclei were analyzed with a histogram PI-A. The 1C

DNA peak was determined by the analysis of king's testis (sperm), allowing the identification

of the 2C, 4C, and 8C peaks of the others samples.

Statistical analysis for ploidy

To compare percentages of nuclei with different multiples of haploid genomes between queen

stages, we used permutational MANOVA followed by pairwise post-hoc perMANOVA (Holm-

Bonferroni correction for multiple comparisons).

Acknowledgements

The authors would like to thank Dr Christian Bordereau and Dr Jannette Mitchell for helpful

discussions on the termite model. The authors acknowledge the Department of Public Works

and Infrastructure, Pretoria and the University of Pretoria for allowing access to their sites of

collect of *M. natalensis*. The authors also thank Cyril Fresillon and Pierre Deparscau,

photographers for the CNRS images. We thank Qi Li for termite colony maintenance. This

study was supported by the International Human Frontier Science Program RGP0060/2018 to

M.V.C. S.S was also supported by a fellowship from Université de Paris Est-Créteil (UPEC).

A.L. acknowledges financial support from France Génomique (ANR-10-INBS-09-08). We

would like to thank the EcoChim platform of University of Rennes for access to metabolomics

facilities.

Author contributions

M.V.C designed the study. D.S-D., A.R., R.L., L-A.P., Z.W.D.B., and M.V.C. carried out

termite experiments, M.V.C and A.L. transcriptomic, D.R. and R.L. metabolomic and S.A. and

R.L. lipidomic experiments. M.A. and M.V.C. measured DNA contents. S.S., M.H., T.V.D

and M.V.C. analyzed the data, wrote the original draft and were responsible for the figures and

tables presented. All authors contributed with expertise, input, and edits throughout the text.

Competing interests

The authors declare that they have no conflict of interest.

Data availability

The authors declare that all data supporting the findings of this study are available within the

paper and supplementary information files, or are available from a dedicated github repository:

https://github.com/MCH74/Mnat_analyses. RNA-seq reads generated in this study are

available in Sequence Read Archive (BioProject ID: PRJNA685589 and BioSample

accessions: SAMN17088123- SAMN17088147).

Additional informations

Supplementary Information accompanies this paper.

REFERENCES

1. Johnson, A. A. & Stolzing, A. The role of lipid metabolism in aging, lifespan regulation, and age-related disease. *Aging Cell* **18**, e13048 (2019).

2. Balaban, R. S., Nemoto, S. & Finkel, T. Mitochondria, oxidants, and aging. *Cell* **120**, 483–495 (2005).

3. Flatt, T. Survival costs of reproduction in *Drosophila*. *Experimental Gerontology* **46**, 369–375 (2011).

4. Kirkwood, T. B. L. Evolution of ageing. *Nature* **270**, 301–304 (1977).

5. Kirkwood, T. B. L. & Austad, S. N. Why do we age? *Nature* **408**, 233–238 (2000).

6. Maklakov, A. A. *et al.* Antagonistically pleiotropic allele increases lifespan and late-life reproduction at the cost of early-life reproduction and individual fitness. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20170376 (2017).

7. Williams, G. C. Natural Selection, the Costs of Reproduction, and a Refinement of Lack's Principle. *The American Naturalist* **100**, 687–690 (1966).

8. Edward, D. A. & Chapman, T. Mechanisms underlying reproductive trade-offs: Costs of reproduction. *Mechanisms of Life History Evolution* (Oxford University Press, 2011).

9. Hansen, M., Flatt, T. & Aguilaniu, H. Reproduction, fat metabolism, and life span: What Is the Connection? *Cell Metabolism* **17**, 10–19 (2013).

10. Taormina, G. *et al.* Longevity: Lesson from model organisms. *Genes* **10**, 518 (2019).

11. Carey, J. R. Demographic mechanisms for the evolution of long life in social insects. *Experimental Gerontology* **36**, 713–722 (2001).

12. Traniello, J. F. A. & Leuthold, R. H. Behavior and ecology of foraging in termites. In *Termites: Evolution, Sociality, Symbioses, Ecology* (eds. Abe, T., Bignell, D. E. & Higashi, M.) 141–168 (Springer Netherlands, 2000).

13. Keller, L. & Genoud, M. Extraordinary lifespans in ants: a test of evolutionary theories of ageing. *Nature* **389**, 958–960 (1997).

14. Kaib, M., Hacker, M. & Brandl, R. Egg-laying in monogynous and polygynous colonies of the termite *Macrotermes michaelseni* (Isoptera, Macrotermitidae). *Insectes soc* **48**, 231–237 (2001).

15. Kuhn, J. M. M., Meusemann, K. & Korb, J. Disentangling the aging network of a termite queen. Preprint at https://www.biorxiv.org/content/10.1101/2020.12.19.423576v1 (2020).

16. Rodrigues, M. A. & Flatt, T. Endocrine uncoupling of the trade-off between reproduction and somatic maintenance in eusocial insects. *Current Opinion in Insect Science* **16**, 1–8 (2016).

17. von Wyschetzki, K., Rueppell, O., Oettler, J. & Heinze, J. Transcriptomic signatures mirror the lack of the fecundity/longevity trade-off in ant queens. *Molecular Biology and Evolution* **32**, 3173–3185 (2015).

18. Heinze, J. & Schrempf, A. Aging and reproduction in social insects – A Mini-Review. *GER* **54**, 160–167 (2008).

19. Harrison, M. C. *et al.* Gene co-expression network reveals highly conserved, well-regulated anti-ageing mechanisms in old ant queens. *Genome Biology and Evolution* (2021).

20. Korb, J. *et al.* Comparative transcriptomic analysis of the mechanisms underpinning ageing and fecundity in social insects. *Philosophical Transactions of the Royal Society B: Biological Sciences* **376**, 20190728 (2021).

21. Lin, S., Werle, J. & Korb, J. Transcriptomic analyses of the termite, *Cryptotermes secundus*, reveal a gene network underlying a long lifespan and high fecundity. *Communications Biology* **4**, 1–12 (2021).

824   22.   Haroon *et al.* Transcriptomic evidence that insulin signalling pathway regulates the
825   ageing of subterranean termite castes. *Scientific Reports* **10**, 8187 (2020).

826   23.   Pan, H. & Finkel, T. Key proteins and pathways that regulate lifespan. *Journal of*
827   *Biological Chemistry* **292**, 6452–6460 (2017).

828   24.   Teleman, A. A. Molecular mechanisms of metabolic regulation by insulin in
829   *Drosophila. Biochemical Journal* **425**, 13–26 (2009).

830   25.   Bjedov, I. *et al.* Mechanisms of life span extension by rapamycin in the fruit fly
831   *Drosophila melanogaster. Cell Metabolism* **11**, 35–46 (2010).

832   26.   Partridge, L., Alic, N., Bjedov, I. & Piper, M. D. W. Ageing in *Drosophila*: the role of
833   the insulin/Igf and TOR signalling network. *Experimental Gerontology* **46**, 376–381
834   (2011).

835   27.   Giannakou, M. E. *et al.* Long-lived *Drosophila* with overexpressed dFOXO in adult fat
836   body. *Science* **305**, 361–361 (2004).

837   28.   Fontana, L. & Partridge, L. Promoting health and longevity through diet: from model
838   organisms to humans. *Cell* **161**, 106–118 (2015).

839   29.   Álvarez-Rendón, J. P., Salceda, R. & Riesgo-Escovar, J. R. *Drosophila melanogaster*
840   as a model for diabetes type 2 progression. *BioMed Research International* vol. 2018
841   (2018).

842   30.   Arrese, E. L. & Soulages, J. L. Insect fat body: energy, metabolism, and regulation.
843   *Annu. Rev. Entomol.* **55**, 207–225 (2009).

844   31.   Li, S., Yu, X. & Feng, Q. Fat body biology in the last decade. *Annual Review of*
845   *Entomology* **64**, 315–333 (2019).

846   32.   Lepage, M. Development in the laboratory of incipient colonies of *Macrotermes*
847   *michaelseni* (Sjoestedt) (Isoptera, Macrotermitidae). *Annales de la Societe*
848   *Entomologique de France* (1990).

849   33.   Han, S. H. (Universite d'Abidjan & Bordereau, C. From colony foundation to dispersal
850   flight in a higher fungus-growing termite, *Macrotermes subhyalinus*, (Isoptera,
851   Macrotermitinae). *Sociobiology (USA)* (1992).

852   34.   Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network
853   analysis. *BMC Bioinformatics* **9**, 559 (2008).

854   35.   Brina, D. *et al.* eIF6 coordinates insulin sensitivity and lipid metabolism by coupling
855   translation to transcription. *Nature Communications* **6**, 8261 (2015).

856   36.   Buszczak, M., Lu, X., Segraves, W. A., Chang, T. Y. & Cooley, L. Mutations in the
857   *midway* gene disrupt a Drosophila Acyl Coenzyme A: Diacylglycerol Acyltransferase.
858   *Genetics* **160**, 1511–1518 (2002).

859   37.   Wang, Y., Viscarra, J., Kim, S.-J. & Sul, H. S. Transcriptional regulation of hepatic
860   lipogenesis. *Nature Reviews Molecular Cell Biology* **16**, 678–689 (2015).

861   38.   Han, S. H. & Bordereau, C. Origin and formation of the royal fat body of the higher
862   termite queens. *J. Morphol.* **173**, 17–28 (1982).

863   39.   Nozaki, T. & Matsuura, K. Evolutionary relationship of fat body endoreduplication and
864   queen fecundity in termites. *Ecology and Evolution* **9**, 11684–11694 (2019).

865   40.   Herzig, S. & Shaw, R. J. AMPK: guardian of metabolism and mitochondrial
866   homeostasis. *Nature Reviews Molecular Cell Biology* **19**, 121–135 (2018).

867   41.   Jeon, S.-M. Regulation and function of AMPK in physiology and diseases. *Exp Mol*
868   *Med* **48**, e245 (2016).

869   42.   Thompson, S. N.. Trehalose—the insect 'blood'sugar. *Advances in insect*
870   *physiology*, Vol *31*, 205-285. (Elsevier, 2003).

871   43.   Poulsen, M. *et al.* Complementary symbiont contributions to plant decomposition in a
872   fungus-farming termite. *PNAS* **111**, 14500–14505 (2014).

873   44.   Lee, D., Son, H. G., Jung, Y. & Lee, S.-J. V. The role of dietary carbohydrates in

874   organismal aging. *Cellular and Molecular Life Sciences* **74**, 1793–1803 (2017).

45.   Musselman, L. P. *et al.* A high-sugar diet produces obesity and insulin resistance in
wild-type *Drosophila*. *Disease Models & Mechanisms* **4**, 842–849 (2011).

46.   Larsen, W. J. Cell remodeling in the fat body of an insect. *Tissue and Cell* **8**, 73–92
(1976).

47.   Kamakura, M. Royalactin induces queen differentiation in honeybees. *Nature* **473**, 478–
483 (2011).

48.   Ristow, M. & Schmeisser, K. Mitohormesis: Promoting health and lifespan by increased
levels of reactive oxygen species (ROS). *Dose-Response* **12**, dose-response.13-
035.Ristow (2014).

49.   Corona, M., Hughes, K. A., Weaver, D. B. & Robinson, G. E. Gene expression patterns
associated with queen honey bee longevity. *Mechanisms of Ageing and Development*
126, 1230–1238 (2005).

50.   Parker, J. D., Parker, K. M., Sohal, B. H., Sohal, R. S. & Keller, L. Decreased expression
of Cu–Zn superoxide dismutase 1 in ants with extreme lifespan. *PNAS* **101**, 3486–3489
(2004).

51.   Slaidina, M., Delanoue, R., Gronke, S., Partridge, L. & Léopold, P. A *Drosophila*
Insulin-like Peptide Promotes Growth during Nonfeeding States. *Developmental Cell*
17, 874–884 (2009).

52.   Bai, H., Kang, P. & Tatar, M. *Drosophila insulin-like peptide-6* (*dilp6*) expression from
fat body extends lifespan and represses secretion of *Drosophila insulin-like peptide-2*
from the brain. *Aging Cell* **11**, 978–985 (2012).

53.   Wang, Y., Azevedo, S. V., Hartfelder, K. & Amdam, G. V. Insulin-like peptides
(AmILP1 and AmILP2) differentially affect female caste development in the honey bee
(*Apis mellifera* L.). *Journal of Experimental Biology* **216**, 4347–4357 (2013).

54.   Poitout, V. *et al.* Regulation of the insulin gene by glucose and fatty acids. *The Journal*
of Nutrition **136**, 873–876 (2006).

55.   Miluzio, A. *et al.* Translational control by mTOR-independent routes: how eIF6
organizes metabolism. *Biochem Soc Trans* **44**, 1667–1673 (2016).

56.   Wang, X. & Proud, C. G. Nutrient control of TORC1, a cell-cycle regulator. *Trends in*
Cell Biology **19**, 260–267 (2009).

57.   Martin, N. *et al.* Honey bee caste lipidomics in relation to life-history stage and the long
life of the queen. *Journal of Experimental Biology* **222**, (2019).

58.   Munro, D. & Blier, P. U. The extreme longevity of *Arctica islandica* is associated with
increased peroxidation resistance in mitochondrial membranes. *Aging Cell* **11**, 845–855
(2012).

59.   Kuhnlein, R. P. Lipid droplet-based storage fat metabolism in *Drosophila*. *J. Lipid Res.*
jlr. R024299 (2012).

60.   Gilbert, L. I. & Chino, H. Transport of lipids in insects. *J. Lipid Res.* **15**, 439–456 (1974).

61.   Gems, D. & Partridge, L. Genetics of longevity in model organisms: debates and
paradigm shifts. *Annu. Rev. Physiol.* **75**, 621–644 (2013).

62.   De Fine Licht, H. H., Andersen, A. & Aanen, D. K. *Termitomyces* sp. associated with
the termite *Macrotermes natalensis* has a heterothallic mating system and multinucleate
cells. *Mycological Research* **109**, 314–318 (2005).

63.   Uys, V. M. & Plant Protection Research Institute (South Africa). *A guide to the termite*
genera of Southern Africa No. 15. (Plant Protection Research Institute, Agricultural
Research Council, 2002).

64.   Mitchell, J. D. Colony Foundation and the development of incipient laboratory colonies
of *Macrotermes natalensis* (Haviland) (Termitidae: Macrotermitinae). *afen* **28**, 215–224
(2020).

65.   Sillam-Dussès, D. *Trail pheromones and sex pheromones in termites*. (Nova Science
Publishers/Novinka, 2010).

66.   Licht, H. H. D. F., Boomsma, J. J. & Aanen, D. K. Presumptive horizontal symbiont
transmission in the fungus-growing termite *Macrotermes natalensis*. *Molecular Ecology*
15, 3131–3138 (2006).

67.   Roisin, Y. Diversity and Evolution of Caste Patterns. In *Termites: Evolution, Sociality,*
Symbioses, Ecology (eds. Abe, T., Bignell, D. E. & Higashi, M.) 95–119 (Springer
Netherlands, 2000).

68.   Noirot, Ch. The Caste System in Higher Termites. In *Caste Differentiation in Social*
Insects Ch. 6 (eds. Watson, J. A. L., Okot-kotber, B. M. & Noirot, CH.) 75–86
(Pergamon, 1985).

69.   Keller, L. Queen lifespan and colony characteristics in ants and termites. *Insectes soc.*
45, 235–246 (1998).

70.   Folmer, O., Black, M., Hoeh, W., Lutz, R. & Vrijenhoek, R. DNA primers for
amplification of mitochondrial cytochrome c oxidase subunit 1 from diverse metazoan
invertebrates. *Molecular Marine Biology and Biotechnology* (1994).

71.   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*
37, 907–915 (2019).

72.   Anders, S., Pyl, P. T. & Huber, W. HTSeq—a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).

73.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

74.   Shannon, P. *et al.* Cytoscape: A Software environment for integrated models of
biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

75.   Alexa, A. & Rahnenfuhrer, J. topGO: Enrichment analysis for Gene Ontology. R
package version 2.28. 0. *BioConductor. Published online* (2016).

76.   Harrison, M. C. *et al.* Hemimetabolous genomes reveal molecular basis of termite
eusociality. *Nature Ecology & Evolution* **2**, 557–566 (2018).

77.   Li, L., Stoeckert, C. J. & Roos, D. S. OrthoMCL: Identification of ortholog groups for
eukaryotic genomes. *Genome Res.* **13**, 2178–2189 (2003).

78.   Slater, G. S. C. & Birney, E. Automated generation of heuristics for biological sequence
comparison. *BMC Bioinformatics* **6**, 31 (2005).

79.   Katoh, K. & Standley, D. M. MAFFT Multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular Biology and Evolution* **30**, 772–
780 (2013).

80.   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree: Computing large minimum evolution
trees with profiles instead of a distance matrix. *Molecular Biology and Evolution* **26**,
1641–1650 (2009).

81.   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Res* **47**, W256–W259 (2019).

82.   Khodayari, S., Moharramipour, S., Larvor, V., Hidalgo, K. & Renault, D. Deciphering
the metabolic changes associated with diapause syndrome and cold acclimation in the
two-spotted spider mite *tetranychus urticae*. *PLOS ONE* **8**, e54025 (2013).

83.   Genitoni, J. *et al.* Hypomethylation of the aquatic invasive plant, *Ludwigia grandiflora*
subsp. *hexapetala* mimics the adaptive transition into the terrestrial morphotype.
Physiologia Plantarum **170**, 280–298 (2020).

84.   McArdle, B. H. & Anderson, M. J. Fitting multivariate models to community data: a
comment on distance-based redundancy analysis. *Ecology* **82**, 290–297 (2001).

85.   Cajka, T., Smilowitz, J. T. & Fiehn, O. Validating quantitative untargeted lipidomics

across nine liquid chromatography–high-resolution mass spectrometry platforms. *Anal.*
Chem. **89**, 12360–12368 (2017).

86.   Merlier, F., Imatoukene, N., Octave, S., Nicaud, J.-M. & Thomasset, B. A gas
chromatography full scan high resolution Orbitrap mass spectrometry method for
separation and characterization of 3-hydroxymethyl pyridine ester of fatty acids at low
levels. *Journal of Chromatography A* **1575**, 72–79 (2018).

87.   Ulmer, C. Z., Patterson, R. E., Koelmel, J. P., Garrett, T. J. & Yost, R. A. A Robust
Lipidomics workflow for mammalian cells, plasma, and tissue using liquid-
chromatography high-resolution tandem mass spectrometry. in *Lipidomics: Methods*
and Protocols (ed. Bhattacharya, S. K.) 91–106 (Springer, 2017).

88.   Tsugawa, H. *et al.* A lipidome atlas in MS-DIAL 4. *Nature Biotechnology* **38**, 1159–
1163 (2020).

89.   Van den Boogaart, K. G., & Tolosana-Delgado, R. *Analyzing compositional data with*
R Vol. 122 (Heidelberg: Springer 2013).

 FIGURE LEGENDS

Figure 1: Overview of the model system. (a) Timeline of the different stages of *Macrotermes*

natalensis colonies founded from one male and one female imago each (T0). Queens from

incipient laboratory colonies were sampled at 3 months after colony establishment (T1), 9

months (T2) and 31 months (T3). Field termite colonies over 20 years old are added. From field

colonies, queens, workers and kings were sampled. Wild oats were supplied to the laboratory

colonies from 3 months onward and the fungus *Termitomyces sp.* was introduced artificially in

3.5 months old colonies. Wood was supplied from 4.5 months onward. The drawings represent

winged imagoes, workers, kings and queens at the different stages (physogastric queens become

larger). Replication and sampling in our incipient colonies is further described in

Supplementary Table 9. **(b)** Principal component analyses (PCAs) of gene expression across

25 samples, spanning 6 colonies, 3 castes and 5 queen stages (FW, QT0, QT1, QT2, QT3, QT4,

KT4). Scores for the two main PCs are shown for all data (left) and among reproductives (i.e.,

without FW; right). **(c)** Weighted Gene Co-expression Network analysis (WGCN). Nine

modules were detected, each represented by a color (rows). In the left-hand heatmap correlation

coefficients and *p*-values are shown, relating gene expression (first major axis of the expression

matrix) of each module to each caste and queen stage (columns). Positive correlations are red

and negative correlations are blue; significant correlation values ($p < 0.05$) have a green border.

Module size is also shown, reflecting the number of genes assigned to each module. Module

expression profiles show mean expression of each module within each caste and queen stage.

Enriched GO-terms of each module are displayed as tag-clouds, in which font size is inversely

related to *p*-value. A visualization of the network is displayed on the right with nodes colored

by their module membership. This network was created with Cytoscape (version 3.8.0) on a

reduced representation of the WGCN containing the top connected genes.

1013

Figure 2: Caste related changes in gene expression. Heatmap representing standardized gene

expression (blue = low; red = high) in fat bodies of FW, QT4 and KT4. Annotations to the right

of the heatmap include WGCN module (Fig. 1), gene names and gene acronyms in *Drosophila*

melanogaster (right) and *Homo sapiens* (left). The map is restricted to expression of genes

involved in metabolic signaling, lipid metabolism, mitochondrial oxidative phosphorylation

system (OXPHOS), mitochondrial ribosome (mt-ribo), mitochondrial fission (mt-fission),

mitochondrial transport (mt-transport), antioxidant defense (AO-def) and DNA repair.

Expression of all genes differ significantly between FW versus QT4 and between FW versus

KT4. The number of replicates per group is provided in Supplementary Table 9.

1023

Figure 3: Cytoscape during adult queen maturation. Changes of gene expression within the

gene co-expression network (GCN) with queen differentiation. Node color represents the

standardized gene expression within each queen stage (low=blue: high=red). Genes that are

significantly up- or down-regulated at each queen stage have a strong, black border. Enriched

GO-terms within each group of up- or down-regulated genes are displayed as tag clouds below

each network.

1030

Figure 4: Heatmap with the expression of genes in fat bodies during adult queen

maturation involved in lipid metabolism. Heatmap representing standardized gene

expression (blue = low; red = high) at each of the five queen stages (QT0-QT4). Annotations

to the right of the heatmap include WGCN module (Fig. 1), gene name and gene acronyms in

Drosophila melanogaster (right) and *Homo sapiens* (left). Number of replicates per group is

provided in Supplementary Table 9. Greater than or less than symbols (>/<) represent

significant differences in expression.

Figure 5: Changes in DNA content of fat body cells during adult queen sexual maturation. Percentages of nuclear 2C, 4C and 8C cells in the fat bodies of queens in different stages (QT0 to QT4). The ternary graph shows the actual data points with stages distinguished by a colour code. The table lists per stage the average percentages of 2C, 4C, 8C. Number of replicates per group are provided in Supplementary Table 9. The proportions of cells with different DNA content differ between stages (compositional perMANOVA, $p = 0.001$), and in particular stages QT0 and QT2 differ from QT4 (post-hoc perMANOVA, both $p = 0.02$).

Figure 6: Canonical discriminant analysis of concentrations of metabolites. Three canonical functions discriminate ages and castes significantly, of which we show scores for the first two. Average scores for each caste and age are shown, plus canonical structure coefficients of each metabolite as vectors from the origin. These are proportional in length to the magnitudes of the correlations of each metabolite with the scores of the discriminant functions and show how information from each metabolite aids in discriminating castes and ages. Numbers of replicates per group are provided in Supplementary Table 9.

Figure 7: Comparison of lipid profiles between FW and QT4 in hemolymph. (a) Hierarchical clustering heatmap analysis of triglycerides (TG, orange), diglycerides (DG, green), phosphatidylethanolamine (PE, black), phosphatidylcholine (PC, grey), lysophosphatidylcholine (LPC, yellow), sphingomyelin (SM, blue) lipids in hemolymph of FW and QT4 performed in MetaboAnalyst 4.0. Individual lipids are shown in rows and samples displayed in columns, according to cluster analysis (Euclidian distance was used and Ward's clustering algorithm). Each colored cell on the heatmap plot corresponds to a normalized lipid amount above (red) or below (blue) the mean concentration for a given lipid. **(b)** Fatty acid percentages in hemolymph of FW and QT4. Percentage (%) of saturated fatty acids (SFA),

monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA) of total FA in

hemolymph. Ternary graph showing the percentage of SFA (orange), MUFA (black) and PUFA

(blue). A permutational MANOVA found that SFA/MUFA/PUFA proportions in hemolymph

were not significant different between FW and QT4. Numbers of replicates per group are

provided in Supplementary Table 9.

Figure 8: Changes in lipid profiles in fat bodies during adult queen maturation. (a)

Hierarchical clustering heatmap analysis of triglycerides (TG, orange), diglycerides (DG,

green), phosphatidylethanolamine (PE, black), phosphatidylcholine (PC, grey),

lysophosphatidylcholine (LPC, yellow), sphingomyelin (SM, blue) lipids in fat body of

different stage of queen (QT0, QT2 and QT4) performed in MetaboAnalyst 4.0. Individual

lipids are shown per row and mean of lipid amount of each stage displayed in columns,

according to cluster analysis (Euclidean distance and Ward's algorithm). Each colored cell on

the heatmap plot corresponds to a normalized lipid amount above (red) or below (blue) the

mean concentrations for a given lipid. **(b)** Fatty acid percentages of different queen maturation

stages (QT0, QT2, QT4) in fat body. Percentage (%) of saturated fatty acids (SFA),

monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA) of total FA in

fat body. Ternary graph showing the percentage of SFA (orange), MUFA (black) and PUFA

(blue). The table below shows the averages of MUFA, SFA and PUFA for each stage.

Permutational MANOVA demonstrated that SFA/MUFA/PUFA proportions in fat bodies of

QT2 and QT4 were significantly different relative to QT0 (Permutational MANOVA, p-value

= 0.012). **(c)** Box plot illustrating the peroxidation index of fat bodies of different queen

maturation stages (QT0, QT2 and QT4). A box consists of upper and lower hinges and a center

line corresponding to the 25th percentile, the 75th percentile and the median, respectively.

Rhombuses represent the averages. Different letters indicate significantly different values

1088   according to a Kruskal-Wallis test followed by Dunn tests. Statistical differences are given for

1089 $p < 0.05$. Number of replicates per group are provided in Supplementary Table 9.

Figure 1

Figure 2

Figure 3

Figure 4

	QT0	QT1	QT2	QT3	QT4
2C (%)	62.1	55.8	23.3	6.8	5.9
4C (%)	31.8	39.0	76.2	92.7	91.4
8C (%)	6.1	5.2	0.5	0.6	2.6

Figure 5

Figure 6

	FW	QT4
SFA (%)	48.43	44.47
MUFA (%)	28.79	38.18
PUFA (%)	22.78	17.34

caste
● FW
● QT4

Figure 7

Figure 8