1 **Topography of corticopontine projections is controlled by postmitotic**
2 **expression of the area-mapping gene Nr2f1**
3

Chiara Tocco[1]*, Martin Øvsthus[2]*, Jan G. Bjaalie[2], Trygve B. Leergaard[2#], Michèle Studer[1#]

[1]*Université Côte d'Azur, CNRS, Inserm, iBV, France*

[2]*Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway*

*These authors have contributed equally to the study.

#Co-corresponding authors:

Michèle Studer
iBV - Institut de Biologie Valrose
Univ. Côte d'Azur
Centre de Biochimie ; UFR Sciences
Parc Valrose, 28 avenue Valrose
06108 Nice Cedex 2
France
Tel.: +33 489150720
e-mail: Michele.studer@unice.fr
19
Trygve B. Leergaard
Institute of Basic Medical Sciences
Division of Anatomy
Domus Medica
Gaustad
Sognsvannsveien 9
0372 Oslo
Norway
Tel. : +47 22851052
e-mail : t.b.leergaard@medisin.uio.no

Summary statement

Cortical gradient expression of the area patterning gene *Nr2f1* spatially and temporally
controls corticopontine topographic connectivity in layer V projection neurons.

Abstract

Axonal projections from layer V neurons of distinct neocortical areas are topographically organized into discrete clusters within the pontine nuclei during the establishment of voluntary movements. However, the molecular determinants controlling corticopontine connectivity are insufficiently understood. Here, we show that an intrinsic cortical genetic program driven by Nr2f1 graded expression is directly implicated in the organization of corticopontine topographic mapping. Transgenic mice lacking cortical expression of *Nr2f1* and exhibiting areal organization defects were used as model systems to investigate the arrangement of corticopontine projections. Combining three-dimensional digital brain atlas tools, *Cre*-dependent mouse lines, and axonal tracing, we show that *Nr2f1* expression in postmitotic neurons spatially and temporally controls somatosensory topographic projections, whereas expression in progenitor cells influences the ratio between corticopontine and corticospinal fibres passing the pontine nuclei. We conclude that cortical gradients of area patterning genes are directly implicated in the establishment of a topographic somatotopic mapping from the cortex onto pontine nuclei.

Keywords: corticopontine topography, layer V neurons, area mapping genes, Nr2f1, mouse models, *Thy1-eYFP-H* reporter line, pontine nuclei, 3D data analysis, 3D digital brain atlasing, anterograde fluorescent tracing

Introduction

Neuronal populations responsible for fine motor coordination are arranged in topographically organized maps in the neocortex and cerebellum, with different body parts being represented in largely continuous maps in the somatosensory cortex (Chapin and Lin, 1984; Fabri and Burton, 1991; Welker, 1971; Woolsey and Van der Loos, 1970), and discontinuous, fractured maps in the cerebellum (Bower, 2011; Bower et al., 1981; Bower and Kassel, 1990; Leergaard et al., 2006; Nitschke et al., 1996; Shambes et al., 1978). The intercalated regions of this network, the pontine nuclei, deep cerebellar nuclei, and thalamus, receive and integrate signals ultimately resulting in coordinated and seamlessly executed behaviours (Buckner, 2013; Peterburs and Desmond, 2016; Stoodley and Schmahmann, 2010), including fine voluntary movements (Badura et al., 2013; Mottolese et al., 2013).

The pontine nuclei constitute the major synaptic relay for cerebro-cerebellar signals (Brodal and Bjaalie, 1992; Lemon, 2008; Mihailoff et al., 1985). Axonal projections originating from layer V pyramidal neurons across the neocortex are distributed in topographically organized clusters within the pontine nuclei, as shown in monkey (Brodal, 1978; Schmahmann and Pandya, 1997), cat (Bjaalie and Brodal, 1997), rat (Leergaard et al., 2000a; Leergaard et al., 2000b), and to some extent also in mice (Henschke and Pakan, 2020; Inoue et al., 1991; Proville et al., 2014). Within the pontine nuclei, the three-dimensional (3D) arrangement of clustered terminal fields, well described in rats, both preserves the overall topographical relationships of the cortical maps, but also partially overlap and introduce new spatial proximities among projections from different cortical areas (Bjaalie and Brodal, 1989; Leergaard, 2003; Leergaard and Bjaalie, 2007).

To date, the mechanisms responsible for establishing the topographic map between the neocortex and pontine nuclei are poorly understood. The leading proposition, referred to as chrono-architectonic hypothesis, postulates that the complex 3D topography is a product of straightforward spatial-temporal gradients, possibly combined with non-specific chemo-attractive mechanisms (Leergaard, 2003; Leergaard and Bjaalie, 2007; Leergaard et al., 1995). Recent new discoveries open the possibility that other mechanisms are also in action during the establishment of the corticopontine maps. Several lines of evidence point to a

functional role of gradients in gene expression during topography of sensory maps in several

systems (D'Elia and Dasen, 2018; Erzurumlu et al., 2010; Fritzsch et al., 2019; McLaughlin

and O'Leary, 2005), but whether this process is also operative during establishment of

corticopontine topography is not understood. A recent study has shown that postmitotic

graded expression of the HOX gene *Hoxa5* is directly involved in imparting an anterior to

posterior identity to pontine neurons and attract corticopontine axons (Maheshwari et al.,

2020). Whether gradient expression of molecular factors along the antero-posterior or

medio-lateral axes of the cerebral cortex also intrinsically determine the topography of

corticopontine projections is still not known. Layer V neurons from the anterolateral

cerebral cortex project to the central regions of the pontine nuclei, while more medially

located cortical regions project to more external parts; motor area projections are

distributed more medially and rostrally, while somatosensory ones reach the middle and

caudal parts of the pontine nuclei. Finally, auditory and visual cortical projections innervate

the dorsolateral regions of the pontine nuclei (Leergaard et al., 2004; Leergaard and Bjaalie,

2007). The fine-tuned and precise corticopontine topography leaves open the possibility for

cortical neurons being intrinsically programmed to target specific groups of pontine

neurons, possibly coupling both intrinsic and extrinsic mechanisms in directing proper

topographical innervation to the pontine nuclei.

A new theme of cortical patterning has emerged, in which genetic factors direct the spatial

and temporal establishment of topographically organized axonal connections between the

cortex and subcortical brain regions (Cadwell et al., 2019). Area mapping genes are

expressed in gradients along the different axes of the cortical primordium, are known to

modulate the size and position of future cortical areas (Alfano and Studer, 2012; Cadwell et

al., 2019; O'Leary and Sahara, 2008), and to determine areal fate and regulate expression of

downstream molecules that in turn control the topographic organization of synaptic inputs

and outputs of related structures (Assimacopoulos et al., 2012; Greig et al., 2013). These

genes represent good candidates for modulating topographic mapping. In mice, the *Nr2f1*

gradient expression appears to be a particularly strong candidate for having a formative role

during the establishment of topographic maps (Armentano et al., 2007; Liu et al., 2000;

Zhou et al., 2001). For instance, *Nr2f1* is expressed in cortical progenitor cells from

embryonic day E9.0 in a high caudo-lateral (future sensory) to low rostro-medial (future

motor) gradient fashion, which is maintained in postmitotic descendants and postnatally when the cortical area map is completed (Bertacchi et al., 2019; Flore et al., 2017; Tomassy et al., 2010). Previous studies show that Nr2f1 promotes somatosensory (S1) by repressing motor identity in postmitotic neurons and in its absence area size and thalamocortical topography are both affected (Alfano et al., 2014; Armentano et al., 2007; Bertacchi et al., 2019). We thus hypothesized that Nr2f1 could represent one of these factors able to control topographic corticopontine mapping during corticogenesis.

To this purpose, we made use of cortico-specific *Nr2f1* conditional knockout mice as an i*n vivo* model system and a paradigm to investigate the contribution of cortical genetic programs in the establishment of topographic corticopontine projections. Two distinct conditional mouse lines were crossed to the *Thy1-eYFP-H* reporter line (Feng et al., 2000), in which YFP is highly expressed in cortical layer V pyramidal neurons and their axonal projections (Porrero et al., 2010). The distribution of fluorescent YFP signals as well as anterogradely labelled corticopontine projections were evaluated by comparison of spatially corresponding microscopic images and 3D visualization of extracted point-coordinate data representing labelling. Our results indicate that cortical Nr2f1 expression plays a dual role in controlling the spatial-temporal development of corticopontine projections. While early expression in progenitor cells influences the ratio between corticofugal fibres passing the pontine nuclei, loss of postmitotic late expression specifically affects topographic pontine mapping. Overall, our data demonstrate that intrinsic genetic programs and postmitotic graded expression of cortical area mapping genes are implicated in the establishment of area-specific targeting of corticopontine neurons.

Results

Benchmark 3D topographic organization of corticopontine projections in wild-type mice

To first establish a 3D reference of the topographical organization of corticopontine projections in wild-type adult mice, we used tract tracing data from the *Allen Institute Mouse Brain Connectivity Atlas* (Wang et al., 2020), see flowchart in **Figure 1A**. These data allowed us to visualize the spatial distribution of the pontine projections of motor and

somatosensory neocortical areas. Corticopontine projections digitized from sagittally-
oriented microscopic images (matching the orientation used for our experimental data, see
below) were co-visualized as 3D data points in a 3D viewer tool (see e.g. **Figure 2A-D**).
Experimental data were selected and coloured according to tracer injection localizations in
the cerebral cortex, to first visualize and compare the pontine distribution of corticopontine
projections from motor and somatosensory areas, and secondly to determine the
organization of projections arising from progressively more medial and caudal locations in
the cerebral cortex, following the cortical neurogenetic gradient that ripples out from the
anterolateral cortex (Leergaard and Bjaalie, 2007; Smart, 1984). Our findings confirm that
the somatosensory and motor neurons of the mouse cortex project to largely separate parts
of the pontine nuclei (Henschke and Pakan, 2020; Inoue et al., 1991; Proville et al., 2014),
with clustered terminal fields topographically distributed in the same concentric fashion
(**Figure 2L-P**) as previously reported in rats (Leergaard et al., 2000a; Leergaard and Bjaalie,
2007). These 3D point data were also used below as supplementary control data, and as
benchmarks for interpreting YFP expression and tract-tracing results in *Nr2f1* mutant mice.

Area-specific layer V neuron distribution in cortices lacking Nr2f1

To assess the influence of cortical area mapping on the establishment of topographical
organization in mouse corticopontine projections, we used *Nr2f1 conditional knock-out* mice
as an experimental model system and their littermates as controls (Alfano et al., 2014;
Armentano et al., 2007). Two well-established conditional *Nr2f1* mouse mutants were used:
the *Nr2f1$^{fl/fl}$::Emx1-Cre* mouse, in which *Nr2f1* expression is abolished from early cortical
progenitor cells at mouse embryonic (E) age 9.5 (Armentano et al., 2007), and the
Nr2f1$^{fl/fl}$::Nex-Cre mouse in which *Nr2f1* expression is inactivated at later stages (E11.5-E12),
solely in cortical postmitotic neurons (Alfano et al., 2014; Goebbels et al., 2006). Both
mouse lines were crossed to the *Thy1-eYFP-H* reporter line to specifically restrict signal
expression to layer V pyramidal neurons (Harb et al., 2016; Porrero et al., 2010). In these
mice, YFP expression follows the physiological distribution of subcortical layer V projections,
including corticospinal and corticopontine fibres, with a high density in motor cortex and a
gradually decreased distribution in S1 and more caudal areas. For simplicity, the lines are
below named *Emx1-cKO* and *Nex-cKO*, respectively.

In agreement with a previous study (Porrero et al., 2010), we observed substantial YFP
signal expression in the hippocampus, tectum, and pontine nuclei, as well as in the globus
pallidus, claustrum, endopiriform nucleus, nucleus of the lateral olfactory tract, mammillary
nuclei, piriform area, and the substantia innominata in adult mice (**Figure 3A**). Signal
expression was also seen in the vestibular nuclei, deep cerebellar nuclei, and cerebellum.
Although YFP expression was present in almost the same cortical regions in 2-months-old
mutant mice as in controls, detailed analysis of layer V expression revealed some distinct
differences in the spatial distribution of *Emx1-cKO* and *Nex-cKO* brains relative to their
respective controls, and between the two conditional lines. We used an ImageJ macro to
automatically count YFP+ nuclei by area of interest in the brain using a threshold based on
intensity and shape of the elements. This allowed to estimate the number of YFP+ neurons
in seven cortical areas (corresponding to prefrontal, motor, somatosensory, auditory, visual
and retrosplenial cortices in control mice) defined by delineations from spatially-registered
overlay images from the *Allen Mouse Brain Atlas* (Wang et al., 2020); **Figure 1; Figure 3B-D'**).
In control animals, the number of YFP+ neurons follows the physiological distribution of
subcortical layer V projections (**Figure 3B, B'**), as previously reported (Polleux et al., 1997;
Porrero et al., 2010; Shepherd, 2009). Strong staining in motor and somatosensory cortices
resulted in bright signal expression in the cerebral peduncle (CP) and CST (**Figure 3A).**

To understand whether the spatial organization of YFP+ layer V cortical neurons was
affected upon loss of cortical *Nr2f1* gradient expression, we quantified and compared the
number of YFP+ cells across mutant and control brains (**Figure 3E, F**). Since previous reports
showed that the distribution of layer V neurons changes in the absence of Nr2f1 (Alfano et
al., 2014c; Armentano et al., 2007a; Tomassy et al., 2010), we expected that the Thy1-YFP
signal would also be altered, as a read-out of layer V changes. Indeed, the layer V gradient
was disrupted in both mutant strains, and YFP+ cells were more homogenously distributed
along the anteroposterior cortical axis (**Figures 3B-F**). While *Emx1-cKO* brains showed a
significant increase of YFP+ cells in parietal and occipital regions, compared to controls, this
was less the case for *Nex-cKO* brains, which instead revealed a decrease only in frontal areas
(**Figure 3F**). To better quantify the differences between corticopontine projections from
frontal and parietal areas, where normally the motor and somatosensory areas develop
(Leergaard et al., 2000a; Leergaard et al., 2000b), we compared the ratio of YFP+ cells over

the total cell populations counted in the frontal (motor) and parietal (somatosensory) areas

(**Figure 3G**). We found that the percentage of YFP+ cells was significantly decreased in

frontal (motor) and concomitantly increased in parietal (somatosensory) in both *Emx1-cKO*

and *Nex-cKO* cortices compared to controls, but no differences were found between the two

lines (Figure 3E, G). This lack of difference in the distribution of YFP+ layer V neurons

between cortical regions normally containing the motor and somatosensory areas, is in line

with the acquisition of a motor-like identity in somatosensory cortex of Nr2f1-deficient

brains, as previously reported (Alfano et al., 2014). We next asked whether and how

alterations in layer V organization observed in the two mutant lines would be translated into

layer V corticospinal projections and/or corticopontine topographic mapping.

Abnormal corticospinal projections and fasciculation in *Nr2f1* mutant brains

We hypothesized that the disordered cortical distribution of YFP-expressing layer V neurons

in mutant mice might influence the integrity of subcortical axonal projections. In all cases,

strong YFP signal expression was seen bilaterally in the main corticofugal pathways (**Figure**

3A), visible as longitudinally oriented fibre bundles coursing towards the CP (**Figure 4A-D)**,

passing dorsal to the pontine nuclei as the longitudinal fasciculus (**Figure 4B'-D')**, and

continuing through the brain stem towards the spinal cord as the CST. Since a large fraction

of the corticobulbar fibres terminate in the pontine nuclei (Tomasch, 1968; Tomasch, 1969),

we reasoned that abnormal distribution of YFP+ layer V neurons observed in mutant mice

(**Figure 3**) might affect corticopontine innervation, and could be reflected in an abnormal

size of the pontine longitudinal fascicle as it enters the CP and exits the pons in rostral and

caudal positions to the pontine nuclei, respectively. To evaluate this, we measured the

dorsoventral width of the longitudinal fascicle of the pons in sequential sections along the

medio-lateral axis. The measurements were taken at rostral and caudal levels to the pontine

nuclei (**Figure 4A'**). Surprisingly, we found the lateral part of the fascicle to be wider at both

rostral and caudal levels in *Nex-cKO* mice compared to *Emx1-cKO* and controls, while being

narrower medially (see red area chart in **Figure 4E, F**), suggesting that the longitudinal

fascicle of the pons is flattened and expanded laterally upon Nr2f1 inactivation in

postmitotic neurons. Only minor differences were observed in the *Emx1-cKO* fascicles (blue

area chart in **Figure 4E, F**). This is also supported by quantification of the total surface of the

longitudinal fascicle of the pons at rostral and caudal levels, which shows a significant
surface reduction at caudal, but only a tendency at rostral levels in *Emx1-cKO* mice (**Figure**
4G, H). These data indicate that loss of Nr2f1 in cortical progenitors, but not in neurons,
results in fewer YFP+ fibres passing the pontine nuclei towards the brain stem to form the
CST.

Moreover, we observed, caudal to the pontine nuclei, abnormally widespread fibre fascicles
in the CST of mutant animals (arrowheads in **Figure 4C', D'**). To determine whether this was
a significant difference between animal groups, we estimated the degree of fibre bundle
fasciculation in the CST of *Emx1-cKO* and *Nex-cKO* mice. At locations of 250µm and 500µm
caudal to the pontine nuclei (**Figure 4A'**), we measured the total dorsoventral width of the
CST at several medio-lateral levels and subtracted the gaps between the YFP-expressing
fibre bundles at the same levels. The ratio of the total width of the CST and fibres was used
as a measure of the fasciculation index (**Figure 4I**). Notably, in both groups of mutant mice,
we found a lower degree of fasciculation in the CST which was more pronounced at the
most caudal level (**Figure 4I**). Together, these data show that loss of *Nr2f1* expression affects
the diameter, shape, and degree of fasciculation of the CST originating from layer V
neurons.

Dual role of Nr2f1 in targeting corticopontine projections

Next, we evaluated whether topographical organization of corticopontine projections
dependents on proper cortical area mapping and layer V distribution (**Figure 5**). To this
purpose, we assessed the spatial distribution of YFP signal expression within the pontine
nuclei by comparing intensity-normalized microscopic images of spatially corresponding
sagittal sections from the brains of *Emx1-cKO*, *Nex-cKO* and control littermate animals
(**Figure 1**B; **Figure 5A**). A complete documentation of spatially comparable and reproducible
images is provided in **Supplementary Figures 1 and 2**. In control brains, we observed a
strong YFP signal in central parts of the pontine nuclei, with the densest expression tending
to surround a centrally located zone exhibiting less dense signals (**Figure 5B-E**). This region
of the pontine nuclei typically receives strong projections from somatosensory areas (green
dots in **Figure 2G-K**). Some signal expression was also visible in medial parts of the pontine
nuclei (**Figure 5D, E**), which is known to primarily receive projections from the cortical motor

areas (purple dots in **Figure 2G-K**). By contrast, signal expression was lower in rostral and
lateral parts of the pontine nuclei (**Figure 5B, C, E**), which are known to receive projections
from visual and auditory areas of the cerebral cortex (Inoue et al., 1991; Leergaard and
Bjaalie, 2007).

Interestingly, *Emx1-cKO* mice showed a relatively homogeneous signal distribution across all
parts of the pontine nuclei, and displayed more expression in the dorsolateral regions
(**Figure 5F-I**). Signal expression was also present in the medial part of the nuclei (**Figure 5G-**
I). This observation fits well with the finding of a higher number of YFP+ neurons in the
occipital cortex (**Figure 3C,C', F**), which projects to the dorsolateral pontine nuclei. By
comparison, *Nex-cKO* animals showed more constrained signal expression and
predominated in rostrally and caudally located clusters extending from the cerebral
peduncle towards the ventral surface of the pons, and medially surrounding a central core
with low expression (**Figure 5J-M**). These clusters were more peripherally located than the
ones observed in control animals (**Figure 5J-M**). Notably, little signal expression was seen in
the central region of the PN of all *Nex-cKO* cases (unfilled arrowheads in **Figure 5J-L**), despite
the presence of YFP+ layer V neurons in somatosensory cortex (**Figure 3D, D', F, G**). This
central region is normally innervated by projections from the face representations located in
S1 (**Figure 2A-F**).

Taken together, these findings show that corticopontine projections are abnormally
distributed in *Nr2f1* cortical deficient mice, with more homogenously (non-specific)
distributed expression in *Emx1-cKO* mice (schematically summarized in **Figure 5N,0**), and
more peripherally distributed signal expression in *Nex-cKO* mice, that display reduced
expression in the central region of the pontine nuclei normally receiving somatosensory
projections (schematically summarized in **Figure 5N,P; Supplementary Figures 1 and 2**). In
both mutant groups, the signal expression was expanded to dorsolateral regions of the
pontine nuclei that normally are innervated by projections from occipital cortical areas. This
suggests that cortical Nr2f1 graded expression in postmitotic neurons might be directly
involved in the establishment of topographically organized corticopontine projections.

Altered somatosensory topographic projections in *Nex-cKO* adult mutant mice

To further support that *Nr2f1* gradient cortical expression might be directly involved in topographical pontine mapping, we decided to focus on the *Nex-cKO* genetic model and unveil the cortical origin of the innervation defect. Compared to the *Emx1-cKO* mice, in which projections appear more disorganized and abundant, we considered the *Nex-cKO* mouse model, exhibiting externally shifted projections, more suitable for investigating the changes in topographical mapping using experimental tract tracing techniques. We injected the AAV9-CAGtdTomato anterograde viral tracer (Pourchet et al., 2021) in the cortex of 5 days-old (P5) *Nex-cKO* mice and littermate controls, in frontal and parietal locations corresponding to the motor or S1 areas in control mice, respectively (**Figures 6, 7**). The mice were sacrificed at P21 and brain sections analysed microscopically (**Figure 1B**). All histological sections were spatially registered to the *Allen Mouse Brain Atlas* (common coordinate framework, CCF3; (Wang et al., 2020)), and the location of tracer injections sites were mapped in the same atlas space (**Figures 6A, 7A**). For each injection site location in a *Nex-cKO* brain, we selected the most corresponding control experiment or wild-type tract-tracing data from the *Allen Mouse Brain Connectivity Atlas* as additional controls. Representative examples of injections and comparisons between controls and littermate *Nex-cKO* brains are shown in **Figures 6B-C' and 7B-C'**. Complete documentation of all tracing experiments cases is provided in ***Supplementary Figures 3-5***.

In controls, the spatial distributions of corticopontine projections (dark grey point clouds in **Figures 6 and 7**) were highly comparable with the labelling patterns seen in corresponding wild-type tracing data from the *Allen Mouse Brain Connectivity Atlas*. As expected, tracer injections into areas corresponding to the motor cortex in control mice gave rise to labelled axonal clusters located rostrally, caudally, and medially in the pontine nuclei (purple dots in **Figure 2G-K**, **Figure 6D-F; Supplementary Figures 3 and 4**). In all *Nex-cKO* mice receiving tracer injections into areas normally corresponding to the motor cortex, the overall distribution of corticopontine labelling was found to be essentially similar to that observed in control cases (compare grey point with red point clouds in **Figure 6D-F; Supplementary Figures 3-5).**

Tracer injections into S1 areas of control animals gave rise to labelled axonal clusters located

centrally or caudally in the pontine nuclei (grey point clouds in **Figure 7D-I).** The spatial

distribution of labelling varied systematically as a function of the location of the cortical

injection sites, such that the 1) face representation in the lateral part of S1 projects centrally

in the pontine nuclei (**Figure 7G,H; Supplementary Figure 3I,J)**, 2) the whisker

representations lateral and more posterior in S1 project to regions surrounding the central

core (**Figure 7F; Supplementary Figure 3K**), while 3) the more medially located forelimb and

hindlimb representations of S1 projected to medio-laterally oriented regions in the caudal

part of the pontine nuclei (**Figure 7D,E; Supplementary Figure 3G,H**). By contrast, tracer

injections in *Nex-cKO* brains, at locations corresponding to control S1 representations of the

whiskers or upper limb, or into the S1/M1 (sensorimotor) lower limb representation, gave

rise to abnormal distribution of corticopontine fibres (**Figure 7D-G**). Specifically, the

corticopontine projections from medial parts of parietal cortex (corresponding to the S1

forelimb and hindlimb representations in control mice) were shifted towards more rostral

locations in *Nex-cKO* experiments (**Figure 7D-F**, red points), resembling the distributions

observed after tracer injections into motor areas in the control experiments (**Figure 6D-F**).

Notably, tracer injections placed in the anterolateral parietal cortex in *Nex-cKO* mice, in

regions normally representing sensory surfaces of the head, gave rise to labelled axons

distributed in the central part of the pontine nuclei, with more subtle difference to the

matching control experiments (**Figure 7G**). In two cases, projections from regions

corresponding to the S1 head representation in control mice were distinctly shifted towards

medial relative to control experiments (red dots in **Figure 7G, and I**), attaining a distribution

resembling the corticopontine projections from head representations in M1 cortex, located

significantly more anteriorly in the cortex (grey dots in **Figure 7I**). Interestingly, two tracer

injections in similar locations, corresponding to the border between the nose and whisker

representations, gave rise to labelling located in central and rostral parts of the pontine

nuclei (**Figure 7F,G**, red dots, see also **Figure 7C'**) with predominantly rostrally-shifted fibres

in one case (**Figure 7F,C'; Supplementary Figure 5J**), and centrally-located and medially-

shifted fibres in the other case (**Figure 7G; Supplementary Figure 5K**). Despite the

differences in density distributions, both cases represent a distinct shift of fibre distributions

towards regions normally receiving motor projections. Taken together, our observations

indicate that corticopontine projections from anterolateral cortex in *Nex-cKO* mice also

display abnormal topographical distributions resembling the normal projections from

homologous representations in the more anterior and medially located primary motor

cortex of control mice. Finally, one tracer injection placed in the most anterolateral part of

S1 in a *Nex-cKO* mouse, at a location corresponding to the perioral surface representations

in controls, yielded corticopontine labelling which was highly similar to that of a control

experiment (**Figure 7H**).

Our findings thus show that corticopontine projections from frontal (motor) areas and the

most anterolaterally located parts of the parietal (S1) cortex are topographically similar in

Nex-cKO brains and controls, whereas corticopontine projections from most parts of parietal

cortex, where the head, whisker, upper limb, and lower limb representations of S1 are

located in control mice, are abnormally shifted towards rostral and medial regions of the

pontine nuclei, that normally receive projections from cortical motor areas (schematically

summarized in Fig. 7J,K). This is in overall agreement with the spatial and temporal control

of Nr2f1 in area mapping. Indeed, the absence of changes in corticopontine projections

from the frontal (motor) cortex might be due to low Nr2f1 expression in this area (spatial

control), whereas lack of changes in the projections originating from the most anterolateral

part of the parietal (S1) cortex, from which the earliest cortical projections to innervate the

forming pontine nuclei originate, might be explained by the late Nr2f1 genetic inactivation

occurring after the earliest layer V neurons have been produced (temporal control).

Discussion

Our present study questions whether and how spatial-temporal cortical expression

gradients are involved in the establishment of normal topographical organization of

corticopontine projections. By combining genetically-modified mice and public mouse brain

connectivity data with tract-tracing techniques and digital brain atlas tools, we have

provided novel evidence of an intrinsic molecular control of layer V cortical neurons during

the establishment of topographical organization of corticopontine projections. Abnormal

areal organization in the neocortex induced by *Nr2f1* inactivation is reflected in altered

corticopontine projections, as well as impaired structural integrity of the CST. While loss of

Nr2f1 from the early progenitor cell pool leads to increased and abnormal corticopontine

innervation at the expense of corticospinal projections, only late postmitotic *Nr2f1*
inactivation reveals altered topographic pontine mapping from medially located parts of
somatosensory cortex controlling whiskerand limb representations. No shifts in projections
from the earliest generated anterolateral cortical areas were observed in these mice, in line
with a spatial and temporal control of Nr2f1 expression, respectively. Overall, our data show
that proper area mapping of the neocortical primordium is a pre-requisite for preserving the
cortical spatial and temporal segregation within the pontine nuclei, and thus correct
corticopontine topographic organization.

Spatial accuracy of topographical data compared across experiments
To ensure accurate 3D data in wild-type and genetically-modified mice, we relied on spatial
alignment of serial microscopic section images to a common reference atlas achieved
through non-linear image registration method (Puchades et al., 2019). The use of non-linear
registration compensated for minor shape differences among brains and allowed
comparison of distribution patterns among spatially relevant data. The focus on the location
rather than the amount of signal expression/axonal labelling also compensated for the
variation in signal expression intensity and size of tracer injections among cases. By
representing signal expression and axonal labelling as 3D point clouds, it became possible to
directly explore and compare location and distribution patterns in 3D in different
combinations of data sets. For the additional benchmark data extracted from the *Allen*
Mouse Brain Connectivity Atlas, we used the same sagittal image orientation as in our
microscopic data, to facilitate comparison of microscopic images in addition to the 3D
comparisons. The relevance and accuracy of the approach was confirmed by demonstrating
that similarly located cortical tracer injections in control animals gave rise to similarly
distributed labelling patterns in the pontine nuclei. Given the distinct patterns of
topographical organization of corticopontine projections, the interpretation and comparison
of spatial distribution patterns and variability in our tract tracing experiments critically
depended on the analysis of tracer injection locations, and a 3D understanding of the
topographical mapping of the cortical surface onto the pontine nuclei, derived from our

analyses of control experiments and earlier studies of the rat corticopontine system

(Leergaard and Bjaalie, 2007).

Mitotic versus postmitotic Nr2f1 functions in layer V corticofugal projections

Our previous work showed overall areal organization impairments in cortical *Nr2f1* mutant

brains, independent of *Nr2f1* inactivation in progenitors or postmitotic neurons. Here, we

report for the first time that *Nr2f1* drives corticopontine connectivity differently in

progenitors *versus* postmitotic neurons. While *Nr2f1* expressed by progenitor cells

modulates the ratio between corticopontine and corticospinal axonal projections, similarly

to what happens in *C. elegans* with the ortholog *UNC-55* (Petersen et al., 2011; Zhou and

Walthall, 1998), postmitotic *Nr2f1* expression specifically acts on somatosensory

topographic organization of corticopontine neurons (**Figures 4, 5**). This suggests that early

Nr2f1 expression in progenitor cells is mainly required in the initial axonal pathfinding of

layer V subtypes, while later postmitotic expression is more implicated in the later

refinement of corticopontine topographical organization. Since YFP+ cells follow the

physiological distribution of subcortical layer V projections (Porrero et al., 2010), Thy1-

driven YFP fluorescence will be induced in cortical subpopulations related to the specificity

of their axonal target region; this tight relationship which will not be altered in the

conditional *KO* models. Accordingly, a higher number of layer V neurons in parietal *Emx1-*

cKO cortices leads to disorganized corticopontine innervation, in accordance with increased

Lmo4 expression, known to drive layer V neurons *versus* the pontine nuclei (Cederquist et

al., 2013; Harb et al., 2016). Differently, *Nex-cKO* parietal axonal projections reach pontine

targets normally innervated by motor-derived cortical areas. Finally, increased number of

YFP+ cells in visual and auditory areas in the occipital cortex, corresponds with an

augmented innervation in dorsolateral regions of pontine nuclei known to receive

projections from the occipital cortex. Together, these data indicate a dual role for Nr2f1 in

layer V corticofugal connectivity, an early role in subtype specification (corticopontine vs

corticospinal), and a later role in topographical mapping.

Revising the chrono-architectonic hypothesis of cortico-pontine circuit development

Previous data in developing rats have shown that pontine neurons settle in the forming

pontine nuclei in a shell-like fashion according to their birthdate, with early born neurons

forming the central core of the pontine nuclei, and later born neurons consecutively settling

around the earlier born neurons forming concentric rings (Altman and Bayer, 1987). At early

postnatal stages, corticopontine axons are chemotropically attracted as collateral branches

from corticospinal axons (Heffner et al., 1990; O'Leary and Terashima, 1988), innervating

the pontine nuclei in a topographic inside-out pattern (Leergaard et al., 1995). Neurons in

the frontal (motor) cortex project rostrally and medially in the pontine nuclei, neurons in the

parietal (somatosensory) cortex project to central and caudal parts, neurons in the temporal

(auditory) cortex to central and lateral regions, and neurons in the occipital (visual) cortex to

lateral and rostral parts of the pontine nuclei (Leergaard and Bjaalie, 2007). This concentric

organization of corticopontine projections suggests that the birthdate of pontine neurons

and their inside-out genesis is linked to the spatial organization of cortical inputs. However,

intrinsic differences in pontine neurons born at different times might also have an

instructive role for corticopontine innervation. A recent study in mice showed that

postmitotic expression of the HOX gene *Hoxa5* guides pontine neurons to settle caudally

within the pontine nuclei, where they are targeted by projections from limb representations

in the somatosensory cortex (Maheshwari et al., 2020). Moreover, ectopic *Hoxa5* expression

in pontine neurons is sufficient to attract cortical somatosensory inputs, regardless of their

spatial position, showing that pontine neurons can play an instructive and attractive role in

topographic input connectivity of corticopontine neurons (Maheshwari et al., 2020).

Nevertheless, maturational gradients in the pontine nuclei cannot fully explain the

complexity of the fine-grained somatotopic topographic connectivity pattern between

cortical input and pontine neuron targets. Since the establishment of topographic maps

requires multiple processes and structures, it is conceivable that the position and specific

intrinsic molecular programs of both presynaptic afferents and postsynaptic target neurons

contribute to this complex corticopontine connectivity map. Indeed, our data show that

without affecting the development and maturation of pontine neurons, corticopontine

Nr2f1-deficient layer V axons originating from the parietal S1 cortex will abnormally target

the pontine region normally deputed to corticopontine motor axons. By contrast, Nr2f1-

deficient axons originating from the frontal and medial cortex will innervate the expected

pontine region allocated to motor axons. This strongly suggests that during the

establishment of corticopontine topography, both structures, the neocortex and the pons

need to be properly pre-patterned by factors involved in spatial and temporal control of
neurogenesis, such as *Nr2f1* for the cortex, and *Hoxa5* for the pontine nuclei (model in
Figure 8).

Conclusion and outlook
By showing that gradient cortical expressions of transcription factor Nr2f1 is directly
involved in the establishment of corticopontine topographic mapping, we provide new
insights into the development of neural systems. However, other factors regulating the size
and location of cortical areas also are likely also implicated. We conclude that distinct
molecular mechanisms in the source (cerebral cortex) and target (pontine nuclei) regions
must be coordinated during the establishment of corticopontine topography. Identifying the
molecular pathways within the cortex and pontine nuclei, as well as the mechanisms and
molecules governing their interaction remains an open question for further studies.

Limitation of the study
This study shows a limited number of viral injections per cortical area that was partially
compensated by adapting mouse tract-tracing data from the Allen Brain Institute to our
analysis. Even through both groups of samples showed a highly consistent pattern of
topographical organization, the two types of data resulted from different experimental
conditions. Because of some variability of the fluorescent signal among samples, we chose
to focus on 3D spatial distribution patterns that turned to be consistent across all control
experiments. We found the *Emx1-cKO* mouse model less suitable for further topographical
analysis due to the disorganized and chaotic innervation pattern observed in the *Thy1-eYFP-*
H reporter background. Although more sophisticated methods are needed to pinpoint the
cellular and molecular mechanisms involved in the establishment of corticofugal
topography, our study represents a useful starting point and resource for further studies of
the development of corticopontine and corticospinal projections in mice.

505
506

507 **Materials and Methods**

Topographical map of corticopontine projections from somatosensory and motor areas

To establish a 3D benchmark map of corticopontine projections from somatosensory and motor areas in in adult wild type mice, we utilized a selection of public experimental tract-tracing data available from the Allen Institute mouse brain connectivity atlas (http://connectivity.brain-map.org/). We selected 11 experiments in which the anterograde tracer EGFP was injected in the right primary/secondary motor cortex (n =6) or primary somatosensory cortex (n = 5) of wild type C57BL/6J mice (**Supplementary Table 1**). Serial two-photon fluorescence images were interactively inspected using the Projection High Resolution Image viewer of the Allen Institute, and from each case, 5 sagittal oriented images of the right pontine nuclei spaced at ~100 µm were captured by screen shot from the largest 3D multiplane thumbnail viewer. The resolution of the captured images was up-sampled three times original size before their spatial alignment to the CCFv3 was optimized using the tools QuickNII (Puchades et al., 2019) and VisuAlign (RRID: SCR_017978), as described below. These images were used to create 3D representations of the axonal labelling in the pontine nuclei (**Figure 2**; see below).

Animals

All mice used were bred in a C57BL/6J background. Male and female animals at any stage of development were used. All experiments were conducted in accordance with the French Animal Welfare Act and European guidelines for the use of experimental animals, using protocols approved by the French Ministry of Education, Research and Innovation and the local ethics committee (CIEPAL NCE/2019–548, Nice) under authorization #15 349 and #15 350. *Nr2f1/COUP-TFI*$^{fl/fl}$ mice were crossed with *Emx1-Cre-recombinase* mice to inactivate *Nr2f1/COUP-TFI* exclusively in cortical progenitors and their progeny (Armentano et al., 2007) or with *Nex-Cre-recombinase* mice to abolish *Nr2f1/COUP-TFI* expression from postmitotic neurons (Alfano et al., 2014). Littermate *Nr2f1/COUP-TFI*$^{fl/fl}$ mice without the presence of the *Cre-recombinase* gene (*Cre-negatives*) were considered controls (**Supplementary Table 2**). For postnatal (P)21 and adult topographic map analysis, *Emx1-cKO* and *Nex-cKO* animals were further crossed with *Thy1-eYFP-H* mice to specifically label layer V projection neurons, as previously reported (Harb et al., 2016; Porrero et al., 2010). Mice were genotyped as previously described (Alfano et al., 2014; Armentano et al., 2007;

538 Harb et al., 2016). Control and mutant littermates were genotyped as $Nr2f1^{fl/fl}$:Thy1-eYFP-
539 $H^{T/+}$ and $Nr2f1^{fl/fl}$:Emx1-Cre:Thy1-eYFP-H$^{T/+}$ or $Nr2f1^{fl/fl}$:Nex-Cre:Thy1-eYFP-H$^{T/+}$, respectively.
540 For simplicity, mutant mice are named *Emx1-cKO* and *Nex-cKO* throughout the text. Midday
541 of the day of the observed vaginal plug was considered as embryonic day 0.5 (E0.5).

Anterograde tracing of corticospinal axons in early postnatal mice

P4-P5 animals were anesthetized on ice for 5 min and kept on ice during the whole
procedure. Viral particles were produced from the AAV9-CAGtdTomato plasmid by the
Alexis Bemelmans (CEA, France) Company, and diluted 1:50 in TE-Buffer (Qiagen, #1018499)
to a final concentration of 1.75e12 vg/ml (kindly donated by I. Dusart, Pierre and Marie
Curie University, Paris, France). Approximately 0.5/1ul was injected unilaterally in different
rostral-caudal and medio-lateral brain locations of control and *Nex-cKO* pups, as previously
described in (Gu et al. 2017).

Microscopic imaging

Mosaic microscopic images were acquired using an Axio Imager M2 epifluorescence
microscope (Carl Zeiss Microscopy GmbH, Jena, Germany) equipped with a halogen lamp, a
MCU 2008 motorized stage, and an EC Plan-Neofluar 10x/0.30 and an AxioCam MRm
camera. ZEN blue software was used for imaging and automatic stitching. Images were
exported in TIFF format and serially ordered from lateral to medial, rotated and if needed,
mirrored to consistent anatomical orientation using Adobe Photoshop CS6 (RRID:
SCR_014199), before being converted to PNG format and resized to 60% of original size
using ImageJ (RRID: SCR_003070) with bilinear interpolation. The resized serial images were
loaded into Adobe Photoshop as a stack, spatially aligned using the ventral surfaces of the
pons and cerebral peduncle as landmarks, cropped and exported as individual PNG files. For
comparative analyses of topographical organization (see below), variations in YFP signal
expression intensity within and between groups were normalized by adjusting the
brightness and contrast of images to equal levels using a custom-made histogram matching
script available for ImageJ (National Institutes of Health; https://imagej.nih.gov/). One
selected, representative case (Experiment 5, Cre-negative, nr: 14250, **Supplementary Table**
2) was used as reference.

Spatial alignment to common 3D reference atlas

Serial sectional images were spatially registered to the *Allen Mouse Common Coordinate Framework*, version 3, 2017 edition of the delineations (CCFv3, (Wang et al., 2020) using the QuickNII software tool (RRID:SCR_016854; (Puchades et al., 2019). Multiple anatomical landmarks (hippocampus, caudate-putamen, inferior and superior colliculus, and external surface of the neocortex) were used to determine the medio-lateral position and orientation of the sagittal section images. For each section image, custom atlas diagrams were aligned to anatomical landmarks in the experimental images using affine transformations, with emphasis on matching the ventral surface of the pons and white matter tracts close to the pontine nuclei and exported as PNG images. As a secondary step, to further optimize registration, the custom atlas images were non-linearly transformed using the software tool VisuAlign v0.8 (RRID:SCR_017978), with particular focus on fitting the template to the outer brain surface, subcortical white matter, and the outer boundaries of the pontine nuclei. To co-display images and the spatially registered custom atlas images, we used the software tool LocaliZoom, which is embedded in the Navigator3 image management system (bit.ly/navigator3), developed and hosted by the Neural Systems Laboratory at the University of Oslo, Norway.

Cortical distribution analysis in *Emx1-cKO* and *Nex-cKO* mutants

Serial section images from *Nex-cKO* and *Emx1-cKO* mutants co-registered to the *Allen Mouse Brain Connectivity Atlas* were analysed using an ImageJ macro allowing automatic counting of spots by area of interest in the brain. The spots are considered as specific staining using a threshold based on intensity and shape of the elements and the composite RGB segmentation atlas as a mask for the region of interest. The macro pre-processes an atlas plate to delineate the regions of interest (ROI), based upon their unique colour-code and corrects the corresponding raw image by subtracting the background generating binary images of the signal and combining it to the ROI maps. Finally, all objects within circularity range of 0.5-1 were counted per ROI using the *find maxima* tool. The process, reiterated for each atlas plate- raw image combination, produced a summary table containing the quantification of particles per area per section, shown as graphs in Figure 3.

596   Corticospinal tract morphometric analysis in *Emx1-cKO* and *Nex-cKO* mutants

Serial section images from *Nex-cKO* and *Emx1-cKO* mutants were analysed using the *Fiji-*
ImageJ Software tool (Schindelin et al., 2015) to determine the total dorsoventral width of
the bundle expressing fluorescent signal in the descending fibre tract in different positions:
rostrally and caudally to the pontine nuclei, and 250 µm and 500 µm caudal to the nuclei.
The width of separate fibre fascicles was also measured 250 µm and 500 µm from the
terminal edge of the pontine nuclei (**Figure 4A'**).

603   Analysis of tracer injection sites

Serial section images of cortical tracer injections in *Nex-cKO* brains (**Supplementary Table 3**)
and experiments taken from the *Allen Mouse Brain Connectivity Atlas*, were spatially aligned
using QuickNII and VisuAlign, as described above. The centre positions of the injection sites
were annotated as a point-coordinate using LocaliZoom, co-displayed with the CCFv3 atlas
in the 3D viewer tool MeshView (RRID:SCR_017222). These visualizations were used to
select spatially corresponding injection site locations for analyses of spatial distribution of
corticopontine projections.

611   Histology, immunohistochemistry and *in situ* hybridization

At P21 and adulthood, animals were anesthetized by intraperitoneal injection of a mixture
of Tiletamine-Zolazepam-Xylazine-Buprenorphine and intracardially perfused with PB Saline
(PBS) followed by 4% paraformaldehyde (PFA) in PBS. Volumes were 15, 20 and 30 ml,
respectively. Brains were removed from the skull and postfixed for 4 h at 4°C in 4% PFA,
before being vibratome-sectioned in 100µm (adult samples) or 150µm (P21 samples) thick
sagittal sections. All sections were incubated overnight at 4°C in a solution of 0.5% Triton X-
100, 3% BSA, 10% goat serum in PBS, for permeabilization and reduction of non-specific
binding of antibodies. For immunofluorescence (IF), sections were incubated for 2 days at
4°C with primary antibodies in a solution of 0.5% Triton X-100, 10% goat serum in PBS, and
then overnight at 4°C with relative secondary antibodies and HOECHST diluted in PBS. For
the complete list of primary and secondary antibodies, see **Supplementary Table 5**. Sections
were   washed   several   times   in   PBS,   then   transferred   on   Superfrost   plus   slides

(ThermoScientific), covered and dried for 30 min to 1 h, and finally mounted with the
Mowiol (Sigma-Aldrich) mounting medium.

Semi-quantitative recording and 3D visualization of spatial distribution patterns

To investigate and compare the 3D distributions of YFP signal expression or anterograde
axonal labelling within the pontine nuclei, we used the annotation functionality in the
LocaliZoom tool to semi-quantitatively record YFP signal expression or labelled axons in all
sections through the pontine nuclei as point coordinates (specified in the coordinate system
of the reference atlas, CCFv3), reflecting the overall density of signal/labelling observed in
the images (**Figure 2C-D**). To compensate for the spacing between sections and allow
inspection of point distributions perpendicularly to the section angle, the z-coordinate of
each point was randomly displaced within the thickness of the gap between sections using a
custom Python script. The point-coordinates were co-displayed in the MeshView 3D viewer
tool (**Figures 2, 5, 6, 7**).

Acknowledgements

We thank Isabelle Dusart in Paris for providing us with AAV9-CAG-tdTomato viral particles
produced by Alexis Bemelmans (CEA, France), Nicolaas Groeneboom and Gergely Csucs for
expert technical assistance, Baptiste Monterroso and Sameh Ben Aicha of the iBV PRISM
microscopy platform for the set-up and optimization of automatic image analysis tools used
in this work, and Sharon Yates, and Maja A. Puchades for useful discussions.

Competing interests:

The authors declare no financial and non-financial competing interests.

Funding

This work was funded by the "Fondation Recherche Médicale; Equipe FRM 2015"
#DEQ20150331750 to M.S.; by an ''Investments for the Future'' LabEx SIGNALIFE (grant
ANR-11- LABX-0028-01) to M.S., by a PhD contract from Région PACA/Inserm and FRM 4[th]
year PhD for C.T, with additional funding from the European Union's Horizon 2020

Framework Program for Research and Innovation under the Specific Grant Agreement No.

785907 (Human Brain Project SGA2), Specific Grant Agreement No. 945539 (Human Brain

Project SGA3), and The Research Council of Norway under Grant Agreement No. 269774

(INCF Norwegian Node) to JGB and TBL.

Data and Code Availability

All microscopic images generated in this project will be shared via the EBRAINS research

infrastructure (https://search.kg.ebrains.eu/) as high-resolution TIFF images under a CC-BY

license, with a unique DOI for each dataset. All tract tracing image data downloaded from

the Allen Institute mouse brain connectivity atlas (Supplementary Table 1) are publicly

available from http://connectivity.brain-map.org/. The custom Python script (spread.py)

used to randomly displace data points within the thickness of a section is available upon

request (to author JGB). The QuickNII (v2.2, RRID: SCR_016854; Puchades et al., 2019)

LocaliZoom and Meshview (RRID: SCR_0170222) tools are available from EBRAINS

(https://search.kg.ebrains.eu/).

References

Alfano, C., Magrinelli, E., Harb, K., Hevner, R. F. and Studer, M. (2014). Postmitotic
control of sensory area specification during neocortical development. *Nature*
communications **5**, 5632.
Alfano, C. and Studer, M. (2012). Neocortical arealization: Evolution, mechanisms and
open questions. *Dev Neurobiol*.
Altman, J. and Bayer, S. A. (1987). Development of the precerebellar nuclei in the rat: IV.
The anterior precerebellar extramural migratory stream and the nucleus reticularis
tegmenti pontis and the basal pontine gray. *J Comp Neurol* **257**, 529-552.
Armentano, M., Chou, S. J., Tomassy, G. S., Leingartner, A., O'Leary, D. D. and Studer,
M. (2007). COUP-TFI regulates the balance of cortical patterning between
frontal/motor and sensory areas. *Nat Neurosci* **10**, 1277-1286.
Assimacopoulos, S., Kao, T., Issa, N. P. and Grove, E. A. (2012). Fibroblast growth factor
8 organizes the neocortical area map and regulates sensory map topography. *J*
Neurosci **32**, 7191-7201.
Badura, A., Schonewille, M., Voges, K., Galliano, E., Renier, N., Gao, Z., Witter, L.,
Hoebeek, F. E., Chedotal, A. and De Zeeuw, C. I. (2013). Climbing fiber input
shapes reciprocity of Purkinje cell firing. *Neuron* **78**, 700-713.
Bertacchi, M., Parisot, J. and Studer, M. (2019). The pleiotropic transcriptional regulator
COUP-TFI plays multiple roles in neural development and disease. *Brain Res* **1705**,
75-94.
Bjaalie, J. G. and Brodal, P. (1989). Visual pathways to the cerebellum: segregation in the
pontine nuclei of terminal fields from different visual cortical areas in the cat.
Neuroscience **29**, 95-107.
Bjaalie, J. G. and Brodal, P (1997). Cat pontocerebellar network: numerical capacity and
axonal collateral branching of neurones in the pontine nuclei projecting to individual
parafloccular folia. *Neurosci Res* **27**, 199-210.
Bower, J. M. (2011). Functional implications of tactile projection patterns to the lateral
hemispheres of the cerebellum of the albino rat: the legacy of Wally Welker. *Annals*
of the New York Academy of Sciences **1225**, 130-141.
Bower, J. M., Beermann, D. H., Gibson, J. M., Shambes, G. M. and Welker, W. (1981).
Principles of organization of a cerebro-cerebellar circuit. Micromapping the
projections from cerebral (SI) to cerebellar (granule cell layer) tactile areas of rats.
Brain, behavior and evolution **18**, 1-18.
Bower, J. M. and Kassel, J. (1990). Variability in tactile projection patterns to cerebellar
folia crus IIA of the Norway rat. *J Comp Neurol* **302**, 768-778.
Brodal, P. (1978). The corticopontine projection in the rhesus monkey. Origin and principles
of organization. *Brain* **101**, 251-283.
Brodal, P. and Bjaalie, J. G. (1992). Organization of the pontine nuclei. *Neurosci Res* **13**,
83-118.
Buckner, R. L. (2013). The cerebellum and cognitive function: 25 years of insight from
anatomy and neuroimaging. *Neuron* **80**, 807-815.
Cadwell, C. R., Bhaduri, A., Mostajo-Radji, M. A., Keefe, M. G. and Nowakowski, T. J.
(2019). Development and Arealization of the Cerebral Cortex. *Neuron* **103**, 980-1004.
Cederquist, G. Y., Azim, E., Shnider, S. J., Padmanabhan, H. and Macklis, J. D. (2013).
Lmo4 establishes rostral motor cortex projection neuron subtype diversity. *J Neurosci*
33, 6321-6332.
Chapin, J. K. and Lin, C. S. (1984). Mapping the body representation in the SI cortex of
anesthetized and awake rats. *J Comp Neurol* **229**, 199-213.
D'Elia, K. P. and Dasen, J. S. (2018). Topographic Maps: Motor Axons Wait Their Turn.
Curr Biol **28**, R86-R88.
Erzurumlu, R. S., Murakami, Y. and Rijli, F. M. (2010). Mapping the face in the
somatosensory brainstem. *Nat Rev Neurosci* **11**, 252-263.

Fabri, M. and Burton, H. (1991). Ipsilateral cortical connections of primary somatic sensory
cortex in rats. *J Comp Neurol* **311**, 405-424.
Feng, G., Mellor, R. H., Bernstein, M., Keller-Peck, C., Nguyen, Q. T., Wallace, M.,
Nerbonne, J. M., Lichtman, J. W. and Sanes, J. R. (2000). Imaging neuronal
subsets in transgenic mice expressing multiple spectral variants of GFP. *Neuron* **28**,
41-51.
Flore, G., Di Ruberto, G., Parisot, J., Sannino, S., Russo, F., Illingworth, E. A., Studer,
M. and De Leonibus, E. (2017). Gradient COUP-TFI Expression Is Required for
Functional Organization of the Hippocampal Septo-Temporal Longitudinal Axis.
Cereb Cortex **27**, 1629-1643.
Fritzsch, B., Elliott, K. L. and Pavlinkova, G. (2019). Primary sensory map formations
reflect unique needs and molecular cues specific to each sensory system. *F1000Res*
8.
Goebbels, S., Bormuth, I., Bode, U., Hermanson, O., Schwab, M. H. and Nave, K. A.
(2006). Genetic targeting of principal neurons in neocortex and hippocampus of NEX-
Cre mice. *Genesis* **44**, 611-621.
Greig, L. C., Woodworth, M. B., Galazo, M. J., Padmanabhan, H. and Macklis, J. D.
(2013). Molecular logic of neocortical projection neuron specification, development
and diversity. *Nat Rev Neurosci* **14**, 755-769.
Harb, K., Magrinelli, E., Nicolas, C. S., Lukianets, N., Frangeul, L., Pietri, M., Sun, T.,
Sandoz, G., Grammont, F., Jabaudon, D., et al. (2016). Area-specific development
of distinct projection neuron subclasses is regulated by postnatal epigenetic
modifications. *eLife* **5**.
Heffner, C. D., Lumsden, A. G. and O'Leary, D. D. (1990). Target control of collateral
extension and directional axon growth in the mammalian brain. *Science* **247**, 217-
220.
Henschke, J. U. and Pakan, J. M. (2020). Disynaptic cerebrocerebellar pathways
originating from multiple functionally distinct cortical areas. *eLife* **9**.
Inoue, K., Terashima, T. and Inoue, Y. (1991). Postnatal development of the pontine
projections from the visual cortex of the mouse. *Okajimas Folia Anat Jpn* **67**, 479-
492.
Leergaard, T. B. (2003). Clustered and laminar topographic patterns in rat cerebro-pontine
pathways. *Anat Embryol (Berl)* **206**, 149-162.
Leergaard, T. B., Alloway, K. D., Mutic, J. J. and Bjaalie, J. G. (2000a). Three-
dimensional topography of corticopontine projections from rat barrel cortex:
correlations with corticostriatal organization. *J Neurosci* **20**, 8474-8484.
Leergaard, T. B., Alloway, K. D., Pham, T. A., Bolstad, I., Hoffer, Z. S., Pettersen, C. and
Bjaalie, J. G. (2004). Three-dimensional topography of corticopontine projections
from rat sensorimotor cortex: comparisons with corticostriatal projections reveal
diverse integrative organization. *J Comp Neurol* **478**, 306-322.
Leergaard, T. B. and Bjaalie, J. G. (1995). Semi-automatic data acquisition for quantitative
neuroanatomy. MicroTrace--computer programme for recording of the spatial
distribution of neuronal populations. *Neurosci Res* **22**, 231-243.
Leergaard, T. B. and Bjaalie, J. G. (2007). Topography of the complete corticopontine
projection: from experiments to principal Maps. *Front Neurosci* **1**, 211-223.
Leergaard, T. B., Lakke, E. A. and Bjaalie, J. G. (1995). Topographical organization in the
early postnatal corticopontine projection: a carbocyanine dye and 3-D computer
reconstruction study in the rat. *J Comp Neurol* **361**, 77-94.
Leergaard, T. B., Lillehaug, S., De Schutter, E., Bower, J. M. and Bjaalie, J. G. (2006).
Topographical organization of pathways from somatosensory cortex through the
pontine nuclei to tactile regions of the rat cerebellar hemispheres. *Eur J Neurosci* **24**,
2801-2812.
Leergaard, T. B., Lyngstad, K. A., Thompson, J. H., Taeymans, S., Vos, B. P., De
Schutter, E., Bower, J. M. and Bjaalie, J. G. (2000b). Rat somatosensory
cerebropontocerebellar pathways: spatial relationships of the somatotopic map of the

774    primary somatosensory cortex are preserved in a three-dimensional clustered
775    pontine map. *J Comp Neurol* **422**, 246-266.
776 **Lemon, R. N.** (2008). Descending pathways in motor control. *Annu Rev Neurosci* **31**, 195-
777    218.
778 **Lillehaug, S., Oyan, D., Leergaard, T. B. and Bjaalie, J. G.** (2002). Comparison of semi-
779    automatic and automatic data acquisition methods for studying three-dimensional
780    distributions of large neuronal populations and axonal plexuses. *Network* **13**, 343-
781    356.
782 **Liu, Q., Dwyer, N. D. and O'Leary, D. D.** (2000). Differential expression of COUP-TFI,
783    CHL1, and two novel genes in developing neocortex identified by differential display
784    PCR. *J Neurosci* **20**, 7682-7690.
785 **Maheshwari, U., Kraus, D., Vilain, N., Holwerda, S. J. B., Cankovic, V., Maiorano, N. A.,**
786    **Kohler, H., Satoh, D., Sigrist, M., Arber, S., et al.** (2020). Postmitotic Hoxa5
787    Expression Specifies Pontine Neuron Positional Identity and Input Connectivity of
788    Cortical Afferent Subsets. *Cell reports* **31**, 107767.
789 **McLaughlin, T. and O'Leary, D. D.** (2005). Molecular gradients and development of
790    retinotopic maps. *Annu Rev Neurosci* **28**, 327-355.
791 **Mihailoff, G. A., Lee, H., Watt, C. B. and Yates, R.** (1985). Projections to the basilar
792    pontine nuclei from face sensory and motor regions of the cerebral cortex in the rat. *J*
793    *Comp Neurol* **237**, 251-263.
794 **Mottolese, C., Richard, N., Harquel, S., Szathmari, A., Sirigu, A. and Desmurget, M.**
795    (2013). Mapping motor representations in the human cerebellum. *Brain* **136**, 330-
796    342.
797 **Nitschke, M. F., Kleinschmidt, A., Wessel, K. and Frahm, J.** (1996). Somatotopic motor
798    representation in the human anterior cerebellum. A high-resolution functional MRI
799    study. *Brain* **119 (Pt 3)**, 1023-1029.
800 **O'Leary, D. D. and Sahara, S.** (2008). Genetic regulation of arealization of the neocortex.
801    *Curr Opin Neurobiol* **18**, 90-100.
802 **O'Leary, D. D. and Terashima, T.** (1988). Cortical axons branch to multiple subcortical
803    targets by interstitial axon budding: implications for target recognition and "waiting
804    periods". *Neuron* **1**, 901-910.
805 **Peterburs, J. and Desmond, J. E.** (2016). The role of the human cerebellum in
806    performance monitoring. *Curr Opin Neurobiol* **40**, 38-44.
807 **Petersen, S. C., Watson, J. D., Richmond, J. E., Sarov, M., Walthall, W. W. and Miller,**
808    **D. M., 3rd** (2011). A transcriptional program promotes remodeling of GABAergic
809    synapses in Caenorhabditis elegans. *J Neurosci* **31**, 15362-15375.
810 **Polleux, F., Dehay, C. and Kennedy, H.** (1997). The timetable of laminar neurogenesis
811    contributes to the specification of cortical areas in mouse isocortex. *J Comp Neurol*
812    **385**, 95-116.
813 **Porrero, C., Rubio-Garrido, P., Avendano, C. and Clasca, F.** (2010). Mapping of
814    fluorescent protein-expressing neurons and axon pathways in adult and developing
815    Thy1-eYFP-H transgenic mice. *Brain Res* **1345**, 59-72.
816 **Pourchet, O., Morel, M. P., Welniarz, Q., Sarrazin, N., Marti, F., Heck, N., Gallea, C.,**
817    **Doulazmi, M., Roig Puiggros, S., Moreno-Bravo, J. A., et al.** (2021). Loss of floor
818    plate Netrin-1 impairs midline crossing of corticospinal axons and leads to mirror
819    movements. *Cell reports* **34**, 108654.
820 **Proville, R. D., Spolidoro, M., Guyon, N., Dugue, G. P., Selimi, F., Isope, P., Popa, D.**
821    **and Lena, C.** (2014). Cerebellum involvement in cortical sensorimotor circuits for the
822    control of voluntary movements. *Nat Neurosci* **17**, 1233-1239.
823 **Puchades, M. A., Csucs, G., Ledergerber, D., Leergaard, T. B. and Bjaalie, J. G.** (2019).
824    Spatial registration of serial microscopic brain images to three-dimensional reference
825    atlases with the QuickNII tool. *PLoS One* **14**, e0216796.
826 **Schindelin, J., Rueden, C. T., Hiner, M. C. and Eliceiri, K. W.** (2015). The ImageJ
827    ecosystem: An open platform for biomedical image analysis. *Molecular reproduction*
828    *and development* **82**, 518-529.

Schmahmann, J. D. and Pandya, D. N. (1997). Anatomic organization of the basilar
pontine projections from prefrontal cortices in rhesus monkey. *J Neurosci* **17**, 438-
458.
Shambes, G. M., Gibson, J. M. and Welker, W. (1978). Fractured somatotopy in granule
cell tactile areas of rat cerebellar hemispheres revealed by micromapping. *Brain,*
behavior and evolution **15**, 94-140.
Shepherd, G. M. (2009). Intracortical cartography in an agranular area. *Front Neurosci* **3**,
337-343.
Smart, I. H. (1984). Histogenesis of the mesocortical area of the mouse telencephalon. *J*
Anat **138 (Pt 3)**, 537-552.
Stoodley, C. J. and Schmahmann, J. D. (2010). Evidence for topographic organization in
the cerebellum of motor control versus cognitive and affective processing. *Cortex* **46**,
831-844.
Tomasch, J. (1968). The overall information carrying capacity of the major afferent and
efferent cerebellar cell and fiber systems. *Confin Neurol* **30**, 359-367.
Tomasch, J. (1969). The numerical capacity of the human cortico-pontocerebellar system.
Brain Res **13, 476-84.**
Tomassy, G. S., De Leonibus, E., Jabaudon, D., Lodato, S., Alfano, C., Mele, A.,
Macklis, J. D. and Studer, M. (2010). Area-specific temporal control of corticospinal
motor neuron differentiation by COUP-TFI. *Proc Natl Acad Sci U S A* **107**, 3576-
3581.
Wang, Q., Ding, S. L., Li, Y., Royall, J., Feng, D., Lesnar, P., Graddis, N., Naeemi, M.,
Facer, B., Ho, A., et al. (2020). The Allen Mouse Brain Common Coordinate
Framework: A 3D Reference Atlas. *Cell* **181**, 936-953 e920.
Welker, C. (1971). Microelectrode delineation of fine grain somatotopic organization of (SmI)
cerebral neocortex in albino rat. *Brain Res* **26**, 259-275.
Woolsey, T. A. and Van der Loos, H. (1970). The structural organization of layer IV in the
somatosensory region (SI) of mouse cerebral cortex. The description of a cortical
field composed of discrete cytoarchitectonic units. *Brain Res* **17**, 205-242.
Zhou, C., Tsai, S. Y. and Tsai, M. J. (2001). COUP-TFI: an intrinsic factor for early
regionalization of the neocortex. *Genes Dev* **15**, 2054-2059.
Zhou, H. M. and Walthall, W. W. (1998). UNC-55, an orphan nuclear hormone receptor,
orchestrates synaptic specificity among two classes of motor neurons in
Caenorhabditis elegans. *J Neurosci* **18**, 10438-10444.
863
864

Figure Legends:

Figure 1. Experimental and analytic workflow. (A) Workflow for generating a 3D

topographic map of corticopontine projections in wild-type mice using tract tracing data

from the Allen Mouse Brain Connectivity Atlas, mapped and compared in a 3D reference

atlas space. Steps performed by the Allen institute are indicated with light grey text in white

boxes. (B) Middle and right columns represent the two paradigms investigated in cKO

models, with the analytic steps performed in adult control, Emx1-cKO and Nex-cKO mutant

animals (middle column), and the tract tracing of the 3D topography of motor and

somatosensory corticopontine projections in young control and Nex-cKO mutants (right

column). All images were spatially registered to the Allen mouse brain atlas (CCFv3; Wang et

al., 2020) prior to analyses, to facilitate comparison of images and spatial distribution

patterns. Results are shown in Figures 2-7.

Figure 2. Topographical organization of corticopontine projections in wild-type mice. (A-D)

Representative example illustrating the data acquisition of corticopontine projections

labelled by viral tracer injection in the S1 face representation of a wild-type mouse from the

Allen Mouse Brain Connectivity Atlas. The tracer injection site centre (A) and anterogradely

labelled axons in the pontine nuclei (B, C) are shown in sagittal microscopic images. (D) Dot

map representation showing semi-quantitatively recorded points corresponding to the

density of labelling observed in the section shown in (C). Panels E,F show the 3D point

populations recorded from the example case, together with a transparent surface rendering

of the right pontine nuclei, seen in view from ventral (E) and medial (F). The S1

corticopontine projections are distributed in dense clusters located centrally in the pontine

nuclei. (G-K and L-P) Differently coloured 3D visualizations of point clouds representing

spatial distribution of anterogradely labelled corticopontine axons derived from 11

experiments available in the Allen Mouse Brain Connectivity Atlas, injected with the

anterograde tracer EGFP in the primary (M1)/secondary (M2) motor cortex (purple dots) or

primary somatosensory (S1) cortex (green dots), at locations indicated with colour coded

circles in G and L. (H, M) 3D visualizations of all points in the right pontine nuclei, colour

coded as indicated in G and L. A grid of transparent grey planes indicate the position and

*orientation of ~100 µm thick digital slices cut with sagittal and frontal orientations through the point clouds, shown in **I-K**, and **N-P**. The red circle in G indicates the tracer injection site shown in **A**. (**G-K**) Point clouds showing that motor and somatosensory areas largely target different parts of the pontine nuclei, with projections from M1 and M2 (purple dots) located more peripherally towards rostral, ventral, and medial than projections from S1 (green dots) as indicated by arrowheads in **I-K**, but also that motor and sensory projections overlap caudally in the pontine nuclei (double arrowheads in **I-K**). (**L-P**) 3D co-visualization of all data points colour-coded in red, yellow or blue according to the location of the cortical injection sites from anterolateral (red) progressively towards medial or posterior (yellow, blue). The slices through the point clouds reveal a concentric arrangement in the pontine nuclei, with projections from the anterolateral parts of the M1/M2 and S1 located centrally and medially (arrowheads in **O-P**), and projections from more medial and posterior cortical locations progressively shifted towards rostral, caudal, and lateral (unfilled arrowheads in (**O-P**). Abbreviations: bfd, barrel field; C, caudal; D, dorsal; ll, lower limb; m, mouth; M1, primary motor cortex; M2, secondary motor cortex; n, nose; R, rostral; S1, primary somatosensory cortex; S2, secondary somatosensory cortex, tr, trunk; ul, upper limb; V, ventral. Scale bars, 1 mm (**A, B**), 200 µm (**C,I**).*

*__Figure 3 – Distribution of YFP-positive layer V pyramidal neurons in cerebral cortex of control, Emx1-cKO and Nex-cKO adult brains.__ (**A**) Fluorescence microscopy image of a representative sagittal section from control brain, showing widespread YFP expression. (**B-D'**) Fluorescence microscopy images of laterally (**B-D**) and medially (**B'-D'**) sagittal sections in Ctrl (**B,B'**), Emx1-cKO (**C,C'**), and Nex-cKO (**D,D'**) brains, with spatially corresponding CCFv3 atlas diagrams superimposed to indicate the location of cortical areas. (**E**) Pie charts illustrating the distribution of YFP+ layer V neurons in different cortical areas along the antero-posterior axis of controls and mutant brains. (**F**) Column graph showing the number of YFP+ neurons across cortical areas in adult Ctrl (grey), Emx1-cKO (blue) and Nex-cKO (red) mice normalized for the total sections analyzed. (**G**) Column graph showing a subsampling of the analysis in **E**. Here, the number of YFP+ layer V neurons in frontal (motor; M) and parietal (somatosensory; S) areas are normalized for the sum of YFP+ cells of M and S regions only, and the values are presented as percentages. Data were analyzed with 2way-ANOVA test*

*(see also **Supplementary Table 4**), and are illustrated as mean ± SEM (Ctrl, n=6; Emx1-cKO,*
n=4, Nex-cKO, n=4). Abbreviations: A, auditory cortex; CA1, cornu ammonis area 1; CB,
cerebellum; CP, cerebral peduncle; CST, corticospinal tract; Cx, cortex; fi, fimbria; GC,
gustatory cortex; LFP, longitudinal fascicle of the pons; M, motor areas (includes primary and
secondary motor cortices); MB, mammillary body; OB, olfactory bulb; PFC, prefrontal cortex;
PN, pontine nuclei; RSC, retrosplenial cortex; S somatosensory areas; SC, superior colliculus;
*Sb, subiculum; Th, thalamus; V, visual cortex. Scale bars, 1mm (**A**); 500μm (**B-D; B'-D'**).*

Figure 4 – Loss of Nr2f1 function leads to abnormal corticospinal projections and
fasciculation. *(**A**) Schematic diagram of a sagittal mouse brain section showing the location*
of the pontine nuclei (PN) and descending fibre tracts (yellow) in the cerebral peduncle (CP),
*longitudinal fascicle of the pons (LFP) and corticospinal tract (CST). (**A'**) Diagram illustrating*
*the different measurements shown in E-I. (**B-D** and **B'-D'**) Fluorescence microscopy images of*
*sagittal sections showing the corticospinal tract entering the LFP (**B-D**) and continuing*
caudal to the pons as the CST. White arrowheads point to the sites of measurement plotted
*in **E** and **F**, respectively. Rostrally, the labelled LFP has similar thickness in the three*
genotypes. Caudally, the CST features defasciculation of fibre bundles (full arrowheads in
***C',D'** point to thinner and more dispersed bundles in Emx1-cKO and Nex-cKO mutants),*
*empty arrowheads point to empty spaces between bundles. (**E, F**) Plots showing LFP*
diameter measurements obtained from lateral to medial before and after innervating the PN
(rostral and caudal respectively). Each measurement represents the average value of
corresponding sections among distinct animals and each position on the x-axis represents a
*specific section of the series. (**G-H**) Column graphs showing average values of the area under*
*the curves in **E-F**. A comparable number of fibres reach the CP in the three genotypes (**G**). In*
Emx1-cKO brains fewer fibres are seen to exit the level of the pons compared to control- and
*Nex-cKO brains (**H**). (**I**) Column graph showing CST fasciculation index, based on*
*measurements of total thickness and fibre thickness (green and purple line respectively in **A'**)*
performed at 250 and 500 μm from the terminal edge of the PN. A ratio between the two
measurements was calculated for each position. Data are represented as mean ± SEM. Data
*were analysed with 2way-ANOVA test (**E-F**) or ordinary one-way ANOVA test (**G-I**) and*
corrected for multiple comparison with the Bonferroni test (see also Supplementary Table 4).
*Ctrl, n=8; Emx1-cKO, n=6; Nex-cKO, n=6. #<0.05 (**E-F**) * < 0.05, **< 0.01, ***<0.005 (**G-I**).*

Abbreviations: CB, cerebellum; CX, cortex; OB, olfactory bulb; SC, spinal cord. Scale bar,
500μm (B').

Figure 5 - Distribution of YFP signal expression in the pontine nuclei of knock-out and
control mice. (A) *3D representation of the outer surface of the brain (transparent grey) and*
pontine nuclei (PN) (transparent brown) in view from ventral. Coloured lines indicate the
*location and orientation of the sagittal sections shown in **B-M**. (**B-D, F-H, J-L**) Fluorescence*
microscopy images of sagittal sections from corresponding medio-lateral levels of the PN,
showing the spatial distribution of YFP signal expression in control, Emx1-cKO, and Nex-cKO
*mice, respectively. (**E, I, M**) 3D visualization of the PN in an oblique view from ventro-medial,*
with point coded representations of signal expression from each of the three sagittal sections
*shown in **B-D**, **F-H**, and **J-L**, respectively. Filled yellow or black arrowheads point to regions*
with increased signal expression in mutant mice, while non-filled arrowheads indicate
*regions with decreased signal expression. In control mice (**B-E**), signal expression is primarily*
*seen in central and caudal parts of the pontine nuclei, while in Emx1-cKO mice (**F-I**) signal*
expression is more widespread and diffuse throughout the entire PN, including more
*peripheral parts towards rostral, ventral and caudal positions (filled arrowheads in **G-I**). In*
*Nex-cKO mice (**J-M**), signal expression is reduced in the central core region of the PN (non-*
*filled arrowheads in **K-M**), while being increased in peripheral (rostral and caudal) regions.*
*(**N-P**) Diagrams summarizing observed changes in corticopontine connectivity upon Nr2f1*
*inactivation. In control mice (**N**), YFP$^+$ corticopontine projections primarily target central part*
of the PN and a substantial number of fibres continue towards the spinal cord. In Emx1-cKO
*mutants (**O**), fewer fibres reach the SC and more projections target the PN (down- and up-*
pointing grey arrows, respectively) and show a pattern of innervation more diffused and
*widespread that in controls. In Nex-cKO animals (**P**), no difference between corticospinal and*
corticopontine projections are detected compared to controls (grey equal sign), but
corticopontine topography appears to be affected, whereby fibres reach more lateral, motor-
receiving PN regions instead of targeting the S1-receiving core (illustrated by grey divergent
arrows). Abbreviations: C, caudal; D, dorsal; L, lateral; M, medial, PN, R, rostral; SC, spinal
cord. Scale bar, 200 μm.

Figure 6. Anterograde tracing of corticopontine projections from frontal (motor) areas in
Nex-cKO mice. *(A) Overview of injection sites in corresponding locations in the right frontal*
*cortex in Nex-cKO (red) and control (grey) brains. The atlas diagrams shown in **A**, **D**, **E**, and **F***
*indicate the control cortical area identities. (**B**, **B'** and **C**, **C'**) Representative microscopic*
*images of the injection site localization (**B-C**) and pontine nuclei innervation (**B'**, **C'**) in the*
*control and Nex-cKO cases reported in panel **F**. (**D-F**) 3D coloured point clouds representing*
axonal labelling in corresponding pairs of Nex-cKO (red) or control/wild-type (dark grey)
mice, shown within a transparent surface representation of the right pontine nuclei in
ventral and medial views. Inset drawings of the brains seen from dorsal show the location of
tracer injection sites for each combination of point clouds. Tracer injections in corresponding
locations in frontal cortex of both Nex-cKO and control/wild-type mice give rise to similar
corticopontine labelling in rostrally located clusters, curving towards ventral and caudal
along the surface of the pontine nuclei. Abbreviations: C, caudal; D, dorsal; M, medial; M1,
primary motor cortex; M2, secondary motor cortex; S1, primary somatosensory cortex; S2,
*secondary somatosensory cortex. Scale bars, 1mm (**B**) 200 μm (**B'**,**D**).*

1004

Figure 7. Anterograde tracing of corticopontine projections from somatosensory areas in
Nex-cKO mice. *(A) Overview of injection sites in corresponding locations in the right parietal*
(primary somatosensory; S1) cortex in Nex-cKO (red), control (grey), and wild-type (grey with
*white cross) brains. (**B**, **B'** and **C**, **C'**) Representative microscopic images of the injection site*
*localization (**B-C**) and pontine nuclei innervation (**B'**, **C'**) in the control and Nex-cKO cases*
*reported in panel **F**. (**D-I**) 3D coloured point clouds representing axonal labelling in*
corresponding pairs of Nex-cKO (red) or control/wild-type (grey) mice, shown within a
transparent surface representation of the right pontine nuclei in ventral and medial views.
Inset drawings of the brains seen from dorsal show the location of tracer injection sites for
each combination of point clouds. Corresponding tracer injections in corresponding locations
in the medial parietal cortex of Nex-cKO and control/wild-type mice give rise to labelling in
different parts of the pontine nuclei, with corticopontine projections in control mice
*distributed in elongated curved clusters located caudally (grey points in **D,E**) or laterally in*
*the pontine nuclei (grey points in **F**), while projections from the same locations in Nex-cKO*
mice are shifted to more peripheral rostral and lateral parts of the pontine nuclei (red points

in **D-F**). All tracer injections in anterolateral part of the parietal cortex of Nex-cKO and control mice, in regions corresponding to the S1 face representation in wild-type mice, gave rise to labelling in the central region of the pontine nuclei, however with a subtle medial shift of projections in Nex-cKO brains (**G**, see also **I**). Corresponding tracer injections in the most anterolateral part of S1 in a Nex-cKO and control gave rise to highly similar labelling, centrally in the pontine nuclei. (**I**) Tracer injections in widely separated locations in S1 (Nex-cKO) and M1 (wild-type control) gave rise to largely corresponding labelling in the medial part of the central core region of the pontine nuclei, albeit with additional rostral and medial labelling in the Nex-cKO experiment. (**J-K**) Diagrams summarizing observed changes in corticopontine connectivity upon Nr2f1 postmitotic inactivation. In control mice (**J**), projections from motor areas (M2/M1, pink) and S1 (purple) target largely segregated parts of the pontine nuclei, with somatosensory projections targeting the central core region, while motor projections target more peripheral rostral, caudal, and medial parts of the pontine nuclei. In Nex-cKO animals, corticopontine topography of S1 is affected, whereby fibres reach lateral, motor-receiving PN regions, instead of targeting the core (illustrated by grey divergent arrows). Abbreviations: C, caudal; D, dorsal; M, medial; M1, primary motor cortex; M2, secondary motor cortex; PN, pontine nuclei; S1, primary somatosensory cortex; S2, secondary somatosensory cortex, SC, spinal cord. Scale bars, 1mm (**B**), 200 μm (**G**).

Figure 8.   Model of corticopontine topography establishment and changes upon Nr2f1 cortical abolishment. (A) Proposed schematic model of how the neocortex (Ncx) and the pontine nuclei (PN) might interact during development in driving corticopontine topography. Both structures need to be pre-patterned by gradient expression of transcription factors. While postmitotic pontine graded expression of Hoxa5 will impart an anterior to posterior identity to pontine neurons (graded blue circles) (Maheshwari et al., 2020), postmitotic cortical gradient expression of Nr2f1 (purple gradient in Ncx) will intrinsically instruct corticopontine neurons to innervate their topographically proper targets (this study). (**B**) In the absence of postmitotic Nr2f1 gradient expression in the neocortex, but maintenance of Hoxa5 expression in the PN, axons from motor and somatosensory cortex will both innervate rostrally-located clusters within the PN, since somatosensory corticopontine projections in

1050 *Nex-cKO mice are changed to resemble motor-like projections. Abbreviations: Cb,*
1051 *cerebellum; CP, cerebral peduncle; SC, spinal cord.*

A

CX

OB

CP — LFP

PN

CST

SC

CB

A'

a (E-G)

b (F-H)　c, c' (K)

d, d' (K)

250μm　500μm

······ Guide lines
━━━ Measurement lines

B *Ctrl*

CP

LFP

PN

D↑→C

Lateral

C *Emx1-cKO*

D *Nex-cKO*

B'

CP

LFP

CST

PN

Medial

C'

D'

E **Rostral**

a

LFP diameter (μm)

Lateral　　　Medial

Ctrl
Emx1-cKO
Nex-cKO

F **Caudal**

b

LFP diameter (μm)

Lateral　　　Medial

G **Rostral area**

a

μm²

H **Caudal area**

b

μm²

***

I **Fasciculation index**

c'/c　d'/d

*　*

*　***

250μm　　500μm

Ctrl
Emx1-cKO
Nex-cKO

A Control, Emx1-cKO, Nex-cKO

B,F,J C,G,K D,H,L

Control

Emx1-cKO

Nex-cKO

Lateral ←——————————→ Medial

B **C** **D** **E**

Control

F **G** **H** **I**

Emx1-cKO

J **K** **L** **M**

Nex-cKO

D → C

D, L → C

N *Control* **O** *Emx1-cKO* **P** *Nex-cKO*

PN   SC

A

Gradient expression of *Nr2f1*

Somatosensory areas

Motor areas

Ncx

Tectum

CP

Cb

PN

SC

Ctrl

Gradient expression of *Hoxa5*

B

Postmitotic inactivation of *Nr2f1*

Somatosensory areas

Motor areas

Ncx

Tectum

CP

Cb

PN

SC

Nex-cKO

Gradient expression of *Hoxa5*

▲ *Nr2f1+* corticopontine projection neurons
▲ *Nr2f1-* corticopontine projection neurons
● *Hoxa5+* pontine neurons