Transient exposure of a buried phosphorylation site in an autoinhibited protein

Simone Orioli[1,2] **and Kresten Lindorff-Larsen**[2*]

*For correspondence:
lindorff@bio.ku.dk (KLL)

[1]Structural Biology and NMR Laboratory & Linderstrøm-Lang Centre for Protein Science, Department of Biology, University of Copenhagen, Denmark; [2]Structural Biophysics, Niels Bohr Institute, Faculty of Science, University of Copenhagen, Denmark

Abstract Autoinhibition is a mechanism used to regulate protein function, often by making functional sites inaccessible through the interaction with a cis-acting inhibitory domain. Such autoinhibitory domains often display a substantial degree of structural disorder when unbound, and only become structured in the inhibited state. This structural dynamics makes it difficult to study the structural origin of regulation, including effects of regulatory post-translational modifications. Here, we study the autoinhibition of the Dbl Homology domain in the protein Vav1 by the so-called acidic inhibitory domain. We use molecular simulations to study the process by which a mostly unstructured inhibitory domain folds upon binding and how transient exposure of a key buried tyrosine residue makes it accessible for phosphorylation. We show that the inhibitory domain, which forms a helix in the bound and inhibited stated, samples helical structures already before binding and that binding occurs via a molten-globule-like intermediate state. Together, our results shed light on key interactions that enable the inhibitory domain to sample a finely-tuned equilibrium between an inhibited and a kinase-accessible state.

Introduction

Regulation of protein activity is crucial for cellular function, and the activity of a large number of proteins is regulated by *cis*-acting elements or modules (*Pufall and Graves, 2002*). When such modules interact with a protein in a way to inhibit its function, this is termed autoinhibition. This mechanism is widespread and can be employed to attenuate DNA-protein or protein-protein interactions (*Dombroski et al., 1993*; *Ko and Prives, 1996*; *Stöven et al., 2000*; *Lee et al., 2005*), to regulate enzymatic activity (*Hubbard, 2004*) or even to impact protein localisation (*Pearson et al., 2000*). Schematically, autoinhibited proteins are usually composed of two main modules, an inhibitory module (IM) and a functional domain (FD), which is regulated by the IM itself (*Trudeau et al., 2013*). The inhibition can either be allosteric in nature or occur by physically hindering the access to the FD's binding or catalytic site. As a consequence, autoinhibited proteins populate at least two distinct states: the inactive (autoinhibited) state, where the IM is bound to the FD, and an active state where the FD and IM do not interact directly.

Conformational transitions between the inactive and the active states of an autoinhibited protein often requires some degree of flexibility in the region connecting the IM with the FD (*Trudeau et al., 2013*), and in some cases the IM itself has been reported to be intrinsically disordered in the active state, adopting a stable structure only upon binding to the FD (*Kim et al., 2000*; *Pufall et al., 2005*). Recent studies have also shown how autoinhibited proteins are enriched in intrinsic

disorder (*Trudeau et al., 2013*). Among other important implications, this fact makes these systems useful examples to study so-called folding-upon-binding equilibria in intrinsically disordered proteins (IDPs) or regions (IDRs), a central challenge in structural biology (*Robustelli et al., 2020*; *Malagrinò et al., 2020*).

IDPs and IDRs populate several structurally heterogeneous conformations when unbound (*Van Der Lee et al., 2014*; *Habchi et al., 2014*), but are nonetheless often able to form highly specific interactions with their binding partners, often (but not always (*Schuler et al., 2020*)) through a process called folding-upon-binding (*Dyson and Wright, 2002*). The characterisation of the binding process at the atomic level would help provide insight into the process and a deeper understanding of how the physical and chemical properties of the IM and FD (sequence, charge distribution etc.) influence their biology. While e.g. NMR and single-molecule fluorescence spectroscopy has provided a wealth of details on the structural properties of bound and unbound states, and the kinetics of interconversion, detailed structural studies of the actual binding process are difficult to achieve by experiments (*Kim et al., 2018*; *Sturzenegger et al., 2018*). On the other hand, all-atom molecular dynamics (MD) simulations have the ability to resolve binding processes at high spatial and temporal resolution, and have been used before to characterize folding-upon-binding processes (*Piana et al., 2013*; *Robustelli et al., 2020*; *Henriques and Lindorff-Larsen, 2020*). In this context it is worth noting that the tethering of an ID to a FD may increase the local concentration of a disordered region (*Sgourakis et al., 2010*), thus making *cis*-acting regulatory regions particularly attractive targets to study such processes.

Here we focus on the Dbl Homology (DH) domain of the protein Vav1 as an example of a protein that is regulated via an intrinsically disordered IM. Vav proteins are large multi-domain proteins whose main function is to act as a guanine nucleotide exchange factor (GEF) for Rho family GTPases (*Bustelo, 2001*; *Tybulewicz, 2005*). Vav proteins contain eight domains, of which the DH domain holds the GEF activity (*Abe et al., 1999*), with the remaining domains regulating function either through modulating activity or localization (*Bustelo, 2001*; *Tybulewicz, 2005*). Just N-terminally to the DH domain is found the so-called acidic (Ac) domain, and the structure of Vav1 (*Aghazadeh et al., 2000*) revealed that part of the Ac domain blocks the active site in the DH domain forming an autoinhibited state (Fig. 1A). This autoinhibition is recapitulated in a minimal construct, termed AD, consisting of a short helical region of the Ac (the inhibitory module) and the DH (the functional domain) domains. This construct has a ∼10 fold decreased GEF activity compared to the DH domain alone (*Li et al., 2008*). The presence of additional domains in Vav decreases activity by an additional 10 fold, but NMR and biochemical experiments have shown that the AD construct captures the main mechanism of inhibition and activation, and that the remaining domains merely modulate these (*Yu et al., 2010*; *Li et al., 2008*).

Activation of Vav1 occurs by phosphorylation of Y174 (Fig. 1A), located centrally in the Ac region. In the autoinhibited state, Y174 is however buried in the interface between the Ac and DH domains (Fig. 1B), and thus inaccessible to the Src-family kinase that phosphorylates it. Through a series of NMR and biochemical experiments, it was previously shown that the Ac region—including Y174— exists in a dynamic equilibrium between a buried state and an exposed, kinase accessible state with a 10:1 population ratio (*Li et al., 2008*). When Y174 is phosphorylated, the active state becomes dominant, thus explaining the 10-fold increase in activity upon phosphorylation. These observations strongly suggest a regulatory mechanism in which the transient exposure of at least part of the Ac region leads to access of the kinase to Y174; once phosphorylated, the Ac domain remains dissociated and the protein is active (Fig. 1B). The dynamics of the Ac helix in the protein's active state and the details of the conformational change, however, remain elusive and unexplained.

We have here examined the transient exposure of the autoinhibitory domain in Vav1 through MD simulations. In particular, we have asked the question whether it is possible to observe such slow and finely tuned motions through all-atom simulations with the goal to unveil the mechanism behind them. The NMR experiments showed that the closed/open equilibrium is fast on the NMR timescale, yet sufficiently slow to give rise to substantial relaxation dispersion, suggesting that the

Figure 1. (A) Cartoon representation of the crystal structure of Vav1, coloured from the N-terminus (red) to C-terminus (blue). The box shows details with the short helix in the Ac domain, highlighting key residues in the interaction between the Ac and DH domains. (B) A schematic representation of the equilibrium between bound and unbound structures of the Ac inhibitory region. Y174 is depicted as blue sticks. (C) A time-series of the three collective variables (CV1–3) that we used to analyse the 1 μs-long MD simulation started from the Vav1 crystal structure. CV1 quantifies the fraction of native contacts between the Ac-helix and the DH domain, CV2 probes the native contacts in the Ac-helix, and CV3 measures the fraction of native contacts between Y174 and the surrounding residues (within a 6Å radius in the crystal structure).

process occurs in the microsecond to millisecond timescale (*Li et al., 2008*). Such slow processes
are often important biologically, but remain difficult to study by simulations. It has previously been
shown that enhanced sampling simulations can be used to capture slow motions on the microsec-
ond to millisecond timescale. In particular, MD simulations coupled with metadynamics (*Bussi and*
Laio, 2020) were able to capture motions observed by NMR in both Cyclophilin A (*Papaleo et al.,*
2014) and the L99A variant of T4 lysozyme (*Wang et al., 2016*). In both cases, however, detailed
structural information was available for both the dominant state as well as the transiently popu-
lated state, and this information was used to guide the simulations. Similarly to the recently studied
ternary complex p27 (*Henriques and Lindorff-Larsen, 2020*), however, for Vav1 less is known about
the structural origin underlying the dynamic motions, so no conformation is available for the active
state of Vav1 that can be employed to simplify the sampling.

We collected over 60 μs of sampling combining unbiased MD and metadynamics simulations
to reveal how the binding of the inhibitory module to the DH domain occurs through a dominant
pathway where the short AC-helix (residues 169-177) forms most of its internal contacts before
binding to its hydrophobic binding pocket. This provides an indication that, in the active state, the
inhibitory module of Vav1 lives in an equilibrium between a helical and an unfolded state and it is
therefore able to retain secondary structure without binding to the DH domain. Moreover, we show
through short MD simulations that the AD interface is dynamic on the microsecond timescale and
the Ac-helix is in a fast equilibrium between a tightly bound and a molten globule-like pre-bound
state.

Results and Discussion

The Ac helix is bound to the DH domain during microsecond MD simulations

We first performed multiple 1 μs-long MD simulation of the Vav1 AD construct, consisting of the Ac
region and the DH domain, and monitored the structure and dynamics of the two domains and the
interface between them. We carried out two simulations (with two different force fields) starting
from the crystal structure and two from the NMR structure. We observe substantial dynamics in
simulations initiated from the NMR structure (Fig. S1, Fig. S2 and discussion in the supplementary
material), in apparent disagreement with NMR experiments suggesting the inhibited state to be
stable on this time scale. In contrast, simulations initiated from the crystal structure were stable,
and we thus focus on those below.

Calculations of the root-mean-square fluctuations, the fraction of native contacts and visual in-
spection of the trajectory revealed the protein to be very stable with only substantial motion in a
loop region and in the tip of the N-terminal region (Fig. S1). The Ac region and the AD interface
were also overall very stable as monitored by three distinct collective variables (see Supporting
Information for a detailed description): CV1 which quantifies the fraction of native contacts at the
interface between the Ac and DH domains in AD, CV2 quantifies the amount of helical structure
in the Ac-helix, and finally CV3 quantifies the interactions between the side chain in residue Y174
(whose phosphorylation leads to activation of Vav1) and residues in the binding site (T205, Y209,
V328, Q331 and R332). As expected from the NMR experiments (*Li et al., 2008*), the behavior of
these collective variables (especially CV3) confirms that Y174 remains firmly bound during the sim-
ulation (Fig. 1C).

Metadynamics simulations show the AD complex is formed through a dominant binding pathway

135 To be able to probe the free energy landscape describing the motions involved in the transient
136 exposure of Y174, we turned to enhanced sampling simulations. Indeed, the NMR experiments (*Li*
137 *et al., 2008*) suggest that the motions of the inhibitory module occur on the μs to ms scale: this
138 goes beyond the timescales that can be easily explored using standard MD simulations. Based on
139 previous experience studying processes on similar time scales (*Papaleo et al., 2014*; *Wang et al.,*

140 *2016*; *Henriques and Lindorff-Larsen, 2020*) we opted for metadynamics as an enhanced sampling
141 tool to accelerate the exploration of the Ac-helix binding-unbinding equilibrium. Metadynamics
142 employs a time-dependent potential, defined along a set of predetermined CVs, to accelerate the
143 sampling of rare events. The choice of CVs is critical and far from trivial, and here is further com-
144 plicated by the fact that little information is available on the unbound state of the Ac helix. For
145 our first metadynamics run, the natural choice was therefore to bias along CV1 and CV2, as a simi-
146 lar choice of CVs has been previously employed to analyse folding-upon-binding dynamics of IDPs
147 onto folded domains (*Robustelli et al., 2020*). During this simulation (see Fig. S3A-C) we were able
148 to observe multiple binding-unbinding events during the first 4 μs, but the Ac-helix did not rebind
149 correctly for the remaining 6 μs of the run, and therefore the simulation failed to converge to a
150 meaningful free energy landscape (see also Fig. S3D and the related discussion).

151 In order to enhance the sampling along slow degrees of freedom that were not included in this
152 first metadynamics run, we used parallel bias metadynamics (PBMetaD, *Pfaendtner and Bonomi*
153 (*2015*)) to enable us to use more than two CVs. Indeed, it has been previously shown that this ap-
154 proach may be used to capture slow motions in proteins when many CVs need to be employed
155 (*Heller et al., 2017*; *Prakash et al., 2018*; *Fu and Pfaendtner, 2018*; *Henriques and Lindorff-Larsen,*
156 *2020*). In contrast to standard well-tempered metadynamics, where Gaussian hills are deposited
157 on a multi-dimensional grid with dimension of the number of CVs used, PBMetaD works by intro-
158 ducing multiple one-dimensional and time-dependent biases, one per CV. In this way, sampling is
159 enhanced but the computational efficiency of the simulation is not severely affected by the num-
160 ber of CVs used. At the same time, one dimensional free energies can be calculated from the
161 sum of the deposited Gaussian hills, while the multidimensional free-energy surface can be de-
162 termined via reweighting techniques (*Pfaendtner and Bonomi, 2015*). We employed 5 CVs (CV1–3
163 and two others, capturing different aspects of the Y174 binding and Ac-helix folding and unfolding
164 dynamics; see the supplementary information for further details) and two parallel walkers in our
165 PBMetaD run in order to enhance the sampling of unbinding and rebinding conformational tran-
166 sitions. We collected a total of 10 μs of sampling (5 μs per walker), during which we were able to
167 sample multiple unbinding-rebinding events (Fig. S4 and S5).

168 We performed a block error analysis (*Flyvbjerg and Petersen, 1989*; *Bussi and Tribello, 2019*) in
169 order to assess the convergence of the CVs in the PBMetaD simulation, and such analysis shows
170 that the estimated errors plateau at block sizes around 0.5 μs (Fig. S6A) and indicate the conver-
171 gence of the simulation. This is furthermore corroborated by the fact that all the binding-unbinding
172 transitions were sampled while the heights of the deposited Gaussians were already very small
173 (Fig. S6B–F).

174 Our PBMetaD simulations of the Vav1 AD construct resulted in a complex free energy surface
175 with multiple minima and distinct states between fully unbound and bound structures (Fig. 2A).
176 The main binding pathway, as indicated by the free energy landscape along CV1 and CV2, starts
177 from an extended state where the the Ac-helix is unbound and unfolded (Fig. 2A, **1**) and progresses
178 through a state with increased helical content in the 169-178 region (Fig. 2A, **2**). Binding of the Ac
179 region to the DH domain is initiated by the insertion of I173 into the hydrophobic pocket formed
180 by F320, L325 and V328 (Fig. 2A, **3** and inset), an interaction which stabilises the positionining of the
181 helix close to the binding site. The formation of the fully bound AD complex progresses through
182 the establishment of native contacts between Y174 and T205, Y209, V328, Q331 and R332 (Fig. 2B, **3**
183 and **4**), forming a molten globule-like state with the helix locked in place (Fig. 2A, **4**). The final bound
184 conformation (Fig. 2A, **5**) is reached through the consolidation of the remaining contacts, especially
185 in the elbow region (M178, T205 and K208), while Y174 ultimately settles into the hydrophobic
186 binding pocket (Fig. 2B, **5**).

187 While examining the free energy landscape along CV1 and CV2 provides an intuitive overview
188 of the binding process, the binding mechanism of the Ac inhibitory module is a complex process
189 on a the high-dimensional surface. For this reason, it is important examine the process from the
190 angle of multiple CVs. In the free energy landscape projected onto CV1/CV3 (Fig. 3A) and CV2/CV3

Figure 2. Free energy landscape and binding mechanism for the Ac helix to the DH domain. (A) Free energy surface, projected along CV1 and CV2, obtained from the dual-walker PBMetaD simulation. The minimum of the free energy surface has been set to zero energy, and low energy regions are indicated in blue, whereas high energy regions are shown in red. The ensembles (**1**–**5**) with 10 protein structures were extracted from the free energy region indicated by the dashed arrows (details in supporting information) and are used to help visualize the process of binding. The inset highlight residue I173 in representative configurations extracted from ensemble **3**. (B) Zoom in on residue Y174 (blue) for representative configurations extracted from ensembles **3**–**5**.

Figure 3. Free energy surface, projected along (A) CV1 and CV3 and (B) CV2 and CV3, obtained from the dual-walkers PBMetaD simulation. The minimum of the free energy surface has been set to zero energy, and low energy regions are indicated in blue, whereas high energy regions are shown in red. The coloured circles on the two plots indicate the positions of the representative configurations in Fig. 2 projected on to the corresponding free energy landscapes. The dashed arrows highlight the main binding pathways and are meant only as a guide for the eyes.

(Fig. 3B) we observe that Y174 binding is neither strongly related to the perfect binding of the Ac-
helix nor to its helical state. This means that pathways exist (the leftmost ones on the CV1/CV3 and
CV2/CV3 landscapes) through which the AD complex can be rendered inactive, by sequestering
Y174 from the solvent, without the need for the inhibitory module to reach a fully bound confor-
mation. On the other hand, the projection of the representative configurations along the Ac-helix
binding pathway (Fig. 2A) onto these free energy landscapes (Fig. 3) clearly shows that Ac-helix bind-
ing appears to follow a well defined pathway where CV3 changes cooperatively with CV1, while CV2
only changes initially and its value remains almost constant for the rest of binding dynamics. For
sake of completeness, we also report the one-dimensional free energies along the remaining two
CVs employed in the PBMetaD simulation (Fig. S7).

Summarising, visual inspection of the free energy landscape (Fig. 2A) suggests that the forma-
tion of the bound AD complex mainly progresses through a pathway where the Ac-helix consoli-
dates most of its helical contacts while the the inhibitory module and the DH domain do not interact,
and then binds to the hydrophobic patch formed by T205, Y209, F320, V328, Q331 and R332 (see
supplementary video 1). Our simulations suggest that the inhibitory helix may fold independently
of the interaction with the binding partner, i.e. the DH domain, and thus does not appear to follow
an induced fit/folding-upon-binding mechanism. Indeed, such a mechanism would appear in the
free energy landscape as either diagonal or L-shaped pathways in the bottom-right hand side of
the CV1-CV2 plot. Moreover, the analysis of the free energy landscapes show that multiple binding
pathways exist that make Y174 inaccessible to phosphorylation by Src kinase (Fig. 3).

Short MD simulations confirm the binding pathway predicted by PBMetaD

As an enhanced sampling technique, the convergence of metadynamics simulations depends on
the choice of CVs. If important CVs are not included one may still drive the transition via the bias
potential, but simulations will not converge and transitions will not be observed as the bias is
decreased. Presumably, this is what occurred in our initial set of simulations (see Fig. S3A-C). In
contrast, our PBMetaD simulations appear to converge and we observe transition also towards
the end of the simulations when the bias had decreased (Fig. S6). Nevertheless, to validate our
PBMetaD simulations and the conformational states, we selected several configurations along the
binding pathway and used them as a starting point for multiple unbiased MD simulations (between
300 and 500 ns in length, see supporting information for further details). The behaviour of these
shorter MD simulations, when projected onto the PBMetaD landscape along CV1 and CV2, strikingly

Figure 4. Projection of multiple unbiased MD simulations, shown as white dots, on to the free energy landscape from PBMetaD and defined by CV1 and CV2. The MD simulations are grouped as follows: (A) simulations sampling the (A) bound state, (B) the molten globule-like state or the (C) unbound state; simulations performing a transition towards the (D) unbound state, (E) bound state or (F) the molten globule state from different locations along the free energy landscape. The red dots indicate the starting points of the simulations; (G) Zoom in on residue Y174 (blue) for the selected frames along an unbiased binding trajectory reported in (E).

agrees with our expectations. In particular, they either sample the local and stable conformational
states predicted by PBMetaD (Fig. 4A-C) or follow low free energy pathways to sample multiple
states (Fig. 4D-F). These simulations also showed a full unbinding event starting from a partially
bound structure, or even the transition to a fully bound structure (Fig. 4D,E). Visual inspection of
these simulations (supplementary videos 2 and 3), further consolidate our interpretation of the
binding pathway provided by PBMetaD. One of these unbiased simulations (Fig. 4E) resulted in the
binding of the the Ac helix and Y174 to the binding pocket (Fig. 4G): At $t = 0$ ns, the helix is held
close to the binding site by the interaction of I173 with the F320, L325 and V328 hydrophobic pocket,
while Y174 is tilted pointing outwards with respect to the binding site; a rotamer jump around $t = 90$
ns brings Y174 closer to the native pocket and the corresponding interaction between Y174 and
R332 is established shortly after ($t = 150$ ns). Finally, around $t = 220$ ns, the N-terminus of the helix
tilts upwards to reach the native conformation. In addition to this structural view of the binding
process, we conclude that the overall behaviour of MD simulations projected onto CV1 and CV2 is
in agreement with the shape of the free energy profile obtained from PBMetaD, further validating
that it represents the dynamics of Vav1.
Inspection of the free energy landscape from the PBMetaD simulations reveals a barrier ($\sim$

3.5 kT, approximately symmetric on both sides) in the CV1-CV2 free energy surface, which sepa-
rates the molten globule state **4** from the fully bound state, **5** (CV1~0.95). We hypothesized that
this would correspond to a transition state towards the fully bound structure, and thus selected
five conformations with CV1~0.95 and CV2 values ranging from 14.3 to 15.4 and ran 5 indepen-
dent unbiased MD simulations starting from each of these structures. We let each of these 25
trajectories run until they either reached state 5 (CV1 $\geq$ 0.98) or state 4 (CV1 $\leq$ 0.82) and evaluated
the binding probability (p_{binding}) as the ratio between the number of simulations reaching state **5**
and the total number of trajectories. In good agreement with the PBMetaD predictions, we found
$p_{\text{binding}} = 0.48 \pm 0.05$ (see supporting material for the estimation of the uncertainty), consistent with
the value $p_{\text{binding}}^{\text{opt}} = 0.5$ expected for a transition state (*Du et al., 1998*).

We also used the simulations initiated from the transition state region to estimate the barrier-
crossing (transition path, TP) time and, based on this, the mean first passage (MFP) time between
the two states. To this end, we considered only the reactive portions of the 25 simulations and
estimated the forward, $\langle \tau_{\text{TP}}^{\text{F}} \rangle$, and backward, $\langle \tau_{\text{TP}}^{\text{B}} \rangle$, mean transition path times by averaging the time
lengths of the, respectively, **4**→**5** and **5**→**4** transition paths. We found $\langle \tau_{\text{TP}} \rangle = \langle \tau_{\text{TP}}^{\text{F}} \rangle + \langle \tau_{\text{TP}}^{\text{B}} \rangle = 50 \pm 13$
ns. We then estimated the MFP time based on the relationship between the mean TP time and the
MFP time: (*Hummer, 2004*; *Chung et al., 2009*)

$$\tau_{\text{MFP}} = \frac{2\pi \langle \tau_{\text{TP}} \rangle}{\log(3\beta \Delta G^{\dagger})} e^{\beta \Delta G^{\dagger}}, \tag{1}$$

Here $\Delta G^{\dagger}$ is the height of the free energy barrier and $\beta = (k_{\text{B}}T)^{-1}$. Using $\Delta G^{\dagger} \sim 3.5$kT from the
PBMetaD simulations and the fact that the height of the barrier is approximately the same on both
sides of the transition, we obtain $\tau_{\text{MFP}}^{\textbf{4}\rightarrow\textbf{5}} = \tau_{\text{MFP}}^{\textbf{5}\rightarrow\textbf{4}} = 4 \pm 1$ μs.

The analyses above enable us to draw some additional conclusions about the dynamics of the
inhibitory module in the inactive state of the complex. Our simulations show that the Ac domain
is in a fast equilibrium between a tightly bound state (state **5**) and one were the Ac-helix is more
relaxed (state **4**) but still inhibits the function of the DH domain by burying residue Y174. This
is coherent with the observation that the the Ac-domain binds to the DH one mostly through hy-
drophobic interactions and establishes a hydrogen bond only between Y174 and R332. The exis-
tence of this fast equilibrium is compatible with experimental results, which suggest the dynamics
of the inhibitory module to occur at multiple timescales (*Li et al., 2008*). Ultimately, the fact that the
state around that CV1~0.95 approximates well a transition state also indicates that CV1 is a good
guess for the reaction coordinate that drives the transition from the bound state to the molten
globule-like state.

The binding equilibrium
The free energy landscape (Fig. 2A) provides an overall picture of the binding dynamics of the in-
hibitory module to the DH domain, which is further corroborated by the unbiased MD simulations
(Fig. 4). Previously, the binding of the Ac helix to the DH domain has been studied using two types
of NMR experiments (*Li et al., 2008*), and it would be relevant to compare our simulations with
these.

First, relaxation dispersion measurements of a number of methyl groups revealed the presence
of microsecond–millisecond dynamics at a number of sites in the Vav1 AD complex (*Li et al., 2008*).
Residues experiencing such dynamics were located both in the Ac helix and in the Ac helix binding
site, but also throughout the rest of the protein. Dispersion measurements on the phosphory-
lated protein (pY174) showed quenched dynamics in those residues in the Ac helix that displayed
exchange dynamics in the unphosphorylated protein. However, extensive dynamics dynamics re-
mained in many other residues. Thus, the dispersion experiments suggest multiple dynamical
processes (with at least four states) on the microsecond–millisecond time scales, with only some
of these related directly to the binding of the Ac helix to the DH domain. This complicated exchange
dynamics prevented quantitative fitting and modelling of the relaxation dispersion NMR data (*Li*
et al., 2008).

In a second set of experiments, the authors studied a series of mutant variants of Vav1 that shift the equilibrium between the bound and unbound Ac helix. The NMR measurements show that the binding-unbinding reaction is fast on the NMR timescale and whre used to estimate the populations of the unbound state, which for the wild-type protein was estimated to be 9% corresponding to a $\Delta G_{\text{exp}} = -2.3$ k_BT. The suggestion that this equilibrium corresponds to opening of the protein and exposure of Y174 to solvent was corroborated by a high correlation between the estimated population of the unbound state for the mutants and the rate of phosphorylation by the Lck kinase.

Together, the NMR experiments thus suggests a complex dynamical landscape between multiple states, making it difficult to compare our simulations directly to experiments. In particular, the interpretation of both sets of NMR experiments are dependent on both the rates of conformational exchange as well as the differences in chemical shifts in the different substates. Since both experiments and simulations revealed that the Vav1 dynamics is not a simple two-state system it remains somewhat unclear which structures should be considered part of the bound state and which would be unbound.

We thus proceeded by estimating the population of bound and unbound structures by integrating the populations from the free energy surface along CV1 and CV2 using different definitions of the bound state (from CV1> 0.5 to CV1> 0.95) and defining the unbound state as those that were not bound (i.e. as a two-state system; *Lindorff-Larsen and Teilum* (*2021*)). The resulting free energy differences range from $\Delta G_{\text{sim}} = 0.6 \pm 0.3$ to 3 ± 1 k_BT (where the uncertainties have been determined via block error analysis) depending on the definition of the bound state (Fig. S8).

These results are in apparent disagreement with the experimentally determined free energy difference between the two states ($\Delta G_{\text{exp}} = -2.3$ k_BT) (*Li et al., 2008*). Possible reasons behind this inconsistency include (i) the possible lack of convergence in the estimation of ΔG_{sim}, which may in part be due to difficulties in overcoming the entropic barrier separating the bound from the unbound states, and (ii) possible imperfections in the force field that we used. These points are discussed below.

Concerning the issue of convergence, we performed a block error analysis (*Flyvbjerg and Petersen, 1989*) to estimate the error of ΔG_{sim} (Fig. S9). The monotonously increasing behavior of the the calculated errors for almost every definition of the unbound state suggests that the values are not fully converged. While the difference between experimental and calculated values of ΔG are greater than the error, the block analysis suggests that the error estimates themselves might be uncertain. This implies that the discrepancy could, at least partially, be explained by the difficulty in converging this quantity.

One of the reasons that the system is difficult to sample is that binding events may occur through 'entropic funnels' that are usually not simply discernible by projecting the dynamics on a small number of collective variables. In the context of the Ac helic binding to the DH domain, there appears to be an entropic barrier separating the unbound state **2** from the partially bound state **3** (Fig. 2). Although such a barrier is not clearly visible in the free energy landscape projected onto CV1 and CV2, a strong indication of the difficulty in observing this transition comes from the fact that the multiple shorter MD simulations are able to interconnect all the metastable states (Fig. 2) except for **2** and **3** (Fig. 4C,D). This makes it particularly hard to observe multiple, independent binding-unbinding events, and even enhanced sampling methods are difficult to apply because of the inherent degeneracy of the unbound state. While methods such as by Funnel Metadynamics (*Limongelli et al., 2013*) can be used to enhance sampling of binding events, they have mostly been employed to relatively rigid ligands.

Finally, lack of agreement between experiment and simulation can be due to inaccuracies in the force field. In our simulations we used the Amber a99SB-*disp* force field (*Robustelli et al., 2018*) as one of the only force fields that has been extensively tested and found to describe both folded and disordered proteins relatively well. This ability is crucial to study systems such as Vav1 that contain both disordered and ordered regions (*Robustelli et al., 2020*). A key aspect of constructing

337 a force field with these properties was to obtain a better balance between protein-protein and
338 protein-water interactions (*Piana et al., 2015*; *Robustelli et al., 2018*). Nevertheless, it is possible
339 that remaining imbalances in both solvent and other interactions can explain the difference in
340 affinity for the Ac helix. Indeed, a recent analysis has shown that a99SB-*disp* underestimates the
341 stability of several complexes between folded proteins (*Piana et al., 2020*).

Conclusions

343 Comparisons of phosphoproteomics data with structural information has revealed the intriguing
344 observation that a number of phosphorylations occur at positions that are buried inside the protein
345 structure in the unphosphorylated state (*Jiménez et al., 2007*). One hypothesis for how these sites
346 may become accessible to kinases is that they exist in a dynamic equilibrium between buried and
347 exposed states, and such dynamics have been shown for example for both Vav1 (*Li et al., 2008*)
348 and p27 in complex with its targets (*Tsytlonok et al., 2019*; *Henriques and Lindorff-Larsen, 2020*).
349 From a functional perspective, these proteins are particularly interesting because both states are
350 essential for function: the major state is an inhibited stated, and the minor state is necessary for
351 the ability of the kinase to relieve inhibition. One would thus expect that the balance between the
352 two states is finely tuned during evolution. From a biophysical perspective, such systems are good
353 model systems to study folding upon binding of intrinsically disordered proteins.
354 We have investigated the binding dynamics of the Ac domain to the DH domain and used both
355 standard MD and metadynamics simulations to probe the transient exposure of Y174 in Vav1. De-
356 spite the disagreement with NMR experiments in terms of the ratio between the active and the
357 inhibited states populations, we have found a complex binding pathway that proceeds by folding
358 the Ac-helix while unbound and accommodating it in the hydrophobic pocket formed by its bind-
359 ing site. This leads to the conclusion that the binding mechanism involves association between a
360 partially helical structure and the DH domain, which then proceeds via a molten-globule like inter-
361 mediate and a major transition before reaching the final bound state.

Methods

Protein structures

364 Two structures are available for the AD complex: one solved by NMR (PDB: 1F5X, *Aghazadeh et al.*
365 (*2000*)) and one by crystallography (PDB: 3KY9, *Yu et al.* (*2010*)). While the NMR structure corre-
366 sponds to the 168-365 construct, similar to the one used to study dynamics in Vav1 (*Li et al., 2008*),
367 the crystal structure is composed by 5 domains (in a dimeric form) and lacks coordinates for the
368 181-188 loop. In order to prepare the crystal structure for MD simulations, we isolated the 169-365
369 fragment of the first monomer and modelled the missing loop using Modeller (*Eswar et al., 2006*).
370 In particular, we generated 50 different models and selected the one with the best Modeller score.
371 In the case of the NMR structure, instead, we removed residue A168 to match the sequence used
372 in the NMR experiments (*Li et al., 2008*).

Molecular dynamics

374 We used GROMACS v.2018.6 (*Abraham et al., 2015*) for our MD simulations. The protein structures
375 were solvated in a dodecahedral box of water molecules with periodic boundary conditions. We
376 set the dimension of the box in a such a way that the protein was 1 nm away from any side of
377 the solvent box ($d = 8.3$ nm for the native, autoinhibited state) resulting in approximately 12,000
378 water molecules. Chloride and sodium ions were included to neutralize the protein and to set the
379 salt concentration to 50 mM. In most of the simulations we employed the a99SB-*disp* force field
380 (*Robustelli et al., 2018*) with its corresponding TIP4PD-like water model, while we conducted some
381 other control simulations using CHARMM36m force field (*Huang et al., 2017*) with the CHARMM-
382 modified TIP3P water model.

The Van der Waals interactions were set to zero at 1.0 and 1.2 nm cutoff, for AMBER and
CHARMM respectively, and the long-range electrostatics was taken into account by means of the
particle mesh Ewald algorithm with a 0.16 nm mesh spacing and cubic interpolation. The bonds
involving hydrogen atoms were constrained using the LINCS algorithm. Equations of motion were
integrated using the leapfrog algorithm with a 2 fs time step. We employed the Bussi thermostat
(*Bussi et al., 2007*) to control the temperature, with $\tau_T = 0.1$ ps^{-1} coupling time constant. Before
production, we used steepest descent to minimize the energy, and then equilibrated the box first
for 5 ns in the NVT ensemble at 300K and then in the NPT ensemble, employing in both cases po-
sition restraints of 1000 kJ mol^{-1} nm^{-2} on all interatomic bonds. We fixed the pressure to 1 atm in
the NPT ensemble through the Parrinello-Rahman barostat (*Parrinello and Rahman, 1981*), setting
$\tau_P = 2$ ps^{-1} as coupling time constant and an isothermal compressibility of 4.5×10^{-5} bar^{-1}.

Well-tempered and parallel bias metadynamics

We used the PLUMED v.2.5 plugin (*Tribello et al., 2014*) together with Gromacs to perform both well-
tempered and multiple-walker parallel-bias metadynamics (respectively, WTMetaD and PBMetaD)
simulations (*Barducci et al., 2008*; *Pfaendtner and Bonomi, 2015*). In these simulations we used
2–5 collective variables (CVs), described below and in the supplementary material. Gaussians were
deposited every 1 ps, with an initial height of 1 kJ/mol, and a bias factor $\gamma^{-1} = 10$ for WTMetaD and
$\gamma^{-1} = 44$ for PBMetaD. In the case of PBMetaD, we employed two walkers that were simultaneously
depositing Gaussians on the five one-dimensional CVs. Input files for PLUMED can be found in
PLUMED-NEST (*Bonomi et al., 2019*) at https://www.plumed-nest.org/eggs/21/017/ .

The initial configurations for metadynamics simulation were prepared as described above apart
from the box size. To avoid self-interaction across the periodic boundaries in the unbound, active
state (which is substantially more extended than the inhibited state), we adopted the following
strategy. We first performed a (non well-tempered) metadynamics simulation with Gaussian hills
heights of 3 kJ/mol in order to induce a rapid transition to an extended state of the Ac-helix. We let
the simulation run for 50 ns and selected the frame corresponding to the most extended config-
uration, which was then re-solvated. This resulted in a substantially larger box ($d = 11.4$ nm) with
respect to the one used for standard MD simulations described above (34,000 water molecules
and a total of 130,000 particles). We then solvated the bound (crystal) structure in a box of this size
and used this as starting point for the WTMetaD. This structure was also used as starting point for
one of the replicas in the PBMetaD simulations, and the other replica was started from an unbound
structure extracted from the WTMetaD simulation.

Analysis

All analyses were conducted using GROMACS, PLUMED, VMD (*Humphrey et al., 1996*), MDTraj
(*McGibbon et al., 2015*) and in-house developed Python tools using Numpy (*Harris et al., 2020*)
and Scipy (*Virtanen et al., 2020*); see https://github.com/KULL-Centre/papers/tree/master/2021/
Vav1-Orioli-et-al for the simulations, data, supplementary videos and the scripts employed to gen-
erate the paper figures and https://github.com/fpesceKU/BLOCKING for the code employed to
perform block error analysis). We obtained free energy surfaces for WTMetaD by employing the
sum_hills tool in PLUMED, while we reweighted PBMetaD simulations following the prescription in
Pfaendtner and Bonomi (*2015*) after discarding the first 1 μs as equilibration (20% of the simulation
length) from both the walkers. All plots were made using Matplotlib (*Hunter, 2007*), while all the
renderings and supplementary videos were obtained using VMD and the Tachyon software (*Stone
et al., 2013*).

Acknowledgments

428 We acknowledge work by Fabio Doro who contributed to the early stages and simulations in this
429 project, and to Micha B. A. Kunze and Elena Papaleo for fruitful discussions. We are grateful to

Francesco Pesce for inspiring discussions and for providing his scripts to perform block error analysis. We acknowledge support from the NordForsk Nordic Neutron Science Programme (grant number 81912) and the Lundbeck Foundation BRAINSTRUC structural biology initiative (R155-2015-2666). We also acknowledge access to computational resources from the ROBUST Resource for Biomolecular Simulations (supported by the Novo Nordisk Foundation grant no. NF18OC0032608) and the Danish National Supercomputer for Life Sciences (Computerome).

References

Abe K, Whitehead IP, O'Bryan JP, Der CJ. Involvement of NH2-terminal sequences in the negative regulation of Vav signaling and transforming activity. Journal of Biological Chemistry. 1999; 274(43):30410–30418.

Abraham MJ, Murtola T, Schulz R, Páll S, Smith JC, Hess B, Lindahl E. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX. 2015; 1:19–25.

Aghazadeh B, Lowry WE, Huang XY, Rosen MK. Structural basis for relief of autoinhibition of the Dbl homology domain of proto-oncogene Vav by tyrosine phosphorylation. Cell. 2000; 102(5):625–633.

Barducci A, Bussi G, Parrinello M. Well-tempered metadynamics: a smoothly converging and tunable free-energy method. Physical review letters. 2008; 100(2):020603.

Best RB, Hummer G, Eaton WA. Native contacts determine protein folding mechanisms in atomistic simulations. Proceedings of the National Academy of Sciences. 2013; 110(44):17874–17879.

Bonomi M, Bussi G, Camilloni C, Tribello GA, Banáš P, Barducci A, Bernetti M, Bolhuis PG, Bottaro S, Branduardi D, et al. Promoting transparency and reproducibility in enhanced molecular simulations. Nature methods. 2019; 16(8):670–673.

Bussi G, Donadio D, Parrinello M. Canonical sampling through velocity rescaling. The Journal of chemical physics. 2007; 126(1):014101.

Bussi G, Laio A. Using metadynamics to explore complex free-energy landscapes. Nature Reviews Physics. 2020; p. 1–13.

Bussi G, Tribello GA. Analyzing and biasing simulations with PLUMED. In: *Biomolecular Simulations* Springer; 2019.p. 529–578.

Bustelo XR. Vav proteins, adaptors and cell signaling. Oncogene. 2001; 20(44):6372–6381.

Chung HS, Louis JM, Eaton WA. Experimental determination of upper bound for transition path times in protein folding from single-molecule photon-by-photon trajectories. Proceedings of the National Academy of Sciences. 2009; 106(29):11837–11844.

Dombroski AJ, Walter WA, Gross CA. Amino-terminal amino acids modulate sigma-factor DNA-binding activity. Genes & development. 1993; 7(12a):2446–2455.

Du R, Pande VS, Grosberg AY, Tanaka T, Shakhnovich ES. On the transition coordinate for protein folding. The Journal of chemical physics. 1998; 108(1):334–350.

Dyson HJ, Wright PE. Coupling of folding and binding for unstructured proteins. Current opinion in structural biology. 2002; 12(1):54–60.

Eswar N, Webb B, Marti-Renom MA, Madhusudhan M, Eramian D, Shen My, Pieper U, Sali A. Comparative protein structure modeling using Modeller. Current protocols in bioinformatics. 2006; 15(1):5–6.

Flyvbjerg H, Petersen HG. Error estimates on averages of correlated data. The Journal of Chemical Physics. 1989; 91(1):461–466.

Fu CD, Pfaendtner J. Lifting the curse of dimensionality on enhanced sampling of reaction networks with parallel bias metadynamics. Journal of chemical theory and computation. 2018; 14(5):2516–2525.

Habchi J, Tompa P, Longhi S, Uversky VN. Introducing protein intrinsic disorder. Chemical reviews. 2014; 114(13):6561–6588.

Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, Wieser E, Taylor J, Berg S, Smith
NJ, Kern R, Picus M, Hoyer S, van Kerkwijk MH, Brett M, Haldane A, del R'ıo JF, Wiebe M, Peterson P, G'erard-
Marchant P, et al. Array programming with NumPy. Nature. 2020 Sep; 585(7825):357–362. https://doi.org/
10.1038/s41586-020-2649-2, doi: 10.1038/s41586-020-2649-2.

Heller GT, Aprile FA, Bonomi M, Camilloni C, De Simone A, Vendruscolo M. Sequence specificity in the
entropy-driven binding of a small molecule and a disordered peptide. Journal of molecular biology. 2017;
429(18):2772–2779.

Henriques J, Lindorff-Larsen K. Protein dynamics enables phosphorylation of buried residues in Cdk2/Cyclin
A-bound p27. bioRxiv. 2020; .

Huang J, Rauscher S, Nawrocki G, Ran T, Feig M, de Groot BL, Grubmüller H, MacKerell AD. CHARMM36m: an
improved force field for folded and intrinsically disordered proteins. Nature methods. 2017; 14(1):71–73.

Hubbard SR. Juxtamembrane autoinhibition in receptor tyrosine kinases. Nature Reviews Molecular Cell Biol-
ogy. 2004; 5(6):464–471.

Hummer G. From transition paths to transition states and rate coefficients. The Journal of chemical physics.
2004; 120(2):516–523.

Humphrey W, Dalke A, Schulten K, et al. VMD: visual molecular dynamics. Journal of molecular graphics. 1996;
14(1):33–38.

Hunter JD. Matplotlib: A 2D graphics environment. Computing in Science & Engineering. 2007; 9(3):90–95. doi:
10.1109/MCSE.2007.55.

Jiménez JL, Hegemann B, Hutchins JR, Peters JM, Durbin R. A systematic comparative and structural analysis
of protein phosphorylation sites based on the mtcPTM database. Genome biology. 2007; 8(5):1–20.

Kim AS, Kakalis LT, Abdul-Manan N, Liu GA, Rosen MK. Autoinhibition and activation mechanisms of the
Wiskott–Aldrich syndrome protein. Nature. 2000; 404(6774):151–158.

Kim JY, Meng F, Yoo J, Chung HS. Diffusion-limited association of disordered protein by non-native electrostatic
interactions. Nature communications. 2018; 9(1):4707.

Ko LJ, Prives C. p53: puzzle and paradigm. Genes & development. 1996; 10(9):1054–1072.

Lee GM, Donaldson LW, Pufall MA, Kang HS, Pot I, Graves BJ, McIntosh LP. The structural and dynamic basis of
Ets-1 DNA binding autoinhibition. Journal of Biological Chemistry. 2005; 280(8):7088–7099.

Li P, Martins IR, Amarasinghe GK, Rosen MK. Internal dynamics control activation and activity of the autoinhib-
ited Vav DH domain. Nature structural & molecular biology. 2008; 15(6):613–618.

Limongelli V, Bonomi M, Parrinello M. Funnel metadynamics as accurate binding free-energy method. Pro-
ceedings of the National Academy of Sciences. 2013; 110(16):6358–6363.

Lindorff-Larsen K, Teilum K. Linking thermodynamics and measurements of protein stability. Protein Eng Des
Sel. 2021 Feb; 34.

Malagrinò F, Visconti L, Pagano L, Toto A, Troilo F, Gianni S. Understanding the Binding Induced Folding of
Intrinsically Disordered Proteins by Protein Engineering: Caveats and Pitfalls. International Journal of Molec-
ular Sciences. 2020; 21(10):3484.

McGibbon RT, Beauchamp KA, Harrigan MP, Klein C, Swails JM, Hernández CX, Schwantes CR, Wang LP, Lane TJ,
Pande VS. MDTraj: A Modern Open Library for the Analysis of Molecular Dynamics Trajectories. Biophysical
Journal. 2015; 109(8):1528 – 1532. doi: 10.1016/j.bpj.2015.08.015.

Papaleo E, Sutto L, Gervasio FL, Lindorff-Larsen K. Conformational changes and free energies in a proline
isomerase. Journal of Chemical Theory and Computation. 2014; 10(9):4169–4174.

Parrinello M, Rahman A. Polymorphic transitions in single crystals: A new molecular dynamics method. Journal
of Applied physics. 1981; 52(12):7182–7190.

Pearson MA, Reczek D, Bretscher A, Karplus PA. Structure of the ERM protein moesin reveals the FERM domain
fold masked by an extended actin binding tail domain. Cell. 2000; 101(3):259–270.

520 **Pfaendtner J**, Bonomi M. Efficient sampling of high-dimensional free-energy landscapes with parallel bias
521 metadynamics. Journal of chemical theory and computation. 2015; 11(11):5062–5067.

522 **Piana S**, Donchev AG, Robustelli P, Shaw DE. Water dispersion interactions strongly influence simulated struc-
523 tural properties of disordered protein states. The journal of physical chemistry B. 2015; 119(16):5113–5123.

524 **Piana S**, Lindorff-Larsen K, Shaw DE. Atomistic description of the folding of a dimeric protein. The Journal of
525 Physical Chemistry B. 2013; 117(42):12935–12942.

526 **Piana S**, Robustelli P, Tan D, Chen S, Shaw DE. Development of a force field for the simulation of single-chain
527 proteins and protein–protein complexes. Journal of chemical theory and computation. 2020; 16(4):2494–
528 2507.

529 **Pietrucci F**, Laio A. A collective variable for the efficient exploration of protein beta-sheet structures: application
530 to SH3 and GB1. Journal of Chemical Theory and Computation. 2009; 5(9):2197–2201.

531 **Prakash A**, Sprenger K, Pfaendtner J. Essential slow degrees of freedom in protein-surface simulations: A
532 metadynamics investigation. Biochemical and biophysical research communications. 2018; 498(2):274–281.

533 **Pufall MA**, Graves BJ. Autoinhibitory domains: modular effectors of cellular regulation. Annual review of cell
534 and developmental biology. 2002; 18(1):421–462.

535 **Pufall MA**, Lee GM, Nelson ML, Kang HS, Velyvis A, Kay LE, McIntosh LP, Graves BJ. Variable control of Ets-1
536 DNA binding by multiple phosphates in an unstructured region. science. 2005; 309(5731):142–145.

537 **Robustelli P**, Piana S, Shaw DE. Developing a molecular dynamics force field for both folded and disordered
538 protein states. Proceedings of the National Academy of Sciences. 2018; 115(21):E4758–E4766.

539 **Robustelli P**, Piana S, Shaw DE. The mechanism of coupled folding-upon-binding of an intrinsically disordered
540 protein. Journal of the American Chemical Society. 2020; .

541 **Schuler B**, Borgia A, Borgia MB, Heidarsson PO, Holmstrom ED, Nettels D, Sottini A. Binding without folding–the
542 biomolecular function of disordered polyelectrolyte complexes. Current opinion in structural biology. 2020;
543 60:66–76.

544 **Sgourakis NG**, Patel MM, Garcia AE, Makhatadze GI, McCallum SA. Conformational dynamics and structural
545 plasticity play critical roles in the ubiquitin recognition of a UIM domain. Journal of molecular biology. 2010;
546 396(4):1128–1144.

547 **Stone JE**, Vandivort KL, Schulten K. GPU-accelerated molecular visualization on petascale supercomputing
548 platforms. In: *Proceedings of the 8th International Workshop on Ultrascale Visualization*; 2013. p. 1–8.

549 **Stöven S**, Ando I, Kadalayil L, Engström Y, Hultmark D. Activation of the Drosophila NF-kappaB factor Relish by
550 rapid endoproteolytic cleavage. EMBO Rep. 2000; 1(4):347–352.

551 **Sturzenegger F**, Zosel F, Holmstrom ED, Buholzer KJ, Makarov DE, Nettels D, Schuler B. Transition path times of
552 coupled folding and binding reveal the formation of an encounter complex. Nature communications. 2018;
553 9(1):4708.

554 **Tribello GA**, Bonomi M, Branduardi D, Camilloni C, Bussi G. PLUMED 2: New feathers for an old bird. Computer
555 Physics Communications. 2014; 185(2):604–613.

556 **Trudeau T**, Nassar R, Cumberworth A, Wong ET, Woollard G, Gsponer J. Structure and intrinsic disorder in
557 protein autoinhibition. Structure. 2013; 21(3):332–341.

558 **Tsytlonok M**, Sanabria H, Wang Y, Felekyan S, Hemmen K, Phillips AH, Yun MK, Waddell MB, Park CG,
559 Vaithiyalingam S, et al. Dynamic anticipation by Cdk2/Cyclin A-bound p27 mediates signal integration in
560 cell cycle regulation. Nature communications. 2019; 10(1):1–13.

561 **Tybulewicz VL**. Vav-family proteins in T-cell signalling. Current opinion in immunology. 2005; 17(3):267–274.

562 **Van Der Lee R**, Buljan M, Lang B, Weatheritt RJ, Daughdrill GW, Dunker AK, Fuxreiter M, Gough J, Gsponer
563 J, Jones DT, et al. Classification of intrinsically disordered regions and proteins. Chemical reviews. 2014;
564 114(13):6589–6631.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P, Weckesser
W, Bright J, van der Walt SJ, Brett M, Wilson J, Millman KJ, Mayorov N, Nelson ARJ, Jones E, Kern R, Larson E,
Carey CJ, et al. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. Nature Methods. 2020;
17:261–272. doi: 10.1038/s41592-019-0686-2.

Wang Y, Papaleo E, Lindorff-Larsen K. Mapping transiently formed and sparsely populated conformations on
a complex energy landscape. Elife. 2016; 5:e17505.

Yu B, Martins IR, Li P, Amarasinghe GK, Umetani J, Fernandez-Zapico ME, Billadeau DD, Machius M, Tomchick
DR, Rosen MK. Structural and energetic mechanisms of cooperative autoinhibition and activation of Vav1.
Cell. 2010; 140(2):246–256.

Supporting Information

Collective Variables

We used one or more of five different CVs in this manuscript.

The first, CV1, quantifies the extent to which the Ac-helix is bound in its native position
to the DH-domain. Specifically, we define CV1 via a continuous function quantifying the
fraction of a set of native contacts that are formed (***Best et al., 2013***):

$$581 \quad \text{CV1}(X) = \sum_{i \in S_1} \sum_{j \in S_2} \frac{1}{1 + \exp\left[\beta\left(r_{ij}(X) - \lambda r_{ij}(X_0)\right)\right]} \tag{2}$$

Here, X represents the instantaneous configuration of the protein, S_1 and S_2 are two sets
of atoms, $r_{ij}(X)$ is the distance between atoms i and j in configuration X, r_{ij}^0 is the distance
between the same two atoms in a reference conformation X_0, $\beta = 5\text{Å}^{-1}$ and $\lambda = 2$. Set S_1
comprises of all the heavy atoms in the Ac-helix (residue 169 to 178), while set S_2 is defined
by all the heavy atoms j in a reference structure found closer than 6Å from any atom $i \in S_1$
such that the amino acids to which i and j belong are at least 3 residues apart in the protein's
sequence. In the the simulations started from the NMR structure, we used this structure as
the reference for defining the CVs, and similarly used crystal structure as reference confor-
mation for the simulations started from that structure. With this choice, some intra-helix
contacts are included in the definition of CV1 and this is useful to distinguish between un-
bound states of the helix with folded helix (CV1$\sim$ 0.25) and unbound states with unfolded
594
595 helix (CV1$\sim$ 0).

596 The second collective variable CV2, instead, measures the amount of helical structure in
the 169-178 portion of the Ac domain. This is achieved by measuring the cosine distance
597 between the instantaneous value of the dihedral angles in the Ac-helix with those of a ref-
598 erence structure X_0:

$$600 \quad \text{CV2}(X) = \frac{1}{2} \sum_{i=1}^{N} \left[1 + \cos(\phi(X) - \phi(X_0))\right] \tag{3}$$

The reference structure for CV2 is the same as for CV1.

The third collective variable, CV3, is defined identically to CV1 but with different choices
for the atomic sets S_1 and S_2. In particular, S_1 contains the heavy atoms in the side chain for
Y174 and the corresponding C_α, while S_2 is defined by all the heavy atoms in the sidechains
of T205, Y209, V328, Q331 and R332. Also in the case of S_2, the alpha-carbons of each
residue have been included as well. In this way, CV3 can be interpreted as a measure of the
binding of Y174 to its binding site, irregardless of the conformation of the Ac-helix. Similarly,
CV4 is defined as the fraction of contacts between the alpha-carbon and sidechain heavy
atoms in I173, Y174 and L177 and those in Y209, T212, P320, L325 and V328. Despite its
similarity with CV3, CV4 introduces contacts relevant for the appropriate positioning of the
Ac-helix, especially at the beginning of the binding process.

Finally, the last collective variable CV5 is based on ***Pietrucci and Laio*** (***2009***) who intro-
613 duced a different way to estimate the helical content. In particular, we used

$$615 \quad \text{CV5}(X) = \sum_{i=1}^{4} \frac{1 - \left(\frac{r_i}{r_0}\right)^6}{1 - \left(\frac{r_i}{r_0}\right)^{12}} \tag{4}$$

618 where r_i measure the distances from an alpha-helical configuration (for residues 169–178),
619 $r_0 = 0.1$ nm and the sum runs over the four 6-residue long stretches of the helix.

Differences between simulations started from the NMR and crystal structures

The two protein structures, respectively resolved through NMR and crystallography and here labelled as *NMR* and *Crystal*, show clear differences in particular in two regions: (i) In the NMR structure the docking pose of the Ac-helix is almost perfectly horizontal with respect to the DH domain, while in the case of the crystal the acidic domain is slightly tilted, with the N-terminus pointing downwards; (ii) Residues 191–196 show a kink that splits the corresponding helix in two parts, while such a kink is absent in the crystal structure. These structural differences also affect the side chain packing of Y174, which forms a seemingly stable hydrogen bond with R332 in the crystal structure. The same interaction is not present in the NMR structure. We performed 1 μs-long simulations starting from either the crystal or the NMR structure and using two different force fields (a99sb-*disp* and CHARMM36m). In Fig. S1 we show the behavior of the fraction of native contacts with respect to the first simulation frame (Q), RMSD to initial frame (RMSD), CV1, CV2 and CV3 on the different MD simulations. Examining all of these metrics in the simulations obtained with both the force fields, we find that the NMR structure is less stable than the crystal structure on the μs timescale. Indeed, the NMR structure looses approximately the 20% of its overall contacts within 500 ns of simulation and the binding of the Ac-helix in the simulation started from the NMR structure appears to be less stable than the one started from the crystal configuration ($\langle CV1_{CRY} \rangle = 0.96 \pm 0.02$ and $\langle CV1_{NMR} \rangle = 0.83 \pm 0.02$ for AMBER and $\langle CV1_{CRY} \rangle = 0.95 \pm 0.02$ and $\langle CV1_{NMR} \rangle = 0.71 \pm 0.02$ for CHARMM). This suggests that a substantial rearrangement of the interface between the DH and the Ac domains occurs in a short timescale in the case of the simulation started from the NMR structure, as also underlined by the significant change in the packing of Y174 during the simulation. Moreover, we stress the fact that the hydrogen bond between Y173 and R332 is transiently broken and reformed during both the a99SB-*disp* and CHARMM36m simulations started from the crystal structure (Fig. S1), while it is not formed in the runs started from the NMR structure. All these differences are evidently due to a convergence issue, as with infinite sampling the two structures should provide the same equilibrium description. However, the collective variables we employ for biasing need a reference template in order to be properly defined, and for the these reasons we decided to proceed with the crystal structure rather than the NMR structure.

Estimation of uncertainties in the calculation of the transition path time

In order to estimate the uncertainty on the calculation of the transition path time and on the committor value we adopted a leave-1-out approach. We split our set of simulations in 5 subsets of 20 simulations each, by removing each time the 5 trajectories started from one of the five selected starting points. Then we computed the value of the binding probability and the transition path time from each of the subsets and averaged the results. Finally, we estimated the uncertainty as the standard error on the sample mean.

Definition of states in the PBMetaD free energy landscape

We selected the 5 states along the possible Ac-DH binding pathway by visual inspection of the PBMetaD free energy landscape along CV1 and CV2. In particular, approximating the

states as rectangles and using the notation **state** $= (x_0, x_1, y_0, y_1)$, we have:

$$\mathbf{1} = (0.03, 0.12, 7.5, 10)$$
$$\mathbf{2} = (0.175, 0.225, 12.1, 14.2)$$
$$\mathbf{3} = (0.56, 0.66, 12.8, 14.3) \tag{5}$$
$$\mathbf{4} = (0.75, 0.83, 12.8, 15.55)$$
$$\mathbf{5} = (0.95, 1, 13.5, 16)$$

The conformations shown in Fig. 2 and 3 have been extracted from the five states among the ones having the highest metadynamics weights.

List of the simulations

Here we list all the unbiased MD simulations we conducted to qualitatively assess the shape of the free energy landscape along CV1 and CV2.

Starting point	# simulations	Length [μs]
CV1 = 0.07, CV2 = 9.81	3	0.5
CV1 = 0.10, CV2 = 8.52	2	0.5
CV1 = 0.18, CV2 = 13.47	3	0.5
CV1 = 0.20, CV2 = 12.04	4	0.5
CV1 = 0.34, CV2 = 15.38	5	0.5
CV1 = 0.43, CV2 = 14.22	5	0.3
CV1 = 0.44, CV2 = 13.63	1	0.5
CV1 = 0.50, CV2 = 14.48	6	0.3
CV1 = 0.53, CV2 = 13.74	2	0.5
CV1 = 0.53, CV2 = 13.74	2	0.5
CV1 = 0.60, CV2 = 13.20	1	0.5
CV1 = 0.61, CV2 = 14.08	2	0.5
CV1 = 0.77, CV2 = 13.53	3	0.5
CV1 = 0.77, CV2 = 15.19	1	0.5
CV1 = 0.79, CV2 = 13.53	3	0.5
CV1 = 0.81, CV2 = 14.16	1	0.5
CV1 = 0.81, CV2 = 15.32	1	0.5
CV1 = 0.82, CV2 = 12.02	1	0.5
CV1 = 0.97, CV2 = 14.20	4	0.3
CV1 = 0.97, CV2 = 15.70	1	0.5
CV1 = 0.98, CV2 = 15.34	1	0.5

Supporting Figure 1. Comparison between the 1 μs MD simulation of the NMR (red solid line) and crystal (blue solid line) Vav1 AD construct using two different force fields: a99SB-*disp* (left) and CHARMM36m (right).

Supporting Figure 2. Behaviour of the Y174-R332 hydrogen bond during the MD simulations using, respectively, the a99SB-*disp* and the CHARMM36m force fields started from the crystal configuration of the AD construct. The solid vertical black lines show the time points where the hydrogen bond is formed.

WTMetaD simulation

We performed WTMetaD simulation with a bias along CV1 and CV2 only. During the approximately 10 μs of simulation, we were able to sample multiple binding and unbinding events, both for the whole helix (CV1, black line in Fig. S3 (a)) and for Y174 (Fig. S3B). However, after the potential reached a stationary state, i.e. approximately around 5.5 μs (see the Gaussian heights in Fig. S3A) the Ac-helix was not able to rebind properly anymore until the very end of the simulation. This is usually a symptom of poor convergence, and indeed the free energy landscape obtained by summing the Gaussian hills (Fig. S3D) confirms this expectation. The free energy surface along CV1 and CV2 shows a sharp minimum around CV1~ 0.4 and CV2~ 10, which appears to be an artefact introduced by the fact that the protein mostly explored that region of CV space during the last 4 μs of simulation.

Supporting Figure 3. Time evolution of the (A) hills heights, (B) CV1 and CV2 and (C) CV3 in the WTMetaD simulation. (D) Final free energy surface along CV1 and CV2 obtained from the WTMetaD simulation.

The occurrence of such artefacts is often caused by one or more of these issues: (i) the choice of the bias factor γ is not optimal and, as a consequence, the bias potential reaches a stationary state before the exploration of the free energy space has been completed; (ii) the choice of the biasing collective variables is not optimal; (iii) the force field is not accurate enough and, when tested on complex conformational transitions and long simulations, it prefers configurations which are not biologically relevant. None of these issues can be easily solved, as both the correct CVs and the correct bias factor can be known precisely only if, respectively, the free energy barriers and the dynamical process under study are known a priori. To mitigate this problem, in the main text we have employed Parallel Bias Metadynamics with 5 collective variables and we have consequently increased the value of the bias factor from 10 to 44.

Further details on PBMetaD simulations

Supporting Figure 4. PBMetaD simulations of reversible binding of the Ac helix to the Vav1 DH domain. Dynamics of CV1 and CV2 the (A) first and (B) second walkers. (C) Dynamics of CV3 for both walkers. CV1 and CV2 describe, respectively, the fraction of native contacts between the Ac-helix and the DH domain and the helical content of the Ac helix, while CV3 measures the fraction of native contacts between Y174 and the surrounding residues within a 6Å distance. Values of CV2 have been divided by 16 to bring them onto the same scale as CV1.

Supporting Figure 5. Behaviour of the Y174-R332 hydrogen bond during the PBMetaD simulation. The solid black and red lines show, respectively, the time behaviour of CV1 and CV3 while the blue vertical lines indicate the time points where the hydrogen bond is formed in the simulation.

Supporting Figure 6. (A) Block error estimates for all CVs used in the PBMetaD simulation; (B)–(F) Time evolution of the Gaussian heights during the PBMetaD simulation for, respectively, CVs from 1 to 5.

Supporting Figure 7. Time evolution of CV4 and CV5 for both walkers in the PBMetaD simulations and the corresponding free energies after reweighting.

Supporting Figure 8. Block error analysis of the population of the unbound state. Probability density of CV1 in the PBMetaD simulation (black solid curve) and the different cutoffs used to determine the unbound state (dashed grey lines). For each cutoff, we report the value of (A) the ΔG_{sim} (red dot) and (B) the populations of the bound and unbound states (respectively, green and blue dots). Uncertainties were determined using block error analysis and, since the simulations were not fully converged, the reported values correspond to the error estimate using the two largest possible blocks (4 μs).

Supporting Figure 9. Error curves obtained by applying block error analysis to the (A) ΔG_{sim}, (B) bound state population and (C) unbound state population obtained from different definitions of the unbound state. The upper threshold in CV1 used to determine the unbound state is reported in the figure labels.