Mec1/Tel1-independent role of protein phosphatase 4 in Hop1 assembly to

promote meiotic chromosome axis formation in budding yeast

Ke Li[1,2], Miki Shinohara[1,3,4,*]

[1]Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan

[2]Department of Bioscience, Graduate School of Science, Osaka University, Toyonaka,

Osaka 560-0043, Japan

[3]Department of Advanced Bioscience, Graduate School of Agriculture, Kindai

University, Nara city, Nara 631-8505, Japan

[4]Agricultural Technology and Innovation Research Institute, Kindai University, Nara city,

Nara 631-8505, Japan

∗Correspondence:

Miki Shinohara

Department of Advanced Bioscience, Graduate School of Agriculture, Kindai University

3327-204 Nakamachi, Nara, Nara 631-8505, Japan

Phone: +81-742-43-6518

FAX: +81-742-43-8976

E-mail: mikis@nara.kindai.ac.jp

ORCID ID: 0000-0002-7413-8235

Abstract

Dynamic changes in chromosomal structure that occur during meiotic prophase play an important role in the progression of meiosis. Among them, meiosis-specific chromosomal axis-loop structures are important as a scaffold for integrated control between the meiotic recombination reaction and the associated checkpoint system to ensure accurate chromosome segregation. However, the molecular mechanism of the initial step of chromosome axis-loop construction is not well understood. Here, we showed that, in budding yeast, a Tel1/Mec1-related protein phosphatase 4 (PP4) is required to promote the assembly of a chromosomal axis components Hop1 and Red1 onto meiotic chromatin via interaction with Hop1. PP4 did not affect Rec8 assembly. Notably, unlike the previously known function of PP4, this novel function was independent of Tel1/Mec1 kinase functions. The defect in Hop1/Red1 assembly in the absence of PP4 function was not suppressed by Pch2 dysfunction. Since Pch2 is a conserved AAA+ ATPase and facilitates eviction of Hop1 protein from the chromosome axis, suggesting that PP4 is required for the initial step of chromatin loading of Hop1 rather than stabilizing Hop1 on the chromosome axis. These results indicate phosphorylation/dephosphorylation-mediated regulation of Hop1 recruitment onto chromatin during chromosome axis construction before meiotic double-strand break formation.

(194 words)

Introduction

Meiotic recombination is essential for proper segregation of homologous chromosomes

during the first meiotic division (meiosis-I) and is initiated by Spo11-dependent

programmed meiotic DNA double-strand break (DSB) formation at meiotic

recombination hot spots (KEENEY 2001). Meiotic DSB formation and subsequent

meiotic recombination require a specific scaffold, an axis-loop structure of chromatin

(SHINOHARA *et al.* 2000; BLAT *et al.* 2002; PANIZZA *et al.* 2011). A highly conserved

meiotic chromosome axis structure contains core factors Red1, Hop1, and meiosis-

specific cohesin complexes, including the meiosis-specific subunit Rec8 in budding

yeast (ROCKMILL AND ROEDER 1988; HOLLINGSWORTH AND BYERS 1989; KLEIN *et al.*

1999). After meiotic DSB formation, recombinases Rad51/Dmc1 are recruited to single-

stranded DSB ends (BISHOP *et al.* 1992; BISHOP 1994; SHINOHARA *et al.* 1997). The

single-stranded DSB ends promote the assembly of Zip-Msh-Mer/synapsis initiation

complex (ZMM/SIC) proteins, including the transverse element component Zip1, to

construct the synaptonemal complex (SC) and promote crossover (CO) formation (SYM

et al. 1993; NOVAK *et al.* 2001; BORNER *et al.* 2004; SHINOHARA *et al.* 2008; SHINOHARA

et al. 2015; DE MUYT *et al.* 2018). In the early leptotene, Zip1 with Zip3 localizes to the

centromere and mediates centromere pairing as an initial process to promote

chromosome pairing (TSUBOUCHI AND ROEDER 2005; FALK *et al.* 2010).

Hop1 is an axis-associated HORMA-domain protein with multiple roles during

meiotic recombination. Hop1 is required for efficient meiotic DSB formation, inter-

homolog bias in meiotic recombination, and meiotic prophase checkpoint activation

and inactivation (HOLLINGSWORTH *et al.* 1990; SCHWACHA AND KLECKNER 1997; NIU *et

al.* 2005; CARBALLO *et al.* 2008; PANIZZA *et al.* 2011; WEST *et al.* 2018). Hop1 together

with Red1 promotes DSB formation. Hop1 is phosphorylated by Tel1/Mec1 (yeast

ATM/ATR) kinases, which are sensor kinases of DSBs that activate the meiotic

prophase checkpoint signaling pathway via Mek1/Mre4 kinase in response to meiotic

recombination as well as synapsis defects (NIU *et al.* 2005; BORNER *et al.* 2008). Hop1,

Red1, and Mek1/Mre4 form the RHM complex involved in forming a stable axis

structure as a scaffold to integrate recombination and the required regulatory

responses associated with it. After meiotic DSB repair, Zip1-mediated SC formation

and a conserved AAA+ ATPase Pch2 promote Hop1 removal from the synapsed

chromosome axis and attenuate the checkpoint signal (BORNER *et al.* 2008). Recently,

Pch2 was demonstrated to be involved in the activation as well as the inactivation of

meiotic prophase checkpoint through Hop1 (RAINA AND VADER 2020). Thus, Hop1

phosphorylation is a key step to complete proper meiotic recombination.

Tel1/Mec1 is also required for the integrated meiotic recombination and

checkpoint activation. Mec1/Tel1 kinase is involved in intra-chromosomal regulation of

the distribution of meiotic DSB formation sites (ANDERSON *et al.* 2015; JOSHI *et al.*

2015; CHALLA *et al.* 2019) and signal transduction of the positive/negative feedback

loop of meiotic DSB formation through Rec114 phosphorylation (SASANUMA *et al.* 2007;

CARBALLO *et al.* 2008; ZHANG *et al.* 2011). In addition Mec1/Tel1 mediates DSB

interference as well as CO interference which is required for non-random distribution

(ZHANG *et al.* 2011; GARCIA *et al.* 2015; SHINOHARA *et al.* 2019).

Protein phosphatase 4 (PP4) is a conserved putative serine/threonine protein

phosphatase that functions specifically in response to DNA damage by attenuating the

phosphorylation signal by Mec1 and Tel1 during mitosis and meiosis (HOFFMANN *et al.*

1994; KEOGH *et al.* 2006; O'NEILL *et al.* 2007; FALK *et al.* 2010; LAI *et al.* 2011;

HUSTEDT *et al.* 2015). Mec1/Tel1 phosphorylates many target proteins including Rad53,

an ortholog of Chk2 kinase in budding yeast. In the mitotic DNA damage response

pathway, PP4 dephosphorylates these targets in the process of recovery from DNA

damage-induced signal transduction (O'NEILL *et al.* 2007). Several studies have

revealed the meiotic DSB-associated function of PP4 during meiotic recombination

(FALK *et al.* 2010; CHUANG *et al.* 2012). During meiosis, centromere pairing is regulated

by the balance of phosphorylation and dephosphorylation of Zip1 through Mec1/Tel1

kinase and PP4 phosphatase (FALK *et al.* 2010).

Here, we focused on the novel function of PP4 which is independent of

Mec1/Tel1 function in early prophase-I, especially in a proper meiotic chromosome axis

construction via physical interaction with Hop1 protein.

Results

PP4 activity is required for timely meiotic DSB formation and efficient CO

formation

Pph3 is a catalytic subunit of the yeast PP4 complex. Pph3 contains several

serine/threonine phosphatase motifs (HOFFMANN *et al.* 1994). We constructed a

catalytic mutation in the yeast *PPH3* gene by substitution of the conserved histidine

residue 112 required for phosphate binding with asparagine (SHI 2009) to generate the

pph3-H112N mutant (Figure 1A). The mitotic phenotypes of cells with this mutation

were analyzed. Pph3 is required for Rad53 dephosphorylation and mitotic DNA

damage repair (O'NEILL *et al.* 2007). The *pph3-H112N* mutation did not affect the

expression of another PP4 subunit Psy2, with conjugation of a 13-repeated Myc

epitope tag (Psy2-13Myc) (Figure S1A). In addition, we analyzed the assembly of

Psy2-13Myc on to meiotic chromatin in pachytene cells (Figure 1B). Consistent with a

previous study (FALK *et al.* 2010), we detected Psy2-13Myc signals as punctate foci on

meiotic chromatin. Importantly, the *pph3-H112N* mutation did not affect the Psy2-

13Myc localization. Pph3 and Psy2 form a very stable PP4 complex (O'NEILL *et al.*

2007), this suggesting that the catalytic activity of PP4 is not required for its meiotic

chromatin localization. Folk et al. reported the meiotic phenotypes of the *pph3*-null

(*pph3Δ*) mutant (FALK *et al.* 2010). Therefore, we characterized two *pph3* mutants,

pph3Δ and *pph3-H112N*, to understand the function of PP4 and its phosphatase

activity during meiosis in more detail. First, it was revealed that both *pph3* mutants

showed similarly delayed meiosis progression compared to the wild type (Figure S1B).

This result was consistent with a previous study on meiosis progression in the *pph3Δ*

and *psy2Δ* mutants (FALK *et al.* 2010).

To confirm whether the delayed meiosis progression in *pph3* mutants is

caused by meiotic recombination defects, we analyzed programmed meiotic DSB

formation as an initial event of meiotic recombination by Southern blotting at the *HIS4-*

LEU2 hot spot in chromosome III in the *pph3* mutants, although meiotic crossover

formation defects in *pph3Δ* have already been reported (FALK *et al.* 2010). The *HIS4-*

LEU2 hot spot is a well-analyzed artificial meiotic DSB hotspot that includes two known

sites, DSB I and II (Figure 1C). We detected and quantified DSB I and showed its

frequencies at each time point after meiosis entry (Figure 1D). Meiotic DSBs became

detectable 3 h after meiosis entry, and showed a peak at 4 h, and disappeared by 6 h

in the wild type (Figure 1D). On the other hand, both *pph3-H112N* and *pph3Δ* showed

an approximately 1 h delay in DSB formation compared to the wild type at the time

when the DSB reached half of the peak height (11%; Figure 1D, dashed line). In

addition, the *pph3* mutants showed an accumulation of meiotic DSBs with hyper-

resected ends until 6 h after meiosis entry, resulting in an approximately 3 h delay in

DSB disappearance (Figure 1D). These observations indicated the necessity of PP4

phosphatase activity for efficient DSB formation and turnover. PP4 is a Mec1/Tel1-

related protein phosphatase. Previous studies revealed that PP4 is required for

centromere pairing through Zip1 dephosphorylation for Mec1-mediated

phosphorylation at S75, or dephosphorylation of many other Mec1/Tel1 targets during

meiosis (KEOGH *et al.* 2006; FALK *et al.* 2010; HUSTEDT *et al.* 2015). Thus, we

constructed mitotic cyclin *CLB2* promoter-driven *MEC1* (p*CLB2-MEC1*) *tel1Δ* allele to

determine the effect of Mec1 and Tel1 on delayed DSB formation and/or turnover in the

pph3 mutants. p*CLB2* induces mitosis-specific gene expression. Thus, p*CLB2-MEC1*

causes meiosis-specific shutdown of *MEC1* transcription (LEE AND AMON 2003). The

p*CLB2-MEC1 tel1*Δ cells showed a reduced level of steady-state meiotic DSB

formation (7.7 ± 1.9% of total DNA) compared to that in the wild type (22.7 ± 3.1%).

p*CLB2-MEC1 tel1*Δ and p*CLB2-MEC1 tel1*Δ *pph3*Δ triple mutants showed more

severe reduction of DSB formation (5.3 ± 0.45% and 7.7 ± 3.2%, respectively; Figure

1D, graph). At later time points, in contrast to the *pph3* mutants, the p*CLB2-MEC1*

*tel1*Δ and p*CLB2-MEC1 tel1*Δ *pph3*Δ mutants showed the timely disappearance of

meiotic DSBs by 6 h in meiosis (Figure 1D). These findings suggested that the

accumulation of DSBs and the resulting delay in DSB repair, which was observed in

the *pph3* mutants at later time points (at 5–6 h) was dependent on Mec1/Tel1 (Figure

1E). In contrast, in the initiation of DSB formation, the p*CLB2-MEC1 tel1*Δ *pph3*Δ triple

mutant showed a peak at 4 h, whereas the p*CLB2-MEC1 tel1*Δ double mutant peaked

between 3 and 4 h after onset of meiosis entry (Figure 1E, graph). Since a delay in

meiotic DSB formation in the absence of Tel1/Mec1 activity was observed, the delayed

DSB formation caused by pph3 mutations appeared to be independent of Mec1/Tel1.

Next, we analyzed hetero-duplex (HD) formation using Southern blot to

analyze meiotic inter-homolog recombination products by using heteroallelic restriction

enzyme sites *Mlu*I and *Bam*HI located near the DSB I site and polymorphic *Xho*I sites

around the *HIS4-LEU2* hot spot (Figure 1C) (STORLAZZI *et al.* 1996). Corresponding to

the delayed meiotic DSB appearance and disappearance at the hot spot, the formation

of both HDs corresponding to intermediates for crossover (CO) and non-crossover

(NCO) were delayed by 3 h in the *pph3* mutants relative to that in the wild type (Figure

1E). In addition, an increase of NCO intermediates (HD1 and 4) at 8 h and a slight

reduction of CO intermediates (HD2 and 3) compared to wild type were observed not

only in the *pph3Δ*, as reported previously (FALK *et al.* 2010), but also in the *pph3-*

H112N mutant (Figure 1E).

After DSB end resection, meiotic DSBs are recognized by recombinases

Rad51 and Dmc1 to facilitate strand exchange for the formation of COs and NCOs.

The cytological analysis revealed that Rad51 appear transiently as punctate foci on

meiotic chromatin structure in the wild type (SHINOHARA *et al.* 2000). Thus, we next

examined the kinetics of Rad51 assembly and disassembly in the *pph3* mutants using

cytological analysis (Figure 1F). In the *pph3* mutants there was a 1 h delay in the

Rad51 assembly compared to the wild type. In addition, accumulation of Rad51 foci

was evident by 5 h after meiotic entry (Supplemental Figure 1C). On the other hand,

the number of Rad51 foci per nucleus at an early time point (3 h) in the *pph3Δ* and

pph3-H112N mutants was significantly lower than that in the wild type (64 ± 2.7, 56 ±

2.7, and 49 ± 3.3, respectively; Figure 1F). This also indicated the involvement of PP4,

as well as its phosphatase activity, for efficient DSB formation. In the disassembly

phase of Rad51 foci, a significant reduction in the number of Rad51 foci per nucleus

was observed in the wild type. However, continuous accumulation of Rad51 foci was

observed in the *pph3* mutants corresponding to inefficient meiotic DSB repair.

PP4 is involved in efficient Hop1 assembly onto chromatin in a Mec1/Tel1-

independent manner

Hop1 is a meiosis-specific phosphorylation target of Mec1 and Tel1, and PP4 is

involved in the dephosphorylation of these targets (CARBALLO *et al.* 2008; FALK *et al.*

2010). Furthermore, Hop1 is a HORMA-domain protein and a component of the axis

structure as a complex with Red1 to form the SC. Hop1 is essential for partner choice

of meiotic recombination (known as inter-homolog bias) and for efficient DSB formation

(HOLLINGSWORTH *et al.* 1990; NIU *et al.* 2005; PANIZZA *et al.* 2011; SUBRAMANIAN *et al.*

2016). Since it was observed (Figure 1C and D) that PP4 is required for efficient

initiation of meiotic DSB formation, we analyzed the kinetics of Hop1 assembly during

meiotic prophase-I (Figure 2A). In wild type Hop1, assembly began 2 h after the onset

of meiosis, corresponding to leptotene, and occurred prior to DSB formation.

We also analyzed the localization of Hop1 and Hop1 phosphorylated at

threonine 318 (Hop1-pT318) on meiotic chromatin. Hop1 phosphorylation is dependent

on Mec1/Tel1 kinase activation by Spo11-induced meiotic DSB formation (CARBALLO *et*

al. 2008). Hop1-pT318 has been observed as punctate foci at the end of short patches

of Hop1 signal in the wild type (CARBALLO *et al.* 2008) (Figure 2A and B）. In the wild

type, 50% of the cells displayed Hop1 foci 2 h after meiosis entry, but low levels of

Hop1-pT318 were observed at the same time. Hop1-pT318 foci began to appear after

3 h (Figure 2A and B). Hop1-pT318 focus formation requires single-stranded region at

meiotic DSB ends (IWASAKI *et al.* 2016). The time lag of approximately 2 h that was

evident between the Hop1 loading and the Hop1-pT318 appearance in the present

study indicated that Hop1 assembly occurs prior to the DSB formation and/or DSB-end

resection in wild type. In the *pph3* mutants, Hop1 assembly was rarely seen at 2 h

compared to the wild type. After 2 h, Hop1 staining gradually increased and reached

the wild-type level frequency (80%) at 5 h. Notably, pT318 staining appeared at almost

the same time (3 h) as the appearance of Hop1 foci (Figure 2A and B) and then

gradually increased, similar to Hop1. Hop1 loading seemed to be the rate-limiting factor

for Hop1 phosphorylation. These results showed that Pph3 and its activity can promote

the efficient loading of Hop1 on chromosomes. We confirmed that this was not due to

delayed expression of Hop1 in the mutant (see Figure 2C).

Next, we examined the effect of Mec1 and Tel1 on Hop1 assembly in the

pph3Δ mutant. While the p*CLB2-MEC1 tel1Δ* mutant cells did not show any defects in

the Hop1 assembly, the p*CLB2-MEC1 tel1Δ pph3Δ* mutant displayed delayed Hop1

assembly, similar to the *pph3* mutants (Figure 2A and B). In contrast, very few Hop1-

pT318 foci were observed in the p*CLB2-MEC1 tel1Δ* mutation background.

Next, we analyzed the phosphorylation status of target proteins of PP4 during

meiosis using western blot. PP4 is known to be involved in dephosphorylation of

phosphorylated S129 of histone H2A (γH2A) after DNA damage in vegetative growth

cells (KEOGH *et al.* 2006), and in Hop1-pT318 during meiosis (CHUANG *et al.* 2012).

Thus, we first confirmed that γH2A signal accumulated in the *pph3* mutants during

meiosis in a Mec1/Tel1 dependent manner, similar to mitotic cells, with markedly

reduced γH2A signals in the p*CLB2-MEC1 tel1Δ pph3Δ* mutant (Figure 2C). Moreover,

accumulation of Hop1-pT318 signal was also observed until 8 h in the *pph3Δ* mutant.

This phosphorylation was dependent on Mec1/Tel1 (Figure 2C). Although Hop1 protein

expression was detectable from 2 h after meiosis entry in both wild type and *pph3*

mutants, initiation of Hop1 phosphorylation at T318 was delayed by 1 h in the *pph3*

mutants compared to the wild type to reach maximum levels. Notably, despite the

timely expression of Hop1 protein in the *pph3* mutants (Figure 2D), the appearance of

Hop1 foci on meiotic nuclear spread was significantly delayed in the *pph3* mutants

(Figure 2A). This result suggested that PP4 might not be involved in Hop1 expression,

but rather acted to promotes Hop1 loading onto chromatin. Moreover, the expression

kinetics of Hop1 protein were also normal in the p*CLB2-MEC1 tel1Δ* or p*CLB2-MEC1*

tel1Δ pph3Δ mutants. In contrast to the γH2A phosphorylation, residual Hop1-pT318

signal was not detected in the p*CLB2-MEC1 tel1Δ* background, consistent with a

previous report showing that Hop1 phosphorylation largely depends on Mec1 and Tel1

activities (CARBALLO *et al.* 2008). Overall, the findings indicate a defect in Hop1

assembly on chromatin, even though a sufficient amount of Hop1 protein is expressed.

PP4 is involved in meiotic chromosome axes formation

Next, we examined the Red1 and Rec8 meiotic chromosome axis components

(ROCKMILL AND ROEDER 1988; KLEIN *et al.* 1999). Red1 forms a complex with Hop1

(NIU *et al.* 2005) and Rec8 is a component of the meiosis-specific sister chromatid

cohesin complex. Both proteins are required for the proper formation of axial elements

in meiosis and a highly-ordered specific chromosome structure, termed the

synaptonemal complex (SC) (KLEIN *et al.* 1999). Red1 and Rec8 are recruited to the

chromatin independent of each other (SUN *et al.* 2015). We examined Red1 and Rec8

staining of the meiotic chromatin (Figure 3A). In the wild type, Red1 and Rec8

colocalized with each other and formed linear structures along the 4′,6-diamidino-2-

phenylindole (DAPI) nuclear signal on meiotic chromatin 3 h after meiosis entry. In

contrast, in the *pph3* mutants, Red1 and Rec8 were observed as many fine signals

throughout the chromatin spreads. The number of Rec8 signals was clearly higher than

that of Red1. Time course experiments revealed the significant delay in the assembly

of Red1 protein in the *pph3* mutants, similar to the assembly of Hop1 shown in Figure

2A and 2B. In contrast, the timing of Rec8 assembly, despite of its unique structure,

was not affected in the presence or absence of Pph3 activity (Figure 3B). In addition, a

delay in the disassembly of both Red1 and Rec8 was observed in the *pph3* mutants.

Next, we analyzed Red1 and Rec8 protein expression during meiosis using

western blot (Figure 3C). Rec8 phosphorylation via Dbf4-dependent kinase (DDK) and

polo-like kinase 1 (PLK1) is associated with cleavage-independent cohesion release

from chromatin during late prophase-I (CHALLA *et al.* 2019). While the expression of

Rec8 occurred at almost the same time in the *pph3* mutants as that in the wild type, the

phosphorylation of Rec8 was delayed in the *pph3* mutants compared to that in the wild

type. Corresponding to the delayed phosphorylation of Rec8, Rec8 cleavage that

occurs during the meiotic divisions was also delayed in the *pph3* mutants. These

results were consistent with the delayed meiosis progression of meiosis in *pph3*

mutants (Figure S1). In addition, Red1 protein expression was quite normal in the *pph3*

mutants (Figure 3C). These results suggest that PP4 promotes Hop1-Red1 loading but

not Rec8-cohesin, which is required for the construction of a proper axis structure.

PP4 physically interacts with Hop1 protein

We next examined the physical interaction between the Hop1-Red1 and PP4 using a

co-immunoprecipitation assay (Figure 4A). Psy2-13Myc protein was precipitated using

anti-Myc antibody in meiotic cell extracts of *PSY2-13MYC* derivative *PPH3* (wild type)

cells. The Hop1 signal was detected in the precipitate from the cells at 2 to 6 h during

meiosis, indicating the interaction of Pys2 with Hop1. The Hop1 signal was also

detected in the precipitates of *pph3-H112N* mutant cells, indicating that the interaction

did not involve the phosphatase activity of PP4. However, the Hop1 signal was weaker

at 4 h than that in the *PPH3* background and was undetectable at 6 h in meiosis. This

might be caused by reduced amounts of the Psy2-13Myc protein signal in the

precipitates. However, the total amount of Psy2-13Myc protein in the *pph3-H112N*

mutant whole cell extracts of the *pph3-H112N* mutant from trichloroacetic acid-fixed

cells was indistinguishable from that in wild type-cells (Supplemental Figure 1A). A

mutation in *PPH3* affected the interaction between Hop1 and Psy2-13Myc, indicating

the interaction between Hop1 and the PP4 complex, rather than Psy2 alone (as an

independent complex).

Since Hop1-PP4 physical interaction was detected, we analyzed other axis

components Red1 and Rec8 proteins in the precipitates using western blot. After

immunoprecipitation in *PSY2-13MYC* cell extracts, as described above, the Hop1

protein signal was observed at 2 and 4 h after meiosis entry. However, the signal was

not evident at 6 h, even though Hop1 protein was expressed in the extract. In contrast,

Red1 and Rec8 were not observed in the precipitates, even though both proteins were

present in whole cell extracts (WCE). The findings indicates that PP4 can interacts with

Hop1 mainly, but not Red1 or Rec8. Given that Hop1 binds to Red1 as a complex (NIU

et al. 2005), there might be two distinct populations of Hop1, Hop1-PP4 and Hop1-

Red1.

Defective Hop1 assembly onto chromosome axes in the *pch2Δ* background in

the *pph3* mutant

Pch2 is a AAA+ ATPase that removes Hop1 protein from meiotic chromosome axes to

stabilize the transverse element Zip1 during formation of the SC structure (CHEN *et al.*

2014). Pch2 is also involved in a PP4-independent Hop1 dephosphorylation (CHEN *et*

al. 2014). Since we observed a defect in Hop1 assembly and delayed phosphorylation

of Hop1 in the *pph3* mutants, we analyzed the effects of the *pph3* mutations in the

Hop1 assembly on the meiotic chromosome axis in the *pch2Δ* background, which

features the over-loading of Hop1.

First, spore viability and distribution of viable spores were analyzed in the

pph3Δ pch2Δ double mutant by using tetrad analysis. Spore viability in the wild type,

pph3Δ, and *pch2Δ* was 97%, 88%, and 91%, respectively. Both *pph3Δ* and *pch2Δ* had

slightly reduced spore viability, but there was no significant bias in the distribution of

viable spores in asci, as reported previously (SAN-SEGUNDO AND ROEDER 1999; FALK *et*

al. 2010). In contrast, the pore viability of the *pph3Δ pch2Δ* double mutant was only

25%, with distributions of 0-, 2-, and 4-viable spores. This vias was indicative of the

non-disjunction of homologous chromosomes in the first division of meiosis.

In Hop1 and Hop1-pT318 assembly/disassembly kinetics, Hop1-pT318

assembly was delayed in the *pch2Δ* mutant, consistent with a delayed meiotic DSB

formation (FARMER *et al.* 2012). This delay was similar to that in the *pph3Δ* and the

pph3Δ pch2Δ double mutant (Figure 4D and E), indicating the absence of additive

defects to *pph3Δ* in Hop1 assembly/disassembly in the *pch2Δ* mutation. The findings

suggest that Pch2 and PP4 function in the same pathway in Hop1

assembly/disassembly.

Next, we examined the effect of the *pch2Δ* mutation on Hop1 localization by

using cytological analysis. As indicated previously (IWASAKI *et al.* 2016) and in Figure

2A, Hop1-pT318 foci were observed as punctate foci in the wild type and were affected

in the *pph3Δ* in early prophase-I (approximately 3 h in meiosis) (Figure 4E). In the

pch2Δ mutant, Hop1 aggregates colocalized with Hop1-pT318 signals. Similar

aggregates were also observed in the *pph3Δ pch2Δ* double mutant at an early time

point.

Notably, at the later time point (6 h during meiosis), Hop1-pT318 signals were

observed as a beaded chain on the fully elongated Hop1 signal on meiotic

chromosomes in the *pch2Δ* mutant cells (Figure 4E). In the *pph3Δ pch2Δ* double

mutant, some Hop1 signals were observed as foci of dots and some colocalized with

the Hop1-pT318 signal, similar to that in *pph3Δ*. These observations were evidence of

the upstream location of the *pph3Δ* mutation relative to the *pch2Δ* mutation in Hop1

localization process. This further suggests that PP4 may function in the promotion of

Hop1 assembly onto the meiotic chromatin to construct the chromosome axis before

Pch2 function in unloading of Hop1 from the chromosome axis. In contrast, in the

pph3Δ pch2Δ double mutant, some Hop1-pT318 signals were observed as beaded

chain on fully elongated Hop1. This indicates that the *pch2Δ* mutation stabilized Hop1

protein in the *pph3Δ* on some chromosome axis, but not on others. Therefore, there

could be other pathways that promote Hop1 assembly in the absence of PP4.

Multiple pathways to promote Hop1 assembly

On the meiotic chromosome axes, Hop1 is dynamic and is removed from

chromosomes through Pch2 activity. Finally, Hop1 is replaced by the transverse

element component Zip1 in the pachytene (BORNER *et al.* 2008). Given these facts, we

examined Zip1 localization in the *pph3Δ pch2Δ* double mutant. In wild type cells, as

previously reported, Zip1 was observed as dotty foci in the leptotene (2 h) and Zip1

begins to elongate after DSB-end resection in the zygotene, followed by the extension

of Zip1 throughout each chromosome in the pachytene (SYM *et al.* 1993; STORLAZZI *et*

al. 1996; SHINOHARA *et al.* 2015). Significant delays in both the appearance of dotty

Zip1 signals and the elongation of Zip1 were evident for *pph3Δ* (Figure 5A), as

reported previously (FALK *et al.* 2010). The *pch2Δ* mutant showed a delay in the Zip1

elongation and also dissociation from the chromosomes after full elongation. However,

the elongated Zip1 became the major population at 5 h after meiosis. In the *pph3Δ*

pch2Δ double mutant, after the delayed appearance of dotty Zip1 signals, similar to

pph3Δ (2 to 3 h), compromised elongation of Zip1 was observed. Complementary

localization of Zip1 and Hop1 on chromosomes from the zygotene to the pachytene

was observed in the wild type. A similar localization was observed in the *pph3Δ* mutant.

In contrast, *pch2Δ* mutant cells showed uniform colocalization of Zip1 and Hop1

(Figure 5B), as previously reported (BORNER *et al.* 2008). *Saccharomyces cerevisiae*

haploid cells have 16 chromosomes. Theoretically, 16 bivalents should be observed in

the pachytene stage: however, the number of linear Zip1 signals per cell was 12 ± 2.7

(mean ± 95% confidence interval) in the *pch2Δ*. The number was significantly reduced

in the *pph3Δ pch2Δ* double mutant in 68% of *pch2Δ* (8.2 ± 2.6) (Figure 5B and C).

While several elongated Zip1 signal colocalized with Hop1 in *pph3Δ pch2Δ*, Hop1-Zip1

free DAPI signal was also often observed in *pph3Δ pch2Δ* (Figure 5B). Next, we

compared the length of Zip1-Hop1 positive signals in *pch2Δ* with in *pph3Δ pch2Δ* to

determine whether there is chromosome size bias in the residual elongation of Zip1-

Hop1 observed in *pph3Δ pch2Δ*. No significant difference was evident ($P = 0.19$,

Figure 5D).

Since the *pph3Δ* mutation showed a chromosome-dependent effect in Hop1-

Zip1 elongation in the *pch2Δ* mutant background, we analyzed Rec8 and Red1

localization, as other axis components, to determine the status of the Hop1-Zip1 free

region observed in the *pph3Δ pch2Δ* double mutant. Unlike the Hop1 protein on the

axes, Red1 is not removed from the axis by Pch2. Rec8 is recruited to chromatin

independent of Hop1-Red1 (SUN *et al.* 2015). Furthermore, the Rec8 focus formation

was not affected in the *pph3* mutants (Figure 3A). In the *pch2Δ* mutant cell, Rec8 and

Red1 obviously colocalized on the chromosome axes (Figure 5E). On the other hand,

in the *pph3Δ pch2Δ* double mutant, Rec8 and Red1 were colocalized only on the

synapsed chromosome axes, and many of the Rec8 were observed as foci located in

the Red1 free region. This indicates the possibility that 68% of the chromosomes axes

with linearized Hop1 in the *pph3Δ pch2Δ* harbored a defect in the Pch2 sensitive step

following Hop1 loading, while the defect in other axes was prior to Hop1 assembly.

Delayed meiosis progression in the *pph3Δ* occurs independent of Pch2

Hop1 is an adaptor of meiosis-specific prophase checkpoint. Hop1

phosphorylation is affected in the *pch2Δ* mutant (LO *et al.* 2014) and Mec1/Tel1-

dependent phosphorylation of Hop1 affects Hop1 activity (CARBALLO *et al.* 2008). In the

wild type, Hop1 phosphorylation was detectable at 3 h during meiosis, continued until 5

h, and then disappeared (Figure 5F). The timing of Hop1-pT318 disappearance

matched the timing of the initiation of meiosis-I (Figure 5G). This was also observed

with *pph3Δ* mutant cells, in which the Hop1-pT318 signal disappeared at 8 h with the

simultaneous appearance of post-meiotic cells. In *pch2Δ* mutant cells, consistent with

previous findings, Hop1 protein was accumulated until 10 h and Hop1 phosphorylation

was compromised. On the other hand, in the *pph3Δ pch2Δ* double mutant, Hop1

protein accumulated until 10 h and Hop1-pT318 signal also accumulated until 8 h and

then disappeared. Notably, corresponding to the delayed disappearance of the Hop1-

pT318 signal, the *pph3Δ pch2Δ* double mutant showed a severe delay in meiosis

progression (Figure 5E and F). The *pch2Δ* mutation can suppress pachytene arrest in

the *zip1* mutant (SAN-SEGUNDO AND ROEDER 1999; BORNER *et al.* 2008). Since the

pch2Δ mutation cannot suppress the delayed progression of meiosis in the *pph3Δ*

mutant, this delay in the *pph3Δ* would be caused by a different mechanism from that in

the *zmm* mutant.

Discussion

During meiotic recombination, the chromosome axis is an essential scaffold. Some

components, including Mer2 and Rec8, are associated with pre-meiotic DNA

replication. On the other hand, the molecular mechanism underlying the formation of a

highly organized chromosomal axis structure, especially, the initial step of Hop1

recruitment onto meiotic chromatin, is not well understood. The results of the present

study support the suggestions that PP4 may function in Hop1 recruitment onto

chromatin in the initiation step of chromosome axis formation in early meiotic

prophase-I and that PP4 promotes Hop1 recruitment.

PP4 is involved in Hop1 assembly independent from Tel1/Mec1 kinase activity

During the formation of the meiotic SC structure, PP4 is required for centromere-

mediated chromosome pairing through Zip1 phosphorylation in a Mec1/Tel1-dependent

manner (FALK *et al.* 2010). We also observed reduced dotty Zip1 signal, which is

known to localize to the centromere (TSUBOUCHI AND ROEDER 2005). The reduction

was not suppressed even in the *pch2Δ* background (Figure 5A). Both *pph3* mutants,

pph3Δ and *pph3-H112N,* displayed significant delays in Hop1 assembly onto chromatin,

but this delay was Tel1/Mec1-independent (Figure 2A and B). Therefore, this delay in

Hop1 assembly would not be caused by compromised centromere pairing. In *pph3*

mutants, the appearance of γH2A, which is phosphorylated by Mec1/Tel1 kinase in

response to programmed meiotic DSBs, was observed at the same time as that in wild

type. The Hop1 phosphorylation (Hop1-pT318) by Tel1/Mec1 was delayed as well. This

result supports the idea that delayed Hop1 phosphorylation is caused by the delayed

assembly of Hop1 onto the axes. Hop1 is one of the rate-determining factors that

promotes meiotic DSB formation (PANIZZA *et al.* 2011). In addition, we revealed the

physical interaction between the Hop1 and PP4 components also in the *pph3-H112N*,

a catalytic mutation, and Hop1 protein in the precipitates was not always

phosphorylated (Figure 4A and B). This findings suggest that the Hop1-PP4 interaction

might not occur only for enzymatic activity, but may also regulate Hop1 assembly.

There is still another possibility that Hop1 localization on the chromosome

axes in the absence of PP4 is unstable and is rapidly removed from the axes. To

assess this, we examined Hop1 localization in the absence of Pch2. Proper Hop1

assembly did not occur in 32% of the chromosomes in each nucleus of *pph3Δ* cells.

This strongly suggests that PP4 is involved in the assembly phase of Hop1 and not in

the stabilization of Hop1 on the chromosome axes. Conversely, 68% of chromosomes

in the same nucleus displayed stable Hop1 assembly by *pch2Δ*, even in *pph3Δ*. These

observations indicate that there are multiple pathways other than the PP4 that promote

Hop1 assembly.

What is the molecular mechanism to promote Hop1 assembly?

The molecular mechanism by which PP4 promotes Hop1 assembly through physical

interactions is unknown. Hop1 is a HORMA domain protein. The dynamics of C-

terminal "closure-motif" may be important for the chromosome axis assembly and

disassembly through Red1 interaction. In addition, substitution of K593 of Hop1 with

alanine (*hop1-K593A*) results in severe defects in the Red1 interaction (WEST *et al.*

2018). We performed phospho-proteomics analysis in extracts from yeast cells

arrested in meiotic prophase-I. The finding revealed the phosphorylation of S595 of

Hop1 during meiosis, possibly by Mek1/Mre4 kinase (Shinohara M., unpublished

result). We previously analyzed Hop1 assembly onto chromatin in the

unphosphorylated mutation background of *hop1-S595L*. Normal Hop1 assembly was

observed in the *hop1-S595L* mutant.

In addition, a delay in Hop1 assembly in the *pph3Δ* mutant was also observed

in the p*CLB2-MEC1 tel1Δ* background, and Pph3 catalytic activity was required for the

timely assembly of Hop1 (Figure 2A and B). These observations indicate that

dephosphorylation of the target protein through PP4 is required for efficient assembly

of Hop1, that Tel1/Mec1 is dispensable from the phosphorylation of the target(s), and

that this phosphorylation may occur before meiotic DSB formation (Figure 6). Hop1

may be the most probable target for dephosphorylation by PP4. On the other hand, the

candidate protein kinase(s) that regulate Hop1 assembly are DDK and/or CDK.

However, the ATP analog sensitive *cdc7* allele (*cdc7-as3*) reportedly showed normal

Hop1 expression but no Hop1 phosphorylation in the presence of an ATP analog (WAN

et al. 2006). The yeast Cdc28, a component of CDK, focus formation on meiotic

chromatin are dependent on Hop1, Red1 and Rec8, but not on Mek1/Mre4. The analog

sensitive allele *cdc28-as1* reportedly displayed compromised Rad51 focus formation

and severe defects in Zip1 elongation, but did not affect Red1 loading in the presence

of an ATP analog (ZHU *et al.* 2010). These findings indicate that CDK and DDK

functions are associated with Hop1 function. However, direct phosphorylation of Hop1

by CDK and/or DDK has not been reported. There are two predicted CDK

phosphorylation motifs (S/T-P) in Hop1: S494 and S531 (ZHU *et al.* 2010). The

presence of sequential phosphorylation by CDK followed by DDK in the sequence of

serine/threonine residues in addition to the CDK phosphorylation motif has been

described for related target proteins such as Mer2 (WAN *et al.* 2006) and Mps3 (RAO *et*

al. 2020), and the S494 of Hop1 overlaps this motif (T-S-^{494}S-P).

PP4 has an independent role from ZMM/SIC components in the prophase

checkpoint

Pch2 activates the meiotic prophase checkpoint when meiotic DSB repair or

chromosome synapsis is compromised in *zip1*, *zip2*, or *dmc1* mutant cells (SAN-

SEGUNDO AND ROEDER 1999; BORNER *et al.* 2008). It has been reported that the PP4

mutant shows a *zmm* mutant-like phenotype in meiotic recombination, NCO/CO ratio

aberration, temperature-sensitive meiosis progression, CO formation, and

compromised CO interference (BORNER *et al.* 2004; FALK *et al.* 2010). In addition,

defective CO formation in *pph3Δ* is epistatic to *msh4Δ* (FALK *et al.* 2010). We also

observed compromised CO and elevated NCO formation through HD analysis (Figure

1C). These are often observed in *zmm* mutants (SHINOHARA *et al.* 2008). *pch2Δ* could

not suppress the delayed meiosis progression of the *pph3Δ* mutant. These findings

suggest that delayed meiosis progression is caused by a different mechanism from that

of *zmm* or *dmc1* mutants. Recently, it was reported that Pch2 in collaboration with

Hop1 mediates the meiotic prophase checkpoint (RAINA AND VADER 2020). In the *pph3*

mutants, compromised synapsis as well as inefficient meiotic recombination easily

activated the meiotic prophase checkpoint because of enhanced Mec1/Tel1 signaling,

but the checkpoint signal could not be restored even in the absence of Pch2. This

could be caused by insufficient Hop1 loading on the chromosome axis or accumulation

of unbound inactive form of Hop1 protein (Figure 6).

Materials and Methods

Yeast strains and media

All the yeast strains and their genotypes are listed in Supplemental table 1. We used

isogenic *S. cerevisiae* SK1 background NKY1551 derivatives (STORLAZZI *et al.* 1996).

PPH3-13MYC was constructed using a PCR-based tagging method (BAHLER *et al.*

1998). *pph3-H112N* was constructed vis two-step gene replacement. YPAD (1% yeast

extract, 2% Bacto peptone, 0.01% adenine, 2% glucose), SPM (0.3% potassium

acetate, 0.03% raffinose), and DISS plates (YPAD medium containing 2.4% Bacto

Agar) were used.

Yeast meiosis time course

The experiments were performed as described previously (SHINOHARA *et al.* 2003).

Meiosis progression was analyzed by counting the number of nuclei in each ascus at

each time point. Nuclei were visualized by staining with 4', 6-diamidino-2-phenylindole,

dihydrochloride (DAPI) and observed under an epifluorescence microscope

(AxioSkop2, Carl Zeiss). More than 100 nuclei were analyzed at each time point.

Cytological analysis

Immunostaining of yeast meiotic nuclear spreads was performed as previously

described (SHINOHARA *et al.* 2015). Stained samples were observed under

epifluorescence using the Axioskop 2 microscope equipped with light-emitting diode

fluorescence light sources (X-Cite; Excelitas Technologies) and a 100× objective

(AxioPlan, NA1.4, Carl Zeiss). Images were captured with a CCD camera (Retiga;

Qimaging) and processed using iVision (BioVision Technologies) and Photoshop

(Adobe). The antibodies used in these assays were anti-Hop1 (IWASAKI *et al.* 2016)

(guinea pig, 1:500) and anti-Hop1-pT318(IWASAKI *et al.* 2016) (rabbit, 1:500), anti-Red1

(chicken, 1:500) (SHINOHARA *et al.* 2008) and anti-Rec8 (rabbit, 1:1000) (ZHU *et al.*

2010). Nuclei containing more than five foci were counted as focus-positive.

Southern blot to detect meiotic DSBs and recombination intermediates

Genomic DNAs prepared from yeast meiosis time course experiments were digested

with *Mlu*I, *Bam*HI, and *Xho*I for heteroduplexes (HDs), and *Pst*I for meiotic DSBs.

Probes for Southern blotting were Probe 155 for HD and Probe 291 for DSB detection

(STORLAZZI *et al.* 1996; SHINOHARA AND SHINOHARA 2013). They were labeled with α-

^{32}P- dATP (NEG512H, PerkinElmer) using the random primer method (Random Primer

6, NEB S1230S, Klenow fragment (3'-5' exo-, M0210) and visualized by using

FLA7000 (Cytiva). The visualized signals were analyzed by using Image Quant TL

software (Cytiva).

Western blotting

Western blotting was performed as described previously (SHINOHARA *et al.* 2015).

Meiotic yeast cell extracts were prepared by cell disruption using a Multi-Beads Shocker

(YASUIKIKAI) after trichloroacetic acid treatment. The primary antibody signal was

visualized using Alexa Fluor 680–labeled secondary antibodies (Thermo Fisher

Scientific) or IRDye 800CW–labeled secondary antibodies (LI-COR Biosciences) using

an Odyssey infrared imaging system (LI-COR Biosciences). The primary antibodies

used in these assays were anti-γH2A (anti-γH2A, AF2288, R&D Systems, 1:1000),

anti-Myc (MC045, Nacalai Tesque, 1/1000), anti-Hop1 (guinea pig, 1/5000)(IWASAKI *et*

al. 2016), anti-Hop1-pT318 (rabbit, 1/5000) (IWASAKI *et al.* 2016), anti-Rec8 (rabbit,

1/10000) (ZHU *et al.* 2010), anti-Red1 (chicken, 1/3000) (SHINOHARA *et al.* 2008) and

anti-α-tubulin (YL1/2, Santa Cruz Biotechnology, 1/3000). Tubulin signals were

visualized using the LI-COR Bioscience system or alkaline phosphatase stain (BCIP-

NBT solution kit, Nacalai Tesque) on the same membrane on which other meiotic

proteins were detected.

Immunoprecipitation

The immunoprecipitation assay was performed as previously described with minor

modifications (SHINOHARA *et al.* 2008). Psy2-13Myc was immunoprecipitated by using

anti-Myc antibody (MC045, Nacalai Tesque) antibody which were bound to Dynabeads

Protein G (Thermo Fisher Scientific) from meiotic yeast cell lysate prepared using a

Multi-Beads shocker (YASUIKIKAI) in lysis buffer (50mM HEPES-NaOH, pH7.5,

150mM NaCl, 20% glycerol, 1mM Sodium orthovanadate, 60mM β-glycerophosphate,

0.1% Nonidet P-40, 1:100 dilution of protease inhibitor cocktail (SIGMA-Aldrich) at the

indicated time points in Figure 4A and 4B. Psy2-13Myc, Hop1, Red1 and Rec8 proteins

in the precipitates and WCEs were detected as described in the western blot section.

Tetrad analysis

Tetrad analysis was performed as described previously (CHALLA *et al.* 2019). Parental

haploid strains were mated for 12 h on YPAD plates at 30°C and then transferred onto

SPM plates. After incubation at 30°C for 48 h, tetrads were dissected onto DISS plates

and incubated for 48 h.

Acknowledgements

We are grateful to Dr. K. Matsuzaki, Dr. F. Klein, Dr. S. Gasser, and the members of

the Shinohara lab for helpful discussions. We thank Ms. A. Murakami and Ms. A.

Kitada for their excellent technical assistance. K.L. was supported by a Fellowship from

the Institute for Protein Research, Osaka University. Special thanks to Dr. A. Shinohara

for critical reading of the manuscript. This work was supported by JSPS KAKENHI

(#15H05973, #19K22402) and Takeda Science Foundation, and was performed in part

under the Cooperative Research Program of the Institute for Protein Research, Osaka

University, CR-17-05.

References

Anderson, C. M., A. Oke, P. Yam, T. Zhuge and J. C. Fung, 2015 Reduced Crossover
Interference and Increased ZMM-Independent Recombination in the Absence of
Tel1/ATM. PLoS Genet 11**:** e1005478.

Bahler, J., J. Q. Wu, M. S. Longtine, N. G. Shah, A. McKenzie, 3rd *et al.*, 1998
Heterologous modules for efficient and versatile PCR-based gene targeting in
Schizosaccharomyces pombe. Yeast 14**:** 943-951.

Bishop, D. K., 1994 RecA homologs Dmc1 and Rad51 interact to form multiple nuclear
complexes prior to meiotic chromosome synapsis. Cell 79**:** 1081-1092.

Bishop, D. K., D. Park, L. Xu and N. Kleckner, 1992 DMC1: a meiosis-specific yeast
homolog of E. coli recA required for recombination, synaptonemal complex
formation, and cell cycle progression. Cell 69**:** 439-456.

Blat, Y., R. U. Protacio, N. Hunter and N. Kleckner, 2002 Physical and functional
interactions among basic chromosome organizational features govern early steps
of meiotic chiasma formation. Cell 111**:** 791-802.

Borner, G. V., A. Barot and N. Kleckner, 2008 Yeast Pch2 promotes domainal axis
organization, timely recombination progression, and arrest of defective
recombinosomes during meiosis. Proc Natl Acad Sci U S A 105**:** 3327-3332.

Borner, G. V., N. Kleckner and N. Hunter, 2004 Crossover/noncrossover differentiation,
synaptonemal complex formation, and regulatory surveillance at the
leptotene/zygotene transition of meiosis. Cell 117**:** 29-45.

Carballo, J. A., A. L. Johnson, S. G. Sedgwick and R. S. Cha, 2008 Phosphorylation of
the axial element protein Hop1 by Mec1/Tel1 ensures meiotic interhomolog
recombination. Cell 132**:** 758-770.

Challa, K., V. G. Fajish, M. Shinohara, F. Klein, S. M. Gasser *et al.*, 2019 Meiosis-
specific prophase-like pathway controls cleavage-independent release of cohesin
by Wapl phosphorylation. PLoS Genet 15**:** e1007851.

Chen, C., A. Jomaa, J. Ortega and E. E. Alani, 2014 Pch2 is a hexameric ring ATPase
that remodels the chromosome axis protein Hop1. Proc Natl Acad Sci U S A
111**:** E44-53.

Chuang, C. N., Y. H. Cheng and T. F. Wang, 2012 Mek1 stabilizes Hop1-Thr318
phosphorylation to promote interhomolog recombination and checkpoint
responses during yeast meiosis. Nucleic Acids Res 40**:** 11416-11427.

De Muyt, A., A. Pyatnitskaya, J. Andreani, L. Ranjha, C. Ramus *et al.*, 2018 A meiotic
633   XPF-ERCC1-like complex recognizes joint molecule recombination
634   intermediates to promote crossover formation. Genes Dev 32: 283-296.

635   Falk, J. E., A. C. Chan, E. Hoffmann and A. Hochwagen, 2010 A Mec1- and PP4-
636   dependent checkpoint couples centromere pairing to meiotic recombination. Dev
637   Cell 19: 599-611.

638   Farmer, S., E. J. Hong, W. K. Leung, B. Argunhan, Y. Terentyev *et al.*, 2012 Budding
639   yeast Pch2, a widely conserved meiotic protein, is involved in the initiation of
640   meiotic recombination. PLoS One 7: e39724.

641   Garcia, V., S. Gray, R. M. Allison, T. J. Cooper and M. J. Neale, 2015 Tel1(ATM)-
642   mediated interference suppresses clustered meiotic double-strand-break
643   formation. Nature 520: 114-118.

644   Hoffmann, R., S. Jung, M. Ehrmann and H. W. Hofer, 1994 The Saccharomyces
645   cerevisiae gene PPH3 encodes a protein phosphatase with properties different
646   from PPX, PP1 and PP2A. Yeast 10: 567-578.

647   Hollingsworth, N. M., and B. Byers, 1989 HOP1: a yeast meiotic pairing gene. Genetics
648   121: 445-462.

649   Hollingsworth, N. M., L. Goetsch and B. Byers, 1990 The HOP1 gene encodes a
650   meiosis-specific component of yeast chromosomes. Cell 61: 73-84.

651   Hustedt, N., A. Seeber, R. Sack, M. Tsai-Pflugfelder, B. Bhullar *et al.*, 2015 Yeast PP4
652   interacts with ATR homolog Ddc2-Mec1 and regulates checkpoint signaling.
653   Mol Cell 57: 273-289.

654   Iwasaki, D., K. Hayashihara, H. Shima, M. Higashide, M. Terasawa *et al.*, 2016 The
655   MRX Complex Ensures NHEJ Fidelity through Multiple Pathways Including
656   Xrs2-FHA-Dependent Tel1 Activation. PLoS Genet 12: e1005942.

657   Joshi, N., M. S. Brown, D. K. Bishop and G. V. Borner, 2015 Gradual implementation
658   of the meiotic recombination program via checkpoint pathways controlled by
659   global DSB levels. Mol Cell 57: 797-811.

660   Keeney, S., 2001 Mechanism and control of meiotic recombination initiation. Curr Top
661   Dev Biol 52: 1-53.

662   Keogh, M. C., J. A. Kim, M. Downey, J. Fillingham, D. Chowdhury *et al.*, 2006 A
663   phosphatase complex that dephosphorylates gammaH2AX regulates DNA
664   damage checkpoint recovery. Nature 439: 497-501.

665   Klein, F., P. Mahr, M. Galova, S. B. Buonomo, C. Michaelis *et al.*, 1999 A central role
666   for cohesins in sister chromatid cohesion, formation of axial elements, and
667   recombination during yeast meiosis. Cell 98: 91-103.

Lai, Y. J., F. M. Lin, M. J. Chuang, H. J. Shen and T. F. Wang, 2011 Genetic
requirements and meiotic function of phosphorylation of the yeast axial element
protein Red1. Mol Cell Biol 31**:** 912-923.
Lee, B. H., and A. Amon, 2003 Role of Polo-like kinase CDC5 in programming meiosis
I chromosome segregation. Science 300**:** 482-486.
Lo, Y. H., C. N. Chuang and T. F. Wang, 2014 Pch2 prevents Mec1/Tel1-mediated
Hop1 phosphorylation occurring independently of Red1 in budding yeast
meiosis. PLoS One 9**:** e85687.
Niu, H., L. Wan, B. Baumgartner, D. Schaefer, J. Loidl *et al.*, 2005 Partner choice
during meiosis is regulated by Hop1-promoted dimerization of Mek1. Mol Biol
Cell 16**:** 5804-5818.
Novak, J. E., P. B. Ross-Macdonald and G. S. Roeder, 2001 The budding yeast Msh4
protein functions in chromosome synapsis and the regulation of crossover
distribution. Genetics 158**:** 1013-1025.
O'Neill, B. M., S. J. Szyjka, E. T. Lis, A. O. Bailey, J. R. Yates, 3rd *et al.*, 2007 Pph3-
Psy2 is a phosphatase complex required for Rad53 dephosphorylation and
replication fork restart during recovery from DNA damage. Proc Natl Acad Sci
U S A 104**:** 9290-9295.
Panizza, S., M. A. Mendoza, M. Berlinger, L. Huang, A. Nicolas *et al.*, 2011 Spo11-
accessory proteins link double-strand break sites to the chromosome axis in
early meiotic recombination. Cell 146**:** 372-383.
Raina, V. B., and G. Vader, 2020 Homeostatic Control of Meiotic Prophase Checkpoint
Function by Pch2 and Hop1. Curr Biol 30**:** 4413-4424 e4415.
Rao, H. B. D. P., T. Sato, K. Challa, M. Shinohara and A. Shinohara, 2020
Phosphorylation of luminal region of the SUN-domain protein Mps3 promotes
nuclear envelope localization during meiosis. bioRxiv**:** 2020.2009.2015.297762.
Rockmill, B., and G. S. Roeder, 1988 RED1: a yeast gene required for the segregation
of chromosomes during the reductional division of meiosis. Proc Natl Acad Sci
U S A 85**:** 6057-6061.
San-Segundo, P. A., and G. S. Roeder, 1999 Pch2 links chromatin silencing to meiotic
checkpoint control. Cell 97**:** 313-324.
Sasanuma, H., H. Murakami, T. Fukuda, T. Shibata, A. Nicolas *et al.*, 2007 Meiotic
association between Spo11 regulated by Rec102, Rec104 and Rec114. Nucleic
Acids Res 35**:** 1119-1133.

Schwacha, A., and N. Kleckner, 1997 Interhomolog bias during meiotic recombination:
703   meiotic functions promote a highly differentiated interhomolog-only pathway.
704   Cell 90**:** 1123-1135.
705   Shi, Y., 2009 Serine/threonine phosphatases: mechanism through structure. Cell 139**:**
706   468-484.
707   Shinohara, A., S. Gasior, T. Ogawa, N. Kleckner and D. K. Bishop, 1997
708   Saccharomyces cerevisiae recA homologues RAD51 and DMC1 have both
709   distinct and overlapping roles in meiotic recombination. Genes Cells 2**:** 615-629.
710   Shinohara, M., D. K. Bishop and A. Shinohara, 2019 Distinct Functions in Regulation
711   of Meiotic Crossovers for DNA Damage Response Clamp Loader
712   Rad24(Rad17) and Mec1(ATR) Kinase. Genetics 213**:** 1255-1269.
713   Shinohara, M., S. L. Gasior, D. K. Bishop and A. Shinohara, 2000 Tid1/Rdh54
714   promotes colocalization of rad51 and dmc1 during meiotic recombination. Proc
715   Natl Acad Sci U S A 97**:** 10814-10819.
716   Shinohara, M., K. Hayashihara, J. T. Grubb, D. K. Bishop and A. Shinohara, 2015 DNA
717   damage response clamp 9-1-1 promotes assembly of ZMM proteins for
718   formation of crossovers and synaptonemal complex. J Cell Sci 128**:** 1494-1506.
719   Shinohara, M., S. D. Oh, N. Hunter and A. Shinohara, 2008 Crossover assurance and
720   crossover interference are distinctly regulated by the ZMM proteins during yeast
721   meiosis. Nat Genet 40**:** 299-309.
722   Shinohara, M., K. Sakai, T. Ogawa and A. Shinohara, 2003 The mitotic DNA damage
723   checkpoint proteins Rad17 and Rad24 are required for repair of double-strand
724   breaks during meiosis in yeast. Genetics 164**:** 855-865.
725   Shinohara, M., and A. Shinohara, 2013 Multiple pathways suppress non-allelic
726   homologous recombination during meiosis in Saccharomyces cerevisiae. PLoS
727   One 8**:** e63144.
728   Storlazzi, A., L. Xu, A. Schwacha and N. Kleckner, 1996 Synaptonemal complex (SC)
729   component Zip1 plays a role in meiotic recombination independent of SC
730   polymerization along the chromosomes. Proc Natl Acad Sci U S A 93**:** 9043-
731   9048.
732   Subramanian, V. V., A. J. MacQueen, G. Vader, M. Shinohara, A. Sanchez *et al.*, 2016
733   Chromosome Synapsis Alleviates Mek1-Dependent Suppression of Meiotic
734   DNA Repair. PLoS Biol 14**:** e1002369.
735   Sun, X., L. Huang, T. E. Markowitz, H. G. Blitzblau, D. Chen *et al.*, 2015 Transcription
736   dynamically patterns the meiotic chromosome-axis interface. Elife 4.

Sym, M., J. A. Engebrecht and G. S. Roeder, 1993 ZIP1 is a synaptonemal complex
protein required for meiotic chromosome synapsis. Cell 72**:** 365-378.
Tsubouchi, T., and G. S. Roeder, 2005 A synaptonemal complex protein promotes
homology-independent centromere coupling. Science 308**:** 870-873.
Wan, L., C. Zhang, K. M. Shokat and N. M. Hollingsworth, 2006 Chemical inactivation
of cdc7 kinase in budding yeast results in a reversible arrest that allows efficient
cell synchronization prior to meiotic recombination. Genetics 174**:** 1767-1774.
West, A. M. V., E. A. Komives and K. D. Corbett, 2018 Conformational dynamics of
the Hop1 HORMA domain reveal a common mechanism with the spindle
checkpoint protein Mad2. Nucleic Acids Res 46**:** 279-292.
Zhang, L., K. P. Kim, N. E. Kleckner and A. Storlazzi, 2011 Meiotic double-strand
breaks occur once per pair of (sister) chromatids and, via Mec1/ATR and
Tel1/ATM, once per quartet of chromatids. Proc Natl Acad Sci U S A 108**:**
20036-20041.
Zhu, Z., S. Mori, H. Oshiumi, K. Matsuzaki, M. Shinohara *et al.*, 2010 Cyclin-
dependent kinase promotes formation of the synaptonemal complex in yeast
meiosis. Genes Cells 15**:** 1036-1050.
754
755

Figure legends

Figure1. PP4 dephosphorylation activity is required for proper meiotic recombination.

(A) Schematic of the functional domain structure of yeast Pph3 (*sc*Pph3) protein. Amino acid sequences contributing to phosphate binding and metal coordination are shown in pale blue and red, respectively. The position of the *pph3-H112N* and the substitution of histidine 112 with asparagine is shown.

(B) Localization of Psy2-13Myc was analyzed in *PPH3* (MSY6198/6190) and the *pph3-H112N* (MSY6242/6244) background by immunostaining with anti-Myc antibody (green). Representative images at 6h in meiosis are shown. Scale bar indicates 2 μm.

(C) Schematics of the *HIS4-LEU2* recombination hotspot. Diagnostic P: *Pst*I, X: *Xho*I, B: *Bam*HI and M: *Mlu*I restriction sites are shown. The size of meiotic DSBs (DSB I, DSB II), and parental (Pa), and heteroduplexes (HDs) that are associated with COs (HD2 and HD3) or NCOs (HD1 and HD4), Parental (Pa1 and Pa2) and the positions of detectable probes 291 and 155 are shown.

(D) Representative Southern blotting of DSB detection in wild type (black, NKY1303/1543), *pph3Δ* (red, MSY5632/MSY5634), *pph3-H112N* (blue, MSY6219/6220), p*CLB2-MEC1 tel1Δ* (purple, MSY5544/5546) and p*CLB2-MEC1 tel1Δ pph3Δ* (orange, MSY5760/5762) using probe 291. The graph shows the mean and standard error of the quantified DSB I signals from more than three times trials.

(E) Representative Southern blotting of HD-associated COs and NCOs formed at

the *HIS4-LEU2* hotspot using probe155. The upper graph shows the mean and

standard error of the quantified HD-associated COs (HD2 and HD3). The lower

graph shows the mean and standard error of the quantified HD-associated

NCOs (HD1 and HD4) of wild type, *pph3Δ*, and *pph3-H112N.*

(F) Representative immunostaining images of Rad51 (green) at 3 h on meiotic

nuclear spreads in wild type (NKY1303/1543), *pph3Δ* (MSY5632/MSY5634),

and *pph3-H112N* (MSY6219/6220). Scale bar indicates 2 μm. The dot plot

graph shows the distribution of the number of Rad51 foci in each Rad51 focus

positive nucleus at the indicated time points in wild type (black, NKY1303/1543),

pph3Δ (red, MSY5632/MSY5634), and *pph3-H112N* (blue, MSY6219/6220).

The data are presented as the medians and 95% confidence interval. Asterisks

indicate statistically significant differences between the indicated strains

determined using the Mann-Whitney U-test (* $P<0.05$, ** $P<0.01$, **** $P<0.0001$)

using Prism9 software.

Figure 2. Hop1 assembly is delayed in the *pph3* mutant in early prophase-I.

(A) Graphic depiction of the kinetics of Hop1 focus (upper) and Hop1-pT318 focus

(lower) assembly on meiotic nuclear spreads in the indicated strains, wild type

(black, NKY1303/1543), *pph3Δ* (red, MSY5632/MSY5634), *pph3-H112N* (blue,

MSY6219/6220), p*CLB2-MEC1 tel1Δ* (purple, MSY5544/5546) and p*CLB2-*

MEC1 tel1Δ pph3Δ (orange, MSY5760/5762). More than 100 cells were

analyzed in each time point. Error bars show the mean ± standard deviation

from more than three times trials.

(B) Representative cytological immunostaining images of Hop1 (green) and Hop1-

pT318 (red) on meiotic nuclear spreads at 3 h in wild type (NKY1303/1543),

pph3Δ (MSY5632/MSY5634), and *pph3-H112N* (MSY6219/6220). Scale bar

indicates 2 μm.

(C) Representative western blot images of phosphorylation of histone H2A (γH2A,

red) and α-tubulin expression (green) at the indicated time point during meiosis

(upper panel). Expression of Hop1 (green, top), phosphorylation of Hop1 at

T318 (Hop1-pT318, red, middle), merged images of Hop1 and Hop1-pT318,

and α-tubulin expression (bottom) at the indicated time points during meiosis. M

denotes molecular weight markers.

(D) Grapic depiction of the average of the relative Hop1 protein signals from

western blot at each time point. The numerical values are shown as a

percentage of the peak amount of signal in the wild type. Error bars show the

mean and ± standard deviation from two independent trials.

Figure 3. Rec8 expression is normal in the *pph3* mutant in early prophase I.

(A) Representative cytological immunostaining images of Rec8 (green) and Red1

(red) on meiotic nuclear spreads at 3 h in wild type (NKY1303/1543), *pph3Δ*

(MSY5632/MSY5634), *pph3-H112N* (MSY6219/6220). wild type

(NKY1303/1543), *pch2Δ* (MSY6446/6448), *pph3Δ* (MSY5632/MSY5634) and

pph3Δ pch2Δ (MSY6417/6419)

(B) Graphs show the kinetics of Rec8 (upper) and Red1 (lower) focus assembly on

meiotic nuclear spreads in the indicated strains, wild type (black,

NKY1303/1543), *pph3Δ* (red, MSY5632/MSY5634), and *pph3-H112N* (blue,

MSY6219/6220). More than 100 cells were analyzed at each time point. Error

bars show the mean ± standard deviation from more than three times trials.

(C) Representative western blot images of expression of Rec8 (upper) and Red1

(lower) with α-tubulin expression images as internal controls at the indicated

time points during meiosis. M denotes molecular weight markers. Asterisks

indicate non-specific protein signals.

Figure 4. PP4 interacts with Hop1 during meiosis.

(A) Immunoprecipitation (IP) analysis of Psy2-13Myc from *PPH3* (non-tag,

NKY1303/1543), *PPH3* (*PSY2-13MYC*, MSY6188/6190), *pph3-H112N*

(MSY6219/6220) and *pph3-H112N* (*PSY2-13MYC*, MSY6242/6244) cells at the

indicated time points during meiosis. Immunoprecipitation products were probed

with anti-Myc and anti-Hop1 antibodies.

(B) Immunoprecipitation (IP) analysis of Psy2-13Myc from *PPH3* (non-tag,

NKY1303/1543) and *PPH3* (*PSY2-13MYC*, MSY6188/6190) cells at the

indicated time points during meiosis. Whole cell extracts and

immunoprecipitation products were probed with anti-Myc, anti-Hop1, anti-Red1

and anti-Rec8 antibodies.

(C) Tetrad analysis of spore viability in wild type (NKY1303/1543), *pph3Δ*

(MSY5632/MSY5634), *pch2Δ* (MSY6446/6448) and *pph3Δ pch2Δ*

(MSY6417/6419). The graphs show the fraction of tetrads containing 0-, 1-, 2-,

3- and 4-viable spores. The overall spore viability is shown in parentheses. n:

number of asci analyzed.

(D) Graphic depiction of the kinetics of Hop1 (left) and Hop1-pT318 (right) focus

assembly on meiotic nuclear spreads in the indicated strains, wild type (black,

NKY1303/1543), *pch2Δ* (gray, MSY6446/6448), *pph3Δ* (red, MSY5632/MSY5634) and *pph3Δ pch2Δ* (orange, MSY6417/6419). More than

100 cells were analyzed at each time point. Error bars show the mean ±

standard deviation from more than three times trials.

(E) Representative cytological immunostaining images of Hop1 (green) and Hop1-

pT318 (red) on meiotic nuclear spreads at 3 h (left panel) and 6 h (right panel)

in wild type (NKY1303/1543), *pch2Δ* (MSY6446/6448), *pph3Δ*

(MSY5632/MSY5634) and *pph3Δ pch2Δ* (MSY6417/6419). Scale bar indicates

2 μm.

Figure 5. *pch2Δ* does not completely suppress the defect in Hop1 assembly

of *pph3Δ*.

(A) Zip1 and Hop1 elongation was analyzed in wild type (NKY1303/1543), *pch2Δ*

(MSY6446/6448), *pph3Δ* (MSY5632/MSY5634), and *pph3Δ pch2Δ*

(MSY6417/6419) via immunostaining at each time point. The graphs show the

percentages of each class of Zip1 morphologies; linear (dark gray, pachytene),

partial (pale gray, zygotene) and dotty (black, leptotene). Error bars show the

mean ± standard deviation from more than three trials. More than 100 cells at

each time point were analyzed.

(B) Representative cytological immunostaining images of Hop1 (green) and Zip1

(red) on pachytene-phase nuclear spreads at 4 h in wild type (NKY1303/1543),

and 6 h in the *pch2Δ* (MSY6446/6448), *pph3Δ* (MSY5632/MSY5634) and

pph3Δ pch2Δ (MSY6417/6419). Scale bar indicates 2 μm.

(C) The dot plot graph of the distribution of the number of linearized Hop1 signals in

each pachytene stage nucleus in *pch2Δ* (N = 112) and *pph3Δ pch2Δ* (N = 109)

at 6 h during meiosis from three independent trials. Error bars show the mean

with 95% confidence interval. Statistically significant differences between the

pch2Δ and *pph3Δ pch2Δ* were determined using the Mann-Whitney U-test (****

P<0.0001) using Prism9 software.

(D) The dot plot graph of the distribution of lengths of linearized Hop1 signal of

pachytene stage nucleus in *pch2Δ* (gray, N = 718) and *pph3Δ pch2Δ* (orange,

N = 780) at 6 h in meiosis from three independent trials. Error bars show the

mean with 95% confidence interval. Statistically significant differences between

the *pch2Δ* and *pph3Δ pch2Δ* were determined using the Mann-Whitney U-test

(ns; not significant) using Prism9 software.

(E) Representative cytological immunostaining images of Rec8 (green) and Red1

(red) on pachytene-stage nuclear spreads in wild type (3 h, NKY1303/1543),

pch2Δ (6 h, MSY6446/6448), *pph3Δ* (6 h, MSY5632/MSY5634) and *pph3Δ*

pch2Δ (6 h, MSY6417/6419). Scale bar indicates 2 μm.

(F) Representative western blot images of expression of Hop1 expression (green)

and phosphorylation of Hop1 at pT318 (Hop1-pT318, red) and α-tubulin

expression (bottom) at the indicated time points during meiosis. M denotes

molecular weight markers.

(G) Meiosis progression in wild type (black, NKY1303/1543), *pch2Δ* (gray,

MSY6446/6448), *pph3Δ* (red, MSY5632/MSY5634) and *pph3Δ pch2Δ* (orange,

MSY6417/6419) mutants are shown. Frequencies of nuclei with more than two

DAPI-staining bodies at each time point are plotted. More than 100 cells were

analyzed at each time point. Error bars show the means and standard deviation

from two independent trials.

Figure 6. Model for PP4 involvement in Hop1 loading onto meiotic chromatin.

PP4 plays multiple roles during meiotic prophase. First, PP4 promotes Hop1

loading in the early phase of meiotic prophase in a Mec1/Tel1-independent manner.

Until the chromatin loading, Hop1 exists in an inactive form, which might prevent

miosis progression. After meiotic DSB formation, Hop1 is phosphorylated at the

DSB site by Tel1/Mec1, and PP4 is involved in dephosphorylation of the

phosphorylated Hop1. Meiotic recombination promotes homolog synapsis by

exchanging Hop1 to Zip1 through the function of AAA+ Pch2 function.

Supplemental figure 1. **Supplemental data for Figure 1.**

(A) Representative western blot images showing Psy2-13myc and α-tubulin

expression at the indicated time points during meiosis in wild type (non-tag,

NKY1303/1543), *pph3Δ* (non-tag, MSY5632/5634), PSY2-13myc

(MSY6188/6190), *pph3-H112N* (non-tag, MSY6219/6220) and *PSY2-13MYC*

pph3-H112N (MSY6242/6244).

(B) Meiosis progression in wild type (NKY1303/1543), *pph3Δ* (MSY5632/5634) and

pph3-H112N (MSY6219/6220) mutants. Frequencies of nuclei with more than

two DAPI-stained bodies are plotted.

(C) The graph of the kinetics of Rad51 focus assembly on meiotic nuclear spreads

in wild type (black, NKY1303/1543), *pph3Δ* (red, MSY5632/MSY5634), and

pph3-H112N (blue, MSY6219/6220). More than 100 cells were analyzed at

each time point. Error bars show the mean ± standard deviation from more than

three times trials.

Supplemental Figure 2. **Supplemental data presentation for Figure 4.**

(A) Whole membrane images of western blot of immunoprecipitates (IP) are

shown in Figure 4A and western blot of whole cell extracts (WCEs) before the

immunoprecipitation.

(B) Whole membrane images of western blot of immunoprecipitates (IP) and

whole cell extracts (WCEs) exhibited in Figure 4B,

Li *et al* Figure 1

Li *et al* Figure 2

A

Hop1 assembly

Hop1-pT318 assembly

- ● Wild type
- ● *pph3Δ*
- ● *pph3-H112N*
- ● *pCLB2-mec1 tel1Δ*
- ● *pCLB2-mec1 tel1Δ pph3Δ*

B

Wild type

pph3Δ

pph3-H112N

pCLB2-MEC1 tel1Δ — *PPH3*

pph3Δ

3 h in meiosis

C

D

Hop1 expression

Li *et al* Figure 3

A

3 h in meiosis

B

Red1

Rec8

C

Li *et al* Figure 4

Li *et al* Figure 5

Supplemental Table-1 Strain list

Strain	Genotype	Reference
NKY1303	*MATa, ho::LYS2, lys2, ura3, leu2::hisG, his4B-LEU2(MluI), arg4 bgl*	(STORLAZZI *et al.* 1996)
NKY1543	*MATa, ho::LYS2, lys2, ura3, leu2::hisG, his4X-LEU2(BamHI)-URA3, arg4-nsp*	(STORLAZZI *et al.* 1996)
MSY5632	NKY1303 with *pph3::HygR*	This study
MSY5634	NKY1543 with *pph3::HygR*	This study
MSY6219	NKY1303 with *pph3-H112N::MluI*	This study
MSY6220	NKY1543 with *pph3-H112N::MluI*	This study
MSY6198	NKY1303 with *PSY2-13MYC::TRP1*	This study
MSY6190	NKY1543 with *PSY2-13MYC::TRP1*	This study
MSY6242	NKY1303 with *PSY2-13MYC::TRP1, pph3-H112N::MluI*	This study
MSY6244	NKY1543 with *PSY2-13MYC::TRP1, pph3-H112N::MluI*	This study
MSY5544	NKY1303 with *pCLB2-MEC1, tel1::TRP1*	This study
MSY5546	NKY1543 with *pCLB2-MEC1, tel1::TRP1*	This study
MSY5760	NKY1303 with *pCLB2-MEC1, tel1::TRP1, pph3::HygR*	This study
MSY5762	NKY1543 with *pCLB2-MEC1, tel1::TRP1, pph3::HygR*	This study
MSY6446	NKY1303 with *pch2::TRP1*	This study
MSY6448	NKY1543 with *pch2::TRP1*	This study
MSY6417	NKY1303 with *pph3::HygR, pch2::TRP1*	This study
MSY6519	NKY1543 with *pph3::HygR, pch2::TRP1*	This study

Storlazzi, A., L. Xu, A. Schwacha and N. Kleckner, 1996 Synaptonemal complex (SC) component Zip1 plays a role in meiotic recombination independent of SC polymerization along the chromosomes. Proc Natl Acad Sci U S A 93: 9043-9048.

A

B

C

Supplemental Figure 2, Li *et al*

A

WCEs | IP: anti-Myc

PPH3 *pph3-H112N* | *PPH3* *pph3-H112N*

PSY2-13MYC − − − + + + − − − + + + − − − + + + − − − + + +

Time in meiosis 2 4 6 2 4 6 2 4 6 2 4 6 | 2 4 6 2 4 6 2 4 6 2 4 6

← Psy2-13Myc

← Hop1

B

WCEs | anti-Myc IP

PSY2-13MYC − − − + + + | − − − + + +

PSY2 + + + − − − | + + + − − −

Time in meiosis 2 4 6 2 4 6 | 2 4 6 2 4 6 (h)

← Psy2-13Myc

← Hop1

← Red1

← Rec8