Discovery of novel ID2 antagonists from pharmacophore-based virtual screening as potential therapeutics for glioma

Genshen Zhong[a,1,*], Yichun Wang[a,1], Qi Wang[a], Minna Wu[b], Yichuang Liu[c], Shitao Sun[c], Zhenli Li[c], Jinle Hao[c], Peiyuan Dou[d], Bin Lin[c,*]

[a] *Henan Key Laboratory of Immunology and Targeted Therapy, Henan Collaborative Innovation Center of Molecular Diagnosis and Laboratory Medicine, School of Laboratory Medicine, Xinxiang Medical University, Xinxiang, Henan, 453003, China*

[b] *School of Basic Medicine, Xinxiang Medical University, Xinxiang, Henan, 453003, China*

[c] *School of Pharmaceutical Engineering, Shenyang Pharmaceutical University, Shenyang, Liaoning, 110016, China*

[d] *School of Chemistry, Cardiff University, Cardiff, CF10 3AT, United Kingdom*

[1] These authors contributed equally to this work.

[*] Corresponding Authors at: Henan Key Laboratory of Immunology and Targeted Therapy, Henan Collaborative Innovation Center of Molecular Diagnosis and Laboratory Medicine, School of Laboratory Medicine, Xinxiang Medical University, Xinxiang, Henan, 453003, China (G. Zhong) and School of Pharmaceutical Engineering, Shenyang Pharmaceutical University, Shenyang, Liaoning, 110016, China (B. Lin).

E-mail addresses: zhonggx@xxmu.edu.cn (G. Zhong), randybinlin@gmail.com (B. Lin).

Highlights

- Two pharmacophores were built from the first-in-class pan-ID antagonists AGX51
- A pharmacophore-based virtual screening was carried out and 16 hit compounds were purchased for pharmacological evaluations in glioma inhibition
- Compound AK-778-XXMU was identified to be a potent ID2 antagonist in the low submicromolar range (KD: 159 nM)

Abstract

Glioma, especially the most aggressive type glioblastoma multiforme, is one of the central nervous system malignant cancer with a poor prognosis. Traditional treatments are mainly surgery combined with radiotherapy and chemotherapy, which is still not satisfactory. Therefore, it is of great clinical significance to find new therapeutic agents. Served as an inhibitor of differentiation, protein ID2 (inhibitor of DNA binding 2) plays an important role in neurogenesis, neovascularization and malignant development of gliomas. It has been shown that ID2 affects the malignant progression of gliomas through different mechanisms. In this study, a pharmacophore-based virtual screening was carried out and 16 hit compounds were purchased for pharmacological evaluations on their ID2 inhibitory activities. Based on the cytotoxicity of these small-molecule compounds, two compounds were shown to effectively inhibit the viability of glioma cells in the low micromolar range. Among them, AK-778-XXMU was chosen for further study due to its better solubility in water.

A SPR assay proved the high affinity between AK-778-XXMU and ID2 protein with

the KD value as 129 nM. The plausible binding mode in the biding site of ID2 was

studied by molecular docking. Subsequently, the cancer-suppressing potency of the

compound was characterized both *in vitro* and *in vivo*. The data demonstrated that

compound AK-778-XXMU is a potent ID2 antagonist which has the potential to be

developed as a therapeutic agent against glioma.

Keywords: glioma, inhibitor of differentiation, pharmacophore, virtual screening

Introduction

Malignant glioma is the most aggressive brain tumor with a high rate of fatality and

poor prognosis [1–3]. Despite the enormous efforts invested to improve the efficacy of

classical combination of surgical resection supplemented by chemotherapeutic agents,

the overall survival of the patients is still not promising [2]. The chemotherapeutic

agents currently approved by the FDA for the treatment of malignant glioma include

temozolomide (Temodar, TMZ) and bevacizumab (Avastin). Unfortunately, as the

first-line drug for glioblastoma, which is the most aggressive type of glioma, the

emergence of TMZ resistance has greatly limited the effectiveness of the drug.

Bevacizumab is currently the only targeted drug approved by FDA for the treatment of

recurrent glioblastoma [4]. It has also been reported that bevacizumab can prolong the

progression-free survival of newly diagnosed glioblastoma patients, but the overall

survival time cannot be prolonged [5]. Moreover, the discovery of a subtype of tumor

cell with stem cell-like self-renewal properties [6], along with insufficient tumor

resection due to invasion of normal brain tissue by tumor cells [7] has been shown to

result in relatively ineffective treatment of glioma [8]. Therefore, it is of great urgence

to develop novel potential therapeutics against novel targets that play key roles in the

progression of glioma.

Inhibitors of DNA binding (ID) proteins, commonly expressed in mammals

synergistically, regulate gene expression, cell lineage localization and cell

differentiation in tissues [9,10]. They belong to the family of helix-loop-helix (HLH)

transcription factors. However, the absence of a basic motif which is necessary for

binding to DNA distinguishes it from the basic helix-loop-helix transcription factors

(bHLH), which contributes to its role of transcription factor inhibitor. Ever since the

first ID gene, ID1, was identified in 1990 by Benezra *et al* [9], three more ID groups

were discovered and named ID2, ID3, and ID4 [11–14]. The crucial role of the ID

proteins is due to their dominant-negative effect when they form heterodimers with

other DNA-binding members of the HLH family, therefore disrupting the protein-DNA

interaction. Not surprisingly, strong association of ID proteins with tumor angiogenesis

has been suggested [15,16]. In particular, the emerging role of ID2 in glioma has

attracted much attention [17]. It was found that ID2 was expressed to varying degrees in

different grades of glioma and was significantly increased in mesenchymal astrocytoma

and glioblastoma compared to low-grade malignant astrocytoma, which was not

detected in normal cerebral white matter [18]. Furthermore, the expression level of ID2

was significantly correlated with the survival of glioblastoma cells in a low glucose

metabolic stress environment by influencing mitochondrial function [19,20]. During

the development of myeloid cells，Kinase insert domain receptor (KDR, a type IV

receptor tyrosine kinase), also known as vascular endothelial growth factor receptor 2

(VEGFR-2), a critical gene for VEGF signaling, has the ability to drive differentiation

of hematopoietic progenitor cells (HPCs) into tumor-promoting myeloid-derived

suppressor cells (MDSCs) which could be enhanced by ID2 by upregulating KDR and

then stimulates the formation of an aggressive pro-angiogenic phenotype in gliomas

[17,21]. It is believed that neural precursor cells (NPCs), a type of neural stem cells,

may also be one origin for glioma cell [22,23]. Phosphorylation of ID2 N-terminal seems to affect its protein expression level in NPCs, in the case of this phosphorylated mutant, ID2 appears to be protected from degradation by the ubiquitin-proteasome system (UPS) to perform a range of pro-oncogenic functions [24].

The ID genes/proteins are critically important during embryonic development but are only active in adults when cancer is present [25–27]. It was found that they are expressed in virtually all cancers [25,28]. The overexpression of ID genes in tumors is linked to an aggressive phenotype and poor clinical outcome. It was proposed that targeting ID proteins was an effective strategy to develop anti-cancer drugs [17].

So far, a variety of strategies have been proposed to interfere with ID activity, including the use of targeted antisense delivery, cell-permeable peptides, antagonists of Id expression, and inhibitors of stability-inducing deubiquitinases, antagonists of ID proteins [29–34]. Among them, the first and only small-molecule pan-ID antagonist is AGX51, discovered by Benezra group [35]. It was shown that AGX51 was able to inhibit Id1-E47 interaction, leading to ubiquitin-mediated degradation of IDs, cell growth arrest, and reduced viability. AGX51 was also able to phenocopy the genetic loss of Id expression in age-related macular degeneration (AMD) and retinopathy of prematurity (ROP) models by inhibiting retinal neovascularization [35]. Due to the therapeutical potential of ID antagonists, it is important to develop more such compounds to target ID proteins.

Therefore, we launched a drug discovery project by utilizing a pharmacophore-based virtual screening approach. Docking-based virtual screening was not chosen because

even though ID2 protein has been already solved by crystallography (PDB: 4AYA), the

binding site is located at the loop region. Such a region is very flexible in nature so that

the binding site may not have a well-defined shape [36]. In contrast,

pharmacophore-based virtual screening does not require the structure of the protein

[37–40]. The pharmacophore was extracted from AGX51 and subjected to a series of

small commercial databases for virtual screening.

In the end, a variety of potential ID2 antagonists were obtained and 16 hit

compounds were purchased for pharmacological evaluations for their ID2 inhibitory

activities. Based on the cytotoxicity of these small-molecule compounds, two

compounds were shown to effectively inhibit the viability of glioma cells at low

micromolar concentrations. Among them, AK-778-XXMU was chosen for further

study due to its better solubility in water. A SPR (surface plasma resonance) assay

proved the high affinity between AK-778-XXMU and ID2 protein with the KD value as

129 nM. Subsequently, the cancer-suppressing potency of the compound was

characterized both *in vitro* and *in vivo*. The data demonstrated that compound

AK-778-XXMU is a potent ID2 antagonist which has the potential to be developed as a

therapeutic agent against glioma.

Results and Discussion

1. Results from pharmacophore-based virtual screening

The commercial compound databases used for virtual screening were

CHEMBRIDGE, SPECS, CHEMDIV, ENAMINE, IBS, LIFECHEMICALS,

VITAS-M, kindly provided in their electronic form by Topscience Co., Ldt

(https://www.tsbiochem.com/). They contained 2,281 compounds in total that were

commercially available. Two pharmacophore models (*R*-model and *S*-model) were

built from the two enantiomers of AGX51 and they were subjected for virtual screening

against those databases mentioned above by using LigandScout 4.0 [38,40]. The

screening yielded 24 compounds from the *R*-model and 33 compounds from the

S-model. After consultation with synthetic chemists for the feasibility of future

chemical modifications of those compounds, 16 compounds were purchased. Their

chemical structures are listed in **Fig. 1** with their compound names from the respective

databases.

Fig. 1. Chemical structures of AGX51 and the 16 hit compounds purchased. The chiral center was

labeled with an asterisk. AK-778/43420895 was highlighted in the round rectangle and renamed as

AK-778-XXMU for further study.

2. Determination of the cytotoxicity of the 16 hit compounds by CCK8 assay

CCK8 assays were conducted next to measure the viability of different glioma cell

lines, namely U87, HS683, and GL261, after treated with different concentrations of

the 16 hit compounds *in vitro* to examine the cytotoxicity of these compounds against

glioma. As shown in **Table 1**, all compounds except three demonstrated detectable

cytotoxicity ($IC_{50} < 100$ μM) against all three glioma cell lines, proving the validity of

our pharmacophore-based virtual screening. Among them, compounds

AK-778/43465022 and AK-778/43420895 had the lowest IC_{50} values to glioma cells

($IC_{50} < 30$ μM), indicating they could inhibit cell viability more effectively. Although

compound AK-778/43465022 showed the most potent inhibition in all three glioma cell

lines, it had poor water solubility **(Fig. 2A)**. Therefore, compound AK-778/43420895

was chosen for further studies. Normal cell lines, human normal hepatocytes (L-02)

and mouse embryo fibroblasts (3T3) were exposed to AK-778/43420895 to rule out the

potential toxicity and other possible side effects. It was shown in **Fig. 2C** that

AK-778/43420895 had little cytotoxicity on normal cell lines. The compound was

re-named as AK-778-XXMU for the subsequent studies.

Table 1. The cytotoxicity against various glioma cell lines from CCK8 assay

Compound Names	IC$_{50}$ (μM)		
	U87	HS683	GL261
K784-9509	56.2±3.1	55.3±2.1	48.1±0.8
E755-0111	46.5±1.3	40.8±1.5	47.9±2.7
C141-0292	77.9±3.1	84.5±1.7	90.2±1.5
E642-0121	60.3±1.0	62.3±0.4	56.7±1.6
Z32554393	>100	>100	94.6±1.6
AK-778/43465022	**15.5±0.95**	**18.7±1.3**	**21.3±0.8**
AO-080/42479467	88.5±0.7	91.4±1.3	90.4±3.4

AK-968/40390012	>100	>100	>100
AN-329/42613617	56.7±4.8	53.2±1.7	61.3±0.8
AO-080/43441933	47.4±1.0	41.5±2.6	52.5±4.1
C481-0437	66.3±1.5	69.8±2.6	60.6±3.7
6835-3516	>100	>100	>100
F711-0605	37.8±2.7	41.5±2.2	50.1±1.5
Z384684494	67.3±5.0	70.3±0.6	74.2±1.1
AK-778/43420895	**27.3±3.7**	**34.2±2.5**	**38.2±5.4**
AQ-149/43243694	54.3±1.7	60.8±0.9	57.3±3.1

2 **Fig. 2**. Characterization of compound AK-778/43420895 (AK-778-XXMU) *in vitro*. A:

3 AK-778/43420895 had better water solubility as compared to compound AK-778/43465022. B and C:

4 Compound AK-778/43320895 reduces the viability of glioma cells but not normal cells. The effect of

5 compound AK-778/43320895 on cell viability of Human astroblastoma (U87), mouse glioma (GL261),

human normal hepatocytes (L-02) and mouse embryo fibroblasts (3T3) after being exposed to a range of

AK-778/43320895 in a concentration gradient for 24 h via CCK8 assay. The percent of cell viability is

plotted in a dose-dependent manner. Data points represent percent of cell viability relative to DMSO

control of triplicates and expressed as the mean ± SD.

3. Determination of the binding affinity between AK-778-XXMU and ID2 protein

Biacore is a new biometric sensing technology based on SPR with advantages of

high sensitivity, real-time, fast and label-free, which has been widely used to describe

interaction and measure affinity between two partners [41,42]. To identify the

quantitative affinity between ID2 protein and compound AK-778-XXMU, SPR assay

was performed using a Biacore 8K system (GE Healthcare). ID2 protein was

immobilized on the sensor surface as the ligand which was capable of capturing the

target compound AK-778-XXMU in solution. By following the kinetics of a binding

process of ID2 protein and AK-778-XXMU in real time, the association/disassociation

rate constants (k_a and k_d) and equilibrium disassociation constants (K_D) of interactions

were obtained. To our delight, the results showed the K_D between compound

AK-778-XXMU and ID2 protein was 0.129 μM (**Fig. 3**). The submicromolar K_D

demonstrated that compound AK-778-XXMU was able to bind to ID2 protein with

high affinity to exert its inhibitory effects.

Fig. 3. Change in response as a function of time in SPR and the fitting curves of affinity between

compound AK-778-XXMU and ID2 at various concentrations. Using 1:1 binding model, ka was

determined to be 2.48E+01 (1/Ms), kd was determined to be 3.19E-06 (1/s), and KD was determined to

be 1.29E-07 (M). Rmax was 83.6 (RU). Chi^2 was 0.23 (RU^2).

4. Binding modes of AGX51 and AK-778-XXMU with ID2

Compound AK-778-XXMU was identified from the pharmacophore-based virtual

screening described above. The validity of the screening was firmly established by the

SPR assay that determined the potent binding of AK-778-XXMU to ID2 with high

affinity in the low submicromolar range. The molecular superimposition of AGX51

and AK-778-XXMU as well as their pharmacophores were shown to be nearly perfect,

as they should be (**Fig. 4A**). Molecular docking studies were performed to provide

more insight into the binding modes of both compounds in the ID2 binding site (**Fig.**

4B). It was shown that both AGX51 and AK-778-XXMU were fit quite well into the

hydrophobic crevice in the loop region of ID2 (PDB: 4AYA) [36]. The binding modes

of both AGX51 and AK-778-XXMU with ID2 were similar to that of AGX51 with ID1

[35]. As shown in **Fig. 4C** and **Fig. 4D**, in addition to the apparent hydrophobic

interactions, a hydrogen bond was observed between the benzodioxole moiety of

AGX51 and Gln55. The corresponding hydrogen bond was between the hydroxyl

group of AK-778-XXMU and Gln55. Two more hydrogen bonds for AK-778-XXMU

were formed between the ketone group and Lys47 as well as between the amide group

and Lys58. Pi-cation interactions for both compounds were observed between the two

phenyl groups in their structures and Lys47/Lys58. Such interactions offered possible

explanations at the atomic level that why AK-778-XXMU was a potent ID2 antagonist.

Fig. 4. Comparison between AGX51 and the hit compound AK-778-XXMU. (A) molecular

superimposition between the S enantiomer of AGX51 (purple) and AK-778-XXMU (grey). Yellow

spheres represent hydrophobic centers. (B) AGX51 (orange) and AK-778-XXMU (green) in the

binding site of ID2 (C) the 2D diagram of interactions between AGX51 and the binding site of ID2. (D)

the 2D diagram of interactions between AK-778-XXMU and the binding site of ID2.

5. Effect of compound AK-778-XXMU on cell migration and invasion

To determine whether compound AK-778-XXMU can influence the migration and

invasion of glioma cells, wound healing assay was carried out. The results are shown in

Fig. 5. As expected, compound AK-778-XXMU could slow down glioma cells wound

healing. The wound closure rates were depressed significantly as low as $(13.11\pm1.52)\%$

for GL261 and $(5.59\pm1.04)\%$ for U87 after exposed to AK-778-XXMU at the

concentration of 12.5 µM. By comparison, closure rates in control group were up to

$(63.13\pm0.78)\%$ for GL261 and $(61.50\pm10.21)\%$ for U87 cells. To further verify the

effect, the Transwell assay was carried out. As shown in **Fig. 6,** compound

AK-778-XXMU could inhibit the migration and invasion of GL261 and U87 cells in a

concentration-dependent manner.

 Fig. 5. Wound healing assay. Compound AK-778-XXMU suppresses glioma cell migrate. After

 creating a straight physical "wound", GL261 and U87 were treated with different concentrations (0, 2.5,

 12.5μM) of compound AK-778-XXMU for 24 h. The process of cells migration into the gap were

 monitored by taking typical photos at different time points (0, 12 and 24 h). Data points represent the

 mean ± SD of three independent experiments.

 Fig. 6. Transwell assay. A: representative images of migrated cells treated with different concentration

 of compound AK-778-XXMU and the quantitative analysis of migrated cells. B: representative images

 of invaded cells treated at different concentration of compound AK-778-XXMU and the quantitative

 analysis of invaded cells. X axis represents different concentrations of compound AK-778-XXMU

1 treatment, and Y axis is the average number of migrated cells or invaded cells in three independent

2 experiments at least. Each treatment group was compared with the control group. ($*p<0.05$; $**p<0.01$;

3 $***p <0.001$; $****p <0.0001$ vs. Control).

6. Compound AK-778-XXMU induces apoptosis in glioma cells

To explore the effect of AK-778-XXMU on induction of apoptosis in glioma cells,

Annexin V/PI double stained with flow cytometry was performed *in vitro*. Cells were

treated with 25 µM and 50 µM AK-778-XXMU for 24 h. It was shown in **Fig. 7** that

exposures of U87 and GL261 tumor cells to different concentrations of

AK-778-XXMU resulted in increased apoptosis.

Fig. 7. Compound AK-778-XXMU induces apoptosis in glioma cells. Human astroblastoma (U87)

and mouse glioma (GL261) were treated at different concentrations (0, 25, 50 µM) of compound

AK-778-XXMU for 24 h. Significantly increased apoptosis percentage of GL261 (A) and U87 (B) were

1 detected by Annexin V/PI double stained by flow cytometry. Data represents the mean ± SD of at least

2 three independent experiments. (***p<0.001 vs. Control; ****p<0.0001 vs. Control).

7. Effect of compound AK-778-XXMU on ID2-KDR signaling

Glioma has abundant neovascularization network to support its invasive growth.

KDR, also named VEGFR2, as an endotheliocyte-specific tyrosine kinase receptor, is

critical in VEGF signaling and proangiogenic response in glioma malignant

progression [43,44]. It has been reported that ID2 plays a critical role in KDR activation

[17]. Therefore, western blot was performed to determine whether ID2-KDR signaling

could be impaired by compound AK-778-XXMU. As expected, compared with the

control group (0 μM), three proteins involved in ID2-KDR signaling (ID2, VEGFA,

KDR) were greatly decreased in a dose-dependent manner (**Fig. 8**), and this expression

pattern was validated by qRT-PCR (**Fig. 9**). Moreover, with increasing concentrations

of compound AK-778-XXMU, an accumulation of tumor suppressor gene *PTEN* and

an impaired expression of MMP2，MMP9 were found (**Fig. 8**).

Fig. 8. Compound AK-778-XXMU inhibits ID2-KDR signaling using Western blotting. Human

glioma cells U87 and HS683 were treated at different concentrations of compound AK-778-XXMU for

24 h. The expression of ID2, VEGFA, PTEN, MMP2, MMP9 and KDR were detected in the whole cell

lysates by Western blot. β-actin was used as the internal reference protein for each sample. The data were

normalized to β-actin expression. (*$p<0.05$; **$p<0.01$; ***$p<0.001$; ****$p<0.0001$ vs. Control).

Fig. 9. qRT-PCR analysis of ID2 and KDR mRNA expression level. The mRNA expression levels of

ID2 and KDR (VEGFR2) in U87 under the same conditions were assessed by qRT-PCR. The data were

normalized to GAPDH expression. (**$p<0.01$; ***$p<0.001$; ****$p <0.0001$ vs. Control)

8. Compound AK-778-XXMU slows down tumor growth *in vivo*

Inspired by the potent effect of AK-778-XXMU in inhibiting glioma cell growth *in*

vitro, next we inoculated our NCG mouse with xenograft U87 cells into the right axilla

to determine the antitumor effect of compound AK-778-XXMU *in vivo*. As shown in

Fig. 10, in contrast to control group, tumors in treatment group (AK-778-XXMU:

2.5mg/kg) had significantly smaller volumes at the endpoint with 49.7% tumor

inhibitory rate. Furthermore, the treatment group showed a reduction of tumor weight

of 57.8% compared with control group at day 12. Notably, there was no significant

reduction between the two groups in body weight. Overall, these data confirmed that

compound AK-778-XXMU could inhibit tumor growth, possibly by impairing ID2

functions in glioma without obvious side-effects.

To provide more insight into the experimental results above that compound

AK-778-XXMU could inhibit the growth of the tumor in NCG mice and further

understand the differences of tumor growth between the two groups, we removed tumor

blocks from the armpit of ten mice after euthanasia of them. As a proliferating cell

associated nuclear antigen, Ki67 antigen was expressed in G1, S, G2 and M phases of

cell cycle, but not in quiescent phase G0 [45]. Its function is closely related with cell

proliferation, which means the higher positive ratio of Ki67, the faster tumor growth. A

large number of neovascularization is a typical feature of malignant tumors and often

an important cause of distant metastasis of tumors. A recent study demonstrated that

glioblastoma stem cells could differentiate into endothelial cells which advance

angiogenesis [46,47]. CD31 antigen, also known as platelet endothelial cell adhesion

molecule-1 (PECAM-1), is a classical marker of endothelial cell that is also strongly

related to the malignant progression of glioma [46,48]. Then immunohistochemical

staining was performed on the interior of the tumor. As shown in **Fig. 11A** and **11B**,

after treated with compound AK-778-XXMU, notable reduced positive rates of Ki67

and CD31 were detected inside the tumors with no destructive side effects on normal

tissues (**Fig. 11C**) by staining the heart, liver, lungs, and kidneys with hematoxylin &

eosin (HE) in treatment group, further demonstrating that compound AK-778-XXMU

had a potent inhibitory effect on glioma.

Fig. 10. Inhibitory effects of compound AK-778-XXMU in U87 glioma in vivo. NCG mice bearing U87

tumors were injected with DMSO for control group or compound AK-778-XXMU for treatment group

(2.5mg/kg every other day for 5 days, n=5 for each group). A: Tumor volume growth curve after

compound AK-778-XXMU treatment. B: Body weight change of the mice. C and D: the excised tumor

and the tumor weight graph at the end of experiment. (**$p< 0.01$ vs Control).

Fig. 11. Immunohistological anlaysis. Representative immunohistochemical staining photographs of

endothelial marker CD31 (A) for tumors vessels and proliferation marker Ki67 (left B) for cell

proliferation in U87 tumors from NCG tumor-bearing mice at endpoint. (C) Histological analysis of

different tissues from tumor-bearing NCG mice at endpoint. Different tissues (heart, liver, lung, kidney)

from U87 tumor-bearing mice were fixed with 10 % (v/v) formalin and stained using Hematoxylin and

Eosin (H&E). No histological and morphological heterogeneity of these viscera from tumor-bearing

NCG mice in two groups were detected.

<div align="center">**Conclusions**</div>

Given the poor prognosis of glioma patients, it has become imperative to search for

new targeted therapeutic regimen to control the malignant progression of glioma and

improve the life quality of patients. In view of the strong correlation between ID2

protein and glioma, a pharmacophore-based virtual screening was conducted to obtain

16 small-molecule compounds, of which 13 had IC_{50} values lower than 100 μM from

CCK8 assays. Further studies were focused on compound AK-778-XXMU, which is an

indole derivative. The affinity between AK-778-XXMU and ID2 was determined to be

129 nM by SPR experiment, which is in the low submicromolar range. Its binding

mode was studied by comparison with AGX51 in the binding site of ID2 protein,

providing the explanation of its potency at the atomic level. It was also shown that

AK-778-XXMU could inhibit cell migration and invasion of glioma cell lines, induce

apoptosis, and more importantly, slow down the tumor growth *in vivo*. Therefore,

compound AK-778-XXMU could be a promising agent for the treatment of glioma

with ID2 overexpression in the future.

14   **Material and methods**

15   1. Cell lines and cell culture

Cell lines (Human astroblastoma, U87; human glioma, HS683; mouse glioma,

GL261) used in this study were obtained from ATCC. All these cell lines were cultured

in Dulbecco's Modified Eagle's medium (DMEM High Glucose; Hyclone, USA)

containing 10% fetal bovine serum (FBS) (Gibco, New York, USA), 1% (v/v)

antibiotic solution containing penicillin and streptomycin (Sigma-Aldrich, USA) in a

5% CO_2 atmosphere at 37 °C.

22   2. Molecular modeling and virtual screening

23   Compound AGX51 was used to build the pharmacophores. The 3D model of AGX51

24   was built and fully optimized with Discovery Studio 3.5 (BIOVIA, Dassault Systèmes).

Since AGX51 has one chiral center, both enantiomers were used and two pharmacophores were generated by LigandScout 4.0 software by Inte:Ligand GmbH (http://www.inteligand.com/ligandscout/) [40]. They were subjected to commercial the following databases: CHEMBRIDGE, SPECS, CHEMDIV, ENAMINE, IBS, LIFECHEMICALS, VITAS-M (2,281 compounds in total) provided in their electronic form by Shanghai-based Topscience Co., Ldt (https://www.tsbiochem.com/) for virtual screening also in LigandScout. The pharmacophore models built from the *R* and *S* configurations yielded 24 and 33 hits, respectively. In the end, 16 compounds were purchased after consideration of possible future chemical modifications. Molecular docking was performed using Glide from Schrödinger, Inc using the default XP protocol [49,50].

3. CCK8 assay

CCK8 assay was used to elucidate the cytotoxicity of inhibitor on glioma cells. The cells were seeded into 96-well plate at 4×10^3 cells per well with 100 μL DMEM supplemented with 10% fetal bovine serum and incubated at 37 °C. When the cells were grown to 60-70% confluency, different concentrations (0, 1.56, 3.125, 6.25, 12.5, 25, 50 μM) of inhibitor were added. After incubation for 24 h, 10 μL CCK-8 was added to each well. The absorbance of each well was detected at 450 nm after incubating at 37 °C for 4 hours. Every experiment was performed in triplicates.

4. SPR

In SPR assay, the surface of the Series S Sensor CM5 chip (GE, BR-1005-30) was activated with 400 mM EDC and 100 mM NHS at a flow rate of 10 μL/min for 420 s.

Then human ID2 protein (Cat#PROTQ02363, purchased from Boster Biological

Technology Co. Ltd, China.) was diluted to 50 μg/mL with fixative reagent (10 mM

sodium acetate, pH 4.5) and immobilized on the surface of Chip CM5 with a fixed

volume of approximately 6000 RU prior to the measurement of the compound

AK-778-XXMU interaction. For compounds, they were first diluted 20-fold with

dilution buffer (1 × PBS, 0.05% Tween20) to change the DMSO content to 5%, and

then diluted with running reagent (1 × PBS, 0.05% Tween20, 5% DMSO) at

concentrations of 100, 50, 25, 12.5, 6.25, 3.125, and 0 μM, respectively. The diluted

compounds were injected into the experimental channel and the reference channel at a

flow rate of 30 μL/min with a binding time of 60 s and a dissociation time of 120 s. The

KD value of each antibody was calculated using Biacore 8K (GE Healthcare) analysis

software. All SPR measurements were performed in triplicates.

5. Apoptosis analysis

Apoptosis of U87 and GL261 cells were detected by Annexin V-FITC/PI apoptosis

detection kit (Key Gen Bio TECH). Glioma cells were treated with different

concentrations (0, 25, 50 μM) of inhibitors for 24 h, harvested and washed twice with

PBS and centrifuged. All cells were then suspended in 500 μL binding buffer, and

stained with 5 μL Annexin V–FITC and 5 μL PI for 20 min at room temperature in the

dark. Samples were run on a flow cytometer within 1 h.

6. Wound healing assay

In the migration assay, glioma cells were planted into 6-well plate at the density of

5 × 105 cells per well with 2 mL high-glucose DMEM containing 10% FBS, incubated

at 37 °C in a 5% CO_2 incubator for 24 h. Then a straight physical "wound" within cell

monolayer was created. The cells were washed twice with PBS buffer gently and the

cells that have been scratched out were discarded. 2 mL high glucose medium (DMEM)

containing different concentrations of compound (0 μM, 2.5 μM, 12.5 μM) was added

to each well. Then continue to culture in 37 °C incubator. Monitor the process of cell

migration into the gap by taking photos at different time points (0 h, 12 h, 24 h).

7. Migration assay

In the migration assay, Transwell inserts with an 8 μM pore size were placed in

24-well plates with 500 μL DMEM culture medium added 10% fetal bovine serum. The

U87 and HS683 cells were diluted to 2×10^5 cells/mL with DMEM high glucose

medium with 2% fetal bovine serum. A total of 200 μL of cell suspension was added to

the inserts. Meanwhile different concentrations (0, 12.5, 50 μM) of inhibitors were

added. Then the chambers were cultured in a 5% CO_2 incubator at 37°C for 24 h. The

cells were fixed with 4% paraformaldehyde, stained with crystal violet then take

pictures and count in five different fields with an inverted microscope (200×).

8. Invasion assay

In the invasion assay, 30 μL Matrigel (BD Bioscience) was added to the up chamber

then allowed to polymerize for at least 40 min at 37°C. The U87 and HS683 cells were

resuspended to 2×10^5 cells/mL with DMEM high glucose medium with 2% fetal

bovine serum. A total of 200 μL of cell suspension was added to the inserts. Meanwhile

different concentrations (0, 12.5, 50μM) of inhibitors were added. Then the plates were

cultured in a 5% CO_2 incubator at 37 °C for 24 h. Gently wipe off the Matrigel (Corning,

Cat#356234) in the upper chamber with a cotton swab. Then the cells were fixed with

4% paraformaldehyde, stained with crystal violet (Beyotime) then take pictures and

count in five different fields with an inverted microscope Pierce (200×).

9. Western blotting

Glioma cells were treated with different concentrations (0, 12.5, 25, 50 μM) of

compound for 24 hours before being lysed by RIPA containing protease inhibitor.

Lysates were incubated on ice for 20 min prior to centrifugation at 12000 rpm for 20

min at 4°C. After being determined concentrations using a BCA kit (Pierce), 30 μg total

protein was separated by SDS-PAGE then transferred to the polyvinylidene fluoride

membrane. The primary antibodies used were as follows: anti-KDR (Cell Signaling

Technology, #2479), anti-PTEN (Cell Signaling Technology, #9188), anti-MMP-9

(Cell Signaling Technology, #13667), anti-MMP-2 (Cell Signaling Technology,

#87809), anti-VEGFA (Cell Signaling Technology, #65373), anti-ID2 (Cell Signaling

Technology, #3431), anti-β-actin (Cell Signaling Technology, #4970). The dilutions

for the antibodies were as follows: 1:1000 for KDR, 1:1000 for PTEN, 1:1000 for

MMP-9, 1:1000 for MMP-2, 1:1000 for VEGFA, 1:1000 for ID2, 1:1000 for β-actin,

1:10000 for Goat anti-Rabbit IgG.

10. Real-time PCR

To quantify the genes involved ID2-KDR signaling expression, human glioma cells

HS683 were exposed to different concentrations (0, 12.5, 25, 50μM) of the compound,

after 24 h, cells were lysed by RNAiso Plus (Takara, 9108, Japan). Before performing

RT-PCR, Prime Script RT reagent Kit (Takara, RR047A, Japan) were used to reverse

transcribe the extracted RNA into cDNA according to the given instructions. The

results were assessed by calculating $2^{-\Delta\Delta Ct}$. The sequences of primers are as follows:

GAPDH forward primer: 5′-GGAGCGAGATCCCTCCAAAAT-3′, reverse primer:

5′-GGCTGTTGTCATACTTCTCATGG-3′; ID2 forward primer:

5′-AGTCCCGTGAGGTCCGTTAG-3′, reverse primer:

5′-AGTCGTTCATGTTGTATAGCAGG-3′; VEGFR2 forward primer:

5′-GTGATCGGAAATGACACTGGAG-3′, reverse primer:

5′-CATGTTGGTCACTAACAGAAGCA-3′.

11. Subcutaneous tumorigenesis in NCG mice

To generate mouse models, ten 8-week-old NCG mice were injected i.h. with

human glioma cell U87 into the right axilla according to the number of 1×10^6

cells/mouse. When the tumor volume was visible and reached 100 mm^3, ten mice were

randomly divided into two cohorts: control group (n=5) and treatment group (n=5). The

control group (DMSO, 2.5 ml/kg), AK-778-XXMU (2.5mg/kg) compound was

injected i.p. into the two groups tumor-bearing mice respectively. Control group

received DMSO (1% v/v) on the 1st, 3rd and 5th day. Treatment group were injected

with AK-778-XXMU compound (1% v/v, dissolved in DMSO) on the 1st, 3rd and 5th

day. The body weight and tumor size of mice were measured at the same time of each

administration, and the changes of body weight and activity ability of mice were

dynamically monitored. On the endpoint, the 7th day of the first administration, the

mice were euthanized, the tumor were excised and weighed.

12. IHC (Immunohistochemistry) assay

The expression of micro vessel density marker CD31 and cell proliferation index Ki67 in tumor samples from tumor-bearing NCG mice were tested via IHC. IHC staining was performed as previously described [51]. Each slice was taken five views for photographing at 400 X magnification for Ki67 and at 200 X magnification for CD31. Each obvious area of positive staining for CD31 was counted as a single vessel. Results were expressed as mean number of Ki67 positive cells ± SEM per high-power field (X400) or the mean number of micro vessels for CD31 ± SEM per high-power field (X200). A total of 15 high-power fields was examined and the number of micro vessels for CD31 or Ki67 positive-staining cells in each group was averaged and expressed as the number per high power field.

13. Hematoxylin & eosin (HE) staining

After being euthanized, different tissues (heart, liver, lung, kidney) from ten tumor-bearing NCG mice were fixed with 10% (v/v) formalin over 24 hours. Then all the specimens were washed, dehydrated and embedded with paraffin commonly. 5μm-thick viscera sections were cut randomly, fixed onto the glass slides then, stained using Hematoxylin and Eosin (H&E) (Solarbio,G1121) after being dewaxed to water, the images were taken by light microscopy with Nikon software (Melville, NY, USA) at 400 × magnification at last.

14. Statistical analysis

All measurements were performed in triplicates. The quantitative data were performed with the SPSS 20.0 software package expressed as the means ± standard

deviation (SD). The significance of the difference from the controls for each

experimental test condition was assayed using unpaired t-test. ANOVAs with the post

hoc Dunnett's test were used for tumor growth in comparison to the control. *P* values <

0.05 were considered statistically significant.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This work was supported by National Natural Science Foundation of China (No.

21406143) to Bin Lin, NSFC-Henan Union grant (No. U1904131) to Genshen Zhong.

Bin Lin also would like to acknowledge the faculty startup funding from Shenyang

Pharmaceutical University.

References:

[1]   E.G. Van Meir, C.G. Hadjipanayis, A.D. Norden, H.K. Shu, P.Y. Wen, J.J.

18   Olson, Exciting new advances in neuro-oncology: the avenue to a cure for

19   malignant glioma, CA. Cancer J. Clin. 60 (2010) 166–193.

20   https://doi.org/10.3322/caac.20069.

1   [2]   A. Omuro, L.M. DeAngelis, Glioblastoma and other malignant gliomas: a

2   clinical review, JAMA. 310 (2013) 1842–1850.

3   https://doi.org/10.1001/jama.2013.280319.

4   [3]   T.A. Dolecek, J.M. Propp, N.E. Stroup, C. Kruchko, CBTRUS statistical report:

5   primary brain and central nervous system tumors diagnosed in the United States

6   in 2005-2009, Neuro-Oncol. 14 (2012) v1-49.

7   https://doi.org/10.1093/neuonc/nos218.

8   [4]   P.Y. Wen, E.Q. Lee, D.A. Reardon, K.L. Ligon, W.K.A. Yung, Current clinical

9   development of PI3K pathway inhibitors in glioblastoma, Neuro-Oncol. 14

10   (2012) 819–829. https://doi.org/10.1093/neuonc/nos117.

11   [5]   H. Malkki, Neuro-oncology: bevacizumab prolongs progression-free survival

12   but not overall survival in newly diagnosed glioblastoma, Nat. Rev. Neurol. 10

13   (2014) 179. https://doi.org/10.1038/nrneurol.2014.47.

14   [6]   S. Bao, Q. Wu, R.E. McLendon, Y. Hao, Q. Shi, A.B. Hjelmeland, M.W.

15   Dewhirst, D.D. Bigner, J.N. Rich, Glioma stem cells promote radioresistance by

16   preferential activation of the DNA damage response, Nature 444 (2006)

17   756–760. https://doi.org/10.1038/nature05236.

18   [7]   I. Paw, R.C. Carpenter, K. Watabe, W. Debinski, H.W. Lo, Mechanisms

19   regulating glioma invasion, Cancer Lett. 362 (2015) 1–7.

20   https://doi.org/10.1016/j.canlet.2015.03.015.

[8]   L.C. Hou, A. Veeravagu, A.R. Hsu, V.C. Tse, Recurrent glioblastoma

multiforme: a review of natural history and management options, Neurosurg.

Focus. 20 (2006) E5. https://doi.org/10.3171/foc.2006.20.4.2.

[9]   R. Benezra, R.L. Davis, D. Lockshon, D.L. Turner, H. Weintraub, The protein Id:

a negative regulator of helix-loop-helix DNA binding proteins, Cell. 61 (1990)

49–59. https://doi.org/10.1016/0092-8674(90)90214-y.

[10]   H.A. Sikder, M.K. Devlin, S. Dunlap, B. Ryu, R.M. Alani, Id proteins in cell

growth and tumorigenesis, Cancer Cell. 3 (2003) 525–530.

https://doi.org/10.1016/s1535-6108(03)00141-7.

[11]   X.H. Sun, N.G. Copeland, N.A. Jenkins, D. Baltimore, Id proteins Id1 and Id2

selectively inhibit DNA binding by one class of helix-loop-helix proteins, Mol.

Cell. Biol. 11 (1991) 5603–5611. https://doi.org/10.1128/mcb.11.11.5603.

[12]   J. Biggs, E. V Murphy, M.A. Israel, A human Id-like helix-loop-helix protein

expressed during early development, Proc. Natl. Acad. Sci. 89 (1992)

1512–1516. https://doi.org/10.1073/pnas.89.4.1512.

[13]   V. Riechmann, I. van Crüchten, F. Sablitzky, The expression pattern of Id4, a

novel dominant negative helix-loop-helix protein, is distinct from Id1, Id2 and

Id3, Nucleic Acids Res. 22 (1994) 749–755.

https://doi.org/10.1093/nar/22.5.749.

[14]   B.A. Christy, L.K. Sanders, L.F. Lau, N.G. Copeland, N.A. Jenkins, D. Nathans,

An Id-related helix-loop-helix protein encoded by a growth factor-inducible

gene, Proc. Natl. Acad. Sci. 88 (1991) 1815–1819.

https://doi.org/10.1073/pnas.88.5.1815.

[15] J. Perk, A. Iavarone, R. Benezra, Id family of helix-loop-helix proteins in cancer,

Nat. Rev. Cancer. 5 (2005) 603–614. https://doi.org/10.1038/nrc1673.

[16] Z. Dong, S. Liu, C. Zhou, T. sumida, H. Hamakawa, Z. Chen, P. Liu, F. Wei,

Overexpression of Id-1 is associated with tumor angiogenesis and poor clinical

outcome in oral squamous cell carcinoma, Oral Oncol. 46 (2010) 154–157.

https://doi.org/10.1016/j.oraloncology.2009.11.005.

[17] Y. Huang, P. Rajappa, W. Hu, C. Hoffman, B. Cisse, J.H. Kim, E. Gorge, R.

Yanowitch, W. Cope, E. Vartanian, R. Xu, T. Zhang, D. Pisapia, J. Xiang, J.

Huse, I. Matei, H. Peinado, J. Bromberg, E. Holland, B.S. Ding, S. Rafii, D.

Lyden, J. Greenfield, A proangiogenic signaling axis in myeloid cells promotes

malignant progression of glioma, J. Clin. Invest. 127 (2017) 1826–1838.

https://doi.org/10.1172/JCI86443.

[18] D.A. Vandeputte, D. Troost, S. Leenstra, H. Ijlst-Keizers, M. Ramkema, D.A.

Bosch, F. Baas, N.K. Das, E. Aronica, Expression and distribution of id

helix-loop-helix proteins in human astrocytic tumors, Glia. 38 (2002) 329–338.

https://doi.org/10.1002/glia.10076.

[19] Z. Zhang, G.J. Rahme, P.D. Chatterjee, M.C. Havrda, M.A. Israel, ID2 promotes

survival of glioblastoma cells during metabolic stress by regulating

mitochondrial function, Cell Death Dis. 8 (2017) e2615.

https://doi.org/10.1038/cddis.2017.14.

[20] A.J. Levine, A.M. Puzio-Kuter, The control of the metabolic switch in cancers

by oncogenes and tumor suppressor genes, Science 330 (2010) 1340–1344.

https://doi.org/10.1126/science.1193494.

[21] C. Stockmann, A. Doedens, A. Weidemann, N. Zhang, N. Takeda, J.I.

Greenberg, D.A. Cheresh, R.S. Johnson, Deletion of vascular endothelial growth

factor in myeloid cells accelerates tumorigenesis, Nature 456 (2008) 814–818.

https://doi.org/10.1038/nature07445.

[22] R. Chen, M.C. Nishimura, S.M. Bumbaca, S. Kharbanda, W.F. Forrest, I.M.

Kasman, J.M. Greve, R.H. Soriano, L.L. Gilmour, C.S. Rivers, Z. Modrusan, S.

Nacu, S. Guerrero, K.A. Edgar, J.J. Wallin, K. Lamszus, M. Westphal, S. Heim,

C.D. James, S.R. VandenBerg, J.F. Costello, S. Moorefield, C.J. Cowdrey, M.

Prados, H.S. Phillips, A hierarchy of self-renewing tumor-initiating cell types in

glioblastoma, Cancer Cell 17 (2010) 362–375.

https://doi.org/10.1016/j.ccr.2009.12.049.

[23] S.M. Pollard, K. Yoshikawa, I.D. Clarke, D. Danovi, S. Stricker, R. Russell, J.

Bayani, R. Head, M. Lee, M. Bernstein, J.A. Squire, A. Smith, P. Dirks, Glioma

stem cell lines expanded in adherent culture have tumor-specific phenotypes and

are suitable for chemical and genetic screens, Cell Stem Cell 4 (2009) 568–580.

https://doi.org/10.1016/j.stem.2009.03.014.

[24] C. Roschger, C. Cabrele, The Id-protein family in developmental and

cancer-associated pathways, Cell Commun Signal. 15 (2017) 7.

https://doi.org/10.1186/s12964-016-0161-y.

[25]   D. Lyden, A.Z. Young, D. Zagzag, W. Yan, W. Gerald, R. O'Reilly, B.L. Bader,

R.O. Hynes, Y. Zhuang, K. Manova, R. Benezra, Id1 and Id3 are required for

neurogenesis, angiogenesis and vascularization of tumour xenografts, Nature

401 (1999) 670–677. https://doi.org/10.1038/44334.

[26]   M.B. Ruzinova, R.A. Schoer, W. Gerald, J.E. Egan, P.P. Pandolfi, S. Rafii, K.

Manova, V. Mittal, R. Benezra, Effect of angiogenesis inhibition by Id loss and

the contribution of bone-marrow-derived endothelial cells in spontaneous

murine tumors, Cancer Cell 4 (2003) 277–289.

https://doi.org/10.1016/s1535-6108(03)00240-x.

[27]   A. Lasorella, R. Benezra, A. Iavarone, The ID proteins: master regulators of

cancer stem cells and tumour aggressiveness, Nat. Rev. Cancer 14 (2014) 77–91.

https://doi.org/10.1038/nrc3638.

[28]   D. Gao, D.J. Nolan, A.S. Mellick, K. Bambino, K. McDonnell, V. Mittal,

Endothelial progenitor cells control the angiogenic switch in mouse lung

metastasis, Science 319 (2008) 195–198.

https://doi.org/10.1126/science.1150224.

[29]   J. Anido, A. Sáez-Borderías, A. Gonzàlez-Juncà, L. Rodón, G. Folch, M.A.

Carmona, R.M. Prieto-Sánchez, I. Barba, E. Martínez-Sáez, L. Prudkin, I.

Cuartas, C. Raventós, F. Martínez-Ricarte, M.A. Poca, D. García-Dorado, M.M.

Lahn, J.M. Yingling, J. Rodón, J. Sahuquillo, J. Baselga, J. Seoane, TGF-β

receptor inhibitors target the CD44[high]/Id1[high] glioma-initiating cell population in

human glioblastoma, Cancer Cell 18 (2010) 655–668.

https://doi.org/10.1016/j.ccr.2010.10.023.

[30] E. Henke, J. Perk, J. Vider, P. de Candia, Y. Chin, D.B. Solit, V. Ponomarev, L.

Cartegni, K. Manova, N. Rosen, R. Benezra, Peptide-conjugated antisense

oligonucleotides for targeted inhibition of a transcriptional regulator *in vivo*, Nat.

Biotechnol. 26 (2008) 91–100. https://doi.org/10.1038/nbt1366.

[31] D.S. Mern, J. Hasskarl, B. Burwinkel, Inhibition of Id proteins by a peptide

aptamer induces cell-cycle arrest and apoptosis in ovarian cancer cells, Br. J.

Cancer. 103 (2010) 1237–1244. https://doi.org/10.1038/sj.bjc.6605897.

[32] H. Mistry, G. Hsieh, S.J. Buhrlage, M. Huang, E. Park, G.D. Cuny, I. Galinsky,

R.M. Stone, N.S. Gray, A.D. D'Andrea, K. Parmar, Small-molecule inhibitors of

USP1 target ID1 degradation in leukemic cells, Mol. Cancer Ther. 12 (2013)

2651–2662. https://doi.org/10.1158/1535-7163.Mct-13-0103-t.

[33] R. Murase, R. Kawamura, E. Singer, A. Pakdel, P. Sarma, J. Judkins, E.

Elwakeel, S. Dayal, E. Martinez-Martinez, M. Amere, R. Gujjar, A. Mahadevan,

P.-Y. Desprez, S.D. McAllister, Targeting multiple cannabinoid anti-tumour

pathways with a resorcinol derivative leads to inhibition of advanced stages of

breast cancer, Br. J. Pharmacol. 171 (2014) 4464–4477.

https://doi.org/10.1111/bph.12803.

[34] L. Soroceanu, R. Murase, C. Limbad, E. Singer, J. Allison, I. Adrados, R.

Kawamura, A. Pakdel, Y. Fukuyo, D. Nguyen, S. Khan, R. Arauz, G.L. Yount,

D.H. Moore, P.-Y. Desprez, S.D. McAllister, Id-1 is a key transcriptional

regulator of glioblastoma aggressiveness and a vovel therapeutic target, Cancer

Res. 73 (2013) 1559–1569. https://doi.org/10.1158/0008-5472.Can-12-1943.

[35] P.M. Wojnarowicz, R. Lima e Silva, M. Ohnaka, S.B. Lee, Y. Chin, A. Kulukian,

S.-H. Chang, B. Desai, M. Garcia Escolano, R. Shah, M. Garcia-Cao, S. Xu, R.

Kadam, Y. Goldgur, M.A. Miller, O. Ouerfelli, G. Yang, T. Arakawa, S.K.

Albanese, W.A. Garland, G. Stoller, J. Chaudhary, L. Norton, R.K. Soni, J.

Philip, R.C. Hendrickson, A. Iavarone, A.J. Dannenberg, J.D. Chodera, N.

Pavletich, A. Lasorella, P.A. Campochiaro, R. Benezra, A small-molecule

pan-Id antagonist inhibits pathologic ocular neovascularization, Cell Rep. 29

(2019) 62-75. https://doi.org/10.1016/j.celrep.2019.08.073.

[36] M.V. Wong, S. Jiang, P. Palasingam, P.R. Kolatkar, A divalent ion is crucial in

the structure and dominant-negative function of ID proteins, a class of

helix-loop-helix transcription regulators, PLoS One. 7 (2012) e48591.

https://doi.org/10.1371/journal.pone.0048591.

[37] S.-Y. Yang, Pharmacophore modeling and applications in drug discovery:

challenges and recent advances, Drug Discov. Today. 15 (2010) 444–450.

https://doi.org/10.1016/j.drudis.2010.03.013.

[38] G. Wolber, A.A. Dornhofer, T. Langer, Efficient overlay of small organic

molecules using 3D pharmacophores, J. Comput. Aided. Mol. Des. 20 (2006)

773–788. https://doi.org/10.1007/s10822-006-9078-7.

[39]   A.R. Leach, V.J. Gillet, R.A. Lewis, R. Taylor, Three-dimensional

2   pharmacophore methods in drug discovery, J. Med. Chem. 53 (2010) 539–558.

3   https://doi.org/10.1021/jm900817u.

4   [40]   G. Wolber, T. Langer, LigandScout: 3-D pharmacophores derived from

5   protein-bound ligands and their use as virtual screening filters, J. Chem. Inf.

6   Model. 45 (2005) 160–169. https://doi.org/10.1021/ci049885e.

7   [41]   C. Boozer, G. Kim, S. Cong, H. Guan, T. Londergan, Looking towards label-free

8   biomolecular interaction analysis in a high-throughput format: a review of new

9   surface plasmon resonance technologies, Curr. Opin. Biotechnol. 17 (2006)

400–405. https://doi.org/10.1016/j.copbio.2006.06.012.

[42]   M.A. Cooper, Optical biosensors in drug discovery, Nat. Rev. Drug Discov. 1

(2002) 515–528. https://doi.org/10.1038/nrd838.

[43]   A.K. Olsson, A. Dimberg, J. Kreuger, L. Claesson-Welsh, VEGF receptor

signalling — in control of vascular function, Nat. Rev. Mol. Cell Biol. 7 (2006)

359–371. https://doi.org/10.1038/nrm1911.

[44]   E. Krzywinska, C. Kantari-Mimoun, Y. Kerdiles, M. Sobecki, T. Isagawa, D.

Gotthardt, M. Castells, J. Haubold, C. Millien, T. Viel, B. Tavitian, N. Takeda, J.

Fandrey, E. Vivier, V. Sexl, C. Stockmann, Loss of HIF-1α in natural killer cells

inhibits tumour growth by stimulating non-productive angiogenesis, Nat.

Commun. 8 (2017) 1597. https://doi.org/10.1038/s41467-017-01599-w.

[45] M.J. Beresford, G.D. Wilson, A. Makris, Measuring proliferation in breast

cancer: practicalities and applications, Breast Cancer Res. 8 (2006) 216.

https://doi.org/10.1186/bcr1618.

[46] X. Mei, Y.S. Chen, F.R. Chen, S.Y. Xi, Z.P. Chen, Glioblastoma stem cell

differentiation into endothelial cells evidenced through live-cell imaging,

Neuro-Oncol. 19 (2017) 1109–1118. https://doi.org/10.1093/neuonc/nox016.

[47] D. Chroscinski, D. Sampey, N. Maherali, Reproducibility Project: Cancer

Biology, Registered report: tumour vascularization via endothelial

differentiation of glioblastoma stem-like cells, eLife. 4 (2015) e04363.

https://doi.org/10.7554/eLife.04363.

[48] S. Das, P.A. Marsden, Angiogenesis in glioblastoma, N. Engl. J. Med. 369 (2013)

1561–1563. https://doi.org/10.1056/NEJMcibr1309402.

[49] R.A. Friesner, J.L. Banks, R.B. Murphy, T.A. Halgren, J.J. Klicic, D.T. Mainz,

M.P. Repasky, E.H. Knoll, M. Shelley, J.K. Perry, D.E. Shaw, P. Francis, P.S.

Shenkin, Glide: A new approach for rapid, accurate docking and scoring. 1.

method and assessment of docking accuracy, J. Med. Chem. 47 (2004)

1739–1749. https://doi.org/10.1021/jm0306430.

[50] T.A. Halgren, R.B. Murphy, R.A. Friesner, H.S. Beard, L.L. Frye, W.T. Pollard,

J.L. Banks, Glide: A new approach for rapid, accurate docking and scoring. 2.

enrichment factors in database screening, J. Med. Chem. 47 (2004) 1750–1759.

https://doi.org/10.1021/jm030644s.

[51]   G. Zhong, S. Zhang, Y. Li, X. Liu, R. Gao, Q. Miao, Y. Zhen, A tandem

2   scFv-based fusion protein and its enediyne-energized analogue show intensified

3   therapeutic efficacy against lung carcinoma xenograft in athymic mice, Cancer

4   Lett. 295 (2010) 124–133. https://doi.org/10.1016/j.canlet.2010.02.020.

> 2,000 compounds

AGX51

Pharmacophore Generation

Pharmacophore-based Virtual Screening

57 hit compounds
16 compounds purchased

CCK8 assay

AK-778-XXMU
KD = 129 nM (SPR)

why so potent?

A overlay of pharmacophores

B AGX51 & AK-778-XXMU in ID2 binding site

C AGX51 interaction mode

D AK-778-XXMU interaction mode

H-Bond —— Carbon Hydrogen Bond ------ Pi-Cation ------ Amide-Pi Stacking ------