Implicit Laplacian of Enhanced Edge: An Unguided Algorithm for Accurate and Automated Quantitative Analysis of Cytoskeletal Images

Pai Li,[1,2,*,#] Ze Zhang[3,*,#], Yiying Tong[3], Bardees M.Foda[4,5], and Brad Day[1]

Affiliations

[1] Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing, Michigan 48824, USA. [2] Department of Plant Biology, Michigan State University, East Lansing, Michigan 48824, USA. [3] Department of Computer Science and Engineering, Michigan State University, East Lansing, MI 48824, USA. [4] Department of Pharmacology and Toxicology, Michigan State University, East Lansing, MI 48824, USA. [5] Molecular Genetics and Enzymology Department, National Research Centre, Dokki, Egypt.

*These two authors contributed equally to this work.
#Corresponding author. Email: lipai@msu.edu and zhangze6@msu.edu

Keywords
Cytoskeleton, cytoskeleton quantitative analysis, unguided image analysis, 3D, implicit Laplacian, density, bundling.

Abstract

The eukaryotic cytoskeleton plays essential roles in cell signaling and trafficking, which is broadly associated with immunity and diseases of human and plants. To date, most analyses aiming at defining the temporal and spatial dynamics of the cytoskeleton have relied on qualitative and quantitative analysis of fluorescence images to describe cytoskeletal function. While state-of-the-art, these approaches have limitations: the diverse shape and brightness of the cytoskeleton cause considerable inaccuracy in both human-driven and automated approaches, and the widely adopted image projection process (3D to 2D) leads to bias and information loss. Here, we describe the development and application of Implicit Laplacian of Enhanced Edge (ILEE), an unguided approach that uses a 2D/3D-compatible local thresholding algorithm for the quantitative evaluation of cytoskeletal status and organization at high performance. Using ILEE, we constructed a Python library to enable automated cytoskeletal image analysis, providing numerous biologically-interpretable indices measuring the density, bundling, severing, branching, and directionality of the cytoskeleton. The data presented herein demonstrate that ILEE resolves the defects of classic cytoskeleton analysis approaches, enables the measurement of novel cytoskeletal features, and yields quantitatively descriptive data with superior accuracy, stability, and robustness. We released the ILEE algorithm as an open-source library and further developed a Google Colab interface as a community resource.

Introduction

Higher eukaryotes have evolved a complex suite of cellular signaling mechanisms to regulate many biological processes, such as growth, development, movement, reproduction, and response to environmental stimuli. Not surprisingly, to maintain the integration and sustainability of a conceptual signaling network, another "physical network" is deployed as the framework to organize all subcellular structures and the conduit to transport molecules and information[1] – namely, the cytoskeleton. Shared by plants and animals are two types of cytoskeleton, actin and microtubule, which can be generalized as a cytoplasmic web-like matrix, physically integrating the plasma membrane, vesicles, and organelles and functionally connecting the intercellular signaling to the extracellular environments through a highly dynamic series of temporally- and spatially-regulated changes of architecture[2,3]. Altogether, the cytoskeleton controls numerous cellular processes such as movement, shaping, cellular trafficking, and intercellular communication[4]. It also provides the mechanical force required for chromosome separation and plasma membrane division during mitosis and meiosis[5]. Besides its role within the cytoplasm, the cytoskeleton also participates in a variety of processes within the nucleus, including RNA polymerase recruitment, transcription initiation, and chromosome scaffolding[6].

As the cytoskeleton is vital for cell activities and life, the dysfunction of cytoskeletal dynamics generally leads to severe disease. For example, in the development of Alzheimer's disease, amyloid development triggers the dysregulation of actin dynamics within dendritic spines, leading to synaptotoxicity[7]. Similarly, the dysfunction of microtubule dynamics can trigger neuropsychiatric disorders[8]. In breast cancer, the cancer cell can deploy aberrant actin aggregation to resist cytotoxic natural killer (NK) cells from the immune system[9]. In the case of plant diseases, the function of the cytoskeleton is similarly required: recent work has demonstrated that the dynamics of actin and microtubule of the host are specifically manipulated by pathogens to paralyze the plant immunity during the infection[10]. Indeed, the eukaryotic cytoskeleton not only serves as an integrated cell signaling and transportation platform, but also take risks as "a focus of disease development" at the molecular level for human and plants. Hence, understanding the architecture and dynamics of the cytoskeleton is broadly associated with life, health, and food security, attracting significant interests across different fields of biology and beyond.

Over the past several decades, confocal microscopy-based methods using fluorescence markers have been developed to monitor changes in cytoskeletal organization[11]. While showing advantages in real-time observation and intuitive visual presentation, these approaches possess critical limitations – namely, they are subject to human interpretation, and as a result, often suffer from bias. As a step to reduce such limitation(s), the emergence of computational, algorithm-based analyses offers a solution to describe the quantitative features of cytoskeletal architecture with reduced human bias. However, while early studies introduced the concept of using generalizable image processing pipelines[12,13] to transfer the task of evaluation away from the user and into a series of computer-based quantitative indices, several key bottlenecks were still prevalent. First, most of the quantitative algorithms that have been described to date are limited to 2D images. As a result, these approaches require users to manually generate z-axis projections from raw data, a process that results in an incredible amount of information loss, especially within the perpendicular portion of the cytoskeleton. Second, many current approaches require users to set thresholds manually to segment cytoskeletal components from the images, a task that results in sampling bias. Lastly, the performance of existing algorithms varies significantly depending on the sample source. This hurdle imposes a considerable disparity in the algorithm performance for plants – which possess a dominance of curvy and fluctuating filaments – compared to the animal cytoskeletal organization, which is generally straight and complanate[14–16]. In fact, while sample source dramatically impacts our ability to evaluate the features of cytoskeletal function across all eukaryotes, the vast majority of current approaches are developed based on cytoskeletal images from animal cells, which indicates potential systemic bias when applied to other types of image samples, such as plants.

Previous work has described the development of a global-thresholding-based pipeline to define and evaluate two key features of cytoskeleton filament organization in living plant cells: cytoskeletal density, defined by *occupancy*, and bundling, defined by statistical *skewness* of fluorescence[17]. While this approach utilizes manual global thresholding (MGT), which can potentially introduce a certain level of user bias, it still outperforms most standardized adaptive/automatic global or local thresholding approaches, such as Otsu[18] and Niblack[19]. More recent advances in MGT-based approaches, such as those by Higaki and colleagues, include additional analysis indices, such as the introduction of coefficient of variation (*CV*) of fluorescence to quantify the degree of filament bundling – an improvement that advances the overall robustness and utility of the original algorithm[20]. However, this analysis pipeline still consumes a considerable amount of time and effort from users for massive sample processing and leaves unaddressed two critical issues of rigor in image processing and analysis: information loss and human bias.

In the current study, we developed implicit Laplacian of enhanced edge (ILEE), a 2D/3D compatible unguided local thresholding algorithm for cytoskeletal segmentation and analysis, which is based on the native brightness, first-order derivative (i.e., gradient), and second-order derivative (i.e., Laplacian) of the cytoskeleton image altogether (see Fig. 1). The study described herein supports ILEE as a superior image quantitative analytic platform that overcomes current limitations such as information loss through dimensional reduction, human bias, and inter-sample instability. As shown, ILEE can accurately recognize cytoskeleton from images with a high variation of filament brightness and thickness, such as live plant samples.

As a key advance in the development of ILEE, we constructed an ILEE-based Python library, namely ILEE_CSK, for the fully automated quantitative analysis of cytoskeleton images. In order to improve ILEE_CSK's capability to explore the features of cytoskeleton, we proposed several novel indices – *linear density*, *diameter_TDT*, *diameter_STD*, *static severing activity*, and *static branching activity* to enable/enhance the measurement of (de-)/polymerization, bundling, severing, and branching dynamics of the cytoskeleton. Together with other classic indices transplanted from other studies, ILEE_CSK totally supports 12 cytoskeletal indices within 5 primary classes: density, bundling, connectivity, branching, and directionality. Our data suggested that ILEE outcompetes other classic algorithms by its superior accuracy, stability, and robustness over the computation of cytoskeleton indices. Furthermore, we provided evidence demonstrating higher fidelity and reliability of 3D-based cytoskeletal computational approaches over traditional 2D-based approaches. In addition, using a series of experiment-based images from pathogen-infected plant cells, we demonstrated that ILEE has an improved sensitivity to

128 distinguish biological differences and owns the potential to reveal novel biological features previously
129 omitted due to insufficient approaches. In sum, this platform not only enables the efficient acquisition and
130 evaluation of key actin filament parameters with high accuracy from both projected 2D and native 3D
131 images, but also enables accessibility to a broader range of cytoskeletal status for biological interpretation.
132
133 The library, ILEE_CSK, is publicly released at GitHub (https://phylars.github.io/ILEE_CSK/). We also
134 developed the ILEE Google Colab pipelines for data processing, visualization, and statistical analysis,
135 which is a convenient and user-friendly interface that requires no programing experience or particular
136 computational device.
137
Results
The ILEE pipeline
Raw images generated by laser scanning confocal microscopy are typically obtained through detecting
in-focus photons by a sensor from each resolution unit on a given focal plane. Since the cytoskeleton is
a 3D structure that permeates throughout the cell, current approaches used to capture the filament
architecture rely on scanning the pixels of each plane along the z-axis at a given depth of step; finally,
these stacks are reconstructed to yield a 3D image. However, due to limited computational biological
resources, most studies have exclusively employed the z-axis-projected 2D image, which results in
substantial information loss and systemic bias in downstream analyses.
147
In our newly developed algorithm, we integrated both 2D and 3D data structures into the same processing
pipeline to ameliorate the aforementioned conflict (Fig. 1a). In short, this pipeline enabled automatic
processing and evaluation of both traditional 2D and native 3D z-stack image analysis. As shown in Fig.
1b, cytoskeleton segmentation using ILEE requires three inputs: an edge-enhanced image, a global
gradient threshold that recognizes the edges of potential cytoskeletal components, and the Laplacian
smoothing coefficient K (described below). With these inputs, a local threshold image is generated via
ILEE, and the pixels/voxels with values above the threshold image at the same coordinates are classified
as cytoskeletal components. The output of this is the generation of the binary image (Fig. 1c). Once
acquired, the binary image is further skeletonized[21] to enable the downstream calculation of numerous
cytoskeleton indices, the sum of which comprises the quantitative features of cytoskeletal dynamics (Fig.
1c). Additionally, because the 2D and 3D modes share a common workflow, all of the calculated
cytoskeleton indices also share the definition for both modes, regardless of the difference in dimensional
spaces. This additional feature enables a horizontal comparison of both modes by the user, which we
assert will significantly contribute to the community by providing massive image datasets for further

examination and comparison through the open-source library. In general, the ultimate goal of this approach, and resultant algorithm, is to construct a pipeline that enables the automated detection of the eukaryotic cytoskeleton from complex biological images in an accurate and unbiased manner.

Image decomposition and processing strategy

One of the central problems of automated cytoskeletal image processing is how to accurately recognize cytoskeletal components – a task that is highly challenging because object pixels (i.e., cytoskeleton components) generally have a high dynamic range of signal intensity within and among individual samples, due to varied bundle thickness, concentration of fluorescent dye, and its binding efficiency. As a framework to further understand this challenge, an image from confocal microscopy is conceptually comprised of three components: (1) the true fluorescence, that which is emitted by the dye molecules within the pixel, (2) the diffraction signal transmitted from neighboring space, and (3) the ground noise generated by the sensor of the imaging system (Fig. 2a). During confocal imaging, the ground noise fluctuates around a constant due to the fixed setting of photon sensors, while the diffraction signal is positively correlated with the local true fluorescence. Therefore, an ideal cytoskeleton segregation algorithm will be an adaptive local thresholding approach that refers to both ground noise and local signal intensity.

Identification of coarse background

To identify the approximate level of the ground noise for downstream fine thresholding, a general strategy is to define a highly representative "feature value" of ground noise, based on which a global threshold can be estimated to segment the "coarse background". Guided by this strategy, we designed an algorithm that calculates a global threshold using the morphological features of the ground noise – namely, non-connected negative element scanning (NNES; Fig. 2b). In brief, NNES scans the total number of non-connected negative elements at different global thresholds, resulting in the identification of the representative value (i.e., the peak) with a maximum non-connected negative element count (Fig. 2b, (i)). The global threshold for the coarse background (Fig. 2b, (iii)) will be determined using a linear model trained by the representative value rendered by NNES and manual global thresholding (MGT), a global threshold determined by researchers experienced in manual cytoskeleton image analysis (Supplemental Fig. 1 and 2). NNES can maintain stability and accuracy over different samples that vary in the brightness distribution, because ground noise is the image component with the lowest value that is subject to a normal distribution and generally does not interfere with the actual fluorescence signal. Another accessible method is to directly use the peak-of-frequency brightness of the image as a representative value to train a model. However, this approach is less accurate because the interval near the theoretical

peak is always turbulent and non-monotone, a limitation potentially due to the pollution of diffracted light

(Supplemental Fig. 1; also see *Discussion*).

Cytoskeleton segmentation by ILEE

The core strategy of ILEE is to accurately identify the edge of all cytoskeletal filament components and

apply an implicit Laplacian smoothing[22] on the selected edge. This leads to the generation of a threshold

image where high gradient areas (i.e., the edge of cytoskeleton filaments) smoothly transit to each other

(Fig. 3a). As illustrated in Supplemental Fig. 3a, the general local threshold trend changes as a function

of the baseline values of the cytoskeleton edges. This is because ILEE selectively filters out high-

frequency noise while preserving salient geometric features of individual actin filaments by leveraging

the spectral characteristics of Laplacian operators (Supplemental Fig. 3b,c). Thus, ILEE can filter the

background and noise regardless of the general local brightness level, demonstrating that its function

does not require an operating kernel which would otherwise restrict the performance when evaluating

filaments of varying thickness. Moreover, the edge of the cytoskeletal component can be smoothed and

elongated using a significant difference filter (SDF; Supplemental Fig. 4) and a Gaussian filter, the sum

of which serves to enhance the continuity of the edge and contributes to the accuracy of edge detection

(Fig. 1b and 3a). For this reason, we name our algorithm as indicated as "Implicit Laplacian of Enhanced

Edge"

For computation, the ILEE algorithm builds a linear system based on Laplacian operators to achieve local

adaptive thresholding for edges of cytoskeletal components (see *Methods* for detail). In brief, we first

used the information from the coarse background to estimate a global threshold of the gradient magnitude

(g_{thres}), and classified pixels/voxels above g_{thres} as boundary elements (I_{edge}) (Fig. 3a, 2.1). Following

the boundary identification, we constructed a $n \times n$ (n is the total number of pixel/voxel in the image)

selection matrix *S*, a sparse diagonal matrix whose diagonal corresponds to the flattened pixel/voxel in

order; the diagonal element was 1 if the pixel/voxel possessed a norm of the gradient above g_{thres}. As

the pixels/voxels belonging to the boundary of the cytoskeleton filament were marked, I_{edge} can be

mathematically defined as shown in Fig. 3a, equations 2.2.1 and 2.2.2, where *L* is the Laplacian matrix

and *K*, or implicit Laplacian smoothing coefficient, is a weight that adjusts the influence of the Laplacian

operator. The local threshold image can therefore be rendered by solving the linear equation shown in

Fig. 3b, 2.2.3.

For a given image input, the performance of ILEE depends on two parameters: g_{thres}, which defines the

edge, and *K*, which determines the weight of detail (i.e., high-frequency components) to be filtered. To

calculate the g_{thres} for an input image, we used brightness values of the area identified as coarse
background by NNES. Since the ground noise is approximately subject to a normal distribution, we
hypothesized a deducible statistical relationship between the image gradient, defined by Scharr
operator[23], and the native brightness of pixels/voxels within the coarse background. Using a normal
random array that simulates the noise with a 2D or 3D data structure, we demonstrate that the distribution
of the background gradient magnitude is also normal-like, and both mean (μ_G) and standard deviation
(σ_G) of the gradients are directly proportional to the standard deviation of their native pixel values (σ_{cbg}),
and we calculate the proportionality coefficient (see Supplemental Fig. 5). For the 3D mode, since the
voxel size on the x- and y-axis is different from that of the z-axis, the proportionality coefficient of μ_G and
σ_G over σ_{cbg} will vary for different ratios of the x-y unit : z unit (see Supplemental Fig. 6). To solve this
problem, we simulated artificial 3D noise images and trained a multi-interval regression model that
accurately ($R^2 > 0.999$) calculates the proportionality coefficient of μ_G and σ_G over σ_{cbg} for different x-y :
z ratios of the voxel. Finally, using this approach and randomly selected actin image samples, we trained
a model, $g_{thres} = \mu_G + k(\sigma_{cbg}) * \sigma_G$, to determine the g_{thres} as ILEE input (Supplemental Fig. 7).
244
To determine the appropriate setting of *K*, we first tested how different *K* values influenced the result of
the local threshold image (I_{thres} of Fig. 3a). As shown in Supplemental Figure 8a, at the optimal g_{thres},
a low value of *K* generated an I_{thres} that was highly consistent with the selected edge; when *K* increases,
the total threshold image shifted towards the average value of the selected edges with increasing loss of
detail. As for the resultant binary image, a lower *K* enables the accurate recognition of thin and faint actin
filament components, yet cannot cover the entire width of thick filaments. Conversely, a high *K* value
covers thick actin filaments with improved accuracy, resulting in a binary image that is less noisy; however,
thin and/or faint filaments tend to be omitted as pseudo-negative pixels (Fig. 3b, Supplemental Fig. 8a).
To overcome this dilemma, we applied a strategy using a lower K_1 and a higher K_2 to compute two
different binary images that focus on thin/faint components and thick components, respectively. Then,
we generated a full outer-join image that contains all cytoskeleton components in these two binary images.
This approach led to improved recognition of cytoskeleton with varying morphologies (see Fig. 3b).
257
As described above, K_1 controls the performance of thin and faint filaments. Since the theoretical
minimum thickness of a distinguishable cytoskeletal component is approximately equal to one pixel/voxel
unit, K_1 can be fixed to a constant to recognize the finest cytoskeletal components from a complex and
heterogeneous set of input samples. Using this approach, we identified an empirical optimal K_1 of 2.5.
However, since different image samples have different distributions of cytoskeleton thickness, K_2, which

controls the performance over thick filaments, must be guided according to the maximum thickness

among all samples in an experiment batch. To ensure that the algorithm described herein is fully unguided,

our strategy was to estimate an appropriate K_2 from an estimated maximum thickness using all samples

from a single batch of experiments, including multiple biological replicates (if applicable). To do this, we

used Niblack thresholding first to generate a coarse binary image (which is sufficiently accurate for the

thickest portion of the filament); from this, we calculated the mean of the top 5% of the Euclidian distance

transformation (DT) values of all positive pixels (see *Methods* for additional information). Next, the top 5%

means of all individual images were averaged to estimate K_2 via a trained model using a sample set with

manual portraited binary ground truth (Fig. 3c and Supplemental Fig. 8b, c, d). Therefore, individual

images from all groups in the same batch of an experiment can be processed by K_2 as estimated using

the model above. In this, the bias of human input was avoided. When processing 3D images, we

additionally provided a parameter that allows using a single K, rather than K_1 and K_2, which balances the

accuracy over thin/faint and thick/bright filaments while saving certain amount of time of computation.

Computational analysis of cytoskeleton indices

Through the image processing pipeline, cytoskeletal indices were automatically calculated from the

binary image generated by ILEE. As a substantial expansion from three previously defined cytoskeletal

indices (e.g., *occupancy*, *skewness*, and *CV*)[17,20], we totally introduced 12 indices (Fig. 1a); particularly,

we focused on 9 of the 12 indices in this study, as they are normalized and ready for biological

interpretation (see Supplementary Fig. 9 for visualized demonstration for each index). These indices fall

within five classes – density, bundling, connectivity, branching, and directionality, which describe the

quantifiable features and functions of cytoskeletal morphology and dynamics. It is worth noting that many

of these indices require a certain level of image post-processing (e.g., oversampling) to further enhance

the accuracy of the output. The detailed mathematical definition of each index is described in *Methods*.

We have defined a novel set of cytoskeletal indices to enable the measurement of certain cytoskeletal

features. For the class "density", we developed *linear density*, a feature that measures filament length

per unit of 2D/3D space. For the class "bundling", we developed two new highly robust indices, referred

to as diameter by total DT (*diameter_TDT*) and diameter by skeleton DT (*diameter_SDT*). Both of them

quantify the physical thickness of filament bundles, in addition to the indirect indices *skewness* and *CV*,

which estimate the relative bundling level based on the statistical distribution of fluorescence intensity.

For the class "connectivity", we defined *static severing activity*, which estimates the severing events within

per length unit of the cytoskeleton. *static severing* activity assumes that a severing event generates two

visible cytoskeletal filaments, which is distinguishable from filament depolymerization. This is an

important consideration in terms of the biological behavior of the cytoskeleton, as it enables the
decoupling of the impact of filament depolymerization and filament severing, key activities facilitated by
the eukaryotic actin depolymerizing factor (ADF) and cofilin family of proteins[24]. For the class "branching",
our algorithm is based on Skan, a recently developed Python library for the graph-theoretical analysis of
the cytoskeleton[25]. To further explore the relationship between filament morphology and the biological
activity of branching, we specifically designed a novel index, *static branching activity*, which we define as
the total number of additional branches emerging from any non-end-point node in per unit length of
filament. In total, this index measures the abundance/frequency of cytoskeletal branching. Finally, our
library supports estimating the level of directional cytoskeletal growth by the index *local anisotropy*, which
measures how local filaments tend to be directional or chaotic. This approach is transplanted from an
ImageJ plug-in FibrilTool[26], but we expanded this algorithm to 3D space (Fig. 1c).

It is noteworthy that we named the two indices describing severing and branching above as "*static*"
severing activity and "*static*" *branching activity*, because, by the most rigorous mathematical criteria, a
dynamic status cannot be directly defined without introducing the time dimension. However, the
introduction of time dimension (using a video sample) generally sacrifices spatial resolution due to the
technical limit of microscopy devices. While it is available to use TIRF (total internal reflection
fluorescence microscopy) or VAEM (variable-angle epifluorescence microscopy) to collect 2D videos[27],
such methods may be limited by the insufficient quantity of filaments in the field of vision and manual
counting of the dynamic events, both of which may introduce bias compared to a larger 3D data structure.
Alternatively, we introduce *static severing activity* and *static branching activity* as indirect indices
measuring the intensity of severing and branching. Here, we assume that the occurrence of
severing/branching events and the net regeneration speed of cytoskeleton has established (or is very
near to) their dynamic equilibrium; at this point, the quantity of pieces/branches of the filament is positively
correlated to the speed of severing and branching. In practice, most experimental schemes set an interval
between the treatment (if any) and imaging, which is long enough to reach the dynamic equilibrium.
Therefore, we expect these two novel indices can provide interpretable and reliable data to estimate the
intensity of severing and branching, especially for confocal images.

ILEE displays high accuracy and stability over actin image samples.
To evaluate the performance of ILEE in terms of its accuracy and compatibility over diverse samples, we
selected a dataset of actin images from Arabidopsis leaves with diverse morphology, and compared ILEE
with numerous traditional global and local thresholding algorithms, including MGT. First, to evaluate the
accuracy of each algorithm in terms of filament segregation, we manually generated the ground truth

binary image from each of the samples using a digital drawing monitor (Fig. 4a, ground truth). Next, we used each ground truth binary image as a reference and compared the filament architecture obtained by ILEE, MGT, and 6 additional adaptive thresholding algorithms. These additional thresholding algorithms include Otsu[18], Triangle[28], Li[29], Yan[30], Niblack[19], and Sauvola[31] (Fig. 4). As an additional concern of rigor, we anticipated that pseudo-positive pixels may be obtained due to operator's bias during the generation of each of the ground truth images (even when the operator is experienced in the actin imaging field). Therefore, we further analyzed each non-connected component of pseudo-positive pixel by its shape and connectivity to the matched elements and identified the actin-like pseudo-positive pixels as possible real actin components.

As shown in Fig. 4a (visualized demonstration), 4b (quantitative analysis), and 4c (bias analysis), ILEE offers superior performance, with improved accuracy, reduced pseudo-positive/negative occurrence, and the lowest bias over local filament thickness compared to current approaches. It is noteworthy, however, that the adaptive global thresholding approaches (from Otsu to Yan) tend to be relatively accurate when judging the thick and bright bundles of the cytoskeleton. However, these approaches are unable to capture faint filaments, and as a result, generate a high pseudo-negative rate. Conversely, both adaptive local thresholding approaches, Niblack and Sauvola, generate numerous block-shaped pseudo-positive elements, and fail to capture the near-edge region of thick filaments. For MGT and Li method, although they showed satisfactory match rate and lower averaged pseudo-positive/negative rates, their performance is far less stable than ILEE (Fig. 4b).

As a next step, we evaluated the accuracy and stability of cytoskeletal indices using ILEE versus other commonly used imaging algorithms. To do this, we first computed the ground truth indices from the manually generated binary images; then, quantitative measurements were collected from all methods and normalized by the relative fold to the result generated from the corresponding ground truth image. As shown in Fig. 4d, ILEE showed the best overall accuracy and cross-sample stability compared to all other quantitative approaches, the highest accuracy for *occupancy*, *skewness*, CV, and *diameter_TDT*. However, we did observe that in terms of the morphology-sensitive indices (i.e., *linear density*, *severing activity*, and *branching activity*), the ILEE algorithm did not fully conform with data collected from the ground truth binary images. Upon further inspection, we determined that this is because the manually portrayed ground truth images and ILEE results showed different tendencies in judging the pixels in the narrow areas between two bright filaments (see *Discussion*). While other approaches displayed obvious, somewhat predictable, inaccuracies, the MGT and Li methods still generated satisfactory results, which

echoes their performance in actin segmentation. However, the performance of these two algorithms
among diverse and complex biological samples was not as stable as ILEE.

In order to further evaluate the stability and robustness of ILEE performance, we continued to analyze
the variance coefficient of all groups (Supplemental Fig. 10), uncovering that ILEE is the only approach
that simultaneously maintained high accuracy and stability. Next, we tested the robustness of ILEE and
other approaches against noise signal disturbance by adding different levels of Gaussian noise to the
image dataset (Supplemental Figs. 11-13). Using this approach, we observed that ILEE is still the best
performing algorithm, maintaining stable and accurate image binarization and cytoskeleton indices
against increasing noise. Taken together, these results demonstrate that ILEE has superior accuracy,
stability, and robustness over MGT and other classic automated image thresholding approaches in terms
of both cytoskeleton segmentation and index computation.

3D ILEE preserves more information and gains higher fidelity over 2D ILEE.
While the 2D ILEE generally outcompetes MGT and other commonly used automated algorithms (Fig.
4), the comparison of cytoskeleton segmentation approaches using the manually portraited ground truth
is currently limited to 2D images, because it is extremely difficult and error-prone to select the "3D ground
truth" voxel by voxel. However, as the 3D mode theoretically preserves information from the z-axis and
renders higher accuracy, it is necessary to circumvent the challenge and verify these merits. Therefore,
we turned to a different strategy using synthetic artificial 3D actin images with certain ground truth to
investigate the performance of 3D versus 2D mode (Fig. 5a).

First, we constructed an actin image simulation model, which mimics three different fractions of image
brightness (real actin, ground noise, and diffraction noise; see Fig. 2a) using three independent statistical
solutions (Supplemental Fig. 19; also see *Methods*). In brief, we utilized a training dataset of 31 diverse
3D samples to generalize the principles that describe the position-based brightness distributions of the
voxels for the three fractions independently. Next, we adopted only the skeleton image generated by 3D
ILEE from the training samples as the topological frame and refilled the whole image with new brightness
values via the actin image simulation model. The advantage of this, rather than generating it *de novo*,
was to ensure that the topological structure of the artificial filaments maximumly mimic the shape of
genuine actin. Accordingly, we generated 31 3D artificial images and processed them using the ILEE 2D
and 3D modes.

Our primary question was how 3D ILEE structurally improves the information loss through the 2D pipeline,
and moreover, yields improved accuracy of the cytoskeletal indices. As shown in Fig. 5b, we observed
that the 2D pipeline resulted in considerable information loss with reduced accuracy. Indeed, while the
3D segmentation almost accurately covers the filament voxels, the 2D pipeline only captured 8% of the
data points as pixels and 42% of the total length of filaments, suggesting potentially biased sampling.
Consequently, 3D ILEE generally gains index values closer to the ground truth than 2D ILEE, which
indicates the 3D mode indeed possesses a better absolute accuracy/fidelity.

In some scenarios, studies are interested in the relative difference between experimental groups for
biological interpretations, rather than the absolute accuracy of quantifiable features. Therefore, we also
measured the "comparative sensitivity" of 3D and 2D ILEE, which reflects their capability to determine
relatively high and low index values. To learn this, the index values of the ground truth images, as well
as those computed through 3D and 2D modes, were first normalized to the fold to the minimum value of
the corresponding group, followed by linear correlation analysis between the ground truth and ILEE
outputs of each sample. As shown in Fig. 5c, the 3D mode has a higher correlation for seven of the nine
indices, among which *diameter (TDT)*, *static severing activity*, and *anisotropy* got a considerable
improvement. Interestingly, we observed that the 2D mode performed slightly better for indices of the
density class (e.g., *occupancy* and *linear density*). We posit that this is because z-axis projection may
increase the contrast between samples with low and high filament density. In conclusion, our data
suggest that the 3D mode of ILEE indeed eliminates the issues related to image projection and provides
reliable results with higher accuracy and comparative sensitivity.

ILEE leads to the discovery of new features of actin dynamics in response to bacteria.
The primary impetus for creating the ILEE algorithm was to develop a method to define cytoskeleton
organization from complex samples, including those during key transitions in cellular status. For example,
previous research has demonstrated that the activation of immune signaling is associated with specific
changes in cytoskeletal organization[27,32–34]. Complementary to these studies, other research identified
the temporal and spatial induction of changes in the cytoskeletal organization as a function of the
pathogen (e.g., *Pseudomonas syringae*) infection and disease development[35–37]. The sum of these
studies, which broadly applied MGT-based quantitative analysis of cytoskeleton architecture, concluded
that virulent bacterial infection triggers elevated density (by *occupancy*) yet induced no changes in
filament bundling (by *skewness*) in the early stages of infection. Since one of our major motivations herein
was to develop an ILEE-based toolkit – supported by novel cytoskeletal indices – to investigate the
process of pathogen infection and immune signaling activation, we collected raw data from a previous

study[34] describing a bacterial infection experiment using Arabidopsis expressing an actin fluorescence

marker (i.e., GFP-fABD2), followed by confocal imaging and data analysis by ILEE as well as MGT

conducted by three independent operators with rich experience in actin quantificational analysis (Fig. 6).

Additionally, because researchers sometimes apply a universal global threshold to all images from a

batch of biological experiments to avoid tremendous labor consumption, we included this approach and

aimed to observe its performance as well. In this experiment, the only categorical variant is whether

sampled plants are treated with bacteria (EV) or not (mock). In total, nine indices that cover features of

density, bundling, severing, branching, and directional order are measured and compared.

Our first question is whether the operator's bias generated by MGT will influence the result and conclusion

generalized from the experiment. We thereby analyzed the correlation of MGT results set by the three

operators and found only a weak correlation between different operators (Fig. 6b), which indicates MGT

indeed introduces bias and potentially impacts quantitative results. Interestingly, while minor statistical

discrepancies between MGTs by different operators are found in some indices (i.e., skewness and

severing activity), most of the MGT results (both adaptive or fixed) shows the same trend as 2D ILEE,

yet with far higher standard deviation, or lower stability (Supplemental Fig. 14a) over a certain biological

treatment. This indicates that the historical data based on MGT should be majorly trustworthy despite the

biased single data points, but an accurate conclusion must be based on a high sampling number that

hedges the deviation of individual data points. Since ILEE gains less systemic error over biological

repeats, we also tested whether ILEE renders higher statistical power to identify potential significant

differences. As suggested by Supplemental Fig. 14b, we that ILEE indeed has the lowest p-values among

t-tests conducted for indices with a trend of difference. We believe these advantages demonstrate ILEE

as a better choice for actin quantification.

Next, we attempted to understand whether different indices of the same class, particularly density and

bundling, can reflect the quantitative level of the class in accordance, or instead show inconsistency. For

density, we correlated the *occupancy* and *linear density* values of all methods over actin images of both

mock and EV groups, and found that *occupancy* and *linear density* measurements are in high conformity,

with a Pearson coefficient at 0.98 (Supplemental Fig. 15). For bundling indices, we were interested in

their level of conformity because direct indices (based on binary shape) and indirect indices (based on

brightness frequency distribution) are entirely different strategies to measure bundling. Using the same

approach of correlation analysis, we found that *diameter_TDT* and *diameter_SDT* display strong positive

correlation, while *skewness* and *CV* have merely medium-low correlation, which echoes the previous

report demonstrating *skewness* and *CV* have different performance on the bundling evaluation[20].

Unexpectedly, we also found that *CV* (as a representative of indirect indices) and *diameter-SDT* (as a
representative of direct indices) have a striking zero correlation. This is perplexing, as it raises the
question on whether *skewness* or *CV* should be regarded as an accurate measurement of bundling (see
Discussion). This discrepancy is also observed by 3D ILEE, whose *CV* and *diameter-SDT* over mock
versus EV revealed the converse results at significant differences. In general, we believe the biological
conclusion that *Pst* DC3000 treatment renders increased actin bundling should be reconsidered with
further inspection.

Lastly, we asked if additional features of plant actin cytoskeletal dynamics in response to virulent bacterial
infection can be identified by the newly introduced indices and enhanced performance of ILEE. As shown
in Fig. 6, we observed significantly regulated *static severing activity*, *local anisotropy*, and static *branching*
activity triggered by the bacteria. At a minimum, these discoveries potentially lead to new biological
interpretations, and as a result, may contribute to the identification of additional immune-regulated
processes as a function of actin dynamics. However, while most 2D approaches were consistent and in
agreement with the other indices, the *static severing activity* estimated by 3D ILEE indicates a significant
but opposite conclusion. As suggested by the ground truth-based comparison on accuracy between 3D
and 2D ILEE (Fig. 5), we believed this discrepancy was due to the information loss and misinterpretation
by z-axis projection during the 2D pipeline. Hence, we have higher general confidence in the results and
conclusion by the 3D mode.

ILEE has broad compatibility with various sample types
Cytoskeleton imaging from live plant samples is arguably one of the most challenging types of images to
evaluate due to the dynamic topology and uneven brightness of actin filaments. While we demonstrated
that ILEE shows superior performance over plant actin samples, ILEE and the ILEE_CSK library are
generally designed for non-specific confocal images of the cytoskeleton, therefore applicable to other
types of samples. To investigate the compatibility of ILEE to other types of image samples, we tested
ILEE on both plant microtubules[38] and animal cell actin images (Supplemental Fig. 16). Importantly, we
found that ILEE and ILEE_CSK library can process both plant and animal images with satisfying
performance. This is encouraging, as ILEE can substitute or improve Hough transform, a straight-line
detection algorithm commonly used for animal cytoskeleton (generally straight and thick), but with some
limitation in neglecting and miscalculating curvy cytoskeleton fractions[14,15]. With the advancement of ILEE,
Hough transform-based analysis may not be essential, and the potential cytoskeleton indices that
rigorously require Hough transform can still utilize ILEE as a provider of binary image input for more
accurate results.

Discussion

We describe the creation of ILEE, an accurate and robust filament segregation algorithm for the unguided quantitative analysis of the organization of the eukaryotic cytoskeleton. As presented, this approach supports the *in vivo* analysis of both 2D and native 3D data structures, enabling an unbiased evaluation of cytoskeletal organization and dynamics. In addition to the development of key methods, we also generated a publicly available Python library that supports the automated computation of 12 filament indices of 5 classes of morphological features of the cytoskeleton. As described above, our data indicate that ILEE shows superior accuracy, robustness, and stability over existing cytoskeleton image analysis algorithms, including the widely adapted MGT approaches[17,39]. As a result of these newly developed approaches, we have developed an open-access library to conduct ILEE based cytoskeleton analysis, which eliminates limitations imposed by the classic 2D MGT approaches, including the introduction of user bias and different types of information loss.

Robustness of NNES-based prediction of global gradient thresholds

The gradient threshold (g_{thres}) defines the selected edge of actin filaments for implicit Laplacian transformation, the appropriateness of which greatly determines the performance of ILEE. To calculate g_{thres} without imposing a user-biased input, our strategy utilized specific feature values collected from the NNES curve and human-guided MGT to train a prediction model for the rapid rendering of a coarse area of the image background. Through this approach, we were able to deduce the corresponding g_{thres} by the mathematical relationship between the statistical distribution of the native pixel values and Scharr gradient magnitude of the coarse background (Supplemental Figs. 5, 6, 7, and *Methods*). This step might first appear unnecessary, since, alternatively, the most straight-forward strategy was to directly train a prediction model using the image gradient histogram and the human-determined g_{thres}. However, as the gradient operator (see *Methods*) for any given object pixel is influenced by the values of surrounding pixels, the calculated gradient on the edge of the background is highly impacted (overestimated) by the contrast of foreground (cytoskeleton) versus the background. In other words, the brightness frequency distribution of the background gradient will change at elevated cytoskeleton fluorescence levels, even though the background *per se* does not. The outcome of this is a significant decrease in the accuracy of gained g_{thres}. For this reason, we assert that g_{thres} should be mathematically deduced from a pre-determined region of background, rather than directly predicted via human-trained models or calculated from the frequency distribution of gradients even using the pre-determined background.

ILEE and visual inspection have different inclinations on the topology between two bright filaments

Our data demonstrate that ILEE generally shows dominant robustness and accuracy for most indices,
compared to the manually portrayed ground truth binary image. However, there are three indices (*linear*
density, *static severing activity*, and *static branching activity*) where ILEE renders stable yet dramatically
lower output than those derived from the ground truth images (Fig. 4d). Since the result is very stable,
we anticipate there may be certain systemic inclinations in either the human-portrayed ground truth or
the ILEE algorithm. After inspecting the binary images generated by ILEE compared with the ground truth,
we identified a potentially critical reason: ILEE is less likely to presume the ambiguous, dubious thin
connections inside the narrow space between two bright filaments. As demonstrated by Supplemental
Fig. 17, while the ground truth and ILEE binary image look very similar, their skeleton images – which
represent their topological structure – show a discrepancy between two bundles of bright filaments.
Considering their procedure of generation, we speculate this is because: (1), ILEE fully outer-joins the
binary results by a lower K_1 and a higher K_2 (Fig. 3c), among which K_2 sacrifices the sensitivity to local
filaments at the high signal region to improve the sensitivity at the low signal region, including the edge
of thick filaments; and (2), human eyes tend to "hallucinate" imaginary filaments that do not statistically
exist. According to our knowledge, there is no overwhelming evidence suggesting either 2D ILEE or the
human eye is more accurate, but 2D ILEE is indeed more stable and conservative. However, 3D ILEE
may solve this paradox because many "adjacent" bright bundles are artifacts out of z-axis projection,
which is distant enough in 3D space to offer ILEE with a satisfactory resolution to split.

The 2D and 3D modes of ILEE may lead to different conclusions
For the analysis over bacteria treated leaf sample (Fig. 6), 2D ILEE generally agreed with MGT while
offering higher robustness and stability. Interestingly, this is not always the case for 3D ILEE, since it led
to different statistical conclusions from 2D approaches in *linear density*, *skewness*, *diameter (SDT)*, *static*
severing activity, and *static branching activity*. However, such difference is understandable, because the
3D mode indeed computes indices with higher accuracy and overcomes many limitations by the 2D data
structure. As demonstrated by Fig. 5b, c, 3D ILEE provide more or less improved absolute fidelity and
comparative sensitivity for *skewness*, *diameter (SDT)*, *static severing activity*, and *static branching*
activity; therefore, our current data support that 3D ILEE is more trustworthy for these indices. Similarly,
the discrepancy of *linear density* over 2D approaches versus 3D ILEE in Fig. 6d also agrees with our
observation that 2D ILEE has higher comparative sensitivity for indices reflecting density. Hence, in terms
of any potential discrepancy between the 2D and 3D modes, we offer a general recommendation: for
occupancy only, 2D ILEE has better capability to distinguish biological difference; for all other
cytoskeleton features, 3D ILEE is more trustworthy and dependable. Although it is currently difficult to
make a definite conclusion whether the current 2D or 3D mode is more accurate across all scenarios, it

is predictable that 3D computation will gradually substitute the classic 2D data structure, with the
mainstream personal computers gaining increasing computational power in the future.
569

Future development and applications
While ILEE has already remedied many disadvantages of traditional methods such as MGT, we are still
working to advance further the ILEE approaches presented herein. Our goal is to ultimately establish
algorithms and the toolbox that provide cytoskeletal indices at perfect accuracy and effectively answer
the specific demands in this field of study. As such, we offer the following as a list of potential upgrades
and applications to be integrated into the library:

- Deep learning-based cytoskeleton segmentation algorithm with "foreign object" removal function.
As presented herein, ILEE enables the generation of trustworthy binary images on large scale,
which enables the construction of deep learning models to identify cytoskeleton components from
confocal images with potentially better performance. The deep learning-based approach is also
the key to solving the ultimate problem of all current cytoskeleton segmentation algorithms
(including ILEE), which is the inability to detect and erase non-cytoskeleton objects with high
fluorescence, such as the nucleus and cytoplasm. As one approach to circumvent this limitation,
free fluorescence proteins can be used as a cell-permeable false signal. This will enable us to
train the model to recognize and exclude the non-cytoskeleton-like foreign objects, to render
ideally pure cytoskeletal datasets.

- Vectorized intermediate image. After generating the difference image (i.e., I_{dif}, Fig. 3a) using
ILEE, one computational limitation of our Python-based algorithm is the tradeoff between the
demand for unlimited high-resolution imaging versus limited computational power. Accordingly,
an ideal strategy is to transfer the pixel/voxel image to a 2D vector image or 3D polygon mesh for
index calculation. This will further enhance the accuracy of ILEE at an acceptable requirement of
computational power.

- Regions of interest and organelle segmentation. There is currently a high demand in research of
plant cell biology to quantify cytoskeletal parameters in stomatal guard cells, as well as additional
cellular and subcellular architectures. In future releases of ILEE, we will develop approaches to
enable automated recognition and selection of regions of interest, such as stomata, for various
demands by the community.

- Compatibility to *x-y-t* and *x-y-z-t* data, where *t* represents time. We are in the process of
developing a 4D-compatible analysis of cytoskeletal dynamics that tracks filament organization
over time. This approach will provide a temporal evaluation of supported indices with high
accuracy and robustness.

Methods

Plant genotypes and growth. *Arabidopsis thaliana* Col-0 expressing the actin cytoskeletal marker GFP-

fABD2[36] was used in this study. Arabidopsis seeds were stratified for 2 d in the dark at 4°C then sown

onto the soil. All plants were grown in a BioChambers model FLX-37 walk-in growth chamber

(BioChambers, Manitoba, Canada) at 20°C under long-day conditions (16 h of light/8 h of dark) with 60%

relative humidity and a light intensity of approximately 120 μmol photons $m^{-2}s^{-1}$.

Bacteria growth and plant inoculation. *Pseudomonas syringae* pv. *tomato* DC3000 (*Pst* DC3000)

strains were grown as previously described[36]. Antibiotics used in this study include kanamycin (GoldBio

#K-120-5, 50 μg mL^{-1}), and rifampicin (GoldBio #R-120-1, 100 μg mL^{-1}). Bacterial treatments for actin

dynamics analysis were conducted following previously described methods[36]. Briefly, 2-week-old

Arabidopsis Col-0/GFP-fABD2 was dipped for 30s in Dip-inoculation solution (10 mM MgCl2 + 0.02%

Silwet-77) with DC3000 with empty vector (EV) at a concentration OD_{600} = 0.2 (ca. 2 x 10^7 colony forming

units/mL). Confocal images were collected at 24 hours post-inoculation.

Mouse cancer cells sample. Yale University Mouse Melanoma line YUMMER1.7D4 cells (EMD

Millipore, #SCC243) were cultured in DMEM (ATCC # 30-2006) supplemented with 10% FBS (Gibco

#10437-028), 1% Pen-strep (ThermoFisher, #15140122), and 1% NEAA (Gibco, #11140035). For

staining actin stress fiber, approximately 10,000 cells were seeded onto a glass coverslip maintained in

a six-well plate overnight. First, semiconfluent cells were fixed with 3.7% formaldehyde for 15 min at RT,

washed three times with PBS, then blocked in PBS supplemented with 2% BSA and 0.1% Triton for one

hour at RT. Next, cells were incubated with 100 nM rhodamine-phalloidin (Cytoskeleton Inc, #PHDR1) in

blocking buffer for 30 minutes at RT in the dark, then washed with PBS three times each for 5 min with

gentle shaking at RT. Stained cells on the coverslip were mounted in ProLong Glass Antifade Mountant

with DAPI (ThermoFisher, #P36982). Slides were cured overnight at room temperature and were then

imaged.

Confocal microscopy. For plant leaf actin images, 2-week-old Col-0/GFP-fABD2 plants were used for

data collection and analysis. Images of epidermal pavement cells and guard cells were collected using

laser confocal scanning microscope (Olympus FV1000D) by obtaining z-series sections at 0.5 μm

intervals. Optical setting: 65x/1.42 PlanApo N objective with a 488 nm excitation laser and 510-575 nm

emission filter. Images were collected at a resolution of 800 x 800 x 25 (*x-y-z*) and a 12-bit dynamic range.

Voxel size was 0.132 μm at the *x*- and *y*-axis and 0.5 μm at the *z*-axis. For animal cell actin images,

YUMMER1.7D4 stained by rhodamine-phalloidin were sampled by the same confocal system. Optical

setting: 100x/1.40 UPLSAPO objective with a 559 nm excitation laser and 570-670 nm emission filter.

Images were collected at a resolution of 800 x 800 x 10 (*x-y-z*) with voxel size of 0.039 μm at the *x*- and

y-axis and 0.16 μm at the z-axis

Manually portrayed ground truth binary image. Seven raw projected images of 400x400 pixel size

with diverse visual appearance (e.g., actin density, shape, thickness, fluorescence brightness) were

selected from our actin image database. Using a pen-drawing display (HUION KAMVAS 22 Plus, GS2202)

and GIMP, we enhanced the brightness of low-value pixels to clarify the actin structure and carefully

portrayed the actin component of the selected image sample. The portrayed layer was extracted and

transferred to binary format for further evaluation.

Double-blind MGT analysis. For the mock versus *P. syringae*-inoculated sample pool, we erased the

sample name and randomized the order and distributed them to three independent scientists with rich

experience in cytoskeleton analysis (referred to as OA, OB, and OC) to let them determine the global

threshold value of each sample manually using the approach described previously[35]. Once completed,

we restored the grouping of the samples for batch analysis. We use Python to mimic the MGT pipeline

generally conducted by ImageJ after determining a specific thresholding value. OA also provided a

universal threshold value (referred to as OA_fix) that applies to all samples as a commonly used fast

MGT approach.

Statistical analysis and data visualization. All data analysis was conducted in the Python 3.8

environment and was described in figure legends.

Determination of K_2 for sample batches. For both 2D and 3D modes of ILEE, each 12-bit single-

channel 3D image $I(x, y, z)$ in a batch of samples was transferred into 2D image $I_{proj}(x, y)$ by z-axis

maximum projection, where each pixel $I_{proj}(x, y) = \max\{I(x, y, z) | z \in N^+, z < z_{max}\}. I_{proj}$ was

processed by a Niblack thresholding function[16] (library API reference [1]) to render $I_{nibthres}(x, y)$, with

parameter k = -0.36 and $window\_size = 2 \operatorname{int}(25\, l) + 1$, where l is the mean of x and y resolution of I_{proj}.

A binary image defined as $I_{binary}(x, y) = \{1, if\ I_{proj}(x, y) > I_{nibthres}(x, y); 0, else\}$ was generated. The

binary image was processed through Euclidean distance transformation (library API reference [2]), and

the mean of the highest 5% values was used as the input of the K_2 calculation function (see Fig. 3) that

outputs individual recommended K_2. Finally, the mean of all individual K_2 will be output as the recommend

K_2 for the total batch.

ILEE. An abbreviated workflow of ILEE is illustrated in Fig. 3b. Here, we describe the overall process in

further detail. For 2D mode, input image structure is $I(pix) = I_{proj}(x,y) = \max\{I(x,y,z)|z \in N^+, z <$

$z_{max}\}$. For 3D mode, $I(pix) = I(x,y,z)$. First, I is treated by a significant difference filter (SDF;

Supplemental Fig. 4) to render I_{SDF}, where:

$$I_{surround} = \{I(x \pm 1, y(,z)), \; I(x, y \pm 1(,z)), \; (0.293\, I(x, y(,z)) + 0.707\, I(x \pm 1, y \pm 1(,z)))\}$$

$$I_{SDF}(x, y(,z)) = \begin{cases} I(x, y(,z)), & if\ |Mean(I_{surround}) - I(x, y(,z))| < 2\, STD(I_{surround}) \\ Mean(I_{surround}), & else \end{cases}$$

In other words, it substitutes a pixel by the mean of the 8 adjacent pixels if the absolute difference

between it and that mean is higher than 2-fold of the standard deviation of the surrounding pixels. Then,

I_{SDF} is input to a discrete gaussian filter with a 3x3(x3) weighting kernel at σ = 0.5, to render $I_{pre} =$

$Gauss(I_{SDF})$, which is the smoothed pre-processed image. Since confocal microscopy has different

resolutions on x/y and z axis (hereby named as U_{xy} and U_z), we adjusted the weighting kernel from

O_{Gauss} to O'_{Gauss} in 3D mode particularly by a scaling operator, as below:

$$f = U_{xy}/U_z$$

$$O_{scalar} = \frac{3}{1 + 2f} \begin{bmatrix} \begin{bmatrix} f & f & f \\ f & f & f \\ f & f & f \end{bmatrix} \\ \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \\ \begin{bmatrix} f & f & f \\ f & f & f \\ f & f & f \end{bmatrix} \end{bmatrix}$$

$$O'_{Gauss} = O_{Gauss} \circ O_{scalar}$$

684   From I_{pre}, the gradient magnitude image *G* is rendered through Scharr operator as:

$$G_x = \begin{bmatrix} 3 & 0 & -3 \\ 10 & 0 & -10 \\ 3 & 0 & -3 \end{bmatrix} * I_{pre}$$

$$G_y = \begin{bmatrix} 3 & 10 & 3 \\ 0 & 0 & 0 \\ 3 & -10 & -3 \end{bmatrix} * I_{pre}$$

$$G = \frac{\sqrt{G_x^{\circ 2} + G_y^{\circ 2}}}{32\sqrt{2}}$$

685   for 2D mode, or:

$$G_y = \begin{bmatrix} \begin{bmatrix} 9 & 30 & 9 \\ 0 & 0 & 0 \\ 9 & -30 & -9 \end{bmatrix} \\ \begin{bmatrix} 30 & 100 & 30 \\ 0 & 0 & 0 \\ -30 & -100 & -30 \end{bmatrix} \\ \begin{bmatrix} 9 & 30 & 9 \\ 0 & 0 & 0 \\ -9 & -30 & -9 \end{bmatrix} \end{bmatrix} \circ O_{scalar} * I_{pre}$$

$$G_z = \begin{bmatrix} \begin{bmatrix} 9 & 30 & 9 \\ 30 & 100 & 30 \\ 9 & 30 & 9 \end{bmatrix} \\ \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \\ \begin{bmatrix} -9 & -30 & -9 \\ -30 & -100 & -30 \\ -9 & -30 & -9 \end{bmatrix} \end{bmatrix} \circ O_{scalar} * I_{pre}$$

$$G = \frac{\sqrt{G_x^{\circ 2} + G_y^{\circ 2} + G_z^{\circ 2}}}{256}$$

687 for 3D mode. Next, to calculate the gradient threshold (g_{thres}) as an input for ILEE, a global threshold

688 t_{cbg} to determine the coarse background (I_{cbg}, as flattened image) is calculated by the NNES (non-

689 connected negative element scanning) function that satisfies:

$$I_{binary.NNE}(x) = \{I(pix)|I(pix) > x\}$$

$$Count.of.NNE\left(I_{binary.NNE}(x_{peak})\right) = Count.of.NNE\left(I_{binary.NNE}(x)\right)_{max}$$

$$t_{cbg} = \begin{cases} 1.163\,x_{peak} + 101.68, & if\ I\ is\ 2D\ (see\ Sup.\ Fig\ 1\mathbf{b}) \\ 7.33\,x_{peak}, & if\ I\ is\ 3D\ (see\ Sup.\ Fig\ 2\mathbf{b}) \end{cases}$$

$$I_{cbg} = \{I(pix)|I(pix) \le t_{cbg}\}$$

692 If I is processed using the 2D mode, as demonstrated previously (see Supplemental Figure 5), the

693 statistical mean and STD of gradient magnitude of I_{cbg} ($\mu_{G.cbg}$ and $\sigma_{G.cbg}$, respectively) are univariate

694 proportional functions of the STD of I_{cbg}, σ_{cbg}, not influenced by the mean of I_{cbg}. Particularly, μ_G and σ_G,

695 of the coarse background area of the gradient image is:

697            $$\mu_{G.cbg} = 0.8542\,\sigma_{cbg}$$

698            $$\sigma_{G.cbg} = 0.4469\,\sigma_{cbg}$$

700 With these, we established a g_{thres} estimation model, where g_{thres} = $\mu_{G.cbg}$ + $k\,(\sigma_{cbg})\,\sigma_G$. Using the

701 optimized parameters described in Supplemental Figure 7, g_{thres} is deduced as:

703            $$k(\sigma_{cbg}) = 0.040018\,\sigma_{cbg}$$

704            $$g_{thres} = \mu_{G.cbg} + k(\sigma_{cbg})\,\sigma_{G.cbg}$$

706 If I is processed by the 3D mode, the inconformity of U_{xy} and U_z will not only impact the weighting of

707 Scharr operator, but also influencing the proportional coefficient of σ_{cbg} to μ_G and σ_G. We simulated the

708 accurate mathematical relationship between the proportional coefficient k_s and U_z/U_{xy} (see

709 Supplemental Figure 6), so g_{thres} for 3D can be calculated as:

711            $$\mu_{G.cbg} = k_{s.\mu}\,\sigma_{cbg}$$

712            $$\sigma_{G.cbg} = k_{s.\sigma}\,\sigma_{cbg}$$

713            $$k(\sigma_{cbg}) = 0.040519\,\sigma_{cbg}$$

714
$$g_{thres} = \mu_{G.cbg} + k(\sigma_{cbg})\,\sigma_{G.cbg}$$

The total process above to computer g_{thres} is referred as function $g_{thres}.estimate(I_{pre}, I_{cbg})$ in Fig. 3a

(1.3).

The critical step to generate the threshold image of the input sample I is implicit Laplacian smoothing.

This algorithm builds a linear system using the Laplacian operator to achieve local adaptive thresholding

based on edges of cytoskeletal components. Leveraging the spectral characteristics of discrete Laplacian

operators, we could filter out high-frequency components (aka, high fluorescence fractions of

cytoskeleton) while preserving the low-frequency salient geometric features of background fluorescence.

First, an edge image is defined as $I_{edge}(pix) = I_{pre}(pix), if\, G(pix) > gthres; 0, else,$ but we transform

I_{edge} into a flattened image vector $\overrightarrow{I_{edge}}$, and hence it can be further be represented as:

727
$$\overrightarrow{I_{edge}} = S\,\overrightarrow{I_{pre}}$$

729 where S is a (sparse) selection matrix, which is a diagonal matrix with i-th diagonal entry being 1 if the i-

730 th pixel has a gradient above g_{thres}.

732 Next, according to the concept of Laplacian operator, we constructed the (sparse) Laplacian matrix L that

733 satisfies:

$$L = \begin{cases} L_4\ \overrightarrow{I_{flt}} = \triangle_{p\_4} * I(x,y) & (2D\ alternative) \\ L_8\ \overrightarrow{I_{flt}} = \triangle_{p\_8} * I(x,y) & (2D\ default) \\ L_6\ \overrightarrow{I_{flt}} = \triangle_{p\_6} * I(x,y,z) & (3D) \end{cases}$$

736   where the Laplacian operators in different modes are defined as:

$$\triangle_{p\_4} = \begin{bmatrix} 0 & -1 & 0 \\ -1 & 4 & -1 \\ 0 & -1 & 0 \end{bmatrix}$$

$$\triangle_{p\_8} = \begin{bmatrix} -\sqrt{2}/2 & -1 & -\sqrt{2}/2 \\ -1 & 4+2\sqrt{2} & -1 \\ -\sqrt{2}/2 & -1 & -\sqrt{2}/2 \end{bmatrix}$$

$$\triangle_{p\_6} = \begin{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \\ \begin{bmatrix} 0 & -1 & 0 \\ -1 & 6 & -1 \\ 0 & -1 & 0 \end{bmatrix} \\ \begin{bmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \end{bmatrix}$$

Therefore, we can establish an implicit Laplacian linear system that targets the edge:

$$(L + KS)\,\overrightarrow{I_{thres}} = \overrightarrow{I_{edge}} = S\,\overrightarrow{I_{pre}}$$

where $\overrightarrow{I_{thres}}$ is the unknown to be solved by conjugate gradient method and restored to 2D/3D array I_{thres},

and K is a weight that adjusts the influence Laplacian operator has on the result (see Fig. 3B).

Finally, a difference image I_{dif} is calculated as $I_{dif} = I_{pre} - I_{thres}$. To further clean the noise, a temporary

binary image $I_{binary.temp}(pix) = \{1, if\ I_{dif}(pix) > 0; 0, else\}$ is generated and the sizes of all positive

connected component (element) are counted. An adjusted difference image I_{dif_adj} is generated by

lowering the values of potential noise into the mean of negative pixels:

$$I_{dif\_adj} = \begin{cases} I_{dif}(pix), & if\ elment\ size \geq 3 \\ Mean\big(I_{dif}(pix)\,\big|\,I_{dif}(pix) < 0\big), & else \end{cases}$$

Using I_{dif_adj}, the various versions of the binary image for computation of different indices are calculated

as:

$$I_{binary}(pix) = \{1, if\ I_{dif\_adj}(pix) > 0; 0, else\}$$

$$I_{binary.ovsp}(pix) = \{1, if\ Interp_{3,3}(I_{dif\_adj}(pix)) > 0; 0, else\}$$

for the 2D mode, where $Interp_{3,3}\ (I_{dif})$ is an oversampled image by bicubic interpolation whose resolution

on both x- and y-axis are upscaled for 3 folds, or

$$I_{binary.ori}(pix) = \{1, if\ I_{dif}(pix) > 0; 0, else\}$$

$$I_{binary}(pix) = \{1, if\ Interp_{1,1,1/f}(I_{dif}(pix)) > 0; 0, else\}$$

$$I_{binary.ovsp}(pix) = \{1, if\ Interp_{3,3,3/f}(I_{dif}(pix)) > 0; 0, else\}$$

for the 3D mode, where $Interp_{1,1,1/f}(I_{dif}(pix))$ aims to restore the voxel to cubic shape by only interpolate

z axis to 1/f folds of its original resolution and $Interp_{3,3,3/f}(I_{dif})$ additionally enhance 3 folds of resolution

for all three axes.

Computation of cytoskeletal indices

•   **_Occupancy_**: the frequency of the positive pixels in the computed binary image, or

$\sum I_{binary}(pix)/N$ for 2D mode and $\sum I_{binary.ori}(pix)/N$ for 3D mode, where _N_ is the number of

total pixels.

•   **_Linear density_**: the length of the skeletonized filament per unit of 2D or 3D space. For 2D mode,

$I_{binary.ovsp}$ is skeletonized using Lee's approach[18] to render the skeletonized image I_{sk}. Then,

linear density is calculated as:

$$linear\ density = \sum length(x, y)/N.$$

$$length(x, y) = \begin{bmatrix} \sqrt{2}/2 & 1/2 & \sqrt{2}/2 \\ 1/2 & 0 & 1/2 \\ \sqrt{2}/2 & 1/2 & \sqrt{2}/2 \end{bmatrix} * I_{sk}(x, y)$$

For the 3D mode, I_{sk} is rendered by I_{binary}, and we use the sum of the Euclidean lengths of all

(graph theory defined) branches obtained by Skan library[22] as the total length of skeletonized

filament and divide it by _N_. This is because sphere-cavities structures existent in the 3D

skeletonized images are not applicable to the concept of length.

• **Skewness:** the (statistical) skewness of the fluorescence value of positive pixels, or
mathematically:

$$skewness = \frac{1}{N_{pos}} \sum_{I_{binary}(pix)==1} \left(\frac{I(pix) - \mu_{pos}}{\sigma_{pos}}\right)^3$$

783
$$skewness = \frac{1}{N_{pos}}$$

785 where N_{pos}, μ_{pos}, and σ_{pos} represents the count, mean, and standard deviation of positive pixels
786 in the raw image I.

787 • **CV:** the (statistical) coefficient of variance of the fluorescence value of positive pixels, or
788 mathematically:

789
$$CV = \frac{\sigma_{pos}}{\mu_{pos}}$$

791 **Diameter_TDT:** average filament diameter estimated by Euclidian distance transformation of total
792 binary image. The Euclidian distance transformation map I_{dis} is calculated as an image with the
793 same shape of $I_{binary.ovsp}$, but the value of each pixel is:

794
$$I_{dis}(x, y(,z)) = \begin{cases} \sqrt{\sum_{i=x,y(,z)} (i - i_{nnp})^2}, & if\ I_{binary.ovsp}(x, y(,z)) = 1 \\ 0, & if\ I_{binary.ovsp}(x, y(,z)) = 0 \end{cases}$$

where i_{nnp} is the coordinates of the nearest negative pixel to (*x,y,z*). Therefore, the
Diameter_TDT is mathematically defined as:

$$Diameter\_TDT = \text{Mean } \{I_{dis}(pix)|I_{dis}(pix) > 0\}$$

• **Diameter_SDT:** average filament diameter estimated by Euclidian distance transformation values
of $I_{binary.ovsp}$ *but sampled using only* positive pixels on I_{sk}, or mathematically as:

$$Diameter\_SDT = \text{Mean } \{I_{dis}(pix)|I_{sk}(pix) = 1\}$$

• **Static severing activity:** the count of connected components in the binary image per unit length
of skeletonized filament. For images captured in 2D mode, it is $Count.of.NNE(I_{binary.ovsp})/$

$\sum length(x,y)$. In 3D mode, both the count of non-connected elements and the length of
skeletonized filament are called from the Skan library using I_{sk} computed from I_{binary} as the input.

• **_Static branching activity_:** the branching point count per unit length of skeletonized filament. I_{sk}
is obtained from $I_{binary.ovsp}$ for 2D mode or I_{binary} for 3D mode respectively and is next input into
Skan library. The total number of type-3 and type-4 branches[22] is collected and then divided by
the length of the skeletonized filament.

• **_Local anisotropy_:** We performed a local averaging of filament alignment tensor, which is
constructed as follows. First, we calculate the unit direction vector for each straight filament
segment g_i. Then, the covariance matrix for each segment is obtained from the following equation:

$$t_i = g_i g_i^t$$

This rank-2 tensor is independent of the orientation of the line segment, and can thus be averaged
over a region containing a collection of unoriented line segments. We weighed each filament
tensor in a circular/spherical neighborhood by the length of every filament to produce a smoothed
tensor field. The eigenvector corresponding to the largest eigenvalue indicated the primary
orientation of filaments in this region. The difference between the maximum and the minimum
eigenvalues is an indicator of the anisotropy in this region. If all the eigenvalues are the same,
the indicator is 0, which implies an isotropic region. If the eigenvalues other than the maximum
are all nearly 0, all the filaments in this region are parallel to each other. In this case, they are all
aligned with the maximum eigenvector, the dominant filament direction of this region.

Simulation model of 3D artificial actin image. For each of the 31 training samples $I(x,y,z)$, the binary
image I_{binary} and skeleton image I_{sk} were previously obtained via ILEE, as described above. Three
independent fractions reflecting artificial actin ($I_{atf.a}$), artificial ground noise ($I_{atf.gn}$), and artificial
diffraction noise ($I_{atf.d}$) are simulated independently.

To calculate $I_{atf.a}$, the brightness of the real actin (I_a) of all training sample is collected as:

$$I_a = I \circ I_{binary}.$$

Here, we define the "reference voxel" of a non-zero voxel in I_a as its nearest voxel whose position has a

positive value in I_{sk}. In other words, for each non-zero voxel of $I_a(x, y, z)$, the distance from (x, y, z) to its

nearest reference voxel $(P_R = (x_R, y_R, z_R))$ on the skeleton image is defined as $D_R(x, y, z)$, where

843
$$D_R(x, y, z) = \min_{I_{sk}(x_i, y_i, z_i)=1} \sqrt{(x - x_i)^2 + (y - y_i)^2 + (z - z_i)^2}, \text{ and}$$

844
$$P_R(x, y, z) = \underset{I_{sk}(x_i, y_i, z_i)=1}{\text{argmin}} \sqrt{(x - x_i)^2 + (y - y_i)^2 + (z - z_i)^2}.$$

846 Meanwhile, the distance from the reference voxel to the nearest non-actin voxel, $D_{R2N}(x, y, z)$, is

847 calculated as:

849
$$D_N(x, y, z) = \min_{I_a(x_i, y_i, z_i)=0} |P_R(x, y, z) - (x_i, y_i, z_i)|.$$

851 Next, the "relative eccentric distance", $D_{rlt}(x, y, z)$, and the relative brightness comparing to the reference

852 voxel, $I_{rlt}(x, y, z)$ are defined as:

854
$$D_{rlt}(x, y, z) = D_R(x, y, z)/D_{R2N}(x, y, z), \text{ and}$$

855
$$I_{rlt}(x, y, z) = I_a(x, y, z)/I_a(P_R(x, y, z)).$$

857 Note that D_{rlt} and I_{rlt} default to 0 whenever the denominator is 0. Because D_{rlt} is discrete, each possible

858 value of D_{rlt} can be counted as $n(D_{rlt})$. Using the I_{rlt} and D_{rlt} of all voxels with $D_{rlt} \leq 1$ in the entire 31

859 images, a polynomial regression model between D_{rlt} and I_{rlt} is fitted using $\log_{10} n(D_{rlt})$ as weights and

860 adjusted into a piecewise polynomial function:

862
$$\hat{\mu}_{I.rlt}(D_{rlt}) = \begin{cases} 1, & (D_{rlt} \leq 0.2) \\ -0.647D_{rlt}^2 + 0.226D_{rlt} + 0.989, & (0.2 < D_{rlt} \leq 1.4) \end{cases}$$

864 Similarly, the STD of the I_{rlt} is also fitted as below:

866
$$\hat{\sigma}_{I.rlt}(D_{rlt}) = -0.562D_{rlt}^2 + 0.778D_{rlt} + 0.065, (D_{rlt} \leq 1.4).$$

868 Finally, the artificial actin is generated as random values:

870
$$I_{atf.a}(x,y,z) = \begin{cases} 0, & \text{if } D_{rlt}(z,x,y) > 1.4 \text{ or } I_{binary}(z,x,y) = 0 \\ I(z_R, x_R, y_R)\,\hat{I}_{rlt}(D_{rlt}(x,y,z)) \mid \hat{I}_{rlt} \sim N(\hat{\mu}_{I.rlt}(D_{rlt}(x,y,z)),\, \hat{\sigma}^2_{I.rlt}(D_{rlt}(x,y,z))), & \text{otherwise} \end{cases},$$

where N means normal distribution. The random values are clamped to [0,4095].

To calculate $I_{atf.gn}$, two different strategies are utilized to generate two groups of voxels. For the voxels

positions in the coarse background of I, we apply a fold of change that is subject to a normal distribution

to each $I(x,y,z)$ as the new ground noise, plus a Gaussian filter (O'_{Gauss}, see ILEE section of *Method*) to

neutralize the magnified gradient; for the rest beyond the coarse background, we gave random values

that subject to a Burr-12 distribution (Library API reference [3]) fitted by voxels in the coarse background

of $I(x,y,z)$, that is:

881
$$I_{atf.gn}(x,y,z) = \begin{cases} 2^k\,I(x,y,z) * O'_{Gauss} \mid k \sim N(0, 0.25^2), & \text{if } I(x,y,z) < t_{cbg} \\ b \mid b \sim Burr12(c, d, \mu_b, \sigma_b), & \text{otherwise} \end{cases},$$

883 where c, d, μ_b, and σ_b are the fitted parameters of the Burr-12 distribution using the training voxels

884 $\{(x,y,z) \mid I(x,y,z) \le t_{cbg}\}$. Similarly, the random values are clamped to [0,4095].

886 Third, for I_{atf_d}, voxels of the positions in I that are neither coarse background nor real actin are used to

887 fit an exponential distribution:

889
$$I_{atf\_d}(x,y,z) = \begin{cases} e \mid e \sim Exp(\mu_b, \sigma_b), & \text{if } I_{atf.a}(x,y,z) = 0 \text{ and } I(x,y,z) \ge t_{cbg} \\ 0, & \text{otherwise} \end{cases},$$

891 where μ_e and σ_e are the fitted parameters of the exponential distribution using the training voxels

892 $\{(x,y,z) \mid I(x,y,z) \ge t_{cbg}\}$. Similarly, the random values are clamped to [0,4095].

894 Finally, these fractions of brightness are summed to generate the complete artificial image I_{atf}, as:

896
$$I_{atf} = I_{atf.a} + I_{atf.gn} + I_{atf.d}.$$

Methods Library API reference:

[1] https://scikit-image.org/docs/stable/api/skimage.filters.html#skimage.filters.threshold_niblack

[2]https://docs.scipy.org/doc/scipy-

0.14.0/reference/generated/scipy.ndimage.morphology.distance_transform_edt.html

[3] https://docs.scipy.org/doc/scipy/reference/generated/scipy.stats.burr12.html

References

1. Lian, N., Wang, X., Jing, Y. & Lin, J. Regulation of cytoskeleton-associated protein activities:
Linking cellular signals to plant cytoskeletal function. *Journal of Integrative Plant Biology* **63**, 241–
250 (2021).

2. Blanchoin, L., Boujemaa-Paterski, R., Sykes, C. & Plastino, J. Actin Dynamics, Architecture, and
Mechanics in Cell Motility. *Physiol Rev* **94**, 29 (2014).

3. Brouhard, G. J. Microtubule dynamics: an interplay of biochemistry and mechanics. 13 (2018).

4. Nick, P. Mechanics of the Cytoskeleton. in *Mechanical Integration of Plant Cells and Plants* (ed.
Wojtaszek, P.) 53–90 (Springer, 2011). doi:10.1007/978-3-642-19091-9_3.

5. Carlton, J. G., Jones, H. & Eggert, U. S. Membrane and organelle dynamics during cell division.
Nature Reviews Molecular Cell Biology **21**, 151–166 (2020).

6. Kristó, I., Bajusz, I., Bajusz, C., Borkúti, P. & Vilmos, P. Actin, actin-binding proteins, and actin-
related proteins in the nucleus. *Histochem Cell Biol* **145**, 373–388 (2016).

7. Rush, T. *et al.* Synaptotoxicity in Alzheimer's Disease Involved a Dysregulation of Actin
Cytoskeleton Dynamics through Cofilin 1 Phosphorylation. *J. Neurosci.* **38**, 10349–10361 (2018).

8. Marchisella, F., Coffey, E. T. & Hollos, P. Microtubule and microtubule associated protein
anomalies in psychiatric disease. *Cytoskeleton* **73**, 596–611 (2016).

9. Absi, A. A. *et al.* Actin Cytoskeleton Remodeling Drives Breast Cancer Cell Escape from Natural
Killer–Mediated Cytotoxicity. *Cancer Res* **78**, 5631–5643 (2018).

10. Li, P. & Day, B. Battlefield Cytoskeleton: Turning the Tide on Plant Immunity. *MPMI* **32**, 25–34
(2019).

11. Melak, M., Plessner, M. & Grosse, R. Correction: Actin visualization at a glance. *Journal of Cell
Science* **130**, 1688–1688 (2017).

12. Lichtenstein, N., Geiger, B. & Kam, Z. Quantitative analysis of cytoskeletal organization by digital
fluorescent microscopy. *Cytometry Part A* **54A**, 8–18 (2003).

13. Shah, S. A., Santago, P. & Rubin, B. K. Quantification of biopolymer filament structure.
Ultramicroscopy **104**, 244–254 (2005).

14. Liu, Y., Nedo, A., Seward, K., Caplan, J. & Kambhamettu, C. Quantifying Actin Filaments in
Microscopic Images using Keypoint Detection Techniques and A Fast Marching Algorithm. in *2020*

IEEE International Conference on Image Processing (ICIP) 2506–2510 (2020).

doi:10.1109/ICIP40778.2020.9191337.

15. Liu, Y., Mollaeian, K. & Ren, J. An Image Recognition-Based Approach to Actin Cytoskeleton

Quantification. *Electronics* **7**, 443 (2018).

16. Alioscha-Perez, M. *et al.* A Robust Actin Filaments Image Analysis Framework. *PLoS Comput Biol*

12, e1005063 (2016).

17. Higaki, T., Kutsuna, N., Sano, T., Kondo, N. & Hasezawa, S. Quantification and cluster analysis of

actin cytoskeletal structures in plant cells: role of actin bundling in stomatal movement during

diurnal cycles in Arabidopsis guard cells. *The Plant Journal* **61**, 156–165 (2010).

18. Otsu, N. A Threshold Selection Method from Gray-Level Histograms. *IEEE Transactions on*

Systems, Man, and Cybernetics **9**, 62–66 (1979).

19. Niblack, W. *An Introduction to Digital Image Processing*. (Prentice Hall, 1985).

20. Higaki, T., Akita, K. & Katoh, K. Coefficient of variation as an image-intensity metric for cytoskeleton

bundling. *Scientific Reports* **10**, 22187 (2020).

21. Lee, T. C., Kashyap, R. L. & Chu, C. N. Building Skeleton Models via 3-D Medial Surface Axis

Thinning Algorithms. *CVGIP: Graphical Models and Image Processing* **56**, 462–478 (1994).

22. Desbrun, M., Meyer, M., Schröder, P. & Barr, A. H. Implicit fairing of irregular meshes using

diffusion and curvature flow. in *Proceedings of the 26th annual conference on Computer graphics*

and interactive techniques - SIGGRAPH '99 317–324 (ACM Press, 1999).

doi:10.1145/311535.311576.

23. Scharr, H. *Optimal operators in digital image processing*. (2000).

24. Tanaka, K. *et al.* Structural basis for cofilin binding and actin filament disassembly. *Nat Commun* **9**,

1860 (2018).

25. Nunez-Iglesias, J., Blanch, A. J., Looker, O., Dixon, M. W. & Tilley, L. A new Python library to

analyse skeleton images confirms malaria parasite remodelling of the red blood cell membrane

skeleton. *PeerJ* **6**, e4312 (2018).

26. Boudaoud, A. *et al.* FibrilTool, an ImageJ plug-in to quantify fibrillar structures in raw microscopy

images. *Nature Protocols* **9**, 457–463 (2014).

27. Henty-Ridilla, J. L., Li, J., Day, B. & Staiger, C. J. ACTIN DEPOLYMERIZING FACTOR4 Regulates

Actin Dynamics during Innate Immune Signaling in *Arabidopsis*. *Plant Cell* **26**, 340–352 (2014).

28. Zack, G. W., Rogers, W. E. & Latt, S. A. Automatic measurement of sister chromatid exchange

frequency. *J Histochem Cytochem.* **25**, 741–753 (1977).

29. Li, C. H. & Tam, P. K. S. An iterative algorithm for minimum cross entropy thresholding. *Pattern*

Recognition Letters **19**, 771–776 (1998).

30. Jui-Cheng Yen, Fu-Juay Chang, & Shyang Chang. A new criterion for automatic multilevel
thresholding. *IEEE Transactions on Image Processing* **4**, 370–378 (1995).

31. Sauvola, J. & Pietikäinen, M. Adaptive document image binarization. *Pattern Recognition* **33**, 225–
236 (2000).

32. Henty-Ridilla, J. L. *et al.* The Plant Actin Cytoskeleton Responds to Signals from Microbe-
Associated Molecular Patterns. *PLOS Pathogens* **9**, e1003290 (2013).

33. Li, J., Cao, L. & Staiger, C. J. Capping Protein Modulates Actin Remodeling in Response to
Reactive Oxygen Species during Plant Innate Immunity. *Plant Physiology* **173**, 1125–1136 (2017).

34. Lu, Y.-J. *et al.* Arabidopsis calcium-dependent protein kinase 3 regulates actin cytoskeleton
organization and immunity. *Nature Communications* **11**, 6234 (2020).

35. Guo, M., Kim, P., Li, G., Elowsky, C. G. & Alfano, J. R. A Bacterial Effector Co-opts Calmodulin to
Target the Plant Microtubule Network. *Cell Host & Microbe* **19**, 67–78 (2016).

36. Kang, Y. *et al.* HopW1 from Pseudomonas syringae Disrupts the Actin Cytoskeleton to Promote
Virulence in Arabidopsis. *PLoS Pathog* **10**, e1004232 (2014).

37. Shimono, M. *et al.* The *Pseudomonas syringae* Type III Effector HopG1 Induces Actin Remodeling
to Promote Symptom Development and Susceptibility during Infection. *Plant Physiol.* **171**, 2239–
2255 (2016).

38. Faulkner, C. *et al.* An automated quantitative image analysis tool for the identification of
microtubule patterns in plants. *Traffic* **18**, 683–693 (2017).

39. Lu, Y.-J. & Day, B. Quantitative Evaluation of Plant Actin Cytoskeletal Organization During Immune
Signaling. in *Plant Pattern Recognition Receptors: Methods and Protocols* (eds. Shan, L. & He, P.)
207–221 (Springer, 2017). doi:10.1007/978-1-4939-6859-6_17.

Acknowledgements

We would like to thank Dr. Yi-Ju Lu (Michigan State University), and Dr. Silke Robatzek (Ludwig-
Maximilians University - Munich) for generously providing raw images of actin and tubulin for
experimental analysis and proof-of-concept analysis, respectively. We would like to thank Dr. Yi-Ju Lu
and Dr. Masaki Shimono for providing individual MGT evaluations for our experiments. We would like to
thank Rongzi Liu (University of Florida) for providing critical advice on statistical analysis and modeling.
We would like to thank Dr. Richard Neubig (MSU) and Dr. Noel Day (Zoetis) for critical advice on
manuscript preparation. We would like to thank Sarah Khoja for improving the reading experience of
ILEE_CSK documentation website. Research in the laboratory of B.D. was supported by grants from the
National Science Foundation (MCB-1953014) and the National Institutes of General Medical Sciences

(1R01GM125743). Research in the laboratory of Y.T. is supported by grant from National Science
Foundation (III-1900473).

1004

Author contributions

1006 P.L., Z.Z., B.D., and Y.T. participated in the design and conception of ILEE pipeline and experiments.
1007 P.L. and Z.Z. developed the ILEE algorithm and ILEE_CSK library. P.L., Z.Z., and B.M.F. performed the
1008 experiments. P.L., Z.Z., B.D., and Y.T. analyzed and interpreted the data. P.L. and B.D. and wrote the
1009 manuscript. All authors actively edited and agreed on the manuscript before submission.

1010

Competing interests

1012 The authors declare no competing interests.

1013

Supplemental information

1015 - S1. Supplemental Figures.pptx

1016

Figures

Fig. 1: ILEE pipeline and the demonstration of cytoskeletal indices. a, ILEE is an adaptive local thresholding approach that
applies to both 2D and 3D data structures, with an output of 12 cytoskeletal indices. Please note that *total connected element*,
total branch, and *total node* (marked "*") are not normalized indices for direct biological interpretation. **b**, Schematic diagram of
the ILEE algorithm. ILEE requires a sample image with enhanced edge gradients, a computed gradient threshold, and an implicit
Laplacian smoothing coefficient, K_2, to generate a binary image and skeletonized image for index computation. Z-axis maximum
projection (red box) is only conducted in the 2D mode. **c**, Visualized demonstration of ILEE performance. The raw data, binary
image generated by ILEE, and demonstration of computed features by both 2D and 3D modes are shown. TDT, total distance
transformation map, used to compute all diameter indices; skeleton NCE, non-connected elements of skeleton image with each
element by different colors, used to estimate severing activity; branching, skeleton image with each branch in different color and
each node by a black cross; anisotropy, local anisotropy level, shown as length of red lines, and direction of the first eigenvector,
shown as the direction of red lines. Scale bar = 20μm.
1029

1030

1031   **Fig. 2: NNES global thresholding. a,** The conceptual decomposition of a confocal fluorescence image of the cytoskeleton. An
1032   Arabidopsis leaf confocal microscopic image of actin, as an example of the eukaryotic cytoskeleton, comprises of 3 components:
1033   ground noise, the mechanical noise of the sensor regardless of the true fluorescence signal, diffraction light, the unavoidable
1034   diffraction signal of fluorescence component around, and true actin signal. They correspond to the noise filtered by coarse
1035   background, noise additionally filtered by ILEE, and segmented actin components in the algorithm. **b,** The scheme and
1036   demonstration of NNES. The curve reflects the NNE (negative non-connected component) count when certain global
1037   thresholding is applied to the raw images of 30 randomly selected samples. Their extremely smooth shape makes it easy to
1038   detect the peak as a feature value as the input for the coarse background estimation model. The demonstration of the filtered
1039   background at positions (i), (ii), (iii), and (iv) are shown above, where (iii) is adopted. The black area surrounded by colored area
1040   is the foreground information to be further processed.
1041

Fig. 3: Cytoskeletal identification by ILEE. a, Visual demonstration and summarized mathematical process of ILEE. On the left, the visualized intermediate images of ILEE process are presented; on the right, an abbreviated mathematical process of ILEE is shown (grey, image pre-processing; blue, ILEE in a narrow sense; yellow, post-processing; also see *Method* for the detailed computational algorithm). **b**, The value of implicit Laplacian smoothing coefficient K influences ILEE performance. When K is small (e.g., 2.5), the rendering of faint and thin filaments is accurate, but the edge of thick filaments tends to be omitted. Conversely, when K is large (e.g., 200), rendering of thick filaments is accurate but thin and faint filaments are omitted. A solution is applied by creating a full outer join threshold image by a fixed $K_1 = 2.5$ and an estimated universal K_2 for an entire biological batch of samples. **c**, K_2 estimation model. A regression model to compute any universal K_2 for a given sample batch (see Supplemental Figure 8 for detail). To augment the training samples, 7 images with manually portrayed binary ground truth are interpolated by a new resolution of different folds to the original into 42 samples (shown as single dots) that cover the general range of actin thickness. A non-linear regression estimation model is trained using the highest 5% DT (distance transform) values in the ground truth (representing filament thickness) and an anticipated K_2 with a specific satisfactory capability to detect thick filament (see Supplemental Figure 8 for detail). In the standard ILEE pipeline, the mean of top 5% DT of all input samples will be calculated by Niblack thresholding to obtain the universal K_2 for ILEE.

1059

Fig. 4: ILEE shows superior accuracy and stability over classic thresholding approaches. The manually portrayed binary ground truth of 7 visually diverse samples were compared with binary images rendered by ILEE, MGT, four global thresholding algorithms (Otsu, Triangle, Li, Yan), and two local thresholding algorithms (Niblack and Sauvola). **a**, Visualized comparison of

ILEE versus other approaches. Pixels with different colors are defined as green: match of rendered binary image and ground truth; blue: pseudo-negative, the pixels omitted by the algorithm; red, noise-like pseudo-positive, the pixels that are rendered by the algorithm but not in the ground truth without a shape of filament; brown, actin-like pseudo-positive, the pseudo-positive pixels within a filament-like component, which cannot be judged by high confidence. ILEE has the most accurate cytoskeleton segmentation. **b**, Quantitative comparison of pixel rendering demonstrated that ILEE has superior accuracy and stability across diverse samples. ILEE has the highest match rate with low and stable error rate; MGT and Li also have acceptable performance. **c**, Comparison of distribution of distance transformation error. Single pixel errors of all 7 samples were merged and summarized as a violin plot. Red dashed line indicates no error, or results identical to the ground truth. ILEE has a symmetric and centralized distribution, indicating an accurate and unbiased filament segmentation. **d**, ILEE has accurate and stable computation of cytoskeletal indices. Nine biologically-interpretable cytoskeleton indices computed using binary images rendered by different algorithms were compared with the ground truth. The index values were normalized to the fold of ground truth. Red dashed line indicates 1-fold, or identical to the ground truth. Additional results are shown in Supplemental Figure 10-13.

a

| Original 3D images | Simulation model | Artificial 3D images with ground truth | Comparison: 2D ILEE vs 3D |

b

Result (Relative to ground truth)	Ground truth	3D ILEE	Projected ground truth	2D ILEE
Filament coverage (by voxel)	100%	108.0±14.4%	10.7±0.7%	8.0±0.7%
Filament coverage (by length)	100%	105.1±11.6%	59.4±4.5%	42.0±4.7%
Match rate	100%	81.5±13.0%	NC	NC
Pseudopositive rate	0	26.5±7.1%	NC	NC
Pseudonegative rate	0	2.9±1.4%	NC	NC
Occupancy	100%	127.7±11.6%	1017.0±67.6% (!)	766.8±71.5% (!)
Linear density	100%	106.6±11.6%	1792.9±121.5% (!)	1218.2±135.6% (!)
Brightness Skewness	100%	103.6±3.2%	100.4±4.3%	96.2±5.1%
Brightness CV	100%	109.3±3.3%	99.8±2.9%	91.3±3.6%
Diameter (TDT)	100%	109.0±1.4%	144.7±6.9%	115.6±3.8%
Diameter (SDT)	100%	116.3±2.6%	128±4.0%	117.5±4.3%
Static severing activity	100%	67.7±8.0%	43.3±8.0%	25.6±3.8%
Static branching activity	100%	97.1±7.0%	137.1±4.1%	91.3±0.7%
Anisotropy	100%	97.4±9.4%	34.1±8.0% (!)	58.5±10.2% (!)

c

1075

1076

1077

Fig. 5: The 3D mode of ILEE renders cytoskeleton indices at much higher accuracy than the 2D mode, due to advanced
data structure. We utilized image samples (N = 31, 25*800*800, Fig.2) to train an actin image simulation model that generates
31 artificial 3D actin images with ground truth. By comparing the performance of 3D and 2D ILEE over the artificial images, it
was demonstrated that the 3D mode displays high fidelity to the ground truth while the 2D mode suffered from loss of information
and systemic bias. **a**, The general experimental scheme. The raw 3D images are analyzed to train an actin image simulation
model which uses statistical approaches to mimic the genuine actin filament, diffraction noise, and ground noise. Then 31
artificial actin images are regenerated with specific ground truth in terms of both actin segmentation and cytoskeleton indices.
The 3D and 2D ILEE were applied to the artificial images, whose results were then compared to their corresponding ground
truth. **b**, a table comparing the absolute accuracy of ILEE by the 3D and 2D mode. It was demonstrated that 2D mode losses
~90% of pixels(voxels) and ~50% of the total length of actin samples, and is less accurate for most of the indices. Exclamation
mark (!): contrast difference due to different definitions of dimensional space in their units; not rigorously comparable. **c**, the
linear correlation levels between ILEE results and the ground truth for all indices, by 2D and 3D mode. The absolute values of
each index of the ground truth, 3D ILEE, and 2D ILEE are normalized by making their relative folds to the minimum data point
of each group. A higher correlation coefficient r indicates a better capability to differentiate indices at different levels relatively,
regardless of their absolute fidelity. The 3D mode has a generally better capability of differentiating high and low values.

1093

1094
Fig. 6: ILEE library enables the discovery of actin dynamic features of bacteria infected leaf tissue. Leaves of the
Arabidopsis actin marker line Col-0/GFP-fABD2 were inoculated with mock or a virulent bacterial pathogen, *Pseudomonas*
syringae pv. tomato DC3000 with empty vector (EV, identical to wild type); images (*n* = 28 for mock, *n* = 31 for EV) of epidermal

cells were captured by laser scanning confocal microscopy at 24 hours post-inoculation (hpi). MGT and ILEE were applied to
generate binary images and all indices were computed using ILEE_CSK. Double-blinded samples were provided to 3 operators
(experienced researchers) OA, OB, and OC for comparison. OA additionally provides data using a universal threshold (OA-fix).
a, Experimental schematic diagram. **b**, Corelative comparison of MGTs of individual samples determined by different operators.
A very low correlation between each pair of operators indicates MGT has an increased risk of bias and is potentially inaccurate.
c, d, e, f, g, h, l, m, n, Output indices *occupancy*, *linear density*, *skewness*, *CV*, *diameter_TDT*, *diameter_SDT*, *static severing*
activity, *static branching activity*, and *local anisotropy*, respectively. Blue, mock; orange, EV. Multiple comparisons are conducted
using a t-test without family-wise error correction (because image samples themselves are ground truth for all methods). Groups
without overlapping letters have p_value lower than 0.05. For **n**, a visual illustration of the concept of local anisotropy is attached,
where each red line segment points to the "averaged" direction of actin growth in the local area and the length shows the intensity
of consistency of the direction. i**, j, k,** Comparisons of different indices in bundling class. *Skewness* and *CV* have medium-weak
correlation; *diameter_TDT* and *diameter_SDT* have strong correlation; *diameter_SDT*, as a representative of direct indicator,
and *CV*, as a representative of indirect indicator, have literally no correlation.
1111