Kinesin-3 mediated delivery of presynaptic neurexin stabilizes growing dendritic spines and postsynaptic components *in vivo*

Devyn Oliver[1], Shankar Ramachandran[1], Alison Philbrook[1], Christopher M. Lambert[1], Ken C. Q. Nguyen[2], David H. Hall[2], *Michael M. Francis[1]

[1]Department of Neurobiology, University of Massachusetts Medical School, Worcester, MA 01605

[2] Department of Neuroscience, Albert Einstein College of Medicine, Bronx, NY 10461

 *Corresponding author: michael.francis@umassmed.edu

Key words:

dendritic spines, neurexin, kinesin, *C. elegans,* synapse development, synaptic maintenance, active zone, protein trafficking

Contributions Summary

D.O. generated strains, transgenic lines, molecular constructs, confocal and electron microscopy images, and data analysis. S.R. performed all calcium imaging experiments and analysis. C.M.L. generated destination vectors and *nrx-1* constructs. D.H.H. and K.C.Q.N. aided in electron microscopy experiments and analysis. M.M.F and D.O designed and interpreted results of all experiments and wrote the manuscript.

Acknowledgements

Nematode strains were provided by the *Caenorhabditis* Genetics Center which is funded by the NIH National Center for Research Resources. Thank you to Dori Schafer for access to 3D rendering Imaris software. Thank you to Dirk Albrecht's laboratory for advice with hydrogel immobilization. Thank you to the laboratories of: Dong Yan, Kang Shen, Andres Maricq, Michael Nonet, Peri Kurshan, and Gert Jansen for providing reagents. Thank you to William Joyce and Michael Gorczyca for technical assistance. Thank you to Steve Cook for help with analysis of electron micrographs and John White for use of N2U electron micrographs. Lastly, we thank members of the Francis laboratory for manuscript comments.

Funding

This research was supported by NIH NINDS RO1NS064263 (MMF), R21NS101649 (MMF), NIH OD 01943 (DHH), and F31NS103365 (DO). The Philips CM10 TEM electron microscope used in this study was acquired through a NIH Shared Instrumentation Grant (1S10OD016214-01A1) at Albert Einstein College of Medicine.

Abstract

A high degree of cell and circuit-specific regulation has complicated efforts to precisely define roles for synaptic adhesion proteins in establishing circuit connectivity. Here, we take advantage of the strengths of *C. elegans* for cell-specific analyses to investigate molecular coordination of pre- and postsynaptic development. We show that developing dendritic spines emerge from the dendrites of wild type GABAergic motor neurons following the localization of active zone proteins and the formation of immature synaptic vesicle assemblies in presynaptic terminals. Similarly, clusters of postsynaptic receptors and F-actin are visible in GABAergic dendrites prior to spine outgrowth. Surprisingly, these developmental processes occur without a requirement for synaptic activity. Likewise, the initial stages of spine outgrowth and receptor clustering are not altered by deletion of the *C. elegans* ortholog of the transsynaptic adhesion protein, neurexin/NRX-1. Over time, however, dendritic spines and postsynaptic receptor clusters are destabilized in the absence of presynaptic NRX-1/neurexin and collapse prior to adulthood. The kinesin-3 family member, UNC-104, delivers NRX-1 to presynaptic terminals and ongoing UNC-104 delivery is required into adulthood for the maintenance of postsynaptic structure. Our findings provide novel insights into the temporal order of synapse formation events *in vivo* and demonstrate a requirement for transsynaptic adhesion in stabilizing mature circuit connectivity.

Introduction

The capabilities of neural circuits to perform specific functions arise from the patterns of synaptic connections between their partnering neurons. The organization of these connections is circuit-specific and established through a complex process that involves the coordinated assembly and maturation of specialized pre- and postsynaptic structures on appropriate partnering neurons, and their maintenance in the mature nervous system. We now have a generalized understanding of synapse structure. Active zone (AZ) proteins and neurotransmitter-filled synaptic vesicles position near the presynaptic membrane for rapid release while neurotransmitter receptors are clustered at high density in apposition to these sites in order to ensure the fidelity of synaptic communication. Genetic studies have identifed numerous mutations that alter circuit connectivity or affect the overall structural organization of synapses (Hendi et al., 2019) (Zhen & Jin, 1999) (Richmond et al., 1999). However, for many of the synapse-associated proteins affected by these mutations, we do not yet have a mechanistic understanding of their roles in establishing synaptic connections, or how their disruption may lead to alterations in synapse development and structure. Gaining an enhanced understanding of the sequence of events involved in synapse assembly, maturation, and maintenance, and their relative timing *in vivo* is critical for addressing these questions.

Several prior studies have examined molecular events during synaptogenesis. *In vivo* studies in *C. elegans* and *Drosophila* have largely focused on the formation of the presynaptic active zone. Collectively, these studies provide compelling evidence for a model where active zone assembly occurs sequentially. While there is some variability across synapse type, the early stages of this process are generally organized by the highly conserved synaptic scaffolds SYD-2/Liprin-α and SYD-1/Rho GTPase, which then recruit additional key conserved AZ proteins such as ELKS-1/Bruchpilot, Piccolo family members, and UNC-10/RIM for subsequent stages of assembly, including clustering of Ca^{2+} channels and the recruitment of synaptic

vesicles (Zhen & Jin, 1999) (Bohme et al., 2016) (Fouquet et al., 2009) (Patel et al., 2006) (Dai et al., 2006). Remarkably, nascent AZs can assemble quite rapidly (within minutes) but then often undergo a more extended period of maturation that can last for several hours (Nagerl et al., 2007) (Lipton et al., 2018) (Fouquet et al., 2009) (McDonald et al., 2020). The relationship of postsynaptic development to presynaptic assembly remains less clear. Live imaging studies of cultured mouse hippocampal neurons have suggested that the assembly of presynaptic components precedes recruitment of postsynaptic receptors and scaffolds (Friedman et al., 2000). More recently, 2-photon imaging of organotypic slice cultures from rat hippocampus showed that new spines accumulate glutamate receptors concurrently with their growth and are competent to participate in transmission within a few hours after outgrowth (Zito et al., 2009). Comparatively few *in vivo* studies have investigated coordinated pre- and postsynaptic development and significant questions remain about molecular links between these processes.

Evolutionarily conserved synaptic adhesion proteins, such as neurexins, are prime candidates for coordinating pre- and postsynaptic events during synaptogenesis. Their importance is underscored by the fact that neurexin alterations are associated with cognitive disease (Sudhof, 2008). For example, mutations in human Neurexin1 are associated with several neuropsychiatric disorders, including schizophrenia and autism spectrum disorders (Kasem et al., 2018). Neurexins are typically localized presynaptically and linked to postsynaptic binding partners such as dystroglycans, LRRTMs, neuroligins, and cerebellins, through extracellular laminin and EGF-like repeats to establish transsynaptic connections. Neurexin mediated transsynaptic signaling has been implicated in key aspects of synapse development and function. For example, loss of presynaptic neurexin at the fly neuromuscular junction increases the length of the presynaptic density, and also alters the size and molecular composition of apposed muscle glutamate receptor clusters (J. Li et al., 2007) (Owald et al., 2012). The broad brain distribution of neurexins and the many neurexin isoform variants in the

nervous system enable complex, cell-type specific functions for neurexins at synapses (Missler & Sudhof, 1998) (Sudhof, 2017). For example, conditional deletion of mouse neurexins revealed strikingly different functions across synapses formed by cortical inhibitory interneurons and those formed by cerebellar climbing fibers onto Purkinje neurons (L. Y. Chen et al., 2017). Genetic tools that enable cell-type specific analysis are therefore critical for uncovering precise functional roles of synaptic adhesion proteins within the context of individual neural circuits. Importantly, molecular mechanisms for their trafficking and delivery to synapses also remain incompletely defined.

We previously identified finger-like dendritic protrusions from the dendritic processes of *C. elegans* DD GABAergic motor neurons **(Figure 1A)** (Philbrook et al., 2018). Characterization of these structures by our laboratory and others pointed towards the idea that they receive synaptic input from presynaptic cholinergic motor neurons and serve analogous roles to dendritic spines in the mammalian brain (J. G. White et al., 1976) (Cuentas-Condori et al., 2019). Deletion of *nrx-1*, the sole *C. elegans* ortholog of neurexin, disrupts these dendritic spines and impairs proper localization of cholinergic receptor clusters to postsynaptic sites on GABAergic dendrites (Philbrook et al., 2018). Here we define the order and timing of pre- versus postsynaptic development at spine-associated synapses and elucidate the role of neurexin in the coordination of these processes *in vivo*. We find that clusters of both presynaptic proteins and postsynaptic receptors are clearly visible prior to spine outgrowth. In the absence of neurexin, immature spines and receptor clusters form initially but then disappear, indicating that neurexin is required for synapse stabilization and maturation rather than synaptogenesis. These maintenance and maturation processes are supported by kinesin-3/UNC-104 motor dependent transport of NRX-1 to presynaptic terminals. Together, our results suggest that early and ongoing axonal delivery of neurexin to sites of synapse formation is critical for stabilization of postsynaptic structures. Our findings cleanly dissect a key role for neurexin mediated adhesion

in sculpting connectivity, specifically in promoting the stabilization and maturation of growing synapses and spines.

Results

We initially explored morphological features of mature spines using 3D rendering of spines from confocal imaging. We identified several morphological classes **(Figure 1B)** based on the number of spine heads and the ratio between the length of the spine to the width of the spine head and spine neck. We found that a majority of DD spines can be categorized into four morphological classes (percentage ± SD): mushroom (5 ± 1%), branched (5 ± 1%), stubby (73 ± 3%), and thin (15 ± 1%). The morphological diversity we observed by light microscopy was also evident in ultrastructural studies using serial electron microscopy (**Figure S1.1).** Clusters of small synaptic vesicles were also evident in neurites (of cholinergic VA/VB neurons, presumably) immediately adjacent to the tips of DD spines, indicating putative sites of synaptic input.

To explore how the neuronal cytoskeleton may contribute to spine morphology, we examined the localization of F-actin and tubulin in DD dendrites using DD neuron-specific expression GFP::UtrCH (GFP fused to the Utrophin calponin homology domain) (Chia et al., 2014) and TBA-1::GFP (GFP-tagged α-tubulin) (Yan et al., 2013) respectively. We found that the two markers occupy distinct territories. Coexpression studies showed that F-actin is strongly enriched within dendritic spines **(Figure 1C-E)**, while tubulin occupies the dendritic shaft **(Figure 1F,G)** with more variable localization to the spine base (**Figure S1.2**). Previous findings for vertebrate neurons indicate a similar segregation of actin and tubulin to the spines and shaft of the dendrite respectively (Gu et al., 2008) (Hu et al., 2008) (Jaworski et al., 2009).

DD spines are positioned immediately adjacent to the axon terminals of cholinergic neurons that are predicted to be their presynaptic partners by electron microscopy (**Figure S1.1**) (J. G. White et al., 1976) (J. G. White et al., 1986) (Cuentas-Condori et al., 2019). We previously showed stimulation of cholinergic motor neurons elicited Ca^{2+} responses in

GABAergic motor neurons (Philbrook et al., 2018). To confirm that spines on DD GABAergic motor neurons represent sites of functional synaptic contacts, we asked whether stimulation of presynaptic cholinergic neurons was sufficient to elicit calcium responses in DD spines. We performed *in vivo* calcium imaging of evoked responses in spines using combined expression of a membrane-associated GCaMP6f calcium sensor in DD GABAergic neurons and a red shifted channelrhodopsin, Chrimson, in cholinergic neurons (Philbrook et al., 2018) (**Figure 1H, I, J) (Figure S1.3)**. Following 5 seconds of baseline recording (488 nm, 100 ms exposure), we measured calcium responses to presynaptic depolarization (5 s, 625 nm, 30 mW/cm^2) (**Figure 1I**). We noted significant fluorescence increases in spines that were timed with the onset of stimulation and were not present in the absence of retinal. These typically occurred simultaneously across multiple spines and returned to baseline within 5 s following stimulation. Importantly, mutation of the cholinergic vesicular transporter/VAChT *unc-17*, reduced evoked calcium responses by 81%, indicating the Ca^{2+} responses we observed in dendrites were dependent on presynaptic acetylcholine release **(Figure 1H, I, J**). Our findings are consistent with those of another recent study (Cuentas-Condori et al., 2019), and indicate that spines on DD GABAergic neurons are primary sites of synaptic contact.

Mitochondria and other organelles are often localized near postsynaptic specializations in order to sustain synaptic function (Vos et al., 2010) (Karagas & Venkatachalam, 2019). To examine the distribution of organelles within DD dendrites and spines, we used fluorescent reporters labeling the Golgi apparatus (AMAN-2::GFP) (Edwards et al., 2013), mitochondria (Pre-Su9::GFP) (Sarasija & Norman, 2015), and endoplasmic reticulum (ER) (RFP::TRAM-1) (Rolls et al., 2002) (Edwards et al., 2013) (**Figure S1.4**). We found that Golgi is exclusively labeled in the DD cell bodies (**Figure S1.4A**). In contrast, mitochondria and rough ER are dispersed throughout the main dendritic shaft and near the base of spines (**Figure S1.4B-E**), perhaps suggesting involvement in spine function (Perez-Alvarez et al., 2020) (Z. Li et al.,

2004). Consistent with our light microscopy observations, we also observed mitochondria in the main dendritic process of DD GABAergic neurons near spines by electron microscopy (**Figure S1.1B,C**).

The development of synapses at DD GABAergic spines

Using the synapses at DD spines as a model, we next investigated the relative timing of pre- and postsynaptic events during synapse formation. *C. elegans* progress through four larval stages of development (L1-L4) prior to adulthood. DD neurons undergo a well-characterized program of synaptic remodeling such that the mature circuit organization is established during the transition from L1 to L2 stage (J. G. White et al., 1978). During this period of remodeling, newly born cholinergic neurons form new synaptic connections with the ventral dendrites of DD GABAergic neurons. This synaptic remodeling event offers a well-defined temporal window to investigate *de novo* formation of neuron-neuron synaptic connections *in vivo*. We focused on understanding the time course of spine formation relative to 2 key events in synaptogenesis: 1) the formation of presynaptic release sites, and 2) the clustering of postsynaptic receptors and F-actin **(Figure 2).**

The appearance of newly born ventral cholinergic motor neuron cell bodies and processes progressed anteriorly to posteriorly starting roughly 16 hours after hatch, consistent with the birth and integration of cholinergic neurons during this time frame (**Figure S2.1**) (Kurup & Jin, 2016). We analyzed the distribution of pre- and postsynaptic markers at timepoints before (12 hours after hatch), during (16, 18, 20 hours after hatch) and following (24, 32 hours after hatch, L4 stage/42-50 hours) DD synaptic remodeling **(Figures 2, S2.1, and S2.2)**. We first assessed the formation of the active zone and presynaptic specialization by analyzing clustering of the Piccolo-like active zone protein, CLA-1, (GFP::CLA-1e) and the synaptic vesicle-associated protein synaptobrevin/SNB-1 (SNB-1::GFP) **(Figure 2A, D-E)** (Xuan et al., 2017).

Overall, we found CLA-1 is more discretely localized to putative synapses than SNB-1, both at earlier timepoints and throughout development, consistent with its specific localization to active zones. Immediately prior to remodeling (~12 hours after hatch), we did not observe significant localization of either presynaptic CLA-1 or SNB-1 adjacent to the DD process. Shortly after (16 hours after hatch), we noted the initial appearance of SNB-1- and CLA-1-associated fluorescence in presynaptic cholinergic processes. Over the next 4 hours (20 hours after hatch) individual CLA-1 clusters were more clearly distinguishable, suggesting CLA-1 association with developing active zone structures **(Figure 2A, D, L)**. Synaptic vesicle fluorescence was initially diffuse at 16 hours after hatch and became more clearly organized into discrete puncta over a similar time course to CLA-1 **(Figure 2A, E, L).** We did not observe the emergence of DD dendritic spines until 24-32 hours after hatch, well after initial active zone formation and recruitment of synaptic vesicles (**Figure S2A, C-E**). Spines continued to mature through L4 stage, increasing in both length and number (**Figure 2.2C**).

We next examined the clustering of postsynaptic acetylcholine receptors in DD dendrites (ACR-12::GFP). Prior to remodeling (12 hours after hatch), we did not observe detectable levels of cholinergic receptors in the DD process. Surprisingly, we noted immature receptor clusters faintly visible in the dendritic shaft by 16 hours after hatch, well prior to the emergence of dendritic spines **(Figure 2B, F, L)**. These receptor clusters increased in number and redistributed towards the tips of growing spines by 32 hr **(Figure 2B, F, L)**. We found a similar developmental trend using DD neuron-specific labeling of the LEV-10 transmembrane auxiliary protein (Gally et al., 2004), previously shown to concentrate in spines (He et al., 2019) (**Figure S2.3**). By L4 stage, receptor and LEV-10 clusters are clearly visible at the tips of mature spines. Together, our analysis indicates that the initial stages of development of both pre- (CLA-1 and SNB-1 clusters) and postsynaptic (AChR and LEV-10 clusters) structures occur prior to spine outgrowth, raising questions about how these initial processes may be regulated.

To begin to address this question, we analyzed the distribution of F-actin in dendrites of the DD neurons during synapse formation **(Figure 2B, G-L)**. We noted that clusters of F-actin were evident in the ventral DD process prior to the completion of synaptic remodeling (12 hours after hatch), prior to presynaptic CLA-1 and synaptic vesicle accumulation, and before the clustering of postsynaptic receptors. Dendritic F-actin-based structures became more abundant coincident with increases in the number of presynaptic CLA-1 and postsynaptic receptor clusters. To investigate this process in real-time, we used live imaging to examine the dynamics of postsynaptic F-actin (GFP::UtrCH) in the developing DD dendrite. We found that F-actin is highly dynamic during early developmental stages (16-20 hours after hatch) compared to L4 stage where the circuit has completed maturation **(Figure S2.4, Video 1, and Video 2).** In young animals, GFP::UtrCH clusters often shuttled out of the cell body to the main dendritic process, perhaps indicating delivery of postsynaptic F-actin to sites of postsynaptic assembly. By 24 hours after hatch, we observed clear co-localization of F-actin with newly formed AChR clusters in the dendritic shaft that are positioned for developmental translocation into growing dendritic spines. In mature animals, AChR clusters are stably sequestered at the tips of spines, while F-actin occupies the spine head and neck **(Figure 2H-K)**.

Dendritic spines form in the absence of presynaptic activity

Our above analysis indicated that the localization of synaptic vesicles and active zone proteins, such as CLA-1, occur prior to spine outgrowth. In the rodent brain, spine morphogenesis is clearly regulated by presynaptic activity (Sala & Segal, 2014) (Saneyoshi et al., 2010) (Engert & Bonhoeffer, 1999) (De Roo et al., 2008), but recent evidence suggests initial spine outgrowth may proceed independently of synaptic activity (Lu et al., 2013) (Sigler et al., 2017) (Sando et al., 2017). We therefore next asked whether presynaptic cholinergic activity is important for the formation of DD dendritic spines. To address this question, we analyzed the

number of dendritic spines (P*flp-13*::mCherry or P*flp-13*::myrGFP) and cholinergic receptors (P*flp-13*::ACR-12::GFP) at L4 stage in strains carrying mutations that affect various aspects of synaptic function and neuronal excitability **(Table 1, Figure S3.1)**.

Surprisingly, the abundance of spines and receptor clusters at L4 stage were not significantly affected by disruption of ACh synthesis in cholinergic motor neurons (mutation of the vesicular acetylcholine transporter *unc-17*) or by a strong reduction in synaptic vesicle exocytosis (mutation of the syntaxin binding protein *unc-18*). Similarly, loss-of-function mutations in genes required for dense core vesicle release (CAPS/*unc-31*) and Ca^{2+} signaling (VGCC/*unc-2,* CaM Kinase II/*unc-43*) did not significantly reduce the number of spines or receptor clusters in the mature circuit **(Table 1, Figure S3.1)**. Our data indicate that alterations in synaptic activity do not affect the abundance of L4 stage dendritic spines. However, we cannot exclude the possibility of developmental delays in spine formation (Cuentas-Condori et al., 2019). Notably, many of the mutations we examined significantly altered spine length and the size of receptor clusters **(Table 1)**, suggesting that activity may influence aspects of spine morphology and post-synaptic structure. Together, our findings point toward a model where initial spine formation and postsynaptic development proceeds without a strong requirement for synaptic activity, while subsequent spine morphogenesis may be affected.

Presynaptic NRX-1 stabilizes postsynaptic components

In previous work, we found that mutations in *nrx-1,* the sole *C. elegans* ortholog of neurexin, eliminate DD dendritic spines and disrupt cholinergic receptor clusters in the mature *C. elegans* circuit (Philbrook et al., 2018), indicating alternate activity-independent mechanisms may be critical for spine formation. Consistent with these findings, we found that mutations of *nrx-1* disrupted dendritic calcium responses evoked by presynaptic stimulation of cholinergic motor neurons **(Figure S3.2)**. To examine possible early involvement of presynaptic NRX-1 in

the establishment of spines we asked when NRX-1 first localizes to presynaptic terminals of cholinergic motor neurons. Axonal clusters of NRX-1::GFP are present by 16 hours after hatch (**Figure 3A, B**), approximately the same time at which assemblies of the presynaptic scaffold CLA-1 become visible (**Figure 2**). Diffuse synaptic vesicle material is also visible at this time **(Figure 2)** but has yet to be organized into discrete clusters.

The early arrival of NRX-1 to synaptic terminals could indicate a role in either initial synapse formation or subsequent stabilization and maturation. To distinguish between these possibilities, we analyzed the development of dendritic spines and cholinergic receptor clusters in *nrx-1(wy778)* null mutants. We quantified spine and AChR cluster number over a similar time course as previously completed for wild type. Surprisingly, we noted that immature spines and receptor clusters were clearly evident in *nrx-1* mutants during early development, albeit at slightly reduced density relative to wild type **(Figures 3C-F, S3.3)**. The density of spines in *nrx-1* mutants increased significantly over the next several hours until 24 hours after hatch (**Figure 3C,D).** However**,** after this time, spine density decreased dramatically such that *nrx-1* DD dendrites were almost completely devoid of spines by L4 stage, consistent with our previous work (Philbrook et al., 2018). Measurements of spine formation and disassembly in live imaging studies of wild type and *nrx-1* mutants offered further support for this conclusion. Mature wild type spines (L4) were remarkably stable over 1-2 hours of recording, but were more dynamic in the developing circuit (16-20 hrs after hatch) **(Figure S3.4, Videos 3-5)**. Developing spines in *nrx-1* mutants were also highly dynamic. Notably, heightened spine dynamics persisted into more advanced developmental stages for *nrx-1* mutants compared with wild type animals. For example, at 21-24 hours after hatch, almost 90% of wild type spines were stable over the recording period, while less than 50% of *nrx-1* mutant spines remained stable, indicating decreased stability in the absence of NRX-1.

Receptor clustering followed a similar trend to spine formation. The number and fluorescence intensity of ACR-12 clusters at 16 hours after hatch in *nrx-1(wy778)* animals (6.8 ± 0.8 clusters/25 μm) were similar to wild type (5.6 ± 0.7 clusters/25 μm) **(Figure 3E-F, S3.3)**. Within a few hours however, ACR-12 cluster number and intensity decreased significantly in *nrx-1* mutant dendrites, and remained low throughout the remainder of development **(Figure 3E-F, S3.3)**. Our time course analysis indicated that both spines and receptor clusters initially form normally in *nrx-1* mutants, but rapidly disappear as animals proceed through development in the absence of NRX-1. Remarkably, we found that the organization of dendritic F-actin was similarly affected by *nrx-1* deletion. GFP::UtrCH was arranged similarly in *nrx-1* mutants and wild type in early development (16 hrs after hatch), positioned in discrete clusters along the main dendritic process (**Figure 3G**). By L4 stage, wild type F-actin was almost entirely associated with dendritic spines. This organization was strikingly altered in L4 stage *nrx-1* mutants. GFP::UtrCH was diffusely localized along the length of *nrx-1* mutant dendrites, indicating that presynaptic NRX-1 promotes dendritic F-actin assemblies (**Figure 3G**). Taken together, our results indicate that presynaptic NRX-1 is dispensable for the earliest stages of postsynaptic assembly and spine formation, but is critical for stabilizing dendritic spines and AChR clusters and promoting their maturation.

Kinesin-3/UNC-104 transports presynaptic neurexin to cholinergic terminals in order to stabilize postsynaptic components

To investigate this model further, we sought to determine how presynaptic neurexin is transported to active zones *in vivo*. Prior work showed that both synaptic vesicles and CLA-1 depend on the Kinesin-3 motor UNC-104 for their delivery to synapses (Hall & Hedgecock, 1991) (Xuan et al., 2017) **(Figures S4.1 and 4)**. In particular, disruption of *unc-104* causes an accumulation of synaptic vesicles within neuronal somata and a corresponding loss of synaptic vesicles within axons (Hall & Hedgecock, 1991) **(Figure S4.1).** Consistent with potential

involvement of UNC-104 in NRX-1 trafficking, we found that UNC-104 and NRX-1 are partially colocalized in cholinergic axons (**Figure S4.2**). *unc-104(e1265)* carries a D1497N mutation in the PH domain of UNC-104 that impairs cargo binding and leads to a severe reduction in UNC-104 abundance in axons (Kumar et al., 2010). NRX-1 endogenously tagged with GFP is strongly visible in neuronal processes within the nerve ring and nerve cords of wild type animals (**Figure S4.3**). In contrast, NRX-1::GFP clusters were strikingly decreased in the nerve cords of *unc-104(e1265)* mutants **(Figures 4A,C, S4.4)**. A similar deficit in axonal NRX-1::GFP localization was evident with specific cholinergic expression of NRX-1::GFP in *unc-104* mutants **(Figures 4B,C, S4.4)**. By contrast, NRX-1::GFP fluorescence was significantly increased (5.4-fold) in *unc-104* mutant cholinergic somata compared to wild type **(Figure S4.4)**. Accumulation of NRX-1 in cholinergic cell bodies, coupled with a decrease in axons, indicates a failure of NRX-1 transport in the absence of functional UNC-104. Consistent with this, cholinergic-specific expression of wild type *unc-104* in *unc-104* mutants restored axonal NRX-1::GFP localization, indicating a cell autonomous requirement **(Figure 4 B,C)**. In contrast, mutation of the Kinesin-1 motor *unc-116* did not produce significant accumulation of NRX-1::GFP in cholinergic somata and caused comparatively modest decreases in axonal NRX-1::GFP, demonstrating preferential involvement of Kinesin-3 for NRX-1 transport (**Figure S4.5**). Live imaging studies offered additional evidence that synaptic vesicle and NRX-1 trafficking may share a common dependence on UNC-104. We found that NRX-1::GFP trafficking events occurred with similar anterograde and retrograde velocities to SNB-1::GFP (labeling synaptic vesicles) trafficking events, though less frequent **(Figure S4.6, Videos 6 and 7)**. Prior studies of active zone protein transport have noted these events occur at reduced frequency compared with synaptic vesicle trafficking events (Lipton et al., 2018). Together, our analyses point to a specific requirement for the UNC-104/KIF1A motor in delivery of NRX-1 to presynaptic terminals.

The density of spines and AChR clusters were also severely reduced in L4 stage *unc-104(e1265)* mutants (**Figure 4 G-I**), similar to *nrx-1* mutants (**Figure 4J**). These effects were rescued by either native or cholinergic expression of wild type *unc-104* in *unc-104(e1265)* mutants, but not by GABA- or muscle-specific *unc-104* expression (**Figure 4 G-I**), demonstrating that presynaptic UNC-104-mediated transport is critical in cholinergic axons for postsynaptic spine development and receptor localization.

Notably, we observed that several other presynaptic AZ components also require UNC-104 for their delivery. In addition to CLA-1, transport of both ELKS-1/ELKS/CAST and UNC-10/RIM were severely disrupted by mutation of *unc-104*, indicating that Kinesin-3 cargo selection and transport is key for the synaptic delivery and assembly of several key active zone constituents in these neurons (**Figure 4C-F**). In contrast, the synaptic localization of these active zone proteins was not appreciably affected by *nrx-1* deletion (**Figure S4.7**). We next asked how a failure in delivery of AZ components may contribute to the severe reductions in spines and receptor clusters we observed in *unc-104* mutants. We compared postsynaptic organization in strains carrying deletion mutations in either *nrx-1, cla-1, unc-10* or *elks-1*. Remarkably, only mutation of *nrx-1* produced significant decreases in dendritic spines (**Figure 4J**), demonstrating that presynaptic ELKS, UNC-10 and CLA-1 are dispensable for postsynaptic development. Our results indicate a specific requirement for NRX-1 in the stabilization and maturation of postsynaptic structures and provide evidence that a failure in synaptic delivery of NRX-1 is a primary causal factor in the postsynaptic structural defects of *unc-104* mutants.

The stabilization of mature dendritic spines requires ongoing synaptic delivery of NRX-1

Our results suggest UNC-104 mediated transport positions NRX-1 at the presynaptic terminal in the early stages of synapse formation where it acts to stabilize growing postsynaptic structures, including spines and receptor clusters. We next sought to address whether there is

an ongoing requirement for presynaptic UNC-104-dependent transport in the stabilization of mature spines. To address this question, we used the previously characterized temperature-sensitive allele, *unc-104(ce782). unc-104(ce782)* animals carry a G105E missense mutation in the motor domain of UNC-104 (Edwards et al., 2015). When grown at permissive temperature (13.5°C), *unc-104(ce782)* axonal synaptic vesicle abundance and animal motility are modestly reduced compared to wild type. In contrast, synaptic vesicles are completely absent from *unc-104(ce782)* axons following growth at restrictive temperature (20-25°C), and animal motility is severely compromised within 12 hours of a shift to 23°C (Edwards et al., 2015). We raised *unc-104(ce782)* animals at the permissive temperature (13.5°C) to L4 stage **(Figure 5)**. We then shifted L4 stage animals to the restrictive temperature (25°C) for 16-20 hrs, and quantified NRX-1::GFP localization and spine density immediately following this shift. Axonal NRX-1::GFP clusters were strikingly decreased in *unc-104(ce782)* mutants subjected to the temperature shift compared with control animals subjected to the same shift or *unc-104(ce782)* mutants raised continuously at the permissive temperature **(Figure 5B,C)**. *unc-104(ce782)* mutants grown at the permissive temperature had a slightly reduced number of spines overall **(Figure 5E)**, compared with wild type. A shift to the restrictive temperature at L4 stage also produced a striking reduction in spine density for *unc-104(ce782)* mutants **(Figure 5D, E)**, but not for wild type animals subjected to the same temperature shift. We obtained similar results for *unc-104(ce782)* mutants using an earlier shift to the restrictive temperature (at L3 stage) **(Figure S5.1)**. Taken together, our findings indicate an ongoing requirement for UNC-104 transport that extends well beyond the period of initial synapse formation and spine outgrowth. Among the UNC-104 cargoes we investigated, only deletion of *nrx-1* produces a significant reduction in spine density at L4 stage. We therefore propose a model where ongoing UNC-104 delivery of presynaptic NRX-1 is critical for postsynaptic maturation and maintenance of mature spines.

Discussion

***C. elegans* GABAergic motor neurons are decorated with functional dendritic spines**

Dendritic spines are known to act as specialized sites for compartmentalizing neurotransmission and are widely observed across various classes of neurons in both vertebrates and invertebrates (Tonnesen & Nagerl, 2016). Though suggested by prior EM studies (J. G. White et al., 1976) (J. G. White et al., 1986), the presence of functional dendritic spines on *C. elegans* neurons had not been fully appreciated until recently (Philbrook et al., 2018) (Cuentas-Condori et al., 2019). Here we examined the developmental trajectory of postsynaptic structures (dendritic spines and cholinergic receptor clusters) located on GABAergic motor neurons relative to presynaptic release sites and showed that the stabilization of growing dendritic spines requires presynaptic NRX-1/neurexin.

Neural circuit organization and performance is dictated by the locations of synaptic connections and the identities of the interconnected neurons. Elucidation of the processes governing the construction of synapses and their coordination across presynaptic axons and postsynaptic dendrites are therefore fundamental for understanding circuit assembly and function. However, relatively few studies have monitored in real-time the coordinated assembly of pre- and postsynaptic specializations *in vivo*. Our work tackled this important question, taking advantage of the GABAergic DD spine model and the cellular precision offered by the *C. elegans* motor circuitry. We find that these dendritic spines share many of the hallmark features of vertebrate dendritic spines, but also have some key differences. **(1)** *C. elegans* spines have morphological similarities with mammalian spines and can be classified into comparable morphological categories. **(2)** F-actin is localized to spines while tubulin is mainly present in the dendritic shaft. F-actin is a major component of mammalian dendritic spines and is important for spine structural dynamics (reviewed in (Borovac et al., 2018)). **(3)** Mitochondria and rough

endoplasmic reticulum are positioned near spines, either in the spine neck or at the base of spines in the dendritic shaft. Dendritic organelles contribute to key processes involved in spine formation and spine metabolism (Liu & Shio, 2008). Ribosomes and mitochondria positioned near or within dendritic spines may aid in modifying dendritic spine activity, local protein synthesis, and act as synaptic ATP and Ca^{2+} sources (Steward & Levy, 1982) (Aakalu et al., 2001) (Chirillo et al., 2019) (Z. Li et al., 2004). **(4)** Presynaptic stimulation elicits calcium responses in spines on GABAergic dendrites. DD spines are located opposite cholinergic release sites and calcium responses are eliminated by mutation of the vesicular acetylcholine transporter. **(5)** Mature *C. elegans* spines display limited overt activity-dependent dynamics in wild type animals and appear comparatively stable relative to spines in rodents. This raises the important question of what is the functional role of spines in GABAergic neurons? Prior studies have provided evidence for compartmentalized calcium signals in *C. elegans* neurons (Hendricks et al., 2012). Therefore, one intriguing possibility is that, similar to the situation in mammals, spines serve to compartmentalize calcium and perhaps other biochemical signals in GABAergic neurons. Spines may also be required in order to achieve the unusual dyadic arrangement of synapses in the ventral nerve cord, where presynaptic specializations of cholinergic motor neurons are positioned for transmission onto both GABA motor neuron and muscle postsynaptic partners (Hall & Russell, 1991). In this case, GABAergic spines may have developed to intercept cholinergic release sites onto muscles, as suggested previously (J. White, 2018).

Ventrally-directed (VA/B) cholinergic neurons are born and integrated into the motor circuit post-embryonically, forming new synaptic connections with both ventral muscles and dorsally-directed (DD) GABAergic motor neurons. We found that F-actin is compartmentalized to discrete regions within the dendritic shafts of DD neurons very early in the development of the circuit (prior to the appearance of spines) and later becomes exclusively localized to dendritic

spines. F-actin labeling of spines is apparent as soon as they are detectable, suggesting that F-actin assemblies may participate in the earliest stages of spine development. Prior to remodeling of DD neurons, sites of GABA release are located on the ventral processes of DD neurons. As F-actin based structures are typically associated with release sites (Chia et al., 2012) (Meng et al., 2015), our finding that F-actin is clustered in ventral DD processes prior to remodeling may suggest that F-actin initially associated with release sites persists in the DD neurite through remodeling and is then repurposed for postsynaptic development, perhaps acting as a landmark or an actin-based mechanism for stimulating post-synaptic maturation.

Notably, we also observed punctate cholinergic receptor fluorescence in GABAergic dendrites several hours prior to the formation of dendritic spines. These immature receptor clusters colocalize with dendritic F-actin assemblies suggesting that receptors are trafficked into dendrites and positioned with F-actin to rapidly populate growing spines. Indeed, we observed that receptor clusters are visible in growing spines as soon as spines can be clearly resolved. This is consistent with recent findings from time-lapse 2-photon imaging of hippocampal organotypic slice cultures showing that receptor accumulation in spines occurs concurrently with spine outgrowth (Zito et al., 2009). Similarly, we found that synaptic material accumulates in presynaptic axons prior to the emergence of spines from dendritic processes. In particular, initial accumulations of the presynaptic scaffold CLA-1 and the synaptic vesicle marker synaptobrevin/SNB-1 were visible in the axon with a similar time-course to the appearance of dendritic cholinergic receptor clusters. Presynaptic material became more clearly localized to discrete puncta over the next 4-6 hours, occurring roughly coincident with receptor accumulation at the tips of growing spines. Notably, our findings do not support a strong requirement for presynaptic neurotransmitter release in spine formation. The abundance of spines was either mildly affected or unaffected across several mutant strains with severely impaired presynaptic function.

Transsynaptic NRX-1/Neurexin signaling stabilizes dendritic F-actin assemblies to promote postsynaptic maturation

Numerous studies across a variety of model systems have examined roles for neurexins in synapse development and function. However, no single consensus view of neurexin function has emerged. Instead, the function of neurexin at specific synapses are thought to be dictated by cellular and molecular context. Mammalian genomes encode three Nrxn genes that can each give rise to α-, β-, and γ-Nrxn isoforms. These isoforms share common intracellular and transmembrane regions but differ in their extracellular domains. α-Nrxn has six extracellular LNS domains interleaved with three EGF-like repeats. β-Nrxn has only one LNS domain, while γ-Nrxn lacks all identifiable extracellular domains. Conditional triple knockout of mouse neurexin indicated synapse loss in parvalbumin-positive prefrontal cortex interneurons, but not somatostatin-positive interneurons (L. Y. Chen et al., 2017). At mouse calyx synapses, conditional triple neurexin panneuronal knockout indicated a role in active zone organization and Ca^{2+} and BK channel regulation, but no requirement in synapse formation or maintenance (Luo et al., 2020). The multi-faceted roles for neurexins at synapses likely emerge as a consequence of the many neurexin isoforms that are generated by alternative splicing in vertebrates, the complexity of their cellular expression, and the potential for these isoforms to selectively interact with distinct postsynaptic partners.

The *C. elegans* genome encodes a single neurexin gene, *nrx-1* which generates two isoforms, a long α and a short γ isoform (Haklai-Topper et al., 2011). As is the case in other systems, *C. elegans* NRX-1 has roles in presynaptic organization, for example in calcium channel clustering at the active zone (Kurshan et al., 2018). Interestingly, the short γ-NRX-1 isoform that lacks identifiable ectodomains performs these roles (worms do not encode β-NRX-

1), but even in the absence of both isoforms, a level of presynaptic functionality is maintained, as indicated by the presence of evoked responses in muscle cells following presynaptic motor neuron stimulation (Philbrook et al., 2018). Our work here and our prior studies pointed toward the importance of NRX-1 at synapses between cholinergic and GABAergic motor neurons (Philbrook et al., 2018). Surprisingly, our time course studies here revealed that initial spine outgrowth and AChR clustering occurs normally in *nrx-1* mutants. However, spines and AChRs in *nrx-1* mutants become destabilized within hours and collapse, such that GABAergic dendrites are almost completely devoid of these postsynaptic components by L4 stage. Dendritic F-actin assemblies, that normally show discrete localization to spines, are also disorganized in *nrx-1* mutants, and are diffusely distributed throughout GABAergic dendritic processes in *nrx-1* mutants. We propose a model where presynaptic NRX-1 is required for the stabilization of dendritic spines and other postsynaptic structures in GABAergic dendrites. We suggest that presynaptic NRX-1 directs the organization of dendritic F-actin to stimulate maturation and stabilization of spines and postsynaptic receptor clusters. A similar form of F-actin based reorganization is a key step in presynaptic differentiation (Chia et al., 2012) (Chia et al., 2014).

Our work has parallels with recent studies in *Xenopus* (S. X. Chen et al., 2010) and *Drosophila* (Constance et al., 2018). In embryonic *Xenopus* brain, presynaptic β-neurexin stabilizes dendritic filopodia through an adhesive partnership with neuroligin to direct the development of dendritic arbors (S. X. Chen et al., 2010). Similarly, during fly metamorphosis, neurexin/neuroligin-based adhesion promotes the growth of neurite branches independently of synaptic activity (Constance et al., 2018). In our studies, NRX-1 stabilization of dendritic spines also occurs independently of synaptic activity and without involvement of neuroligin (Philbrook et al., 2018), suggesting interaction with an alternate binding partner that to date remains unidentified.

UNC-104/Kinesin 3-mediated synaptic delivery of NRX-1/Neurexin is an essential early step required for synapse maturation and stabilization

Efficient trafficking of synaptic and active zone cargoes to presynaptic terminals is essential for synapse formation and neuronal communication. Long-range axonal transport of synaptic vesicles and active zone proteins is carried out by anterograde kinesin and retrograde dynein motors. However, significant questions remain about which synaptic proteins are trafficked together and their dependence on specific motors. We found a specific requirement for the kinesin-3 UNC-104/KIF1A motor to deliver NRX-1 to presynaptic terminals. The anterograde transport of synaptic and dense core vesicles are also strongly dependent upon UNC-104 (Hall & Hedgecock, 1991) (Zahn et al., 2004). In contrast, several studies suggest that other AZ proteins exhibit weaker requirements for UNC-104 in their delivery. For example, ELKS-1 and other AZ proteins localize to *C. elegans* HSN synapses in an UNC-104-independent manner (Patel et al., 2006). Similarly, UNC-10/RIM properly concentrates within the nerve cord of *unc-104(e1265)* mutants, while synaptic vesicles accumulate in neuronal cell bodies (Koushika et al., 2001). Here, we found that NRX-1, UNC-10/RIM, ELKS-1/ELKS and CLA-1/piccolo all require UNC-104 for anterograde transport to cholinergic motor neuron synapses suggesting cell-specific functions for UNC-104 mediated delivery of active zone molecules.

Prior studies of rodent cultured neurons also suggested a requirement for the Kinesin-3 motor in neurexin transport, indicating that mechanisms for NRX-1 delivery to synapses are conserved (Neupert et al., 2015). The effects of a failure in delivery on synapse structure and function were not previously assessed. We found that dendritic spines collapse and AChR clusters disperse when UNC-104 delivery of synaptic cargoes is disrupted. Of the UNC-104 cargoes we analyzed, only mutation of *nrx-1* produced significant disruption of dendritic spines. We therefore propose that NRX-1 is a key UNC-104 cargo required for spine stabilization. Our

live imaging points to similar rates of transport for synaptic vesicles and NRX-1. However, it remains unclear whether NRX-1 is delivered as a component of synaptic vesicles or may be segregated into a distinct vesicular population, such as synaptic vesicle protein transport vesicles (STVs). Our findings that mutations which severely impair synaptic vesicle release do not affect spine density point toward the latter possibility.

The prior identification of a temperature-sensitive allele of *unc-104* allowed us to explore the temporal requirements for UNC-104 delivery of NRX-1. Importantly, we found that UNC-104 delivery was not solely required during early synaptogenesis but was critical throughout the developmental progression of the circuit toward maturity. These findings argue that ongoing delivery of presynaptic NRX-1 is required to maintain postsynaptic structure long after circuit assembly is complete. This raises interesting questions about the relationship between NRX-1 transport and synapse stability, perhaps suggesting that alterations in the rate of axonal NRX-1 transport may directly impact synaptic connectivity in mature animals. More broadly, our studies of the development and stabilization of synapses at dendritic spines in *C. elegans* provide a new view of the role of adhesive mechanisms in circuit connectivity and highlight a novel role for neurexin in the stabilization of mature synapses and dendritic spines.

Materials and Methods

Strains

All strains are N2 Bristol strain derivatives (wild type) and were maintained at room temperature (20-24°C) on nematode growth media plates (NGM) seeded with *E. coli* strain OP50. Transgenic strains were obtained by microinjection to achieve transformation (Mello et al., 1991) and identified using co-injection markers. Integrated lines were produced with X-ray irradiation and outcrossed to wild type/N2 Bristol eight times. Only hermaphrodites (L1-L4, young adults) were used in this study. A complete list of all strains used in this study is found in **Supplemental File 1**. Worms used for time course studies were staged by transferring embryos to seeded OP50 plates and transferring to 25°C (time point 0).

Molecular Biology

Plasmids were constructed using the two-slot Gateway Cloning system (Invitrogen) and confirmed by restriction digest and sequencing.

Utrophin/F-actin reporter. Sequence coding for mCherry was amplified from pDest-16 (and ligated into AgeI-HF/BspEI-digested PNYL183 (P*flp-13*::GFP::UtrCH, gift from Dong Yan's laboratory) to make pCL87 (P*flp-13*::mCherry::UtrCH).

TBA-1/tubulin reporter. Sequence coding for GFP::TBA-1 was amplified from plasmid pYJ128 (gift from Kang Shen laboratory) and ligated into a destination vector to create pDest-173. pDest-173 was recombined with pENTR-5'-*flp-13* to generate pDO61 (P*flp-13::*GFP::TBA-1).

Golgi apparatus reporter. pDest-161 (AMAN-2::GFP) was recombined with pENTR-3'-*flp-13* to generate pDO47 (P*flp-13*::AMAN-2::GFP). pDest-161 was created by HindIII-HF digestion of P*gpa-4*::AMAN::GFP (gift from Gert Jansen's laboratory) and ligated with a destination vector.

Mitochondria reporter. pre-su9::GFP was isolated from XmaI/SphI-HF digested pDM1389 (gift from A.V. Maricq lab) and ligated into a destination vector to create pDest-91 (pre-su9::GFP). pDest-91 was recombined with pENTR-5'-*flp-13* to generate pAP102 (P*flp-13*::pre-Su9::GFP).

ER reporter. Tag-RFP::TRAM-1 was amplified from plasmid pCT27 (gift from A.V. Maricq laboratory) and ligated to a KpnI-HF/NgoMIV digested destination vector to generate pDest-169. pDest-169 was recombined with pENTR-3'-*flp-13* to generate pDO59 (P*flp-13*::RFP::TRAM-1).

Presynaptic vesicle reporter. acr-5 promoter was amplified from plasmid pDM806 and ligated into pENTR-D-TOPO to generate pENTR-40. pENTR-40 was then recombined with pDest-5 to generate pAP264 (P*acr-5*::SNB-1::GFP).

B-type specific motor neuron reporter. pDO45 *(Pacr-5::GFP)* was created by recombining pDest-94 (GFP) and pENTR-40 (*acr-5* promoter).

myr-GCamp6f. myr-GCaMP6f was amplified from plasmid pDest-164 and ligated into a destination vector to create pDest-180. pDest-180 was recombined with pENTR3'-*flp-13* to create pDO69 (P*flp-13*::SL2::myr-GCamp6f).

CLA-1 reporter. 3XnovoGFP::CLA-1e was isolated from SphI-HF/AgeI-HF digested pKP85 (gift from Peri Kurshan) and ligated with a destination vector to create pDest-307 pDest-307 was recombined with pENTR-3'-*unc-17β* to generate pSR58 (P*unc-17β*::3XnovoGFP::CLA-1e cDNA).

unc-104 rescue constructs. UNC-104cDNA::mCherry was isolated from SphI-HF/NheI-HF digested pLL48 (gift from Kang Shen lab) and ligated with a destination vector to make pDest-308. pDest-308 was recombined with pENTR-*unc-17β* to generate pDO126 (P*unc-17β*::UNC-104cDNA::mCherry), pENTR-3'-*unc-47* to generate pDO134 (P*unc-47*::UNC-104cDNA::mCherry), and pENTR-3'-*myo-3* to generate pDO136 (P*myo-3*::UNC-104cDNA::mCherry).

Punc-17β::NRX-1::GFP. pAP112 was generated by recombining pENTR-3'-*unc-17β* with pDest-99 (NRX-1::GFP).

ELKS-1::mCherry. mCherry::ELKS-1cDNA was isolated from SphI-HF/AscI digested pK043 (gift from Peri Kurshan laboratory) and ligated with a destination vector to create pDest-325. pDest-

325 was recombined with pENTR-3'-*unc-129* to generate pDO156 (P*unc*-129::ELKS-1::mCherry).

UNC-10::GFP. UNC-10::GFP was isolated from BamHI-HF/SbfI digested pRIM3 (gift from Michael Nonet laboratory) and ligated with a destination vector to generate pDest-304. pDest-304 was recombined with pENTR-3'-*unc-129* to generate pDO128 (P*unc-129*::UNC-10::GFP) .

CRISPR/Cas-9 endogenously tagged NRX-1::GFP

Strain PHX3578 *nrx-1(syb3578)* was generated in N2 animals by SunyBiotech. Linker sequence and GFP were inserted after exon 27 at the 3' end of C29A12.4a1/*nrx-1.* Sequence flanking the GFP is below.

5': CGGGAAT<u>GG**A**GT**C**</u>GCAAAGAAAAAGGATTTTAAAGAGTGGTACGTAaaggtaccgcgggccc gggatccaccggtcgccaccatggtg

3': TCCATTTCTTCAATCAAAACTCAATACAATGATGATTAAAAAATTCACTTTTGTCTGCAAA TTGCCACAACTTCAAAACGGGTAC

PAM sequence is <u>underlined</u>

Synonymous mutations are **bolded**

Linker is in lower case letters

nrx-1 gene is CAPTIALIZED

Hydrogel solution

Hydrogel was prepared as an aqueous solution of polyethylene glycol (20 % w/v) and the photoinitiator Irgacure (0.5% stock, 0.1% working concentration), and stored at 4°C in the dark.

Confocal microscopy

All strains were immobilized with sodium azide (0.3 M) on a 2% or 5% agarose pad. Images were obtained using an Olympus BX51WI spinning disk confocal equipped with a 63x objective. For time course analyses, newly hatched larvae were transferred to a seeded OP50 plate and maintained at 25°C.

Long term live imaging of dendritic spines and F-actin. Nematodes were immobilized using 2 μL 50 mM muscimol in 10 μL 20% PEG hydrogel solution. Once paralyzed, hydrogel was curated using a handheld UV Transilluminator (312 nm, 3 minutes). Z-stacks were acquired every 5 minutes for at least one hour.

Live imaging of synaptic vesicle and NRX-1 trafficking. Nematodes were immobilized using 50 mM muscimol on a 5% agarose pad. Cholinergic commissures were imaged using Perkin Elmer spinning disk confocal equipped with a 63x objective using 100 ms exposure for 30 seconds.

Confocal microscopy analysis

All image analysis was conducted using ImageJ software (open source) within defined ROIs using intensity threshold values determined from control experiments. ROIs were located 15-30 μm anterior to DD1, DD2 or DD3 somata

Spine analysis. Mature spines were quantified as protrusions from the dendrite >0.3 μm in length (measuring from the base to the tip of the protrusion). Spine density was defined as the number of spines/unit length within a selected ROI.

3D rendering was conducted using Imaris/bitplane 3D image analysis software. Morphological categories were determined based on criteria used in (Harris et al., 1992). The ImageJ line tool was used to measure the length and width of the spine 3D rendering. Based on these calculations, spines were placed into one of four morphological categories based on the

following criteria: Stubby: middle width>total length; Thin: middle width=tip width; Mushroom: tip width>middle width; Branched: >1 spine head.

Fluorescence intensity analysis

F-actin/spine and F-actin/tubulin. The DD neurite was extracted using the line and line straighten functions. Total fluorescence intensity (utrophin/UtrCH, tubulin/TBA-1, ER/TRAM-1) on the dendritic shaft (line width 4) and below the dendritic shaft (line width 8) were measured. For soma fluorescence intensity, ROI was drawn around the perimeter of the soma.

Synaptic marker and receptor cluster. Background fluorescence was subtracted, and the number and size of synaptic/receptor puncta were measured using the 'analyze particles' function. Confocal montages were assembled using the 'straighten to line' function in a 50 μm region of the dorsal nerve cord.

Calcium Imaging

Animals were grown on plates seeded with OP50 containing 2.7 mM All-Trans Retinal (ATR). Plates were stored at 4°C under dark conditions and used within one week. Imaging was carried out using L4 animals immobilized in hydrogel (Burnett et al., 2018). Animals were transferred to 7.5 µL of the hydrogel mix placed on a silanized glass slide and covered with a glass slide. Hydrogel was cured using a handheld UV Transilluminator (312 nm, 3 minutes). Post-curing, the covering slide was removed and replaced with a coverslip. Imaging was carried out using a Yokogawa CSU-X1-A1N spinning disk confocal system (Perkin Elmer) equipped with EM-CCD camera (Hamamatsu, C9100-50) and 63X oil immersion objective. Chrimson photoactivation (~30 mW/cm^2) was achieved using a TTL-controlled 625 nm light guide coupled LED (Mightex Systems). A 556 nm BrightLine single-edge short-pass dichroic beam splitter was positioned in the light path (Semrock) (**Figure S1.3**). Data were acquired at 10 Hz for 15 s using Volocity software and binned at 1×1 during acquisition. Analysis was performed using ImageJ. The DD

neurite process in each time series was extracted using the straighten function, background subtracted and photobleaching correction was carried out by fitting an exponential function to the data (CorrectBleach plugin). A smoothing function was applied to the data to enhance signal-to-noise. Individual spine ROIs were identified using the mCherry fluorescence. Post imaging processing, pre-stimulus baseline fluorescence (F_0) was calculated as the average of the data points in the first 4 s of the recording. Data was normalized to prestimulus baseline as $\Delta F/F_0$, where $\Delta F = F - F_0$. Peak $\Delta F/F_0$ was determined by fitting a Gaussian function to the $\Delta F/F_0$ time sequence using Multi peak 2.0 (Igor Pro, WaveMetrics). All collected data were analyzed, including failures (no response to stimulation). Control recordings were carried out in the absence of Retinal.

Electron microscopy

Staged L4 wild type (P*flp-13*::ACR-12::GFP) hermaphrodites were subjected to high-pressure freeze fixation using a HPM10 high-pressure freezer. Hermaphrodites were slowly freeze substituted in 2% osmium tetroxide (OSO4), 0.1% uranyl acetate, 2% H2O in acetone fixative (Weimer, 2006) (Muller-Reichert et al., 2010). Samples were embedded in blocks, sectioned at 70 nm-thick serial sections and subsequently collected on copper slot grids. Samples were post-stained in 2.5% uranyl acetate in 70% methanol for 20 minutes and 1:6 lead citrate in H_2O for 2 minutes at room temperature. Images were acquired using the Philips CM10 TEM.

We identified and characterized 6 DD1 spines from a 35 µm region of the ventral head region where the neurite emerges from the cell body of DD1 GABAergic motor neuron.

Temperature shift experiments

Wild type and *unc-104(ce782)ts* animals eggs were raised at 13.5°C until L3 or L4 stage as indicated, then shifted to 25°C for 16-20 hrs prior to imaging. Animals were age-matched at the time of the temperature shift to account for developmental delays of *unc-104(ce782)ts* mutants.

References

Aakalu, G., Smith, W. B., Nguyen, N., Jiang, C., & Schuman, E. M. (2001). Dynamic visualization of local protein synthesis in hippocampal neurons. *Neuron, 30*(2), 489-502. doi:10.1016/s0896-6273(01)00295-1

Bohme, M. A., Beis, C., Reddy-Alla, S., Reynolds, E., Mampell, M. M., Grasskamp, A. T., . . . Sigrist, S. J. (2016). Active zone scaffolds differentially accumulate Unc13 isoforms to tune Ca(2+) channel-vesicle coupling. *Nat Neurosci, 19*(10), 1311-1320. doi:10.1038/nn.4364

Borovac, J., Bosch, M., & Okamoto, K. (2018). Regulation of actin dynamics during structural plasticity of dendritic spines: Signaling messengers and actin-binding proteins. *Mol Cell Neurosci, 91*, 122-130. doi:10.1016/j.mcn.2018.07.001

Burnett, K., Edsinger, E., & Albrecht, D. R. (2018). Rapid and gentle hydrogel encapsulation of living organisms enables long-term microscopy over multiple hours. *Commun Biol, 1*, 73. doi:10.1038/s42003-018-0079-6

Chen, L. Y., Jiang, M., Zhang, B., Gokce, O., & Sudhof, T. C. (2017). Conditional Deletion of All Neurexins Defines Diversity of Essential Synaptic Organizer Functions for Neurexins. *Neuron, 94*(3), 611-625 e614. doi:10.1016/j.neuron.2017.04.011

Chen, S. X., Tari, P. K., She, K., & Haas, K. (2010). Neurexin-neuroligin cell adhesion complexes contribute to synaptotropic dendritogenesis via growth stabilization mechanisms in vivo. *Neuron, 67*(6), 967-983. doi:10.1016/j.neuron.2010.08.016

Chia, P. H., Chen, B., Li, P., Rosen, M. K., & Shen, K. (2014). Local F-actin network links synapse formation and axon branching. *Cell, 156*(1-2), 208-220. doi:10.1016/j.cell.2013.12.009

Chia, P. H., Patel, M. R., & Shen, K. (2012). NAB-1 instructs synapse assembly by linking adhesion molecules and F-actin to active zone proteins. *Nat Neurosci, 15*(2), 234-242. doi:10.1038/nn.2991

Chirillo, M. A., Waters, M. S., Lindsey, L. F., Bourne, J. N., & Harris, K. M. (2019). Local resources of polyribosomes and SER promote synapse enlargement and spine clustering after long-term potentiation in adult rat hippocampus. *Sci Rep, 9*(1), 3861. doi:10.1038/s41598-019-40520-x

Constance, W. D., Mukherjee, A., Fisher, Y. E., Pop, S., Blanc, E., Toyama, Y., & Williams, D. W. (2018). Neurexin and Neuroligin-based adhesion complexes drive axonal arborisation growth independent of synaptic activity. *eLife, 7*. doi:10.7554/eLife.31659

Cuentas-Condori, A., Mulcahy, B., He, S., Palumbos, S., Zhen, M., & Miller, D. M., III. (2019). C. elegans neurons have functional dendritic spines. *eLife, 8*, e47918. doi:10.7554/eLife.47918

Dai, Y., Taru, H., Deken, S. L., Grill, B., Ackley, B., Nonet, M. L., & Jin, Y. (2006). SYD-2 Liprin-alpha organizes presynaptic active zone formation through ELKS. *Nat Neurosci, 9*(12), 1479-1487. doi:10.1038/nn1808

De Roo, M., Klauser, P., Garcia, P. M., Poglia, L., & Muller, D. (2008). Spine dynamics and synapse remodeling during LTP and memory processes. *Prog Brain Res, 169*, 199-207. doi:10.1016/S0079-6123(07)00011-8

Edwards, S. L., Yorks, R. M., Morrison, L. M., Hoover, C. M., & Miller, K. G. (2015). Synapse-Assembly Proteins Maintain Synaptic Vesicle Cluster Stability and

Regulate Synaptic Vesicle Transport in Caenorhabditis elegans. *Genetics, 201*(1), 91-116. doi:10.1534/genetics.115.177337

Edwards, S. L., Yu, S. C., Hoover, C. M., Phillips, B. C., Richmond, J. E., & Miller, K. G. (2013). An organelle gatekeeper function for Caenorhabditis elegans UNC-16 (JIP3) at the axon initial segment. *Genetics, 194*(1), 143-161. doi:10.1534/genetics.112.147348

Engert, F., & Bonhoeffer, T. (1999). Dendritic spine changes associated with hippocampal long-term synaptic plasticity. *Nature, 399*(6731), 66-70. doi:10.1038/19978

Fouquet, W., Owald, D., Wichmann, C., Mertel, S., Depner, H., Dyba, M., . . . Sigrist, S. J. (2009). Maturation of active zone assembly by Drosophila Bruchpilot. *J Cell Biol, 186*(1), 129-145. doi:10.1083/jcb.200812150

Friedman, H. V., Bresler, T., Garner, C. C., & Ziv, N. E. (2000). Assembly of new individual excitatory synapses: time course and temporal order of synaptic molecule recruitment. *Neuron, 27*(1), 57-69. doi:10.1016/s0896-6273(00)00009-x

Gally, C., Eimer, S., Richmond, J. E., & Bessereau, J. L. (2004). A transmembrane protein required for acetylcholine receptor clustering in Caenorhabditis elegans. *Nature, 431*(7008), 578-582. doi:10.1038/nature02893

Gu, J., Firestein, B. L., & Zheng, J. Q. (2008). Microtubules in dendritic spine development. *J Neurosci, 28*(46), 12120-12124. doi:10.1523/JNEUROSCI.2509-08.2008

Haklai-Topper, L., Soutschek, J., Sabanay, H., Scheel, J., Hobert, O., & Peles, E. (2011). The neurexin superfamily of Caenorhabditis elegans. *Gene Expr Patterns, 11*(1-2), 144-150. doi:10.1016/j.gep.2010.10.008

Hall, D. H., & Hedgecock, E. M. (1991). Kinesin-related gene unc-104 is required for axonal transport of synaptic vesicles in C. elegans. *Cell, 65*(5), 837-847. doi:10.1016/0092-8674(91)90391-b

Hall, D. H., & Russell, R. L. (1991). The posterior nervous system of the nematode Caenorhabditis elegans: serial reconstruction of identified neurons and complete pattern of synaptic interactions. *J Neurosci, 11*(1), 1-22. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/1986064

Harris, K. M., Jensen, F. E., & Tsao, B. (1992). Three-dimensional structure of dendritic spines and synapses in rat hippocampus (CA1) at postnatal day 15 and adult ages: implications for the maturation of synaptic physiology and long-term potentiation. *J Neurosci, 12*(7), 2685-2705. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/1613552

He, S., Cuentas-Condori, A., & Miller, D. M., 3rd. (2019). NATF (Native and Tissue-Specific Fluorescence): A Strategy for Bright, Tissue-Specific GFP Labeling of Native Proteins in Caenorhabditis elegans. *Genetics, 212*(2), 387-395. doi:10.1534/genetics.119.302063

Hendi, A., Kurashina, M., & Mizumoto, K. (2019). Intrinsic and extrinsic mechanisms of synapse formation and specificity in C. elegans. *Cell Mol Life Sci, 76*(14), 2719-2738. doi:10.1007/s00018-019-03109-1

Hendricks, M., Ha, H., Maffey, N., & Zhang, Y. (2012). Compartmentalized calcium dynamics in a C. elegans interneuron encode head movement. *Nature, 487*(7405), 99-103. doi:10.1038/nature11081

Hu, X., Viesselmann, C., Nam, S., Merriam, E., & Dent, E. W. (2008). Activity-dependent dynamic microtubule invasion of dendritic spines. *J Neurosci, 28*(49), 13094-13105. doi:10.1523/JNEUROSCI.3074-08.2008

Jaworski, J., Kapitein, L. C., Gouveia, S. M., Dortland, B. R., Wulf, P. S., Grigoriev, I., . . . Hoogenraad, C. C. (2009). Dynamic microtubules regulate dendritic spine morphology and synaptic plasticity. *Neuron, 61*(1), 85-100. doi:10.1016/j.neuron.2008.11.013

Karagas, N. E., & Venkatachalam, K. (2019). Roles for the Endoplasmic Reticulum in Regulation of Neuronal Calcium Homeostasis. *Cells, 8*(10). doi:10.3390/cells8101232

Kasem, E., Kurihara, T., & Tabuchi, K. (2018). Neurexins and neuropsychiatric disorders. *Neurosci Res, 127*, 53-60. doi:10.1016/j.neures.2017.10.012

Koushika, S. P., Richmond, J. E., Hadwiger, G., Weimer, R. M., Jorgensen, E. M., & Nonet, M. L. (2001). A post-docking role for active zone protein Rim. *Nat Neurosci, 4*(10), 997-1005. doi:10.1038/nn732

Kumar, J., Choudhary, B. C., Metpally, R., Zheng, Q., Nonet, M. L., Ramanathan, S., . . . Koushika, S. P. (2010). The Caenorhabditis elegans Kinesin-3 motor UNC-104/KIF1A is degraded upon loss of specific binding to cargo. *PLoS Genet, 6*(11), e1001200. doi:10.1371/journal.pgen.1001200

Kurshan, P. T., Merrill, S. A., Dong, Y., Ding, C., Hammarlund, M., Bai, J., . . . Shen, K. (2018). gamma-Neurexin and Frizzled Mediate Parallel Synapse Assembly Pathways Antagonized by Receptor Endocytosis. *Neuron, 100*(1), 150-166 e154. doi:10.1016/j.neuron.2018.09.007

Kurup, N., & Jin, Y. (2016). Neural circuit rewiring: insights from DD synapse remodeling. *Worm, 5*(1), e1129486. doi:10.1080/21624054.2015.1129486

Li, J., Ashley, J., Budnik, V., & Bhat, M. A. (2007). Crucial role of Drosophila neurexin in proper active zone apposition to postsynaptic densities, synaptic growth, and synaptic transmission. *Neuron, 55*(5), 741-755. doi:10.1016/j.neuron.2007.08.002

Li, Z., Okamoto, K., Hayashi, Y., & Sheng, M. (2004). The importance of dendritic mitochondria in the morphogenesis and plasticity of spines and synapses. *Cell, 119*(6), 873-887. doi:10.1016/j.cell.2004.11.003

Lipton, D. M., Maeder, C. I., & Shen, K. (2018). Rapid Assembly of Presynaptic Materials behind the Growth Cone in Dopaminergic Neurons Is Mediated by Precise Regulation of Axonal Transport. *Cell Rep, 24*(10), 2709-2722. doi:10.1016/j.celrep.2018.07.096

Liu, Q. A., & Shio, H. (2008). Mitochondrial morphogenesis, dendrite development, and synapse formation in cerebellum require both Bcl-w and the glutamate receptor delta2. *PLoS Genet, 4*(6), e1000097. doi:10.1371/journal.pgen.1000097

Lu, W., Bushong, E. A., Shih, T. P., Ellisman, M. H., & Nicoll, R. A. (2013). The cell-autonomous role of excitatory synaptic transmission in the regulation of neuronal structure and function. *Neuron, 78*(3), 433-439. doi:10.1016/j.neuron.2013.02.030

Luo, F., Sclip, A., Jiang, M., & Sudhof, T. C. (2020). Neurexins cluster Ca(2+) channels within the presynaptic active zone. *EMBO J, 39*(7), e103208. doi:10.15252/embj.2019103208

McDonald, N. A., Fetter, R. D., & Shen, K. (2020). Assembly of synaptic active zones requires phase separation of scaffold molecules. *Nature, 588*(7838), 454-458. doi:10.1038/s41586-020-2942-0

Mello, C. C., Kramer, J. M., Stinchcomb, D., & Ambros, V. (1991). Efficient gene transfer in C.elegans: extrachromosomal maintenance and integration of transforming sequences. *EMBO J, 10*(12), 3959-3970. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/1935914

Meng, L., Mulcahy, B., Cook, S. J., Neubauer, M., Wan, A., Jin, Y., & Yan, D. (2015). The Cell Death Pathway Regulates Synapse Elimination through Cleavage of Gelsolin in Caenorhabditis elegans Neurons. *Cell Rep, 11*(11), 1737-1748. doi:10.1016/j.celrep.2015.05.031

Missler, M., & Sudhof, T. C. (1998). Neurexins: three genes and 1001 products. *Trends Genet, 14*(1), 20-26. doi:10.1016/S0168-9525(97)01324-3

Muller-Reichert, T., Mancuso, J., Lich, B., & McDonald, K. (2010). Three-dimensional reconstruction methods for Caenorhabditis elegans ultrastructure. *Methods Cell Biol, 96*, 331-361. doi:10.1016/S0091-679X(10)96015-9

Nagerl, U. V., Kostinger, G., Anderson, J. C., Martin, K. A., & Bonhoeffer, T. (2007). Protracted synaptogenesis after activity-dependent spinogenesis in hippocampal neurons. *J Neurosci, 27*(30), 8149-8156. doi:10.1523/JNEUROSCI.0511-07.2007

Neupert, C., Schneider, R., Klatt, O., Reissner, C., Repetto, D., Biermann, B., . . . Heine, M. (2015). Regulated Dynamic Trafficking of Neurexins Inside and Outside of Synaptic Terminals. *J Neurosci, 35*(40), 13629-13647. doi:10.1523/JNEUROSCI.4041-14.2015

Owald, D., Khorramshahi, O., Gupta, V. K., Banovic, D., Depner, H., Fouquet, W., . . . Sigrist, S. J. (2012). Cooperation of Syd-1 with Neurexin synchronizes pre- with postsynaptic assembly. *Nat Neurosci, 15*(9), 1219-1226. doi:10.1038/nn.3183

Patel, M. R., Lehrman, E. K., Poon, V. Y., Crump, J. G., Zhen, M., Bargmann, C. I., & Shen, K. (2006). Hierarchical assembly of presynaptic components in defined C. elegans synapses. *Nat Neurosci, 9*(12), 1488-1498. doi:10.1038/nn1806

Perez-Alvarez, A., Yin, S., Schulze, C., Hammer, J. A., Wagner, W., & Oertner, T. G. (2020). Endoplasmic reticulum visits highly active spines and prevents runaway potentiation of synapses. *Nat Commun, 11*(1), 5083. doi:10.1038/s41467-020-18889-5

Philbrook, A., Ramachandran, S., Lambert, C. M., Oliver, D., Florman, J., Alkema, M. J., . . . Francis, M. M. (2018). Neurexin directs partner-specific synaptic connectivity in C. elegans. *eLife, 7*. doi:10.7554/eLife.35692

Richmond, J. E., Davis, W. S., & Jorgensen, E. M. (1999). UNC-13 is required for synaptic vesicle fusion in C. elegans. *Nat Neurosci, 2*(11), 959-964. doi:10.1038/14755

Rolls, M. M., Hall, D. H., Victor, M., Stelzer, E. H., & Rapoport, T. A. (2002). Targeting of rough endoplasmic reticulum membrane proteins and ribosomes in invertebrate neurons. *Mol Biol Cell, 13*(5), 1778-1791. doi:10.1091/mbc.01-10-0514

Sala, C., & Segal, M. (2014). Dendritic spines: the locus of structural and functional plasticity. *Physiol Rev, 94*(1), 141-188. doi:10.1152/physrev.00012.2013

Sando, R., Bushong, E., Zhu, Y., Huang, M., Considine, C., Phan, S., . . . Maximov, A. (2017). Assembly of Excitatory Synapses in the Absence of Glutamatergic

Neurotransmission. *Neuron, 94*(2), 312-321 e313. doi:10.1016/j.neuron.2017.03.047

Saneyoshi, T., Fortin, D. A., & Soderling, T. R. (2010). Regulation of spine and synapse formation by activity-dependent intracellular signaling pathways. *Curr Opin Neurobiol, 20*(1), 108-115. doi:10.1016/j.conb.2009.09.013

Sarasija, S., & Norman, K. R. (2015). A gamma-Secretase Independent Role for Presenilin in Calcium Homeostasis Impacts Mitochondrial Function and Morphology in Caenorhabditis elegans. *Genetics, 201*(4), 1453-1466. doi:10.1534/genetics.115.182808

Sigler, A., Oh, W. C., Imig, C., Altas, B., Kawabe, H., Cooper, B. H., . . . Brose, N. (2017). Formation and Maintenance of Functional Spines in the Absence of Presynaptic Glutamate Release. *Neuron, 94*(2), 304-311 e304. doi:10.1016/j.neuron.2017.03.029

Steward, O., & Levy, W. B. (1982). Preferential localization of polyribosomes under the base of dendritic spines in granule cells of the dentate gyrus. *J Neurosci, 2*(3), 284-291. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/7062109

Sudhof, T. C. (2008). Neuroligins and neurexins link synaptic function to cognitive disease. *Nature, 455*(7215), 903-911. doi:10.1038/nature07456

Sudhof, T. C. (2017). Synaptic Neurexin Complexes: A Molecular Code for the Logic of Neural Circuits. *Cell, 171*(4), 745-769. doi:10.1016/j.cell.2017.10.024

Tonnesen, J., & Nagerl, U. V. (2016). Dendritic Spines as Tunable Regulators of Synaptic Signals. *Front Psychiatry, 7*, 101. doi:10.3389/fpsyt.2016.00101

Vos, M., Lauwers, E., & Verstreken, P. (2010). Synaptic mitochondria in synaptic transmission and organization of vesicle pools in health and disease. *Front Synaptic Neurosci, 2*, 139. doi:10.3389/fnsyn.2010.00139

Weimer, R. M. (2006). Preservation of C. elegans tissue via high-pressure freezing and freeze-substitution for ultrastructural analysis and immunocytochemistry. *Methods Mol Biol, 351*, 203-221. doi:10.1385/1-59745-151-7:203

White, J. (2018). Clues to basis of exploratory behaviour of the C. elegans snout from head somatotropy. *Philos Trans R Soc Lond B Biol Sci, 373*(1758). doi:10.1098/rstb.2017.0367

White, J. G., Albertson, D. G., & Anness, M. A. (1978). Connectivity changes in a class of motoneurone during the development of a nematode. *Nature, 271*(5647), 764-766. doi:10.1038/271764a0

White, J. G., Southgate, E., Thomson, J. N., & Brenner, S. (1976). The structure of the ventral nerve cord of Caenorhabditis elegans. *Philos Trans R Soc Lond B Biol Sci, 275*(938), 327-348. doi:10.1098/rstb.1976.0086

White, J. G., Southgate, E., Thomson, J. N., & Brenner, S. (1986). The structure of the nervous system of the nematode Caenorhabditis elegans. *Philos Trans R Soc Lond B Biol Sci, 314*(1165), 1-340. doi:10.1098/rstb.1986.0056

Xuan, Z., Manning, L., Nelson, J., Richmond, J. E., Colon-Ramos, D. A., Shen, K., & Kurshan, P. T. (2017). Clarinet (CLA-1), a novel active zone protein required for synaptic vesicle clustering and release. *eLife, 6*. doi:10.7554/eLife.29276

Yan, J., Chao, D. L., Toba, S., Koyasako, K., Yasunaga, T., Hirotsune, S., & Shen, K. (2013). Kinesin-1 regulates dendrite microtubule polarity in Caenorhabditis elegans. *eLife, 2*, e00133. doi:10.7554/eLife.00133

Zahn, T. R., Angleson, J. K., MacMorris, M. A., Domke, E., Hutton, J. F., Schwartz, C., & Hutton, J. C. (2004). Dense core vesicle dynamics in Caenorhabditis elegans neurons and the role of kinesin UNC-104. *Traffic, 5*(7), 544-559. doi:10.1111/j.1600-0854.2004.00195.x

Zhen, M., & Jin, Y. (1999). The liprin protein SYD-2 regulates the differentiation of presynaptic termini in C. elegans. *Nature, 401*(6751), 371-375. doi:10.1038/43886

Zito, K., Scheuss, V., Knott, G., Hill, T., & Svoboda, K. (2009). Rapid functional maturation of nascent dendritic spines. *Neuron, 61*(2), 247-258. doi:10.1016/j.neuron.2008.10.054

Figures

Mushroom 5.49 ± 0.66% Branched 5.49 ± 0.66% Stubby 73.63 ± 3.09% Thin 15.38 ± 1.21%

Figure 1. *C. elegans* GABAergic DD neuron dendritic spines share characteristics with dendritic spines in vertebrates.

(A)   Representative fluorescent images and cartoon representation of the ventral nerve cord/dendrite of a single L4 stage GABAergic DD (DD2) motor neuron with ventrally projecting dendritic spines. CB, cell body.

(B)   Fluorescent image (top, inverted LUT) and 3D rendering (middle and insets) of DD dendritic spines from an animal expressing P*flp-13*::mCherry, shows diverse spine morphologies. DD dendritic spines share similar morphologies with mammalian dendritic spines: mushroom (6.02 ± 0.66%), branched (6.02 ± 0.66%), stubby (80.72 ± 3.09%), and thin (16.87 ± 1.21%). Arrowheads, spine insets. n=83 dendritic spines from 11 animals, measurements, percentage ± SD.

(C)   Representative images of DD dendritic spines from animal co-expressing P*flp-13*::mCherry (labeling dendrite, red) and P*flp-13*::GFP::UtrCH (labeling F-actin, green). F-actin is highly enriched in DD dendritic spines.

(D)   Scatterplot of F-actin/GFP::UtrCH fluorescence in dendritic shaft compared to dendritic spines, normalized to total fluorescence within the ROI. Bars indicate mean ± SEM. Student's t-test, ***$p<0.001$, n=21 animals.

(E)   Scatterplot of the percentage of spines with or without F-actin. Bars, mean ± SEM, n=21 animals.

(F)   Fluorescent images of tubulin (P*flp-13*::TBA-1::GFP) (green) and actin (P*flp-13*::mCherry::UtrCH) (red) in DD neurons. Dendritic spines are highly enriched with F-actin while tubulin occupies the main dendritic process.

(G)   Quantification of fluorescence ratio of actin/tubulin in the dendritic shaft versus dendritic spines. Student's t-test, ****p<0.0001, n=31 spines from 6 animals.

(H)   Calcium transients recorded from wild type GABAergic DD motor neuron dendrites expressing myrGCaMP6f (P*flp-13*::myrGCaMP6f::SL2::mCherry) during stimulation of presynaptic cholinergic motor neurons (P*acr-2::*Chrimson). Top, representative image showing dendritic spines identified by mCherry fluorescence (inverted LUT). Representative maximum intensity projection heat maps showing calcium fluxes immediately prior to stimulation (prestimulus, middle panel) and during stimulus (ON, bottom panel). Heat maps were generated by maximum intensity projection of myrGCaMP6f fluorescence from the first 4 seconds of recording (prestimulus) and the 5 s period of stimulation (red sharded bar in I). Numbering corresponds to representative traces in Figure 1I.

(I)   Representative evoked responses for indicated dendritic spines in wild type animals (wild type 1, 2 and 3, Figure 1H), animals grown in the absence of retinal (-Retinal), and *unc-17(e113)* mutants. Calcium responses were continuously recorded at 10 Hz for 15 s. Prestimulus calcium fluxes were recorded for 5 s, followed by stimulation (625 nm, ~30 mW/cm^2) of cholinergic motor neurons for 5 s. All data was normalized to prestimulus (ΔF/F$_0$). Red shaded bar indicates duration of stimulation. Pink traces indicate Gaussian fits to responses (ΔF/F$_0$) during stimulation.

(J)   Scatter plot showing peak ΔF/F$_0$ of responses during stimulation in wild type or *unc-17(e113)* mutants. Bars, mean ± SEM. Statistical analysis, Student's t-test, ****p<0.0001. n ≥ 10 animals.

A hours post-hatch | 12 hrs | 16 hrs | 20 hrs | 24 hrs | 32 hrs | L4

Presynapse

CLA-1
GABA::mCherry
ACh::
novoGFP::CLA-1e

ACh synaptic vesicles
GABA::mCherry
ACh::SNB-1::GFP

B hours post-hatch | 12 hrs | 16 hrs | 20 hrs | 24 hrs | 32 hrs | L4

Postsynapse

AChRs
GABA::mCherry
GABA::ACR-12::GFP

actin
GABA::mCherry
GABA::GFP::UtrCH

2 μm

C Spines

20 hrs

L4

5 μm
GABA::mCherry (inverted)

D CLA-1/Clarinet
CLA-1 clusters
spines
Average Number
12 17 13 13 10 8 24
Hours/Age post-hatch

E SNB-1/SVs
presynaptic vesicles
spines
Average Number
12 15 12 13 10 15
Hours/Age post-hatch

F AChRs
ACR-12 clusters
spines
Average Number
17 13 29 28 29 16 12
Hours/Age post-hatch

G actin
F-actin clusters
spines
Average Number
22 22 19 19 20 13 15
Hours/Age post-hatch

H 24 hours post-hatch
GABA::ACR-12::GFP
GABA::mCherry::UtrCH
AChR
F-actin
merge
1 μm

I
1 μm
AChR F-actin

J L4 animals
GABA::ACR-12::GFP
GABA::mCherry::UtrCH
AChR
F-actin
merge
1 μm

K
1 μm
AChR F-actin

L
F-actin localizes along DD neuron

AChRs traffic along the dendritic shaft. Presynaptic vesicles/proteins form along axon of newly born ACh neurons

AChRs and F-actin migrate into immature spiny protrusions and presynaptic proteins begin to cluster

F-actin enriched dendritic spines elongate near sites of mature active zones

development

ACh axon | GABA dendritic shaft | F-actin | CLA-1 | AChR | SNB-1 | GABA dendrite with spines

Figure 2. Temporal order of molecular events during synapse and spine development.

(A)   Fluorescent images of GABAergic dendrites at 12, 16, 20, 24, and 32 hours after hatch and at L4 stage (~42-50 hours post-hatch). Animals express a postsynaptic dendritic spine marker (P*flp-13*::mCherry) with either presynaptic active zone (P*unc-17β*::GFPnovo2::CLA-1e) (top) or synaptic vesicle marker (P*acr-5*::SNB-1::GFP) (bottom). Images are pseudo colored to indicate spines (magenta) and presynaptic CLA-1 or vesicle clusters (cyan).

(B)   Fluorescent images of GABAergic dendrites at 12, 16, 20, 24, and 32 hours after hatch and L4 animals (~42-50 hours post-hatch). Animals express a postsynaptic dendritic spine marker (P*flp-13*::mCherry) with either postsynaptic AChR (P*unc-47*::ACR-12::GFP, top) or postsynaptic actin (P*flp-13*::GFP::UtrCH, bottom) markers. Images are pseudo colored to indicate spines (magenta) and postsynaptic AChR or F-actin (cyan). White asterisks (*) indicate images used in Figure 2C.

(C)   Fluorescent images (inverted LUT) of GABAergic dendrites at 20 hours after hatch and L4 stage (~42-50 hours post-hatch) as indicated. Animals express a DD dendritic spine marker (P*flp-13*::mCherry). Dashed red boxes indicate regions used in Figure 2B (asterisks).

(D-G) Quantification of spine number (purple) relative to clusters of active zone protein/CLA-1 (D), synaptic vesicle/SNB-1 (E), AChR/ACR-12 (F), and F-actin/UtrCH (G) (black) through development. Clusters of each of these markers are visible well prior to spine outgrowth. Blue shading indicates the approximate

period of DD remodeling. For this and subsequent figures, numbering indicates animals quantified at each timepoint for each genotype. Data points and bars indicate mean ± SEM.

(H)   Fluorescent confocal images of DD1 AChR/ACR-12 (P*flp-13*::ACR-12::GFP, magenta) and F-actin clusters (P*flp-13*::mCherry::UtrCH, cyan) 24 hours after hatch. Dotted rectangle indicates high magnification image shown in 2I. Expanded view of DD1 dendrites 24 hours after hatch from Figure 2H.

(J)   Fluorescent confocal images of DD1 AChR/ACR-12 (P*flp-13*::ACR-12::GFP, magenta) and F-actin clusters (P*flp-13*::mCherry::UtrCH, cyan) at L4 stage. Dotted rectangle indicates high magnification image shown in 2K.

(K)   Expanded view of DD1 dendrites 24 hours after hatch from Figure 2J.

(L)   Cartoon summary of the order of events during development of synapses onto DD neurons. In early development, presynaptic vesicles (SNB-1) and active zone protein (CLA-1) localize along presynaptic terminals while postsynaptic F-actin and postsynaptic receptors cluster along the postsynaptic process. Following their formation, dendritic spines begin to extend. Spines continue to elongate throughout development and into the mature circuit (L4).

Table 1. Analysis of dendritic spines and cholinergic receptors in mutant strains with altered synaptic activity.

Dendritic spine number, spine length, ACR-12/AChR cluster number, and AChR cluster size at L4 stage normalized to the same measurements from wild type are shown (mean ± SEM) for each genotype (ACR-2/nAChR UNC-2/CaV2α, UNC-13/MUNC-13, UNC-17/VAChT, UNC-18/MUNC-18, UNC-31/CAPS). Mutations that affected synaptic activity had little effect on the density of spines or AChR clusters, but had variable effects on spine length and AChR cluster size. One-way AVOVA, Dunnett's multiple comparisons test, *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

Genotype	Spines						AChR/receptors					
	Normalized spine number ± SEM (%)	N	P value	Normalized spine length ± SEM (%)	N	P value	Normalized ACR-12 cluster number ± SEM (%)	N	P value	Normalized ACR-12 cluster size ± SEM (%)	N	P value
wild type	100.0 ± 4.1	58	ns	100.0 ± 2.1	404	ns	100.0 ± 6.3	17	ns	100.0 ± 5.1	220	ns
acr-2(ok1887)	86.2 ± 9.1	24	ns	77.1 ± 2.4	141	****	94.6 ± 10.4	20	ns	109.8 ± 5.9	245	ns
acr-2(n2420)gf	52.9 ± 5.9	26	****	83.8 ± 2.6	135	**	75.6 ± 6.3	18	ns	81.3 ± 5.4	178	ns
unc-2(e55)	81.8 ± 7.6	29	ns	90.9 ± 3.7	122	ns	77.2 ± 9.5	18	ns	83.9 ± 5.7	172	ns
unc-2(zf35)gf	81.7 ± 7.2	22	ns	114.7 ± 4.8	117	*	120.0 ± 7.2	20	ns	130.1 ± 7.2	312	**
unc-13(e51)	62.1 ± 8.1	18	**	121.5 ± 7.4	78	***	85.0 ± 10.6	19	ns	74.8 ± 4.4	210	ns
unc-17(e113)	104.5 ± 6.8	12	ns	94.8 ± 4.4	83	ns	101.6 ± 7.1	24	ns	119.8 ± 5.7	317	ns
unc-18(e234)	76.9 ± 10.0	11	ns	98.5 ± 3.9	59	ns	96.4 ± 9.9	19	ns	113.3 ± 6.5	238	ns
unc-31(e169)	97.6 ± 5.7	22	ns	106.3 ± 3.6	166	ns	107.0 ± 8.8	21	ns	123.9 ± 5.9	291	*

A

ACh::NRX-1

GABA:: mCherry
ACh:: NRX-1::GFP

16 hrs 20 hrs 24 hrs L4

2 µm

B

NRX-1 cluster
Spine

14
16
18
15

Average Number

16 18 20 24 L4

Hours/Age post-hatch

C

wild type   *nrx-1(wy778)*

16 hrs
18 hrs
20 hrs
24 hrs
L4

5 µm

GABA::myrGFP(inverted)

D

spines

wild type
nrx-1(wy778)

% animals with 2 or more spines

100
75
50
25

16 18 20 24 L4

Hours/Age post-hatch

E

wild type   *nrx-1(wy778)*

16 hrs
18 hrs
20 hrs
24 hrs
L4

5 µm

GABA::ACR-12::GFP

F

AChR

WT
nrx-1(wy778)

*
***
***
**

Number of ACR-12::GFP clusters

16 18 20 24 L4

Hours/Age post-hatch

G

F-actin (*GABA*::UtrCH::GFP)
spines (*GABA*::mCherry)

16 hours after hatch   L4 stage

wild type   wild type

5 µm   5 µm   *nrx-1(wy778)*
nrx-1(wy778)

Figure 3. Dendritic spines and postsynaptic components form initially but subsequently collapse without the synaptic adhesion protein, NRX-1.

(A)   Fluorescent confocal images of dendritic spines (P*flp-13*::mCherry, magenta)and cholinergic expression of NRX-1::GFP (P*unc-17β*::NRX-1::GFP, cyan) at 16, 20, and 24 hours after hatch and L4 stage (~42-50 hours post-hatch). Presynaptic clusters of NRX-1::GFP are visible by 16 hours after hatch and increase in size until roughly 24 hours after hatch.

(B)   Quantification of the average number of spines (purple) and NRX-1 clusters (black) at the time points indicated. Blue shading indicates the approximate period of DD remodeling. Numbers indicate the number of animals quantified at each timepoint. Data points indicate mean ± SEM.

(A)   Fluorescent images (inverted LUT) of DD1 dendritic spines (P*flp-13*::myrGFP) in wild type (left) and *nrx-1(wy778)* (right) animals at 16, 18, 20, and 24 hours after hatch and at L4 stage. Immature dendritic spines clusters form initially but then collapse in *nrx-1* mutants.

(B)   Quantification of the percentage of wild type (black) and *nrx-1(wy778)* (purple) animals with 2 or more DD dendritic spines at the timepoints indicated. Numbers indicate the number of animals quantified at each timepoint.

(C)   Fluorescent images of ACR-12/AChR clusters (P*flp-13*::ACR-12::GFP) in wild type (left) and *nrx-1(wy778)* mutants (right) 16, 20, and 24 hours after hatch and at L4 stage (~42-50 hours post-hatch). Immature AChR clusters initially form but then disperse in *nrx-1* mutants.

(D)   Quantification of the number of ACR-12::GFP clusters in wild type (black) and *nrx-1(wy778)* mutants (purple). Two-way ANOVA, Sidak's multiple comparisons, *$p<0.05$, **$p<0.01$, ***$p<0.001$.

(E)   Fluorescent confocal images of dendritic spines (P*flp-13*::mCherry) and F-actin (P*flp-13*::GFP::UtrCH) in wild type and *nrx-1(wy778)* animals at 16 hours after hatch and L4 stage.

A NRX-1::GFP (knock-in)

wild type

unc-104(e1265)

10 μm

B *ACh*::NRX-1::GFP

wild type

unc-104(e1265)

+ P*ACh*::unc-104

10 μm

C

% of control (number of puncta)

****

NRX-1::GFP (knock-in)
P*ACh*::NRX-1::GFP
+ P*ACh*::unc-104 rescue
CLA-1::GFP
UNC-10::GFP
ELKS-1::mCherry

unc-104(e1265)

D *ACh*::CLA-1::GFP

wild type

unc-104(e1265)

10 μm

E *ACh*::UNC-10::GFP

wild type

unc-104(e1265)

10 μm

F *ACh*::ELKS-1::mCherry

wild type

unc-104(e1265)

10 μm

G Spines (*GABA*::myrGFP)

wild type

unc-104(e1265)

+ P*ACh*::unc-104

5 μm

H AChR (*GABA*::ACR-12::GFP)

wild type

unc-104(e1265)

+ P*ACh*::unc-104

5 μm

I

Normalized number (%)

**** **** ****

wild type
unc-104(e1265)
native
ACh
GABA
Muscle

unc-104(e1265)

spines
rescued lines/total

native:	3/3
ACh:	3/3
GABA:	0/4
muscle:	0/3

AChR/ACR-12 clusters
rescued lines/total

native:	4/4
ACh:	3/4
GABA:	0/3
muscle:	0/3

J Spines
(*GABA*::mCherry or *GABA*::myrGFP)

wild type

nrx-1(wy778)

cla-1(ok937)

unc-10(md1117)

elks-1(ok2762)

5 μm

Figure 4. UNC-104/KIF1A motor-dependent transport of NRX-1 to presynaptic terminals is required for stabilization of postsynaptic structures.

(A)   Representative confocal images of NRX-1::GFP (endogenous *nrx-1* knock-in) from dorsal cords (inverted LUT) of wild type and *unc-104(e1265)* mutants. Images on each line are from different animals (5 shown for each genotype). Mutation of *unc-104* impairs axonal NRX-1:GFP localization.

(B)   Representative confocal images of cholinergic NRX-1::GFP (P*unc-129*::NRX-1::GFP, inverted LUT) from the dorsal nerve cord of wild type, *unc-104(e1265)* and *unc-104(e1265)* mutants rescued by cholinergic expression of wildtype *unc-104* (*Punc-17*β). Images on each line are from different animals (5 shown for each genotype).

(C)   Scatterplot of the average number of NRX-1::GFP (endogenous or transgenic), CLA-1::GFP, UNC-10::GFP, or ELKS-1::mCherry clusters per 50 μm of dorsal nerve cord in *unc-104(e1265)* mutants or rescue as indicated. Values are normalized to their respective controls. Red dotted line indicates 50% of wild type values. Bars indicate mean ± SEM. Student's t-test, ****$p<0.0001$. n ≥ 11 animals. Raw values with respective controls are shown in Figure S4.6.

(D)   Representative confocal images of cholinergic CLA-1 (P*unc-17*β::CLA-1::GFP, inverted LUT) from the dorsal nerve cord of wild type or *unc-104(e1265)* mutants. Images on each line are from different animals (5 shown for each genotype).

(E)   Representative confocal images of cholinergic UNC-10 (P*unc-129*::UNC-10::GFP, inverted LUT) from dorsal nerve cords of wild type and *unc-104(e1265)* mutants. Images on each line are from different animals (5 shown for each genotype).

(F) Representative confocal images of cholinergic ELKS-1 (P*unc-129*::ELKS-1::mCherry, inverted LUT) from dorsal nerve cords of wild type and *unc-104(e1265)* mutants. Images on each line are from different animals (5 shown for each genotype).

(G) Fluorescent images (inverted LUT) of DD dendritic spines (P*flp-13*::myrGFP) from wild type, *unc-104(e1265)*, and *unc-104(e1265)* mutants rescued by cholinergic expression of wild type *unc-104*.

(H) Fluorescent images of AChR/ACR-12 clusters in DD dendrites (P*flp-13*::ACR-12::GFP) of wild type, *unc-104(e1265)*, and *unc-104(e1265)* mutants rescued by cholinergic expression of wild type *unc-104*.

(I) Scatterplot showing quantification of DD dendritic spines per 15 μm (P*flp-13*::myrGFP) (purple) and DD cholinergic receptor clusters (P*flp-13*::ACR-12::GFP) (green) in wild type, *unc-104(e1265),* and indicated cell-specific rescue lines. Inset, number of rescuing lines/total transgenic lines tested for each rescue construct. Bars, mean ± SEM. One-way ANOVA, Dunnett's multiple comparisons test, ****$p < 0.0001$. n ≥ 10 animals.

(J) Fluorescent images of DD dendritic spines (P*flp-13*::myrGFP or P*flp-13*::mCherry) from wild type, *nrx-1(wy778)*, *cla-1(ok937)*, *unc-10(md1117)*, or *elks-1(ok2762)* mutants. Only mutation of *nrx-1* affects dendritic spines.

Figure 5. UNC-104/KIF1A transport of NRX-1 is required in the mature circuit to maintain postsynaptic structure.

(A)   Cartoon depiction of experimental timeline. Animals were grown at 13.5°C until L4 stage (approx. 144 hours in *unc-104(ce782)ts* mutants, 120 hours in wild type

animals) before shifting animals to the restrictive temperature of 25°C for 16-20 hours and imaging.

(B)   Line scan fluorescent images of NRX-1::GFP (knockin to endogenous locus, inverted LUT) from dorsal cords of wild type and *unc-104(ce782)* mutants grown continuously at the permissive temperature (13.5°C) (middle panel) or shifted to the restrictive temperature (25°C) at L4 stage for 16-20 hours before imaging (bottom panel). Each line scan indicates an individual animal.

(C)   Quantification of NRX-1::GFP clusters along a 50 μm region of the dorsal nerve cord in wild type and *unc-104(ce782)ts* animals. Two-way ANOVA, Tukey's multiple comparisons test, ns, not significant, *$p<0.05$, n ≥ 11 animals. Data points indicate mean ± SEM.

(D)   Fluorescent images of DD spines in wild type and *unc-104(ce782)ts* animals grown continuously at the permissive temperature (13.5°C) (middle panel) or shifted to the restrictive temperature (25°C) at L4 stage for 16-20 hours before imaging.

(E)   Quantification of the number of spines per 15 μm in wild type and *unc-104(ce782)ts* animals. Two-Way ANOVA, Tukey's multiple comparisons test, ns, not significant, **$p<0.001$. n ≥ 13 animals. Data points indicate mean ± SEM.

Supplemental Figures

Figure S1.1 Ultrastructure of DD dendritic spines.

(A)   Cartoon depiction of the anterior serial cross-sections used for electron microscopy studies of the ventral nerve cord, focusing on the dendrite of the DD1 neuron.

(B)   Serial cross-sections (225/491, 226/491, 227/491, 228/491) of the ventral nerve cord. Teal indicates DD1 dendrite and dendritic spines (226/491 and 227/491) dipping into the ventral nerve cord to meet presynaptic cholinergic terminals (presumably VA/VB neurons, magenta). Asterisks indicates mitochondria within the dendritic shaft, see Figure S1.4. Scale bar, 100 nm.

(C)   Serial cross-sections (443/597, 444/597, 445/597, 446/597) of the ventral nerve cord of John White's N2U series (J. G. White et al., 1976). Teal indicates DD1 dendrite and dendritic spines dipping into the ventral nerve cord to meet presynaptic cholinergic terminals (magenta) (sections 444-446). Asterisks indicates mitochondria within the dendritic shaft, see Figure S1.4.

(D)   Ventral nerve cord electron micrographs. Dendritic spines (teal) have varied morphology and sizes. Two representative micrographs from serial sections in which the entire extent of the spine is visible are shown. Asterisk indicates mitochondrion within the dendritic shaft. Also see Figure S1.4B. Scale bar, 100 nm.

Figure S1.2. Microtubules localize at the base of spines.

Fluorescent images of tubulin (P*flp-13*::TBA-1::GFP) and F-actin (P*flp-13*::mCherry::UtrCH) in DD neurons. White arrows indicate dendritic regions where tubulin dips into the base of a dendritic spine.

Figure S1.3. Cartoon illustration of calcium imaging recording platform.

Imaging was carried out using a Yokogawa CSU-X1-A1N spinning disk confocal system (Perkin Elmer) equipped with EM-CCD camera (Hamamatsu, C9100-50) and 63X oil immersion objective. Chrimson photoactivation (~30 mW/cm^2) was achieved using a TTL-controlled 625 nm light guide coupled LED (Mightex Systems), permitting illumination of the entire immobilized animal, while simultaneously recording GCaMP6f fluorescence (excitation 488 nm, emission 525 nm). 625 nm light was prevented from reaching the

camera using a 556 nm edge BrightLine single-edge short-pass dichroic beam splitter positioned in the light path (Semrock).

A **Golgi**

GABA::mCherry

GABA::AMAN-2::GFP

merge

5 μm

B **Mitochondria**

GABA::mCherry

GABA::pre-Su9::GFP

merge

5 μm

C

pre-Su9::GFP fluorescence intensity / total fluorescence (AU)

****

dendritic shaft   dendritic spines

D **Rough ER**

GABA::myrGFP

GABA::RFP::TRAM-1

merge

5 μm

E

TRAM-1::RFP fluorescence intensity / total fluorescence (AU)

***

dendrite shaft   dendritic spines

Figure S1.4. Examination of fluorescent organelle markers and F-actin interacting proteins in DD dendrites and dendritic spines.

(A)   Fluorescent images of DD1 dendritic spines (P*flp-13*::mCherry) and Golgi marker (P*flp-13*::AMAN-2::GFP) in DD neurons. Golgi apparatus is primarily localized to the DD1 soma. Dotted white line traces the neuronal cell body.

(B)   Fluorescent images of DD1 dendritic spines (P*flp-13*::mCherry) and mitochondria (P*flp-13*::pre-Su9::GFP). Mitochondria are localized within the main dendritic process of DD neurons near spines.

(C)   Quantification of mitochondria fluorescence intensity in dendritic shaft and dendritic spines. Bars, mean ± SEM. Student's t-test, ****$p<0.0001$.

(D)   Fluorescent images of DD1 dendritic spines (P*flp-13*::myrGFP) and rough endoplasmic reticulum (ER) (P*flp-13*::TRAM::RFP). Dotted white line traces the neuronal cell body. ER is primarily localized within the main dendritic process of DD neurons and occasionally at the spine base.

(E)   Quantification of rough ER fluorescence intensity (P*flp-13*::RFP::TRAM-1) in dendritic shaft and dendritic spines. Bars, mean ± SEM. Student's t-test, ***$p<0.001$. n=9 animals.

A neuron reporter

16 hours

DD1
DD2
VB2
DB1 DB2
DB3
VB1

DD neurons::mCherry (P*flp-13*::mCherry)
ACh::GFP (P*acr-5*::GFP)
post-embryonic born
VB cholinergic neurons

24 hours

DD1
DB1
VB3
DB3
DD2
VB2
VB1 DB2

5 μm

B synaptic vesicles

12 hours

DD1
DB1
DB2
DB3
DD2

DD neurons::mCherry (P*flp-13*::mCherry)
ACh synaptic vesicles::GFP
(P*acr-5*::SNB-1::GFP)
post-embryonic born
VB cholinergic neurons

16 hours

DD1
VB1
DB2
DB1
DB3
DD2

24 hours

DB2 DD1
DB1
VB3
DD2
DB3

5 μm

C

12 hours after hatch

DB2 DD1
DB1
DB3
DD2

16 hours after hatch

VB2
VB1
DB2 DD1
DB1
DB3
DD2

24 hours after hatch

VB2
VB1
DB2
DD1
DB1
VB3
DB3
DD2

synaptic
vesicles

dendritic
spines

embryonic
born ACh
neurons

embryonic
born GABAergic
neurons

post-embryonic ACh
neurons

Figure S2.1. Development of post-embryonic born ventral cholinergic motor neurons.

(A)   Fluorescent images of B-type cholinergic neurons (DB/VB) (P*acr-5*::GFP) (green) and DD GABAergic neurons (P*flp-13*::mCherry) (red) at 16 and 24 hours after hatch. VB cholinergic neurons are born post-embryonically in an anterior to posterior order (blue). Note that at 16 hours after hatch P*acr-5*::GFP fluorescence indicating VB1 and VB2 cell bodies is visible. By 24 hours after hatch P*acr-5*::GFP fluorescence indicating VB3 is visible. White dotted circles outline the neuronal cell bodies.

(B)   Fluorescent images of the anterior ventral nerve cord in animals co-expressing the synaptic vesicle marker P*acr-5*::SNB-1::GFP in B-type cholinergic neurons (DB/VB) (green) with P*flp-13*::mCherry labeling DD GABAergic neurons (red) at 12,16, and 24 hours after hatch. At 12 hours, little SNB-1::GFP fluorescence is visible in the ventral nerve cord. Embryonic born, dorsally directed B-type (DB) cholinergic motor neurons predominantly make synaptic contacts in the dorsal nerve cord. Ventrally-directed, B-type (VB) cholinergic motor neurons are born post-embryonically and have not yet completed their maturation at this time. SNB-1::GFP fluorescence in the ventral nerve cord is faintly visible by 16 hrs after hatch and becomes more prominent by 24 hrs after hatch, coincident with maturation of VB motor neurons. White dotted circles indicate outlines of neuronal cell bodies. Yellow arrows indicated presynaptic vesicle clusters, SNB-1.

(C)   Cartoon representation of the developmental timing of post-embryonic born ventral cholinergic motor neurons, spine outgrowth, and cholinergic synaptic vesicle localization.

Figure S2.2. Spine outgrowth progresses anteriorly to posteriorly.

(A and B) Fluorescent images (inverted LUT) of DD1 and DD2 dendritic spines 24 hours

after hatch (A) and at L4 (~42-50 hours after hatch) stage (B). Animals express P*flp-*

13::mCherry to label dendritic spines. Red dashed rectangle indicates inset anterior

to DD1 soma. Blue dashed rectangle indicates inset anterior to DD2 soma. Note that

DD1 spines form prior to DD2 spines.

(C) Quantification of the length of growing dendritic spines at 24 and 32 hours after hatch

and at L4 (~42-50 hours after hatch) stage. Data points indicate mean ± SEM.

Figure S2.3 LEV-10 localizes to putative postsynaptic sites prior to spine formation.

(A)   Fluorescent confocal images showing dendritic spines (magenta) and the postsynaptic CUB transmembrane domain LEV-10 (cyan). Animals express P*flp-13*::mCherry with DD neuron specific LEV-10::GFP11$_{x7}$ using a strategy for cell-specific labeling of endogenous LEV-10 (NATF) (He et al., 2019)

(B)   Quantification of the number of DD dendritic spines and LEV-10 clusters at 16, 20, and 24 hours after hatch, and L4 stage (~42-50 hours post-hatch).

Figure S2.4 Actin dynamics during development.

(A,C,E) Line scans displaying relative fluorescence intensity of F-actin (P*flp-13*::GFP::UtrCH) at 16 hours post-hatch (A), 24 hours post-hatch (C), and L4 (~42-50 hours post-hatch) animals (E). Each color indicates a line scan of fluorescence intensity for the same DD dendrite ROI acquired at 5-minute intervals. Note the variable distribution of fluorescence intensities across line scans from images acquired near 16 hours after hatch compared to later time points, indicating increased F-actin dynamics during early development.

(B,D,F) Confocal images (inverted LUT) showing P*flp-13*::GFP::UtrCH fluorescence (labeling F-actin) in the DD dendrite of 16 (B), or 24 (D) hours after hatch, or at L4 stage (~42-50 hours post-hatch) (F). For each, sequences of fluorescent images separated by 30 minutes are shown.

Video 1: Confocal live imaging video of F-actin dynamics (labeled by P*flp-13*::GFP::UtrCH) in the DD1 dendrite of wild type at 16 hours after hatch. For display, movies are shown at 3 fps. Images were acquired every 5 minutes.

Video 2: Confocal live imaging video of F-actin dynamics (labeled by P*flp-13*::GFP::UtrCH) in the DD1 dendrite of wild type at L4 stage. For display, movies are shown at 3 fps. Images were acquired every 5 minutes.

Figure S3.1. Spines and AChR localize independently of presynaptic activity.

(A,B) Fluorescent confocal images (inverted LUT) of DD1 dendritic spines (P*flp-13*::mCherry (A) or P*flp-13*::myrGFP (B)) in L4 stage wild type and selected mutant strains where synaptic activity is affected.

(C) Fluorescent confocal images of AChR clusters (P*flp-13*::ACR-12::GFP) in the DD dendrite of L4 stage wild type and selected mutant strains where synaptic activity is affected.

Figure S3.2. Dendritic calcium responses are severely reduced in *nrx-1* mutants.

Scatter plot showing peak $\Delta F/F_0$ responses measured from GABAergic DD motor neuron dendrites during a 5s period cholinergic photostimulation stimulation in wild type, *acr-16(ok789); acr-12(ok367)*, *nrx-1(wy778), nrx-1(nu485),* and *nrx-1(ok1649)* mutants. All genotypes co-express P*flp-13*::myrGCaMP6f::SL2::mCherry for measurement of dendritic calcium responses with P*acr-2::*Chrimson for cholinergic neuron depolarization. Bars indicate mean ± SEM. One-way ANOVA, Dunnett's multiple comparisons, ****$p<0.0001$. Wild type control is the same as Figure 1J. n ≥ 10 animals.

Figure S3.3. Spine and AChR localization in developing wild type and *nrx-1(wy778)* mutants.

(A)   Quantification of the number of DD spines/15 μm in wild type (black) and *nrx-1(wy778)* (purple) animals 16, 20, and 24 hours after hatch and at L4 stage (~42-50 hours post-hatch). Two-way ANOVA, Sidak's multiple comparisons test, **p<0.01, ****p<0.0001. Data points indicate mean ± SEM. Numbers indicate animals quantified for each timepoint.

(B)   Quantification of ACR-12::GFP fluorescence intensity in wild type (black) and *nrx-1(wy778)* (purple) animals at 16, 20, and 24 hours after hatch and L4 stage (~42-

50 hours post-hatch). Two-way ANOVA, Sidak's multiple comparisons test, **$p<0.01$, ***$p<0.001$. Data points indicate mean ± SEM.

Figure S3.4. Spine dynamics during development in wild type and *nrx-1(wy778)* mutants.

(A)   Quantification of the percentage of spine events in wild type and *nrx-1(wy778)* animals at 16-20 hours after hatch, 21-24 hours after hatch, and at L4 compared to the total number of events. An event is defined as an incidence of spine destabilization/pruning, elongation/formation, or a stable spine during an entire live imaging recording. Z-stacks of DD1 dendrites were acquired every 5 minutes for

60-160 minutes. Numbers indicates number of events for each genotype at each timepoint.

(B) Fluorescent image (inverted LUT) of DD dendritic spines at timepoint zero in a wild type animal 16-20 hours after hatch. Dashed red box indicates region shown in Figure S3.4C.

(C) Fluorescent images (inverted LUT) of an individual wild type DD dendritic spine (P*flp-13*::mCherry) at 0, 45, and 70 minute timepoints during live imaging of area indicated by red box in Figure S3.4B. Red dotted line outlines spine dynamics.

(D) Fluorescent image (inverted LUT) of *nrx-1(wy778)* mutant DD1 dendritic spines (P*flp-13*::mCherry) at timepoint zero acquired 16-20 hours after hatch. Dashed red box indicates region shown in Figure S3.4E.

(E) Fluorescent images (inverted LUT) of an individual *nrx-1(wy778)* mutant DD dendritic spine (P*flp-13*::mCherry) at 90. 95, and 100 minute timepoints during live imaging of area indicated by red box in Figure S3.4D. Red dotted lines outline spine dynamics.

(F) Fluorescent image (inverted LUT) of wild type DD dendritic spines (P*flp-13*::mCherry) at timepoint zero at L4 stage. Dashed red box indicates region shown in Figure S3.4G.

(G) Fluorescent images (inverted LUT) of an individual wild type DD dendritic spine (P*flp-13*::mCherry) at 0, 30, and 65 minute timepoints during live imaging of the area indicated by red box in Figure S3.4F.

Video 3. Wild type DD dendritic spines (P*flp-13*::myrGFP) 16 hours after hatch. A z-stack was acquired every 5 minutes for at least 1 hr of imaging duration. For display, movies are shown at 3 fps

Video 4. *nrx-1(wy778)* mutant DD dendritic spines (P*flp-13*::myrGFP) 16 hours after hatch. A z-stack was acquired every 5 minutes for at least 1 hr of imaging duration. For display, movies are shown at 3 fps

Video 5. Wild type DD dendritic spines (P*flp-13*::myrGFP) at L4 stage. A z-stack was acquired every 5 minutes for at least 1 hr of imaging duration. For display, movies are shown at 3 fps

Figure S4.1. Mutation of *unc-104* impairs synaptic vesicle delivery to cholinergic axons in the dorsal nerve cord.

(A)   Fluorescent images of cholinergic synaptic vesicles (P*acr-2*::mCherry::RAB-3) in the dorsal nerve cord of young adult wild type and *unc-104(e1265)* mutants. Images on each line are from different animals (4 are shown for each genotype).

(B)   Quantification of cholinergic synaptic vesicles (P*acr-2*::mCherry::RAB-3) per 50

        µm of the dorsal nerve cords of wild type and *unc-104(e1265)* mutants. Student's

        t-test, ****$p<0.0001$. Bars, mean ± SEM.

(C)   Fluorescent images of the soma in wild type and *unc-104(e1265)* mutants

        expressing cholinergic vesicle reporter (P*acr-2*::mCherry::RAB-3). Dotted white

        lines outline the cell body.

(D)   Quantification of soma cholinergic synaptic vesicle fluorescence intensity (AU)

        (P*acr-2*::mCherry::RAB-3) of wild type and *unc-104(e1265)* mutants. Student's t-

        test, ****$p<0.0001$. Bars, mean ± SEM.

Figure S4.2. NRX-1 colocalizes with UNC-104 in cholinergic axon terminals.

(A)   Line scans depicting relative fluorescent intensity of NRX-1::GFP (green) and

UNC-104::mCherry (red) for a 50 µm region of the dorsal nerve cord. Gray dotted

rectangles indicate corresponding puncta in Figure S4.2B.

(B)   Fluorescent images of the dorsal nerve cord in an adult animal expressing NRX-

1::GFP (P*unc-129*::NRX-1::GFP) and UNC-104::mCherry (P*unc-129*::UNC-

104::mCherry). Colocalization is indicated by white carets.

Figure S4.3. NRX-1 endogenously tagged with GFP localizes to neuronal processes.

NRX-1::GFP localizes within neuronal processes of the nerve ring and shows punctate

localization in processes of the ventral and dorsal nerve cords of L4 stage worms.

NRX-1::GFP (knock-in)

ACh::NRX-1::GFP

Figure S4.4. NRX-1 localization to cholinergic axons is decreased in _unc-104_ mutants.

(A)   Quantification of NRX-1::GFP (endogenous knock-in) clusters in a 50 μm region of the dorsal nerve cord in wild type and _unc-104(e1265)_ animals. Bars, mean ± SEM. Student's t-test, ****_p_<0.0001. n ≥ 11 animals.

(B)   Quantification of NRX-1::GFP (endogenous knock-in) axon fluorescence intensity in a 50 μm region of the dorsal nerve cord. Bars, mean ± SEM. Student's t-test, ****_p_<0.0001. n ≥ 11 animals.

(C)   Fluorescent images of NRX-1::GFP (endogenous knock-in) in somas of wild type and *unc-104(e1265)* mutants. Dotted white lines outline the neuronal cell body.

(D)   Quantification of NRX-1::GFP soma fluorescence intensity. Bars, mean ± SEM. Student's test, ****$p<0.001$.

(E)   Quantification of the number of NRX-1::GFP clusters in a 50 μm region of the dorsal nerve cord in wild type, *unc-104(e1265)*, and *unc-104(e1265)* mutants rescued with cholinergic expression of wild type *unc-104*. Bars, mean ± SEM. One-way ANOVA, Dunnett's multiple comparisons test, ****$p<0.0001$. n ≥ 11 animals.

(F)   Quantification of NRX-1::GFP fluorescence intensity in cholinergic axons of a 50 μm region of the dorsal nerve cord. Bars, mean ± SEM. One-way ANOVA, Dunnett's multiple comparisons test, ****$p<0.0001$. n ≥ 11 animals.

(G)   Fluorescent images of NRX-1::GFP in cholinergic somata of wild type, *unc-104(e1265)* and *unc-104(e1265)* mutants rescued with cholinergic expression of wild type *unc-104*. Dotted white lines outline the neuronal cell body.

(H)   Quantification of NRX-1::GFP fluorescence intensity in cholinergic somata. Bars, mean ± SEM. One-way ANOVA, Dunnett's multiple comparisons test, ****$p<0.001$.

Figure S4.5. NRX-1 localization in *unc-116*/KIF5A/C mutants.

(A)   Fluorescent images (inverted LUT) of NRX-1::GFP (P*unc-129*::NRX-1::GFP) from the dorsal nerve cord of wild type and *unc-116(e2310)* mutants. Images on each line are from different animals (5 are shown for each genotype).

(B)   Quantification of the number of NRX-1 clusters (P*unc-129*::NRX-1::GFP) in a 50 μm region of the dorsal nerve cord of wild type and *unc-116(e2310)* animals. Bars, mean ± SEM. Student's t-test, ***$p<0.001$. n ≥ 14 animals.

(C)   Fluorescent images of NRX-1 (P*unc-129*::NRX-1::GFP) from cholinergic neuron somata of wild type and *unc-116(e2310)* mutants. Dotted white lines outline cell bodies.

(D)   Quantification of fluorescence intensity of NRX-1 (*Punc-129*::NRX-1::GFP) from cholinergic somas of wild type and *unc-116(e2310)* mutants. Bars, mean ± SEM. Student's t-test, ns, not significant.

Figure S4.6. NRX-1 and SNB-1 have similar anterograde and retrograde velocities.

(A, B) Kymographs of synaptic vesicle (P*unc-129*::SNB-1::GFP) **(A)** and NRX-1 (P*unc-129*::NRX-1::GFP) **(B)** trafficking events recorded from cholinergic neuron commissures.

(C)   Quantification of the anterograde velocity (μm/second) of SNB-1::GFP (blue) and

NRX-1::GFP (orange) along the axonal commissure. ns, not significant. n ≥ 10

animals for all panels.

(D)   Quantification of the retrograde velocity (μm /second) of SNB-1::GFP (blue) and

NRX-1::GFP (orange) along the axonal commissure. ns, not significant.

(E)   Quantification of the total number of SNB-1::GFP (blue) and NRX-1::GFP

(orange) trafficking events binned into retrograde and anterograde directions.

Note that NRX-1 trafficking events occur significantly less frequently than SNB-1

events.

Video 6: NRX-1::GFP trafficking in a cholinergic commissure of a young adult animal (P*unc-129*::NRX-1::GFP). Videos were recorded at 100 ms exposure for 30 seconds. For display, movie is shown at 25 fps.

Video 7: SNB-1::GFP trafficking in a cholinergic commissure of a young adult animal (P*unc-129*::SNB-1::GFP). Videos were recorded at 100 ms exposure for 30 seconds. For display, movie is shown at 25 fps.

Figure S4.7. Active zone protein localization in *nrx-1(wy778)* and *unc-104(e1265)* mutants.

(A)   Line scan fluorescent images (inverted, LUT) of active zone protein CLA-1 (*Punc-17β*::CLA-1-1::GFP, inverted) from cholinergic axons of dorsal nerve cords in wild type, *nrx-1(wy778)*, and *unc-104(e1264)* mutants. Images on each line are from different animals (5 are shown for each genotype). These data correspond to quantification in Figure 4C.

(B)   Scatterplot of CLA-1::GFP puncta in a 50 μm region of the dorsal nerve cord in wild type, *nrx-1(wy778)* and *unc-104(e1265)* animals. One-way ANOVA, Dunnett's multiple comparisons test, ****$p<0.0001$. Bars, mean ± SEM.

(C)   Scatterplot of CLA-1::GFP fluorescence intensity in a 50 μm region of the dorsal nerve cord in wild type, *nrx-1(wy778)* and *unc-104(e1265)* animals. One-way ANOVA, Dunnett's multiple comparisons test **$p<0.01$, ****$p<0.0001$. Bars, mean ± SEM.

(D)   Line scan fluorescent images (inverted, LUT) of active zone protein UNC-10 (*Punc-129*::UNC-10::GFP) from cholinergic axons of dorsal nerve cords in wild type, *nrx-1(wy778)* and *unc-104(e1265)* mutants. Images on each line are from different animals (5 are shown for each genotype). These data correspond to quantification in Figure 4C.

(E)   Quantification of UNC-10::GFP puncta in a 50 μm region of the dorsal nerve cord in wild type, *nrx-1(wy778)* and *unc-104(e1265)* animals. One-way ANOVA, Dunnett's multiple comparisons test, **$p<0.001$, ****$p<0.0001$. Bars, mean ± SEM.

(F)   Quantification of UNC-10::GFP fluorescence intensity in a 50 μm region of the dorsal nerve cord in wild type, *nrx-1(wy778)* and *unc-104(e1265)* mutants. One-way ANOVA, Dunnett's multiple comparisons test, ****$p<0.0001$. Bars, mean ± SEM.

(G)   Line scan fluorescent images (inverted, LUT) of active zone protein ELKS-1 (P*unc-129*::ELKS-1::mCherry, inverted) from cholinergic axons of dorsal nerve cords in wild type, *nrx-1(wy778)* and *unc-104(e1265)* mutants. Images on each line are from different animals (5 are shown for each genotype). These data correspond to quantification in Figure 4C.

(H)   Quantification of ELKS-1::mCherry puncta per 50 μm of the dorsal nerve cord in indicated genotypes. One-way AVOVA, Dunnett's multiple comparisons test, ****$p<0.0001$, ns, not significant. Bars, mean ± SEM.

(I)   Quantification of ELKS-1::mCherry fluorescence intensity in a 50 μm region of the dorsal nerve cords of wild type, *nrx-1(wy778)* and *unc-104(e1265)* mutants. One-way ANOVA, Dunnett's multiple comparisons test, ***$p<0.001$, ns, not significant. Bars, mean ± SEM.

Figure S5.1 Shift to restrictive temperature during L3 in wild type and *unc-104(ce782)ts* animals.

(A)   Cartoon depiction of experimental timeline. Animals were grown at 13.5°C until L3

stage (~approx. 120 hours in *unc-104(ce782)* mutants, 96 hours in wild type

animals) before shifting animals to their restrictive temperature of 25°C for 16-20 hours and imaging.

(B)   Fluorescent images of DD spines from wild type (top) or *unc-104(ce782)ts* animals grown continuously at the permissive temperature (13.5°C) (middle panel) or shifted to the restrictive temperature (25°C) for 16-20 hours before imaging.

(C)   Quantification of DD spines per 15 μm from wild type and *unc-104(ce782)ts* mutants. Two-way ANOVA, Tukey's multiple comparisons test, ****$p<0.0001$, n ≥ 12 animals. Data points indicate mean ± SeM.