1

Machine learning algorithms can predict tail biting outbreaks in pigs using feeding behaviour records

4
5

Catherine Ollagnier[1][*], Claudia Kasper[2], Anna Wallenbeck[3], Linda Keeling[3], Giuseppe Bee[1], Siavash A.

Bigdeli[4]

[1] Swine Research Unit, Agroscope, Posieux, Fribourg, Switzerland

[2] Animal GenoPhenomics, Agroscope, Posieux, Fribourg, Switzerland

[3] Department of Animal Environment and Health, Swedish University of Agricultural Sciences,

Uppsala, Sweden

[4] Edge Vision and AI, CSEM, Neuchâtel, Switzerland

*E-mail: catherine.ollagnier@agroscope.admin.ch

Abstract

Tail biting is a detrimental behaviour that impacts the welfare and health of pigs. Early detection of

tail biting precursor signs allows for preventive measures to be taken, thus avoiding the occurrence

of the tail biting event. This study aimed to build a machine-learning algorithm for real time

detection of upcoming tail biting outbreaks, using feeding behaviour data recorded by an electronic

feeder. Prediction capacities of seven machine learning algorithms (e.g., random forest, neural

networks) were evaluated from daily feeding data collected from 65 pens originating from 2 herds of

grower-finisher pigs (25-100kg), in which 27 tail biting events occurred. Data were divided into

training and testing data, either by randomly splitting data into 75% (training set) and 25% (testing

set), or by randomly selecting pens to constitute the testing set.

The random forest algorithm was able to predict 70% of the upcoming events with an accuracy of

94%, when predicting events in pens for which it had previous data. The detection of events for

unknown pens was less sensitive, and the neural network model was able to detect 14% of the

upcoming events with an accuracy of 63%. A machine-learning algorithm based on ongoing data

collection should be considered for implementation into automatic feeder systems for real time

prediction of tail biting events.

Keywords

32   Tail biting, pig, behavioural change, prediction, monitoring behaviour, animal welfare

Background

Tail biting (TB) is an abnormal behaviour in pigs that is thought to have a multi-factorial origin. It can

be triggered by factors like the lack of enrichment material, unfavourable environmental conditions,

an unbalanced diet or poor health status. In addition to the welfare and ethical concerns associated

with this cannibalistic behaviour, TB events cause pain, trigger infections, impair growth and devalue

the carcasses.[1-5]

Routine tail docking is prohibited in Switzerland (455.1 OPAn, 2008) and in the EU (Council Directive

2008/120/EC, Annex 1), and farmers are requested to set up measures to prevent TB outbreaks. One

strategy is to pinpoint the farm-specific risk factors for TB and to find solutions to reduce them.[4, 6]

Another strategy is to monitor animals' behaviours to detect early signs of forthcoming outbreaks.[7-

9] Early identification of TB indicators are important for efficient intervention. The behavioural

monitoring can be done at the pen and at the individual animal level. Identification at the individual

animal level can support preventive measures like for instance, by removing the biter or the bitten

pigs. Observations at the pen level are more efficient to detect the TB event.[1]

To date, only a few behavioural indicators were studied at the pen level. These early indicators, like

changes in activity level, tail posture and exploratory and feeding behaviour are present up to 63

days before the outbreak starts. Statham et al. reported that pigs spend less time lying and more

time standing and sitting within 4 days before an outbreak[10], but this difference in postures was

not observed by Wedin et al.[7] Lower tail position seems also to be indicative of an outbreak, and

several authors have reported increased incidence of tucked or hanging tails in pens before and

during TB outbreaks.[7-10,12] Lately, an automated system was developed that uses 3D cameras to

detect predictive indicators of TB.[8] With this new tool, the authors were able to automatically

detect low tail posture, which was positively associated with more tail damage. However, the tail

posture may not be very specific to a TB event, as a low tail posture has also been connected to

negative emotional status in pigs [7], which can be caused by other factors like sickness. Exploratory

behaviours could also be used as early indicators of TB events. For Zonderland et al., the amount of

explorative behaviours decreased 6 days before an event[11], whereas for Statham et al., the

environmental manipulations increase one day before the TB events.[10] However, Ursinus et al.

couldn't find any change in the frequency of explorative behaviours prior to TB events.[14]

Electronic feeding systems with single-spaced feeders enable automatic monitoring of the feeding

behaviours of each pig individually. Recording the identity of the pig, feeder entry and exit times and

the amount of food consumed allow the calculation of the frequency of feeder visits per day, feeding

rates, mean feeder occupation time, mean food intake per feeder visit, total food intake and total

feeder duration per day for each pig. In 1994, Young and Lawrence found that pigs housed in groups

and fed from automatic feeders showed a temporal patterning of feeding behaviour.[15] They also

suggested that the feeding behavior might also be altered by social conditions. It has been later

described that changes in feeding behaviours with automatic feeders were associated with negative

events like aggressive behaviour or disrupted social dynamics.[8] If a TB event can be predicted from

behaviour, as postulated by Statham et al. [9], then data from electronic feeders could be used to

monitor in real time the feeding behaviour of the pigs. Indeed, the feeding behaviour of pigs

assessed by electronic feeders appears to change before a TB event. Wallenbeck and Keeling,

Munsterhjelm et al. and Tessier et al. describe changes in feeding behaviours before TB events.[17-

19] These findings suggest that feeding behaviours recorded by electronic feeders could be a valid

tool to detect early signs of TB event. Therefore, we aimed to use the feeding behaviour data

combined with machine learning (ML) algorithms for real time prediction of TB outbreak. If

successful, this will be the first demonstration that a model can predict TB in real time. Such a

monitoring tool would open up new opportunities for the farmer to take targeted action in specific

pens to prevent the TB event. Being able to prevent TB would serve the welfare of the animals and

provide economic benefits to the farmer. Since the tool requires only data that are already available

from pig farms equipped with automatic feeders, it could be easily implemented in commercial

practice as an additional management tool.

Material and methods

Data collection

This study analyses the feeding behaviours of two herds of grower-finisher pigs weighing between 25

and 100 kilograms. One data set originates from a testing boar station in Sweden and contains data

collected from October 2004 to July 2007. The data set comes from a previous retrospective study

that was published by Wallenbeck and Keeling.[10] The second data set contains data from the

experimental pig farm of Agroscope and comprises recordings from November 2018 to April 2020. As

tail docking is prohibited in Sweden and in Switzerland, the data are from pigs with intact tails.

The Swedish data set includes data from 42 pens (21 TB and 21 CTL) of boars (purebred Yorkshire,

Landrace or Hampshire), recorded 70 days before and after the TB date. Boars were housed in

groups of 7 to 14 animals per pen. Each pen measured 15.7 m^2 and had a slatted floor and plain

resting area. All pigs had *ad libitum* access to the pelleted feed, which was optimised according to the

Swedish nutrition norms for fattening pigs.[11] Water was provided *ad libitum* and straw was offered

daily.

The Swiss data set consisted of 23 pens (6 tail biting and 17 control) of females and castrated male

pigs (Swiss Large White), recorded 100 days before and after the TB date. Twenty pens (18 m^2)

contained 11 to 15 pigs and were equipped with two automatic feeders; three pens (78 m^2) were

equipped with eight automatic feeders for 31 to 55 pigs. All pens had straw in racks and sawdust on

the floor. Water was available *ad libitum* through nipple drinkers. The pelleted finisher diet was

formulated to have 20% lower dietary crude protein and essential amino acids compared to a

standard diet formulated according to the Swiss feeding recommendations for pigs.[12]

For both study sites, data were collected by individual automatic feeders (ACEMO 48, Acemo, France;

or MLP, Agrotronic Schauer, Austria) that recorded the number of visits to the feeder and the

amount of feed consumed. The feeders were 0.6 m wide and 1.5-2.2 m long. Only one pig could

enter the feeder at a time, and other pigs could not dislodge the pig feeding inside the feeder. Each

pig had access to only one feeder.

A pen was assigned to the TB category if at least one pig had to be treated for tail damages. The TB

date (day 0) was defined as the date at which the first treatment was recorded. For the Swedish data

set, each pen in the TB category was paired to a pen in the control (CTL) category. For the Swiss data

set, all the 23 farrowing batches reared under the same housing and feeding conditions were

considered for analysis.

Definitions of the observations, analysis in time series and missing value imputation

The frequency of daily feeder visits (DFV), the daily feed consumption (DFC) and the standard

deviation of the feed consumption at each visit (StdFC) were calculated per day and per pig (Table 1).

These parameters were considered as 'observations' to predict TB events at the pen level.

Table 1: Observations used for tail biting predictions
The frequency of daily feed visit, daily feed consumption and standard deviation of the feed consumption were the observations used to predict tail biting (TB) or control (CTL) classes with the ML models. These parameters were derived from the data collected by the automatic feeder.

	definition	units	abbreviated
Frequency of daily feeder visits	Number of visits to the feeder (from 0:00 to 23:59:59 that date)	n	DFV
Daily feed consumption	Total feed consumption (from 0:00 to 23:59:59 that date)	g	DFC
Standard deviation of the feed consumption	Daily standard deviation of the feed consumption at each visit	g	StdFC

The time dependency of the observations was taken into account by analysing the data by groups of

consecutive data points, called the 'analysis window'. The prediction model considered the analysis

window to achieve a prediction at the pen level. The analysis window was first defined to contain

observations from 14 consecutive days (Fig. 1). The A_date was defined as the first day of the

analysis window. The analysis window slides along the timeline and the A_date defines the class of

the analysis window, i.e., "TB" or "CTL". Analysis windows from control pens were always classified

as "CTL", whereas analysis windows from TB pens were considered as "TB" class only between day -

35 and day 10 (TB window). The analysis window of a TB pen was classified as "TB" if the A_date was

inside the TB window and "CTL", if outside the TB window. Missing values were computed using

median imputation (by data set) and a principal component analysis was performed before ML

analysis.

In each pen, the analysis window contained observations from 10 pigs for 14 days, to standardize the

size of the analysis window. Observations from 10 pigs was considered to give enough information

on the pen, without creating too many missing data points, for the few occurrences that contained

fewer than 10 pigs.

At the end of the data framing, each analysis window contained 420 observations {3 variables (DFV,

DFC, StdFC) × 10 pigs × 14 days}, and one outcome (the class of the window: "TB" or "CTL"). The

Swiss and the Swedish data set were merged into a third data set, called Swedish+Swiss, to

incorporate more diverse observations and further increase the model's generalizability for unseen

data (pen/country). In total, the combined data set (Swedish and Swiss data) contained 6605 analysis

windows, with 5479 and 1126 "CTL" and "TB" windows, respectively. The characteristics of the TB

windows compared to CTL are presented in Table 2.

Table 2: Characteristics of the data sets. Mean and standard deviation (SD) of the observation are
calculated for each data set depending on the class (TB, tail biting, or CTL, control) of the analysis
window. The total data set (Swedish+Swiss data set) contained 6605 analysis windows, with 5479
and 1126 CTL and TB windows, respectively. Pigs characteristics (age, and bodyweight) are not
balanced between analysis windows, thus the differences between CTL and TB windows should not
be considered as traits depicting the change in feeding behavior between a control and a TB pen.

		Swedish		Swiss	
Observations		**CTL**	**TB**	**CTL**	**TB**
DFC[1]	**Mean**	2337.1*	2005.7	2282.3*	2438.8
	SD	757.4	708.6	600.2	678.2
DFV[2]	**Mean**	24.9*	24.0	12.3*	12.0
	SD	19.4	18.0	6.0	7.2
StdFC[3]	**Mean**	128.6*	113.8	166.8*	195.6
	SD	85.9	71.2	96.3	129.7

148 *significant difference between CTL and TB analysis window classes (p<0.0001)
149 DFC[1]: frequency of daily feeder visits
150 DFV[2]: daily feed consumption
151 StdFC[3]: standard deviation of the feed consumption at each visit
152

Models

All data were analysed with R3.6.3, using the caret package to build ML models.[13]

Commonly used classification ML methods were first tested on all three data sets (i.e., Swedish, Swiss

and Swedish+Swiss). Table 3 presents a list of ML methods considered and implemented with R

packages in this study.

Table 3: The seven machine learning methods used to predict tail biting events from feeding
behavior data
Common ML approaches were tested in this classification problem with a binary outcome

Machine Learning	R Package	Function
Generalized Linear Model with Stepwise Feature Selection	glmnet	glmnet
Random forest	ranger	rf
Support Vector Machines with Radial Basis Function Kernel	kernlab	svmRadial
Bayesian Generalized Linear Model	arm	bayesglm
Neural network	nnet	nn
K-nearest neighbor	caret	kNN
Partial Least Squares Discriminant Analysis	pls	pls

For each data set, predictive models were first trained on a subset of data (training set), and the

models' performances on this training set were then compared. The predictive performances of the

models were further compared using the unseen data (test set). The test set contained either new

unseen analysis windows (cross validation, or CV, approach) or new pens (leave one out cross

validation, or LOOP, approach).[14] The LOOP approach gives estimate metrics that are valuable

when a new pen is presented to the model for prediction, as there is no pen overlapping between

the training and testing data set. This represents the situation where the farmer tries to predict a TB

event in a hitherto unknown pen. However, the CV resampling approach predicts TB events based on

data previously recorded in the pen. This approach is correct when the model can be continuously

updated with previous records of the pen, so that the prediction model already knows the feeding

behaviour of the pen and tries to classify the analysis window of the testing set based on previous

knowledge on this pen.

Model evaluation: performances metrics

This is a classification problem with binary outcomes (TB or CTL), and performances of the models

should be assessed on parameters calculated with a confusion matrix.[15] Performance metrics

definitions and confusion matrix are presented in Table 4. The sensitivity (rate of predicted TB class

given the actual TB class) assesses the capacity of the model to detect an upcoming TB event. The

positive predictive value (PPV) evaluates the capacity of the model to correctly predict a TB class. All

models were optimized to maximize the sensitivity, as this study aimed to detect early warnings of

TB events. The specificity (rate of predicted CTL given the actual CTL class) assesses the predictive

performances of CTL in CTL class. The kappa statistic assesses how the model outperforms a random

model that simply always predicts "CTL". According to Landis and Koch, a kappa of 0-0.20 is slight,

0.21-0.40 is fair, 0.41-0.60 is moderate, 0.61-0.80 is substantial, and 0.81-1 is almost perfect.[25] The

p-value assesses the statistical significance of the difference in accuracy between the random model

and the tested model.

Table 4: The confusion matrix and performances metrics used to assess the performances of the models

A confusion matrix was applied to evaluate the prediction performances of the ML models for this classification problem with a binary outcome (tail biting, "TB" or control "CTL" class). The definition of the performances metrics are presented.

Confusion matrix

		Actual	
		TB	CTL
Predicted TB		**TP** True positive predicted TB class that are actually TB class	**FP** False positive predicted TB class that are actually CTL class
CTL		**FN** False negative TB class not detected by the model (predicted as CTL class)	**TN** True negative predicted CTL class that are actually CTL class

Performances metrics

$$Sensitivity = \frac{TP}{TP + FN}$$

$$Positive\ Predicted\ Value\ (PPV) = \frac{TP}{TP + FP}$$

$$Specificity = \frac{TN}{TN + FP}$$

$$Accuracy = \frac{TN + TP}{TN + FP + TP + FN}$$

P value: statistical significance of the difference with a random model always predicting CTL class.

$\kappa = \dfrac{p_o - p_e}{1 - p_e}$ where ρ_o is the observed accuracy, and ρ_e is the expected accuracy of a random model always predicting CTL class.

Results

Model performances

Table 5 presents the model performances on the three training data sets. All models performed

significantly better than the random prediction model (that simply always predict "CTL"), with kappas

ranging from 0.30 to 1.00. Even if the criteria for optimization was the sensitivity, this performance

criterion was always lower or equal to the specificity, which is most likely due to the imbalance

between the numerous CTL and the rare TB.

Table 5: Models performances on <u>training data sets</u> for the Swiss, Swedish and Swedish+Swiss data sets. Seven machine learning models were trained on the training data set using a cross validation resampling approach. Performance metrics derived from the confusion matrix (table 4) are used to assess the models' performances. Models in bold are considered as the best predictive models.

	Accuracy	Sensitivity	Specificity	PPV	Kappa	p-value
Swedish						
glmnet[1]	0.88	0.63	0.94	0.76	0.61	<0.0001
rf[2]	**1.00**	**1.00**	**1.00**	**1.00**	**1.00**	**<0.0001**
svmRadial[3]	0.94	0.80	0.98	0.90	0.81	<0.0001
bayesglm[4]	0.88	0.63	0.94	0.76	0.61	<0.0001
nn[5]	**1.00**	**1.00**	**1.00**	**1.00**	**1.00**	**<0.0001**
kNN[6]	**0.99**	**0.99**	**0.99**	**0.98**	**0.98**	**<0.0001**
pls[7]	0.87	0.52	0.96	0.79	0.56	<0.0001
Swiss						
glmnet	0.92	0.40	0.99	0.79	0.50	<0.0001
rf	**1.00**	**1.00**	**1.00**	**1.00**	**1.00**	**<0.0001**
svmRadial	0.99	0.88	1.00	1.00	0.93	<0.0001
bayesglm	0.93	0.41	0.99	0.78	0.50	<0.0001
nn	**0.99**	**0.98**	**0.99**	**0.96**	**0.97**	**<0.0001**
kNN	0.95	0.54	1.00	0.98	0.67	<0.0001
pls	0.91	0.19	1.00	0.96	0.30	<0.0001
Swedish + Swiss						
glmnet	0.86	0.30	0.97	0.67	0.35	<0.0001
rf	**1.00**	**1.00**	**1.00**	**1.00**	**1.00**	**<0.0001**
svmRadial	1.00	0.99	1.00	1.00	1.00	<0.0001
bayesglm	0.86	0.30	0.97	0.67	0.35	<0.0001
nn	0.97	0.87	0.99	0.97	0.90	<0.0001
kNN	0.97	0.82	1.00	0.98	0.88	<0.0001
pls	0.85	0.13	0.99	0.75	0.18	<0.0001

[1]Generalized Linear Model with Stepwise Feature Selection; [2]Random forest; [3]Support Vector Machines with Radial Basis Function Kernel; [4]Bayesian Generalized Linear Model; [5]Neural network; [6]K-nearest neighbor; [7]Partial Least Squares Discriminant Analysis

 Model prediction performances

Tables 6 and 7 summarize the performances of the models to predict unseen analysis windows (CV)

or unseen pens (LOOP), respectively. For the Swedish+Swiss data set, the performances of the model

on CV and LOOP were assessed on the combined testing set (Swedish+Swiss) and on subsets of the

Swedish or Swiss data set separately. The random forest (RF) model showed the best predictive

performances on both the unseen analysis windows and the unseen pens, with an average accuracy

of 84% and a sensitivity of 38% on all data sets. Predictive performances were higher on unseen

analysis windows than on unseen pens, and always lower than the model performances on the

training set. In the Swedish data set, the predictive performances of the RF model on unseen pens

were poor (kappa<0). The models trained on the Swedish data were unable to predict Swiss data

(kappa<0.06), and vice versa (data not shown). Models trained on the Swedish+Swiss data set

showed a poorer performance (low kappa) in predicting new analysis windows of a Swedish or Swiss

subset data set than the same model trained on the Swedish or Swiss data sets individually. Lower

kappas were also obtained for prediction on new pens, except for glmnet, bayesglm and pls models,

that had improved predictive performances (kappa) on the Swiss subset.

Table 6: Performances of models to predict <u>unseen analysis windows</u> (Cross Validation (CV) approach) of the Swedish, Swiss and Swedish+Swiss testing data sets. Seven machine learning models were trained on the training data set and their prediction performances to predict unseen analysis windows (testing set) were evaluated. Performance metrics derived from the confusion matrix (table 4) are used to assess the models' performances. Models in bold are considered as the best predictive models.

		Swedish					
		Accuracy	Sensitivity	Specificity	PPV	Kappa	p value
glmnet[1]		0.80	0.45	0.90	0.57	0.38	0.04
rf[2]		**0.95**	**0.80**	**0.99**	**0.98**	**0.85**	**<0.0001**
svmRadial[3]		0.86	0.54	0.95	0.75	0.54	<0.0001
bayesglm[4]		0.81	0.45	0.90	0.57	0.39	0.04
nn[5]		0.87	0.70	0.91	0.70	0.62	<0.0001
kNN[6]		**0.97**	**0.94**	**0.98**	**0.94**	**0.92**	**<0.0001**
pls[7]		0.81	0.36	0.93	0.60	0.35	-
		Swiss					
		Accuracy	Sensitivity	Specificity	PPV	Kappa	p value
glmnet		0.85	0.22	0.94	0.30	0.17	-
rf		**0.94**	**0.50**	**0.99**	**0.95**	**0.63**	**<0.0001**
svmRadial		0.91	0.28	1.00	0.92	0.39	0.006
bayesglm		0.85	0.22	0.93	0.30	0.17	-
nn		**0.89**	**0.47**	**0.95**	**0.54**	**0.44**	**-**
kNN		**0.93**	**0.41**	**1.00**	**1.00**	**0.55**	**<0.0001**
pls		0.89	0.11	0.99	0.56	0.15	-
		Swedish + Swiss					
		Accuracy	Sensitivity	Specificity	PPV	Kappa	p value
Glmnet[a]		0.82	0.15	0.96	0.43	0.15	-
	Swedish[b]	0.77	0.21	0.93	0.46	0.18	-
	Swiss[c]	0.88	0.02	0.99	0.20	0.02	-
rf[a]		**0.94**	**0.70**	**0.99**	**0.96**	**0.78**	**<0.0001**
	Swedish[b]	**0.95**	**0.80**	**0.99**	**0.95**	**0.83**	**<0.0001**
	Swiss[c]	**0.94**	**0.48**	**1.00**	**1.00**	**0.62**	**<0.0001**
svmRadial[a]		0.88	0.38	0.98	0.80	0.46	<0.0001
	Swedish[b]	0.86	0.48	0.97	0.80	0.52	<0.0001
	Swiss[c]	0.90	0.16	1.00	0.76	0.23	-
bayesglm[a]		0.82	0.16	0.96	0.43	0.16	-
	Swedish[b]	0.77	0.21	0.93	0.46	0.19	-
	Swiss[c]	0.82	0.16	0.96	0.43	0.15	-
nn[a]		**0.88**	**0.55**	**0.94**	**0.67**	**0.53**	**<0.0001**
	Swedish[b]	**0.87**	**0.63**	**0.94**	**0.75**	**0.61**	**<0.0001**
	Swiss[c]	**0.88**	**0.37**	**0.95**	**0.47**	**0.35**	**-**
kNN[a]		**0.96**	**0.78**	**1.00**	**0.98**	**0.84**	**<0.0001**
	Swedish[b]	**0.96**	**0.87**	**0.98**	**0.94**	**0.88**	**<0.0001**
	Swiss[c]	**0.92**	**0.32**	**1.00**	**1.00**	**0.46**	**0.0006**
pls[a]		0.83	0.06	0.99	0.62	0.09	-
	Swedish[b]	0.79	0.09	0.98	0.62	0.11	-
	Swiss[c]	0.89	0.00	1.00	0.00	0.00	-

[1]Generalized Linear Model with Stepwise Feature Selection; [2]Random forest; [3]Support Vector Machines with Radial Basis Function Kernel; [4]Bayesian Generalized Linear Model; [5]Neural network; [6]K-nearest neighbor; [7]Partial Least Squares Discriminant Analysis
[a]Prediction performances of the models on testing data set containing Swedish and Swiss data.
[b]Prediction performances of the models on the Swedish subset of the testing data set.
[c]Prediction performances of the models on the Swiss subset of the testing data set.

Table 7: Performances of models to predict <u>unseen pens</u> (Leave one out cross validation (LOOP) approach) of the Swedish, Swiss and Swedish+Swiss data sets.

Seven machine learning models were trained on the training data set and their prediction performances to predict unseen pens (testing set) were evaluated. Performance metrics derived from the confusion matrix (table 4) are used to assess the models' performances. Bold models are considered as the most promising models, according to the authors. Models in bold are considered as the best predictive models.

Swedish	Accuracy	Sensitivity	Specificity	PPV	Kappa	p value
glmnet[1]	0.56	0.14	0.71	0.14	<0.00	-
rf[2]	**0.63**	**0.11**	**0.81**	**0.17**	**<0.00**	**-**
svmRadial[3]	0.54	0.20	0.66	0.17	<0.00	-
bayesglm[4]	0.56	0.14	0.71	0.14	<0.00	-
nn[5]	0.56	0.05	0.73	0.06	<0.00	-
kNN[6]	0.62	0.00	0.83	0.00	<0.00	-
pls[7]	0.58	0.10	0.75	0.12	<0.00	-

Swiss	Accuracy	Sensitivity	Specificity	PPV	Kappa	p value
glmnet	0.82	0.18	0.96	0.50	0.19	-
rf	**0.84**	**0.18**	**0.99**	**0.75**	**0.24**	**-**
svmRadial	0.81	0.02	0.99	0.25	0.01	-
bayesglm	0.82	0.18	0.95	0.47	0.18	-
nn	0.75	0.04	0.90	0.09	<0.00	-
kNN	**0.84**	**0.12**	**0.99**	**0.86**	**0.18**	**-**
pls	0.82	0.08	0.99	0.57	0.10	-

Swedish + Swiss		Accuracy	Sensitivity	Specificity	PPV	Kappa	p value
Glmnet[a]		0.73	0.12	0.90	0.27	0.03	-
	Swedish[b]	0.62	0.00	0.84	0.00	<0.00	-
	Swiss[c]	0.87	0.37	0.99	0.86	0.45	0.008
rf[a]		0.74	0.00	0.96	0.00	<0.00	-
	Swedish[b]	0.69	0.00	0.93	0.00	<0.00	-
	Swiss[c]	0.82	0.00	1.00	0	0	-
svmRadial[a]		0.71	0.03	0.91	0.08	<0.00	-
	Swedish[b]	0.63	0.03	0.84	0.06	<0.00	-
	Swiss[c]	0.82	0.02	1.00	1.00	0.03	-
bayesglm[a]		0.73	0.13	0.91	0.28	0.04	-
	Swedish[b]	0.63	0.01	0.84	0.02	<0.00	-
	Swiss[c]	0.87	0.37	0.99	0.86	0.46	0.009
nn[a]		0.63	0.14	0.78	0.16	<0.00	-
	Swedish[b]	0.56	0.21	0.68	0.18	<0.00	-
	Swiss[c]	0.74	0.00	0.91	0.00	<0.00	-
kNN[a]		0.74	0.00	0.95	0.00	<0.00	-
	Swedish[b]	0.69	0.00	0.93	0.00	<0.00	-
	Swiss[c]	0.82	0.00	1.00	0.00	0.00	-
pls[a]		0.77	0.08	0.97	0.41	0.07	-
	Swedish[b]	0.70	0.00	0.94	0.00	<0.00	-
	Swiss[c]	0.86	0.24	0.99	0.92	0.34	0.04

[1]Generalized Linear Model with Stepwise Feature Selection; [2]Random forest; [3]Support Vector Machines with Radial Basis Function Kernel; [4]Bayesian Generalized Linear Model; [5]Neural network; [6]K-nearest neighbor; [7]Partial Least Squares Discriminant Analysis
[a]Prediction performances of the models on testing data set containing Swedish and Swiss data.
[b]Prediction performances of the models on the Swedish subset of the testing data set.
[c]Prediction performances of the models on the Swiss subset of the testing data set.

Impact of TB and analysis windows

We have tested the influence of the size of the analysis window and the size of the TB window on

predictive performances for the Swedish data set. Accuracy of models with analysis windows of 7 or

21, or with different TB windows (-49 to +10 days, or -10 to +5 days) were compared with the default

windows size (analysis window: 14 days; TB window: -35 to +10 days). The RF model was chosen for

this analysis, as it was the model with the best predictive performances over the three data sets and

for the two (CV and LOOP) approaches.

The predictive performances of the RF model increases when the analysis window contains 21 days

(instead of 14 days) or when the TB window was larger (-49 to +10 days) (Table 8). Interestingly, the

RF model never predicted TB class in CTL pens, thus ML models were able to discriminate CTL pen

behaviour from TB pen behaviour. In addition, the prediction of a TB event for a TB pen was almost

exclusively within the TB window (high specificity). The detection of an upcoming TB event was not

better near the TB date (day 0) than at the beginning of the TB window (day -35, for default TB

window size).

Table 8: Prediction performances of the random forest (rf) model depending on the TB and analysis
window size on the Swedish data set.
The prediction performances of the rf model were assessed while varying the size of the TB window
and of the analysis window. The prediction were made for unseen analysis windows or for unseen
pens. The rf model was chosen as it seems to the authors that it is the most promising model for this
ML prediction, based on previous results (Tables 6 and 7).

	Analysis window size	TB window size	Accuracy	Sensitivity	Specificity	PPV	Kappa	p value
					Swedish			
Unseen analysis windows (CV)	7 days	[-35;10]	0.94	0.76	0.99	0.96	0.81	<0.0001
	14 days	**[-35;10]**	**0.94**	**0.70**	**0.99**	**0.96**	**0.78**	**<0.0001**
		[-49;10]	**0.97**	**0.88**	**0.99**	**0.97**	**0.90**	**<0.0001**
		[-10;5]	0.95	0.45	1.00	0.94	0.58	<0.0001
	21 days	**[-35;10]**	**0.97**	**0.87**	**1.00**	**0.99**	**0.91**	**<0.0001**
Unseen pens (LOOP)	7 days	[-35;10]	0.64	0.00	0.87	0.00	<0.00	-
	14 days	**[-35;10]**	**0.63**	**0.11**	**0.81**	**0.17**	**<0.00**	**-**
		[-49;10]	**0.61**	**0.19**	**0.82**	**0.35**	**0.02**	**-**
		[-10;5]	0.91	0.00	1.00	0.00	0.00	-
	21 days	[-35;10]	0.66	0.00	0.88	0.00	<0.00	-

Discussion

In this study, TB events could be detected up to 35 days in advance using an ML model that analysed feeding behaviours recorded by electronic feeders. Early indicators of TB events were more easily identified when the model had access to previous records of the pen. Thus, one should consider performing continuous analysis of the data of each pen, even in the absence of TB events. To our knowledge this is the first time that an ML algorithm has been used to predict TB in pigs.

In the following discussion we compare our results using an ML model with statistical models from other articles attempting to characterize feeding behaviours before and after a TB event.

Munsterhjelm et al. previously investigated feeding behaviours in relation to TB, 70 days before to 28 days after a first wound was detected in the pen. Feeding behaviours in TB pens were compared to matched CTL pens. Pigs in TB pens tend to visit less frequently the feeder than pigs in CTL pens. The frequency of daily feeder visits (DFV) reduced with the approach of the TB date. The feeding behaviour in TB pens significantly changed 14 days before the detection of the TB event. Time spent at the feeder and daily feed consumption (DFC) were also reduced in TB pens compared to CTL pens. This tendency persisted after the TB date, and pigs continued to spend less time in the feeder and visit less frequently, even if the difference from the control group decreased with time. Pigs also tended to eat faster (more intake per second). They concluded that the rapid change in feeding behaviour suggests that TB behaviour escalates 14 days before the TB date.[18]

The change in feeding behaviour in TB pen was also observed by Tessier et al. during a TB outbreak in a pen. They studied the evolution of DFV, the DFC and the feeding time – 7 days before (pre-injury phase), 7 days during (acute phase) and 7 days after (recovery phase) the TB outbreak. The DFV decreased before the TB date, reached a minimum during the outbreak, and increased during the recovery phase. Consumption time (for a given amount of feed eaten) decreased while the TB outbreak was evolving (during the pre-injury and acute phases) and remained low during the recovery phase. This study corroborates that a change in feeding behaviour in a pen can indicate a

future TB event. This also confirms the gradual change of the feeding behaviour over time, reaching

its maximum at the TB date.[19]

The statistical analysis by Wallenbeck and Keeling on the Swedish data set also demonstrated a

significant decrease in DFV in TB pens compared to matched CTL pens, 42 to 63 days before the TB.

In addition, the DFC was always reduced in TB pens compared to CTL pens.[17] This difference in

feeding behaviour between CTL and TB pens must have been also detected in the present analysis by

the ML models that never predict TB class in CTL pens.

In contrast, Zonderland et al. did not find any change in feeding behaviour, especially in the

difference in time spent feeding between CTL and TB pens.[26] But they studied pigs fed *ad libitum*

from dry feeders with two feeding spaces, where pigs could feed at the same time from the same

feeder. The feeding behaviour in pens with feeder troughs is different from those with automatic

feeders.[16] The upcoming TB event may be responsible for this difference.

The comparison between TB and CTL windows in the present study (Table 2) doesn't reflect a general

tendency of the change in feeding behaviour. Indeed, the characteristics of the pigs in CTL and TB

windows are not balanced, so the differences in age and weight between the two categories could

explain the differences in DFV, DFC and StdFC and also the inconsistencies observed between the

Swiss and Swedish data sets.

Wallenbeck and Keeling described a change in DFV 42 to 63 days before a TB event, comparing TB

pens to CTL pens.[17] In the present study, the detection was limited to 35 days before the event (TB

window: -35 to +10 days). Nevertheless, increasing the size of the TB window to (-49 to +10 days)

improves the accuracy and the sensitivity of the RF model. In addition, the two analyses did not use

the same reference: the Wallenbeck analysis compared TB pens to matched CTL pens, trying to

characterize a feeding behaviour that would be typical of a future TB event.[17] For the present

analysis, each pen was also its own control, as a TB pen could be a TB class when inside the TB

window but a CTL when outside. The model could not only detect feeding behaviour that is typical of

a future TB event, but also detect changes in feeding behaviour over time that are indicative of TB

events.

Indeed, as demonstrated by Munsterhjelm et al. and Tessier et al., a drastic change in feeding

behaviour in each pen is also indicative of an upcoming TB event.[18-19] In addition, the present

analysis considers a combination of three feeding behaviour parameters (DFV, DFC, StdFC), thus

increasing the possibility to detect TB events. In a comprehensive book chapter on TB, Valros

describes the gradual change in feeding behaviour until the TB date. DFV can start decreasing

months to weeks before, whereas DFC seems to be affected only six days before the TB date.[5]

Thus, the combination of these two indicators should improve the specificity of detection.

With modified feeding behaviours being present one to several months before the event, the TB

events observed in the present study seemed to result from a long process and therefore to belong

to the two-stage type, as described by Taylor et al.[17] This type is described as starting as a harmless

behaviour that escalates to actual TB outbreak. A suggested cause was chronic but moderate stress.

In general, TB appears to be connected to an increased stress level in pigs.[18, 19]

In the different studies, the first changes in feeding behaviour linked to TB occurred 42 to 63

days[10], or 70 days[20] before the first TB event. Depending on the monitoring frequency, the TB

event may be detected sooner or later. Therefore the TB date may be determined when the TB

process has just begun or when it has become an outbreak. This would also explain the discrepancies

between how long before the TB event the feeding behaviour is described to have changed.

In addition, it seems that the feeding behaviour is not only specific to a pen, but each site has its own

characteristics (e.g., breeds, climates and feeding compositions). The model trained on one data set

(Swedish or Swiss) was unable to detect events in the other data set, and the combination of the

data sets (Swedish + Swiss) did not greatly improve model performance. Thus, the increase in size

resulting from the combination of the Swedish and Swiss data did not provide much advantage. One

explanation could be linked to site-specific risks factors. As acknowledged by the European Food

Safety Authority 2014, one of the difficulties in preventing TB resides in the fact that every farm is

different and has its own risk factors. Prevention strategies need therefore to be designed at a farm-

specific level.[3] Depending on the chronic risk factors of the farm (e.g., breed, sex, feeding, access to

manipulable material, space allowance and group size), the feeding behaviour associated with a TB

event may vary. This observation was already acknowledge by Taylor et al. and Valros, when they

defined the four types of TB (two-stage, sudden-forceful, obsessive and epidemic), associated with

four putative causations. The two-stage type was brought about by chronic and moderate stress. The

sudden-forceful type was suggested to be caused by competition for resources. The epidemic type

stemmed from a sudden change in the pig's life (e.g., feeding disturbance or temperature).[5,28]

In conclusion, an ML model can be implemented in an automatic feeder system on a farm to detect

early signs of TB behaviours at least 35 days before the actual TB event. Thanks to these early

warnings, farmers could implement measures to prevent the occurrence of the TB event – for

example, by adding more straw as occupational material. Farmers could also start a TB risk analysis

to identify the reasons why pigs are disturbed. Continuous implementation of the model on farms

would also lead to improved prevention of TB events, serving the welfare of the pigs and bringing an

economic benefit to the farmers. Finally, this ML approach could also be a useful tool by allowing the

systematic study of the effectiveness of different intervention strategies under controlled conditions.

References

1.   Larsen MLV, Andersen HML, Pedersen LJ. Can tail damage outbreaks in the pig be predicted

by behavioural change? Vet J. 2016;209:50-6. doi: 10.1016/j.tvjl.2015.12.001.

2.   Brunberg EI, Rodenburg TB, Rydhmer L, Kjaer JB, Jensen P, Keeling LJ. Omnivores going

astray: A review and new synthesis of abnormal behavior in pigs and laying hens. Front Vet Sci.

2016;3(57). doi: 10.3389/fvets.2016.00057.

3.   European Food Safety Authority. Scientific opinion concerning a multifactorial approach on

the use of animal and non-animal-based measures to assess the welfare of

pigs. EFSA J. 2014:12(5):3702. doi: 10.2903/j.efsa.2014.3702.

4.   European Food Safety Authority. Pig welfare risks associated with tail biting in pigs and

possible means to reduce the need for tail docking considering the different housing and husbandry

systems - scientific opinion of the panel on animal health and welfare. EFSA J. 2007;611:1-13. doi:

10.2903/j.efsa.2007.611.

5.   Valros A. Chapter 5 - Tail biting. In: Špinka M, editor. Advances in Pig Welfare. Woodhead

Publishing; 2018. p. 137-66.

6.   D'Eath RB, Arnott G, Turner SP, Jensen T, Lahrmann HP, Busch ME, et al. Injurious tail biting

in pigs: how can it be controlled in existing systems without tail docking? Animal. 2014;8(9):1479-97.

Epub 2014/08/19. doi: 10.1017/s1751731114001359. PubMed PMID: 25130712.

7.   Wedin MB, Baxter EM, Jack M, Futro A, D'Eath RB. Early indicators of tail biting outbreaks in

pigs. Appl Anim Behav Sci. 2018;208:7-13. doi: 10.1016/j.applanim.2018.08.008.

8.   D'Eath RB, Jack M, Futro A, Talbot D, Zhu Q, Barclay D, et al. Automatic early warning of tail

biting in pigs: 3D cameras can detect lowered tail posture before an outbreak. PLoS One.

2018;13(4):e0194524. doi: 10.1371/journal.pone.0194524.

9.   Lahrmann HP, Hansen CF, D'Eath R, Busch ME, Forkman B. Tail posture predicts tail biting

outbreaks at pen level in weaner pigs. Appl Anim Behav Sci. 2018;200:29-35. doi:

10.1016/j.applanim.2017.12.006.

10.   Statham P, Green L, Bichard M, Mendl M. Predicting tail-biting from behaviour of pigs prior

to outbreaks. Appl Anim Behav Sci. 2009;121:157-64. doi: 10.1016/j.applanim.2009.09.011.

11.   Zonderland JJ, van Riel JW, Bracke MBM, Kemp B, den Hartog LA, Spoolder HAM. Tail posture

predicts tail damage among weaned piglets. Appl Anim Behav Sci. 2009;121(3):165-70. doi:

10.1016/j.applanim.2009.09.002.

12.   Wallgren T, Larsen A, Gunnarsson S. Tail posture as an indicator of tail biting in undocked

finishing pigs. Animals (Basel). 2019;9(1):18. doi: 10.3390/ani9010018. PubMed PMID: 30626058.

13.   Reimert I, Bolhuis JE, Kemp B, Rodenburg TB. Indicators of positive and negative emotions

and emotional contagion in pigs. Physiol Behav. 2013;109:42-50. Epub 2012/11/20. doi:

10.1016/j.physbeh.2012.11.002. PubMed PMID: 23159725.

14.   Ursinus WW, Van Reenen CG, Kemp B, Bolhuis JE. Tail biting behaviour and tail damage in

pigs and the relationship with general behaviour: predicting the inevitable? Appl Anim Behav Sci.

2014;156:22-36. doi: 10.1016/j.applanim.2014.04.001.

15.   Young R, Lawrence A. Feeding behaviour of pigs in groups monitored by a computerized

feeding system. Anim Prod. 1994;58:145-52. doi: 10.1017/S0003356100007182.

16.   Brunberg E, Wallenbeck A, Keeling L. Tail biting in fattening pigs: associations between

frequency of tail biting and other abnormal behaviours. Appl Anim Behav Sci. 2011;133. doi:

10.1016/j.applanim.2011.04.019.

17.   Wallenbeck A, Keeling LJ. Using data from electronic feeders on visit frequency and feed

consumption to indicate tail biting outbreaks in commercial pig production. J Anim Sci.

2013;91(6):2879-84. Epub 2013/03/13. doi: 10.2527/jas.2012-5848. PubMed PMID: 23478818.

18.   Munsterhjelm C, Nordgreen J, Heinonen M, Janczak A, Valros A, editors. Feeding behaviour

and performance in relation to injurious tail biting in boars – a longitudinal study. 7th Eurpoean

Symposium of Porcine Health Management (ESPHM), Dublin, Ireland; Jun 2016.

19.   Tessier F, Maikoff G, Bee G, Ollagnier C, editors. Tail biting in switzerland: a restrospective

study. 51eme Journées de la Recherche Procine (JRP), Paris; Feb 2019.

20.   Andersson C, Lindberg JE. Forages in diets for growing pigs 1. Nutrient apparent digestibilities

and partition of nutrient digestion in barley-based diets including lucerne and white-clover meal.

Animal Science. 1997;65(3):483-91. Epub 2010/09/02. doi: 10.1017/S1357729800008687.

21.   Agroscope. Apports alimentaires recommandés pour les porcs (livre jaune) 2005 [cited 2020].

22.   Kuhn M. Building predictive models in R using the caret package. J Stat Softw. 2008;1(5). doi:

10.18637/jss.v028.i05.

23.   Molinaro AM, Simon R, Pfeiffer RM. Prediction error estimation: a comparison of resampling

methods. Bioinformatics. 2005;21(15):3301-7. Epub 2005/05/21. doi: 10.1093/bioinformatics/bti499.

PubMed PMID: 15905277.

24.   Tharwat A. Classification assessment methods. Applied Computing and Informatics.

2020;17(1):168-192. doi: 10.1016/j.aci.2018.08.003.

25.   Landis JR, Koch GG. An application of hierarchical kappa-type statistics in the assessment of

majority agreement among multiple observers. Biometrics. 1977;33(2):363-74. doi:

10.2307/2529786.

26.   Zonderland J, Schepers F, Bracke MBM, Hartog LA, Kemp B, Spoolder HAM. Characteristics of

biter and victim piglets apparent before a tail-biting outbreak. Animal. 2011;5:767-75. doi:

10.1017/S1751731110002326.

27.   Nielsen BL, Lawrence AB, Whittemore CT. Feeding behaviour of growing pigs using single or

multi-space feeders. Appl Anim Behav Sci. 1996;47(3):235-46. doi: 10.1016/0168-1591(95)00649-4.

28.   Taylor NR, Main DC, Mendl M, Edwards SA. Tail-biting: a new perspective. Vet J.

2010;186(2):137-47. Epub 2009/10/07. doi: 10.1016/j.tvjl.2009.08.028. PubMed PMID: 19804997.

29.   Munsterhjelm C, Brunberg E, Heinonen M, Keeling L, Valros A. Stress measures in tail biters

and bitten pigs in a matched case-control study. Anim Welf. 2013;22(3):331-338. doi:

10.7120/09627286.22.3.331.

30.   Zupan M, Janczak AM, Framstad T, Zanella AJ. The effect of biting tails and having tails bitten

in pigs. Physiol Behav. 2012;106(5):638-44. doi: 10.1016/j.physbeh.2012.04.025.

Figure 1: Analysis of the time dependency of the data thanks to the analysis window approach.
Analysis window slides along the time line (Blue arrow). The tail biting (TB) pen will be classified as TB class, when the A_date (first day of the analysis window) enters the TB window (orange block) and as "CTL" class when it is outside the TB window. Control pen will always be classified as "CTL".

Figure 1