The recent evolutionary rescue of a staple crop depended on over half a century of global germplasm exchange

Kebede T. Muleta[1*], Terry Felderhoff[1*], Noah Winans[1], Rachel Walstead[2], Jean Rigaud Charles[3], J. Scott Armstrong[4], Sujan Mamidi[2], Chris Plott[2], John P. Vogel[5], Peggy G. Lemaux[6], Todd C. Mockler[7], Jane Grimwood[2], Jeremy Schmutz[2,5], Gael Pressoir[3], Geoffrey P. Morris[1,8]

[1] Department of Agronomy, Kansas State University, Manhattan, Kansas, USA 66502
[2] Genome Sequencing Center, HudsonAlpha Institute for Biotechnology, Huntsville, AL 35806
[3] Chibas and Faculty of Agriculture and Environmental Sciences, Quisqueya University, Port-au-Prince, Haiti
[4] U.S. Department of Agriculture, Agricultural Research Service, Wheat, Peanut and Other Field Crops Research Unit, 1301 North Western Rd., Stillwater, OK 74075
[5] Department of Energy Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, CA 94720
[6] Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720
[7] Donald Danforth Plant Science Center, St. Louis, MO 63132
[8] Department of Soil & Crop Science, Colorado State University, Fort Collins, Colorado, 80526
[*] Equal contributions

Corresponding author:
Geoffrey P. Morris
Geoff.Morris@colostate.edu

ABSTRACT

Rapid environmental change can lead to extinction of populations or evolutionary rescue via genetic adaptation. In the past several years, smallholder and commercial cultivation of sorghum (*Sorghum bicolor*), a global cereal and forage crop, has been threatened by a global outbreak of an aggressive new biotype of sugarcane aphid (SCA; *Melanaphis sacchari*). Here we characterized genomic signatures of adaptation in a Haitian sorghum breeding population, which had been recently founded from admixed global germplasm, extensively intercrossed, and subjected to intense selection under SCA infestation. We conducted evolutionary population genomics analyses of 296 post-selection Haitian lines compared to 767 global accessions at 159,683 single nucleotide polymorphisms. Despite intense selection, the Haitian population retains high nucleotide diversity through much of the genome due to diverse founders and an intercrossing strategy. A genome-wide fixation (F_{ST}) scan and geographic analyses suggests that adaptation to SCA in Haiti is conferred by a globally-rare East African allele of *RMES1*, which has also spread to other breeding programs in Africa, Asia, and the Americas. *De novo* genome sequencing data for SCA resistant and susceptible lines revealed putative causative variants at *RMES1*. Convenient low-cost markers were developed from the *RMES1* selective sweep and successfully predicted resistance in independent U.S. × African breeding lines and eight U.S. commercial and public breeding programs, demonstrating the global relevance of the findings. Together, the findings highlight the potential of evolutionary genomics to develop adaptive trait breeding technology and the value of global germplasm exchange to facilitate evolutionary rescue.

INTRODUCTION

Ongoing processes of global change, encompassing climate change, nutrient cycles, and pest outbreaks, are shaping the evolution of natural and agricultural ecosystems (*1*, *2*). Intense selection pressure following environment changes may lead to the rapid decline or extinction of populations (*3*, *4*). If a population is to persist under such strong selection, adaptive standing genetic variation must exist or adaptive *de novo* variation must arise on a sufficiently fast timescale (*5*). This population genetic phenomenon, evolutionary rescue, has become a focus of considerable empirical and theoretical study in ecology and conservation biology, since the current rate of global change could exceed the capacities of many populations to adapt (*6*, *7*). Still, there is a lack of examples of evolutionary rescue occuring in the field and at large geographic scales (*4*). In agricultural systems, the spread of pests or emergence of new aggressive biotypes may lead to a reduction of crop diversity or a total loss of crop cultivation (*8*). Therefore, understanding and facilitating evolutionary rescue in agricultural systems is critical for global food security.

Populations of crops or wild species subjected to strong selection pressure may experience a major population bottleneck, resulting in a loss of genetic diversity (*9*). The level of diversity preserved in a population recovering from strong selection depends on the number of backgrounds on which the adaptive alleles emerge (*10*), which can determine the potential for future adaptation or genetic gain. Conversely, adaptation conferred by a beneficial variant derived from a single progenitor causes the removal of genetic diversity from the surviving population (*10*, *11*). Evolutionary population genomics approaches using genome-wide polymorphism data from diverse germplasm can identify candidate loci for adaptive traits (*12*). While genome scans for selection have been widely used to identify putative adaptive alleles in crops (*9*, *13*), they have not yet been used to identify trait-predictive markers for molecular breeding of stress-resilient varieties (*14*).

Sorghum (*Sorghum bicolor* L. [Moench]) is among the world's most important staple crops for smallholder farmers in semiarid regions, as well as a commercial grain and forage crop in industrialized nations (*15*). Since 2013 an aggressive biotype of the sugarcane aphid (SCA; *Melanaphis sacchari*) has become a major threat to global sorghum production, with widespread and substantial yield loss (*16*, *17*). The *M. sacchari* superclone has been rapidly expanding (*18*), putting >90% of the sorghum-producing areas of North America at risk and threatening to end sorghum cultivation in some areas (*16*). In Haiti, a Caribbean nation with one of the world's highest rates of food insecurity, sorghum is among the most important staple crops (*19*). However, heavy infestations by *M. sacchari* since 2015 have caused the loss of over 70% of sorghum production in the country and prevented production of most local landraces (*20*). Shortly before the SCA outbreak, a new Haitian breeding population had been launched by Chibas using global admixed germplasm, rapid-cycling intercrossing, and selection under smallholder conditions (i.e. no insecticidal treatment) (*21*). Selecting from a small number of breeding lines that survived SCA infestation, a new SCA resistant sorghum variety, Papèpichon, was developed and distributed nationally (*19*), and intercrossing and advancement of resistant

breeding lines has continued.

Here we used a retrospective genomic analysis of the Haitian sorghum breeding
population that was subjected to strong selection under SCA infestation, to understand the
genetic basis of the evolutionary rescue following the SCA outbreak, as well as the origins of the
SCA resistance alleles. We find that the rapid adaptation of the Haitian breeding population to
the SCA outbreak was due to selection for a globally-rare Ethiopian allele at the *RMES1* SCA
resistance locus, which is shared across programs in Africa, Asia, and the Americas because of
>50 years of global germplasm exchange prior to the SCA outbreak. Further, we developed a
convenient low-cost molecular marker based on the evolutionary genome scan and validated it in
eight commercial and public sorghum breeding programs, demonstrating the value of leveraging
global germplasm exchange and evolutionary population genomics to improve crop resilience.

RESULTS

Genome-wide polymorphism and nucleotide diversity
To understand the evolutionary rescue of sorghum following the SCA outbreak, we conducted a
retrospective genomic analysis of the Haitian breeding population (HBP) in comparison to a
global diversity panel (GDP). Genotyping-by-sequencing of 296 HBP and 767 GDP (Supp. Fig.
S1; Supp. File S1) sorghum lines generated 159,683 polymorphic SNPs with an average SNP
density of 75 and 229 per Mb in the HBP and GDP, respectively (Supp. Fig. S2). The GDP had a
higher proportion of low-frequency minor alleles (<5% MAF) compared to the HBP (Supp. Fig.
S3). Average inbreeding coefficients (F_{IS}) in HBP and the GDP was estimated at 0.7 and 0.9,
respectively (Supp. Table S1). The effect of selection on genetic diversity in HBP was assessed
based on genome-wide nucleotide diversity (π) in the HBP in comparison to (i) the GDP and (ii)
a major public program in the US (Texas A&M pre-breeding lines, TAM-PBL, $N = 35$). Average
nucleotide diversity in the HBP was estimated at 2.3×10^{-5}. In the GDP and TAM-PBL, estimates
of average π were 5.8×10^{-5} and 4.8×10^{-5}, respectively (Fig. 1A-C, Supp. Table S2). In the HBP,
31% of 1 Mb windows have negative average Tajima's D values, while in the GDP
predominantly positive values of Tajima's D were observed (Supp. Fig. S4).

Contributions of global sorghum diversity to the Haitian breeding population
The genetic ancestry of the HBP from global germplasm was inferred based on population
structure analyses. In a neighbor joining analysis, the HBP clusters with caudatum accessions
(Fig. 2A), specifically caudatums from East Africa. Similarly, in principal coordinate analysis,
the HBP cluster swith East African caudatum accessions (Fig. 2C). To estimate ancestry
coefficients for HBP lines, we used Bayesian model-based clustering in ADMIXTURE,
projecting HBP lines onto ancestral populations and allele frequencies defined using only GDP
(with HBP lines omitted). With the GDP, the lowest cross-validation error was observed at $K = 8$
(Supp. Fig. S5) and accessions clustered by ecogeographic region and botanical type, as
expected. ADMIXTURE projection analysis suggests that the HBP is admixed, largely
consisting of caudatum haplotypes (>80% of the genome) with a remaining small percentage
being contributed by durra and guinea sorghums (Fig. 2D).

Evidence of a selective sweep in the Haitian breeding population at *RMES1*

To identify genome regions implicated in the evolutionary rescue of the HBP, genome-wide
scans for outlier loci were performed based on an F_{ST} test. Overall, the HBP is moderately
differentiated from the global diversity panel, with an average genome-wide F_{ST} of 0.16 (Fig.
3A, Supp. File S2). Based on a Bonferroni-adjusted *P*-value < 0.01, F_{ST} analysis identified 171
outlier genomic regions, which are candidate selective sweep regions. Several genomic regions
with F_{ST} outlier regions co-localized with candidate genes for traits under selection by the Chibas
breeding program, including photoperiodic flowering, inflorescence architecture, stay☐green,
stem sugar content, and SCA resistance (Fig 1D; Supp. File S3). Interestingly, the most extreme
F_{ST} outliers were observed on chromosome 6, precisely colocalizing with *RMES1*, a locus
previously shown to underlie SCA resistance in a Chinese sorghum line of unknown pedigree
(*22*) (Fig. 3A-B). To characterize the prevalence of the putative selected haplotype and identify
its geographic origin, we mapped the allelic distribution of the highest F_{ST} SNP S6_2995581 in
global georeferenced sorghum landraces (Fig. 3C) and compared these distributions to the allele
frequency in US and Haitian breeding germplasm (Fig. 3C, inset left). Globally, the allele is rare
(<2%), found only in Ethiopian caudatum landraces and a few breeding lines from West Africa
and the US. However, the sweep-associated allele is common (~40%) in Ethiopian caudatum
accessions (Fig. 3C; Supp. Table S3). The high local frequency of the sweep-associated allele in
Ethiopia suggests a likely origin of the SCA resistance allele in the Ethiopian highlands (Fig. 3C,
inset right).

Comparative genomic analysis to identify candidate causative variants

To identify candidate causative variants for the *RMES1* locus, we used whole-genome
resequencing and *de novo* genome sequencing of sorghum accessions with known SCA
reactions. The *RMES1* interval previously defined based on biparental linkage mapping (*22*)
includes seven gene models (Sobic.006G017000, Sobic.006G017100, Sobic.006G017200,
Sobic.006G017332, Sobic.006G017266, Sobic.006G017400, and Sobic.006G017500) that were
candidates for the causative gene. Comparative genomic analyses based on local multiple
sequence alignment (MSA) of *de novo* genome sequence of the resistant accession (PI 276837,
the Ethiopian progenitor of SCA resistant line SC170) and three sorghum reference genomes of
SCA susceptible lines (BTx623, Tx430, and BTx642) were used to identify potential causative
variants. No sequence variants were identified in the exons of three of the seven genes
(Sobic.006G017000, Sobic.006G017100, and Sobic.006G017266). A total of 35, 32, and 29
nonsynonymous SNPs were detected in the exons of Sobic.006G017200, Sobic.006G017400,
and Sobic.006G017500, when comparing the sequences of the resistant PI 276837 and the three
susceptible accessions. In addition, three insertion-deletion variations resulting in frame-shift
were detected in Sobic.006G017500. (Supp. File S4). To further refine the set of candidate
causative variants, we performed a localized association analysis for SCA resistance ("resistant"
or "susceptible", based on literature classification) around *RMES1* with resequencing data for
diverse sorghum accessions (Fig. 4, Supp. File S5) that detected 101 highly significant
associations (*P*-value > 0.0001). Annotations of the variants within the *RMES1* locus indicate

that only ten of 101 associated variants are nonsynonymous (5 of 10 in Sobic.006G017200 and
the remaining 5 of 10 in Sobic.006G017500.

Development and validation of a molecular marker based on the selective sweep
Next, we sought to test the hypothesis that the genome region identified by the F_{ST} scan
underlies variation for SCA resistance in other global sorghum germplasm. Therefore we
developed a kompetitive allele specific PCR (KASP) marker based on the SNPs at the *RMES1*
locus identified in the F_{ST} scan. Of the candidate SNPs (Supp. File S2, Supp. Table S3, Supp.
File S6), SNP 06_02892438 was determined to have the best combination of linkage, LD, and
technical KASP functionality of the SNPs. Alternative SNPs were also developed into markers
(Supp. File S6), and while the markers are often used as technical checks, testing has confirmed
the priority of the marker based on SNP 06_02892438 (Sbv3.1_06_02892438R). Initial
validation of the Sbv3.1_06_02892438R KASP marker using DNA samples from known
resistant lines (SC110, Tx2783, and IRAT204), susceptible lines (BTx623 and BTx642) (*23*),
and multiple F_2 families segregating for SCA resistance demonstrated that the KASP marker
Sbv3.1_06_02892438R was in complete agreement with historical phenotypes of inbred lines
and segregated within F_2 populations (Supp. File S7).
An F_4 population derived from a cross between IRAT204 (resistant African variety) and
Tx430 (susceptible US breeding line) was used to further validate the broader utility and
predictiveness of the KASP marker for marker-assisted selection (Fig. 5A-B). A total of 50 F_4
lines together with resistant (IRAT204 and SC110) and susceptible controls (RTx430) were
genotyped with the KASP marker Sbv3.1_06_02892438R. Both resistant controls and 23 F_4 lines
were homozygous for the resistant allele. The susceptible control and 9 F_4 lines were
homozygous for the susceptible allele, and the remaining 18 F_4 lines were heterozygous at the
SNP. Twenty-three selected F_4 lines with three resistant and three susceptible control lines were
tested for SCA reaction in a free-choice flat screen assay in the greenhouse, scoring aphid
damage rating, leaf greenness (SPAD), and seedling height. The SCA reaction phenotypes match
the KASP marker genotypes, demonstrating the reliability and predictability of using KASP
markers in marker-assisted selection for SCA resistance breeding (Fig. 5A-B; Supp. File S7).

Multi-program validation and deployment in commercial and public breeding programs
To further validate the utility of the SCA resistance KASP markers, we tested them with five US
commercial seed company breeding programs and three US public sector breeding programs,
representing a large fraction of the US sorghum breeding community (Fig. 5C). (The programs
are anonymized to avoid disclosing proprietary information.) Under the hypothesis that (i)
RMES1 underlies SCA resistance in US breeding programs and (ii) the KASP marker
(Sbv3.1_06_02892438R) tags the relevant resistant vs. susceptible haplotypes, the breeders'
phenotype-based classification of SCA resistance should largely match the KASP marker
genotype-based prediction. As expected, the match between the phenotype-based breeder
classification and KASP marker genotypes is high, ranging from ~60-100%, with most
germplasm sets (9/12) have >80% matching (Fig. 5C; Fig. S6). Less than 0.5% of mismatches
(5/1100) were observed among technical replicates (independent tissue samples from the same

plant), so mismatches are unlikely to be due to KASP genotyping errors. Mismatches may be due
to differences among programs of SCA resistant or susceptible haplotypes, or errors in the
phenotype-based resistance classifications (some of which are based on visual ratings under
natural field infestations, which are prone to false positives (*24*)). There were also some
genotype-phenotype mismatches in public germplasm checks used by commercial and public
programs (Fig. 5C). In nearly all cases, further investigation revealed that mismatches were due
to unexpected heterogeneity in public germplasm within or among breeding programs (Supp.
Table S4).

DISCUSSION

***RMES1* is a major resistance gene underlying evolutionary rescue of sorghum worldwide**
Understanding the genetics of evolutionary rescue, including the genetic architecture and
molecular basis, could contribute to more resilient conservation and breeding strategies (*25*).
Here we hypothesized, parsimoniously, that a single Mendelian SCA resistance locus *RMES1*
could underlie the global evolutionary rescue of sorghum to the new *M. sacchari* superclone.
Previous studies had suggested that a single dominant locus is responsible for SCA resistance in
families derived from resistant Chinese grain sorghum variety Henong 16 (H16) and susceptible
BTx623, or families derived from US breeding lines, resistant RTx2738 and susceptible CK60
(*22*, *26*). The H16 resistance was mapped to a ~130 kb region at 2.7 Mb on chromosome 6
(*RMES1*) (*22*). Consistent with the *RMES1* evolutionary rescue hypothesis, the genome region
with the highest F_{ST} in the HBP colocalized precisely with *RMES1* (Fig. 3). Together, the
evolutionary genome scan (Fig. 3) and multi-program marker validation (Fig. 5) provides strong
evidence that *RMES1* is the major SCA resistance locus globally, shared across the Americas,
Asia, and Africa. However, our findings do not preclude the hypothesis that other SCA
resistance loci were selected in Haiti and were required for the evolutionary rescue. In particular,
other F_{ST} scan peaks on chromosome 2, 7, 8, and 9 (Fig. 3) could correspond to other SCA
resistance loci. Given that SCA resistance is fixed in the Haitian program, further population
development and quantitative trait locus mapping for SCA resistance will be necessary to test
this hypothesis.
Identifying the causal variant underlying SCA resistance would advance our
understanding of aphid resistance mechanisms in plants (*27*) and facilitate development of
perfectly-predictive molecular markers for SCA resistance breeding (*28*). Our comparative
genomic analysis between the resistant PI 276837 and the three susceptible reference genomes
identified four candidate genes with putative functional variants within the *RMES1* locus
(Sobic.006G017200, Sobic.006G017332, Sobic.006G017400 and Sobic.006G017500; Supp. File
S4). Three of the four genes in the candidate region encode leucine-rich repeat (LRR) proteins, a
gene family involved in immune responses to invading pathogens and insects (*29*). Given that
some LRR genes mediate plant resistance to aphids and other phloem-feeding insects (*27*) these
genes represent promising candidates for the *RMES1* causative gene. Functional annotation and
sequence comparison between the resistant and susceptible accession identified non-synonymous
variants only in Sobic.006G017200 and Sobic.006G017500 (Fig. 4), suggesting these NLR are

promising candidates for the *RMES1* gene. Fine-mapping and positional cloning will be needed
to test these hypotheses and positively identify the causative variant.

Evolutionary rescue of sorghum depended on a half century of global germplasm exchange
In the twentieth century, sorghum genebanks and breeding programs exchanged germplasm
widely (*30*, *31*). Based on pedigree records and morphology we hypothesized that the Haitian
breeding population originated from global admixed germplasm with a primary contribution of
Ethiopian caudatum of the zerazera working group. Consistent with this hypothesis, HBP
genotypes clustered with caudatum sorghum of East Africa (Fig. 2), but admixture analysis
identified a contribution from durra and guinea sorghum from West Africa (Fig. 2D). Combining
population genomics findings (Fig. 2, 3) with genebank and pedigree records (*32*, *33*), we can
map the history of global germplasm exchange that led to the evolutionary rescue of sorghum in
Haiti following the SCA outbreak (Fig. 6A), as well as the spread of the SCA resistance allele
from Ethiopia to breeding programs around the world (Fig. 6B). Notably, the evolutionary rescue
of sorghum in the Americas (Haiti and US) involved germplasm and knowledge exchange over a
period of >50 years, involving nine countries on three continents.
In the case of the SCA outbreak, the global sorghum improvement community was
fortunate that the rare SCA resistance allele originated in East African caudatums, since this
germplasm is preferred by many sorghum breeders worldwide and widely used by breeding
programs in Africa, Asia, and the Americas (*30*, *34*). The SCA resistance allele appears to have
been inadvertently spread across sorghum breeding programs across the three continents long
before the recent SCA outbreak (Fig. 6). For example, SC110, a converted version of an Ethiopia
caudatum landrace (PI 257599/IS 12610) identified as SCA resistant in several world regions
(*23*, *35*), is a major contributor to the pedigrees of most SCA-resistant breeding lines in the US
(Fig. 6B) (*36*). The same progenitor line (IS 12610) was used by breeding programs in West
Africa (Fig. 6B) as a parent of IRAT204 (CE151-262; PI 656031), a widely-adopted variety (*37*)
and key progenitor of current West African breeding programs (*34*).
Another potential benefit of germplasm exchange is the maintenance of diversity in
breeding programs following strong selection, including evolutionary rescue. Given the strong
selection on the HBP during the SCA outbreak, it might be expected that the post-selection HBP
no longer retains sufficient diversity for future adaptation and genetic gain (*7*). However, the
HBP was founded with diverse admixed global germplasm (Fig. 2) and extensively intercrossed,
so it appears to have retained sufficient genetic diversity for future adaptation and crop
improvement. We observe only a modest reduction in nucleotide diversity observed throughout
the genome of the HBP relative to global accessions, East African caudatums, or a major public
pre-breeding program (Fig. 1E; Supp. Fig. S7). Recombination during intercrossing cycles (prior
to the SCA outbreak) presumably reshuffled the SCA resistance allele onto many backgrounds,
suggesting that the intercrossing approach was critical to allow the Haitian program to retain
diversity for future genetic gain and adaptation.

Rapid discovery and deployment of a global trait-predictive molecular marker using evolutionary population genomics

Molecular marker development based on phenotype-to-genotype mapping of trait loci (e.g. linkage or association mapping) is limited by availability of suitable mapping populations, phenotyping capacity, and genotyping resources, which can take years to develop (*13*, *38*). For instance, spatial and temporal variability of SCA pressure in field trials limits the effectiveness of field phenotyping (*24*), while greenhouse assays can be complicated and time-consuming for lower-resourced programs. Thus, an evolutionary genomics approach, which leverages a history of selection by smallholder farmers or plant breeders, could have advantages for marker discovery. Despite wide use of evolutionary genome scans in crops, the hypotheses generated on adaptive loci are rarely, if ever, tested by independent experimental approaches (e.g. with near isogenic lines) (*39*). To our knowledge, this is the first example where an evolutionary or population genomic scan led directly to molecular breeding technology in use in commercial and public varietal development (Fig. 3, 5).

Here we demonstrated the effectiveness of the evolutionary population genomic approach, showing that a marker discovered in a single developing-country breeding program (Chibas-Haiti) can link crop improvement efforts across three continents (North America, Africa, Asia; Fig. 5A, 6) and across the commercial and public sector (Fig. 5A, C). Thus, our findings establish the value of evolutionary population genomics to facilitate and guide global crop improvement. The KASP marker developed and validated in this study can facilitate the rapid conversion of existing farmer-preferred varieties for SCA resistance (e.g. by marker-assisted introgression) (*40*). While the *RMES1* resistance allele is currently conferring effective resistance, a further biotype shift in the aphid could overcome this gene. Several biotype shifts occurred in the 1960-1980s for the greenbug aphid *Schizaphis graminum* (*41*) and slowed genetic gain in sorghum for many years (*42*). The markers developed here could facilitate identification of new SCA resistance genes, via by counterselection of *RMES1* allele to reveal novel SCA resistance. These outsourced KASP markers are convenient for breeding programs, since they require no laboratory labor or facilities, and are low cost relative to dedicated field or greenhouse phenotyping capacity, at ~$2 per sample for DNA extraction and marker genotyping (*43*).

Synergy of long-standing germplasm exchange practices with new genomics technologies

In this study, we integrated evolutionary population genomic analyses and historical records on global germplasm exchange to show that the recent evolutionary rescue of sorghum depended on >50 years of germplasm exchange. Germplasm exchange led to global diffusion of a rare SCA resistance allele, sometimes purposely and sometimes inadvertently, from smallholder farmers in Ethiopian highlands across breeding programs in Africa, the Americas, and Asia. Over the past several decades, movement of crop genetic resources through international cooperation of germplasm exchange has provided access to adaptive genetic variation for crop improvement (*31*). However, germplasm exchange is increasingly restricted due to commercial or institutional interests asserting intellectual property (IP) rights and governments asserting national or local

sovereignty over genetic resources (*44*). While IP rights and sovereignty are important
considerations, the question remains how to balance these aims with the benefits of free
exchange of global public goods (*45*, *46*). While we are not in a position to resolve these societal
tradeoffs, our study does highlight the global food security benefits of germplasm exchange and
the opportunities that could be lost due to restrictions on exchange.
Taken together, our findings suggest that new genomic technologies will be most
powerful when leveraged with global exchange of germplasm and knowledge. No matter how
powerful new genomic technologies are in terms of accuracy or throughput, their utility will
depend on the germplasm assayed, since all genetic mapping approaches require effective
recombination and allelic diversity (*47*, *48*). Global germplasm exchange vastly increases both
these parameters, providing a "bank" of historical recombinations and allelic variants that can be
rapidly leveraged with new genomic tools (Fig. 3, 4). Therefore, our best opportunity to address
challenges of global change may be to leverage new genomic technologies with long-standing
practices of global germplasm exchange.

MATERIALS AND METHODS

Sorghum breeding and production in Haiti
The Chibas sorghum breeding program was launched in 2013 using admixed global germplasm,
including heterogeneous breeding material from West Africa carrying *ms3* nuclear male sterility,
and inbred global accessions. During 2015–2018, the material was selected in breeding nurseries
under low-input conditions (approximating local smallholder practices) and extensively
intercrossed using the *ms3* sterility system. No insecticides were used to limit SCA infestations
in breeding nurseries in this period and natural SCA infestations were intense during this period
(e.g. Fig. 1C). Note, selection pressure on sorghum by SCA in Haiti is expected to be greater
than in temperate zone (e.g. U.S.) because the SCA infestation occurs year-round in this tropical
environment. Annual sorghum production estimates for Haiti are based on FAOSTAT (2009–
2014 and 2018) (*49*) and the USDA forecast for 2019-2020 (*19*). FAOSTAT data for 2015–2017
and 2019 was not used because it was based on imputation ("FAO data based on imputation
methodology") that did not account for the known effects of SCA (e.g. "this aphid spread
throughout the country and decimated Haiti sorghum production") (*19*). Production for the
missing years of SCA outbreak was inferred based on 2009 agriculture survey acreage prior to
infestation in each region and assessment of sorghum production levels compared to pre-
infestation levels, adjusted to FAOSTAT (1990–2014) production averages for each region.

Plant genetic resources
The HBP (N = 296) are inbred lines derived from a recurrent selection breeding population
developed by intercrossing germplasm that survived natural SCA infestation. For genomic DNA
extraction, fresh leaf tissue of each accession was collected from two weeks old seedlings raised
in a greenhouse. Tissue was lyophilized for two days and then grounded up using a 96-well plate
plant tissue grinder (Retsch Mixer Mill). Genomic DNA was extracted using the BioSprint 96
DNA Plant Kit (QIAGEN), quantified using Quant-iT™ PicoGreen® dsDNA Assay Kit, and

normalized to 10 ng/uL. An additional set of global accessions (GDP, $N = 767$) was assembled
based on a published data set (*50*, *51*) including sorghum accessions from 52 countries on five
continents and all major botanical types (Supp. Fig. S1, Supp. File S1). The GDP accessions
included 164 caudatum, 96 guinea, 81 durra, 57 bicolor, and 47 kafir accessions, along with 288
of other botanical types and 34 accessions of unknown botanical type.

Genotyping-by-sequencing
Genotypes for the 296 Haitian breeding lines were generated with genotyping-by-sequencing.
Genomic DNA digestion, ligation and PCR amplification processes were performed according to
the methods previously described (*50*). The libraries were sequenced using the single-end 100-
cycle sequencing by Illumina HiSeq2500 (Illumina, San Diego CA, USA) at the University of
Kansas Medical Center, Kansas City, MO, USA. A total of 220 million reads for the HBP were
combined with published data for the GDP (*50*) for SNP calling. TASSEL 5 GBS v2 pipeline
(*52*) was used to perform the SNP calling of the sequence data obtained from Illumina
sequencing. Reads were aligned to the BTx623 sorghum reference genome v.3.1 (*53*) with the
Burrows-Wheeler Alignment (*54*). The SNPs were filtered for 20% missingness, then missing
data were imputed using BEAGLE 4.0 (*55*). Genotyping data are available at Dryad [*accession*
to be added following acceptance].

Population genomic analyses
Genome-wide nucleotide diversity (π) and Tajima's *D* statistics for HBP and GDP were
estimated based on a non-overlapping sliding window of 1 Mbp across the genome using
VCFtools (*56*). The characterization of the population structure of the HBP was based on a
discriminant analysis of principal components (DAPC) in the Adegenet package in R (*57*). A
distance matrix calculated based on a modified Euclidean distance model was used to create a
cladogram based on a neighbor-joining algorithm in TASSEL (*58*). Neighbor-joining analysis
was visualized using the APE package in R (*59*). The population structure of the germplasm
panel was further assessed by the Bayesian model-based clustering method implemented in the
ADMIXTURE program (*60*). Pairwise SNP differentiation (F_{ST}) between the HBP and the GDP
were calculated and outlier loci were detected based on an inferred distribution of neutral F_{ST}
using the R Package OutFLANK (*61*).

Whole genome resequencing
Around the 130 kb mapped interval in BTx623, SNPs from 10 sorghum accessions with known
SCA resistance status were examined to search for functional mutations responsible for SCA
resistance. Six of the 10 resequenced accessions represent known susceptible lines, which
include RTx430 (PI 655996), BTx623 (PI 564163), Tx7000 (PI 655986), Tx2737 (PI 655978),
BTx642, and RTx436. The remaining four resequenced accessions represent known resistant
lines, which includes PI 257599 (SC110 original exotic parent), PI 276837 (SC170 original
exotic parent), PI 534157 (SC170), and IS 36563 (IRAT204). These samples were used pre-
publication for this interval analysis with permission from TERRA-REF (Mockler), JGI
Sorghum Pan-genome project (Mockler), BMFG Sorghum Genomic Toolbox (Mockler and

Morris), JGI Sorghum Diversity project (John Mullet), and the JGI EPICON project (Vogel).
The reads were mapped to *Sorghum bicolor* v3.1 using bwa-mem. The bam file was filtered for
duplicates using Picard (http://broadinstitute.github.io/picard) and realigned around indels using
GATK (*62*). Multi-sample SNP calling was done using SAMtools mpileup and Varscan V2.4.0
with a minimum coverage of 8 and a minimum alternate allele frequency of four. Repeat content
of the genome was masked using 24 bp kmers. Kmers that occur at a high frequency, up to 5%,
were masked. SNPs around 25 bp of the mask were removed for further analysis. A SNP was
included for further analysis only when it has coverage in 75% of the samples, and a MAF >
0.005. Functional annotation of the variants within the *RMES1* locus was performed using
SNPEff.

De novo genome sequencing

De novo genome assembly of the resistance sorghum line PI 276837 was used to perform
comparative genomic analysis to identify the causative variant for SCA resistance at the *RMES1*
locus. PI 276837 main assembly consisted of 101.47x of PACBIO coverage with an average read
size of 11,931 bp. The genome was assembled using Canu 1.8, a fork of the Celera Assembler
designed for high-noise single-molecule sequencing. The resulting sequence was polished using
ARROW. The assembled genome resulted in contig N50 sizes ranging from 14 to 19 kb and
scaffold N50 sizes ranging from 5 to 65 kb. Sequence variations at *RMES1* locus between the *de*
novo sequence of PI 276837 were compared to the reference genomes of BTx623, Tx430, and
BTx642.

KASP marker development

SNPs from the F_{ST} genomic selection scan were selected for development into markers based on
several factors: LOD score of the F_{ST} analysis, proximity to *RMES1* locus, and suitability of the
flanking sequence for KASP assay development. The KASP assays were developed utilizing a
third-party genotyping service provider, Intertek AgriTech (Alnarp, Sweden), who designed the
KASP assays via the Kraken software. All genomic DNA extraction and KASP genotyping were
performed by Intertek using two 6 mm leaf punches dried with silica beads. Initial technical
validation of the KASP marker was performed using known resistant (SC110, IRAT204 and
Tx2783) and susceptible (KS585 and Tx7000) sorghum lines. Further validation was performed
by genotyping a panel of 10 known resistant and 28 known susceptible lines, along with multiple
F1 crosses of each of the lines. The KASP markers developed for SCA resistance selection
(Supp. File S6) are publicly available through the third-party genotyping service provided by
Intertek. For further information on accessing markers contact the corresponding author.

Marker validation in public and commercial breeding programs

To test the predictiveness of the marker, a population segregating for SCA resistance was
developed by crossing the susceptible Tx430 and resistant IRAT204. F_3 and F_4 lines of the
Tx430 x IRAT204 population were genotyped with the KASP marker together with the
susceptible and resistant parents. The same population was evaluated for SCA reaction using a
free-choice flat-screen trial in the greenhouse. Tx2783 and SC110 were included as known

resistant genotypes, along with the known susceptible genotypes, KS 585 and Tx7000 (*63*, *64*).
Free-choice flat-screen assay, data collection (damage rating, SPAD score, and plant height
difference), and analysis were conducted as previously described (*24*). Validation of the KASP
marker across different breeding programs was performed in eight breeding programs, five
commercial and three public in the US. Each program collected tissue samples from known
tolerant and susceptible parental breeding lines, F_1s of the parental lines, and later generation
lines from their SCA tolerance breeding populations; the SCA reaction phenotypes of the late
generation lines may or may not have been known. For the parental breeding lines, both
technical replicates (tissue samples from the same plant) and biological replicates (tissue samples
from separate plants) were collected in order to test both the technical function of the markers
and the reliability of the germplasm, respectively. Additionally, most programs included public
sources (e.g. Tx2783) of known SCA tolerance as checks. Tissue samples were sent to Intertek,
who extracted DNA and performed the KASP genotyping.

ACKNOWLEDGMENTS

This study is made possible by the support of the American People provided to the Feed the
Future Innovation Lab for Collaborative Research on Sorghum and Millet through the United
States Agency for International Development (USAID) under Associate Award No. AID-OAA-
LA-16-00003, "Feed the Future Innovation Lab for Genomics-Assisted Sorghum Breeding". The
contents are the sole responsibility of the authors and do not necessarily reflect the views of
USAID or the United States Government. The work conducted by the US Department of Energy
Joint Genome Institute is supported by the Office of Science of the US Department of Energy
under Contract No DE-AC02-05CH11231. We thank the Joint Genome Institute and
collaborators for pre-publication access to the genomes of Tx430 and Tx642 for use in this study.
Additional support was provided by the Bill & Melinda Gates Foundation under the "Sorghum
Genomics Toolkit" project. We thank the breeding programs that participated in the marker
testing. We thank Matt Davis for technical support and Jesse Lasky for comments on the
manuscript. Such use does not constitute an official endorsement or approval by the United
States Department of Agriculture or the Agricultural Research Service of any product or service
to the exclusion of others that may be suitable. USDA is an equal opportunity provider and
employer. The authors declare that they have no competing interests. All data needed to evaluate
the conclusions in the paper are present in the Supplementary Materials or Dryad.

Figure 1: Evolutionary rescue following a continental outbreak of a sorghum pest
(A) Infestation of sugarcane aphid (SCA), *M. sacchari*, on a commercial hybrid in the US
sorghum-growing production region (Kansas). (B) SCA infestation on a traditional sorghum
variety on a smallholder farm in Haiti (brown plant in foreground; green leaves in background
are maize and wild grasses). (C) Reaction of susceptible (brown plants; foreground) and resistant
(green plants; background) sorghum breeding lines under natural SCA infestation during
breeding trials in Haiti. (D) Estimates of annual sorghum production in Haiti (2006-2019),
indicating the start of the SCA outbreak (1, red arrow) and the start of national distribution of
SCA resistant variety, Papépichon (2, blue arrow). (E) Genome-wide nucleotide diversity (π) in
the Haitian breeding population (red line) compared to a global diversity panel (blue bars).
Nucleotide diversity was calculated for a non-overlapping sliding window of 1 Mbp across the
genome. The grey vertical dashed lines indicate the position of *a priori* candidate genes for
breeding targets of the Haiti program which colocalized with genomic regions of reduced π (see
Supp. File S3 for details).

Figure 2: **Population structure of the Haitian breeding population in relation to global sorghum diversity reflects its derivation from East African germplasm**. Genetic relatedness of the Haitian breeding population (HBP) to the global diversity assessed by neighboring joining method, color-coded by botanical type (A) or highlighting the close relationship between the HBP and East African caudatum (EAC) germplasm (C) Scatterplot of the first two principal component (PC) of genome wide SNP variation, demonstrating the clustering of HBP within EAC germplasm. (D) Bayesian hierarchical clustering of the HBP and GDP with the probability of membership (*Q*) in each of K = 8 ancestral populations. The Q-value bar plots are arranged by botanical types to reflect the relationship of the HBP to the GDP. Note, color-coding of the bar plots in panel D is arbitrary and does not reflect the color-code in panels A-C. BR = Bicolor, CG = caudatum-guinea, CK= caudatum-kafir, BC = bicolor-caudatum, BD = bicolor-durra, CD = caudatum durra, O = others (includes botanical types containing less than 10 individuals), W= wild.

512
Figure 3: **Genome scan for selection identifies the major aphid resistance allele at *RMES1***
originating in Ethiopia. (A) Genome-wide scan for selection in the Haitian breeding population
using fixation index (F_{ST}) with the -log(F_{ST} p-value) (y-axis) plotted against position on the
chromosome (x-axis). (B) Detailed view (5 Mb) of top F_{ST} peak on chromosome 6 that
colocalizes with the *RMES1* locus. The ~130 kb region from 2,667,082 to 2,796,847 bp
corresponding to the published *RMES1* interval is denoted with the red bar. (C) Global allele
distribution of the SNP that showed the highest F_{ST} value (S6_2995581), which colocalized with
the *RMES1* locus. Allelic state for georeferenced global germplasm is denoted with points. Allele
frequencies in the United States (C=151, T=2, T/C=1) and Haiti (C=6, T=287, T/C =3) breeding
germplasm, denoted in pie charts with area proportional to number of accessions, show the allele
is almost fixed in Haitian breeding germplasm and rare in U.S. breeding germplasm.
524
525

527
Figure 4: Whole-genome resequencing and local association mapping identifies potential
causative variants at *RMES1*. Functional annotation and association mapping of nucleotide
polymorphisms within the *RMES1* locus across a set of 13 diverse sorghum accessions with
known SCA resistance or susceptibility. The -log of *p*-values of local marker-trait association
scan plotted against the chromosomal positions at the *RMES1* locus on chromosome 6. Variants
are color-coded by annotation generated by the SnpEff program. Blue bars represent the seven
annotated genes within the *RMES1* interval (A = Sobic.006G017000, B = Sobic.006G017100, C
= Sobic.006G017200, D = Sobic.006G017332 and Sobic.006G017266, E = Sobic.006G017400,
F = Sobic.006G017500.v3.1). Grey bars indicate genes outside of the *RMES1* interval as
originally defined (*22*). 3'UTR: 3 prime UTR variant, 5'UTR: 5 prime UTR variant, downstream:
Downstream gene variant, intergenic: intergenic region, intron: Intron variant, missense:
Missense variant, SAI_SRI: splice acceptor/intron or splice region intron variants, SG_SL: stop
gained or stop loss variant, synonymous: Synonymous variant upstream: Upstream gene variant
541
542

Figure 5: Multi-program evaluation of a molecular marker developed based on the selective sweep validates its global utility.

(A) The KASP marker predicts SCA resistance in independent US × Senegal breeding lines. The histogram represents the aphid damage ratings of F_4 lines from a Tx430 × IRAT204 ($N = 22$) family under infestation by *M. sacchari* at the seedling stage in a choice greenhouse assay. The cyan bars represent the aphid damage ratings for lines carrying the +/+ genotypes at the SNP 06_02892438, while the red bars represent aphid damage ratings of the lines carrying the -/- genotypes. The blue dashed lines represent the average aphid damage rating of the resistant checks Tx2783, IRAT204, and SC110 (green, blue, and purple dashed lines, respectively) while red dashed line represents the average damage rating of both susceptible checks, Tx7000 and Tx430. (B) Representative SCA reaction from the choice greenhouse assay for an F4 line carrying the +/+ genotype (left) versus the susceptible parent Tx430 (right). (C) Evaluation of the same marker in eight US breeding programs. Percent match of KASP marker genotyping prediction with breeder-provided SCA resistance classification for five commercial breeding programs and three public breeding programs. ND = Not determined.

Figure 6: Evolutionary rescue of sorghum through >50 years of global exchange of germplasm and knowledge. (A) Germplasm and knowledge exchange inferred from pedigree records and genomic analyses. Germplasm exchange is denoted by solid lines. Knowledge exchange through scientific literature is denoted in dashed lines. The star indicates the inferred origin of the SCA resistance allele in the Ethiopian highlands, with at least two paths to the Americas, via IS 23572 (yellow line) or PI 257599 (red line). (B) Pedigree relationships among global accessions, breeding lines, breeding programs, or varieties, color-coded by inferred SCA resistance or susceptibility. Note, with respect to US commercial hybrids, the diagram is illustrative and is not meant to imply that all US commercial hybrids used Tx2783 as the SCA resistance donor. Some known pedigree information has been omitted from the diagram for clarity.

REFERENCES

1. S. R. Palumbi, Humans as the world's greatest evolutionary force. *Science*. **293**, 1786–1790 (2001).
2. T. Wheeler, J. von Braun, Climate change impacts on global food security. *Science*. **341**, 508–513 (2013).
3. Gonzalez, Ronce, Ophélie, Ferriere, Regis, Hochberg, Michael E., Evolutionary rescue: an emerging focus at the intersection between ecology and evolution. *Philos. Trans. R. Soc. B Biol. Sci.* **368**, 20120404 (2013).
4. G. Bell, Evolutionary Rescue. *Annu. Rev. Ecol. Evol. Syst.* **48**, 605–627 (2017).
5. B. M. Peter, E. Huerta-Sanchez, R. Nielsen, Distinguishing between Selective Sweeps from Standing Variation and from a De Novo Mutation. *PLOS Genet.* **8**, e1003011 (2012).
6. H. K. Alexander, G. Martin, O. Y. Martin, S. Bonhoeffer, Evolutionary rescue: linking theory for conservation and medicine. *Evol. Appl.* **7**, 1161–1179 (2014).
7. H. A. Orr, R. L. Unckless, The population genetics of evolutionary rescue. *PLOS Genet.* **10**, e1004551 (2014).
8. R. S. Turner, After the famine: Plant pathology, Phytophthora infestans, and the late blight of potatoes, 1845–1960. *Hist. Stud. Phys. Biol. Sci.* **35**, 341–370 (2005).
9. J. M. Burke, J. C. Burger, M. A. Chapman, Crop evolution: from genetics to genomics. *Curr. Opin. Genet. Dev.* **17**, 525–532 (2007).
10. B. A. Wilson, P. S. Pennings, D. A. Petrov, Soft Selective Sweeps in Evolutionary Rescue. *Genetics*. **205**, 1573–1586 (2017).
11. P. S. Pennings, J. Hermisson, Soft Sweeps III: The Signature of Positive Selection from Recurrent Mutation. *PLOS Genet.* **2**, e186 (2006).
12. J. J. Vitti, S. R. Grossman, P. C. Sabeti, Detecting natural selection in genomic data. *Annu. Rev. Genet.* **47**, 97–120 (2013).
13. C. R. Cavanagh, S. Chao, S. Wang, B. E. Huang, S. Stephen, S. Kiani, K. Forrest, C. Saintenac, G. L. Brown-Guedira, A. Akhunova, D. See, G. Bai, M. Pumphrey, L. Tomar, D. Wong, S. Kong, M. Reynolds, M. L. da Silva, H. Bockelman, L. Talbert, J. A. Anderson, S. Dreisigacker, S. Baenziger, A. Carter, V. Korzun, P. L. Morrell, J. Dubcovsky, M. K. Morell, M. E. Sorrells, M. J. Hayden, E. Akhunov, Genome-wide comparative diversity uncovers multiple targets of selection for improvement in hexaploid wheat landraces and cultivars. *Proc. Natl. Acad. Sci.* (2013), doi:10.1073/pnas.1217133110.
14. M. B. Hufford, J. C. Berny Mier y Teran, P. Gepts, Crop Biodiversity: An Unfinished Magnum Opus of Nature. *Annu. Rev. Plant Biol.* **70**, 727–751 (2019).
15. R. Monk, C. Franks, J. Dahlberg, in *Yield Gains in Major US Field Crops* (Crop Science Society of America, 2014; https://dl.sciencesocieties.org/publications/books/abstracts/cssaspecialpubl/yieldgainsinmaj/293), pp. 293–310.
16. R. D. Bowling, M. J. Brewer, D. L. Kerns, J. Gordy, N. Seiter, N. E. Elliott, G. D. Buntin, M. O. Way, T. A. Royer, S. Biles, E. Maxson, Sugarcane Aphid (Hemiptera: Aphididae): A New Pest on Sorghum in North America. *J. Integr. Pest Manag.* **7** (2016), doi:10.1093/jipm/pmw011.
17. J. W. Gordy, M. J. Brewer, R. D. Bowling, G. D. Buntin, N. J. Seiter, D. L. Kerns, F. P. F. Reay-Jones, M. O. Way, Development of Economic Thresholds for Sugarcane Aphid (Hemiptera: Aphididae) in Susceptible Grain Sorghum Hybrids. *J. Econ. Entomol.* **112**, 1251–1259 (2019).

18.　S. Nibouche, L. Costet, J. R. Holt, A. Jacobson, A. Pekarcik, J. Sadeyen, J. S. Armstrong,
G. C. Peterson, N. McLaren, R. F. Medina, Invasion of sorghum in the Americas by a new
sugarcane aphid (Melanaphis sacchari) superclone. *PLOS ONE*. **13**, e0196124 (2018).

19.　USDA-FAS, Grain and Feed Annual Report: Haiti (HA2020-0001) (2020), (available at
https://apps.fas.usda.gov/newgainapi/api/Report/DownloadReportByFileName?fileName=
Grain%20and%20Feed%20Annual_Port-au-Prince_Haiti_04-15-2020).

20.　W. Calvin, Effects of intercropping and biological insecticides on sugarcane aphid
(Hemiptera: aphididae) infestations on sorghum, and identification of natural enemies and
alternate hosts in Haiti, 77 (2019).

21.　K. T. Muleta, G. Pressoir, G. P. Morris, Optimizing genomic selection for a sorghum
breeding program in Haiti: a simulation study. *G3 Genes Genomes Genet.* **9**, 391–401
(2019).

22.　F. Wang, S. Zhao, Y. Han, Y. Shao, Z. Dong, Y. Gao, K. Zhang, X. Liu, D. Li, J. Chang, D.
Wang, Efficient and fine mapping of RMES1 conferring resistance to sorghum aphid
Melanaphis sacchari. *Mol. Breed.* **31**, 777–784 (2013).

23.　J. S. Armstrong, W. L. Rooney, G. C. Peterson, R. T. Villenueva, M. J. Brewer, D. Sekula-
Ortiz, Sugarcane Aphid (Hemiptera: Aphididae): Host Range and Sorghum Resistance
Including Cross-Resistance From Greenbug Sources. *J. Econ. Entomol.* **108**, 576–582
(2015).

24.　J. S. Armstrong, L. Mbulwe, D. Sekula-Ortiz, R. T. Villanueva, W. L. Rooney, Resistance
to Melanaphis sacchari (Hemiptera: Aphididae) in forage and grain sorghums. *J. Econ.
Entomol.* **110**, 259–265 (2017).

25.　C. Burgarella, A. Barnaud, N. A. Kane, F. Jankowski, N. Scarcelli, C. Billot, Y. Vigouroux,
C. Berthouly-Salazar, Adaptive Introgression: An Untapped Evolutionary Mechanism for
Crop Adaptation. *Front. Plant Sci.* **10** (2019), doi:10.3389/fpls.2019.00004.

26.　H. M. Tetreault, S. Grover, E. D. Scully, T. Gries, N. A. Palmer, G. Sarath, J. Louis, S. E.
Sattler, Global Responses of Resistant and Susceptible Sorghum (Sorghum bicolor) to
Sugarcane Aphid (Melanaphis sacchari). *Front. Plant Sci.* **10** (2019),
doi:10.3389/fpls.2019.00145.

27.　Y. Jiang, C.-X. Zhang, R. Chen, S. Y. He, Challenging battles of plants with phloem-
feeding insects and prokaryotic pathogens. *Proc. Natl. Acad. Sci.* **116**, 23390–23397
(2019).

28.　J. R. Andersen, T. Lübberstedt, Functional markers in plants. *Trends Plant Sci.* **8**, 554–560
(2003).

29.　L. McHale, X. Tan, P. Koehl, R. W. Michelmore, Plant NBS-LRR proteins: adaptable
guards. *Genome Biol.* **7**, 212 (2006).

30.　C. W. Smith, R. A. Frederiksen, in *Sorghum: Origin, History, Technology, and Production*,
C. W. Smith, R. A. Frederiksen, Eds. (Wiley, NY, 2000;
https://books.google.com/books?hl=en&lr=&id=b7vxU44v794C&oi=fnd&pg=PA191&dq=
History+of+Cultivar+Development+in+the+United+States:+From+%22Memoirs+of+A.+B
.+Maunder--
Sorghum+Breeder%22&ots=UuB7XqgYWP&sig=cVMz_e6MAD5WTEKprcSdYOThAs0
), vol. 2, p. 191.

31.　R. R. Klein, F. R. Miller, D. V. Dugas, P. J. Brown, A. M. Burrell, P. E. Klein, Allelic
variants in the PRR37 gene and the human-mediated dispersal and diversification of
sorghum. *Theor. Appl. Genet.*, 1–15 (2015).

32. G. C. Peterson, J. W. Johnson, G. L. Teetes, D. T. Rosenow, Registration of Tx2783 Greenbug Resistant Sorghum Germplasm Line. *Crop Sci.* **24**, 390 (1984).

33. République du Sénégal, Catalogue officiel des espèces et des variétés cultivées au Sénégal [Official catalog of cultivated species and varieties in Senegal] (2012), (available at http://www.fao.org/pgrfa-gpa-archive/sen/docs/senegal_varietes/Catalogue_%20varietes.htm).

34. J. M. Faye, F. Maina, E. A. Akata, B. Sine, C. Diatta, A. Mamadou, S. Marla, S. Bouchet, N. Teme, J.-F. Rami, D. Fonceka, N. Cisse, G. P. Morris, A genomics resource for genetics, physiology, and breeding of West African sorghum. *Plant Genome*. **n/a**, e20075 (2021).

35. L. Mbulwe, G. C. Peterson, J. Scott-Armstrong, W. L. Rooney, Registration of sorghum germplasm Tx3408 and Tx3409 with tolerance to sugarcane aphid [Melanaphis sacchari (Zehntner)]. *J. Plant Regist.* **10**, 51–56 (2016).

36. G. C. Peterson, J. S. Armstrong, B. B. Pendleton, M. Stelter, M. J. Brewer, Registration of RTx3410 through RTx3428 sorghum germplasm resistant to sugarcane aphid [Melanaphis sacchari (Zehntner)]. *J. Plant Regist.* **12**, 391–398 (2018).

37. J. Ndjeunga, K. Mausch, F. Simtowe, in *Crop improvement, adoption, and impact of improved varieties in food crops in sub-Saharan Africa*, T. S. Walker, J. Alwang, Eds. (CABI, Wallingford, 2015; http://www.cabi.org/cabebooks/ebook/20153367543), pp. 123–147.

38. S. V. Nuzhdin, T. L. Turner, Promises and limitations of hitchhiking mapping. *Curr. Opin. Genet. Dev.* **23** (2013), doi:10.1016/j.gde.2013.10.002.

39. S. Hoban, J. L. Kelley, K. E. Lotterhos, M. F. Antolin, G. Bradburd, D. B. Lowry, M. L. Poss, L. K. Reed, A. Storfer, M. C. Whitlock, Finding the genomic basis of local adaptation: pitfalls, practical solutions, and future directions. *Am. Nat.* (2016), doi:10.1086/688018.

40. T. Yohannes, T. Abraha, D. Kiambi, R. Folkertsma, C. T. Hash, K. Ngugi, E. Mutitu, N. Abraha, M. Weldetsion, C. Mugoya, C. W. Masiga, S. de Villiers, Marker-assisted introgression improves Striga resistance in an Eritrean farmer-preferred sorghum variety. *Field Crops Res.* **173**, 22–29 (2015).

41. T. A. Royer, B. B. Pendleton, N. C. Elliott, K. L. Giles, Greenbug (hemiptera: aphididae) biology, ecology, and management in wheat and sorghum. *J. Integr. Pest Manag.* **6** (2015), doi:10.1093/jipm/pmv018.

42. F. R. Miller, Y. Kebede, in *Genetic Contributions to Yield Gains of Five Major Crop Plants* (John Wiley & Sons, Ltd, 1984; https://acsess.onlinelibrary.wiley.com/doi/abs/10.2135/cssaspecpub7.c1), pp. 1–14.

43. M. J. Thomson, High-throughput SNP genotyping to accelerate crop improvement. *Plant Breed. Biotechnol.* **2**, 195–212 (2014).

44. P. Heisey, K. Day-Rubenstein, "Using crop genetic resources to help agriculture adapt to climate change: economics and policy" (SSRN Scholarly Paper ID 2709190, Social Science Research Network, Rochester, NY, 2015), , doi:10.2139/ssrn.2709190.

45. J. Kloppenburg, D. L. Kleinman, The plant germplasm controversy. *BioScience*. **37**, 190–198 (1987).

46. J. Kotschi, B. Horneburg, The Open Source Seed Licence: A novel approach to safeguarding access to plant germplasm. *PLOS Biol.* **16**, e3000023 (2018).

47. R. Bernardo, *Molecular markers and selection for complex traits in plants: Learning from the last 20 years, Crop Science 48 (2008): 1649-1664.* (2008), vol. 48.

709   48.   B. Brachi, G. P. Morris, J. O. Borevitz, Genome-wide association studies in plants: the
710   missing heritability is in the field. *Genome Biol.* **12**, 232 (2011).

711   49.   FAO, FAOSTAT (2021), (available at http://www.fao.org/faostat/en/).

712   50.   G. P. Morris, P. Ramu, S. P. Deshpande, C. T. Hash, T. Shah, H. D. Upadhyaya, O. Riera-
713   Lizarazu, P. J. Brown, C. B. Acharya, S. E. Mitchell, J. Harriman, J. C. Glaubitz, E. S.
714   Buckler, S. Kresovich, Population genomic and genome-wide association studies of
715   agroclimatic traits in sorghum. *Proc. Natl. Acad. Sci.* **110**, 453–458 (2013).

716   51.   J. R. Lasky, H. D. Upadhyaya, P. Ramu, S. Deshpande, C. T. Hash, J. Bonnette, T. E.
717   Juenger, K. Hyma, C. Acharya, S. E. Mitchell, E. S. Buckler, Z. Brenton, S. Kresovich, G.
718   P. Morris, Genome-environment associations in sorghum landraces predict adaptive traits.
719   *Sci. Adv.* **1**, e1400218 (2015).

720   52.   J. C. Glaubitz, T. M. Casstevens, F. Lu, J. Harriman, R. J. Elshire, Q. Sun, E. S. Buckler,
721   TASSEL-GBS: A high capacity genotyping by sequencing analysis pipeline. *PLoS ONE.* **9**,
722   e90346 (2014).

723   53.   R. F. McCormick, S. K. Truong, A. Sreedasyam, J. Jenkins, S. Shu, D. Sims, M. Kennedy,
724   M. Amirebrahimi, B. D. Weers, B. McKinley, A. Mattison, D. T. Morishige, J. Grimwood,
725   J. Schmutz, J. E. Mullet, The Sorghum bicolor reference genome: improved assembly, gene
726   annotations, a transcriptome atlas, and signatures of genome organization. *Plant J.* **93**, 338–
727   354 (2018).

728   54.   H. Li, R. Durbin, Fast and accurate short read alignment with Burrows–Wheeler transform.
729   *Bioinformatics.* **25**, 1754–1760 (2009).

730   55.   B. L. Browning, S. R. Browning, Genotype imputation with millions of reference samples.
731   *Am. J. Hum. Genet.* **98**, 116–126 (2016).

732   56.   P. Danecek, A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E.
733   Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, G. McVean, R. Durbin, 1000 Genomes
734   Project Analysis Group, The variant call format and VCFtools. *Bioinformatics.* **27**, 2156–
735   2158 (2011).

736   57.   T. Jombart, S. Devillard, F. Balloux, Discriminant analysis of principal components: a new
737   method for the analysis of genetically structured populations. *BMC Genet.* **11**, 94 (2010).

738   58.   P. J. Bradbury, Z. Zhang, D. E. Kroon, T. M. Casstevens, Y. Ramdoss, E. S. Buckler,
739   TASSEL: software for association mapping of complex traits in diverse samples.
740   *Bioinformatics.* **23**, 2633–2635 (2007).

741   59.   E. Paradis, J. Claude, K. Strimmer, APE: analyses of phylogenetics and evolution in R
742   language. *Bioinformatics.* **20**, 289–290 (2004).

743   60.   D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in
744   unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).

745   61.   M. C. Whitlock, K. E. Lotterhos, Reliable detection of loci responsible for local adaptation:
746   inference of a null model through trimming the distribution of FST. *Am. Nat.* **186**, S24–S36
747   (2015).

748   62.   G. A. V. der Auwera, M. O. Carneiro, C. Hartl, R. Poplin, G. del Angel, A.
749   Levy☐Moonshine, T. Jordan, K. Shakir, D. Roazen, J. Thibault, E. Banks, K. V. Garimella,
750   D. Altshuler, S. Gabriel, M. A. DePristo, *Curr. Protoc. Bioinforma.*, in press,
751   doi:https://doi.org/10.1002/0471250953.bi1110s43.

752   63.   J. S. Armstrong, W. L. Rooney, G. C. Peterson, R. T. Villenueva, M. J. Brewer, D. Sekula-
753   Ortiz, Sugarcane aphid (Hemiptera: Aphididae): host range and sorghum resistance
754   including cross-resistance from greenbug sources. *J. Econ. Entomol.* **108**, 576–582 (2015).

755 64. S. Paudyal, J. S. Armstrong, K. L. Giles, M. E. Payton, G. P. Opit, A. Limaje, Categories of
756   resistance to sugarcane aphid (hemiptera: aphididae) among sorghum genotypes. *J. Econ.*
757   *Entomol.* **112**, 1932–1940 (2019).
758