A sex-specific evolutionary interaction between *ADCY9* and *CETP*

Isabel Gamache[1,2], Marc-André Legault[1,2,3], Jean-Christophe Grenier[1], Rocio Sanchez[1], Eric Rhéaume[1,2], Samira Asgari[4,5], Amina Barhdadi[1,3], Yassamin Feroz Zada[3], Holly Trochet[1,2], Yang Luo[4,5], Leonid Lecca[6,7], Megan Murray[4], Soumya Raychaudhuri[4,5,8,9,10], Jean-Claude Tardif[1,2], Marie-Pierre Dubé[1,2,3], Julie G. Hussin[1,2]*

Affiliations:

[1] Montreal Heart Institute, Montreal, Québec, Canada

[2] Faculty of Medicine, Université de Montréal, Montreal, Quebec, Canada

[3] Université de Montréal Beaulieu-Saucier Pharmacogenomics Centre, Montreal, Canada

[4] Center for Data Sciences, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02115, USA

[5] Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA 02115, USA

[6] Socios En Salud, Lima, Peru

[7] Department of Global Health and Social Medicine, Harvard Medical School

[8] Centre for Genetics and Genomics Versus Arthritis, Manchester Academic Health Science Centre, University of Manchester, Manchester M13 9PL, UK

[9] Department of Biomedical Informatics, Harvard Medical School, Boston, MA 02115, USA

[10] Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02115, USA

* E-mail : julie.hussin@umontreal.ca

Abstract

Pharmacogenomic studies have revealed associations between rs1967309 in the adenylyl cyclase type 9 (*ADCY9*) gene and clinical responses to the cholesteryl ester transfer protein (CETP) modulator dalcetrapib, however, the mechanism behind this interaction is still unknown. Here, we characterized selective signals at the locus associated with the pharmacogenomic response in human populations and we show that rs1967309 region exhibits signatures of positive selection in several human populations. Furthermore, we identified a variant in *CETP*, rs158477, which is in long-range linkage disequilibrium with rs1967309 in the Peruvian population. The signal is mainly seen in males, a sex-specific result that is replicated in the LIMAA cohort of over 3,400 Peruvians. Analyses of RNA-seq data further suggest an epistatic interaction on *CETP* expression levels between the two SNPs in multiple tissues, which also differs between males and females. We also detected interaction effects of the two SNPs with sex on cardiovascular phenotypes in the UK Biobank, in line with the sex-specific genotype associations found in Peruvians at these loci. We propose that *ADCY9* and *CETP* coevolved during recent human evolution due to sex-specific selection, which points towards a biological link between dalcetrapib's pharmacogene *ADCY9* and its therapeutic target *CETP*.

Introduction

Coronary artery disease (CAD) is the leading cause of mortality worldwide. It is a complex disease caused by the accumulation of cholesterol-loaded plaques that block blood flow in the coronary arteries. The cholesteryl ester transfer protein (CETP) mediates the exchange of cholesterol esters and triglycerides between high-density lipoproteins (HDL) and lower density lipoproteins (1,2). Dalcetrapib is a CETP modulator that did not reduce cardiovascular event rates in the overall dal-OUTCOMES trial of patients with recent acute coronary syndrome (3). However, pharmacogenomic analyses revealed that genotypes at rs1967309 in the *ADCY9* gene, coding for the ninth isoform of adenylate cyclase, modulated clinical responses to dalcetrapib (4). Individuals who carried the AA genotype at rs1967309 in *ADCY9* had less cardiovascular events, reduced atherosclerosis progression, and enhanced cholesterol efflux from macrophages when treated with dalcetrapib compared to placebo (4,5). In contrast, those with the GG genotype had the opposite effects from dalcetrapib. Furthermore, a protective effect against the formation of atherosclerotic lesions was seen only in the absence of both *Adcy9* and *CETP* in mice (6), suggesting an interaction between the two genes. However, the underlying mechanisms linking *CETP* and *ADCY9,* located 50 Mb apart on chromosome 16, as well as the relevance of the rs1967309 non-coding genetic variant are still unclear.

Identification of selection pressure on a genetic variant can help shed light on its importance. Adaptation to different environments often leads to a rise in frequency of variants, by favoring survival and/or reproduction fitness. An example is the lactase gene (*LCT*) (7–11), where a positively selected intronic variant in *MCM6* leads to an escape from epigenetic inactivation of

LCT and facilitates lactase persistence after weaning (12). Results of genomic studies for

phenotypes such as adaptation to high-altitude hypoxia in Tibetans (13), fatty acid metabolism in

Inuits (14) or response to pathogens across populations (15) have also been confirmed by

functional studies (16–20). Thus, population and regulatory genomics can be leveraged to unveil

the effect of genetic mutations at a single non-coding locus and reveal the biological mechanisms

of adaptation.

When two or more loci interact during adaptation, a genomic scan will likely be underpowered to

pinpoint the genetic determinants. In this study, we took a multi-step approach on the *ADCY9* and

CETP candidate genes to specifically study their interaction (Figure 1). We used a joint

evolutionary analysis to evaluate the potential signatures of selection in these genes (Step 1), which

revealed positive selection pressures acting on *ADCY9*. Sex-specific genetic associations between

the two genes are discovered in Peruvians (Step 2), a population in which natural selection for

high-altitude was previously found on genes related to cardiovascular health (21). Furthermore,

our know-down experiments and analyses of large-scale transcriptomics (Step 3) as well as

available phenome-wide resources (Step 4) bring further evidence of a sex-specific epistatic

interaction between *ADCY9* and *CETP*.

Results

Signatures of selection at rs1967309 in *ADCY9* in human populations

The genetic variant rs1967309 is located in intron 2 of *ADCY9*, in a region of high linkage

disequilibrium (LD), in all subpopulations in the 1000 Genomes Project (1000G), and harbors

heterogeneous genotype frequencies across human populations (Figure 2a). Its intronic location

makes it difficult to assess its functional relevance but exploring selective signals around intronic

SNPs in human populations can shed light on their importance. In African populations (AFR), the

major genotype is AA, which is the homozygous genotype for the ancestral allele, whereas in

Europeans (EUR), AA is the minor genotype. The frequency of the AA genotype is slightly higher

in Asia (EAS, SAS) and America (AMR) compared to that in Europe, becoming the most frequent

genotype in the Peruvian population (PEL). Using the integrated haplotype score (iHS) (22) (Step

1a, Figure 1), a statistic that enables the detection of evidence for recent strong positive selection

(typically when $|iHS| > 2$), we observed that several SNPs in the LD block around rs1967309

exhibit selective signatures in non-African populations ($|iHS_{SAS}| = 2.66$, $|iHS_{EUR}| = 2.31$), whereas

no signal is seen in this LD block in African populations (Figure 2b, Supplementary Figure 1,

Supplementary text). Our analyses suggest that this locus in *ADCY9* has been the target of recent

positive selection in several human populations, with multiple, possibly independent, selective

signals detectable around rs1967309. However, recent positive selection as measured by iHS does

not seem to explain the notable increase in frequency for the A allele in the PEL population

($f_A=0.77$), compared to the European ($f_A=0.41$), Asian ($f_A=0.44$), and other American populations

($f_A=0.54$ in AMR without PEL).

To test whether the difference between PEL and other AMR allele frequencies at rs1967309 is

significant, we used the population branch statistic (PBS) (Step 1b, Figure 1). This statistic has

been developed to locate selection signals by summarizing differentiation between populations

using a three-way comparison of allele frequencies between a specific group, a closely related

population, and an outgroup (13). It has been shown to increase power to detect incomplete

selective sweeps on standing variation. Applying this statistic to investigate rs1967309 allele

frequency in PEL, we used Mexicans (MXL) as a closely related group and a Chinese population

(CHB) as the outgroup (Methods). Over the entire genome, the CHB branches are greater than

PEL and MXL branches ($mean_{CHB}$=0.020, $mean_{MXL}$=0.008, $mean_{PEL}$=0.009), which reflects the

expectation under genetic drift. However, the estimated PEL branch length at rs1967309 (Figure

2c), which reflects differentiation since the split from the MXL population ($PBS_{PEL,rs1967309}$=0.051,

empirical p-value = 0.014), surpasses the CHB branch length ($PBS_{CHB,rs1967309}$=0.049, empirical p-

value > 0.05), which reflects differentiation since the split between Asian and American

populations, whereas no such effect is seen in MXL ($PBS_{MXL,rs1967309}$=0.026, empirical p-value >

0.05), or for any other AMR populations. Furthermore, the PEL branch lengths at several SNPs in

this LD block (Figure 2c) are in the top 5% of all PEL branch lengths across the whole genome

($PBS_{PEL,95th}$ = 0.031), whereas these increased branch lengths are not observed outside of the LD

block (Figure 2c). These results are robust to the choice of the outgroup and the closely related

AMR population (Methods).

The increase in frequency of the A allele at rs1967309 is also seen in genotype data from Native

American populations (23), with Andeans showing genotype frequencies highly similar to PEL

(f_A=0.77, Figure 2a). The PEL population has a large Andean ancestry (Methods, Supplementary

Figure 2a,b) and almost no African ancestry, strongly suggesting that the increase in AA genotype

arose in the Andean population and not from admixture with Africans. The PEL individuals that

harbor the AA genotype for rs1967309 do not exhibit a larger genome-wide Andean ancestry than

non-AA individuals (p-value=0.30, Mann-Whitney U test). Overall, these results suggest that the

ancestral allele A at rs1967309, after dropping in frequency following the out-of-Africa event, has

increased in frequency in the Andean population and has been preferentially retained in the

Peruvian population's genetic makeup, potentially because of natural selection.

Evidence for co-evolution between *ADCY9* and *CETP* in Peru

The pharmacogenetic link between *ADCY9* and the CETP modulator dalcetrapib raises the question of whether there is a genetic relationship between rs1967309 in *ADCY9* and *CETP*, both located on chromosome 16. Such a relationship can be revealed by analyzing patterns of long-range linkage disequilibrium (LRLD) (24,25), in order to detect whether specific combinations of alleles (or genotypes) at two loci are particularly overrepresented. To do so, we calculated the genotyped-based linkage disequilibrium (r^2) (Step 2a, Figure 1) between rs1967309 and each SNP in *CETP* with minor allele frequency (MAF) above 0.05. In the Peruvian population, there are four SNPs, (including 2 in perfect LD in PEL) that exhibit r^2 values with rs1967309 that are in the top 1% of r^2 values (Figure 3a) computed for all 37,802 pairs of SNPs in *ADCY9* and *CETP* genes with MAF>0.05 (Methods). Despite the r^2 values themselves being low ($r^2_{rs158477}$=0.080, $r^2_{rs158480;rs158617}$=0.089, $r^2_{rs12447620}$=0.090), these values are highly unexpected for these two genes situated 50 Mb apart (*ADCY9/CETP* empirical p-value<0.006, Supplementary Table 1) and thus correspond to a significant LRLD signal. This signal is not seen in other 1000G populations (Supplementary Table 1). We also computed r^2 between the four identified SNPs' genotypes and all *ADCY9* SNPs with MAF above 0.05 (Figure 3b). The distribution of r^2 values for the rs158477 *CETP* SNP shows a clear bell-shaped pattern around rs1967309 in *ADCY9*, which strongly suggests the rs1967309-rs158477 genetic association detected is not simply a statistical fluke, while the signal in the region for the other SNPs is less conclusive. The SNP rs158477 in *CETP* is also the only one that has a PEL branch length value higher than the 95[th] percentile, also higher than the CHB branch length value ($PBS_{PEL,rs158477}$= 0.062, Supplementary Figure 3a), in line with

the observation at rs1967309. Strikingly, this *CETP* SNP's genotype frequency distribution across

the 1000G and Native American populations resembles that of rs1967309 in *ADCY9* (Figure 3c).

Given that the Peruvian population is admixed (26), particular enrichment of genome segments for

a specific ancestry, if present, would lead to inflated LRLD between these segments (27–30), we

thus performed several admixture-related analyses (Step 2b, Figure 1). No significant enrichment

is seen at either locus and significant LRLD is also seen in the Andean source population (Figure

3-figure supplement 1a,b, Supplementary text). Furthermore, we see no enrichment of Andean

ancestry in individuals harboring the overrepresented combination of genotypes, AA at rs1967309

+ GG at rs158477, compared to other combinations (p-value=0.18, Mann-Whitney U test). These

results show that admixture patterns in PEL cannot be solely responsible for the association found

between rs1967309 and rs158477. Finally, using a genome-wide null distribution which allows to

capture the LRLD distribution expected under the admixture levels present in this sample

(Supplementary text), we show that the r^2 value between the two SNPs is higher than expected

given their allele frequencies and the physical distance between them (genome-wide empirical p-

value=0.01, Figure 3d). Taken together, these findings strongly suggest that the AA/GG

combination is being transmitted to the next generation more often (ie. is likely selectively favored)

which reveals a signature of co-evolution between *ADCY9* and *CETP* at these loci.

Still, such a LRLD signal can be due to a small sample size (29). To confirm independently the

association between genotypes at rs1967309 of *ADCY9* and rs158477 of *CETP*, we used the

LIMAA cohort (31,32), a large cohort of 3,509 Peruvian individuals with genotype information,

to replicate our finding. The ancestry distribution, as measured by RFMix (Methods) is similar

between the two cohorts (Supplementary Figure 2a,b), however, the LIMAA cohort population

structure shows additional subgroups compared to the 1000G PEL population sample

(Supplementary Figure 2c-e): to limit confounders, we excluded individuals coming from these

subgroups (Supplementary text). In this dataset (N=3,243), the pair of SNPs rs1967309-rs158477

is the only pairs identified in PEL who shows evidence for LRLD, with an r^2 value in the top 1%

of all pairs of SNPs in *ADCY9* and *CETP* (*ADCY9/CETP* empirical p-value=0.003, Figure 3-figure

supplement 1c,d, Figure 3-figure supplement 2, Supplementary Table 1). The r^2 test used above is

powerful to detect allelic associations, but the net association measured will be very small if

selection acts on a specific genotype combination rather than on alleles. In that scenario, and when

power allows it, the genotypic association is better assessed by with a χ^2 distributed test statistic

(with four degrees of freedom, χ^2_4) comparing the observed and expected genotype combination

counts (25). The test confirmed the association in LIMAA (χ^2_4=82.0, permutation p-value <0.001,

genome-wide empirical p-value=0.0003, Supplementary text). The association discovered

between rs1967309 and rs158477 is thus generalizable to the Peruvian population and not limited

to the 1000G PEL sample.

Sex-specific long-range linkage disequilibrium signal

Because the allele frequencies at rs1967309 were suggestively different between males and

females (Figure 4-figure supplement 1), we performed sex-stratified PBS analyses, which

suggested that the LD block around rs1967309 is differentiated between sexes in the Peruvians

(Figure 4-figure supplement 2, Supplementary text). We therefore explored further the effect of

sex on the LRLD association found between rs1967309 and rs158477 and performed sex-stratified

LRLD analyses. These analyses revealed that the correlation between rs1967309 and rs158477 is

only seen in males in PEL (Figure 4a,b, Supplementary Figure 4a,b, Supplementary Table 1): the

r^2 value rose to 0.348 in males (*ADCY9/CETP* empirical p-value=8.23x10^{-5}, genome-wide

empirical p-value<2.85 x 10^{-4}, N=41) and became non-significant in females (*ADCY9*/*CETP*

empirical p-value=0.78, genome-wide empirical p-value=0.80, N=44). In the Andean population,

the association between rs1967309 and rs158477 is not significant when we stratified by sex

(Supplementary Table 1), but we still see significant association signals with rs158477 at other

SNPs in *ADCY9* LD block in both sexes (Figure 4-figure supplement 3). The LRLD result in PEL

cannot be explained by differences of Andean ancestry proportion between males and females (p-

value=0.27, Mann-Whitney U test). A permutation analysis that shuffled the sex labels of samples

established that the observed difference between the sexes is larger than what we expect by chance

(p-value=0.002, Supplementary Figure 4c, Supplementary text). In the LIMAA cohort, we

replicate this sex-specific result (Figure 4c,d, Supplementary Table 1) where the r^2 test is

significant in males (*ADCY9*/*CETP* empirical p-value=0.003, N=1,941) but not in females

(*ADCY9/CETP* empirical p-value=0.52, N=1,302). The genotypic χ_4^2 test confirms the association

between *ADCY9* and *CETP* is present in males (χ_4^2 = 56.6, permutation p-value=0.001, genome-

wide empirical p-value=0.002, Supplementary text), revealing an excess of rs1967309-AA +

rs158477-GG. This is also the genotype combination driving the LRLD in PEL. In females, the

test also shows a weaker but significant effect (χ_4^2 = 37.0, permutation p-value = 0.017, genome-

wide empirical p-value = 0.001) driven by an excess of a different genotype combination,

rs1967309-AA + rs158477-AA, which is, however, not replicated in PEL possibly because of lack

of power (Supplementary text).

Epistatic effects on *CETP* gene expression

LRLD between variants can suggest the existence of gene-gene interactions, especially if they are

functional variants (29). In order to be under selection, mutations typically need to modulate a

phenotype or an endophenotype, such as gene expression. We have shown previously (6) that

CETP and *Adcy9* interact in mice to modulate several phenotypes, including atherosclerotic lesion

development. To test whether these genes interact in humans, we knocked down (KD) *ADCY9* in

hepatocyte HepG2 cells (Step 3a, Figure 1) and performed RNA sequencing on five KD biological

replicates and five control replicates, to evaluate the impact of decreased *ADCY9* expression on

the transcriptome. We confirmed the KD was successful as *ADCY9* expression is reduced in the

KD replicates (Figure 5a), which represents a drastic drop in expression compared to the whole

transcriptome changes (False Discovery Rate [FDR] = 4.07 x 10^{-14}, Methods). We also observed

that *CETP* expression was increased in *ADCY9-KD* samples compared to controls (Figure 5a), an

increase that is also transcriptome-wide significant (FDR=1.97 x 10^{-7}, ß = 1.257). This increased

expression was validated by qPCR, and western blot also showed increased CETP protein product

(Methods, Figure 5-figure supplement 1a,b, Supplementary text), but its overexpression did not

significatively modulate *CETP* expression (Figure 5-figure supplement 1c). Knocking down or

overexpressing *CETP* did not impact *ADCY9* expression on qPCR (Figure 5-figure supplement 1

d,e). These experiments demonstrate an interaction between *ADCY9* and *CETP* at the gene

expression level and raised the hypothesis that *ADCY9* potentially modulates the expression of

CETP through a genetic effect mediated by rs1967309.

To test for potential interaction effects between rs1967309 and *CETP*, we used RNA-seq data from

diverse projects in humans: the GEUVADIS project (33), the Genotype-Tissue Expression (GTEx

v8) project (34) and CARTaGENE (CaG) (35) (Step 3b, Figure 1). When looking across tissues in

GTEx, *ADCY9* and *CETP* expressions negatively correlate in almost all the tissues (Supplementary

Figure 6, Supplementary text), which is consistent with the effect observed during the *ADCY9-KD*

experiment, showing increased expression of *CETP* expression when *ADCY9* is lowly expressed

(Figure 5a, Figure 5-figure supplement 1a,b). We evaluated the effects of the SNPs on expression

levels of *ADCY9* and *CETP* by modelling both SNPs as continuous variables (additive model)

(Methods). The *CETP* SNP rs158477 was reported as an expression quantitative trait locus (eQTL)

in GTEx v7 and, in our models, shows evidence of being a *cis* eQTL of *CETP* in several other

tissues (Supplementary text), although not reaching genome-wide significance. To test specifically

for an epistatic effect between rs1967309 and rs158477 on *CETP* expression, we included an

interaction term in eQTL models (Methods). We note here that we are testing for association for

this specific pair of SNPs only, and that effects across tissues are not independent, such that we set

our significance threshold at p-value=0.05. This analysis revealed a significant interaction effect

(p-value=0.03, ß= -0.22) between the two SNPs on *CETP* expression in GEUVADIS

lymphoblastoid cell lines (Figure 5b, Supplementary Figure 7a). In rs1967309 AA individuals,

copies of the rs158477 A allele increased *CETP* expression by 0.46 (95% CI 0.26-0.86) on average.

In rs1967309 AG individuals, copies of the rs158477 A allele increased CETP expression by 0.24

(95% CI 0.06-0.43) on average and the effect was null in rs1967309 GG individuals (p-

value$_{GG}$=0.58). This suggests that the effect of rs158477 on *CETP* expression changes depending

on genotypes of rs1967309. The interaction is also significant in several GTEx tissues, most of

which are brain tissues, like hippocampus, hypothalamus and substantia nigra, but also in skin,

although we note that the significance of the interaction depends on the number of PEER factors

included in the model (Supplementary Figure 8). These factors are needed to correct for unknown

biases in the data, but also potentially lead to decreased power to detect interaction effects (36). In

CaG whole blood samples, the interaction effect using additive genetic effect at rs1967309 was

not significant, similarly to results from GTEx in whole blood samples. However, given the larger

size of the dataset, we evaluated a genotypic encoding for the rs1967309 SNP in which the

interaction effect is significant (p-value=0.008, Supplementary Figure 7b) in whole blood,

suggesting that rs1967309 could be modulating rs158477 eQTL effect, in this tissue at least, with

a genotype-specific effect. We highlight that the sample sizes of current transcriptomic resources

do not allow to detect interaction effects at genome-wide significance, however the likelihood of

finding interaction effects between our two SNPs on *CETP* expression in three independent

datasets is unlikely to happen by chance alone, providing evidence for a functional genetic

interaction.

Given the sex-specific results reported above, we stratified our interaction eQTL analyses by sex.

We observed that the interaction effect on *CETP* expression in CaG whole blood samples

(N_{male}=359) is restricted to male individuals, and, despite low power due to smaller sample size in

GEUVADIS, the interaction is also only suggestive in males (Supplementary Figure 7c,d). In

GTEx, most well-powered tissues that showed a significant effect in the sex-combined analyses

also harbor male-specific interactions (Supplementary Figure 9). For instance, GTEx skin male

samples (N_{male}=330) show the most significant male-specific interaction effects, with the

directions of effects replicating the sex-combined result in GEUVADIS (an increase of *CETP*

expression for each rs158477 A allele in rs1967309 AA individuals) albeit with an observable

reversal of the direction in rs1967309 GG individuals (decrease of *CETP* expression with

additional rs158477 A alleles) (Figure 5c, Figure 5-figure supplement 2a). However, significant

effects in females are detected in tissues not previously seen as significant for the interaction in

the sex-combined analysis, in the tibial artery (Figure 5d, Figure 5-figure supplement 2) and the

heart atrial appendage (Supplementary Figure 9). For tissues with evidence of sex-specific effects

in stratified analyses, we also tested the effect of an interaction between sex, rs158477 and

rs1967309 (Methods) on *CETP* expression: the three-way interaction is only significant for tibial

artery (Figure 5-figure supplement 2).

Epistatic effects on phenotypes

The interaction effect of rs1967309 and rs158477 on *CETP* expression in several tissues, found in

multiple independent RNA-seq datasets, coupled with the detection of LRLD between these SNPs

in the Peruvian population suggest that selection may act jointly on these loci, specifically in

Peruvians or Andeans. These populations are well known for their adaptation to life in high

altitude, where the oxygen pressure is lower and where the human body is subjected to hypoxia

(37–40). High altitude hypoxia impacts individuals' health in many ways, such as increased

ventilation, decreased arterial pressure, and alterations of the energy metabolism in cardiac and

skeletal muscle (41,42). To test which phenotype(s) may explain the putative coevolution signal

discovered (Step 4, Figure 1), we investigated the impact of the interaction between rs1967309

and rs158477 on several physiological traits, energy metabolism and cardiovascular outcomes

using the UK Biobank and GTEx cohort (Figure 6-figure supplement 1, Supplementary Table 2).

The UK Biobank has electronic medical records and GTEx has cause of death and variables from

medical questionnaires (34). The interaction term was found to be nominally significant (p-

value<0.05) for forced vital capacity (FVC), forced expiratory volume in 1-second (FEV1) and

whole-body water mass, and suggestive (p-value<0.10) for the basal metabolic rate, all driven by

the effects in females (Figure 6a). For CAD, the interaction is suggestive (p-value<0.10) and, in

this case, driven by males (Figure 6a).

Given this sex-specific result on CAD, the condition targeted by dalcetrapib, we tested the effect

of an interaction between sex, rs158477 and rs1967309 (genotypic encoding, see Methods) on

binary cardiovascular outcomes including myocardial infarction (MI) and CAD. For CAD, we see

a significant three-way interaction effect, meaning that for individuals carrying the AA genotype

at rs1967309, the association between rs158477 and CAD is in the opposite direction in males and

females. In other words, in rs1967309-AA females, having an extra A allele at rs158477, which is

associated with higher *CETP* expression (Figure 5b), has a protective effect on CAD. Conversely,

in rs1967309-AA males, each A allele at rs158477 increases the probability of having an event

(Figure 6a). Little effect is seen in either sex for AG or GG at rs1967309, although the

heterozygotes AG behave differently in females (which further justifies the genotypic encoding of

rs1967309). The beneficial effect of the interaction on CAD thus favors the rs1967309-AA +

rs153477-GG males and the rs1967309-AA + rs153477-AA females, two genotype combinations

which are respectively enriched in a sex-specific manner in the LIMAA cohort (Supplementary

text). Again, observing such a result that concords with the direction of effects in the LRLD sex-

specific finding is noteworthy. A significant interaction between the SNPs is also seen in the GTEx

cohort (p-value=0.004, Figure 6-figure supplement 2, Supplementary text), using questionnaire

phenotypes reporting on MI, but the small number of individuals precludes formally investigating

sex effects.

Among the biomarkers studied (Supplementary Table 2), only lipoprotein(a) [Lp(a)] is suggestive

in males (p-value=0.08) for an interaction between rs1967309 and rs158477, with the same

direction of effect as that for CAD (Figure 6). Again, given the differences observed between the

sexes, we tested the effect of an interaction between sex, rs158477 and rs1967309 (genotypic

coding, Methods) on biomarkers, and only Lp(a) was nominally significant in a three-way

interaction (p-value=0.049). The pattern is similar to the results for CAD, ie. a change in the effect

of rs158477 depending on the genotype of rs1967309 in males, with the effect for AA females in

the opposite direction compared to males (Figure 6b). These concordant results between CAD and

Lp(a) support that the putative interaction effect between the loci under study on phenotypes

involves sex as a modifier.

Discussion

In this study, we used population genetics, transcriptomics and interaction analyses in biobanks to

study the link between *ADCY9* and *CETP*. Our study revealed selective signatures in *ADCY9* and

a significant genotypic association between *ADCY9* and *CETP* in two Peruvian cohorts,

specifically between rs1967309 and rs158477, which was also seen in the Native population of the

Andes. The interaction between the two SNPs was found to be nominally significant for respiratory

and cardiovascular disease outcomes (Figure 6, Figure 6-figure supplement 2). Additionally, a

nominally significant epistatic interaction was seen on *CETP* expression in many tissues, including

the hippocampus and hypothalamus in the brain. Despite brain tissues not displaying the highest

CETP expression levels, CETP that is synthesized and secreted in the brain could play an important

role in the transport and the redistribution of lipids within the central nervous system (43,44) and

has been associated with Alzheimer's disease risk (45,46). These findings reinforce the fact that

the SNPs are likely functionally interacting, but extrapolating on the specific phenotypes under

selection from these results is not straight forward. Identifying the phenotype and environmental

pressures that may have caused the selection signal is complicated by the fact that the UK Biobank

participants, on which the marginally significant associations have been found, do not live in the

same environment as Peruvians. In Andeans from Peru, selection in response to hypoxia at high

altitude was proposed to have effects on the cardiovascular system (21). The hippocampus

functions are perturbed at high altitude (eg. deterioration of memory (47,48)), whereas the

hypothalamus regulates the autonomic nervous system (ANS) and controls the heart and

respiratory rates (49,50), phenotypes which are affected by hypoxia at high altitude (51,52).

Furthermore, high altitude-induced hypoxia (53,54) and cardiovascular system disturbances

(55,56) have been shown to be associated in several studies (57–61), thus potentially sharing

common biological pathways. Therefore, our working hypothesis is that selective pressures on our

genes of interest in Peru are linked to the physiological response to high-altitude, which might be

the environmental driver of coevolution.

The significant interaction effects on *CETP* expression vary between sexes in amplitude and

direction, with most signals driven by male samples, but significant interaction effects observed in

females only, despite sample sizes being consistently lower than for males. Notably, in the tibial

artery and heart atrial appendage, two tissues directly relevant to the cardiovascular system, the

female-specific interaction effect on *CETP* expression is reversed between rs1967309 genotypes

AA and GG, compared to the effects seen in males in skin and brain tissues. Given our *ADCY9-*

KD were done in liver cell lines from male donors, future work to fully understand how rs1967309

and rs158477 interact will focus on additional experiments in cells from both male and female

donors in these relevant tissues. In a previous study, we showed that inhibition of both *Adcy9* and

CETP impacted many phenotypes linked to the ANS in male mice (6), but in the light of our

results, these experiments should be repeated in female mice. The function of ANS is important in

a number of pathophysiological states involving the cardiovascular system, like myocardial

ischemia and cardiac arrhythmias, with significant sex differences reported (62–64).

The interaction effect between the *ADCY9* and *CETP* SNPs on both respiratory and cardiovascular

phenotypes differs between the sexes, with effects on respiratory phenotypes limited to females

(Figure 6a) and cardiovascular disease phenotype associations showing significant three-way sex-

by-SNPs effects (Figure 6). Furthermore, the LRLD signal is present mainly in males (Figure 4),

although the genotype association is also seen in female for a different genotype combination,

suggesting the presence of sex-specific selection. This type of selection is very difficult to detect,

especially on autosomes, with very few empirical examples found to date in the human genome

despite strong theoretical support of their occurrence (65). However, sexual dimorphism in gene

expression between males and females on autosomal genes has been linked to evolutionary

pressures (66–68), possibly with a contribution of epistasis. As the source of selection, we favor

the hypothesis of differential survival over differential ability to reproduce, because the genetic

combination between *ADCY9* and *CETP* has high chances to be broken up by recombination at

each generation. Even in the case where recombination is suppressed in males between these loci,

they would still have equal chances to pass the favored combination to both male and female

offspring, which would not explain the sex-specific LRLD signal. We see an enrichment for the

rs1967309-AA + rs158477-GG in males and rs1967309-AA + rs158477-AA in females, which are

the beneficial combination for CAD in the corresponding sex, possibly pointing to a sexually

antagonistic selection pressure, where the fittest genotype combination depends on the sex.

Such two-gene selection signature, where only males show strong LRLD, can happen if a specific

genotype combination is beneficial in creating males (through differential gamete fitness or in

utero survival, for example) or if survival during adulthood is favored with a specific genotype

combination compared to other genotypes. In the case of age-dependent differential survival, the

genotypic association is expected to be weaker at younger ages, however the LRLD signal between

rs1967309 and rs158477 in the LIMAA cohort did not depend on age neither in males nor in

females (Supplementary text). Since very few individuals were younger than 20 years old, it is

likely that the age range in this cohort is not appropriate to distinguish between the two

possibilities. This age-dependent survival therefore remains to be tested in comparison with

pediatric cohorts of Peruvians: if the LRLD signal is absent in newborns for example, it will

suggest a strong selective pressure acts early in life on boys. To specifically test the in-utero

hypothesis, a cohort of stillborn babies with genetic information could allow to evaluate if the

genotype combination is more frequent in these. Lastly, it may be that the evolutionary pressure is

linked to the sex chromosomes (69,70), and a three-way interaction between *ADCY9*, *CETP* and

Y chromosome haplotypes or mitochondrial haplogroups remains to be explored.

Even though we observed the LRLD signal between rs1967309 and rs158477 in two independent

Peruvian cohorts, reducing the likelihood that our result is a false positive, one limitation is that

the individuals were recruited in the same city (Lima) in both cohorts. However, we show that

both populations are heterogeneous with respect to ancestry (Supplementary Figure 2), suggesting

that they likely represent accurately the Peruvian population. As recent admixture and population

structure can strongly influence LRLD, we performed several analyses to consider these

confounders, in the full cohorts and in the sex-stratified analyses. All analyses were robust to

genome-wide and local ancestry patterns, such that our results are unlikely to be explained by these

effects alone (Supplementary text). Unfortunately, we could not use expression and phenotypic

data from Peruvian individuals, which makes all the links between the selection pressures and the

phenotype associations somewhat indirect. Future studies should focus on evaluating the

phenotypic impact of the interaction specifically in Peruvians individuals, in cohorts such as the

Population Architecture using Genomics and Epidemiology (PAGE) (71), in order to confirm the

marginally significant associations found in European cohorts. Indeed, the Peruvian/Andean

genomic background could be of importance for the interaction effect observed in this population,

which reduces the power of discovery in individuals of unmatched ancestry. Furthermore, not

much is known about the strength of this type of selection, and simulations would help evaluate

how strong selection would need to be in a single generation to produce this level of LRLD.

Another limitation is the low number of samples per tissue in GTEx and the cell composition

heterogeneity per tissue and per sample (72), which can be partially captured by PEER factors and

can modulate the eQTL effects. Therefore, our power to detect tissue-specific interaction effects

is reduced in this dataset, making it quite remarkable that we were able to observe multiple

nominally significant interaction effects between the loci.

Despite these limitations, our results support a functional role for the *ADCY9* intronic SNP

rs1967309, likely involved in a molecular mechanism related to *CETP* expression, but this

mechanism seems to implicate sex as a modulator in a tissue-specific way, which complicates

greatly its understanding. In the dal-OUTCOMES clinical trial, the partial inhibitor of CETP,

dalcetrapib, did not decrease the risk of cardiovascular outcomes in the overall population, but

rs1967309 in the *ADCY9* gene was associated to the response to the drug, which benefitted AA

individuals (4). Interestingly, rs1967309 AA is found in both the male and female beneficial

combinations of genotypes for CAD, the same that are enriched in Peruvians, but without taking

rs158477 and sex into account, this association was masked. The modulation of *CETP* expression

by rs1967309 could impact CETP's functions that are essential for successfully reducing

cardiovascular events. The rs158477 locus could be a key player for these functions, and

dalcetrapib may be mimicking its impact, hence explaining the pharmacogenomics association.

Furthermore, in the light of our results, some of these effects could differ between men and women

(73), which may need to be taken into consideration in the future precision medicine interventions

potentially implemented for dalcetrapib.

In conclusion, we discovered a putative epistatic interaction between the pharmacogene *ADCY9*

and the drug target gene *CETP*, that appears to be under selection in the Peruvian population. Our

approach exemplifies the potential of using evolutionary analyses to help find relationships

between pharmacogenes and their drug targets. We characterized the impact of the *ADCY9/CETP*

interaction on a range of phenotypes and tissues. Our gene expression results in brain tissues

suggest that the interaction could play a role in protection against challenges to the nervous system

caused by stress such as hypoxia. The female-specific eQTL interaction results in arteries and heart

tissues further suggest a link with the cardiovascular system, and the phenotype association results

support further this hypothesis. In light of the associations between high altitude-induced hypoxia

and cardiovascular system changes, the interaction identified in this study could be involved in

both systems: for example, ADCY9 and CETP could act in pathways involved in adaptation to

high altitude, which could influence cardiovascular risk via their interaction in a sex-specific

manner. Finally, our findings of an evolutionary relationship between *ADCY9* and *CETP* during

recent human evolution points towards a biological link between dalcetrapib's pharmacogene

ADCY9 and its therapeutic target *CETP*.

Material and Methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
gene (Homo Sapiens)	CETP	GenBank	HGNC:1869	

gene (Homo Sapiens)	ADCY9	GenBank	HGNC:240	
cell line (Homo Sapiens)	HepG2	ATCC	HB-8065	Hepatocellular carcinome
recombinant DNA reagent	pEZ-M46-AC9 plasmid	GeneCopoeia™	EX-H0609-M46	Methods section
recombinant DNA reagent	pEZ-M50-CETP plasmid	GeneCopoeia™	EX-C0070-M50	Methods section
antibody	anti-CETP (rabbit monoclonal)	Abcam	#ab157183	(1:1000) in 3% BSA, TBS, tween 20 0.5%, O/N 4°C
antibody	Goat anti-rabbit antibody (goat polyclonal)	Abcam	#ab6721	(1:10 000) in 3% BSA 1h at room temperature
sequence-based reagent	Human CETP_F	IDT Technologies	PCR primers	CTACCTGTCTTTCCATAA
sequence-based reagent	Human CETP_R	IDT Technologies	PCR primers	CATGATGTTAGAGATGAC
sequence-based reagent	Human ADCY9_F	IDT Technologies	PCR primers	CTGAGGTTCAAGAACATCC
sequence-based reagent	Human ADCY9_R	IDT Technologies	PCR primers	TGATTAATGGGCGGCTTA
sequence-based reagent	Silencer Select siRNA against human ADCY9	Ambion	Cat. #4390826 ID 1039	CCUGAUGAAAGAUUACUUUtt
sequence-based reagent	Silencer Select siRNA against human CETP	Ambion	Cat. #4392420 ID 2933	GGACAGAUCUGCAAAGAGAtt
sequence-based reagent	Negative Control siRNA	Ambion	Cat. #4390844	
commercial assay or kit	Lipofectamine RNAiMAX reagent	Invitrogen	Cat. #13778	
commercial assay or kit	Lipofectamine 2000 reagent	Invitrogen	Cat. #11668-019	
commercial assay or kit	RNeasy Plus Mini Kit	Qiagen	Cat. #74136	
commercial assay or kit	High-Capacity cDNA Reverse Transcription Kit	Applied Biosystems	Cat. #4368814	
commercial assay or kit	Agilent RNA 6000 Nano Kit for	Agilent Technologies	Cat. #5067-1511	

	Bioanalyzer 2100 System			
commercial assay or kit	SYBR-Green reaction mix	BioRad	Cat. #1725274	
commercial assay or kit	Amicon Ultra 0.5 ml 10 kDa cutoff units	Millipore Sigma	Cat. #UFC501096	
commercial assay or kit	Western Lightning ECL Pro	Perkin Elmer	Cat. #NEL122001 EA	
commercial assay or kit	TGX Stain-Free FastCast Acrylamide 10%	BioRad	Cat# 1610183	
software, algorithm	TrimGalore!	DOI : 10.14806/ej.17.1.200		
software, algorithm	STAR (v.2.6.1a)	DOI : 10.1093/bioinformatics/bts635		
software, algorithm	RSEM (v.1.3.1)	DOI : 10.1186/1471-2105-12-323		
software, algorithm	R statistical software (v.3.6.0/v.3.6.1)	https://www.r-project.org/		
software, algorithm	FlashPCA2	DOI : 10.1093/bioinformatics/btx299		
software, algorithm	Vcftools (v.0.1.17)	DOI : 10.1093/bioinformatics/btr330		
software, algorithm	RFMix (v.2.03)	DOI : 10.1016/j.ajhg.2013.06.020		
software, algorithm	PEER	DOI : 10.1038/nprot.2011.457		
software, algorithm	pyGenClean (v.1.8.3)	DOI : 10.1093/bioinformatics/btt261		
software, algorithm	SAS (v.9.4)	https://www.sas.com/en_us/software/stat.html		
software, algorithm	EPO pipeline (version e59)	DOI : 10.1093/database/bav096		
software, algorithm	Bcftools (v.1.9)	DOI : 10.1093/bioinformatics/btr509		
software, algorithm	GenotypeHarmonizer (v.1.4.20)	DOI : 10.1186/1756-0500-7-901		
software, algorithm	Hapbin (v.1.3.0)	DOI : 10.1093/molbev/msv172		

software, algorithm	SHAPEIT2 (r.837)	DOI : 10.1038/nmeth.1785		
software, algorithm	PBWT	DOI : 10.1093/bioinformatics/btu014		
software, algorithm	Beacon designer software (v.8) (Premier Biosoft)	http://www.premierbiosoft.com/qOligo/Oligo.jsp?PID=1		
Other	1000 Genomes project	DOI : 10.1038/nature15393		
Other	LIMAA	DOI : 10.1038/s41467-019-11664-1	dbGAP :phs002025.v1.p1	dbgap project #26882
Other	Native American genetic dataset	DOI : 10.1038/nature11258		
Other	GEUVADIS	DOI : 10.1038/nature12531		
Other	GTEx (v8)	DOI : 10.1038/ng.2653	dbGAP : phs000424.v8.p2	dbgap project #19088
Other	CARTaGENE biobank	DOI : 10.1093/ije/dys160		CAG project number 406713
Other	UK biobank	DOI : 10.1371/journal.pmed.1001779		UKB project #15357 and #20168
Other	Sanger Imputation Server	DOI : 10.3389/fgene.2019.00034		

Population Genetics Datasets

The whole-genome sequencing data from the 1000 Genomes project (1000G) Phase III dataset (ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/release/20130502/) was filtered to exclude INDELs and CNVs so that we kept only biallelic SNPs. This database has genomic variants of 2,504 individuals across five ancestral populations: Africans (AFR, $n = 661$), Europeans (EUR, $n = 503$), East Asians (EAS, $n = 504$), South Asians (SAS, $n = 489$) and Americans (AMR, $n = 347$) (74). The replication dataset, LIMAA, has been previously published (31,32) and was accessed through dbGaP [phs002025.v1.p1, dbgap project #26882]. This cohort was genotyped with a customized Affymetric LIMAAray containing markers optimized for Peruvian-specific rare and coding

variants. We excluded related individuals as reported previously (31), resulting in a final dataset

of 3,509 Peruvians. We also identified fine-scale population structure in this cohort and a more

homogeneous subsample of 3,243 individuals (1,302 females and 1,941 males) in this cohort was

kept for analysis (Table 1, Supplementary text). The Native American genetic dataset (NAGD)

contains 2,351 individuals from Native descendants from the data from a previously published

study (23). Individuals were separated by their linguistic families identified by Reich and

colleagues (23). NAGD came under the Hg18 coordinates, so a lift over was performed to transfer

to the Hg19 genome coordinates. Pre-processing details for these datasets are described in

Supplementary text.

eQTL Datasets

We used several datasets (Table 1) for which we had both RNA-seq data and genotyping. First,

the GEUVADIS dataset (33) for 1000G individuals was used (available at

https://www.internationalgenome.org/data-portal/data-collection/geuvadis). A total of 287 non-

duplicated European samples (CEU, GBR, FIN, TSI) were kept for analysis. Second, the

Genotype-Tissue Expression v8 (GTEx) (34) was accessed through dbGaP (phs000424.v8.p2,

dbgap project #19088) and contains gene expression across 54 tissues and 948 donors, genetic and

phenotypic information. Phenotype analyses are described in Supplementary text. The cohort

contains mainly of European descent (84.6%), aged between 20 and 79 years old. Analyses were

done on 699 individuals, 66% of males and 34% of females (Supplementary Figure 10a). Third,

we used the data from the CARTaGENE biobank (35) (CAG project number 406713) which

includes 728 RNA-seq whole-blood samples with genotype data, from individuals from Quebec

(Canada) aged between 36 and 72 years old (Supplementary Figure 10b). Genotyping and RNA-

seq data processing pipelines for these datasets are detailed in Supplementary text. To quantify

ADCY9 gene expression, we removed the isoform transcript ENST00000574721.1 (ADCY9-205

from the Hg38) from the Gene Transfer Format (GTF) file because it is a "retained intron" and

accumulates genomic noise (Supplementary text), masking true signals for *ADCY9*. To take into

account hidden factors, we calculated PEER factors (75) on the normalized expressions, on all

samples and stratified by sex (sPEER factors). To detect eQTL effects, we performed a two-sided

linear regression on *ADCY9* and *CETP* expressions using R (v.3.6.0) (https://www.r-project.org/)

with the formula $lm(p \sim rs1967309 * rs158477 + Covariates)$ for evaluating the interaction

effect, $lm(p \sim rs1967309 + rs158477 + Covariates)$ for the main effect of the SNPs and

$lm(p \sim rs1967309 * rs158477 * sex + Covariates)$ for evaluating the three-way interaction

effect. Under the additive model, each SNP is coded by the number of non-reference alleles (G for

rs1967309 and A for rs158477), under the genotypic model, dummy coding is used with

homozygous reference genotype set as reference. The covariates include the first 5 Principal

Components (PCs), age (except for GEUVADIS, information not available), sex, as well as PEER

factors. We tested the robustness of our results to the inclusion of different numbers of PEER

factors in the models and we report them all for GEUVADIS, CARTaGENE and GTEx

(Supplementary Figure 7-9). Reported values in the text are for five PEER factors in GEUVADIS,

ten PEER factors in CARTaGENE, 25 sPEER for skin sun exposed in male and 10 sPEER for

artery tibial in female in GTEx. Covariates specific to each cohort are reported in Supplementary

text.

UK biobank processing and selected phenotypes

The UK biobank (76) contains 487,392 genotyped individuals from the UK still enrolled as of

August 20th 2020, imputed using the Haplotype Reference Consortium as the main reference panel,

and accessed through project #15357 and UKB project #20168. Additional genetic quality control

was done using pyGenClean (v.1.8.3) (77). Variants or individuals with more than 2% missing

genotypes were filtered out. Individuals with discrepancies between the self-reported and genetic

sex or with aneuploidies were removed from the analysis. We considered only individuals of

European ancestry based on PCs, as it is the largest population in the UK Biobank, and because

ancestry can be a confounder of the genetic effect on phenotypes. We used the PCs from UK

Biobank to define a region in PC space using individuals identified as "white British ancestry" as

a reference population. Using the kinship estimates from the UK Biobank, we randomly removed

individuals from kinship pairs where the coefficient was higher than 0.0884 (corresponding to a

3rd degree relationship). The resulting post QC dataset included 413,138 individuals. For the

reported phenotypes, the date of baseline visit was between 2006 and 2010. The latest available

hospitalization records discharge date was June 30th 2020 and the latest date in the death registries

was February 14th 2018. We used algorithmically-defined cardiovascular outcomes based on

combinations of operation procedure codes (OPCS) and hospitalization or death record codes

(ICD9/ICD10). A description of the tested continuous variables can be found in Supplementary

Table 2. We used age at recruitment defined in variable #21022 and sex in variable #31. We

ignored self-reported events for cardiovascular outcomes as preliminary analyses suggested they

were less precise than hospitalization and death records.

In association models, each SNP analyzed is coded by the number of non-reference alleles, G for

rs1967309 and A for rs158477. SNP rs1967309 was also coded as a genotypic variable, to allow

for non-additive effects. For continuous traits (Supplementary Table 2) in the UK Biobank, general

two-sided linear models (GLM) were performed using SAS software (v.9.4). A GLM model was

first performed using the covariates age, sex and PCs 1 to 10. The externally studentized residuals

were used to determine the outliers, which were removed. The normality assumption was

confirmed by visual inspection of residuals for most of the outcomes, except *birthwt* and *sleep*.

For biomarkers and cardiovascular endpoints, regression analyses were done in R (v.3.6.1). Linear

regression analyses were conducted on standardized outcomes and logistic regression was used for

cardiovascular outcomes. Marginal effects were calculated using margins package in R. In both

cases, models were adjusted for age at baseline and top 10 PCs, as well as sex when not stratified.

In models assessing two-way (rs1967309 by rs158477) or three-way (rs1967309 by rs158477 by

sex) interactions, we used a 2 d.f. likelihood ratio test for the genotypic dummy variables'

interaction terms (genotypic model) (Supplementary text).

RNA-sequencing of ADCY9-knocked-down HepG2 cell line

The human liver hepatocellular HepG2 cell line was obtained from ATCC, a cell line derived from

the liver tissue of a 15-year-old male donor (78). Cells were cultured in EMEM Minimum essential

Medium Eagle's, supplemented with 10% fetal bovine serum (Wisent Inc). 250 000 cells in 2 ml

of medium in a six-well plate were transfected using 12.5 pmol of Silencer Select siRNA against

human ADCY9 (Ambion cat # 4390826 ID 1039), Silencer Select siRNA against CETP (Ambion

cat 4392420 ID 2933) or Negative Control siRNA (Ambion cat #4390844) with 5 μl of

Lipofectamine RNAiMAX reagent (Invitrogen cat #13778) in 500 μl Opti-MEM I reduced serum

medium (Invitrogen cat # 31985) for 72h (Supplementary table 3, Supplementary text). The

experiment was repeated five times at different cell culture passages. Total RNA was extracted

from transfected HepG2 cells using RNeasy Plus Mini Kit (Qiagen cat #74136) in accordance with

the manufacturer's recommendation. Preparation of sequencing library and sequencing was

performed at the McGill University Innovation Center. Briefly, ribosomal RNA was depleted

using NEBNext rRNA depletion kit. Sequencing was performed using Illumina NovaSeq 6000 S2

paired end 100 bp sequencing lanes. Basic QC analysis of the 10 samples was performed by the

Canadian Centre for Computational Genomics (C3G). To process the RNA-seq samples, we first

performed read trimming and quality clipping using TrimGalore! (79)

(https://github.com/FelixKrueger/TrimGalore), we aligned the trimmed reads on the Hg38

reference genome using STAR (v.2.6.1a) and we ran RSEM (v.1.3.1) on the transcriptome aligned

libraries. Prior to normalization with limma and voom, we filtered out genes which had less than

6 reads in more than 5 samples. For *ADCY9* and *CETP* gene-level differential expression analyses,

we compared the mean of each group of replicates with a t-test for paired samples. The

transcriptome-wide differential expression analysis was done using limma, on all genes having an

average of at least 10 reads across samples from a condition. Samples were paired in the

experiment design. The multiple testing was taken into account by correcting the p-values with the

qvalue R package (v.4.0.0) (80), to obtain transcriptome-wide FDR values.

Overexpression of *ADCY9* and *CETP* genes in HepG2 cell line

For *ADCY9* and *CETP* overexpression experiments, 500 000 cells in 2 ml of medium in a six-well

plate were transfected using 1 ug of pEZ-M46-AC9 or pEZ-M50-CETP plasmids

(GeneCopoeia^{TM}) with 5 ul of Lipofectamine 2000 reagent (Invitrogen cat # 11668-019) for 72h.

Total RNA was extracted from transfected HepG2 cells using RNeasy Plus Mini Kit (Qiagen cat

#74136) in accordance with the manufacturer's recommendation (Supplementary table 3,

Supplementary text).

Natural selection analyses

We used the integrated Haplotype Statistic (iHS) (22) and the population branch statistic (PBS) (74) to look for selective signatures. The iHS values were computed for the each 1000G population. An absolute value of iHS above 2 is considered to be a genome wide significant signal (22). Prior to iHS computation, ancestral alleles were retrieved from 6 primates using the EPO pipeline (version e59) (81) and the filtered 1000 Genomes vcf files were converted to change the reference allele as ancestral allele using bcftools (82) with the fixref plugin. The hapbin program (v.1.3.0) (83) was then used to compute iHS using per population-specific genetic maps computed by Adam Auton on the 1000G OMNI dataset (ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/technical/working/20130507_omni_recombination_rates). When the genetic map was not available for a subpopulation, the genetic map from the closest sub-population was selected according to their global FST value computed on the phase 3 dataset. We scanned the *ADCY9* and *CETP* genes using the population branch statistic (PBS), using 1000G sub-populations data. PBS summarizes a three-way comparison of allele frequencies between two closely related populations, and an outgroup. The grouping we focused on was PEL/MXL/CHB, with PEL being the focal population to test if allele frequencies are especially differentiated from those in the other populations. The CHB population was chosen as an outgroup to represent a Eurasian population that share common ancestors in the past with the American populations, after the out-of-Africa event. Using PJL (South Asia) and CEU (Europe) as an outgroup, or CLM as a closely related population (instead of MXL) yield highly similar results. To calculate F_{ST} for each pair of population in our tree, we used vcftools (84) by subpopulation. We calculated normalized PBS values as in (21), which adjusts values for positions with large branches in all populations,

for the whole genome. We use this distribution to define an empirical threshold for significance

based on the 95th percentile of all PBS values genome-wide for each of the three populations.

Long-range linkage disequilibrium

Long-range linkage disequilibrium (LRLD) was calculated using the function geno-r2 of vcftools

(v.0.1.17) which uses the genotype frequencies. LRLD was evaluated in all subpopulations from

1000 Genomes Project Phase III, in LIMAA and NAGD, for all biallelic SNPs in *ADCY9*

(chr16:4,012,650-4,166,186 in Hg19 genome reference) and *CETP* (chr16:56,995,835-57,017,756

in Hg19 genome reference). We analyzed loci from the phased VCF files that had a MAF of at

least 5% and a missing genotype of at most 10%, in order to retain a maximum of SNPs in NAGD

which has higher missing rates than the others. We extracted the 99th percentile of all pairs of

comparisons between *ADCY9* and *CETP* genes to use as a threshold for empirical significance and

we refer to these as *ADCY9/CETP* empirical p-values. In LIMAA, we also evaluated the genotypic

association using a χ^2 test with four degrees of freedom (χ^2_4) using a permutation test, as reported

in (25) (Supplementary text).

Furthermore, for both cohorts, we created a distribution of LRLD values for random pairs of SNPs

across the genome to obtain a genome-wide null distribution of LRLD to evaluate how unusual

the genotypic association between our candidate SNPs (rs1967309-rs158477) is while taking into

account the cohort-specific background genomic noise/admixture and allele frequencies. We

extracted 3,513 pairs of SNPs that match rs1967309 and rs158477 in terms of MAF, physical

distance (in base pairs) and genetic distance (in centiMorgans (cM), based on the PEL genetic

map) between them in both cohorts (Supplementary text), and report genome-wide empirical p-

values based on this distribution. For the analyses of LRLD between *ADCY9* and *CETP* stratified

by sex, we considered the same set of SNP pairs that we used for the full cohorts, but separated

the dataset by sex before calculating the LRLD values. To evaluate how likely the differences

observed in LRLD between sex are, we also performed permutations of the sex labels across

individuals to create a null distribution of sex specific effects (Supplementary text).

Local ancestry inference

To evaluate local ancestry in the PEL subpopulation and in the LIMAA cohort, we constructed a

reference panel using the phased haplotypes from 1000 Genomes (YRI, CEU, CHB) and the

phased haplotypes of NAGD (Northern American, Central American and Andean)

(Supplementary text). We kept overlapping positions between all datasets, and when SNPs had the

exact same genetic position, we kept the SNP with the highest variance in allele frequencies across

all reference populations (Supplementary text). We ran RFMix (v.2.03) (85) (with the option

'reanalyze-reference' and for 25 iterations) on all phased chromosomes. We estimated the whole

genome average proportion of each ancestry using a weighted mean of the chromosome specific

proportions given by RFMix based on the chromosome size in cM. For comparing the overall

Andean enrichment inferred by RFMix between rs1967309/rs158477 genotype categories, we

used a two-sided Wilcoxon-t-test. To evaluate the Andean local ancestry enrichment specifically

at *ADCY9* and *CETP*, we computed the genome-wide 95th percentile for proportion of Andean

attribution for all intervals given by RFMix.

Code and source data

Numerical summary data represented as a graph in main figures, as well as the code to reproduce

figures   and   analyses,   can   be   found   here:

https://github.com/HussinLab/adcy9_cetp_Gamache_2021. Raw RNA sequencing data for

knocked down experiments in hepatocyte HepG2 cells are deposited the data on NCBI Gene

Expression Omnibus, accession number GSE174640.

Main manuscript

Table 1

Figures 1-6

Figure 3-figure supplement 1-2

Figure 4-figure supplement 1-3

Figure 5-figure supplement 1-2

Figure 6-figure supplement 1-2

Supplemental Information description

Supplementary figures 1-10

Supplementary tables 1-3

Supplementary methods and texts

Acknowledgements

We thank all members of the Hussin lab for their constructive comments and feedback throughout

this project, as well as the insightful input from three reviewers and reviewing and senior editors

at eLife. This work was completed thanks to computational resources provided by Compute

Canada clusters Graham and Beluga. This work was funded by the *Institut de Valorisation des*

Données (IVADO), Health Collaboration Acceleration Fund from the Ministère de l'Économie et

de l'Innovation of the Government of Quebec and the Montreal Heart Institute (MHI) Foundation.

J.G.H. is a *Fonds de la Recherche en Santé* (FRQS) Junior 1 fellow. I.G. receives a PhD

scholarship from the MHI Foundation and M.A.L. holds a PhD scholarship from Canadian

Institutes of Health Research. M.P.D. holds the Canada Research Chair in Precision Medicine Data

Analysis. J.C.T. holds the Canada Research Chair in Personalized Medicine and the Université de

Montréal endowed research chair in atherosclerosis.

Author Contributions

Conceptualization: I.G., M.P.D and J.G.H.; Data curation: I.G., M.A.L., J.C.G.; Statistical and

bioinformatic analyses: I.G., M.A.L., J.C.G., H.T, S.A, A.B. and Y.F.Z.; Data acquisition: J.C.G.,

J.G.H, Y.L., L.L., M.M. and S.R.; Wet lab experimentation: R.S. and E.R.; Writing – Original

draft: I.G. and J.G.H.; Writing – Review & editing: I.G., M.A.L., J.C.G., R.S., E.R., S.A., H.T.,

Y.L., S.R., J.C.T., M.P.D. and J.G.H.; Supervision: J.C.T., M.P.D. and J.G.H.; Funding

acquisition: J.C.T., M.P.D., J.G.H.

Competing Interests statement

J.G.H. has received speaker honoraria from Dalcor and District 3 Innovation Centre. J.C.T. reports

grants from Government of Quebec, National Heart, Lung, and Blood Institute of the U.S. National

Institutes of Health (NIH), the MHI Foundation, from Bill and Melinda Gates Foundation, Amarin,

Esperion, Ionis, Servier, RegenXBio; personal fees from Astra Zeneca, Sanofi, Servier; and

personal fees and minor equity interest from Dalcor. M.P.D. and J.C.T. have a patent Methods for

Treating or Preventing Cardiovascular Disorders and Lowering Risk of Cardiovascular Events

issued to Dalcor, no royalties received, a patent Genetic Markers for Predicting Responsiveness to

Therapy with HDL-Raising or HDL Mimicking Agent issued to Dalcor, no royalties received, and

a patent Methods for using low dose colchicine after myocardial infarction with royalties paid to

Invention assigned to the Montreal Heart Institute. M.P.D. reports personal fees and other from

Dalcor and personal fees from GlaxoSmithKline, other from AstraZeneca, Pfizer, Servier, Sanofi.

The remaining authors have nothing to disclose.

References

1.   Lagrost L. Regulation of cholesteryl ester transfer protein (CETP) activity: review of in vitro and in vivo studies. Biochim Biophys Acta. 1994 Dec 8;1215(3):209–36.

2.   Shinkai H. Cholesteryl ester transfer-protein modulator and inhibitors and their potential for the treatment of cardiovascular diseases. Vasc Health Risk Manag. 2012;8:323–31.

3.   Schwartz GG, Olsson AG, Abt M, Ballantyne CM, Barter PJ, Brumm J, et al. Effects of dalcetrapib in patients with a recent acute coronary syndrome. N Engl J Med. 2012 Nov 29;367(22):2089–99.

4.   Tardif J-C, Rhéaume E, Lemieux Perreault L-P, Grégoire JC, Feroz Zada Y, Asselin G, et al. Pharmacogenomic determinants of the cardiovascular effects of dalcetrapib. Circ Cardiovasc Genet. 2015 Apr;8(2):372–82.

5.   Tardif J-C, Rhainds D, Brodeur M, Feroz Zada Y, Fouodjio R, Provost S, et al. Genotype-dependent effects of dalcetrapib on cholesterol efflux and inflammation. Circ Cardiovasc Genet. 2016 Aug;9(4):340–8.

6.   Rautureau Y, Deschambault V, Higgins M-È, Rivas D, Mecteau M, Geoffroy P, et al. ADCY9 (Adenylate Cyclase Type 9) inactivation protects from atherosclerosis only in the absence of CETP (Cholesteryl Ester Transfer Protein). Circulation. 2018 Oct 16;138(16):1677–92.

7.   Bersaglieri T, Sabeti PC, Patterson N, Vanderploeg T, Schaffner SF, Drake JA, et al. Genetic signatures of strong recent positive selection at the lactase gene. Am J Hum Genet. 2004 Jun;74(6):1111–20.

8.   Enattah NS, Trudeau A, Pimenoff V, Maiuri L, Auricchio S, Greco L, et al. Evidence of still-ongoing convergence evolution of the lactase persistence T-13910 alleles in humans. Am J Hum Genet. 2007 Sep;81(3):615–25.

9.   Gamba C, Jones ER, Teasdale MD, McLaughlin RL, Gonzalez-Fortes G, Mattiangeli V, et al. Genome flux and stasis in a five millennium transect of European prehistory. Nature Communications. 2014 Oct 21;5(1):5257.

10.   Itan Y, Powell A, Beaumont MA, Burger J, Thomas MG. The origins of lactase persistence in Europe. PLOS Computational Biology. 2009 août;5(8):e1000491.

11.   Poulter M, Hollox E, Harvey CB, Mulcare C, Peuhkuri K, Kajander K, et al. The causal element for the lactase persistence/non-persistence polymorphism is located in a 1 Mb region of linkage disequilibrium in Europeans. Ann Hum Genet. 2003 Jul;67(Pt 4):298–311.

12.   Labrie V, Buske OJ, Oh E, Jeremian R, Ptak C, Gasiūnas G, et al. Lactase non-persistence is directed by DNA variation-dependent epigenetic aging. Nat Struct Mol Biol. 2016 Jun;23(6):566–73.

13.   Yi X, Liang Y, Huerta-Sanchez E, Jin X, Cuo ZXP, Pool JE, et al. Sequencing of fifty human exomes reveals adaptation to high altitude. Science. 2010 Jul 2;329(5987):75–8.

14.   Fumagalli M, Moltke I, Grarup N, Racimo F, Bjerregaard P, Jørgensen ME, et al. Greenlandic Inuit show genetic signatures of diet and climate adaptation. Science. 2015 Sep 18;349(6254):1343.

15.   Hollenbach JA, Thomson G, Cao K, Fernandez-Vina M, Erlich HA, Bugawan TL, et al. HLA diversity, differentiation, and haplotype evolution in Mesoamerican Natives. Human Immunology. 2001 Apr 1;62(4):378–90.

16.   Li P, Zhao J, Kothapalli KSD, Li X, Li H, Han Y, et al. A regulatory insertion-deletion polymorphism in the FADS gene cluster influences PUFA and lipid profiles among Chinese

adults: a population-based study. The American Journal of Clinical Nutrition. 2018 Jun
1;107(6):867–75.
17.   Reynolds LM, Dutta R, Seeds MC, Lake KN, Hallmark B, Mathias RA, et al. FADS
genetic and metabolomic analyses identify the Δ5 desaturase (FADS1) step as a critical control
point in the formation of biologically important lipids. Scientific Reports. 2020 Sep
28;10(1):15873.
18.   Tashi T, Scott Reading N, Wuren T, Zhang X, Moore LG, Hu H, et al. Gain-of-function
EGLN1 prolyl hydroxylase (PHD2 D4E:C127S) in combination with EPAS1 (HIF-2α)
polymorphism lowers hemoglobin concentration in Tibetan highlanders. J Mol Med. 2017 Jun
1;95(6):665–70.
19.   Meyer D, C. Aguiar VR, Bitarello BD, C. Brandt DY, Nunes K. A genomic perspective
on HLA evolution. Immunogenetics. 2018 Jan 1;70(1):5–27.
20.   Blais M-E, Zhang Y, Rostron T, Griffin H, Taylor S, Xu KY, et al. High frequency of
HIV mutations associated with HLA-C suggests enhanced HLA-C-restricted CTL selective
pressure associated with an AIDS-protective polymorphism. J Immunol. 2012 May
1;188(9):4663–70.
21.   Crawford JE, Amaru R, Song J, Julian CG, Racimo F, Cheng JY, et al. Natural selection
on genes related to cardiovascular health in high-altitude adapted Andeans. Am J Hum Genet.
2017 Nov 2;101(5):752–67.
22.   Voight BF, Kudaravalli S, Wen X, Pritchard JK. A map of recent positive selection in the
human genome. PLOS Biology. 2006 Mar 7;4(3):e72.
23.   Reich D, Patterson N, Campbell D, Tandon A, Mazieres S, Ray N, et al. Reconstructing
Native American population history. Nature. 2012 Aug;488(7411):370–4.
24.   Lewontin RC, Kojima K. The evolutionary dynamics of complex polymorphisms.
Evolution. 1960;14(4):458–72.
25.   Rohlfs RV, Swanson WJ, Weir BS. Detecting coevolution through allelic association
between physically unlinked loci. The American Journal of Human Genetics. 2010 May
14;86(5):674–85.
26.   Harris DN, Song W, Shetty AC, Levano KS, Cáceres O, Padilla C, et al. Evolutionary
genomic dynamics of Peruvians before, during, and after the Inca Empire. Proc Natl Acad Sci U
S A. 2018 Jul 10;115(28):E6526–35.
27.   Li W-H, Nei M. Stable linkage disequilibrium without epistasis in subdivided
populations. Theoretical Population Biology. 1974 Oct 1;6(2):173–83.
28.   Nei M, Li WH. Linkage disequilibrium in subdivided populations. Genetics. 1973
Sep;75(1):213–9.
29.   Park L. Population-specific long-range linkage disequilibrium in the human genome and
its influence on identifying common disease variants. Scientific Reports. 2019 Aug 6;9(1):11380.
30.   Slatkin M. Linkage disequilibrium — understanding the evolutionary past and mapping
the medical future. Nat Rev Genet. 2008 Jun;9(6):477–85.
31.   Asgari S, Luo Y, Akbari A, Belbin GM, Li X, Harris DN, et al. A positively selected
FBN1 missense variant reduces height in Peruvian individuals. Nature. 2020 Jun;582(7811):234–
9.
32.   Luo Y, Suliman S, Asgari S, Amariuta T, Baglaenko Y, Martínez-Bonet M, et al. Early
progression to active tuberculosis is a highly heritable trait driven by 3q23 in Peruvians. Nat
Commun. 2019 Aug 21;10(1):3765.
33.   Lappalainen T, Sammeth M, Friedländer MR, 't Hoen PAC, Monlong J, Rivas MA, et al.

Transcriptome and genome sequencing uncovers functional variation in humans. Nature. 2013
Sep;501(7468):506–11.

34.   The Genotype-Tissue Expression (GTEx) project. Nat Genet. 2013 Jun;45(6):580–5.

35.   Awadalla P, Boileau C, Payette Y, Idaghdour Y, Goulet J-P, Knoppers B, et al. Cohort
profile of the CARTaGENE study: Quebec's population-based biobank for public health and
personalized genomics. Int J Epidemiol. 2013 Oct;42(5):1285–99.

36.   Brynedal B, Choi J, Raj T, Bjornson R, Stranger BE, Neale BM, et al. Large-scale trans-
eQTLs affect hundreds of transcripts and mediate patterns of transcriptional co-regulation. Am J
Hum Genet. 2017 Apr 6;100(4):581–91.

37.   Beall CM. Two routes to functional adaptation: Tibetan and Andean high-altitude
natives. Proc Natl Acad Sci U S A. 2007 May 15;104(Suppl 1):8655–60.

38.   Brutsaert TD, Parra EJ, Shriver MD, Gamboa A, Rivera-Ch M, León-Velarde F.
Ancestry explains the blunted ventilatory response to sustained hypoxia and lower exercise
ventilation of Quechua altitude natives. Am J Physiol Regul Integr Comp Physiol. 2005
Jul;289(1):R225-234.

39.   Julian CG, Moore LG. Human genetic adaptation to high altitude: evidence from the
Andes. Genes (Basel). 2019 février;10(2):150.

40.   Moore LG. Human genetic adaptation to high altitudes: current status and future
prospects. Quat Int. 2017 Dec 15;461:4–13.

41.   Milledge JS, West JB, Schoene RB. High altitude medicine and physiology [Internet]. 4th
ed. CRC Press; 2007. Available from: https://books.google.ca/books?id=lj59BgAAQBAJ

42.   Murray AJ. Energy metabolism and the high-altitude environment. Experimental
Physiology. 2016;101(1):23–7.

43.   Albers JJ, Tollefson JH, Wolfbauer G, Albright RE. Cholesteryl ester transfer protein in
human brain. Int J Clin Lab Res. 1992;21(3):264–6.

44.   Yamada T, Kawata M, Arai H, Fukasawa M, Inoue K, Sato T. Astroglial localization of
cholesteryl ester transfer protein in normal and Alzheimer's disease brain tissues. Acta
Neuropathol. 1995 Dec 1;90(6):633–6.

45.   Murphy EA, Roddey JC, McEvoy LK, Holland D, Hagler DJ, Dale AM, et al. CETP
polymorphisms associate with brain structure, atrophy rate, and Alzheimer's disease risk in an
APOE-dependent manner. Brain Imaging Behav. 2012 Mar;6(1):16–26.

46.   Oestereich F, Yousefpour N, Yang E, Ribeiro-da-Silva A, Chaurand P, Munter LM. The
cholesteryl ester transfer protein (CETP) raises cholesterol levels in the brain and affects
presenilin-mediated gene regulation. bioRxiv. 2020 Nov 24;2020.11.24.395186.

47.   Lieberman P, Morey A, Hochstadt J, Larson M, Mather S. Mount Everest: a space
analogue for speech monitoring of cognitive deficits and stress. Aviat Space Environ Med. 2005
Jun;76(6 Suppl):B198-207.

48.   Shukitt-Hale B, Stillman MJ, Welch DI, Levy A, Devine JA, Lieberman HR. Hypobaric
hypoxia impairs spatial memory in an elevation-dependent fashion. Behavioral and Neural
Biology. 1994 Nov 1;62(3):244–52.

49.   Horiuchi J, McDowall LM, Dampney RAL. Vasomotor and respiratory responses evoked
from the dorsolateral periaqueductal grey are mediated by the dorsomedial hypothalamus. The
Journal of Physiology. 2009;587(21):5149–62.

50.   Rahmouni K. Cardiovascular regulation by the arcuate nucleus of the hypothalamus:
neurocircuitry and signaling systems. Hypertension. 2016 Jun;67(6):1064–71.

51.   Bärtsch Peter, Gibbs J. Simon R. Effect of altitude on the heart and the lungs. Circulation.

2007 Nov 6;116(19):2191–202.

52.   Hainsworth R, Drinkhill MJ, Rivera-Chira M. The autonomic nervous system at high
altitude. Clin Auton Res. 2007 Feb;17(1):13–9.

53.   Bigham AW, Lee FS. Human high-altitude adaptation: forward genetics meets the HIF
pathway. Genes Dev. 2014 Oct 15;28(20):2189–204.

54.   Moore LG. Measuring high-altitude adaptation. J Appl Physiol (1985). 2017 Nov
1;123(5):1371–85.

55.   Abe H, Semba H, Takeda N. The roles of hypoxia signaling in the pathogenesis of
cardiovascular diseases. J Atheroscler Thromb. 2017 Sep 1;24(9):884–94.

56.   Lee JW, Ko J, Ju C, Eltzschig HK. Hypoxia signaling in human diseases and therapeutic
targets. Exp Mol Med. 2019 Jun 20;51(6):1–13.

57.   Faeh D, Gutzwiller F, Bopp M, Swiss National Cohort Study Group. Lower mortality
from coronary heart disease and stroke at higher altitudes in Switzerland. Circulation. 2009 Aug
11;120(6):495–501.

58.   Naeije R. Physiological adaptation of the cardiovascular system to high altitude. Progress
in Cardiovascular Diseases. 2010 May 1;52(6):456–66.

59.   Ostadal B, Kolar F. Cardiac adaptation to chronic high-altitude hypoxia: Beneficial and
adverse effects. Respiratory Physiology & Neurobiology. 2007 Sep 30;158(2):224–36.

60.   Riley CJ, Gavin M. Physiological changes to the cardiovascular system at high altitude
and its effects on cardiovascular disease. High Altitude Medicine & Biology. 2017 Mar
15;18(2):102–13.

61.   Savla JJ, Levine BD, Sadek HA. The effect of hypoxia on cardiovascular disease: friend
or foe? High Alt Med Biol. 2018 Jun;19(2):124–30.

62.   Abhishekh HA, Nisarga P, Kisan R, Meghana A, Chandran S, Trichur Raju, et al.
Influence of age and gender on autonomic regulation of heart. J Clin Monit Comput. 2013 Jun
1;27(3):259–64.

63.   Dart AM, Du X-J, Kingwell BA. Gender, sex hormones and autonomic nervous control
of the cardiovascular system. Cardiovascular Research. 2002 Feb 15;53(3):678–87.

64.   Nugent AC, Bain EE, Thayer JF, Sollers JJ, Drevets WC. Sex differences in the neural
correlates of autonomic arousal: a pilot PET study. Int J Psychophysiol. 2011 Jun;80(3):182–91.

65.   Morrow EH, Connallon T. Implications of sex-specific selection for the genetic basis of
disease. Evol Appl. 2013 Dec;6(8):1208–17.

66.   Connallon T, Clark AG. Sex linkage, sex-specific selection, and the role of
recombination in the evolution of sexually dimorphic gene expression. Evolution. 2010
Dec;64(12):3417–42.

67.   Parsch J, Ellegren H. The evolutionary causes and consequences of sex-biased gene
expression. Nature Reviews Genetics. 2013 Feb;14(2):83–7.

68.   Williams TM, Carroll SB. Genetic and molecular insights into the development and
evolution of sexual dimorphism. Nature Reviews Genetics. 2009 Nov;10(11):797–804.

69.   Cox RM, Cox CL, McGlothlin JW, Card DC, Andrew AL, Castoe TA. Hormonally
mediated increases in sex-biased gene expression accompany the breakdown of between-sex
genetic correlations in a sexually dimorphic lizard. The American Naturalist. 2017 Jan
25;189(3):315–32.

70.   McGlothlin JW, Cox RM, Brodie ED III. Sex-Specific Selection and the Evolution of
Between-Sex Genetic Covariance. Journal of Heredity. 2019 Jul 1;110(4):422–32.

71.   Wojcik GL, Graff M, Nishimura KK, Tao R, Haessler J, Gignoux CR, et al. Genetic

analyses of diverse populations improves discovery for complex traits. Nature. 2019
Jun;570(7762):514–8.
72.   Aguet F, Brown AA, Castel SE, Davis JR, He Y, Jo B, et al. Genetic effects on gene
expression across human tissues. Nature. 2017 Oct;550(7675):204–13.
73.   Metzinger Mark P., Saldanha Suzanne, Gulati Jaskeerat, Patel Kershaw V., El-Ghazali
Ayea, Deodhar Sneha, et al. Effect of anacetrapib on cholesterol efflux capacity: a substudy of
the DEFINE trial. Journal of the American Heart Association. 2020 Dec 15;9(24):e018136.
74.   Auton A, Abecasis GR, Altshuler DM, Durbin RM, Abecasis GR, Bentley DR, et al. A
global reference for human genetic variation. Nature. 2015 Oct;526(7571):68–74.
75.   Stegle O, Parts L, Piipari M, Winn J, Durbin R. Using probabilistic estimation of
expression residuals (PEER) to obtain increased power and interpretability of gene expression
analyses. Nat Protoc. 2012 Feb 16;7(3):500–7.
76.   Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, et al. UK biobank: an open
access resource for identifying the causes of a wide range of complex diseases of middle and old
age. PLOS Medicine. 2015 Mar 31;12(3):e1001779.
77.   Lemieux Perreault L-P, Provost S, Legault M-A, Barhdadi A, Dubé M-P. pyGenClean:
efficient tool for genetic data clean up before association testing. Bioinformatics. 2013 Jul
1;29(13):1704–5.
78.   López-Terrada D, Cheung SW, Finegold MJ, Knowles BB. Hep G2 is a hepatoblastoma-
derived cell line. Hum Pathol. 2009 Oct;40(10):1512–5.
79.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal. 2011 May 2;17(1):10–2.
80.   Storey JD. A direct approach to false discovery rates. Journal of the Royal Statistical
Society: Series B (Statistical Methodology). 2002;64(3):479–98.
81.   Herrero J, Muffato M, Beal K, Fitzgerald S, Gordon L, Pignatelli M, et al. Ensembl
comparative genomics resources. Database (Oxford) [Internet]. 2016 Feb 20 [cited 2021 Feb
23];2016. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4761110/
82.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence
alignment/map format and SAMtools. Bioinformatics. 2009 Aug 15;25(16):2078–9.
83.   Maclean CA, Chue Hong NP, Prendergast JGD. hapbin: an efficient program for
performing haplotype-based scans for positive selection in large genomic datasets. Molecular
Biology and Evolution. 2015 Nov 1;32(11):3027–9.
84.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant
call format and VCFtools. Bioinformatics. 2011 Aug 1;27(15):2156–8.
85.   Maples BK, Gravel S, Kenny EE, Bustamante CD. RFMix: a discriminative modeling
approach for rapid and robust local-ancestry inference. Am J Hum Genet. 2013 Aug
8;93(2):278–88.
933
934

935 **Table 1. Cohort information.** Sample sizes are reported after quality control steps.

Cohort/Subpopulation	Abbreviation	Ethnicity	Sample size (% female)	Age	Reference
1000G – Peruvian	PEL*	Peruvian	85 (52%)	NA	(74)
LIMAA/Peruvian	LIMAA	Peruvian	3,243 (40%)	29.6 ± 13.8	(31,32)
Native Amerind/Andean	NAGD/AND	Amerind/Peruvian	88 (40%)	NA	(23)
GEUVADIS	GEUVADIS*	European descent	287 (54%)	NA	(33)
CARTaGENE	CaG	European descent	728 (51%)	53.6 ± 8.7	(35)
GTEx	GTEx	European descent	699 (34%)	52.6 ± 13.1	(34)
UK biobank	Ukb*	European descent	413,138 (54%)	56.8 ± 8.0	(76)

936 *indicates a discovery cohort
937 NA: not available
938

Main figures

941
942
Figure 1. Flowchart of experimental design and main results. The four main steps of the
analyses conducted in this study are reported along with the datasets used for each step and the
genetic loci on which the analyses are performed. Green colored boxes represent analyses for
which sex is considered.
KD = Knock-down
OX = Overexpression

Figure 2. Natural selection signature at rs1967309 in *ADCY9*. (a) Genotype frequency distribution of rs1967309 in populations from the 1000 Genomes (1000G) Project and in Native Americans. (b) Significant iHS values (absolute values above 2) for 1000G continental populations and recombination rates from AMR-1000G population-specific genetic maps, in the *ADCY9* gene. (c) PBS values in the *ADCY9* gene, in CHB (outgroup, left panel), PEL (middle panel) and MXL (right panel). Horizontal lines represent the 95th percentile PBS value genome-wide for each population. Vertical dotted black lines define the LD block around rs1967309 (black circle) from 1000G population-specific genetic maps. Gene plots for *ADCY9* showing location of its exons are presented in blue below each plot. Abbreviations: Altaic from Mongolia and Russia: ALT; Uralic Yukaghir from Russia: URY; Chukchi Kamchatkan from Russia: CHK; Northern American from Canada, Guatemala and Mexico: NOA; Central American from Costal Rica and Mexico: CEA; Chibchan Paezan from Argentina, Bolivia, Colombia, Costa Rica and Mexico: CHP; Equatorial Tucanoan from Argentina, Brazil, Colombia, Gualana and Paraguay: EQT; Andean from Bolivia, Chile, Colombia and Peru: AND. For 1000G populations, abbreviations can be found here https://www.internationalgenome.org/category/population/.

965
966

Figure 3. Long-range linkage disequilibrium between rs1967309 and rs158477 in Peruvians from Lima, Peru. (a) Genotype correlation (r^2) between rs1967309 and all SNPs with MAF>5% in *CETP*, for the PEL population. (b) Genotype correlation between the 3 loci identified in (a) to be in the 99th percentile and all SNPs with MAF>5% in *ADCY9*. The dotted line indicates the position of rs1967309. The horizontal lines in (a,b) represent the threshold for the 99th percentile of all comparisons of SNPs (MAF>5%) between *ADCY9* and *CETP*. Figure 3-figure supplement 1 presents the same plots for Andeans and in the replication cohort (LIMAA) and Figure 3-figure supplement 2 compares the r^2 values between PEL and LIMAA (c) Genotype frequency distribution of rs158477 in 1000G and Native American populations. (d) Genomic distribution of r^2 values from 3,513 pairs of SNPs separated by between 50-60 Mb and 61±10 cM away across all Peruvian chromosomes from the PEL sample, compared to the rs1967309-rs158477 r^2 value (dotted grey line) (genome-wide empirical p-value=0.01). The vertical black line shows the threshold for the 95th percentile threshold of all pairs. Gene plots showing location of exons for *CETP* (a) and *ADCY9* (b) are presented in blue below each plot. Abbreviations: Altaic from Mongolia and Russia: ALT; Uralic Yukaghir from Russia: URY; Chukchi Kamchatkan from Russia: CHK; Northern American from Canada, Guatemala and Mexico: NOA; Central American from Costal Rica and Mexico: CEA; Chibchan Paezan from Argentina, Bolivia, Colombia, Costa Rica and Mexico: CHP; Equatorial Tucanoan from Argentina, Brazil, Colombia, Gualana and Paraguay: EQT; Andean from Bolivia, Chile, Colombia and Peru: AND. For 1000G populations, abbreviations can be found here https://www.internationalgenome.org/category/population/

987
988
Figure 3-figure supplement 1. Long-range linkage disequilibrium in the Andean population
from the Native Population (n=88) (a,b) and in the LIMAA cohort (n=3243) (c,d). (a,c)
Genotype correlation (r^2) between rs1967309 and all SNPs with MAF>5% in *CETP*. (b,d)
Genotype correlation between the 3 loci identified in Figure 3a to be in the 99[th] percentile and all
SNPs with MAF>5% in *ADCY9*. The dotted line indicates the position of rs1967309. The
horizontal lines represent the threshold for the 99[th] percentile of all comparisons of SNPs
(MAF>5%) between *ADCY9* and *CETP*.

996
Figure 3-figure supplement 2. Comparison of genotype correlation between Peruvian from
1000G and from the LIMAA cohort. Comparison of genotype correlation (r^2) between all SNPs
in *ADCY9* and *CETP* with MAF>5% in the Peruvian population (PEL) in 1000G (x axis) and
LIMAA cohort (y axis). Colored dots represent the value for SNPs higher than the 99th percentile
with rs1967309 in PEL identified in Figure 3a. Black lines represent the 99th percentile in both
populations.

1003
1004
Figure 4. Sex-specific long-range linkage disequilibrium. Genotype correlation between the loci
identified in *CETP* in Figure 3a and all SNPs with MAF>5% in *ADCY9* for (a,b) the PEL
population and (c,d) LIMAA cohort in males (a,c) and in females (b,d). Genotype frequencies per
sex are shown in Figure 4-figure supplement 1 and sex-specific PBS values in Figure 4-figure
supplement 2. The horizontal line shows the threshold for the 99th percentile of all comparisons of
SNPs (MAF>5%) between *ADCY9* and *CETP*. The vertical dotted line represents the position of
rs1967309. Blue dots represent the rs158477 SNPs and pink represents the other three SNPs
identified in Figure 3a (rs158480, rs158617 and rs12447620), which are in near-perfect LD. Figure
4-figure supplement 3 shows the same analysis in Andeans from NAGD. Gene plots for *ADCY9*
showing location of its exons are presented in blue below each plot.

1015
1016
Figure 4-figure supplement 1. Genotype frequency distribution per sex. Genotype frequency
distribution of rs1967309 in *ADCY9* (a,b) and rs158477 in *CETP* (c,d) in populations from the
1000 Genomes (1000G) Project, in Native Americans (NAGD) and LIMAA cohorts, in females
(a,c) and males (b,d). Abbreviations: Altaic from Mongolia and Russia: ALT; Uralic Yukaghir
from Russia: URY; Chukchi Kamchatkan from Russia: CHK; Northern American from Canada,
Guatemala and Mexico: NOA; Central American from Costal Rica and Mexico: CEA; Chibchan
Paezan from Argentina, Bolivia, Colombia, Costa Rica and Mexico: CHP; Equatorial Tucanoan
from Argentina, Brazil, Colombia, Gualana and Paraguay: EQT; Andean from Bolivia, Chile,
Colombia and Peru: AND. For 1000G populations, abbreviations can be found here
https://www.internationalgenome.org/category/population/.

1027
1028
1029 **Figure 4-figure supplement 2. PBS values in the *ADCY9* per sex, comparing the CHB**
1030 **(outgroup), MXL and PEL.** Horizontal lines represent the 95th percentile PBS value of the
1031 chromosome 16 for each population for each sex. Vertical black lines represent the LD block
1032 around rs1967309 (shown as a black circle for PEL). Gene plots for *ADCY9* showing location of
1033 its exons are presented in blue below each plot.

1034
1035
Figure 4-figure supplement 3. Sex-specific long-range linkage disequilibrium in the Andean
population (NAGD). Genotype correlation between the loci identified in CETP in Figure 3a and
all SNPs with MAF>5% in *ADCY9* for the Andean population, in males (N=54) and in females
(N=34). The horizontal line shows the threshold for the 95th percentile of all comparisons of SNPs
(MAF>5%) between *ADCY9* and *CETP*. The vertical dotted line represents the position of
rs1967309.

1042
Figure 5. Effect of *ADCY9* on *CETP* expression. (a) Normalized expression of *ADCY9* or *CETP*
genes depending on wild type (WT) and *ADCY9-KD* in HepG2 cells from RNA sequencing on
five biological replicates in each group. P-values were obtained from a two-sided Wilcoxon paired
test. qPCR and western blot results in HepG2 are presented in Figure 5-figure supplement 1. (b,c,d)
CETP expression depending on the combination of rs1967309 and rs158477 genotypes in (b)
GEUVADIS (p-value=0.03, ß= -0.22, N=287), (c) GTEx-Skin Sun Exposed in males (p-
value=0.0017, ß= -0.32, N=330) and in (d) GTEx-Tibial artery in females (p-value=0.026, ß= 0.38,
N=156), for individuals of European descent according to principal component analysis. P-values
reported were obtained from a two-way interaction of a linear regression model for the maximum
number of PEER/sPEER factors considered. Figure 5-figure supplement 2 show the interaction p-
values depending on number of PEER/sPEER factors included in the linear models.

1054
1055
Figure 5-figure supplement 1. *ADCY9*/CETP **interaction in HepG2 cells.** (a) Relative mRNA expression of *CETP* of HepG2 cells 72h post-transfection with siRNA against human *ADCY9* (si1039). qPCR assay was normalized with PGK1 and HBS1L genes, n= 5 independent experiments, (p-value=0.0026 from t-test). (b) Quantification of CETP protein by Western blot assay, 200 ml of cell media (concentrated with Amicon ultra 0.5 ml 10 kDA units) from cells transfected with siRNA against human ADCY9 (si1039), were separated on 10% TGX-acrylamide gel and transferred to PVDF membrane. CETP protein expression was determined using a primary antibody rabbit monoclonal anti-CETP (Abcam, ab157183) 1:1000 (3% BSA, TBS, Tween 20 0.5%) O/N 4°C, followed by HRP-conjugated secondary antibody goat anti-rabbit 1:10 000 (3% BSA) 1h RT. Figure b represents densitometry analysis of n=3 experiments, p-value=0.0029 from t-test. (c,e) Relative mRNA expression of (c) *CETP* and (e) *ADCY9* genes in HepG2 cells post-transfection with pEZ-M50-CETP (overexpression of *CETP*) or pEZ-M46-ADCY9 (overexpression of *ADCY9*) plasmids. qPCR assay was normalized with PGK1 and HBS1L genes, n=4 independent experiments. (d) Relative mRNA expression of *ADCY9* of HepG2 cells 72h post-transfection with siRNA against human *CETP*. qPCR assay was normalized with PGK1 and HBS1L genes, n= 4 independent experiments.

1084
Figure 6. Epistatic association of rs1967309 and rs158477 on phenotypes in the UK biobank.
(a) Significance of the interaction effect between rs1967309 and rs158477 on several physiological traits, energy metabolism and cardiovascular outcomes overall and stratified by sex in the UK biobank. Horizontal lines represent the p-value thresholds at 0.05 (plain) and 0.10 (dotted). Single-SNP p-values are shown in Figure 6-figure supplement 1. (b,c) Sex-stratified effects of rs158477 on (b) cardiovascular phenotypes and (c) biomarkers depending on the genotype of rs1967309 (genotypic encoding). The p-values p_{itx} reported come from a likelihood ratio test comparing models with and without the three-way interaction term between the two SNPs and sex. Sex-combined results using GTEx cardiovascular phenotype data are shown in Figure 6-figure supplement 2. See Supplementary Table 2 for the list of abbreviations.

Phenotypes

1095
1096 **Figure 6-figure supplement 1. Single SNP effects of rs1967309 and rs158477 on phenotypes**
1097 **in the UK biobank.** Significance of the marginal effect of rs1967309 and rs158477, both codded
1098 as additive, on several physiological traits, energy metabolism and cardiovascular outcomes,
1099 overall and stratified by sex in the UK biobank. The dotted line represents the p-value at 0.05. See
1100 Supplementary Table 2 for the list of abbreviations.

1101

Figure 6-figure supplement 2. Epistatic association of rs1967309 and rs158477 on cardiovascular disease in GTEx. (a) Effect of the rs158477 SNP on the cardiovascular phenotype (n=693, cas=120, control=563) depending on the genotype of rs1967309 in GTEx. For both models, rs158477 was codded as additive (GG=0, GA=1, AA=2). For the additive model, rs1967309 was codded as additive (AA=0, AG=1, GG=2). P-value of the interaction (p_{itx}) was obtained using a linear regression in R. For the genotypic model, rs1967309 was codded as a genotypic variable and p-values were obtained from a likelihood ratio test comparing models with and without the interaction term between the SNPs. (b) Proportion of cases for each genotype combinations between rs1967309 and rs158477. The numerator indicates the number of cases and the denominator the number total of individuals (cases+controls). Darker colors show higher proportions of cases.