3   Do host-consumed resources increase endoparasitic but decrease ectoparasitic infections?

Brin Shayhorn[1][*], Chloe Ramsay[2][*][a], Kristi Medina[1], Erin Sauer[3], Jason R. Rohr[2]

[1]University of South Florida, Department of Integrative Biology, Tampa, FL 33620, USA

[2]University of Notre Dame, Department of Biological Sciences, Notre Dame, IN 46556, USA

[3]University of Wisconsin-Madison, Department of Forest and Wildlife Ecology, Madison, WI

53706, USA

*Co-first authorship

[a]-Corresponding Author: Chloe Ramsay, Department of Biological Sciences, 100 Galvin Life

Science Center, University of Notre Dame, IN, 46556; Email: cramsay@nd.edu

Abstract

Habitat loss and disease outbreak play a major role in the decline of biodiversity. Habitat

degradation is often associated with reduced food resources, which can lead to less investment in

host immunity and increased infections. However, pathogens use host resources for replication

and pathogen traits, such as infecting hosts internally or short generation times, might allow

pathogens to rapidly capitalize on host-consumed nutrients. Thus, it is unclear whether increased

food consumption by hosts should reduce or amplify pathogen levels. conducted experiments on

Cuban treefrogs (*Osteopilus septentrionalis*) to test how food availability affects infection levels

of Ranavirus and the fungal pathogen *Batrachochytrium dendrobatidis* (Bd), which are both

associated with mass die-offs of amphibians. Given that Ranavirus is an endoparasite with a

much shorter generation time than the ectoparasitic Bd, we postulated that Ranavirus might be

able to capitalize on host-consumed resources more quickly than Bd. We hypothesized that

increased food availability to hosts might reduce Bd infections more than Ranavirus infections.

As predicted, augmenting food access decreased Bd infection intensity, but increased Ranavirus

infection intensity. Future work should assess whether pathogen traits, such as generation time

and endo- versus ectoparasitism, generally affect whether food resources more positively benefit

hosts or pathogens.

Keywords: Ranavirus, *Batrachochytrium dendrobatidis,* resource, pathogen traits, amphibian

declines, habitat loss

Introduction

Habitat loss and degradation can reduce food resources for wildlife, which can exacerbate

and cause disease (Daszak et al. 1999, Gibbons et al. 2000). This is because mounting an

immune response requires resources and therefore hosts short on resources are unlikely to mount

a completely effective immune response, which could increase disease risk and severity (Gray et

al. 2009, Echaubard et al. 2010). As an example, food restricted bumblebees (*Bombus terrestris*)

experienced a 50% increase in mortality from *Crithidia bombi* infection (protist) when compared

to bees fed *ad libitum* (Brown et al. 2000).

While hosts use resources to combat pathogens, pathogens also utilize host resources.

Thus, positive relationships between host resources and pathogen loads are theoretically possible

(Cressler et al. 2014a). For example, increased resources can increase pathogen density in the

environment and consequently pathogen load in hosts (Johnson et al. 2007, Civitello et al. 2015,

Civitello et al. 2018). Additionally, hosts with more resources are often healthier and larger and

thus might be more capable of tolerating higher infection loads (Pulkkinen and Ebert 2004,

Knutie et al. 2017). Finally, high resource levels can be used by some pathogens to increase their

replication, despite the potential for a subsequent increase in host immune response (Johnson et

al. 2007, Seppälä et al. 2008, Hall et al. 2009, Cressler et al. 2014a, Civitello et al. 2018).

Although there have been in-depth examinations of how host traits and resources affect

whether the host or parasite benefits more from increased resource availability (Pulkkinen and

Ebert 2004, de Roode et al. 2008, Seppälä et al. 2008, Hall et al. 2009, Cressler et al. 2014b, a,

Hite and Cressler 2019), surprisingly, the role of parasite traits on resource-dependent parasitism

remains under tested. Certain parasite traits may facilitate pathogens utilizing host resources.

For example, quickly replicating pathogens could potentially capitalize on high resource

environments faster than slowly replicating pathogens (Arsnoe et al. 2011, Hall et al. 2012), and

perhaps even faster than the host can use these resources to mount an immune response. Also,

pathogens with more direct access to host-consumed resources, such as endoparasites in the

gastrointestinal tract of the host, may be able to access these resources more easily than those

without direct access, such as ectoparasites.

In recent decades, amphibian populations have experienced global declines, partially due

to land use changes that have altered resource availability and disease outbreaks (Gibbons et al.

2000). Two pathogens contributing to population declines in amphibians are *Batrachochytrium*

dendrobatidis (Bd) and Ranavirus (Daszak et al. 1999). Bd is a widespread and pathogenic

fungus that attacks amphibian epithelial tissues, preventing osmotic regulation through the skin

and causing mortality by cardiac arrest (Voyles et al. 2009). It is an ectoparasite, residing on the

skin of amphibians with a ~ 4d replication rate at room temperature (Piotrowski et al. 2004,

Voyles et al. 2009). Ranavirus is a pathogen that results in hemorrhaging and cell death of major

organs (Gray et al. 2009). Ranavirus resides internally and has a comparatively high replication

rate, inducing host death in as few as two days for tadpoles and seven days for adults of

susceptible species (Green et al. 2002, Sutton et al. 2014).

Studies examining how host susceptibility to Bd and Ranavirus are affected by resource

availability have reported mixed results. One study found that increased resources reduced host

susceptibility to Bd (Venesky et al. 2012) and the rest found no effects of resource availability

(Reeve et al. 2013, Cothran et al. 2015, Buck et al. 2016). However, all these Bd studies focused

on nitrogen, phosphorus, carotenoids, or protein levels, addressing food quality rather than

quantity. Also, all these studies were conducted on tadpoles rather than adults, despite adults

being the most susceptible amphibian life stage to Bd (Venesky et al. 2012, Cothran et al. 2015,

McMahon and Rohr 2015, Buck et al. 2016). A study on how resource availability affected

Ranaviral infections in tadpoles, the life stage when hosts are most susceptible to Ranavirus

(Green et al. 2002, Gray et al. 2009), found that resources did not affect host susceptibility to

Ranavirus (Reeve et al. 2013). However, this study only limited host access to resources before

exposing hosts to Ranavirus (Reeve et al. 2013). Here, we examine how host-consumed

resources affect Ranavirus and Bd growth on amphibians to gather evidence in support of or

against the hypothesis that rapidly replicating endoparasites (i.e., Ranavirus) and slowly

replicating ectoparasites (i.e., Bd) benefit more than and less than hosts, respectively, from host-

consumed nutrients.

103   **Methods**

Animal Husbandry

Adult and larval Cuban tree frogs (*Osteopilus septentrionalis*) were collected from the

University of South Florida Botanical Gardens (Tampa, Florida, USA) in November 2015 and

August 2017, respectively. Both life stages were housed in 946 mL sterilized plastic deli cups,

with adults placed on unbleached paper towels wet with artificial spring water (ASW) and larvae

placed in 800 mL of ASW. All individuals were kept at 20^{o}C with a 12hr photoperiod. Adults

and tadpoles were used in Bd and Ranavirus experiments, respectively, because these life stages

are the most susceptible to these pathogens (Gray et al. 2009, McMahon and Rohr 2015).

Experimental Design

Bd experiment: Eighteen adult Cuban tree frogs of equal mass were assigned randomly to

the four treatments (2x2 experiment) for a total of 72 individually housed frogs. All individuals

then received 1 mL of deionized water pipetted on to their dorsal surface that was from rinsed

Petri dishes (15 x 150 mm; 1% agar, 1% tryptone medium) that either were (Bd treatment; $n =$

36) or were not (control; $n = 36$) growing Bd (SRS 812 isolate). Each Bd exposed frog received

10^5 zoospores (Gervasi et al. 2013). After Bd or sham exposure, individuals were placed on

either an *ad libitum* diet, where they were provided with an excess of small house crickets

(*Acheta domestica*; ~0.27g) three times per week or a restricted diet, where they were provided

~0.094g of crickets three times per week.

123   *Ranavirus experiment:* Ten Cuban tree frog tadpoles were assigned randomly to the six

treatments groups (2 x 3 experiment) for a total of sixty tadpoles individually housed in deli cups

with 100mL of ASW. Ranavirus (FV3-like) was cultured in fathead minnow (*Pimephales*

promelas) cells and maintained at -80°C in minimal essential medium (MEM). Tadpoles

received either 77µL (10^5 PFUs) of Ranavirus (FV3-like) in MEM or 77µL of MEM-without

virus as a sham treatment. Previous studies found that doses between 10^2-10^6 PFUs cause

sublethal effects and morbidity in tadpoles (Hoverman et al. 2010). After twenty-four hours,

700mL of ASW was added to each cup for a total of 800 mL/tadpole. Each tadpole was then

provided either one, two, or three 0.6 x 1-cm cylindrical plugs (measured using the first

centimeter of a disposable micropipette tip) of a mix of fish flakes and spirulina (1:1) in 1% agar,

according to their randomly assigned low, medium, or high food treatment.

General Methods

136   All amphibians were weighed or staged (if tadpoles; Gosner 1960) at the beginning of the

experiment and every week thereafter. Clean containers were provided weekly. To quantify

infections, each adult frog was swabbed weekly on their abdomen and hind limbs. The mouth

and cloaca of tadpoles were swabbed at the end of the experiment. All swabs were placed in 1mL

microcentrifuge tubes and stored at -20°C for later processing. After the experiment concluded,

DNA was extracted from the swabs and quantitative polymerase chain reaction (qPCR) was used

to quantify Bd load (Boyle et al. 2004) and Ranavirus load (Picco et al. 2007). Any individuals

surviving to the end of the experiment (three weeks for Bd and two weeks for Ranavirus) were

euthanized with a 0.1% solution of buffered MS-222.

For the Bd experiment, blood from euthanized frogs was extracted using microcapillary

tubes to collect plasma. We then conducted enzyme-linked immunosorbent assays (ELISA) on

the plasma to quantify IgY antibody levels (adapted from Knutie et al. 2017 ; see Supplemental

Methods for details).

Statistical Analysis

All analyses were conducted in R (3.4.2, 2017) and figures were created using the *visreg*

package and *visreg* function (Breheny and Burchett 2019). We used general linear models to

examine the effect of resource availability, parasite exposure, and their interaction on host

growth, development (Gosner stage; Ranavirus experiment only), and immune response (IgY

antibody levels in the Bd experiment only). Host growth was measured as the difference between

pre-exposure and 3-week post-exposure body mass (g) for Bd and as change in mass per week

for Ranavirus.

We also examined the effect of resource availability and Ranavirus exposure on survival

by conducting a survival analysis using the *survival* package and the *coxph* function (Therneau

and Lumley 2019). Individuals that survived to the end of the experiment were right-censored.

No amphibians died in the Bd experiment and consequently survival analyses were not run.

To assess how Bd loads changed over time with resource availability, we analyzed Bd

infection intensity using a generalized linear mixed effects model with a negative binomial error

distribution. Resource availability and weeks since the beginning of infection were used as

interacting independent variables, and individual was included as a random effect. To assess how

Ranavirus intensity changed with resource availability, we used a similar analysis, but without

the inclusion of week since infection or the random effect in the model because intensity was not

measured through time. To assess how resource level affected pathogen prevalence, we ran two

generalized linear regression models (one for each pathogen) with binomial error distributions.

The binomial response variable was host infection status (infected or not) and resource level was

used as a predictor variable. Generalized linear mixed models were run using the *pscl* package

and *glm.nb* function (Jackman et al. 2020). Tukey post-hoc tests were run to test which resource

levels differed from one another using the *multcomp* package and *glht* function (Hothorn 2010).

175   **Results**

176   In both the Bd and Ranavirus experiments, hosts fed more had higher growth rates

177   (g/week; Ranavirus: $F_{2, 51}$=5.72, p<0.006, Fig. 1A; Bd: β=0.208, χ^2_1= 58.196, $p < 0.001$, Fig.

178   1B). However, neither Ranavirus exposure (β= -2.97, $F_{1, 51}$=0.016, p=0.89) nor Bd exposure (β=-

179   0.013, χ^2_1= 0.478, $p = 0.490$) affected host growth rate.

180   Host development rate, as measured by Gosner stages per week, was significantly

affected by an interaction between Ranavirus exposure and resource availability ($F_{2,51}$=7.89,

p=0.001). Ranavirus exposure decreased host development rate for hosts in the low resource

group ($F_{1,17}$=10.92, p=0.004), but did not significantly affect development rate in the medium or

high resource groups (medium resources: $F_{1,17}=1.32$, $p=0.27$; high resources: $F_{1,17}=0.02$, $p=0.89$;

Fig. 1C).

Mortality was not significantly affected by resource availability ($\chi^2_2=0.0003$, $p=0.999$),

Ranavirus exposure ($\beta=2.028$, $\chi^2_1=0.33$, $p=0.567$), or the interaction between the two ($\chi^2_2=3.94$,

$p=0.14$) in the Ranavirus experiment. One individual died in the control-low resource treatment

group and one in the Ranavirus-low and Ranavirus-high resource treatment groups for a total of

three mortalities. There was no mortality in the Bd experiment. Ranavirus intensity was 4.6x

higher in tadpoles assigned to the high resource than the low resource group ($\chi^2_1=4.50$, $p=0.03$;

Fig 2A). In contrast, Bd intensities were 2.5x higher in frogs fed the low resource level than the

high resource level ($\chi^2_1=4.27$, $p<0.05$; Fig 2A), but there were no significant effects of time since

infection ($\chi^2_1=079$, $p=0.78$) or an interaction between resource availability and time ($\chi^2_1=0.51$,

$p=0.47$). Surviving tadpoles provided the low resource treatment had 89% Ranavirus prevalence,

whereas tadpoles in the medium and high resource groups had 70% and 44% Ranavirus

prevalence, respectively ($\chi^2_1=3.57$, $p=0.059$; Fig. 2B). Bd prevalence did not significantly differ

between the high (88%) and low (80%) resource groups ($\chi^2_1=0.50$, $p=0.48$). Additionally, host

IgY was not affected by resource availability ($\beta=-0.12$, $F_{1,32}=2.34$, $p=0.14$), Bd exposure

($\beta=0.059$, $F_{1,32}=0.083$, $p=0.78$), or their interaction ($F_{1,32}=0.20$, $p=0.66$).

202   **Discussion**

Hosts use resources to combat infections and limited resources may force tradeoffs

between immunity and growth or reproduction (Lochmiller and Deerenberg 2000, Rauw 2012).

Pathogens also use host resources making it unclear whether host-consumed resources should be

beneficial or costly to the pathogen. Resource quantity or quality and host traits might determine

whether increased resources are more beneficial for the host or pathogen (Cressler et al. 2014a).

In this experiment, we assessed the level of evidence in support of the hypothesis that pathogen

traits can alter competition for resources between host and pathogen. We found that resource

level was positively associated with pathogen load for Ranavirus, a rapidly replicating

endoparasite, but negatively associated with pathogen load for Bd, a more slowly replicating

ectoparasite (Fig. 2).

In both experiments, increasing food availability increased frog growth, regardless of

whether the frogs were exposed to pathogens or not (Fig. 1A, B), suggesting that hosts in the

high resource groups had sufficient resources to combat pathogens. However, neither Ranavirus

nor Bd infections affected host growth rate (g/week) or survival, regardless of resource

availability. In contrast to growth rates, when hosts were exposed to Ranavirus and had limited

resources, hosts had significantly decreased development rates, whereas Ranavirus exposure did

not alter development rates when hosts were in the medium or high resource treatment groups

(Fig. 1C). Amphibian larvae might increase development to escape a resource-limited

environment, as has been seen with other inhospitable conditions, such as with predators (Relyea

2007), pesticides (Rohr et al. 2004), and in rapidly drying freshwater environments (Bekhet et al.

2014).

While higher resource levels did not alter IgY levels, they did decrease Bd intensity in

hosts, suggesting that these increases in resources allowed hosts to more effectively combat Bd

(Fig. 2A). Increased resources could have led to increases in immunity other than IgY, such as

increases in antimicrobial peptides used to combat Bd, or increases in non-immunological

defenses, such as increased skin shedding rates, which have been shown to decrease Bd intensity

(Rollins-Smith et al. 2002, Cramp et al. 2014). Our results suggest that any increases in Bd

replication from increased resources was not sufficient to counteract subsequent increases in

immunity, leading to the highest Bd loads when the host had the least food. Bd is an ectoparasite

and has a ~4 day replication rate, which could make it challenging for Bd to use host-consumed

resources effectively (Piotrowski et al. 2004). This pattern of increased food resources

decreasing Bd load was shown in a past study on larval southern leopard frogs (Venesky et al.

2012). However, other studies found that the quality of resources do not alter Bd loads in three

different species of larval frogs (Cothran et al. 2015, Buck et al. 2016). Unlike these previous

studies, our study tested resource quantity and not quality, which could account for these

differences. Additionally, amphibian immunity in the larval stage is less complex than in the

adult life stage; thus, how hosts use resources to combat Bd in the larval life stage might be

different than in the adult life stage that we tested (Rollins-Smith 1998).

In contrast to Bd, abundant food increased Ranaviral intensity (Fig. 2B) while decreasing

Ranavirus prevalence. Ranavirus grows best in hematopoietic tissues (Chinchar 2002),

suggesting that Ranavirus may benefit from the increased growth we found in hosts with access

to more resources. Also, Ranavirus replicates rapidly and is an endoparasite, which may allow

this pathogen to more effectively use host resources for replication than some more slowly

replicating ectoparasites, like Bd. In a past study on wood frogs (*Rana sylvatica*), reduced

resources did not affect host susceptibility to Ranavirus (Reeve et al. 2013). However, in this

study frogs had reduced access to resources only before individuals were infected, which means

that Ranavirus likely had similar levels of host-consumed resources to utilize (Reeve et al. 2013).

In our study, Ranavirus prevalence decreased with increasing resources, the opposite of the

intensity pattern. These results together suggest that increased resources may help the host mount

an initial immune response to combat Ranaviral infections, but once established, Ranavirus can

use host consumed resources to rapidly replicate.

Organisms in natural ecosystems often experience resource limitation from natural or

anthropogenic factors, driving the importance of understanding how resources alter disease

progression in hosts. In our study, higher resources increased amphibian growth but altered

pathogen abundance positively or negatively depending on the pathogen. These results support

the hypothesis that pathogen traits may be an important factor in determining whether the host or

pathogen benefit more from increased resource availability. Other studies have found that

resource availability is positively associated with pathogen replication in viruses (Arsnoe et al.

2011, Hall et al. 2012) and in endoparasites (Pulkkinen and Ebert 2004, Tylianakis et al. 2004),

consistent with our results and the notion that rapidly replicating endoparasites might benefit

from increases in host-consumed resources. Future studies should further examine the role that

pathogen traits play in determining whether the host or pathogen benefit more from increased

resource availability.

Acknowledgements: We would like to thank the University of South Florida for allowing access

to use their Botanical Gardens for specimen collection. Thanks also to Jason Hoverman who

supplied the Ranavirus, Joyce Longcore who supplied the Bd used for infecting amphibians, and

Sarah Knutie for developing and assisting with the ELISA assay used in this experiment. JRR

was supported by grants from the National Institutes of Health (R01TW010286-01), and the

National Science Foundation (DEB-2017785, IOS-1754868).

References:

Arsnoe, D. M., H. S. Ip, and J. C. Owen. 2011. Influence of body condition on influenza A virus

infection in mallard ducks: experimental infection data. PLoS One **6**:e22633.

Bekhet, G. A., H. A. Abdou, S. A. Dekinesh, H. A. Hussein, and S. S. Sebiae. 2014. Biological

factors controlling developmental duration, growth and metamorphosis of the larval

green toad, Bufo viridis viridis. The Journal of Basic & Applied Zoology **67**:67-82.

Boyle, D. G., D. B. Boyle, V. Olsen, A. T. Morgan, and A. D. Hyatt. 2004. Rapid quantitative

detection of chytridiomycosis (*Batrachochytrium dendrobatidis*) in amphibian samples

using real-time Taqman PCR assay. Dis Aquat Organ **60**:141-148.

Breheny, P., and W. Burchett. 2019. visreg: Visualization of regression models, R version 2.5-1.

Brown, M. J. F., R. Loosli, and P. Schmid-Hempel. 2000. Condition-depedent expression of

virulence in a trypanosome infecting bumblebees. Oikos **91**:421-427.

Buck, J. C., J. R. Rohr, and A. R. Blaustein. 2016. Effects of nutrient supplementation on host-

pathogen dynamics of the amphibian chytrid fungus: a community approach. Freshw Biol

61:110-120.

Chinchar, V. G. 2002. Ranavirus (family *Iridoviridae*): emerging cold-blooded killers. Archives

of Virology **147**:447-470.

Civitello, D. J., B. E. Allman, C. Morozumi, and J. R. Rohr. 2018. Assessing the direct and

indirect effects of food provisioning and nutrient enrichment on wildlife infectious

disease dynamics. Philos Trans R Soc Lond B Biol Sci **373**.

Civitello, D. J., R. M. Penczykowski, A. N. Smith, M. S. Shocket, M. A. Duffy, and S. R. Hall.

2015. Resources, key traits and the size of fungal epidemics in Daphnia populations. J

Anim Ecol **84**:1010-1017.

Cothran, R. D., S. S. Gervasi, C. Murray, B. J. French, P. W. Bradley, J. Urbina, A. R. Blaustein,

and R. A. Relyea. 2015. Carotenoids and amphibians: effects on life history and

susceptibility to the infectious pathogen, Batrachochytrium dendrobatidis. Conserv

Physiol **3**:cov005.

Cramp, R. L., R. K. McPhee, E. A. Meyer, M. E. Ohmer, and C. E. Franklin. 2014. First line of

defence: the role of sloughing in the regulation of cutaneous microbes in frogs. Conserv

Physiol **2**:cou012.

Cressler, C. E., W. A. Nelson, T. Day, and E. McCauley. 2014a. Disentangling the interaction

among host resources, the immune system and pathogens. Ecol Lett **17**:284-293.

Cressler, C. E., W. A. Nelson, T. Day, and E. McCauley. 2014b. Starvation reveals the cause of

infection-induced castration and gigantism. Proceedings of the Royal Society B

281:20141087.

Daszak, P., L. Berger, A. A. Cunningham, A. D. Hyatt, D. E. Green, and R. Speare. 1999.

Emerging infectious diseases and amphibian population declines. Emerging infectious

diseases **5**:735–748.

de Roode, J. C., A. B. Pedersen, M. D. Hunter, and S. Altizer. 2008. Host plant species affects

virulence in monarch butterfly parasites. J Anim Ecol **77**:120-126.

Echaubard, P., K. Little, B. Pauli, and D. Lesbarreres. 2010. Context-dependent effects of

ranaviral infection on northern leopard frog life history traits. PLoS One **5**:e13723.

Gervasi, S., C. Gondhalekar, D. H. Olson, and A. R. Blaustein. 2013. Host identity matters in the

amphibian-Batrachochytrium dendrobatidis system: fine-scale patterns of variation in

responses to a multi-host pathogen. PLoS One **8**:e54490.

320 Gibbons, J. W., D. E. Scott, T. J. Ryan, K. A. Buhlmann, T. D. Tuberville, B. S. Metts, J. L.

321   Greene, T. Mills, Y. Leiden, S. Poppy, and C. T. Winnie. 2000. The global decline of

322   reptiles, déjà vu amphibians. BioScience **50**:653-666.

323 Gosner, K. L. 1960. A simplified table for staging anuran ambryos and larvae with notes on

324   identification. Herpetologica **16**:183-190.

325 Gray, M. J., D. L. Miller, and J. T. Hoverman. 2009. Ecology and pathology of amphibian

326   ranaviruses. Dis Aquat Organ **87**:243-266.

327 Green, D. E., K. A. Converse, and A. K. Schrader. 2002. Epizootiology of sixty-four amphibian

328   morbidity and mortality events in the USA, 1996-2001. Annals New York Academy of

329   Sciences **969**.

330 Hall, S. R., C. R. Becker, M. A. Duffy, and C. E. Cáceres. 2012. A power-efficiency trade-off in

331   resource use alters epidemiological relationships. Ecology **93**:645-656.

332 Hall, S. R., J. L. Simonis, R. M. Nisbet, A. J. Tessier, and C. E. Caceres. 2009. Resource ecology

333   of virulence in a planktonic host-parasite system: an explanation using dynamic energy

334   budgets. Am Nat **174**:149-162.

335 Hite, J. L., and C. E. Cressler. 2019. Parasite-Mediated Anorexia and Nutrition Modulate

336   Virulence Evolution. Integr Comp Biol **59**:1264-1274.

337 Hothorn, W. 2010. multcomp: simultaneous inference in general parametric models, R package

338   version 1.4-10.

339 Hoverman, J. T., M. J. Gray, and D. L. Miller. 2010. Anuran susceptibilities to ranaviruses: role

340   of species identity, exposure route, and a novel virus isolate. Dis Aquat Organ **89**:97-107.

341 Jackman, S., A. Tahk, A. Zeileis, C. Maimone, J. Fearon, and Z. Meers. 2020. pscl: Political

342   Science Computational Laboratory. R version 1.5.5.

Johnson, P. T., J. M. Chase, K. L. Dosch, R. B. Hartson, J. A. Gross, D. J. Larson, D. R.

Sutherland, and S. R. Carpenter. 2007. Aquatic eutrophication promotes pathogenic

infection in amphibians. Proc Natl Acad Sci U S A **104**:15781-15786.

Knutie, S. A., C. L. Wilkinson, Q. C. Wu, C. N. Ortega, and J. R. Rohr. 2017. Host resistance

and tolerance of parasitic gut worms depend on resource availability. Oecologia

183:1031-1040.

Lochmiller, R. L., and C. Deerenberg. 2000. Trade-offs in evolutionary immunology: just what is

the cost of immunity? Oikos **88**:87-98.

McMahon, T. A., and J. R. Rohr. 2015. Transition of chytrid fungus infection from mouthparts

to hind limbs during amphibian metamorphosis. Ecohealth **12**:188-193.

Picco, A. M., J. L. Brunner, and J. P. Collins. 2007. Susceptibility of the endangered California

tiger salamander, Ambystoma californiense, to ranavirus infection. J Wildl Dis **43**:286-

290.

Piotrowski, J. S., S. L. Annis, and J. E. Longcore. 2004. Physiology of *Batrachochytrium*

dendrobatidis, a chytrid pathogen of amphibians   Mycologia **96**:9-15.

Pulkkinen, K., and D. Ebert. 2004. Host starvation decreases parasite load and mean host size in

experimental populations. Ecology **85**:823-833.

Rauw, W. M. 2012. Immune response from a resource allocation perspective. Front Genet **3**:267.

Reeve, B. C., E. J. Crespi, C. M. Whipps, and J. L. Brunner. 2013. Natural stressors and

ranavirus susceptibility in larval wood frogs (Rana sylvatica). Ecohealth **10**:190-200.

Relyea, R. A. 2007. Getting out alive: how predators affect the decision to metamorphose.

Oecologia **152**:389-400.

Rohr, J. R., A. A. Elskus, B. S. Shepherd, P. H. Crowley, T. M. McCarthy, J. H. Niedzwiecki, T.

Sager, A. Sih, and B. D. Palmer. 2004. Multiple stressors and salamanders: effects of an

herbicide, food limitation, and hydroperiod. Ecological Applications **14**:1028-1040.

Rollins-Smith, L. A. 1998. Metamorphsis and the amphibian immune system. Immunological

Reviews **166**:221-230.

Rollins-Smith, L. A., J. K. Doersam, J. E. Longcore, S. K. Taylor, J. C. Shamblin, C. Cary, and

M. A. Zasloff. 2002. Antimicrobial peptide defenses against pathogens associated with

global amphibian declines. Developmental and Comparitive Immunology **26**:63-72.

Seppälä, O., K. Liljeroos, A. Karvonen, and J. Jokela. 2008. Host condition as a constraint for

parasite reproduction. Oikos **0**:080208022927783-080208022927780.

Sutton, W. B., M. J. Gray, R. H. Hardman, R. P. Wilkes, A. J. Kouba, and D. L. Miller. 2014.

High susceptibility of the endangered dusky gopher frog to ranavirus. Dis Aquat Organ

112:9-16.

Therneau, T. M., and T. Lumley. 2019. survival: Survival analysis, R version 2.44-1.1.

Tylianakis, J. M., R. K. Didham, and S. D. Wratten. 2004. Improved fitness of aphid parasitoids

receiving resource subsidies. Ecology **85**:658-666.

Venesky, M. D., T. E. Wilcoxen, M. A. Rensel, L. Rollins-Smith, J. L. Kerby, and M. J. Parris.

2012. Dietary protein restriction impairs growth, immunity, and disease resistance in

southern leopard frog tadpoles. Oecologia **169**:23-31.

Voyles, J., S. Young, L. Berger, C. Campbell, W. F. Voyles, A. Dinudom, D. Cook, R. Webb, R.

A. Alford, L. F. Skerratt, and R. Speare. 2009. Pathogenesis of Chytridiomycosis, a cause

of catastrophic amphibian declines. Science **326**:582-585.

Figure 1. Effects of varying food levels on Cuban tree frog (*Osteopilus septentrionalis*) weight change

(grams/week) when **A)** exposed or not to Ranavirus or **B)** *Bactrachochytrium dendrobatidis* (Bd) and **C)**

on their development rate (Gosner stage/week) when exposed to Ranavirus or not. Increased food

availability increased host growth rates (Ranavirus: $F_{2,51}$=5.72, p<0.006; Bd: $F_{1,68}$= 59.5, $p < 0.001$),

regardless of pathogen exposure, whereas Ranavirus exposure significantly decreased development rates

in the low resource treatment, but not across the other tested resource levels (resource x exposure:

$F_{2,51}$=7.89, p=0.001; low resource exposure: $F_{1,17}$=10.92, p=0.004). Data are shown as a conditional plot,

with expected values (line), 95% confidence intervals for the expected values (shading), and partial

residuals (points).

Figure 2. Pathogen intensity in infected larval (A) and adult (B) Cuban tree-frogs (*Osteopilus*

septentrionalis) fed a range of resource levels (high, medium, and low). **A)** *Bactrachochytrium*

dendrobatidis (Bd) loads were consistently higher in individuals with low resource availability (χ^2_1=4.27,

p<0.05), which was independent of weeks since infection (time since infection-by-resource availability:

χ^2_1=0.51, p=0.47). **B)** Ranavirus loads were higher when hosts were fed more food (χ^2_1=5.17, p<0.05).

Data in both figures are shown as partial residual plots, with the expected values (line), 95% confidence

bands or intervals for the expected values (shading), and partial residuals (points).

Figure 1

A

B

409 **Figure 2**