mRNA vaccine-induced T cells respond identically to SARS-CoV-2 variants of concern

but differ in longevity and homing properties depending on prior infection status

Jason Neidleman[1, 2, †], Xiaoyu Luo[1, †], Matthew McGregor[1, 2], Guorui Xie[1, 2], Victoria Murray[3],

Warner C. Greene[1, 4], Sulggi A. Lee[5, *], and Nadia R. Roan[1, 2, *, #]

[1] Gladstone Institute of Virology, San Francisco, CA, USA

[2] Department of Urology, University of California, San Francisco, CA, USA

[3] Zuckerberg San Francisco General Hospital and the University of California, San Francisco, CA,

USA

[4] Departments of Medicine, and Microbiology and Immunology, University of California, San

Francisco, CA, USA

[†]Equal contribution

[*]Co-correspondence:

nadia.roan@gladstone.ucsf.edu, sulggi.lee@ucsf.edu

[#]Lead contact

ABSTRACT

While mRNA vaccines are proving highly efficacious against SARS-CoV-2, it is important

to determine how booster doses and prior infection influence the immune defense they elicit,

and whether they protect against variants. Focusing on the T cell response, we conducted a

longitudinal study of infection-naïve and COVID-19 convalescent donors before vaccination and

after their first and second vaccine doses, using a high-parameter CyTOF analysis to phenotype

their SARS-CoV-2-specific T cells. Vaccine-elicited spike-specific T cells responded similarly to

stimulation by spike epitopes from the ancestral, B.1.1.7 and B.1.351 variant strains, both in

terms of cell numbers and phenotypes. In infection-naïve individuals, the second dose boosted

the quantity and altered the phenotypic properties of SARS-CoV-2-specific T cells, while in

convalescents the second dose changed neither. Spike-specific T cells from convalescent

vaccinees differed strikingly from those of infection-naïve vaccinees, with phenotypic features

suggesting superior long-term persistence and ability to home to the respiratory tract including

the nasopharynx. These results provide reassurance that vaccine-elicited T cells respond

robustly to emerging viral variants, confirm that convalescents may not need a second vaccine

dose, and suggest that vaccinated convalescents may have more persistent nasopharynx-

homing SARS-CoV-2-specific T cells compared to their infection-naïve counterparts.

INTRODUCTION

A year and a half since the December 2019 emergence of SARS-CoV-2, the novel betacoronavirus had already infected almost 200 million people and taken the lives of over 4 million, nearly collapsed worldwide health systems, disrupted the global economy, and perturbed society and public health on a scale not experienced within the past 100 years. Fortunately, multiple highly-efficacious vaccines, including the two-dose mRNA-based ones developed by Pfizer/BioNTech and Moderna, which confer ~90% protection against disease, were approved for emergency use before the end of 2020. Although the vaccines provide the most promising route for a rapid exit from the COVID-19 pandemic, concerns remain regarding the durability of the immunity elicited by these vaccines and the extent to which they will protect against the variants of SARS-CoV-2 now spreading rapidly around the world.

The first variant observed to display a survival advantage was the D614G, which was more transmissible than the original strain and quickly became the dominant variant throughout the world [1]. This variant, fortunately, did not evade immunity and in fact appeared to be more sensitive than the original strain to antibody neutralization by convalescent sera [2]. More worrisome, however, was the emergence at the end of 2020 of rapidly-spreading variants in multiple parts of the world, including B.1.1.7, B.1.351, P.1, and B.1.427/B.1.429 (originally identified in United Kingdom, South Africa, Brazil, and California, respectively) [3], followed by additional highly transmissible variants in 2021 including the B.1.61.72 which was first detected in India [4]. Some variants, including B.1.1.7, may be more virulent [5]. While antibodies against the original strain elicited by either vaccination or infection generally remain potent against B.1.1.7, their activity against B.1.351 and P.1 is compromised [6-15]. Antibodies from vaccinees were 14-fold less effective against B.1.351 than against the ancestral strain, and a subset of individuals completely lacked neutralizing antibody activity against B.1.351 9 months or more after convalescence [13].

Reassuringly, early data suggest that relative to antibody responses, T cell-mediated immunity appears to be less prone to evasion by the variants [16-22]. Among 280 CD4+ and 523

CD8+ T cell epitopes from the original SARS-CoV-2, an average of 91.5% (for CD4) and 98.1%

(for CD8) mapped to regions not mutated in the B.1.1.7, B1.351, P.1, and B.1.427/B.1.429

variants. Focusing on just the spike response, the sole SARS-CoV-2 antigen in the mRNA-based

vaccines, then 89.7% of the CD4+ epitopes and 96.4% of the CD8+ epitopes are conserved [17].

In line with this, the magnitude of the response of T cells from convalescent or vaccinated

individuals was not markedly reduced when assessed against any of the variants [17]. The relative

resistance of T cells against SARS-CoV-2 immune evasion is important in light of the critical role

these immune effectors play during COVID-19. T cell numbers display a strong, inverse

association with disease severity [23, 24], and the frequency of SARS-CoV-2-specific T cells predicts

recovery from severe disease [25, 26]. SARS-CoV-2-specific T cells can also provide long-term, self-

renewing immunological memory: these cells are detected more than half a year into

convalescence, and can proliferate in response to homeostatic signals [27, 28]. Furthermore, the

ability of individuals with inborn deficiencies in B cell responses to recover from COVID-19 without

intensive care suggests that the combination of T cells and innate immune mechanisms is

sufficient for recovery when antibodies are lacking [29].

Although T cells against the ancestral strain display a response of similar magnitude and

breadth to the variants [17], to what extent these T cells' phenotypes and effector functions differ

during their response to variant detection is a different question. Small changes in the sequences

of T cell epitopes, in the form of altered peptide ligands (APLs), can theoretically alter how the T

cells respond to stimulation. Indeed, change of a single residue can convert a proliferative, IL4-

secreting effector response into one that continues to produce IL4 in the absence of proliferation

[30]. Furthermore, APLs can activate Th1 cells without inducing either proliferation or cytokine

production, shift Th1 responses into Th2-focused ones, and in some instances even render T

cells anergic or immunoregulatory by eliciting TGFβ production [31].

Another important aspect that hasn't been explored is to what extent vaccine- vs. infection-

induced T cell responses differ phenotypically and functionally, and to what extent convalescent

individuals benefit from vaccination as they already harbor some form of immunity against the

virus. Studies based on the antibody and B cell response suggest that for COVID-19

convalescents, a single dose of the mRNA vaccines is helpful while the additional booster is not

necessary [10, 32, 33]; how this translates in the context of vaccine-elicited T cell immunity is not clear.

To address these knowledge gaps, we conducted 39-parameter phenotyping by CyTOF

on 33 longitudinal specimens from 11 mRNA-vaccinated individuals, 6 of whom had previously

contracted and recovered from COVID-19. For each participant, blood specimens were obtained

prior to vaccination, two weeks following the first dose, and two weeks following the second. For

every specimen, we assessed in depth the phenotypes and effector functions of total CD4+ and

CD8+ T cells, and of CD4+ and CD8+ T cells responding to the original SARS-CoV-2 spike, to

spike from variants B.1.1.7 and B.1.351, and to nucleocapsid. By conducting analyses on the

resulting 165 high-dimensional datasets generated, we find a reassuringly unaltered T cell

response against the variants, an ability of the booster dose to alter the phenotypes of vaccine-

elicited T cells, and a striking impact of prior infection on qualitative features of T cells elicited by

vaccination.

RESULTS

Study Design

To characterize the phenotypic features of mRNA vaccination-elicited SARS-CoV-2-

specific T cells, we procured 33 longitudinal blood samples from the COVID-19 Host Immune

Response and Pathogenesis (CHIRP) cohort. Four of the participants had received the

Moderna (mRNA-1273) vaccine, while the remaining 7 had received the Pfizer/BioNTech

(BNT162b2) one. For all participants, longitudinal specimens were obtained at three timepoints:

prior to vaccination, ~2 weeks (range 13-18 days) after the first vaccine dose, and ~2 weeks

(range 6-38 days) after the second dose. Five of the participants were never infected with

SARS-CoV-2, while the remaining 6 had completely recovered from mild (non-hospitalized)

COVID-19 disease (Table S1). These prior infections had all occurred in the San Francisco Bay

Area between March – July of 2020, when the dominant local strain was the original ancestral

strain. Each specimen was phenotyped using a 39-parameter T cell-centric CyTOF panel (see

Methods and Table S2) at baseline (to establish the overall phenotypes of total CD4+ and CD8+

T cells), and following 6 hours of stimulation with overlapping 15-mer peptides spanning the

entire original (ancestral) SARS-CoV-2-spike, B.1.1.7 spike, B.1.351 spike, or the original

SARS-CoV-2 nucleocapsid (the latter as a control for a SARS-CoV-2-specific response not

boosted by vaccination). Including all the baseline and stimulation conditions, a total of 165

specimens from the 11 participants were analyzed by CyTOF.

SARS-CoV-2-specific T cells elicited by vaccination recognize B.1.1.7 and B.1.351

variants

We first confirmed our ability to identify SARS-CoV-2-specific T cells by stimulating

PBMCs from vaccinated individuals with spike peptides. In line with our prior studies

implementing a 6-hour peptide stimulation [26, 28], spike-specific CD4+ T cells could be specifically

identified through intracellular cytokine staining for IFNγ, and a more robust response was

observed among CD4+ than CD8+ T cells (Fig. 1A). Activation induced markers (AIM) such as

Ox40, 4-1BB, and CD69 could also be identified in T cells after spike peptide stimulation, but

with a higher background in the baseline (no peptide stimulation) specimens relative to the

intracellular cytokine staining approach (Fig. S1). For this reason, in this study we exclusively

used IFNγ positivity in the peptide-stimulated samples as a marker of antigen-specific T cells.

In the infection-naïve participants, the first vaccination dose primed a spike-specific

CD4+ T cell response, which was further boosted with the second dose (Fig. 1B, *top left*). For

each participant and time point, similar numbers of cells were stimulated by exposure to the

ancestral or variant spikes. This finding suggests that vaccine-elicited spike-specific CD4+ T

cells recognize ancestral and variant spike equally well, and is consistent with their recently

reported ability to recognize variant strains [17]. The response of vaccine-elicited CD8+ T cells to

spike peptides was weaker, and mostly apparent only after the second dose (Fig. 1B, *top right*).

As expected, vaccination did not elicit T cells able to respond to nucleocapsid peptides (Fig. 1C,

top panels).

In contrast to the infection-naïve individuals where spike-specific CD4+ T cells were

clearly elicited and then boosted upon the second dose, spike-specific CD4+ T cell responses in

convalescent individuals did not show a consistent upward trend. Convalescent donor PID4112

had a large frequency of pre-vaccination SARS-CoV-2-specific CD4+ T cells that increased to

>1% of the total CD4+ T cell frequency after the first dose and then dampened after dose 2 (Fig.

1B, *bottom left*). PID4112 also exhibited an elevated nucleocapsid-specific CD4+ T cell

response after the first vaccination dose (Fig. 1C, *bottom left*), which may have been due to

bystander effects resulting from the concomitant large spike-specific response. In comparison,

PID4112's spike-specific CD8+ T cell response was low after dose 1, and boosted after dose 2

(Fig. 1B, *bottom right*). In contrast to PID4112, the remaining five convalescent donors exhibited

an overall weak spike-specific T cell response. In fact, when comparing these five donors to the

five infection-naïve donors, there was a significant decrease in the magnitude of the spike-

specific CD4+ T cell response, while the spike-specific CD8+ T cell response was equivalent

between the two groups (Fig. 1D). These results were unexpected and suggest that, when

excluding outlier PID4112, the magnitude of the vaccine-elicited spike-specific CD4+ T cell

response (after full vaccination) was lower in convalescent individuals than in infection-naïve

individuals.

These assessments of the magnitude of the spike-specific T cell response together

suggest that 1) in infection-naïve individuals the CD4+ T cell response is boosted by the second

vaccination dose, 2) convalescent individuals exhibit a more disparate response, with most

donors mounting a weaker response than infection-naïve individuals, and 3) the response is

more robust among CD4+ than CD8+ T cells. As a higher number of SARS-CoV-2-specific

CD4+ T cells were available for analysis, we focused on this subset for our subsequent

analyses.

Vaccine-elicited spike-specific CD4+ T cells responding to B.1.1.7 and B.1.351 spike are

indistinguishable from those responding to ancestral spike

Leveraging our ability to not only assess the magnitude but also the detailed (39-

parameter) phenotypic features of SARS-CoV-2-specific CD4+ T cells, we first determined

whether the ancestral and variant spike epitopes stimulated different subsets of vaccine-elicited

spike-specific CD4+ T cells. Such differences could theoretically result from the fact that ~5-

10% of the spike epitopes differ between variants and ancestral strains [17], and may therefore

act as APLs steering responding cells towards different fates. We isolated the datasets

corresponding to both post-vaccination timepoints for all eleven donors, and then exported the

data corresponding to spike-specific CD4+ T cells (as defined by IFNγ production, Fig. 1). After

reducing the multidimensional single-cell data for each individual specimen to a two-dimensional

datapoint through multidimensional scaling (MDS) [34], we observed the ancestral spike-

stimulated samples to be interspersed among the B.1.1.7- and B.1.351-responding ones (Fig.

2A). We then visualized the spike-specific CD4+ T cells at the single-cell level. When visualized

alongside total (baseline) CD4+ T cells, spike-specific cells occupied a distinct "island" defined

by high expression of IFNγ (Fig. 2B), suggesting unique phenotypic features of these cells. To

better analyze these spike-responding CD4+ T cells, we visualized them in isolation within a

new t-SNE which clearly demonstrated complete mixing of the cells stimulated by the ancestral,

B.1.1.7, and B.1.351 spike proteins (Fig. 2C). Almost all of the responding cells expressed high

levels of CD45RO and low levels of CD45RA (Fig. 2D), suggesting them to be mostly memory

cells. These memory CD4+ T cells included central memory T cells (Tcm), T follicular helper

cells (Tfh), and those expressing multiple activation markers (CD38, HLADR, CD69, CD25) and

receptors known to direct cells to tissues including the respiratory tract (CXCR4, CCR5, CCR6,

CD49d) (Fig. 2E). The expression levels of these and all other antigens quantitated by CyTOF

were not statistically different between CD4+ T cells responding to the three spike proteins (Fig.

S2). To confirm the identical phenotypes of the three groups of spike-responding cells, we

implemented unbiased clustering by flowSOM. Spike-stimulated cells were clustered into 8

subsets, and no subset was preferentially enriched in any one of the three groups (Fig. 2F).

Together, these data suggest that vaccine-elicited spike-specific CD4+ T cells respond in the

same manner to spike epitopes from the ancestral or variant strains, and would probably mount

similar responses *in vivo* to infection by all three virus types.

Phenotypic alterations of spike-specific CD4+ T cells in infection-naïve recipients after

the second vaccine dose

204   We next took advantage of our longitudinal study design to assess for any changes in

the differentiation of spike-specific T cell responses over the course of the 2-dose vaccination.

As the data presented above suggested no phenotypic differences between CD4+ T cells

responding to the ancestral, B.1.1.7, and B.1.351 spike proteins, our subsequent analyses

combined these datasets. We first assessed whether, among infection-naïve individuals, the

phenotypes of spike-specific CD4+ T cells were different after the first and second doses. While

MDS and tSNE visualizations of the data revealed that the cells from the two timepoints were

somewhat interspersed (Fig. 3A, B), flowSOM clustering suggested some differences in cluster

distribution (Fig. 3C, D). Direct comparison of the cluster frequencies revealed a cluster (B8)

significantly enriched after the first dose, and a different cluster (B5) significantly enriched after

the second dose (Fig. 3E). As these two clusters differentially expressed the Tcm markers

CD27 and CCR7 (Fig. 3F), we then assessed whether Tcm cells were differentially represented

among spike-specific CD4+ T cells after each of the vaccination doses. Indeed, Tcm cells were

significantly higher after the first dose (Fig. 3G), consistent with Cluster 8 (enriched after the first

dose) expressing high levels of these two receptors. Assessment of other canonical CD4+ T cell

subsets – in particular naïve (Tn), stem cell memory (Tscm), effector memory RA (Temra),

effector memory (Tem), T transitional memory (Ttm), Tfh, and regulatory T cells (Treg) –

revealed Tn cells, like the Tcm subset, to be decreased after the second dose. By contrast, Ttm

cells were found to be higher after the second dose, while the remaining subsets were not

altered (Fig. 3G, H). Overall, Tcm and Tfh were the most abundant subsets among the spike-

specific CD4+ T cells (Fig. 3G, H). These data together suggest that after receiving the second

dose, infection-naïve individuals' spike-specific CD4+ T cells increase in quantity (Fig. 1B), and

alter their phenotypes as reflected by a decrease Tcm cells and an increase in Ttm cells.

We then conducted a similar analysis in the convalescent individuals. As the pre-

vaccination timepoint included spike-specific CD4+ T cells primed by prior SARS-CoV-2

infection, we included all three timepoints in this analysis. When the data were visualized by

MDS, it was apparent that most of the pre-vaccination specimens localized away from the post-

vaccination specimens, which were interspersed with each other (Fig. 4A). Similar distinctions

between pre-and post-vaccination specimens were visualized at the single-cell level by tSNE,

which was particularly apparent when visualized as contour heatmaps (Fig. 4B, C). Clustering of

the cells by flowSOM revealed that the cluster distribution was markedly skewed among the pre-

vaccination cells (Fig. 4D, E), with one cluster being under-represented (C2) and one over-

represented (C5) as compared to both post-vaccination timepoints (Fig. 4F). Cluster C3 was the

only cluster that was significantly different after 1 vs. 2 doses (Fig. 4F) but as this cluster

comprised only < 5% of the cells it was not analyzed further. To assess what may drive the

differences between the phenotypes of the pre- vs. post-vaccination spike-specific CD4+ T

cells, we assessed for markers differentially expressed between clusters C2 and C5. Cluster C2

cells preferentially expressed the Tcm markers CD27 and CCR7, the Tfh markers PD1 and

CXCR5, and the co-stimulatory receptors ICOS and Ox40, while among these only CD27 was

preferentially expressed in Cluster C5 (Fig. S3). Manual gating confirmed Tcm, Tfh, and

ICOS+Ox40+ cells to be preferentially enriched in the post-vaccination specimens (Fig. 4G, H,

I). None of the canonical subsets were differentially abundant after the first vs. second

vaccination dose. Together, these results suggest that, in contrast to the infection-naïve

individuals, convalescents' spike-specific CD4+ T cells were similar after the first vs. second

vaccination dose; however, in these individuals vaccination drastically altered the phenotypes of

the pre-existing spike-specific CD4+ T cells (presumably elicited from the original infection).

Vaccination-induced spike-specific CD4+ T cells from convalescent individuals exhibit

unique phenotypic features of increased longevity and tissue homing

We next determined whether there were any phenotypic differences between the

vaccine-induced spike-specific CD4+ T cells from the infection-naïve vs. convalescent

individuals. Removal of convalescent outlier PID4112 revealed the magnitude of the spike-

specific CD4+ T cell response to be lower in the convalescents than in infection-naïve

participants after full vaccination (Fig. 1D). But when all donors were included there was no

statistically significant difference in response magnitude (Fig. 5A). However, the spike-specific

CD4+ T cells from the convalescent and infection-naïve individuals exhibited clear phenotypic

differences when assessed by both MDS (Fig. 5B) and tSNE contours (Fig. 5C); this was more

apparent after the second vaccine dose, but could already be observed after the first. Since the

cells after the second dose are more clinically relevant (as they are the ones persisting in

vaccinated individuals moving forward), we focused our subsequent analysis on just this

timepoint. When visualized as a dot plot, it was apparent that the spike-specific CD4+ T cells

from infection-naïve individuals segregated away from those from the convalescents (Fig. 5D).

Clustering of the data also demonstrated differences between the two patient groups (Fig. 5E,

F), which was confirmed by demonstration of a significant difference in Cluster A1 abundance

(Fig. 5G).

To identify these phenotypic differences, we first assessed the relative distributions of

the main canonical CD4+ T cell subsets. Interestingly, the vaccinated convalescents harbored

significantly more spike-specific Tcm and Tn, and less spike-specific Ttm (Fig. 6A). By contrast,

Tfh and Treg frequencies were not different between infection-naïve and convalescent

vaccinees (Fig. 6B). To broaden our analysis, we assessed for unique features of Cluster A1,

which was over-represented in the infection-naïve donors, and Cluster A3, an abundant cluster

which was over-represented in the convalescent donors albeit insignificantly (Fig. 5G).

Interestingly, Cluster A1 expressed low levels of CD127, CXCR4, and CCR7 in contrast to

Cluster A3 (Fig. S4A). As Cluster A1 is enriched among the infection-naïve individuals, these

findings suggest that these three receptors may be expressed at lower levels on the cells from

these individuals, relative to those from vaccinated convalescents. This was confirmed by our

detection of higher expression of CD127, CXCR4, and CCR7 on spike-specific CD4+ T cells

from the convalescents, although for CXCR4 the difference did not reach statistical significance

(Fig. S4B).

We then followed up on each of these three differentially expressed markers. CD127,

the alpha chain of the IL7 receptor, can drive IL7-mediated homeostatic proliferation of SARS-

CoV-2-specific CD4+ T cells [28], and serves as a marker of long-lived precursor memory cells [35].

To assess the potential longevity of the spike-specific CD4+ T cells, we determined the

percentage of CD127+ cells expressing low levels of the terminal differentiation marker CD57.

After the second dose of vaccination, convalescent individuals harbored more long-lived

(CD127+CD57-) spike-specific CD4+ T cells than infection-naïve individuals (Fig. 6C). CXCR4,

the second preferentially-expressed marker among the convalescents' spike-specific CD4+ T

cells, was recently suggested to direct bystander T cells to the lung during COVID-19, and to be

co-expressed with the T resident memory / activation marker CD69 [26]. Interestingly, spike-

specific CD4+ T cells from convalescent donors harbored a highly significantly elevated

proportion of CXCR4+CD69+ cells (Fig. 6D), suggesting a potentially superior ability to migrate

into pulmonary tissues. The last differentially expressed antigen, CCR7, is a chemokine

receptor that directs immune cells to lymph nodes. As CD62L, a selectin that also mediates

lymph node homing, was also on our panel, we assessed whether CCR7+CD62L+ cells were

enriched among the spike-specific CD4+ T cells from the convalescent donors, and found this to

be the case (Fig. 6E).

Our finding that the convalescent donors' spike-specific CD4+ T cells were preferentially

CXCR4+CD69+ and CCR7+CD62L+ suggested that they may preferentially migrate out of the

blood into lymphoid tissues. Supporting this possibility was our observation that, after the

second vaccine dose, the percentages of CCR7+CD62L+ spike-specific cells increased as the

percentages of spike-specific CD4+ T cells decreased (Fig. 6F). This suggests that the low

spike-specific CD4+ T cell response after the second dose of vaccination in some convalescent

donors (Fig. 1D) may have resulted from these cells preferentially leaving the blood

compartment. This was further supported by our finding that the expression levels of CCR7 and

CD62L on spike-specific CD4+ T cells inversely correlated with the magnitude of the spike-

specific CD4+ T cell response (Fig. 6G). To assess whether the CCR7+CD62L+ and

CXCR4+CD69+ CD4+ T cells have the potential to migrate into the nasopharynx, the most

common site of SARS-CoV-2 entry, we obtained paired blood and nasal swabs from one of the

participants (PID4101) and phenotyped total CD4+ T cells isolated from these specimens.

There was a marked enrichment of both CCR7+CD62L+ and CXCR4+CD69+ CD4+ T cells in

the intranasal specimens (Fig. 6H), suggesting that CD4+ T cells expressing these markers

preferentially exit the blood and enter the nasopharynx. Together, these data suggest that after

vaccination, spike-specific CD4+ T cells from convalescent individuals differ from those in

infection-naïve individuals in that they appear to be more long-lived, and may more readily

migrate out of the blood to mucosal sites, thus explaining their overall lower frequencies

measured from the blood.

Phenotypic features of spike-specific CD8+ T cells from vaccinated, convalescent individuals are unique but differ from their CD4+ T cell counterparts

Finally, we assessed to what extent the main similarities and differences observed with spike-specific CD4+ T cells were also seen for spike-specific CD8+ T cells. Similar to the CD4+ T cells, spike-specific CD8+ T cells stimulated by the three different spike proteins (ancestral, B.1.1.7, B.1.351) did not differ in their phenotypic features (Fig. S5A-C). Also similar to the CD4+ T cells, spike-specific CD8+ T cells elicited by vaccination differed phenotypically in the infection-naïve vs. convalescent individuals (Fig. S5D-F). Unlike the CD4+ T cell data, however, these phenotypic differences could not be accounted for by distribution changes among the main canonical subsets Tn, Tscm, Temra, Tcm, Tem, and Ttm (Fig. S5G). Also unlike the CD4+ T cells, these differences were also not explained by differential abundance of the CD127+CD57-, CXCR4+CD69+, or CCR7+CD62L+ subsets (Fig. S5H). Instead, the differences appear to be due to other phenotypic changes, including elevated frequencies of activated cells in the convalescent donors, in particular those co-expressing the Tcm marker CD27 and activation marker CD38, and the checkpoint inhibitor molecule CTLA4 and activation marker 4-1BB (Fig. S5I). These results suggest that vaccine-elicited spike-specific CD8+ T cells, like their CD4+ counterparts, respond equivalently to the B.1.1.7 and B.1.351 variants, and exhibit qualitative differences in convalescent individuals but via different phenotypic alterations than their CD4+ counterparts.

DISCUSSION

T cells are important orchestrators and effectors during antiviral immunity. They may hold the key to long-term memory due to their ability to persist for decades, yet these cells have been disproportionately understudied relative to their humoral immune counterparts in the context of COVID-19. Here, we designed a longitudinal study assessing both the frequency and phenotypic characteristics of SARS-CoV-2-specific T cells in order to address the following

questions: 1) Do SARS-CoV-2-specific T cells elicited by vaccination respond similarly to

ancestral and variant strains?, 2) To what extent is the second dose needed for boosting T cell

responses in infection-naïve and convalescent individuals?, and 3) Do vaccine-elicited memory

T cells differ in infection-naïve vs. convalescent individuals?

To answer the first question, we compared post-vaccination SARS-CoV-2 spike-specific

T cell responses against ancestral vs. the variant B.1.1.7 and B.1.351 strains. Consistent with

recent studies [16-22], we find that vaccination-elicited T cells specific to the ancestral spike protein

also recognize variant spike proteins. We further demonstrate that the phenotypic features of

these cells are identical, whether they are stimulated by ancestral or variant spike proteins. This

was important to establish because of prior reports that effector T cells can respond differently

to APLs by altering their cytokine production or by mounting an immunoregulatory response [30,

31]. APLs could theoretically arise when a variant infects an individual that was previously

exposed to ancestral spike through vaccination or prior infection. That both the quantity and

quality of T cell responses is maintained against the variants may provide an explanation for the

real-world efficacy of the vaccines against variants. Although limited data are available, thus far

all vaccines deployed in areas where the B.1.1.7 or B.1.351 strains dominate have protected

vaccinees from severe and fatal COVID-19 [36]. Given the potentially important role of SARS-

CoV-2-specific T cells in protecting against severe and fatal COVID-19 [26, 27], we postulate that

this protection may have been in large part mediated by vaccine-elicited T cells. In contrast,

efficacy of the vaccines against mild or moderate disease in variant-dominated regions of the

world is more variable. For example, in South Africa where B.1.351 is dominant, the

AstraZeneca ChAdOx1 vaccine only prevented ~10% of mild-to-moderate disease cases [37],

while more recent data from Pfizer/BioNTech's vaccine administered in Qatar, where both

B.1.1.7 and B.1.351 are dominant, revealed that fully vaccinated individuals were 75% less

likely to develop COVID-19 [38]. The overall diminished vaccine-mediated protection against

milder disease in variant-dominated regions of the world might be explained by the likely

important role of antibodies to prevent initial infection by blocking viral entry into host cells

(manifesting as protection against asymptomatic and mildly symptomatic infection), and the

observation that vaccine-elicited antibodies are generally less effective against the variant than

against ancestral spike in lab assays [6-15]. Reassuringly, there is no evidence that vaccinated

individuals mount a weaker immune response to variants than do unvaccinated individuals,

which could theoretically result through a phenomenon termed original antigenic sin (where the

recall response is inappropriately diverted to the vaccination antigen at the expense of a

protective response against the infecting variant strain) [39].

To address the second question of whether a booster dose is needed, we compared the

T cells after the first vs. second vaccination doses, among the infection-naïve and convalescent

individuals. In infection-naïve individuals, spike-specific responses were observed after the first

vaccination dose, and were further boosted after the second. This enhancement of the T cell

response after the second dose is similar to the reported increase in anti-spike IgG levels after a

second dose in infection-naïve individuals [32, 33]. Interestingly, phenotypic changes were also

observed after the second dose in that the B cells producing the anti-spike antibodies

differentiated from IgM-dominant to IgG-dominant producers [32]. We also observed some

phenotypic changes among spike-specific CD4+ T cells after the second dose, as reflected by

an increase in the Ttm response at the expense of the Tcm response. Importantly, however,

after either dose, spike-specific CD4+ T cells were still primarily Tcm and Tfh cells, the latter of

which are important for providing helper function for B cells. The prominence of SARS-CoV-2-

specific Tfh cells after just one dose of vaccination is consistent with prior reports that a single

dose of SARS-CoV-2 mRNA in mice is sufficient to elicit potent B and Tfh cell responses in

germinal centers [40]. These results suggest that with regards to T cells, the booster dose is

necessary for enhancing the magnitude and results in some phenotypic changes although a

robust Tfh response is already established the first dose. Overall, our conclusions are in line

with those drawn from serological studies [32, 33]: that it is important to administer the second

vaccine dose in infection-naïve individuals to boost spike-specific responses.

A different situation appears to be the case for convalescent individuals. Longitudinal

serological studies suggest that the spike-specific antibody response in convalescent individuals

after the first mRNA dose is already equivalent to that of infection-naïve individuals after their

second mRNA dose [32, 33], suggesting that convalescent individuals may only need a single dose

of vaccination. We found no evidence of increased numbers of spike-specific CD4+ T cells after

the second dose, and minimal phenotypic changes between the cells at the two post-

vaccination timepoints. Spike-specific CD4+ T cells from these individuals did however exhibit

marked phenotypic changes as they transitioned from the pre- to the post-vaccination

timepoints. This was expected since the cells from the pre-vaccination timepoint are resting

memory CD4+ T cells that were primed months prior, while the post-vaccination timepoints were

more recently-reactivated memory cells. Interestingly, unlike for the infection-naïve individuals

where all individuals responded similarly to each dose of vaccination, the magnitude of the

CD4+ T cell response differed markedly between different convalescent individuals. PID4112

had a large pool of spike-specific CD4+ T cells prior to vaccination, and their numbers increased

to extremely high levels after the first vaccination dose. Surprisingly, this large peak in the spike-

specific response was accompanied by an increase in the nucleocapsid-specific CD4+ T cells,

which was unexpected since the vaccine does not contain nucleocapsid. We suspect this high

response to nucleocapsid was due to inflammation-mediated bystander activation of T cells in

an antigen-independent manner. Consistent with this hypothesis, the participant reported severe

side effects (severe headache, chills, myalgia, nausea, and diarrhea) after the first dose. The

remaining five convalescent donors, by contrast, never exhibited a robust T cell response, and

in fact after full vaccination actually exhibited a highly significantly lower CD4+ T cell response

than the infection-naïve vaccinees. We speculate on an explanation further below. Overall, our

results suggest that a second SARS-CoV-2 vaccine dose in individuals who have recovered

from COVID-19 may provide less benefit than in individuals who have not previously been

exposed to SARS-CoV-2; these findings are in line with recommendations from previously

published serological studies [10, 32, 33].

427   One of the most striking observations from this study, and the third and final question we

set out to answer, was the remarkably distinct phenotypes of spike-specific CD4+ T cells from

infection-naïve vs. convalescent individuals who were fully vaccinated. The spike-specific CD4+

T cells from the convalescent individuals harbored features suggesting increased potential for

long-term persistence: they were enriched for Tcm cells, which are have longer *in vivo* half-lives

than their Tem and Ttm counterparts [41], and express elevated levels of CD127, a marker of

long-lived memory T cells [35]. Interestingly, CD127 expression on SARS-CoV-2-specific T cells

has been implicated in COVID-19 disease amelioration and in these cells' long-term

persistence. CD127 expression was more frequent on spike-specific CD4+ T cells from ICU

patients who eventually survived severe COVID-19 than in those that did not [26]. IL7, the ligand

for CD127, can drive homeostatic proliferation and expansion of spike-specific CD4+ T cells [28],

and CD127 is not only expressed on SARS-CoV-2-specific memory CD4+ and CD8+ T cells,

but its levels increase further over the course of convalescence [28, 42]. Together, these findings

suggest that after vaccination, spike-specific CD4+ T cells in convalescent individuals may

persist longer than those from infection-naïve individuals, but additional long-term follow-up

studies will be required to directly test whether this indeed is the case.

443   Another interesting characteristic of post-vaccination spike-specific CD4+ T cells from

convalescent individuals relative to infection-naïve individuals was their expression of multiple

tissue-homing receptors. In particular, these cells were preferentially CCR7+CD62L+ and

CXCR4+CD69+. CCR7 and CD62L mediate homing to lymph nodes, while CXCR4 is a

chemokine receptor important in migration of hematopoietic stem cells to bone marrow, but also

able to direct immune cells to the lung during inflammation [43]. Interestingly, we recently

observed co-expression of CXCR4 with CD69 (an activation marker that also identifies T

resident memory cells) in pulmonary T cells from COVID-19 patients [26]. Many of these cells

were bystander (non-SARS-CoV-2-specific) CXCR4+CD69+ T cells whose numbers in blood

increased prior to death from COVID-19. We therefore proposed a model whereby recruitment

of non-SARS-CoV-2-specific T cells into the lungs of severe patients may exacerbate the

cytokine storm and thereby contribute to death [26]. In the case of the vaccinated convalescent

individuals, however, expression of CXCR4 and CD69 on SARS-CoV-2-specific T cells is

expected to be beneficial as it would direct the T cells capable of recognizing infected cells into

the lung. CCR7 and CD62L co-expression would further enable these cells to enter draining

lymph nodes and participate in germinal center reactions. Supporting the hypothesis that the

post-vaccination spike-specific CD4+ T cells from convalescent individuals may better home to

lymphoid tissues is our observation that frequencies of these cells in blood correlated negatively

with the extent to which they co-expressed CCR7 and CD62L. This was further supported by

our finding that CD4+ T cells from the nasopharynx of the upper respiratory tract were

preferentially CCR7+CD62L+ and CXCR4+CD69+ relative to their blood counterparts. All

together, these results imply that compared to infection-naïve individuals, convalescents' spike-

specific CD4+ T cells may be superior in surviving and migrating to the respiratory tract. Directly

testing this hypothesis will require obtaining large numbers of respiratory tract cells from

vaccinated, infection-naïve vs. convalescent individuals (e.g., via bronchoalveolar lavages or

endotracheal aspirates) for quantitation and characterization of SARS-CoV-2-specific T cells. Of

note, vaccination of infection-naïve individuals might not induce a strong humoral immunity in

the respiratory mucosa either, as neutralizing antibodies against SARS-CoV-2 are rarely

detected in nasal swabs from vaccinees [13]. If it turns out that current vaccination strategies do

not ensure robust humoral and cell-mediated immune responses in the respiratory tract, then

strategies that better elicit mucosal-homing SARS-CoV-2-specific B and T cells in infection-

naïve individuals – for example by implementing an intranasal route of mRNA immunization –

may hold a greater chance of achieving sterilizing immunity.

Limitations

As this study was aimed at using in-depth phenotypic characterization as a discovery tool, it

focused on deeply interrogating many different conditions (e.g., spike variants, longitudinal

sampling) rather than many donors. Therefore, although a total of 165 CyTOF specimens were

run, only 11 donors were analyzed. The findings reported here should be confirmed in larger

cohorts. A second limitation of the study was the need to stimulate the specimens in order to

identify and characterize the vaccine-elicited T cells. We limited peptide exposure to 6 hours to

minimize phenotypic changes caused by the stimulation, similar to our prior studies [26, 28]. Finally,

the analysis focused on CD4+ T cells because the overall numbers of detectable spike-specific

CD8+ T cells were low. Nonetheless, the main findings we made with the CD4+ T cells – that

they recognize variants equivalently, and that the phenotypes of the responding cells differ by

prior SARS-CoV-2 natural infection status – were recapitulated among CD8+ T cells. Additional

studies in a larger number of participants testing more cells, and implementing the use of

combinatorial MHC class I tetramers in conjunction with high-parameter phenotyping [44], would

increase the ability to characterize in greater depth the vaccine-elicited CD8+ T cell response.

ACKNOWLEDGEMENTS

This work was supported by the Van Auken Private Foundation, David Henke, and Pamela and Edward Taft (N.R.R.); philanthropic funds donated to Gladstone Institutes by The Roddenberry Foundation and individual donors devoted to COVID-19 research (N.R.R.); the Program for Breakthrough Biomedical Research (N.R.R., S.A.L.), which is partly funded by the Sandler Foundation; and Awards #2164 (N.R.R.), #2208 (N.R.R.), and #2160 (to S.A.L.) from Fast Grants, a part of Emergent Ventures at the Mercatus Center, George Mason University. We acknowledge the NIH DRC Center Grant P30 DK063720 and the S10 1S10OD018040-01 for use of the CyTOF instrument. We thank Stanley Tamaki and Claudia Bispo for CyTOF assistance at the Parnassus Flow Core, Heather Hartig for help with recruitment, Françoise Chanut for editorial assistance, and Robin Givens for administrative assistance.

AUTHOR CONTRIBUTIONS

J.N. designed and performed experiments, and conducted data analyses; X.L. helped develop an analysis plan and conducted data analyses; M.M. processed and banked specimens; G.X. performed experiments; V.M. conducted CHIRP participant interviews, enrollment, and specimen collection; W.C.G. participated in data analysis, performed supervision, and edited the manuscript; S.A.L. established the CHIRP cohort, conducted CHIRP participant interviews, enrollment, and specimen collection, and edited the manuscript; N.R.R. conceived ideas for the study, performed supervision, conducted data analyses, and wrote the manuscript. All authors read and approved the manuscript.

COMPETING FINANCIAL INTERESTS: The authors declare no competing financial interests.

METHODS

Human Subjects

Eleven participants from the COVID-19 Host Immune Pathogenesis (CHIRP) cohort were recruited for this study. Six were previously infected with SARS-CoV-2 as established by RT-PCR, and had fully recovered from a mild course of disease. Importantly, infections of these six individuals had all occurred in the San Francisco Bay Area between March – July of 2020, when the dominant local strain was the original ancestral (Wuhan) strain. The remaining five participants were not previously infected with the virus. All eleven participants were vaccinated with both doses of either of the Moderna or Pfizer/BioNTech mRNA vaccines (Table S1). Blood was drawn from each of the eleven participants prior to vaccination, ~2 weeks after the first vaccine dose, and ~2 weeks after the second vaccine dose (33 specimens total). On the day of each blood draw, PBMCs were isolated from blood using Lymphoprep™ (StemCell Technologies), and then cryopreserved in 90% fetal bovine serum (FBS) and 10% DMSO. For participant PID4101, an additional blood-draw and intranasal swab specimens were obtained for immunophenotyping studies. This study was approved by the University of California, San Francisco (IRB # 20-30588).

Preparation of specimens for CyTOF

Cryopreserved PBMCs were revived and cultured overnight to allow for antigen recovery. The cells were then counted, and then two million cells per treatment condition were stimulated with the co-stimulatory agents 0.5 μg/ml anti-CD49d clone L25 and 0.5 μg/ml anti-CD28 clone L293 (both from BD Biosciences), in the presence of 0.5 μM of overlapping 15-mer SARS-CoV-2 spike peptides PepMix™ SARS-CoV-2 peptides from the original SARS-CoV-2 strain, B.1.1.7, or B.1.351, or overlapping 15-mer SARS-CoV-2 nucleocapsid peptides (all from JPT Peptide Technologies). Stimulations were conducted for 6 hours in RP10 media (RPMI

1640 medium (Corning) supplemented with 10% FBS (VWR), 1% penicillin (Gibco), and 1%

streptomycin (Gibco)), in the presence of 3 µg/ml Brefeldin A Solution (eBioscience) to enable

detection of intracellular cytokines. To establish the phenotypes of total T cells in the absence of

stimulation, two million cells were cultured in parallel with the stimulated samples, but in the

presence of only 3 µg/ml Brefeldin A.

After culture, the cells were treated with cisplatin (Sigma-Aldrich) as a live/dead marker

and fixed with paraformaldehyde (PFA) as previously described [28, 45]. Cisplatin treatment and

fixation was performed as follows: first, cells were resuspended in 2 ml PBS (Rockland) with 2

ml EDTA (Corning), followed by addition of 2 ml PBS/EDTA supplemented with 25 µM cisplatin

(Sigma-Aldrich) for 60 seconds. Cisplatin staining was then quenched with 10 ml of CyFACS

(metal contaminant-free PBS (Rockland) supplemented with 0.1% FBS and 0.1% sodium azide

(Sigma-Aldrich)), centrifuged, and resuspended in 2% PFA in CyFACS. Fixation was allowed to

proceed for 10 minutes at room temperature, after which cells were washed twice with CyFACS,

and then resuspended in CyFACS containing 10% DMSO. Fixed cells were stored at -80°C until

analysis by CyTOF. For paired blood/swab specimens from PID4101, cells were immediately

cisplatin-treated and fixed, without prior cryopreservation.

CyTOF staining and data acquisition

CyTOF staining was conducted in a fashion similar to recently described methods [26, 28,

45-48]. Cisplatin-treated cells were thawed, counted, and each treatment condition was barcoded

using the Cell-ID 20-Plex Pd Barcoding Kit (Fluidigm). After the cells were barcoded and

washed, the barcoded samples were combined and diluted to 6×10^6 cells / 800 µl CyFACS per

well in Nunc 96 DeepWell™ polystyrene plates (Thermo Fisher). Cells were blocked with mouse

(Thermo Fisher), rat (Thermo Fisher), and human AB (Sigma-Aldrich) sera for 15 minutes at

4°C, and then washed twice in CyFACS. Surface CyTOF antibody staining (Table S2) was

conducted for 45 minutes at 4°C, in a volume of 100 μl / sample. Cells were then washed three

times with CyFACS and fixed overnight at 4°C in 100 μl of 2% PFA in PBS. The next day,

samples were washed twice with Intracellular Fixation & Permeabilization Buffer (eBioscience),

and incubated for 45 minutes at 4°C. After two additional washes with Permeabilization Buffer

(eBioscience), samples were blocked for 15 minutes at 4°C in 100 μl of Permeabilization Buffer

containing mouse and rat sera. After one additional wash with Permeabilization Buffer, samples

were stained with the intracellular CyTOF antibodies (Table S2) at 4°C for 45 minutes in a

volume of 100 μl / sample. Cells were then washed once with CyFACS, and stained for 20

minutes at room temperature with 250 nM of Cell-ID^{TM} Intercalator-IR (Fluidigm). Immediately

prior to sample acquisition, cells were washed twice with CyFACS buffer, once with MaxPar®

cell staining buffer (Fluidigm), and once with Cell acquisition solution (CAS, Fluidigm). Cells

were resuspended in EQ™ Four Element Calibration Beads (Fluidigm) diluted in CAS

immediately prior to acquisition on a Helios-upgraded CyTOF2 instrument (Fluidigm) at the

UCSF Parnassus flow core facility.

CyTOF data analysis

CyTOF datasets, exported as flow cytometry standard (FCS) files, were de-barcoded

and normalized according to manufacturer's instructions (Fluidigm). FlowJo software (BD

Biosciences) was used to identify CD4+ T cells (live, singlet CD3+CD19-CD4+CD8-) and CD8+

T cells (live, singlet CD3+CD19-CD4-CD8+) among all analyzed samples. IFNγ+ in the

stimulated samples were considered to be the SARS-CoV-2-responsive cells. For high-

dimensional analyses of SARS-CoV-2-specific T cells among the stimulated samples, we

excluded samples with an insufficient number of events (≤ 3) to limit skewing of the data.

Manual gating analysis was initially performed using FlowJo, and then select populations were

exported as FCS files and then imported into R software as GatingSet objects. Using the

CytoExploreR package, 2D-gates were manually drawn on the imported samples. The 2D dot

plots and statistical results were exported for data visualization, bar-graph generation, and

statistical comparisons as previously described

(https://github.com/DillonHammill/CytoExploreR). High-dimensional analyses (MDS, tSNE, and

flowSOM) were performed using R software by implementing a CyTOF workflow recently

described [49].

For MDS plot generation, we used the plotMDS function from the *limma* package with

default settings. Euclidean distances between all samples were calculated using the arcsinh-

transformed median expression levels with cofactor 5, of the lineage and functional markers

listed below.

CD8	Lineage (Only for CD8 subset)
CD4	Lineage (Only for CD4 subset)
CD161	Lineage
HLADR	Lineage
CD45RO	Lineage
CD69	Lineage
CRTH2	Lineage
PD1	Lineage
CXCR5	Lineage
CD27	Lineage
CD3	Lineage
CD2	Lineage
CD62L	Lineage
CCR6	Lineage
OX40	Lineage
CD28	Lineage
CD127	Lineage
RORγt	Lineage
CXCR4	Lineage
CTLA4	Lineage
NFAT	Lineage
CCR5	Lineage
CD137	Lineage
CD95	Lineage
ICOS	Lineage
CD49d	Lineage
CD7	Lineage
Tbet	Lineage
TIGIT	Lineage

CCR7	Lineage
CD45RA	Lineage
CD57	Lineage
CD38	Lineage
α4β7	Lineage
CD25	Lineage
IFNγ	Function
IL6	Function
IL4	Function
IL17	Function

The first (MDS1) and second (MDS2) MDS dimensions were plotted to show the dissimilarities

between samples from the indicated conditions as described [34].

tSNE was performed using the Trsne function from the *Rtsne* package using arcsinh-

transformed expression of lineage markers (no PCA step, iterations = 1000, perplexity = 30,

theta = 0.5). Events corresponding to unstimulated T cells were down-sampled to 1000 cells per

sample, and SARS-CoV-2-specific cells (cell numbers ranging from 4 to 229 per sample) were

all included in the tSNE analyses without down-sampling. Each cell was displayed in a tSNE

plot for dimension reduction visualization and colored with arcsinh-transformed cell marker

expression as heatmaps, or pseudo-colored by the appropriate group.

Unsupervised cell subset clustering was performed using FlowSOM [50] and

ConsensusClusterPlus packages using arcsinh-transformed expression levels of the lineage

markers indicated above [51]. For clustering of SARS-CoV-2-specific T cells, we set the meta-

cluster number to 8 and cluster number to 40. The frequency of each cluster within each sample

was calculated using the following equation:

(Frequency of cluster in specified sample) = (Cell count of cluster / Total cell count of specified

sample)

This was then converted to a percentage by multiplying by 100. The percentages of each cluster

from the selected samples were plotted as box plots with jittered points, followed by statistical

analysis between the groups. To compare the abundance distribution of clusters between

groups, frequencies of clusters in samples from each group were normalized using the equation

below:

(Normalized frequency of cluster in specified sample) = (Frequency of cluster in specified

sample/ Total number of samples in each group)

This was then converted to a percentage by multiplying by 100, and plotted as stacked bar

charts.

Statistical Analysis

The statistical tests used in comparison of groups are indicated within the figure legends. For 2-

group comparisons, student's t-tests were performed and p-values were adjusted for multiple

testing using the Holm-Sidak method where applicable. For comparisons of 3 or more groups,

significance between groups was first evaluated by one-way ANOVA, and then the p-values

were adjusted for multiple testing using the Holm-Sidak method where applicable. For datasets

with significant ANOVA-adjusted p-values (≤ 0.05), we performed Tukey's honestly significant

difference (HSD) post-hoc test to determine the p-values between individual groups.

Raw Data Availability

For this study, a total of 120 specimens were analyzed by CyTOF. Each specimen included

both CD4+ and CD8+ T cells. For each specimen, we gated separately on events

corresponding to CD4+ T cells (live, singlet CD3+CD4+CD8-) and CD8+ T cells (live, singlet

CD3+CD4-CD8+), and exported the files as 240 individual FCS files. These 240 raw CyTOF

datasets are available for download through the public repository Dryad via the following link:

https://doi.org/10.7272/Q60R9MMK

 MAIN FIGURE LEGENDS

Figure 1. SARS-CoV-2-specific T cells elicited by vaccination recognize variants, and in a

manner that differs among individuals with prior COVID-19. (**A**) Identification of vaccine-

elicited spike-specific T cells. PBMCs before vaccination (Pre-Vac) or 2 weeks after each dose

of vaccination were stimulated with spike peptides and assessed by CyTOF 6 hours later for the

presence of spike-specific (IFNγ-producing) CD4+ (*left*) or CD8+ (*right*) T cells. The "no peptide"

conditions served as negative controls. Shown are longitudinal data from an infection-naïve

(PID4101, *top*) and convalescent (PID4112, *bottom*) individual. **(B)** Quantification of the spike-

specific CD4+ (*left*) and CD8+ (*right*) T cells recognizing the ancestral (squares), B.1.1.7

(triangles), and B.1.351 (circles) spike peptides in infection-naïve (*top*) and convalescent

(*bottom*) individuals before and after vaccination. Note the similar frequencies of T cells

responding to all three spike proteins in each donor, the clear boosting of spike-specific CD4+ T

cell frequencies in infection-naïve but not convalescent individuals, and the overall higher

proportion of responding CD4+ than CD8+ T cells. The dotted line corresponds to the

magnitude of the maximal pre-vaccination response in infection-naïve individuals and is

considered as background. The y-axes are fitted based upon the maximal post-vaccination

response values for each patient group and T cell subset. The *p*-values shown (**p < 0.01, ***p

< 0.001) were calculated by student's t-test. **(C)** As expected, nucleocapsid-specific T cell

responses are generally low over the course of vaccination, with the exception of convalescent

donor PID4112. Shown are the frequencies of nucleocapsid-specific CD4+ (*left*) and CD8+

(*right*) T cells, as measured by IFNγ production upon stimulation with ancestral nucleocapsid

peptides, in infection-naïve (*top*) and convalescent (*bottom*) individuals. The dotted line

corresponds to the magnitude of the maximal pre-vaccination response infection-naïve

individuals, and is considered as the background signal. Y-axes are labeled to match the

corresponding y-axes for spike-specific T cell responses in *panel B*. **(D)** The CD4+ T cell

response is boosted by the second vaccine dose to a greater extent in infection-naïve than

convalescents individuals. Shown are the frequencies of spike-specific CD4+ (*left*) and CD8+

(*right*) T cells stimulated by the three spike proteins (squares: ancestral; triangles: B.1.1.7;

circles: B.1.351) among the infection-naïve (aqua) and convalescent (coral) donors, after

removal of outlier PID4112. ***p < 0.001 comparing the infection-naïve vs. convalescent post-

dose 2 specimens, were calculated using student's t-test.

Figure 2. SARS-CoV-2-specific CD4+ T cells responding to B.1.1.7 and B.1.351 spike have

the same phenotypes as those responding to ancestral spike. (A) Datasets corresponding

to spike-specific CD4+ T cells after vaccination were visualized as a multidimensional scaling

(MDS) plot. Each datapoint reflects the cumulative phenotypes averaged across all the SARS-

CoV-2-specific CD4+ T cells from a single stimulated sample. Data for both infection-naïve and

convalescent individuals, and for both the post-dose 1 and post-dose 2 timepoints, are shown.

The lack of segregation of the cells responding to the ancestral, B.1.1.7, and B.1.351 spike

proteins suggest phenotypic similarities. **(B)** Visualization of the datasets by tSNE dot plots.

CD4+ T cells responding to ancestral or variant spike stimulation by producing high amounts of

IFNγ (*right*) segregate together and away from the total CD4+ T cell population (*left*). Each dot

represents one cell. **(C)** CD4+ T cells responding to ancestral spike and its variants are

phenotypically similar, as shown by their complete mingling on a tSNE dot plot. **(D, E)** Spike-

responding CD4+ T cells are mostly memory cells, as indicated by high CD45RO and low

CD45RA expression levels, and include those expressing high levels of Tcm, Tfh, activation,

and respiratory tract migration markers. Shown is the tSNE depicted in *panel C* displaying the

relative expression levels of the indicated antigens (Red: high; Blue: low). **(F)** CD4+ T cells

responding to ancestral spike and its variants distribute in a similar fashion among the 8 clusters

identified by flowSOM. Shown on the left is the distribution of T cells responding to ancestral or

variant spike peptides on the tSNE depicted in *panel C,* colored according to the flowSOM

clustering. Shown on the right is the quantification of the flowSOM distribution data. No

significant differences were observed between the three groups in the distribution of their cells

among the 8 clusters, as calculated using a one-way ANOVA and adjusted for multiple testing

(n = 8) using Holm-Sidak method ($p > 0.05$).

Figure 3. Phenotypes of spike-specific CD4+ T cells from infection-naïve individuals

following first and second dose of vaccination. (A) MDS plot depicting samples of spike-

specific CD4+ T cells in vaccinated infection-naïve individuals, showing some interspersion of

the cells from the two post-vaccination timepoints. Each dot represents a single specimen. **(B)**

tSNE dot plot of spike-specific CD4+ T cells from vaccinated infection-naïve individuals. Each

dot represents a single cell. **(C)** tSNE plots depicting cells from the two timepoints, colored

according to the cells' cluster classification as determined by flowSOM. **(D)** Distribution among

flowSOM clusters of post-vaccination spike-specific CD4+ T cells from infection-naïve

individuals between the two post-vaccination timepoints. **(E)** Two clusters of spike-specific

CD4+ T cells (B5 and B8) are differentially abundant after the first vs. second vaccination

doses. *$p < 0.05$, *** $p < 0.001$ as determined using student's t-tests adjusted for multiple testing

(n = 8) using Holm-Sidak method. **(F)** The Tcm markers CD27 and CCR7 are differentially

expressed among Clusters B5 and B8, as depicted by histograms. **(G)** The proportions of Tn

(CD45RO-CD45RA+CCR7+CD95-), Tscm (CD45RO-CD45RA+CCR7+CD95+), Temra

(CD45RO-CD45RA+CCR7-), Tcm (CD45RO+CD45RA-CCR7+CD27+), Tem

(CD45RO+CD45RA-CCR7-CD27-), and Ttm (CD45RO+CD45RA-CCR7-CD27+) among spike-

specific CD4+ cells in infection-naive individuals after the first vs. second vaccination doses. *p

< 0.05, ***$p < 0.001$, ns = non-significant as determined by student's t-test. **(H)** The proportions

of Tfh (CD45RO+CD45RA-PD1+CXCR5+) and Treg (CD45RO+CD45RA-CD25+CD127^{low})

among spike-specific CD4+ T cells are similar in infection-naive individuals after the first vs.

second vaccination doses. ns = non-significant as determined by student's t-test.

Figure 4. Differentiation of spike-specific memory CD4+ T cells after vaccination of

convalescent individuals. (A) MDS plot depicting datasets corresponding to spike-specific

CD4+ T cells in convalescent individuals before and after vaccination. **(B)** tSNE contour

heatmaps of spike-specific CD4+ T cells from convalescent individuals emphasizes phenotypic

differences between the pre- and post-vaccination cells. Cell densities are represented by color.

(C) tSNE dot plot of spike-specific CD4+ T cells from convalescent individuals, demonstrating

the distinct localization of the pre-vaccination cells on the right. **(D)** Spike-specific CD4+ T cells

are phenotypically distinct between the pre- and post-vaccination specimens. Shown are tSNE

plots depicting cells from the three indicated timepoints, colored according to the cells' cluster

classification as determined by flowSOM. **(E)** The distribution of spike-specific CD4+ T cells

classified as flowSOM clusters differs between the pre- and post-vaccination timepoints. **(F)**

Multiple clusters of spike-specific CD4+ T cells are differentially abundant between the pre- and

post-vaccination specimens. **p < 0.01, ***p < 0.001, ****p < 0.0001 as determined by one-way

ANOVA and adjusted for multiple testing (n = 8) using the Holm-Sidak method followed by

Tukey's honestly significant difference (HSD) post-hoc test. **(G)** Spike-specific CD4+ Tcm

increase in convalescent individuals after vaccination. Shown are the proportions of Tn, Tscm,

Temra Tcm, Tem, and Ttm among spike-specific CD4+ cells in convalescent individuals before

and after vaccination. **(H)** Spike-specific CD4+ Tfh increase in convalescent individuals after

vaccination. Shown are the proportions of Tfh and Treg among spike-specific CD4+ T cells in

convalescent individuals before and after vaccination. **(I)** Spike-specific CD4+ T cells expressing

ICOS and Ox40 increase in convalescent individuals after vaccination. In panels *G-I*, *p < 0.05,

**p < 0.01, ***p < 0.001, and ****p < 0.0001 as determined by one-way ANOVA followed by

Tukey's HSD post-hoc test.

Figure 5. Phenotypic features of spike-specific CD4+ T cells differ between infection-naïve and convalescent individuals after vaccination. (A) The frequency of spike-specific CD4+ T cells is similar in infection-naïve and convalescent individuals two weeks after the second vaccination dose. Note that when convalescent donor PID4112, who had an unusually high pre-vaccination frequency of spike-specific CD4+ T cells (Fig. 1D), was excluded, the frequency was significantly lower among the convalescents. **(B)** MDS plots of the phenotypes of spike-specific CD4+ T cells in infection-naïve and convalescent individuals after first and second dose vaccinations. **(C)** tSNE contour heatmaps of spike-specific CD4+ T cells from infection-naïve and convalescent individuals, after first and second dose vaccinations, highlighting the phenotypic differences between the two groups of patients. Cell densities are represented by color. **(D)** tSNE dot plot of spike-specific CD4+ T cells from infection-naïve and convalescent individuals after second dose of vaccination, demonstrating the segregation of the cells from the two groups of patients. **(E)** Spike-specific CD4+ T cells are phenotypically distinct between the infection-naïve and convalescent individuals. Shown are tSNE plots depicting cells after the second dose of vaccination, colored according to the cells' cluster classification as determined by flowSOM. **(F)** The distribution of spike-specific CD4+ T cells into flowSOM clusters differs between the infection-naïve and convalescent individuals after the second vaccine dose. **(G)** Cluster A1 is over-represented in infection-naïve relative to convalescent individuals after the second dose of vaccination. **p < 0.01, as determined by student's t-tests adjusted for multiple testing (n = 8) using the Holm-Sidak method.

Figure 6. The post-vaccination spike-specific CD4+ T cells of convalescents harbor phenotypic features of elevated longevity and tissue homing. (A) Spike-specific CD4+ T cells from convalescent vaccinated individuals harbor higher proportions of Tn and Tcm cells and lower proportions of Ttm cells than those from infection-naïve vaccinated individuals. The

proportions of Tn, Tscm, Temra, Tcm, Tem, and Ttm cells among spike-specific CD4+ T cells

were determined by manual gating. **p < 0.01, ***p < 0.001, ****p < 0.0001, ns = non-

significant, as determined by student's t-test. **(B)** The proportions of Tfh and Treg among spike-

specific CD4+ T cells are similar in infection-naïve vs. convalescent individuals after

vaccination. ns = non-significant, as determined by student's t-test. **(C)** Spike-specific CD4+ T

cells expressing the homeostatic proliferation marker CD127 and lacking expression of the

terminal differentiation marker CD57 are more frequent in vaccinated convalescent than

vaccinated infection-naïve individuals. **p < 0.01, as determined by student's t-test. **(D)** Spike-

specific CD4+ T cells expressing CXCR4, which directs cells to tissues including the lung, and

CD69, a marker of T cell activation and tissue residence, are more frequent in convalescent

vaccinated individuals. ***p < 0.001, as determined by student's t-test. **(E)** Spike-specific CD4+

T cells expressing the lymph node homing receptors CCR7 and CD62L are more frequent in

vaccinated convalescent individuals. *p < 0.05, as determined by student's t-test. **(F)** The

proportions of CCR7+CD62L+ cells among spike-specific CD4+ T cells associate negatively

with the frequencies of spike-specific CD4+ T cells after the second dose of vaccination

(correlation coefficient (R) < 0). *P*-values were calculated using t distribution with n-2 degrees

of freedom. **(G)** Expression levels (reported as mean signal intensity, or MSI) of CCR7 and

CD62L among spike-specific CD4+ T cells associate negatively (R < 0) with overall frequencies

of spike-specific CD4+ T cells after the second dose of vaccination. *P*-values were calculated

using t distribution with n-2 degrees of freedom. The 95% confidence intervals of the regression

lines in the scatter plots of *panels F-G* are shaded in grey. **(H)** CCR7+CD62L+ and

CXCR4+CD69+ CD4+ T cells are more frequent in nasopharynx than blood. Unstimulated

CD4+ T cells from the blood (*grey*) or from an intranasal swab (*red*) were obtained on the same

day from PID4101 and then phenotyped by CyTOF. Numbers indicate the percentages of the

corresponding cell population within the gate. Results are gated on live, singlet CD3+CD4+CD8-

cells.

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. Six-hour stimulation with spike peptides does not induce significant expression of activation markers in SARS-CoV-2-specific T cells. (A) CD4+ T cells were assessed for co-expression of the activation-induced markers (AIM) Ox40 and 4-1BB following 6 hours of stimulation with ancestral spike peptides using PBMC specimens from a representative infection-naïve individual (PID4197) before vaccination (Pre-Vac), or two weeks after dose 1 or dose 2 of vaccination. **(B)** CD8+ T cells were assessed for co-expression of the AIM CD69 and 4-1BB following 6 hours of stimulation, using same specimens as *panel A*. Baseline specimens not treated with peptide are shown as a comparison control. Numbers correspond to percentages of cells within the gates. Note that the activated (AIM+) cells that appear in stimulated specimens probably do not reflect peptide-specific stimulation as AIM+ cells are also detected in the baseline specimens.

Figure S2. Expression levels of all CyTOF phenotyping markers are equivalent between CD4+ T cells responding to stimulation by spike from ancestral, B.1.1.7, and B.1.351 spike. Shown are the mean expression levels of each antigen in post-vaccination spike-responding CD4+ T cells quantitated by CyTOF. Each datapoint corresponds to a single specimen. No significant differences were observed in expression levels for any of the antigens between any of the three groups, as assessed by one-way and ANOVA adjusted for multiple testing (n = 39) using the Holm-Sidak method ($p > 0.05$).

Figure S3. Antigens differentially expressed among Clusters C2 and C5, differentially represented among pre- vs. post-vaccination spike-specific CD4+ T cells from convalescent individuals. Shown are histogram depictions of the expression levels of the

indicated activation markers in Cluster C2 **(A)** or C5 **(B)** from convalescent individuals. Cluster

C2 was more abundant post-vaccination, while Cluster C5 was more abundant pre-vaccination.

Figure S4. Cluster A1, enriched among spike-specific CD4+ T cells from infection-naïve

relative to convalescent vaccinees, express low levels of markers of homeostatic

proliferation and tissue homing. (A) Shown are histograms of the expression levels of the

alpha chain of the IL7 receptor (CD127), the chemokine receptor CXCR4, and the lymph node

homing receptor CCR7, among clusters A1 or A3, the former of was enriched in infection-naïve

relative to convalescent individuals after vaccination. Data were concatenated from all clustered

cells. **(B)** Relative expression levels, as depicted by normalized mean signal intensity (MSI), of

CD127, CXCR4, and CCR7 among all specimens of spike-specific CD4+ T cells from infection-

naïve and convalescent individuals, after the second vaccination dose. *p < 0.05, **p < 0.01, ns

= non-significant, as determined using student's t-tests and corrected for multiple testing (n =

39) using the Holm-Sidak method.

Figure S5. Phenotypic features of spike-specific CD8+ T cells from vaccinated,

convalescent individuals are unique and differ from those of their CD4+ T cell

counterparts. (A-C) MDS (*A*) or tSNE (*B, C*) plots demonstrating phenotypic similarities

between spike-specific CD8+ T cells responding to spike from the ancestral, B.1.1.7, or B.1.351

strains. Data are displayed in a format similar to that for CD4+ T cells presented in Fig. 2A-C.

(D) MDS plot depicting specimens of spike-specific CD8+ T cells in infection-naïve and

convalescent individuals after second vaccination dose. **(E)** tSNE contour heatmaps depicting

spike-specific CD8+ T cells from infection-naïve and convalescent individuals, after the second

vaccination dose. Cell densities are represented by color. **(F)** tSNE dot plot of spike-specific

CD8+ T cells from infection-naïve and convalescent individuals after second vaccination dose.

(G) The distribution of spike-specific cells among the main canonical CD8+ T cell subsets (Tn,

Tscm, Temra, Tcm, Tem, Ttm) is similar in infection-naïve vs. convalescent individuals after

second vaccination dose. **(H)** T cell subsets that were differentially enriched in infection-naïve

vs. convalescent individuals among spike-specific CD4+ T cells after second vaccination dose

(Fig. 6C) are not differentially enriched among spike-specific CD8+ T cells. Shown are the

proportions of cells that are CD127+CD57-, CXCR4+CD69+, or CCR7+CD62L+ cells among

spike-specific CD8+ T cells as determined by manual gating. **(I)** Cells co-expressing CD27 and

CD38, and CTLA4 and CD137, are elevated among spike-specific CD8+ T cells from

vaccinated convalescent individuals relative to vaccinated infection-naïve individuals. *p < 0.05,

**p < 0.01 as determined by student's t-test.

SUPPLEMENTARY TABLES

Table S1. Participant Characteristics

Patient ID	Gender	Age	Prior Infection Status	Vaccine	Days post PCR+ test at pre-vaccination timepoint	Days post vaccine dose #1	Days post vaccine dose #2
PID4101	Female	45	Uninfected	Pfizer/BioNT	NA	13	12
PID4109	Male	33	Uninfected	Pfizer/BioNT	NA	12	33
PID4197	Female	76	Uninfected	Pfizer/BioNT	NA	14	13
PID4198	Male	79	Uninfected	Moderna	NA	18	10
PID4199	Female	32	Uninfected	Pfizer/BioNT	NA	14	10
PID4104	Female	33	Convalescent	Moderna	212	14	14
PID4108	Female	20	Convalescent	Pfizer/BioNT	226	13	38
PID4112	Female	59	Convalescent	Moderna	254	16	13
PID4114	Female	46	Convalescent	Moderna	216	16	50
PID4117	Female	51	Convalescent	Pfizer/BioNT	82	16	6
PID4118	Female	39	Convalescent	Pfizer/BioNT	173	18	28

867 **Table S2. List of CyTOF antibodies used in study**. Antibodies were either purchased from
868 the indicated vendor or prepared in-house using commercially available MaxPAR conjugation
869 kits per manufacturer's instructions (Fluidigm).
870

Antigen Target	Clone	Elemental Isotope	Vendor
HLADR	TÜ36	Qdot (112Cd)	Thermofisher
RORγt*	AFKJS-9	115 In	In-house
CD49d (α4)	9F10	141Pr	Fluidigm
CTLA4*	14D3	142Nd	In-house
NFAT*	D43B1	143Nd	Fluidigm
CCR5	NP6G4	144Nd	Fluidigm
CD137	4B4-1	145Nd	In-house
CD95	BX2	146Nd	In-house
CD7	CD76B7	147Sm	Fluidigm
ICOS	C398.4A	148Nd	Fluidigm
Tbet*	4B10	149Sm	In-house
IL4*	MP4-25D2	150Nd	In-house
CD2	TS1/8	151Eu	Fluidigm
IL17*	BL168	152Sm	In-house
CD62L	DREG56	153Eu	Fluidigm
TIGIT	MBSA43	154Sm	Fluidigm
CCR6	11A9	155Gd	In-house
IL6*	MQ2-13A5	156 Gd	In-house
CD8	RPA-T8	157Gd	In-house
CD19	HIB19	157Gd	In-house
CD14	M5E2	157Gd	In-house
OX40	ACT35	158Gd	Fluidigm
CCR7	G043H7	159Tb	Fluidigm
CD28	CD28.2	160Gd	Fluidigm
CD45RO	UCHL1	161Dy	In-house
CD69	FN50	162Dy	Fluidigm
CRTH2	BM16	163Dy	Fluidigm
PD-1	EH12.1	164Dy	In-house
CD127	A019D5	165Ho	Fluidigm
CXCR5	RF8B2	166Er	In-house
CD27	L128	167Er	Fluidigm
IFNγ*	B27	168Er	Fluidigm
CD45RA	HI100	169Tm	Fluidigm
CD3	UCHT1	170Er	Fluidigm
CD57	HNK-1	171Yb	In-house
CD38	HIT2	172Yb	Fluidigm
α4β7	Act1	173Yb	In-house
CD4	SK3	174Yb	Fluidigm
CXCR4	12G5	175Lu	Fluidigm
CD25	M-A251	176Yb	In-house
CD161	NKR-P1A	209 Bi	In-house

871 *Intracellular antibodies
872

REFERENCES

1.   Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, Hengartner N, Giorgi EE, Bhattacharya T, Foley B, et al. (2020). Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. Cell 182, 812-827 e819. 2020/07/23.

2.   Weissman D, Alameh MG, de Silva T, Collini P, Hornsby H, Brown R, LaBranche CC, Edwards RJ, Sutherland L, Santra S, et al. (2021). D614G Spike Mutation Increases SARS CoV-2 Susceptibility to Neutralization. Cell Host Microbe 29, 23-31 e24. 2020/12/12.

3.   Plante JA, Mitchell BM, Plante KS, Debbink K, Weaver SC and Menachery VD. (2021). The variant gambit: COVID-19's next move. Cell Host Microbe 29, 508-515. 2021/04/01.

4.   Callaway E. (2021). Delta coronavirus variant: scientists brace for impact. Nature 595, 17-18. 2021/06/24.

5.   Davies NG, Jarvis CI, Group CC-W, Edmunds WJ, Jewell NP, Diaz-Ordaz K and Keogh RH. (2021). Increased mortality in community-tested cases of SARS-CoV-2 lineage B.1.1.7. Nature. 2021/03/17.

6.   Wang P, Nair MS, Liu L, Iketani S, Luo Y, Guo Y, Wang M, Yu J, Zhang B, Kwong PD, et al. (2021). Antibody Resistance of SARS-CoV-2 Variants B.1.351 and B.1.1.7. Nature. 2021/03/09.

7.   Collier DA, De Marco A, Ferreira I, Meng B, Datir R, Walls AC, Kemp SS, Bassi J, Pinto D, Fregni CS, et al. (2021). Sensitivity of SARS-CoV-2 B.1.1.7 to mRNA vaccine-elicited antibodies. Nature. 2021/03/12.

8.   Muik A, Wallisch AK, Sanger B, Swanson KA, Muhl J, Chen W, Cai H, Maurus D, Sarkar R, Tureci O, et al. (2021). Neutralization of SARS-CoV-2 lineage B.1.1.7 pseudovirus by BNT162b2 vaccine-elicited human sera. Science 371, 1152-1153. 2021/01/31.

9.   Garcia-Beltran WF, Lam EC, St Denis K, Nitido AD, Garcia ZH, Hauser BM, Feldman J,

Pavlovic MN, Gregory DJ, Poznansky MC, et al. (2021). Multiple SARS-CoV-2 variants escape

neutralization by vaccine-induced humoral immunity. Cell. 2021/03/21.

10.   Stamatatos L, Czartoski J, Wan YH, Homad LJ, Rubin V, Glantz H, Neradilek M, Seydoux

E, Jennewein MF, MacCamy AJ, et al. (2021). mRNA vaccination boosts cross-variant

neutralizing antibodies elicited by SARS-CoV-2 infection. Science. 2021/03/27.

11.   Cele S, Gazy I, Jackson L, Hwa SH, Tegally H, Lustig G, Giandhari J, Pillay S, Wilkinson

E, Naidoo Y, et al. (2021). Escape of SARS-CoV-2 501Y.V2 from neutralization by convalescent

plasma. Nature. 2021/03/30.

12.   Hoffmann M, Arora P, Gross R, Seidel A, Hornich BF, Hahn AS, Kruger N, Graichen L,

Hofmann-Winkler H, Kempf A, et al. (2021). SARS-CoV-2 variants B.1.351 and P.1 escape from

neutralizing antibodies. Cell. 2021/04/02.

13.   Planas D, Bruel T, Grzelak L, Guivel-Benhassine F, Staropoli I, Porrot F, Planchais C,

Buchrieser J, Rajah MM, Bishop E, et al. (2021). Sensitivity of infectious SARS-CoV-2 B.1.1.7

and B.1.351 variants to neutralizing antibodies. Nat Med. 2021/03/28.

14.   Edara VV, Norwood C, Floyd K, Lai L, Davis-Gardner ME, Hudson WH, Mantus G, Nyhoff

LE, Adelman MW, Fineman R, et al. (2021). Infection- and vaccine-induced antibody binding and

neutralization of the B.1.351 SARS-CoV-2 variant. Cell Host Microbe 29, 516-521 e513.

2021/04/03.

15.   Kuzmina A, Khalaila Y, Voloshin O, Keren-Naus A, Boehm-Cohen L, Raviv Y, Shemer-

Avni Y, Rosenberg E and Taube R. (2021). SARS-CoV-2 spike variants exhibit differential

infectivity and neutralization resistance to convalescent or post-vaccination sera. Cell Host

Microbe 29, 522-528 e522. 2021/04/01.

16.   Skelly DT, Harding AC, Gilbert-Jaramillo J, Knight ML, Longet S, Brown A, Adele S,

Adland E, Brown H, Team ML, et al. (2021). Vaccine-induced immunity provides more robust

heterotypic immunity than natural infection to emerging SARS-CoV-2 variants of concern.

Research Square.

17.   Tarke A, Sidney J, Methot N, Zhang Y, Dan JM, Goodwin B, Rubiro P, Sutherland A, da

Silva Antunes R, Frazier A, et al. (2021). Negligible impact of SARS-CoV-2 variants on CD4 (+)

and CD8 (+) T cell reactivity in COVID-19 exposed donors and vaccinees. bioRxiv. 2021/03/11.

18.   Redd AD, Nardin A, Kared H, Bloch EM, Pekosz A, Laeyendecker O, Abel B, Fehlings M,

Quinn TC and Tobian AA. (2021). CD8+ T cell responses in COVID-19 convalescent individuals

target conserved epitopes from multiple prominent SARS-CoV-2 circulating variants. medRxiv.

2021/02/18.

19.   Geers D, Shamier MC, Bogers S, den Hartog G, Gommers L, Nieuwkoop NN, Schmitz

KS, Rijsbergen LC, van Osch JAT, Dijkhuizen E, et al. (2021). SARS-CoV-2 variants of concern

partially escape humoral but not T-cell responses in COVID-19 convalescent donors and

vaccinees. Sci Immunol 6. 2021/05/27.

20.   Woldemeskel BA, Garliss CC and Blankson JN. (2021). SARS-CoV-2 mRNA vaccines

induce broad CD4+ T cell responses that recognize SARS-CoV-2 variants and HCoV-NL63. J

Clin Invest 131. 2021/04/07.

21.   Stankov MV, Cossmann A, Bonifacius A, Dopfer-Jablonka A, Ramos GM, Godecke N,

Scharff AZ, Happle C, Boeck AL, Tran AT, et al. (2021). Humoral and cellular immune responses

against SARS-CoV-2 variants and human coronaviruses after single BNT162b2 vaccination. Clin

Infect Dis. 2021/06/17.

22.   Tauzin A, Nayrac M, Benlarbi M, Gong SY, Gasser R, Beaudoin-Bussieres G, Brassard

N, Laumaea A, Vezina D, Prevost J, et al. (2021). A single dose of the SARS-CoV-2 vaccine

BNT162b2 elicits Fc-mediated antibody effector functions and T cell responses. Cell Host Microbe

29, 1137-1150 e1136. 2021/06/17.

23.   Chen G, Wu D, Guo W, Cao Y, Huang D, Wang H, Wang T, Zhang X, Chen H, Yu H, et

al. (2020). Clinical and immunological features of severe and moderate coronavirus disease 2019.

J Clin Invest 130, 2620-2629. 2020/03/29.

24.   Woodruff MC, Ramonell RP, Nguyen DC, Cashman KS, Saini AS, Haddad NS, Ley AM,

Kyu S, Howell JC, Ozturk T, et al. (2020). Extrafollicular B cell responses correlate with

neutralizing antibodies and morbidity in COVID-19. Nat Immunol 21, 1506-1516. 2020/10/09.

25.   Rydyznski Moderbacher C, Ramirez SI, Dan JM, Grifoni A, Hastie KM, Weiskopf D,

Belanger S, Abbott RK, Kim C, Choi J, et al. (2020). Antigen-Specific Adaptive Immunity to SARS-

CoV-2 in Acute COVID-19 and Associations with Age and Disease Severity. Cell 183, 996-1012

e1019. 2020/10/05.

26.   Neidleman J, Luo X, George AF, McGregor M, Yang J, Yun C, Murray V, Gill G, Greene

WC, Vasquez J, et al. (2021). Distinctive features of SARS-CoV-2-specific T cells predict recovery

from severe COVID-19. Cell Rep 36, 109414. 2021/07/15.

27.   Dan JM, Mateus J, Kato Y, Hastie KM, Yu ED, Faliti CE, Grifoni A, Ramirez SI, Haupt S,

Frazier A, et al. (2021). Immunological memory to SARS-CoV-2 assessed for up to 8 months after

infection. Science 371. 2021/01/08.

28.   Neidleman J, Luo X, Frouard J, Xie G, Gill G, Stein ES, McGregor M, Ma T, George AF,

Kosters A, et al. (2020). SARS-CoV-2-Specific T Cells Exhibit Phenotypic Features of Helper

Function, Lack of Terminal Differentiation, and High Proliferation Potential. Cell Rep Med 1,

100081. 2020/08/26.

29.   Soresina A, Moratto D, Chiarini M, Paolillo C, Baresi G, Foca E, Bezzi M, Baronio B,

Giacomelli M and Badolato R. (2020). Two X-linked agammaglobulinemia patients develop

pneumonia as COVID-19 manifestation but recover. Pediatr Allergy Immunol. 2020/04/23.

30.   Evavold BD and Allen PM. (1991). Separation of IL-4 production from Th cell proliferation

by an altered T cell receptor ligand. Science 252, 1308-1310. 1991/05/31.

31.   Sloan-Lancaster J and Allen PM. (1996). Altered peptide ligand-induced partial T cell

activation: molecular mechanisms and role in T cell biology. Annu Rev Immunol 14, 1-27.

1996/01/01.

32.   Goel RR, Apostolidis SA, Painter MM, Mathew D, Pattekar A, Kuthuru O, Gouma S, Hicks

P, Meng W, Rosenfeld AM, et al. (2021). Distinct antibody and memory B cell responses in SARS-

CoV-2 naive and recovered individuals following mRNA vaccination. Sci Immunol 6. 2021/04/17.

33.   Ebinger JE, Fert-Bober J, Printsev I, Wu M, Sun N, Figueiredo JC, Eyk JEV, Braun JG,

Cheng S and Sobhani K. (2021). Prior COVID-19 Infection and Antibody Response to Single

Versus Double Dose mRNA SARS-CoV-2 Vaccination. medRxiv. 2021/03/04.

34.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W and Smyth GK. (2015). limma powers

differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res

43, e47. 2015/01/22.

35.   Kaech SM, Tan JT, Wherry EJ, Konieczny BT, Surh CD and Ahmed R. (2003). Selective

expression of the interleukin 7 receptor identifies effector CD8 T cells that give rise to long-lived

memory cells. Nat Immunol 4, 1191-1198.

36.   Gupta RK. (2021). Will SARS-CoV-2 variants of concern affect the promise of vaccines?

Nat Rev Immunol. 2021/05/01.

37.   Madhi SA, Baillie V, Cutland CL, Voysey M, Koen AL, Fairlie L, Padayachee SD, Dheda

K, Barnabas SL, Bhorat QE, et al. (2021). Efficacy of the ChAdOx1 nCoV-19 Covid-19 Vaccine

against the B.1.351 Variant. N Engl J Med. 2021/03/17.

38.   Abu-Raddad LJ, Chemaitelly H, Butt AA and National Study Group for C-V. (2021).

Effectiveness of the BNT162b2 Covid-19 Vaccine against the B.1.1.7 and B.1.351 Variants. N

Engl J Med. 2021/05/06.

39.   Klenerman P and Zinkernagel RM. (1998). Original antigenic sin impairs cytotoxic T

lymphocyte responses to viruses bearing variant epitopes. Nature 394, 482-485. 1998/08/11.

40.   Lederer K, Castano D, Gomez Atria D, Oguin TH, 3rd, Wang S, Manzoni TB, Muramatsu

H, Hogan MJ, Amanat F, Cherubin P, et al. (2020). SARS-CoV-2 mRNA Vaccines Foster Potent

Antigen-Specific Germinal Center Responses Associated with Neutralizing Antibody Generation.

Immunity 53, 1281-1295 e1285. 2020/12/10.

41.   Bacchus-Souffan C, Fitch M, Symons J, Abdel-Mohsen M, Reeves DB, Hoh R, Stone M,

Hiatt J, Kim P, Chopra A, et al. (2021). Relationship between CD4 T cell turnover, cellular

differentiation and HIV persistence during ART. PLoS Pathog 17, e1009214. 2021/01/20.

42.   Ma T, Ryu H, McGregor M, Babcock B, Neidleman J, Xie G, George AF, Frouard J, Murray

V, Gill G, et al. (2021). Protracted yet coordinated differentiation of long-lived SARS-CoV-2-

specific CD8+ T cells during COVID-19 convalescence. bioRxiv 2021.04.28.441880.

43.   Mamazhakypov A, Viswanathan G, Lawrie A, Schermuly RT and Rajagopal S. (2019).

The role of chemokines and chemokine receptors in pulmonary arterial hypertension. Br J

Pharmacol 178, 72-89. 2019/08/11.

44.   Newell EW, Sigal N, Nair N, Kidd BA, Greenberg HB and Davis MM. (2013). Combinatorial

tetramer staining and mass cytometry analysis facilitate T-cell epitope mapping and

characterization. Nat Biotechnol 31, 623-629. 2013/06/12.

45.   Ma T, Luo X, George AF, Mukherjee G, Sen N, Spitzer TL, Giudice LC, Greene WC and

Roan NR. (2020). HIV efficiently infects T cells from the endometrium and remodels them to

promote systemic viral spread. Elife 9, e55487. 2020/05/27.

46.   Cavrois M, Banerjee T, Mukherjee G, Raman N, Hussien R, Rodriguez BA, Vasquez J,

Spitzer MH, Lazarus NH, Jones JJ, et al. (2017). Mass Cytometric Analysis of HIV Entry,

Replication, and Remodeling in Tissue CD4+ T Cells. Cell Rep 20, 984-998.

47.   Neidleman J, Luo X, Frouard J, Xie G, Hsiao F, Ma T, Morcilla V, Lee A, Telwatte S,

Thomas R, et al. (2020). Phenotypic analysis of the unstimulated in vivo HIV CD4 T cell reservoir.

Elife 9, e55487. 2020/09/30.

48.   Xie G, Luo X, Ma T, Frouard J, Neidleman J, Hoh R, Deeks SG, Greene WC and Roan

NR. (2021). Characterization of HIV-induced remodeling reveals differences in infection

susceptibility of memory CD4(+) T cell subsets in vivo. Cell Rep 35, 109038. 2021/04/29.

49.   Nowicka M, Krieg C, Crowell HL, Weber LM, Hartmann FJ, Guglietta S, Becher B,

Levesque MP and Robinson MD. (2017). CyTOF workflow: differential discovery in high-

throughput high-dimensional cytometry datasets. F1000Res 6, 748. 2017/05/26.

50.   Van Gassen S, Callebaut B, Van Helden MJ, Lambrecht BN, Demeester P, Dhaene T and

Saeys Y. (2015). FlowSOM: Using self-organizing maps for visualization and interpretation of

cytometry data. Cytometry A 87, 636-645. 2015/01/13.

51.   Wilkerson MD and Hayes DN. (2010). ConsensusClusterPlus: a class discovery tool with

confidence assessments and item tracking. Bioinformatics 26, 1572-1573. 2010/04/30.

1032

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

A.

B.

Figure S2

Figure S3

A.

B.

Relative Expression

Cluster C2

All responding cells

Relative Expression

Cluster C5

All responding cells

Figure S4

A.

Cluster A1   Cluster A3

Density

Relative Expression

Cluster A1   Cluster A3

All responding cells   All responding cells

B.

CD127

mean

*

CXCR4

mean

ns

CCR7

mean

**

Infection-naïve   Convalescent

Figure S5