Developmental single-cell transcriptomics of hypothalamic POMC progenitors

reveal the genetic trajectories of multiple neuropeptidergic phenotypes

Hui Yu[1*], Marcelo Rubinstein[1,2,3*], Malcolm J Low[1*]

[1]Department of Molecular and Integrative Physiology, University of Michigan Medical School, Ann Arbor, MI 48109, USA

[2]Instituto de Investigaciones en Ingeniería Genética y Biología Molecular, Consejo Nacional de Investigaciones Científicas y Técnicas, 1428 Buenos Aires, Argentina

[3]Departamento de Fisiología, Biología Molecular y Celular, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, 1428 Buenos Aires, Argentina

* Co-corresponding Authors

Address correspondence to:

Malcolm J Low, MD, PhD, Department of Molecular & Integrative Physiology, University of Michigan Medical School, North Campus Research Complex, Bldg 25, Rm.25-3680, 2800 Plymouth Rd., Ann Arbor, MI 48109-2800. Email: mjlow@umich.edu

Marcelo Rubinstein, PhD, Instituto de Investigaciones en Ingeniería Genética y Biología Molecular, Consejo Nacional de Investigaciones Científicas y Técnicas, 1428 Buenos Aires, Argentina. Email: mrubins@dna.uba.ar

Hui Yu, PhD, Department of Molecular & Integrative Physiology, University of Michigan Medical School, North Campus Research Complex, Bldg 25, Rm.25-3634, 2800 Plymouth Rd., Ann Arbor, MI 48109-2800. Email: huiyuz@umich.edu

Abstract (159 words)

Proopiomelanocortin (POMC) neurons of the hypothalamic arcuate nucleus are essential to regulate food intake and energy balance. However, the ontogenetic transcriptional programs that specify the identity and functioning of these neurons are poorly understood. Here, we use scRNAseq to define the transcriptomes characterizing POMC progenitors in the developing hypothalamus and TRAP-seq to analyze the subsequent translatomes of mature POMC neurons. Our data show that *Pomc*-expressing neurons originate from multiple developmental pathways expressing distinct combinations of transcription factors. The predominant cluster, featured by high levels of *Pomc* and *Prdm12* transcripts represents the precursor of canonical arcuate POMC neurons. Additional clusters of progenitors expressing lower levels of *Pomc* mature into different neuronal phenotypes characterized by distinct combinations of transcription factors, neuropeptides, processing enzymes, cell surface and nuclear receptors. We conclude that the genetic programs specifying the identity and differentiation of arcuate POMC neurons are diverse and generate a heterogeneous repertoire of neuronal phenotypes early in development that continue to evolve postnatally.

Introduction

Proopiomelanocortin (POMC)-expressing neurons located in the arcuate nucleus of the hypothalamus play an essential role in the regulation of food intake by maintaining a melanocortin-dependent anorexigenic tone. The critical importance of central melanocortins, specifically in arcuate neurons, is apparent in mice lacking *Pomc* that develop hyperphagia and early-onset extreme obesity(Bumaschny et al., 2012). This phenotype is closely mirrored in humans carrying biallelic null-mutations in *POMC*(Krude et al., 1998). In addition, arcuate POMC neurons release the *Pomc*-encoded opioid peptide *β*-endorphin, which is critical in mediating normal food intake and stress-induced analgesia(Parikh et al., 2011).

Postmitotic POMC precursors acquire their phenotypic identity in the developing hypothalamus days earlier than any other peptidergic neurons born in the presumptive arcuate nucleus (eg. NPY/AGRP, GHRH, SST, KISSPEPTIN). *Pomc* mRNA is initially observed in the tuberal portion of the prospective mouse hypothalamus of E10.5 embryos, following the concurrent advent of the essential transcription factors (TFs) ISL1(Nasif et al., 2015), NKX2-1(Orquera et al., 2019) and PRDM12(Hael, Rojo, Orquera, Low, & Rubinstein, 2020). In fact, the early ablation of either *Isl1*, *Nkx2-1* or *Prdm12* disrupts the onset of hypothalamic *Pomc* expression in conditional mutant mouse embryos(Hael et al., 2020; Nasif et al., 2015; Orquera et al., 2019). Given that the number of arcuate neurons coexpressing *Nkx2-1*, *Isl1* and *Prdm12* greatly exceeds that of POMC neurons in this area, it is expected that additional and still unknown TFs integrate the complete genetic program that specifies the early identity of POMC neurons.

Other molecular markers of arcuate POMC neurons such as transporters, receptors, channels and co-transmitters exhibit great variability indicating that a heterogeneous pool of diverse subpopulations of these neurons are involved in multiple circuits and functions. In fact,

previous studies have suggested that subsets of arcuate *Pomc*-expressing neurons may act as

precursors of terminally differentiated neurons of alternative neuropeptidergic phenotypes such

as NPY/AGRP and KISSPEPTIN/NEUROKININ-B/DYNORPHIN (KNDY) neurons in the adult

hypothalamus(Padilla, Carmody, & Zeltser, 2010; Sanz et al., 2015). Initial attempts to

characterize the transcriptome of adult POMC neurons at the single-cell level confirmed the

heterogeneous nature of arcuate POMC neurons and revealed the existence of distinct clusters

including those co-expressing the aforementioned neuropeptide genes(Campbell et al., 2017; R.

Chen, Wu, Jiang, & Zhang, 2017; Huisman et al., 2020; Lam et al., 2017). However, the

molecular signatures of the precursors of the various hypothalamic *Pomc*-expressing lineages

and their transition into mature neurons within the postnatal Arc are still lacking.

Here, we track the origin and maturation of hypothalamic POMC precursors in the

developing and early postnatal hypothalamus by performing single-cell RNA-seq transcriptomics

of fluorescently labeled cells taken from *Pomc-TdDiscomaRed-Sv40PolyA* (*Pomc-TdDsRed*)

transgenic mice at different embryonic (E11.5, E13.5, E15.5 and E17.5) and early postnatal (P5

and P12) ages. Our data suggested that the POMC precursors arise from multiple different

transcriptional routes. Some of these precursors further develop into mature POMC neurons

while others follow diverse neuropeptidergic trajectories indicating that a heterogeneous

population of *Pomc*-expressing neurons is established early in the developing arcuate nucleus**.**

Combining translating ribosome affinity purification with RNA-sequencing (TRAP-seq) of *Pomc-*

eGFPL10a transgenic mice at P12 and P60, we defined transcriptional programs that function

distinctly in early embryonic POMC precursors and later in mature POMC neurons. These

datasets provide valuable resources for comprehensively understanding the genetic

development of POMC neurons.

Results

Cluster analysis of gene expression in Pomc positive hypothalamic cells integrated across six development ages

We captured the single cell transcriptomes of hypothalamic *Pomc*-expressing cells at four key embryonic days (E11.5, E13.5, E15.5 and E17.5) and two early postnatal days (P5 and P12) when the maturing neurons already develop axons and innervate key target nuclei distal to the Arc(Bouret, Bates, Chen, Myers, & Simerly, 2012; Bouret, Draper, & Simerly, 2004a, 2004b). Medial basal hypothalami from mice that express the reporter transgene *Pomc-TdDsRed* selectively in POMC cells were pooled from multiple embryos or postnatal pups at the indicated ages. Dissociated single cell suspensions were sorted for DsRed fluorescence and used for library preparation with the 10X genomics platform (Figure 1A). After a rigorous screening procedure to remove substandard scRNA sequencing data (see Methods), we used a combination of computational platforms to analyze the transcriptomes of 13,962 cells that had *Pomc* counts of ≥ 1 unique molecular identifiers (UMI).

A Seurat analysis of all cells integrated across the six-time points revealed 11 distinct cell clusters projected onto a UMAP plot (Figure 1B). The same data are represented in a heat map corresponding to the top 30 expressed genes per cluster (Supplementary Figure 1). Clusters 1-8 (10,828 cells) were all highly enriched for a panel of three neuronal marker gene transcripts. The embedded table in Figure 1B lists the percentages of neurons of each cluster at each developmental age. The other three clusters (3,134 cells) were enriched in transcripts characteristic of either microglia, vascular and leptomeningeal cells (VLMC) or astrocytes (Figure 1C). GO ontology analyses for biological processes further confirmed the neuronal or non-neuronal identities of the 11 individual clusters (data not shown).

By definition, all cells in the neuronal and non-neuronal clusters contained *Pomc* transcripts (UMI ≥ 1), however there were significant differences in the average expression

levels per cluster, ranging from 1.3 to 33.8 [exp(*log Normalized* counts) -1)]. Nomenclature of

the 8 neuronal clusters is based on a combination of relatively high (>20), medium (5-20) or low

(<5) *Pomc* levels and high expression of one differential key feature gene as follows: no. 1,

Pomc^{high}/Prdm12; no. 2 *Pomc^{med}/Ebf1*; no. 3, *Pomc^{med}/Nr4a2*; no. 4, *Pomc^{low}/Otp*; no. 5,

Pomc^{low/}Tac2; no. 6, *Pomc^{low}/Dlx5*; no. 7, *Pomc^{low}/Prdm13*; and no. 8, *Pomc^{low}/Pitx2*. The cluster

distribution and relative expression of those feature genes are represented by individual UMAP

plots (Figure 1D). Average expression levels of all genes, including *Pomc*, in the 11 clusters and

complete lists of all feature genes that define the individual clusters are presented in

Supplemental Tables 1 and 2, respectively.

In order to validate our experimental approach of isolating *Pomc*-expressing cells from

the developing hypothalamus by FACS for the transgenic surrogate marker *Pomc-TdDsRed*, we

performed an independent unsupervised cluster analysis of the transcriptomes from the 5,724

cells that had *DsRed* counts of ≥ 1 UMI (Supplemental Figure 2A-C and Supplemental Table 3).

A projection analysis (Supplemental Figure 2D-F) of the differentially expressed feature genes

defined 9 DsRed clusters (Supplemental Table 4) which closely corresponded to 9 of the 11

clusters based on *Pomc* counts of ≥ 1 UMI (Supplemental Table 2). The two exceptions were

the smallest clusters 7. *Pomc^{low}/Prdm13* and Astrocytes (Supplemental Figure 2D-F).

Furthermore, the percentages of cells at each developmental age from the corresponding

neuronal clusters were remarkably similar in the *Pomc* UMI ≥ 1 data set compared to the *DsRed*

UMI ≥ 1 data set (compare Figure 1B to Supplemental Figure 3A). The different population of

DsRed fluorescent sorted cells showing UMI ≥ 1 for *Pomc* (13,962) or *DsRed* (5,724) may be

due to differential half-lives of the fluorescent protein DSRED used during cell sorting and its

transcript used for barcoding the RNAseq libraries.

Temporal gene expression analysis of the eight neuronal clusters at individual developmental stages

Next, we analyzed temporal gene expression patterns for each of the 8 neuronal clusters at the 6 distinct developmental ages (Supplemental Table 5). The progressive changes in feature genes across time are responsible for the observed within cluster heterogeneity on the heat map in Supplemental Figure 1.

Cluster 1. *Pomc^high^/Prdm12* cells were the most abundant among the eight neuronal clusters and consistently had the highest levels of *Pomc* expression at all six developmental ages. We consider them to be the direct precursors of canonical POMC neurons in the adult hypothalamus (Figure 2A-2C). Other feature genes included a set of four TFs known to be critical for the initiation and/or maintenance of Arc *Pomc* expression, *Isl1, Nkx2-1, Prdm12* and *Tbx3(Hael et al., 2020; Nasif et al., 2015; Orquera et al., 2019; Quarta et al., 2019)* (Figure 2A). They were expressed in similar developmental patterns to each other, with the exception of *Tbx3* that was significantly expressed only from day E15.5. Notably, there was a lack of the pituitary-specific *Tbx19* TF, confirming that no pituitary tissue was included in the hypothalamic dissections. Two homologues of the *Drosophila sine oculus* homeobox gene, *Six3* and *Six6,* which have not been described previously in the context of Arc POMC neuron development, were also expressed robustly at all six time points (Figure 2A). Co-expression of their mRNAs with *Pomc* by RNAScope at ages E12.5 and E16.5 confirmed the scRNAseq analysis, although the two TFs were expressed in additional hypothalamic domains and the pituitary gland (Figure 2C). *Vim1, Sox1, 2, 3 and 14,* and *Myc*, whose expression is typically associated with neuronal precursors or immature neurons, tended to have higher expression at the earlier embryonic ages (Figure 2A). The nuclear receptor gene *Nr5a1*, also known as steroidogenic factor 1 (*Sf1*), was a key feature gene expressed almost exclusively at ages E11.5. (Figure 2A), but only minimally expressed at later embryonic time points when NR5A1 is critically involved in the

development of the hypothalamic ventromedial nucleus (VMN)(Ikeda, Luo, Abbud, Nilson, &

Parker, 1995). This differential spatio-temporal presence of NR5A1 within or outside POMC

neurons was confirmed by immunofluorescence on hypothalamic sections taken from E11.5 and

E17.5 mouse embryos (Figure 2C). Interestingly, *Nr5a1*, *Nr0b1* (*Dax1*) and *Shh*, transcripts

were also present in Cluster 1 at the two earliest ages, and are factors known to interact with

the *Wnt/*β-catenin signaling pathway in early cell fate determination in the adrenal and pituitary

glands(Luo, Ikeda, & Parker, 1994; Meeks, Weiss, & Jameson, 2003; Mizusaki et al., 2003). A

combination of pseudotime and gene ontology analyses for the top expressed genes

(Log2FoldChange >0.25, *P*<0.05) in cluster1 revealed that the genes expressed at the

beginning of pseudotime function in the organization of cell projections, axonogenesis and the

regulation cellular component size, whereas genes expressed at the middle and end of the

pseudotime function are related to neuron maturation and neuron-specified functions such as

the formation of myelin sheath and regulation of synaptic vesicle cycling (Figure 2B).

Cluster 2. *Pomc^{med}/Ebf1* neurons made up the second highest abundance cluster of

POMC neurons, particularly at E11.5 and E13.5. After these time points, the fraction of all

neurons constituting cluster 2 dropped by approximately two thirds from ages E15.5 through

P12. In contrast, the relative percentage of neurons in clusters 3-8 increased over time with

peaks at different embryonic ages (Figure 1B). Cluster 2 neurons, on average, had medium

levels of *Pomc* transcripts that were divided distinctly from their lowest at ages E11.5-E13.5 to

highest at later embryonic and postnatal ages (Figure 2D; Supplemental Table 5). Major cluster

2 feature markers included members of the non-basic helix-loop-helix early B-cell TF family EBF

(Supplemental Table 2). *Ebf1* is typically co-expressed with either *Ebf2* or *Ebf3* and pairs of

these factors were shared not only in cluster 2, but also in clusters 3 and 8 (Supplemental

Figure 4A and D). Dual RNAScope *in situ* hybridization showed overlap of *Ebf1* and *Pomc*

mRNAs in hypothalamic neurons at an early embryonic stage (E13.5), whereas at later

embryonic stage (E17.5), POMC neurons had barely detectable *Ebf1* mRNAs (Figure 2F). A

similar expression pattern was found with another cluster 2 featured gene, *Pou4f1 (POU Class 4 Homeobox 1)*. Dual RNAScope *in situ* hybridization confirmed the overlap of *Pou4f1* and *Pomc* mRNAs at E13.5, but not at E17.5 (Figure 2F). These results, together, suggest that *Pomc* mRNA levels peak only after expression of both *Ebf1* and *Pou4f1* decrease. Two other prominent feature genes in cluster 2 were *Crabp1* and *Crabp2*, which encode cellular retinoic acid binding proteins. A majority of the marker genes for cluster 2 are predicted to be expressed at the beginning of pseudotime and the main functions are associated with the maintenance of basic cell biological processes such as nucleosome organization, axonogenesis, microtubule transport, and intracellular transport (Figure 2E). The genes ordered in the middle and end of pseudotime function for the regulation of synapse structure and activities.

Visual examination of the UMAP plots in Figure 1D suggested that cluster 2 was composed of two compartments, one with high and one with low *Pomc* abundance. Unsupervised reclustering of the differentially expressed genes from integrated cluster 2 confirmed the existence of two subclusters (Figure 2G). Subcluster 2-1 contained two-thirds of the neurons, which were *POMC^{high}* and present only from ages E15.5-P12. Subcluster 2-2 contained the other one-third of cluster 2 neurons, which in contrast were *Pomc^{low}* and detected at ages E11.5 and E13.5. Consistently, known activating TFs for *Pomc* expression such as *Nkx2-1*, *Isl1* and *Tbx3* were limited to subcluster 2-1 while the *Ebf* and *Crabp* transcripts were characteristic of subcluster 2-2 (Supplemental Table 6 and Supplemental Table 7). Conspicuously absent from both subclusters was *Prdm12* mRNA.

Cluster 3. *Pomc^{med}/Nr4a2* neurons displayed high expression of its major feature gene *Nr4a2* (Nuclear Receptor Subfamily 4 Group A Member 2) (Supplemental Figure 4A). *Nr4a2* mRNA was also expressed at relatively high levels in clusters 2. *Pomc^{med}/Ebf1* and 8. *Pomc^{low}/Pitx2*. Originally known as *Nurr1*, *Nr4a2* has been implicated in the regulation of *Pomc* expression in the pituitary corticotropic cell line AtT20(Kovalovsky et al., 2002). Furthermore,

Nr4a2 is a pioneer transcriptional regulator important for the differentiation and maintenance of

meso-diencephalic dopaminergic (mdDA) neurons during development(Perlmann & Wallen-

Mackenzie, 2004). It is crucial for expression of a set of genes including *Slc6a3* (plasma

membrane dopamine transporter, DAT), *Slc18a2* (synaptic vesicular monoamine transporter,

VMAT), *Th* (Tyrosine hydroxylase), *Foxa1* and *Foxa2* (forkhead family TFs) and *Drd2*

(dopamine receptor 2) that together define the dopaminergic neuronal phenotype(Hong et al.,

2014; Jankovic, Chen, & Le, 2005; Saucedo-Cardenas et al., 1998; Smits, Ponnio, Conneely,

Burbach, & Smidt, 2003). An additional key factor involved in the generation and maintenance

of mdDA neurons, *Lmx1a* (Lim- and homeodomain factor 1a) (Andersson et al., 2006; Hong et

al., 2014), was also a key feature gene of clusters 3 and 8 (Supplemental Figure 4A and 4D)

suggesting that POMC and DA neurons share at least part of their genetic differentiation

programs.

227   Similar to cluster 2, reclustering of the differentially expressed genes from cluster 3

revealed two subclusters (Supplemental Figure 4C). Subcluster 3-1 constituted 72% of the main

cluster 3 and was characterized by *Pomc^{low}* neurons with strong *Nr4a2* expression at all

developmental ages. Feature genes for subcluster 3-1 also included *Lmx1a, Slc6a3, Slc18a2,*

Foxa2 and *Th,* all feature genes for developmental subclusters E13.5-10 and E15.5-11 derived

from cluster 3 (Supplemental Figure 7) indicating that a fraction of cluster 3 neurons were

dopaminergic and existed transiently during a three-day developmental window (Supplemental

Tables 2, 5 and 9). In contrast, subcluster 3-2 neurons were *POMC^{high}* together with

characteristic expression of the known *Pomc* transcriptional activator genes and were present

only at ages P5 and P12. The dissociation of high *Pomc* expression with low *Nr4a2* expression

in subcluster 3-2 suggests that the latter is not necessary for hypothalamic *Pomc* expression,

unlike its role in pituitary corticotrophs(Murphy & Conneely, 1997).

Cluster 8. *Pomc^low^/Pitx2* is more closely related to clusters 2. *Pomc^med^/Ebf1* and 3.

Pomc^med^/Nr4a2 than to any of the others (Supplemental Figure 4D). In addition to *Pitx2,* major

feature genes were *Barhl1, Foxa1, Foxp2, Lmx1a/b,* and *Tac1* (Supplemental Table 2).

Intriguingly, virtually all of the top cluster 8 defining genes (Supplemental Table 2) are identical

to those that characterize both the mesencephalic dopamine neurons and glutamatergic, non-

dopaminergic, subthalamic nucleus neurons(Kee et al., 2017). Dual RNAscope in situ

hybridization for *Pitx2* and *Pomc* showed limited overlap at E12.5 (Supplemental Figure 4E).

Cluster 8 is the only cluster with the neuropeptide *Cck* as a feature gene at developmental ages

P5 and P12 (Supplemental Figure S9).

Cluster 4. *Pomc^low^/Otp* cells uniquely expressed *Otp* at all developmental ages (Figure

3A). *Pomc* expression was moderate at E11.5 and then dropped to consistently lower levels

through age P12 despite continued expression of *Isl1, Nkx2-1* and *Tbx3.* Interestingly,

expression of the other transcriptional modulator important for *Pomc* expression, *Prdm12*, was

detected only at E11.5. Other characteristic features of cluster 4 were increasing gradients of

Agrp, Npy and *Calcr* expression over the course of development (Figure 3A). Minimal co-

expression of *Pomc* and *Npy* was confirmed by FISH at age E15.5 (Figure 3C). *Sst* was highly

expressed in this cluster at ages E11.5, E13.5, P5 and P12 with a temporary drop off at E15.5

and E17.5. Altogether, these patterns of gene expression indicate that a subpopulation of *Pomc*

neuronal progenitors differentiates into the canonical AGRP/NPY neurons located in the adult

Arc as suggested previously by studies based on transgenic lineage tracing(Padilla et al., 2010).

Most of the top enriched genes of this cluster are expressed in the middle and end of

pseudotime prediction emphasizing the functions of regulation of inhibitory synapses, GABA-

ergic synapses, ribosomal biogenesis, synaptic vesicle cytoskeletal transport and others (Figure

3B). A map of gene expression in individual cells within the cluster 4 UMAP plot demonstrated

that *Npy* was expressed in virtually all *Agrp* neurons, but few *Sst* neurons. *Sst* and *Agrp*

neurons were essentially separate cell populations (Figure 3D). The GABAergic marker *Slc32a1*

encoding the vesicular inhibitory amino acid transporter was expressed at the second highest

level in cluster 4 relative to the other clusters (Supplemental Table 1), consistent with the

GABAergic phenotype of mature AGRP/NPY neurons (Supplementary Figure 6).

Cluster 5. $Pomc^{low}/Tac2$ represents a distinct neuronal population from cluster 1 in its

patterns of TFs associated with the development of mature POMC neurons (Figure 3E). In

particular, $Isl1$ was highly expressed only at E11.5 and $Prdm12$ at E11.5 and E13.5 in cluster 5.

Starting from E15.5 through P12 there was a gradual onset of expression for a set of genes

including $Tac2, Ar, Esr1, Tacr3, Prlr, Pdyn,$ and $Kiss1/Gm28040$ isoforms (Figure 3E). Most of

the top enriched genes for this cluster are inferred to be expressed at the middle to end along

the pseudotime prediction, and their functions are related to reproductive process, synapse

assembly/organization, pre/post synaptic specialization, and neuron migration (Figure 3F).

Confirmations of co-expression for $Pomc$ with either $Tac2, Tacr3$ and $Sox14$ or TDTOMATO

with ESR1 at several embryonic ages are shown in Figure 3G. Interestingly, $Sox14$ has been

reported to be necessary for expression of $Kiss1$(Huisman et al., 2019). Taken together, this

ensemble of genes is characteristic of adult Arc KNDY neurons that play an essential role in the

control of reproductive physiology. A previous publication based on transgene lineage tracing

also concluded that a significant subpopulation of KNDY neurons is derived from POMC

progenitor cells(Sanz et al., 2015).

A prominent and unique characteristic of **cluster 6.** $Pomc^{low/Dlx5}$ neurons was the

expression of members of the Dlx family of homeobox TFs (Supplementary Figure 5A and 5B).

The six family members are usually expressed in 3 linked pairs of isoforms: 1 and 2, 3 and 4 or

5 and 6. Cluster 6 was enriched in $Dlx1/2,$ and $Dlx5/6$ together with $Dlxos1$. The $Dlx1/2$ pair is

known to activate expression of $Ghrh$ in the mouse Arc(Lee et al., 2018). However, there were

no transcripts for $Ghrh$ expressed in cluster 6 or any of the other $Pomc$ neuronal clusters

suggesting that $Dlx1/2$ are not sufficient to induce $Ghrh$ expression. Moreover, it has been

recently shown that *Dlx1/2* binding to *Otp* regulatory elements inhibits OTP production,

subsequently reducing downstream *Agrp* expression(Lee et al., 2018). Cluster 6 had the highest

expression of the opioid prepronociceptin (*Pnoc*) compared to the other integrated clusters, but

low levels of preprodynorphin (*Pdyn*), similar to cluster 5 KNDY neurons.

Cluster 6 had the highest *Gad1*, *Gad2* and *Slc32a1* (*Vgat,* vesicular inhibitory amino

acid transporter) expression density of all clusters with virtually no *Slc17a6* (*Vglut2,* vesicular

glutamate transporter 2) expression, suggesting that these differentiating neurons were

exclusively GABAergic. A comparison of GABAergic and glutamatergic markers among the

eight neuronal clusters is shown in Supplementary Figure 6. Cluster 4 neurons were also

uniformly GABAergic. Cluster 1 neurons contained a mixture of glutamatergic and GABAergic

markers with the exception of undetectable *Slc32a1.* However, they did express low levels of

Slc18a2 (*Vmat2*, vesicular monoamine transporter 2), which possibly functions as an alternative

vesicular GABA transporter in some dopamine neurons(German, Baladi, McFadden, Hanson, &

Fleckenstein, 2015). This combination of mixed glutamatergic and GABAergic features in cluster

1 is characteristic of postnatal POMC neurons(Jones et al., 2019; Wittmann, Hrabovszky, &

Lechan, 2013) and was also present to a limited extent in cluster 2. The remaining neuronal

clusters 3, 5, 7 and 8 were all uniformly glutamatergic.

Cluster 7. *Pomc^{low}/Prdm13* contained the smallest number of cells among the eight

neuronal clusters, and the cells were primarily limited to developmental ages E13.5 to E17.5

(Figure 1B). Feature genes included *Prdm13, Nr5a1, Adcypap1, Cnr1, Fam19a1, Rbp1* and

Pdyn (Supplementary Figure 5C). *Prdm13* was reported recently(X. Chen et al., 2020) to also

be highly expressed in E15.5 POMC cells.

We further analyzed all of the neurons and non-neuronal cells by reclustering them

based on their individual developmental ages (Supplemental Figure 7) rather than for their gene

expression profiles integrated across all ages. Average gene expression levels for each of these

developmental subclusters are listed in Supplemental Table 8 and the feature genes defining

the developmental subclusters are listed in Supplemental Table 9. Based on these data, it was

then possible to trace the temporal continuity between developmental subclusters from age

E11.5 to P12 relative to the original integrated cell cluster identities (Supplemental Figure 8).

These data are important because they connected the dynamic nature of gene expression

profiles at each embryonic stage leading to the eventual transcriptomes present in the postnatal

time points.

Comparison of transcriptional profiles for neuropeptides, neuropeptide processing

enzymes, neuroendocrine secretory proteins and GPCRs in the eight neuronal POMC

clusters at each developmental age

We identified 19 unique neuropeptide prohormone genes that were clearly differentially

expressed features in at least one of the neuronal clusters at one or more developmental ages

(Supplemental Figure 9, Supplemental Table 5). In addition to the neuropeptide genes already

mentioned as characteristic features of certain clusters, additional neuropeptides were featured

in other clusters primarily at postnatal ages P5 and P12. Notable examples are *Gal, Tac1,*

Adcyap1, Cartpt and *Pthlh.* Unlike the mosaic of neuropeptide gene expression profiles across

clusters and developmental ages, the neuroendocrine secretory proteins, characteristic of

dense core granules,were expressed ubiquitously in all eight neuronal clusters with similar

gradients across all developmental ages (Supplemental Figure S9). These included members of

the chromogranin (*Chga* and *Chgb*) and secretogranin (*Scg2, Scg3* and *Scg5*) families. Among

the prohormone processing enzymes, *Pcsk1* was prominently expressed only at P5 and P12 in

most neuronal clusters while *Pcsk2* was expressed at both embryonic and postnatal ages in the

same clusters as *Pcsk1*. *Pam* was expressed in the same cluster/developmental age pattern as

Pcsk1. Only *Cpe* was expressed in virtually all neurons at all developmental ages.

Split plots of GPCR gene expression revealed a wide range of cluster- and age-specific patterns (Supplemental Figures 10A and 10B). *Cnr1* (Cannabinoid receptor 1) was an important feature primarily of clusters 7 and the two interrelated clusters 3 and 8. The two metabotropic GABA receptors *Gabbr1* and *Gabbr2* were strongly expressed in all eight neuronal clusters but differed in their developmental gradients. *Gabbr1* was expressed at all developmental ages while *Gabbr2* transcripts were largely present only in more mature neurons at postnatal ages P5 and P12. The opioid receptor *Oprl1* that is selectively activated by nociception/orphanin FQ(Toll, Bruchas, Calo, Cox, & Zaveri, 2016) stood out for its consistently strong expression in all eight neuronal clusters, particularly at postnatal days P5 and P12.

The prokineticin receptor 1 (*Prokr1*) was selectively expressed together with the GPCR accessory protein gene *Mrap2* in cluster 1 *Pomc^{high}/Prdm12* neurons. Although MRAP2 was initially identified as an activating modulator of melanocortin receptor signaling, consequently reducing food intake, it was subsequently shown to be promiscuous in its interaction with additional GPCRs(Srisai et al., 2017). Unlike its action on MC4R signaling, MRAP2 inhibits PROKR1 signaling to promote food intake in mice independently of its interaction with MC4R(Chaly, Srisai, Gardner, & Sebag, 2016). Therefore, the co-expression of *Prokr1* and *Mrap2* in anorexigenic POMC neurons suggests an additional mechanism for PROKR1's regulation of energy homeostasis by modulating the release of melanocortins from POMC neurons. Finally, the high expression of *Npy2r* (neuropeptide Y receptor 2) in both clusters 1 and 4 matches well with its expression in adult anorexigenic POMC neurons and orexigenic AGRP/NPY neurons, respectively.

TRAP-seq analysis of Pomc-expressing cells at P12 and P60

To define the important genetic programs that direct the transition from early postnatal to adulthood and to compare the differences of the transcriptional programs that guide embryonic

vs. postnatal / adult development, we performed Translating ribosome affinity purification TRAP-

seq using compound *Pomc-CreERT2; Rosa26-eGFPL10a* transgenic mice where POMC

neurons are labeled with a *eGFPL10a* tag after tamoxifen administration at specific

developmental stages. Colocalization of POMC and GFP immunoreactivity confirms the

expression of *eGFPL10a* in Arc POMC neurons upon tamoxifen injection (Figure 4A). Anti-GFP-

conjugated beads were used to pull-down actively translating RNAs bound to the eGFPL10a

ribosomes from hypothalamic extracts. Both the pull-down mRNA samples and the resultant

supernatant mRNA samples were subjected to RNA sequencing. Principal component analysis

(PCA) shows distinct separation of pull-down and supernatant samples at both ages P12 and

P60 from three independent experiments, suggesting good quality of these samples and the

intrinsic differences between pull-down and supernatant samples (Figure 4B). Both *Pomc* and

GFP were highly expressed in TRAP pull-down samples relative to the supernatants (Figure

4C), further validating the specificity of the TRAP-seq method. Gene enrichment analysis

identified 1143 and 1047 highly enriched genes ($P<0.05$) from *Pomc- eGFPL10a* P12 and P60

TRAP-seq pull-downs, respectively (Figure 4D, Supplemental Tables 10 and 11), from which

653 genes expressed in both datasets (Figure 4E). We hypothesize that the commonly

expressed genes may be associated with the maintenance of POMC functional identity,

whereas the uniquely expressed genes at P12 or P60 may be related to distinct age-dependent

biological phenomena. To test this hypothesis, we performed gene ontology analysis on the 653

co-expressed genes, 394 P60 specifically expressed genes and 490 P12 specifically expressed

genes. The results demonstrate that the commonly expressed genes are associated with the

establishment and maintenance of neuron structure and basic neuronal functions. The top 3

functional annotations include the organization of synapses, regulation of neuron differentiation

and cytoskeleton-dependent intracellular transport. The ontology annotations on the uniquely

expressed genes from P12 and P60 are dramatically different from each other. The functions of

P12 uniquely enriched genes emphasize basic cellular metabolism, cellular modeling and

biochemical changes, whereas, the P60 enriched genes are related to the maturation of POMC

neurons such as neurotransmitter development and POMC specific function establishment

including the responses to nutrient, and the construction of feeding behavioral circuits (Figure

4F). We next compared the TRAP-seq commonly expressed genes to our developmental single

cell RNA-seq data to examine their expression levels in each cluster and each age.

Approximately one third (241) of the genes shown in the heatmaps on the left are abundantly

expressing in the single cell RNA-seq dataset across all the clusters. Notably, compared to the

four embryonic stages, a constantly higher transcript abundance was observed at both P5 and

P12 ages (grey outlined boxes in Figure 4G), indicating great genetic similarities between P5

and P12. Intriguingly, the patterns of expression levels are very similar in both the P12 and P60

TRAP-seq datasets (Figure 4G, right heatmaps). To gain a better visualization of the distribution

of TRAP-seq common genes expression at a single cell level, we grouped the 241 genes as

one module, calculated module scores based on the normalized counts and projected the

scores to a UMAP plot of the eight neuronal clusters (Figure 4H). Most cells from cluster 1

($Pomc^{high}$/$Prdm12$), cluster 5 ($Pomc^{low}$/$Tac2$), cluster 3 ($Pomc^{med}$/$Nr4a2$) and a subcluster of

cluster 2 ($Pomc^{med}$/$Ebf1$) present a higher module score (high gene expression levels).

Assessment together with Figure 5B confirms that most of the high scoring cells are derived

from the two postnatal stages (P5 and P12).

Developmental relationships of gene expression patterns in the eight neuronal POMC

clusters at four embryonic ages

To investigate the origins and trajectories of POMC progenitors, we initially performed RNA

velocity analysis on cells from the 8 neuronal clusters. Since the cells from P5 and P12 are

distantly segregated from the four embryonic stages, which subsequently induces noise for cell

trajectory analysis, we decide to focus on analyzing cells from the four embryonic stages.

Appling cell cluster and age information to RNA velocity analysis reveals multiple progenitor

origins including a major origin from the $Pomc^{high}$/Prdm12 cluster and two minor origins from

distinct portions of the $Pomc^{med}$/Ebf1 cluster (Figure 5A and 5B), which is consistent with the

early appearance of these two clusters at E11.5 (Supplemental Figure 11A). Further

investigation to cells from E13.5, E15.5 and E17.5 suggest dual origins from $Pomc^{high}$/Prdm12

or $Pomc^{med}$/Ebf1 clusters (Supplemental Figure 11B), indicating that a subset of Pomc-

expressing cells may lose Pomc properties during development, which is in agreement with a

previous study (Padilla et al., 2010).

To further validate the hypothesis of multiple origins of POMC progenitors and to better

visualize cell trajectories in three dimensional Euclidian space, we constructed diffusion maps of

all neurons from the four embryonic stages (Figure 5C). The image depicts a triangular beveled

star polygon with rays 1 and 2 derived primarily from distinct groups of neurons at ages E11.5

and E13.5, a center vertex containing individual neurons from the intermediate ages E13.5 and

E15.5 and a third ray with neurons from ages E15.5 and E17.5. The same data are alternatively

color-coded for each neuron's cluster identity (Figure 5D) showing that ray 1 is primarily derived

from cluster 2, ray 2 from cluster 1, the center vertex from a convergence of all clusters 1-8 and

ray 3 from clusters 1, 2, 4, 5 and 6. Compared to Ray 2, cells in Ray 1 are not continuously

ordered, but instead show an abrupt transition at the junction of cells from E11.5 and E13.5. The

diffusion maps analysis appears to confirm the multiple origins of POMC progenitor cells

demonstrated by the RNA velocity study.

We then overlaid the geometric location of each neuron with the expression levels of

selected feature genes from each of the eight clusters (Figure 5E). Consistent with the diffusion

maps analyses, the more granular data for individual genes and neurons illustrates the

essentially identical patterns of high Pomc expression with the two originally discovered cognate

transactivating factors *Isl1* and *Nkx2-1* across embryonic development (Rays 2 and 3).

However, *Isl1* was also expressed in numerous *Pomc^{low}* neurons at E11.5 and E13.5 (Ray 1).

Prdm12 and *Tbx3* exhibited complementary temporal patterns of high expression at early (Ray

2) versus late (Ray 3) embryonic ages, respectively. The combined developmental and

geometric patterns of *Six3, Six6* and *Sox14* expression were all very similar to both *Pomc* and

Nkx2-1, suggesting that all four of these TF genes might be necessary to define the cell fate of a

subpopulation of hypothalamic neuronal progenitors into mature differentiated POMC neurons.

In contrast to those genes with expression patterns that defined the continuum of

neuronal identity from Ray 2 to Ray 3, another set of genes including *Ebf1, Pou4f1* and *Nr4a2*

was similarly highly expressed only at the two earliest embryonic ages in neurons constituting

Ray 1. The expression of *Pitx2,* which characterizes cluster 8 and a portion of cluster 3, was

high in Ray 1 as it merges into the center vertex and was mostly derived from age E13.5.

Finally, the temporal and geometric patterns of high expression levels for *Otp, Prdm13, Tac2*

and *Dlx5* were distinct from each other and the aforementioned genes, consistent with their

ontogeny into the distinct differentiated cell fates of cluster 4. AGRP/NPY/GABA, cluster 5.

KNDY and cluster 6. GABA/PNOC neurons, respectively.

Projection of phenotypically distinct cell clusters identified by developmental

transcriptomic profiles to neuronal clusters in the adult Arc

We next asked how the transcriptomes of developing POMC neuron progenitors were related to

those of neurons in the adult Arc at age P60. The raw single cell sequencing data from

Campbell, et al.(Campbell et al., 2017) was transformed, filtered and then combined with our

data using the standard data sets integration method from the Seurat package as detailed in the

Materials and Methods. Each of the eight *Pomc* neuronal clusters defined from our scRNAseq

data and the 34 neuronal clusters from the adult study of Campbell *et al.* (2017)(Campbell et al.,

2017) were then projected onto combined UMAP plots to predict the cellular relationships

between the two cluster sets (Supplemental Figure 12). A summary of the embryonic to adult

cluster projections is shown in Supplemental Figure 13.

66% of cells in the *Pomc^high^/Prdm12* neuronal cluster projected to cluster n14 *Pomc/Ttr*

from the adult Arc and another 15% projected to n15 *Pomc/Anxa2* neurons. Although, neither of

the adult feature genes *Ttr* or *Anxa2* was expressed significantly in any of the integrated

developmental clusters, other key feature genes such as *Pomc, Nkx2-1, Tbx3, Tox2, Six3, Six6,*

Sox2, and Dlk1 do correlate. All together, these data indicate the continued phenotypic

evolution of POMC neurons between ages P12 to P60.

93% of *Pomc^low^/Tac2* cells projected to the adult cluster n20 *Kiss1/Tac2*, providing

further evidence that at least a subpopulation of embryonic POMC progenitors further

differentiate into mature KNDY neurons. Similarly, there was a 67% correspondence of

Pomc^low^/Otp neurons with cluster n13 *Agrp/GM8773* neurons, while another 27% of the former

projected to n23 *Sst/Unc13c* neurons. These two pairings are consistent with the developmental

subclustering of *Pomc^low^/Otp* neurons into those that co-express *Agrp* and *Npy* and those that

highly express *Sst* but neither of the other two neuropeptides (Figure 3D). 72% of *Pomc^low^/Dlx5*

neurons projected to n22 *Tmem215* neurons. *Tmem215* was identified as a feature gene in

combination with *Dlx5* in subcluster P12-12 from our data set (Supplemental Figure 7 and

Supplemental Table 9).

78% of cluster 7 *Pomc^low^/Prdm13* neurons projected to the adult neuronal cluster n29

Nr5a1/Adcyap1. In the adult hypothalamus, expression of these two genes is characteristic of

the ventromedial hypothalamic nucleus instead of the Arc. Expression of the trio of *Prdm13,*

Nr5a1 and *Adcyap1* genes was a significant defining feature unique to cluster 7

Pomc^low^/Prdm13 neurons from our developmental data set, although *Nr5a1* and *Adcyap1*

individually were also feature genes in additional clusters (Supplemental Table 2). Interestingly,

the other 11% of $Pomc^{low}$/Prdm13 neurons projected to cluster n20 *Kiss/Tac2*. Cluster 5 KNDY

neurons had significant *Tac1* and *Sox14* expression (Figure 3 and Supplemental Table 2),

which may suggest why cluster 7 also projected to the adult n20 *Kiss/Tac2* cluster.

Integrated cluster 2 $Pomc^{med}$/Ebf1 neurons projected primarily to n34 unassigned2 cells,

but 18% projected to n21 *Pomc/Glipr1* neurons and 8% to n14 *Pomc/Ttr* neurons. However,

when cluster 2 was reclustered into two distinct components, subcluster 2-1 clearly projected to

both clusters n14 *Pomc/Ttr* and n15 *Pomc/Anxa2* neurons, exactly the same as integrated

developmental cluster no. 1 $Pomc^{high}$/Prdm12 canonical POMC neurons. Similarly, integrated

cluster 3 $Pomc^{med}$/Nr4a2 neurons projected mostly to n33 unassigned1 cells but 16% of the

neurons projected to n32 *Slc17a6*/Trhr neurons. VGLUT2 encoded by *Slc17a6* was most

prominently expressed in cluster 3 $Pomc^{med}$/Nr4a2 neurons compared to all the other integrated

clusters.

Discussion

In this study, we tracked the genetic programs that give rise to *Pomc*-expressing neurons in the

developing hypothalamus and followed their progression into terminally differentiated POMC

neurons and alternative phenotypic destinies. To this end, we performed a single-cell RNA-seq

study specifically of *Pomc*-expressing cells present in developing mouse hypothalami at four

embryonic time points (E11.5, E13.5, E15.5, E17.5) and two early postnatal days (P5 and P12).

We also performed a complementary TRAP-seq study on affinity purified translating RNAs

derived from hypothalamic POMC cells obtained at ages P12 and P60. These developmental

stages were selected based on the ontogeny of *Pomc* expression in the presumptive

hypothalamus(Japon, Rubinstein, & Low, 1994), peak of neurogenesis of POMC neurons(Nasif

et al., 2015; Orquera et al., 2019), subsequent developmental maturation(Padilla et al., 2010;

Sanz et al., 2015) and terminal differentiation(Quarta et al., 2019).

Our study revealed the existence of a major group of POMC neurons (Cluster 1.

Pomc^high/Prdm12) that showed by far the highest levels of *Pomc* UMIs relative to all other

neuronal clusters at all ages. We believe these developing cells constitute the direct precursors

of canonical POMC neurons in the adult hypothalamus. Cluster 1 was characterized by a unique

combinatorial set of feature TFs that includes *Isl1*, *Nkx2-1* and *Prdm12*. In the last few years,

our laboratories demonstrated that these three TFs are present in neurons that start expressing

Pomc in the developing arcuate nucleus at E10.5, and their expression continues at later

developmental ages and adulthood in hypothalamic POMC neurons. Indeed, we showed that

ISL1, NKX2-1 and PRDM12 are absolutely necessary to specify the identity of POMC

neurons(Hael et al., 2020; Nasif et al., 2015; Orquera et al., 2019). Moreover, we also

demonstrated previously that adult mice lacking any of these three TFs exclusively from POMC

cells display greatly reduced levels of hypothalamic *Pomc* mRNA together with increased

adiposity and body weight. However, our results also indicated that the sole presence of ISL1,

NKX2-1 and PRDM12 at E10.5-E11.5 is insufficient to determine the early identity of

hypothalamic POMC neurons, suggesting that the combinatorial set of TFs necessary for

neuronal-specific *Pomc* expression has additional early components, yet to be discovered. In

this regard, our scRNA-seq study revealed several novel candidates in cluster 1 cells including

Six3, *Six6, Sox1/2/3, Sox14* and other TFs (Figure 2). Future molecular and functional genetic

studies will be needed to assess their potential contribution to Arc *Pomc* expression and the

functioning of POMC neurons.

Six3 and *Six6* are two evolutionarily related TF genes known to play critical roles in the

development of the eyes, forebrain and pituitary(W. Liu & Cvekl, 2017; W. Liu, Lagutin,

Swindell, Jamrich, & Oliver, 2010; Suh et al., 2010). The expression patterns of *Six3* and *Six6*

highly overlap during early embryogenesis but segregate into different territories during late

embryonic development, indicating possible distinct postnatal functions(Geng et al., 2008).

Indeed, *Six3* knockout mice die at birth whereas mice lacking *Six6* exhibit decreased numbers

of GNRH neurons and impaired fertility(Larder, Clark, Miller, & Mellon, 2011). Although *Six3* and

Six6 are abundantly expressed in Arc POMC neurons across all examined developmental

stages (Figure. 2), their participation in the control of *Pomc* expression and the patterning and

function of POMC neurons requires further investigation. Regarding Sox genes, a family of TFs

known to regulate cell fate by transactivating cell-specific genes, *in silico* analysis of conserved

SOX protein binding motifs suggests the potential binding of SOX proteins including SOX1,

SOX2, SOX3, and SOX14 to _Pomc_ neuronal enhancers nPE1 and nPE2(X. Chen et al., 2020).

All four of these *Sox* genes were highly enriched in $Pomc^{high}$/*Prdm12* cluster 1 (Figure 2 and

Supplementary Table 2) as well as in $Pomc^{low}$/*Tac2* cluster 5 and $Pomc^{low}$/*Prdm13* cluster 7.

Furthermore, mice lacking *Sox14* exhibit a great reduction in KNDY neurons(Huisman et al.,

2019).

A particular feature gene present in cluster 1 was *Tbx3*, a T-box TF that has been

recently reported to play a key role in Arc *Pomc* expression(Quarta et al., 2019). Unlike *Isl1*,

Nkx2.1 and *Prdm12*, which are all strictly required at the onset of *Pomc* expression (E10.5),

Tbx3 expression commences several days later (E15.5) in the Arc and, therefore, apparently

does not participate in the early specification of POMC neurons. However, *Tbx3* plays an

important role in the functional maturation of these neurons as a transcriptional booster of

arcuate *Pomc* expression. It was recently demonstrated in mice specifically lacking *Tbx3* from

POMC neurons that *Pomc* expression was diminished leading to hyperphagia and

obesity(Quarta et al., 2019).

The TRAP-seq results add another layer to our understanding of the genetic machinery

that leads to mature functional POMC neurons. Genes that are uniquely enriched at P12 are

associated with the basic cellular biology process and cell metabolism whereas genes

specifically expressed at P60 are involved in the establishment of POMC neuronal functions.

The co-expressed genes from TRAP-seq are highly expressed at P5 and P12 in the single cell

sequencing datasets, suggesting that the genetic programs that dominate embryonic POMC

neuronal development are substantially different from the programs that guide postnatal POMC

neuronal differentiation. Although only one week apart, the cells at P5 share more genetic

similarities to the cells at P12 rather than the cells at E17.5.

An unanticipated finding of our study is that Arc *Pomc*-expressing neurons originate from

multiple distinct genetic pathways. Neuronal precursors in cluster 1 characterized by the

coexpression of feature genes *Isl1*, *Nkx2-1*, *Prdm12*, *Six3*, *Six6*, and the *Sox* genes 1, 2, 3 and

14 (Figure 2) express high *Pomc* levels and give rise to the major group of canonical POMC

neurons. In contrast, cluster 2 characterized by expression of the feature genes *Ebf1* and

Pou4f1, the absence of *Prdm12* and low levels of *Pomc,* differ essentially from cluster 1 POMC

neurons at developmental ages 11.5 through 15.5, which give rise to the alternative POMC

neurons. Further studies are needed to understand the functional significance of this non-

canonical population of *Pomc*-expressing neurons that originate from an alternative route.

The identification of *Pomc^{low}/Tac2* and *Pomc^{low}/Otp* clusters confirmed that some fertility-

regulating KNDY neurons(Sanz et al., 2015) and orexigenic AGRP/NPY neurons(Padilla et al.,

2010) arise from *Pomc* progenitors. Moreover, our data indicate that *Pomc* precursors may also

give rise to four other less characterized neuronal populations noted as *Pomc^{med}/Nr4a2*,

Pomc^{low}/Pitx2, *Pomc^{low}/Dlx5* and *Pomc^{low}/Prdm13*. The *Pomc^{med}/Nr4a2* and *Pomc^{low}/Pitx2* are

related clusters sharing 20% of their top 150 feature genes and both of them projected to the

same two heterogeneous clusters n33 and n34 in adult Arc neurons (Supplemental Figure S13).

Interestingly, many of the top feature genes in cluster 8 *Pomc^{low}/Pitx2*, including *Pitx2, Lmx1a,*

Nr4a2, Foxa1, Foxp2, and *Barhl2* were reported previously as marker genes for the

supramammillary nucleus (SMN) (Kim et al., 2020). This cluster also had the most abundant

Cck transcripts, further indicating that it may represent a group of cells from the SMN

considering the distribution of *Cck*-expressing neurons in the brain(Burgunder & Young, 1989).

Notably, POMC cells are undetectable in the SMN in adult mice, suggesting that early *Pomc*

expression in these cells may disappear during postnatal growth. This holds true when we

compared our datasets to Lam *et. al's* study profiling FACS sorted individual POMC-EGFP cells

from adult hypothalamus(Lam et al., 2017). None of the clusters from their study were

characterized by *Pitx2*, *Lmx1a* or *Foxa1* expression. Similarly, the *Pomc^{low}/Prdm13* cluster failed

to match any clusters from Lam *et. al's* study(Lam et al., 2017), which is not unexpected since

this cluster had the most abundant *Nr5a1* transcripts and corresponded to the n29

Nr5a1/Adcyap1 (Supplemental Figure S13) cluster(Campbell et al., 2017), representing a cell

population in the ventromedial hypothalamus featured by *Nr5a1* (*Sf1*) expression.

Immunohistochemistry showed that NR5A1 is abundantly co-localized in *Pomc*-expressing cells

at early embryonic days (E11.5-15.5) but not later perinatal days (E17.5) (Figure 2C).

Conversely, cluster 2 adult POMC cells (featured with *Fzd5*, *Setd4* and *4732456N10Rik*) from

Lam *et.al's.* study failed to match any of the clusters in our study, indicating that this cluster may

contain cells generated postnatally. The signature genes for our *Pomc^{low}/Dlx5* cluster including

Dlx1, Dlx2, Dlx5, and *Dlx6* were highly enriched and specific for cluster 1 in Lam *et. al's* study,

indicating that the *Pomc^{low}/Dlx5* cluster may become a subset of adult POMC neurons. Other

than the above four less defined clusters, the *Pomc^{high}/Prdm12* cluster is highly similar to cluster

4 in Lam *et. al's* study characterized by *Prdm12, S100a10, Cpne4* and *Calb1* expression,

representing the classic POMC neurons in adult mice.

The *Pomc^{med}/Nr4a2* cluster, particularly subcluster 3-1, is the second cluster enriched

with *Pitx2, Lmx1a,* and *Foxa1* expression and also has the most abundant *Nr4a2*, *Pitx3*, and *Th*

transcripts (Supplemental Table1). The early B cell factors (*Ebf1-3*) were also present in this

subgroup of cells in a temporally restricted expression pattern (Figure 2D). Since the *Ebfs*

control differentiation or migration of MbDA subpopulation in a temporally-dependent

manner(Yin et al., 2009), it is reasonable to infer that $Pomc^{med}/Nr4a2$ subcluster 3-1

(Supplemental Figure S4C) may define a subpopulation of immature MbDA neurons. Indeed,

Nr4a2 is expressed in 18.5% of cells in cluster 4 of adult POMC cells from the Lam *et.al.*

study(Lam et al., 2017).

It is worth mentioning that despite some extent of similarities between embryonic and

adult POMC cells, the majority of cells profiled in our study are distantly related to mature adult

POMC neurons. For example, most adult POMC cells express proprotein convertase PC1/3

(*Pcsk1*) and leptin receptor (*Lepr*), whereas only 22% and 2.7% of cells from our datasets

express the two genes, respectively, indicating dramatic transcriptional signaling changes in

POMC neurons during postnatal maturation after age P12.

Concluding remarks

In summary, this study is the first to comprehensively characterize transcriptomes of Arc

hypothalamic POMC cells during embryonic and early postnatal development. This dataset

extends our understanding of the diversification of early POMC cells and provides a valuable

resource for further elucidating the regulatory mechanism for *Pomc* expression and neuronal

maturation. The identification of new marker genes in each subpopulation from single cell RNA-

seq and uniquely expressed genes at P12 or P60 from the TRAP-seq study will potentially lead

to new directions for future functional studies of POMC neurons.

Acknowledgements

We thank Dr. Graham L. Jones, Dr. Zoe Thompson and Charles Keane for their assistance in

animal dissection, tissue sectioning and data analysis, Dr. Joshua Welch and Tongyu Liu for

advice on data analysis, the Flow Cytometry Core, the Advanced Genomics Core, the

Bioinformatics Core and the Microscopy, Imaging and Cellular Physiology Core (supported by

NIH Grant: P30DK020572) and the University of Michigan Center for Gastrointestinal Research

(supported by NIH grant: P30DK034933) at University of Michigan for data collection.

Author Contributions

640   H.Y., M.R., and M.J.L. jointly conceived the study, designed the experiments, analyzed data and

641   drafted the manuscript. H.Y. conducted the experiments. All of the authors edited and approved

642   the final manuscript.

Funding

645   This study was supported by NIH grant DK068400 (M.J.L. and M.R.)

Competing Interests

648   The authors declare no competing interests.

Methods

Mice

All procedures were performed in accordance with the Institutional Animal Care and Use

Committee (IACUC) at the University of Michigan and followed the Public Health Service

guidelines for the humane care and use of experimental animals. Mice were housed in

ventilated cages under controlled temperature and photoperiod (12-h light/12-h dark cycle, lights

on from 6:00 AM to 6:00 PM), with free access to tap water and laboratory chow (5L0D,

LabDiet). Breeding mice were fed with the breeder chow diet (5008, LabDiet). Transgenic mice

expressing the fluorescent protein Tdimer-Discosoma red (TdDsRed) in POMC neurons were

generated previously[17]. The vast majority of POMC-TdDsRed cells were validated as authentic

POMC neurons based on co-localization of ACTH immunostaining with the POMC-TdDsRed

fluorophore (Hentges, Otero-Corchon, Pennock, King, & Low, 2009) *Pomc-CreERT2* and

Rosa26$^{eGFP-L10a}$ (Stock no. 024750, The Jackson Laboratory) mice were generated as previously

described (J. Liu et al., 2014). To induce POMC neurons specific expression of eGFP-L10a at

P12 or P60, tamoxifen (50 mg/kg) was injected intraperitoneally for five constitutive days from

P6 to P10 or P50 to P54.

Generation of single cell (sc) suspension for scRNAseq

Adult male POMC-TdDsRed/+ mice were bred with female POMC-TdDsRed/+ mice and the day

of copulation plug detection was counted as embryonic day 0.5 (E0.5). Embryonic hypothalami

were dissected at E11.5, E13.5, E15.5, and E17.5. Postnatal hypothalami were isolated from

pups at postnatal day 5 (P5) and P12. In each dissection, hypothalami from at least 6 pups were

pooled together to acquire enough cells for fluorescence-activated cell sorting (FACS). Pooled

tissue samples were digested in papain solution supplemented with DNAse for 20 mins (E11.5

and E13.5), 30 mins (E15.5), 40 mins (E17.5) or 1hr (P5, and P12) with gentle agitation based

on a published protocol[14]. The isolated cells were stained with DAPI as an indicator for cell

viability and sorted by FACS for the transgenic red fluorophore (Supplemental Figure 14). The

collected cells were counted and subjected to scRNA-seq library preparation and sequencing

(10x Genomics).

Single Cell RNA Sequencing Data processing

A total of 37,053 single cells (E11.5: 5,329, E13.5: 7,051, E15.5: 5,148, E17.5: 2,488, P5: 7,551

and P12: 9,486) from six different developmental stages were processed using the 10X

Genomics Chromium system. The libraries were sequenced on Illumina HiSeq 4000 and

NovaSeq platforms. We obtained a total of over 2.6 billion reads with an average of 70,7896

reads per cell. Over 85% of reads mapped confidently to the mouse genome across all six

developmental stages. Raw reads were processed with Cell Ranger (version 2.2 and version

3.0). Seurat package (version 3.1.5)(Butler, Hoffman, Smibert, Papalexi, & Satija, 2018) was

used for downstream analysis. Since the number of genes and UMI counts varied across

developmental stage we applied different criteria at each stage to filter out possible doublets

and low quality cells. Specifically, at E11.5, we removed outlier cells that had UMI counts <

5,000 or >60,000 and gene counts <1,000 or >8,000 (determined by the visualization of UMI

counts and gene count distributions). At E13.5, we removed outlier cells that had UMI counts

<2,500 or >30,000 and gene counts >1,500. At E15.5, we removed outlier cells that had UMI

counts >40,000 and gene counts <1,000 or > 6,000. At E17.5, we removed outlier cells that had

UMI counts >25,000 and gene counts <1.000 or > 5,000. At P5, we removed outlier cells that

had UMI counts <1,500 or >60,000 and gene counts <1,500 or >8,000. At P12, we removed

outlier cells that had UMI counts<2,000 or >60,000and gene counts <1,500 or > 8000.

Moreover, cells with high proportions of mitochondrial genes (>10%) or hemoglobin genes

(>10%) were filtered out. Finally, we removed all the cells without any *Pomc* transcript UMI

counts. A total of 13,953 cells passed the criteria (E11.5: 1,498, E13.5: 1,796, E15.5: 2,078,

E17.5: 1,139, P5: 3,909 and P12: 3,533) for downstream analysis.

The gene expression level for each cell was normalized by the total expression,

multiplied by a scaling factor of 10,000 followed by log-transformation(Butler et al., 2018). The

top 2,000 most variable genes were selected from each age and 60 integration anchors were

used to combine all the datasets together. After scaling and centering the integrated dataset, we

performed principal components (PC) analysis on the data matrix and the top 50 PCs were

selected based on JackStraw and Elbow plots from Seurat Package[19] for data visualization

using Uniform Manifold Approximation and Projection (UMAP) technique. We next constructed a

Shared Nearest Neighbor (SNN) graph by setting an expected number of neighbors to 50. In

order to cluster the cells, a SNN modularity optimization technique within a function

FindClusters was used to group cells together with a resolution parameter 0.1. The identity for

each cluster was assigned based on the level of *Pomc* transcripts and prior knowledge of

marker genes. The similar analysis was applied to define *TdDsRed* cell clusters (Supplementary

Figure 2 and Supplementary Figure 3). In addition to the above criteria applied to remove

unwanted cells, we chose cells with at least one *TdDsRed* transcript UMI count and used the

top 2,000 highly variable genes for principal components analysis. The top 30 PCs were

selected for unsupervised clustering with a resolution of 0.3.

Unsupervised cell clustering from each developmental age (Supplementary Figure 6)

was conducting using the following PC numbers (E11.5: 25; E13.5:24, E15.5: 26, E17.5: 19, P5:

40, P12: 34) with their corresponding resolutions (E11.5: 0.2; E13.5: 0.6, E15.5: 0.5, E17.5: 0.5,

P5: 0.5, P12: 0.5. The UMAP plots, violin plots, feature plots and dot plots were all generated in

the Seurat Package. The average expression levels for each gene within each developmental

subcluster (Supplemental Table 8) were obtained from Seurat and calculated by the average of

[*exp(logNormalized counts) -1*]. The differentially expressed genes in each identified cluster was

identified by the comparison of gene expression levels in a specific cluster to all the other

clusters.

Cluster identity comparisons

We took the following steps to compare cluster identities from developmental POMC cells to 34

adult ARC neuronal clusters(Campbell et al., 2017) (Supplementary Figure 12 and

Supplementary Figure 13). 1) We obtained the adult ARC single cell sequencing data from

(Kiselev, Yiu, & Hemberg, 2018)

and transformed the dataset from SingleCellExperiment format to Seurat object format. 2) We

normalized adult ARC data and acquired the top 2000 most variable gene for further analysis. 3)

The standard datasets integration method from Seurat package was performed with 50

integration anchors and used to combine the two datasets. 4) The integrated data were scaled

and the top 50 PCs were selected for unsupervised clustering implemented in Seurat. 5) We

chose 50 transfer anchors to predict the developmental POMC cells' identities based on adult

ARC cell clusters. A similar approach was applied to compare cluster identities between POMC

cells and DSRED cells (Supplementary Figure 2) with the top 50 anchors for data integration

and 50 anchors to transfer cell identities from POMC cells to DSRED cells. We also applied this

approach to conduct age-pairwise comparisons (Supplementary Figure 8) with default setting

from Seurat V3.1.5 package.

Cell lineage construction

For RNA velocity analysis, spliced and unspliced matrices of reads were summarized using

velocyto (v.0.17.16) with default parameters (La Manno et al., 2018). Low complexity and repeat

regions were downloaded from the UCSC browser (rmsk table from mm10). scVelo was

performed for RNA velocity analysis (v.0.2.2) (Bergen, Lange, Peidli, Wolf, & Theis, 2020). The

cell embedding information was acquired from Seurat analysis. The top 4000 genes, the top 20

principal components and the top 20 neighbors under stochastic mode were used to generate

velocity graph.

Cell lineage was constructed using the Destiny (version 2.14.0) package, which implements

the formulation of diffusion maps(Angerer et al., 2016). Diffusion maps are a spectral method for

non-linear dimension reduction, which is especially suitable for analyzing single-cell gene

expression data from different time-courses. We removed VLMC, glial cells and astrocytes

(cluster 9, 10 and 11 in Figure1) and kept only neurons at embryonic stages (E11.5, E13.5,

E15.5 and E17.5), resulting in a total of 6,005 cells for this analysis. The log-transformed

normalized data from Seurat data slot and the corresponding annotation information were

imported to construct an expression matrix. The diffusion maps were constructed under the

default setting for the Gaussian kernel width sigma (σ) and 300 nearest neighbors. The first two

Diffusion Components (DCs) were used to visualize the results. The 3D plots were produced

using rgl package (version 0.100.50) with the top three DCs.

To perform pseudotime gene expression analysis, cells from cluster 1, cluster 2, cluster 4 and

cluster 5 were extracted from the Seurat object respectively. Raw counts were acquired to

construct Monocle (v.2.4.0)(Qiu, Hill, et al., 2017; Qiu, Mao, et al., 2017; Trapnell et al., 2014)

datasets. Cells were ordered along the pseudotime by setting E11.5 as the root state.

Differentially expressed genes (Log2FoldChange > 0.25 and P<0.05) in each cluster from the

Seurat analysis were plotted along the pseudotime using "plot_pseudotime_heatmap" function.

Gene ontology analysis (Biological Process) were performed using ClusterProfiler (v.3.18.1)

(Yu, Wang, Han, & He, 2012).

TRAP-seq analysis

Mice (*Pomc-CreERT2*; ROSA26[eGFP-L10a]) were euthanized and decapitated. The brain was

removed from the skull and the arcuate nucleus was isolated from 2-mm thick coronal slices

using a brain matrix. Tissues were then homogenized and the subsequent tissue lysate was

subjected to immunopurification steps according to the previous protocol (Heiman, Kulicke,

Fenster, Greengard, & Heintz, 2014). The ribo-depletion kit (RiboGone, Takara, CA) and

SMARTer Stranded total RNA sample prep kit were used to remove excess ribosomal RNA and

synthesize cDNA library. Samples were sequenced on a 50-cycle single end run on a HiSeq

4000 (Illumina) according to manufacturer's protocols.

Raw sequencing data was processed at the University of Michigan Bioinformatics core.

Briefly, the quality of the raw reads data was checked using FastQC (v.0.11.3) and the filtered

reads were aligned to reference genome (UCSC mm10) using TopHat (v.2.0.13) and Bowtie2

(v.2.2.1) with default parameters. The HTSeq/DEseq2 method was used for differential

expression analysis with paired-samples and treatment (pull-down vs. supernatant) as the main

effects.

Tissue collection and Immunofluorescence staining

Depending on developmental stage, embryos at E10.5 – E17.5 were fixed in 4%

paraformaldehyde in phosphate buffer solution (PBS) at 4 °C for a various period of time.

Specifically, embryos at E10.5 to E12.5 were fixed for 1h; heads from embryos at E15.5-E16.6

and E17.5 were fixed for 2 h and 4 h respectively. Tissues were stabilized in 10% sucrose/10%

gelatin in PBS at 37 °C for 30 min prior to embedding in O.C.T. compound as described

previously[9]. Embryonic brains were sectioned sagittally on a cryostat (Leica CM1950) at 20 μm

thickness. For dual immunofluorescence, after washing excess O.C.T. compound with PBS, a

heat-induced antigen retrieval process was performed using citrate buffer (10 mM anhydrous

citric acid and 0.05% Tween-20, pH 6.0) at 80 °C for 30 min followed by two washes in PBS.

Sections were then blocked with 3% normal goat serum and 0.1% Triton X-100 for 1 hour and

incubated with primary antibodies at room temperature overnight. After PBS washes, sections

were incubated with goat secondary antibodies for 2 hours at room temperature. Nuclei were

stained with DAPI (1 mg/l) for 10 min and the slides washed five times with PBS before

mounting with ProLong™ Gold Antifade Mountant (ThermoFisher). Sections were imaged on a

Nikon 90i fluorescence microscope with NIS-Elements software. Information on the sources and

dilution of antibodies are listed in Supplementary Table 12.

***In situ* hybridization**

Fluorescence *in situ* hybridization (FISH) was performed using RNAscope[®] Multiplex

Fluorescent V2 Assay (Advanced Cell Diagnostics) according to the manufacturer's instructions

with slight modifications. Specifically, after washing with PBS, sections were post-fixed with 4%

paraformaldehyde for 20 minutes, and dehydrated in a series of ethanol solutions (50%, 70%

and 100%). After drying, dehydrated sections were incubated in RNAscope[®] Hydrogen Peroxide

solution for 10 mins followed by the protease treatment using RNAscope[®] Protease IV for 20

mins. Sections were then hybridized to target RNA probes for 2 hours in the HybEZ[TM] II

hybridization oven. Hybridized probes were amplified using a cascade of signal amplification

solutions (AMP 1-3) followed by standard signal developing protocol as described in the

manufacturer's brochure. Detailed information on RNA probes and dilutions are listed in

Supplementary Table10. Confocal images were obtained using a Nikon Instruments A1

Confocal Laser Microscope with NIS-Elements software.

 FIGURE LEGENDS

 Figure 1. Single cell sequencing of $Pomc^{DsRed}$ cells from developing mouse hypothalami at

 E11.5, E13.5, E15.5, E17.5, P5 and P12 reveals eight distinct neuronal clusters and three non-

 neuronal clusters. (A) Schematic diagram showing the overall experimental procedure. (B)

 UMAP (Uniform Manifold Approximation and Projection) plot showing the distribution of major

 cell clusters and their corresponding cell% at each developmental stage. (C) Violin plots

 showing the expression of three signature genes and *Pomc* for each cluster. (D) UMAP plots

 showing the enrichment of feature genes in each cluster. Genes representing each cluster were

 colored and highlighted to show the cluster-specific enrichment. The color intensity corresponds

 to the normalized gene expression from Seurat analysis, where gene expression for each cell

 was normalized by the total expression, multiplied by a scale factor of 10,000, and then log-

 transformed. VLMC, vascular and leptomeningeal cell.

Figure 2. Temporal gene expression patterns of the *Pomc^high^/Prdm12* cluster and *Pomc^med^/Ebf1*

clusters. (A) and (D) Violin plots showing the expression of signature genes at each

developmental stage in the *Pomc^high^/Prdm12* cluster and *Pomc^med^/Ebf1* cluster respectively. (B)

and (E) Heatmaps showing the gene ontology analysis based on the top marker genes in the

order of pseudotime in *Pomc^high^/Prdm12* cluster and *Pomc^med^/Ebf1* cluster respectively. (C)

Fluorescence *in situ* hybridization showing the co-localization of *Pomc* (green) and *Six6* (red),

the co-localization of *Pomc* (green) and *Six3* (red) at indicated developmental stages.

Immunofluorescence showing the co-localization TDTOMATO (red) NR5A1 (green) at the

indicated developmental stages. (F) Fluorescence *in situ* hybridization showing the co-

localization of *Pomc* (green) and *Ebf1* (red) and the co-localization of *Pomc* (green) and *Pou4f1*

(red) at indicated developmental stages. (G) Re-clustering cells from the *Pomc^med^/Ebf1* cluster

reveals two sub-clusters denoted as 2-1 *Pomc^high^* and 2-2 *Pomc^low^*. Violin plots showing the

signature gene expression in 2-1 *Pomc^high^* and 2-2 *Pomc^low^* clusters. Insets are magnified views

of the indicated boxes. Arrows indicate co-expressing neurons in the merged panels. Scale bar:

50 μm. Image orientation: left, posterior(P); right, anterior(A). RP: Rathke's pouch; Pit: pituitary

gland. Arc: arcuate nucleus; VMH: ventromedial hypothalamus. Mes: mesenchyme.

Figure 3. Temporal gene expression patterns of the *Pomc^low^/Otp* cluster and *Pomc^low^/Tac2*

cluster. (A) and (E) Violin plots showing the expression of signature genes at each

developmental stage in *Pomc^low^/Otp* cluster and *Pomc^low^/Tac2* cluster respectively. (B) and (F)

Heatmaps showing the gene ontology analysis based on the top marker genes in the order of

pseudotime in *Pomc^low^/Otp* cluster and *Pomc^low^/Tac2* cluster respectively. (C) Fluorescence *in*

situ hybridization showing the co-localization of *Pomc* (green) and *Npy* (red) at E15.5. (D)

UMAP plots showing cells in the *Pomc^low^/Otp* cluster expressing *Sst*, *Agrp*, *Npy*, *Sst*/*Agrp*, *Sst*/

Npy or *Agrp/Npy.* Cells expressing *Sst* and cells expressing both *Agrp*/*Npy* are from two distinct

cell populations. (G) Fluorescence *in situ* hybridization showing the co-localization of *Pomc*

(green) and *Tac2* (red), the co-localization of *Pomc* (green) and *Tacr3* (red), and the co-

localization of *Pomc* (green) and *Sox14* (red) at the indicated developmental stages.

Immunofluorescence showing the co-localization of TDTOMATO (red) and ESR1 (green) at the

indicated developmental stages. Insets are magnified views of the indicated boxes. Arrows

indicate co-expressing neurons in the merged panels. Scale bar: 50 μm. Image orientation: left,

posterior; right, anterior; Pit: pituitary gland, RP: Rathke's pouch.

Figure 4. TRAP-seq showing gene enrichment and gene expression profiles at P12 and P60.

(A) Schematic diagram showing the generation of *Pomc-CreERT2; ROSA26$^{eGFP-L10a}$* mice for

TRAP-seq experiment. Immunofluorescence validated the co-localization of POMC and eGFP.

(B) Principal Component Analysis showing the separation of the RNA sequencing data from

beads pull-down vs. supernatant at P12 (Purple) and P60 (Green) respectively. (C) Heatmap

showing both *Pomc* and eGFP were highly expressed in beads pull-down samples. Rows are

the biological replicates of each sample. Sup: Supernatant samples; PD: Pull-Down samples.

Data were presented as scaled counts per million (CPM). (D) Gene enrichment plots showing

1143 genes and 1047 genes were significantly expressed in beads pull-down samples at P12

and P60 respectively (P<0.05). (E) Venn diagram showing the number of genes highly enriched

in both P12 (Purple) and P60 (Green) beads pull-down samples. (F) Gene ontology analysis

showing the top 15 biological processes that were represented in P12 and P60 co-expressed

genes, P12 uniquely expressed genes and P60 uniquely expressed genes. (G) Expression

profile of the top enriched genes from both P12 and P60 TRAP-seq datasets across 8 neuronal

clusters; grey boxes indicate the higher expression of these genes in the two postnatal stages.

The two heatmaps (right) indicating the top genes expression in P12 and P60 TRAP-seq

datasets. (H) UMAP plot showing the distribution of the top enriched genes from both P12 and

P60 Trap-seq datasets in 8 neuronal cell clusters. CPM: counts per million; org.: organization;

reg.: regulation; pos.:positive; neg.: negative.

Figure 5. RNA velocity and diffusion maps analyses illustrating the developmental trajectories

of hypothalamic neuronal clusters. (A) RNA velocity analysis showing the multiple origins of

embryonic POMC cells in 8 neuronal clusters according to Seurat analysis cell embedding. (B)

RNA velocity analysis showing the multiple origins of POMC cells correspond to the early

embryonic stages. (C-D) Diffusion map showing the cell lineages of POMC neurons during

embryogenesis. Cells are colored on the basis of their developmental stage (C) or based on

their previously defined clusters (D). (E) Diffusion maps showing the expression of selected

genes according to previous characterization of the feature genes representing each cluster.

The color intensity corresponds to the normalized gene expression from Seurat analysis.

 Supplemental Data

Figure S1

Heat map showing the expression of the top 30 marker genes defining each cluster from Figure

1. M: microglia; V: vascular and leptomeningeal cells; A: astrocytes.

Figure S2

Analysis of POMC cells with *DsRed* transcripts reveals 9 distinct cell clusters. (A) UMAP plot

showing the distribution of major cell clusters. (B) Violin plots showing signature gene

expression in each cell type. Three feature genes were chosen to represent each cell type. (C)

UMAP plots showing the enrichment of feature genes in each cluster. Genes representative of

each cluster were colored and highlighted to show the cluster-specific enrichment. The color

intensity corresponds to the normalized gene expression from Seurat analysis. (D) Identification

of similar cell clusters between POMC cells with *DsRed* transcripts and POMC cell clusters.

Red: POMC cell clusters are from Figure 1. Blue: POMC cells with *DsRed* transcripts from

Supplemental Figure 2A. (E) Venn diagram showing the number of cells expressing *Pomc*

transcripts, *DsRed* transcripts or both *Pomc* and *DsRed* transcripts. Pie chart showing the

percentage of POMC cells expressing *DsRed* transcripts at each developmental stage. (F)

Vertical bar-chart showing the percentage of cells with *DsRed* transcripts projected to

corresponding POMC cell clusters. VLMC, vascular and leptomeningeal cell.

Figure S3

Overview of cell clusters from POMC cells with *DsRed* transcripts. (A) UMAP plot showing the

distribution of major cell clusters and their corresponding cell% from each developmental stage.

(B) UMAP plots showing the number of cells at each developmental stage. (C) UMAP plots

showing the splitting of clusters by developmental stages.

Figure S4

Temporal gene expression patterns of the *Pomc^med^/Nr4a2* and *Pomc^low^/Pitx2* clusters. (A) Violin

plots showing the expression of signature genes in the *Pomc^med^/Nr4a2* cluster at each

developmental stage. (B) Fluorescence *in situ* hybridization showing the co-localization of *Pomc*

(green) and *Nr4a2* (red). (C) Re-clustering cells from the *Pomc^med^/Nr4a2* cluster reveals two

sub-clusters denoted as 3-1 *Pomc^low^* and 3-2 *Pomc^high^* clusters. Violin plot showing the signature

gene expression in 3-1 *Pomc^low^* and 3-2 *Pomc^high^* clusters. (D) Violin plots showing the

expression of signature genes at each developmental stage in the *Pomc^low^/Pitx2* cluster. (E)

Fluorescence *in situ* hybridization showing the co-localization of *Pomc* (green) and *Pitx2* (red).

Scale bar: 50 μm. Insets are magnified views of the indicated boxes. Arrows indicate co-

expressing neurons in the merged panels. Scale bar: 50 μm. Image orientation: left, posterior

(P); right, anterior (A). RP: rathke's pouch; Pit: pituitary gland.

Figure S5

Temporal gene expression patterns of the *Pomc^low^/Dlx5* and *Pomc^low^/Prdm13* clusters. (A) Violin

plots showing the expression of signature genes at each developmental stage in the

Pomc^low^/Dlx5 cluster. (B) Fluorescence *in situ* hybridization showing the co-localization of *Pomc*

(green) and *Dlx5* (red). (C) Violin plots showing the expression of signature genes at each

developmental stage in the *Pomc^low^/Prdm13* cluster. (D) Fluorescence *in situ* hybridization

showing the co-localization of *Pomc* (green) and *Prdm13* (red). Insets are magnified views of

the indicated boxes. Arrows indicate co-expressing neurons in the merged panels. Scale bar: 50

μm. Image orientation: left, posterior; right, anterior; Pit: pituitary gland.

Figure S6

Distribution of GABAergic and glutamatergic neurons in POMC^{DsRed} cell clusters (A) UMAP plots

showing the distribution of cells with *Gad1* or *Vglut2* gene expression. (B) Violin plots showing

expression of the GABAergic or glutamatergic marker genes in each neuronal cluster.

Figure S7

UMAP plots showing the distribution of major cell clusters at each developmental stage. (A)

E11.5, (B) E13.5, (C) E15.5, (D) E17.5, (E) P5 and (F) P12. The developmental subclusters

derived from each primary cluster are colored to corresponding cells in the UMAP plots.

Figure S8

POMC^{DsRed} developmental cell subclusters shown in Supplemental Figure 7 are projected to the

original integrated cell clusters from Figure 1. The projections were based on the Seurat

analysis and manual comparisons of feature genes in each subcluster (Supplemental Table 2

and supplemental Table5). The clusters from individual stages leading to the same final

integrated clusters are labeled with the same color. Question marks indicate uncertainties of

individual projections.

Figure S9

Overview of neuropeptide, secretory granule and processing enzyme gene expression patterns

across clusters and developmental ages. The size of dots indicates the percentage of cells

expressing the specific transcripts. The color intensity corresponds to the levels of normalized

gene expression.

Figure S10 A and B

Overview of G-protein coupled receptor (GPCR) gene expression patterns across clusters and

developmental ages. The size of dots indicates the percentage of cells expressing the specific

transcripts. The color intensity corresponds to the levels of normalized gene expression. The list

of genes was acquired from https://www.guidetopharmacology.org/targets.jsp and Gene

Ontology (GO:0005576).

Figure S11

(A) UMAP plots showing the splitting of neuronal clusters by developmental stages. Clusters

were dash-dot circled to show the emerging / evolving progress through development. (B) RNA

velocity analysis showing the multiple origins of POMC cells are mainly from 1.

$Pomc^{high}/Prdm12$ and 2. $Pomc^{med}/Ebf1$ clusters and are mainly from E11.5 and E13.5. (C) RNA

velocity analysis on neurons from three late embryonic stages (E13.5, E15.5, and E17.5)

showing two origins of POMC cells from 1. $Pomc^{high}/Prdm12$ and 2. $Pomc^{med}/Ebf1$ clusters at

E13.5.

Figure S12

Identification of similar clusters between the neuronal POMC^{DsRed} developmental clusters and

the neuronal clusters derived from adult arcuate nucleus(Campbell et al., 2017). (A) UMAP plots

showing the integration of POMC^{DsRed} clusters and adult arcuate nucleus neuronal clusters. (B)

The percentage of POMC^{DsRed} cells projecting to corresponding adult arcuate nucleus neuronal

clusters.

Figure S13

Projection of embryonic POMC^{DsRed} cell subtypes to adult arcuate nucleus neuronal clusters.

Projected cell clusters from the *Campbell et al.* study [14] were labeled with the same color as the

POMC^{DsRed} cells clusters. The adult ARC neuronal clusters highlighted in bold font indicate a

higher percentage of cells projected to this cluster from the corresponding developmental

clusters. The detailed cell embedding and percentage of cells projecting to adult Arc neuronal

clusters are shown in Supplemental Figure S12.

Figure S14

Fluorescence-activated cell sorting (FACS) strategy for collecting the single POMC-TdDsRed

cells used for scRNAseq.

Table S1 (corresponding to Figure 1)

Table S2 (corresponding to Figure 1)

Table S3 (corresponding to Supplemental Figure 2)

Table S4 (corresponding to Supplemental Figure 3)

Table S5 (corresponding to Figures 2 and 3 and Supplementary Figures 4 and 5)

Table S6 (corresponding to Figure 2G, cluster 2 and Supplemental Figure 4C, cluster 3)

Table S7 (corresponding to Figure 2G, cluster 2 and Supplemental Figure 4C, cluster 3)

Table S8 (corresponding to Supplemental Figure 7)

Table S9 (corresponding to Supplemental Figure 7)

Table S10 (corresponding to Figure 4)

Table S11 (corresponding to Figure 4)

Table S12 Table of key resources.

1005

1006

References

Andersson, E., Tryggvason, U., Deng, Q., Friling, S., Alekseenko, Z., Robert, B., . . . Ericson, J. (2006).
Identification of intrinsic determinants of midbrain dopamine neurons. *Cell, 124*(2), 393-405.
doi:10.1016/j.cell.2005.10.037

Angerer, P., Haghverdi, L., Buttner, M., Theis, F. J., Marr, C., & Buettner, F. (2016). destiny: diffusion
maps for large-scale single-cell data in R. *Bioinformatics, 32*(8), 1241-1243.
doi:10.1093/bioinformatics/btv715

Bergen, V., Lange, M., Peidli, S., Wolf, F. A., & Theis, F. J. (2020). Generalizing RNA velocity to transient
cell states through dynamical modeling. *Nat Biotechnol, 38*(12), 1408-1414.
doi:10.1038/s41587-020-0591-3

Bouret, S. G., Bates, S. H., Chen, S., Myers, M. G., Jr., & Simerly, R. B. (2012). Distinct roles for specific
leptin receptor signals in the development of hypothalamic feeding circuits. *J Neurosci, 32*(4),
1244-1252. doi:10.1523/JNEUROSCI.2277-11.2012

Bouret, S. G., Draper, S. J., & Simerly, R. B. (2004a). Formation of projection pathways from the arcuate
nucleus of the hypothalamus to hypothalamic regions implicated in the neural control of feeding
behavior in mice. *J Neurosci, 24*(11), 2797-2805. doi:10.1523/JNEUROSCI.5369-03.2004

Bouret, S. G., Draper, S. J., & Simerly, R. B. (2004b). Trophic action of leptin on hypothalamic neurons
that regulate feeding. *Science, 304*(5667), 108-110. doi:10.1126/science.1095004

Bumaschny, V. F., Yamashita, M., Casas-Cordero, R., Otero-Corchon, V., de Souza, F. S., Rubinstein, M., &
Low, M. J. (2012). Obesity-programmed mice are rescued by early genetic intervention. *J Clin
Invest, 122*(11), 4203-4212. doi:10.1172/JCI62543

Burgunder, J. M., & Young, W. S., 3rd. (1989). Neurons containing cholecystokinin mRNA in the
mammillary region: ontogeny and adult distribution in the rat. *Cell Mol Neurobiol, 9*(2), 281-294.
doi:10.1007/BF00713035

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., & Satija, R. (2018). Integrating single-cell transcriptomic
data across different conditions, technologies, and species. *Nat Biotechnol, 36*(5), 411-420.
doi:10.1038/nbt.4096

Campbell, J. N., Macosko, E. Z., Fenselau, H., Pers, T. H., Lyubetskaya, A., Tenen, D., . . . Tsai, L. T. (2017).
A molecular census of arcuate hypothalamus and median eminence cell types. *Nat Neurosci,
20*(3), 484-496. doi:10.1038/nn.4495

Chaly, A. L., Srisai, D., Gardner, E. E., & Sebag, J. A. (2016). The Melanocortin Receptor Accessory Protein
2 promotes food intake through inhibition of the Prokineticin Receptor-1. *Elife, 5*.
doi:10.7554/eLife.12397

Chen, R., Wu, X., Jiang, L., & Zhang, Y. (2017). Single-Cell RNA-Seq Reveals Hypothalamic Cell Diversity.
Cell Rep, 18(13), 3227-3241. doi:10.1016/j.celrep.2017.03.004

Chen, X., Wyler, S. C., Li, L., Arnold, A. G., Wan, R., Jia, L., . . . Liu, C. (2020). Comparative Transcriptomic
Analyses of Developing Melanocortin Neurons Reveal New Regulators for the Anorexigenic
Neuron Identity. *J Neurosci, 40*(16), 3165-3177. doi:10.1523/JNEUROSCI.0155-20.2020

Geng, X., Speirs, C., Lagutin, O., Inbal, A., Liu, W., Solnica-Krezel, L., . . . Oliver, G. (2008).
Haploinsufficiency of Six3 fails to activate Sonic hedgehog expression in the ventral forebrain
and causes holoprosencephaly. *Dev Cell, 15*(2), 236-247. doi:10.1016/j.devcel.2008.07.003

German, C. L., Baladi, M. G., McFadden, L. M., Hanson, G. R., & Fleckenstein, A. E. (2015). Regulation of
the Dopamine and Vesicular Monoamine Transporters: Pharmacological Targets and
Implications for Disease. *Pharmacol Rev, 67*(4), 1005-1024. doi:10.1124/pr.114.010397

Hael, C. E., Rojo, D., Orquera, D. P., Low, M. J., & Rubinstein, M. (2020). The transcriptional regulator
PRDM12 is critical for Pomc expression in the mouse hypothalamus and controlling food intake,
adiposity, and body weight. *Mol Metab, 34*, 43-53. doi:10.1016/j.molmet.2020.01.007

Heiman, M., Kulicke, R., Fenster, R. J., Greengard, P., & Heintz, N. (2014). Cell type-specific mRNA
purification by translating ribosome affinity purification (TRAP). *Nat Protoc, 9*(6), 1282-1291.
doi:10.1038/nprot.2014.085

Hentges, S. T., Otero-Corchon, V., Pennock, R. L., King, C. M., & Low, M. J. (2009). Proopiomelanocortin
expression in both GABA and glutamate neurons. *J Neurosci, 29*(43), 13684-13690.
doi:10.1523/JNEUROSCI.3770-09.2009

Hong, S., Chung, S., Leung, K., Hwang, I., Moon, J., & Kim, K. S. (2014). Functional roles of Nurr1, Pitx3,
and Lmx1a in neurogenesis and phenotype specification of dopamine neurons during in vitro
differentiation of embryonic stem cells. *Stem Cells Dev, 23*(5), 477-487.
doi:10.1089/scd.2013.0406

Huisman, C., Cho, H., Brock, O., Lim, S. J., Youn, S. M., Park, Y., . . . Lee, J. W. (2019). Single cell
transcriptome analysis of developing arcuate nucleus neurons uncovers their key developmental
regulators. *Nat Commun, 10*(1), 3696. doi:10.1038/s41467-019-11667-y

Huisman, C., Cho, H., Brock, O., Lim, S. J., Youn, S. M., Park, Y., . . . Lee, J. W. (2020). Author Correction:
Single cell transcriptome analysis of developing arcuate nucleus neurons uncovers their key
developmental regulators. *Nat Commun, 11*(1), 1359. doi:10.1038/s41467-020-14964-z

Ikeda, Y., Luo, X., Abbud, R., Nilson, J. H., & Parker, K. L. (1995). The nuclear receptor steroidogenic
factor 1 is essential for the formation of the ventromedial hypothalamic nucleus. *Mol*
Endocrinol, 9(4), 478-486. doi:10.1210/mend.9.4.7659091

Jankovic, J., Chen, S., & Le, W. D. (2005). The role of Nurr1 in the development of dopaminergic neurons
and Parkinson's disease. *Prog Neurobiol, 77*(1-2), 128-138. doi:10.1016/j.pneurobio.2005.09.001

Japon, M. A., Rubinstein, M., & Low, M. J. (1994). In situ hybridization analysis of anterior pituitary
hormone gene expression during fetal mouse development. *J Histochem Cytochem, 42*(8), 1117-
1125. doi:10.1177/42.8.8027530

Jones, G. L., Wittmann, G., Yokosawa, E. B., Yu, H., Mercer, A. J., Lechan, R. M., & Low, M. J. (2019).
Selective Restoration of Pomc Expression in Glutamatergic POMC Neurons: Evidence for a
Dynamic Hypothalamic Neurotransmitter Network. *eNeuro, 6*(2). doi:10.1523/ENEURO.0400-
18.2019

Kee, N., Volakakis, N., Kirkeby, A., Dahl, L., Storvall, H., Nolbrant, S., . . . Perlmann, T. (2017). Single-Cell
Analysis Reveals a Close Relationship between Differentiating Dopamine and Subthalamic
Nucleus Neuronal Lineages. *Cell Stem Cell, 20*(1), 29-40. doi:10.1016/j.stem.2016.10.003

Kim, D. W., Washington, P. W., Wang, Z. Q., Lin, S. H., Sun, C., Ismail, B. T., . . . Blackshaw, S. (2020). The
cellular and molecular landscape of hypothalamic patterning and differentiation from embryonic
to late postnatal development. *Nat Commun, 11*(1), 4360. doi:10.1038/s41467-020-18231-z

Kiselev, V. Y., Yiu, A., & Hemberg, M. (2018). scmap: projection of single-cell RNA-seq data across data
sets. *Nat Methods, 15*(5), 359-362. doi:10.1038/nmeth.4644

Kovalovsky, D., Refojo, D., Liberman, A. C., Hochbaum, D., Pereda, M. P., Coso, O. A., . . . Arzt, E. (2002).
Activation and induction of NUR77/NURR1 in corticotrophs by CRH/cAMP: involvement of
calcium, protein kinase A, and MAPK pathways. *Mol Endocrinol, 16*(7), 1638-1651.
doi:10.1210/mend.16.7.0863

Krude, H., Biebermann, H., Luck, W., Horn, R., Brabant, G., & Gruters, A. (1998). Severe early-onset
obesity, adrenal insufficiency and red hair pigmentation caused by POMC mutations in humans.
Nat Genet, 19(2), 155-157. doi:10.1038/509

La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V., . . . Kharchenko, P. V.
(2018). RNA velocity of single cells. *Nature, 560*(7719), 494-498. doi:10.1038/s41586-018-0414-6

Lam, B. Y. H., Cimino, I., Polex-Wolf, J., Nicole Kohnke, S., Rimmington, D., Iyemere, V., . . . Yeo, G. S. H.
(2017). Heterogeneity of hypothalamic pro-opiomelanocortin-expressing neurons revealed by
single-cell RNA sequencing. *Mol Metab, 6*(5), 383-392. doi:10.1016/j.molmet.2017.02.007

Larder, R., Clark, D. D., Miller, N. L., & Mellon, P. L. (2011). Hypothalamic dysregulation and infertility in mice lacking the homeodomain protein Six6. *J Neurosci, 31*(2), 426-438. doi:10.1523/JNEUROSCI.1688-10.2011

Lee, B., Kim, J., An, T., Kim, S., Patel, E. M., Raber, J., . . . Lee, J. W. (2018). Dlx1/2 and Otp coordinate the production of hypothalamic GHRH- and AgRP-neurons. *Nat Commun, 9*(1), 2026. doi:10.1038/s41467-018-04377-4

Liu, J., Krautzberger, A. M., Sui, S. H., Hofmann, O. M., Chen, Y., Baetscher, M., . . . McMahon, A. P. (2014). Cell-specific translational profiling in acute kidney injury. *J Clin Invest, 124*(3), 1242-1254. doi:10.1172/JCI72126

Liu, W., & Cvekl, A. (2017). Six3 in a small population of progenitors at E8.5 is required for neuroretinal specification via regulating cell signaling and survival in mice. *Dev Biol, 428*(1), 164-175. doi:10.1016/j.ydbio.2017.05.026

Liu, W., Lagutin, O., Swindell, E., Jamrich, M., & Oliver, G. (2010). Neuroretina specification in mouse embryos requires Six3-mediated suppression of Wnt8b in the anterior neural plate. *J Clin Invest, 120*(10), 3568-3577. doi:10.1172/JCI43219

Luo, X., Ikeda, Y., & Parker, K. L. (1994). A cell-specific nuclear receptor is essential for adrenal and gonadal development and sexual differentiation. *Cell, 77*(4), 481-490. doi:10.1016/0092-8674(94)90211-9

Meeks, J. J., Weiss, J., & Jameson, J. L. (2003). Dax1 is required for testis determination. *Nat Genet, 34*(1), 32-33. doi:10.1038/ng1141

Mizusaki, H., Kawabe, K., Mukai, T., Ariyoshi, E., Kasahara, M., Yoshioka, H., . . . Morohashi, K. (2003). Dax-1 (dosage-sensitive sex reversal-adrenal hypoplasia congenita critical region on the X chromosome, gene 1) gene transcription is regulated by wnt4 in the female developing gonad. *Mol Endocrinol, 17*(4), 507-519. doi:10.1210/me.2002-0362

Murphy, E. P., & Conneely, O. M. (1997). Neuroendocrine regulation of the hypothalamic pituitary adrenal axis by the nurr1/nur77 subfamily of nuclear receptors. *Mol Endocrinol, 11*(1), 39-47. doi:10.1210/mend.11.1.9874

Nasif, S., de Souza, F. S., Gonzalez, L. E., Yamashita, M., Orquera, D. P., Low, M. J., & Rubinstein, M. (2015). Islet 1 specifies the identity of hypothalamic melanocortin neurons and is critical for normal food intake and adiposity in adulthood. *Proc Natl Acad Sci U S A, 112*(15), E1861-1870. doi:10.1073/pnas.1500672112

Orquera, D. P., Tavella, M. B., de Souza, F. S. J., Nasif, S., Low, M. J., & Rubinstein, M. (2019). The Homeodomain Transcription Factor NKX2.1 Is Essential for the Early Specification of Melanocortin Neuron Identity and Activates Pomc Expression in the Developing Hypothalamus. *J Neurosci, 39*(21), 4023-4035. doi:10.1523/JNEUROSCI.2924-18.2019

Padilla, S. L., Carmody, J. S., & Zeltser, L. M. (2010). Pomc-expressing progenitors give rise to antagonistic neuronal populations in hypothalamic feeding circuits. *Nat Med, 16*(4), 403-405. doi:10.1038/nm.2126

Parikh, D., Hamid, A., Friedman, T. C., Nguyen, K., Tseng, A., Marquez, P., & Lutfy, K. (2011). Stress-induced analgesia and endogenous opioid peptides: the importance of stress duration. *Eur J Pharmacol, 650*(2-3), 563-567. doi:10.1016/j.ejphar.2010.10.050

Perlmann, T., & Wallen-Mackenzie, A. (2004). Nurr1, an orphan nuclear receptor with essential functions in developing dopamine cells. *Cell Tissue Res, 318*(1), 45-52. doi:10.1007/s00441-004-0974-7

Qiu, X., Hill, A., Packer, J., Lin, D., Ma, Y. A., & Trapnell, C. (2017). Single-cell mRNA quantification and differential analysis with Census. *Nat Methods, 14*(3), 309-315. doi:10.1038/nmeth.4150

Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H. A., & Trapnell, C. (2017). Reversed graph embedding resolves complex single-cell trajectories. *Nat Methods, 14*(10), 979-982. doi:10.1038/nmeth.4402

Quarta, C., Fisette, A., Xu, Y., Collden, G., Legutko, B., Tseng, Y. T., . . . Tschop, M. H. (2019). Functional
identity of hypothalamic melanocortin neurons depends on Tbx3. *Nat Metab, 1*(2), 222-235.
doi:10.1038/s42255-018-0028-1
Sanz, E., Quintana, A., Deem, J. D., Steiner, R. A., Palmiter, R. D., & McKnight, G. S. (2015). Fertility-
regulating Kiss1 neurons arise from hypothalamic POMC-expressing progenitors. *J Neurosci,*
35(14), 5549-5556. doi:10.1523/JNEUROSCI.3614-14.2015
Saucedo-Cardenas, O., Quintana-Hau, J. D., Le, W. D., Smidt, M. P., Cox, J. J., De Mayo, F., . . . Conneely,
O. M. (1998). Nurr1 is essential for the induction of the dopaminergic phenotype and the
survival of ventral mesencephalic late dopaminergic precursor neurons. *Proc Natl Acad Sci U S A,*
95(7), 4013-4018. doi:10.1073/pnas.95.7.4013
Smits, S. M., Ponnio, T., Conneely, O. M., Burbach, J. P., & Smidt, M. P. (2003). Involvement of Nurr1 in
specifying the neurotransmitter identity of ventral midbrain dopaminergic neurons. *Eur J*
Neurosci, 18(7), 1731-1738. doi:10.1046/j.1460-9568.2003.02885.x
Srisai, D., Yin, T. C., Lee, A. A., Rouault, A. A. J., Pearson, N. A., Grobe, J. L., & Sebag, J. A. (2017). MRAP2
regulates ghrelin receptor signaling and hunger sensing. *Nat Commun, 8*(1), 713.
doi:10.1038/s41467-017-00747-6
Suh, C. S., Ellingsen, S., Austbo, L., Zhao, X. F., Seo, H. C., & Fjose, A. (2010). Autoregulatory binding sites
in the zebrafish six3a promoter region define a new recognition sequence for Six3 proteins. *FEBS*
J, 277(7), 1761-1775. doi:10.1111/j.1742-4658.2010.07599.x
Toll, L., Bruchas, M. R., Calo, G., Cox, B. M., & Zaveri, N. T. (2016). Nociceptin/Orphanin FQ Receptor
Structure, Signaling, Ligands, Functions, and Interactions with Opioid Systems. *Pharmacol Rev,*
68(2), 419-457. doi:10.1124/pr.114.009209
Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., . . . Rinn, J. L. (2014). The
dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single
cells. *Nat Biotechnol, 32*(4), 381-386. doi:10.1038/nbt.2859
Wittmann, G., Hrabovszky, E., & Lechan, R. M. (2013). Distinct glutamatergic and GABAergic subsets of
hypothalamic pro-opiomelanocortin neurons revealed by in situ hybridization in male rats and
mice. *J Comp Neurol, 521*(14), 3287-3302. doi:10.1002/cne.23350
Yin, M., Liu, S., Yin, Y., Li, S., Li, Z., Wu, X., . . . Zhou, J. (2009). Ventral mesencephalon-enriched genes
that regulate the development of dopaminergic neurons in vivo. *J Neurosci, 29*(16), 5170-5182.
doi:10.1523/JNEUROSCI.5569-08.2009
Yu, G., Wang, L. G., Han, Y., & He, Q. Y. (2012). clusterProfiler: an R package for comparing biological
themes among gene clusters. *OMICS, 16*(5), 284-287. doi:10.1089/omi.2011.0118

1183

Figure 1A

Timeline of hypothalami collection | **Cell Isolation** | **FACS** | **Library Prep** | **Sequencing** | **Analysis**

E11.5 E13.5 E15.5 E17.5 P5 P12

BD Arial III

10x Genomics

Illumina

B

Microglia

Astrocytes

VLMC

UMAP_2 / UMAP_1

Percentage of Total Neurons

	E11.5	E13.5	E15.5	E17.5	P5	P12
1. *Pomc^high^/Prdm12*	50.8%	40.6%	34.7%	33.3%	44.1%	47.2%
2. *Pomc^med^/Ebf1*	43.4%	31.1%	11.8%	13.7%	13.8%	14.4%
3. *Pomc^med^/Nr4a2*	0.5%	3.4%	5.9%	1.1%	5.6%	9.1%
4. *Pomc^low^/Otp*	2.4%	6.0%	20.7%	20.9%	2.8%	4.2%
5. *Pomc^low^/Tac2*	1.2%	3.3%	10.7%	24.6%	29.1%	16.0%
6. *Pomc^low^/Dlx5*	0.5%	1.0%	6.2%	4.9%	3.9%	5.7%
7. *Pomc^low^/Prdm13*	0.1%	1.4%	3.3%	1.1%	0.2%	0.4%
8. *Pomc^low^/Pitx2*	1.0%	13.2%	6.8%	0.4%	0.6%	2.9%

C

Neuron: *Tubb3*, *Syt1*, *Mapt*

Pomc

Microglia: *Aif1*, *Fcrls*, *P2ry12*

VLMC: *Pdgfra*, *Col3a1*, *Lum*

Astrocytes: *Agt*, *Sox9*, *Aqp4*

D

Pomc *Nr4a2* *Tac2*

Prdm12 *Otp* *Dlx1*

Dlx2 *Dlx5* *Prdm13*

High / Low

Ebf1 *Pitx2* *Lmx1a*

UMAP_2 / UMAP_1