Dual origin and multiple neuropeptidergic trajectories of hypothalamic POMC

progenitors revealed by developmental single-cell transcriptomics

Hui Yu[1*], Marcelo Rubinstein[1,2,3*], Malcolm J Low[1*]

[1]Department of Molecular and Integrative Physiology, University of Michigan Medical School,

Ann Arbor, MI 48109, USA

[2]Instituto de Investigaciones en Ingeniería Genética y Biología Molecular, Consejo Nacional de

Investigaciones Científicas y Técnicas, 1428 Buenos Aires, Argentina

[3]Departamento de Fisiología, Biología Molecular y Celular, Facultad de Ciencias Exactas y

Naturales, Universidad de Buenos Aires, 1428 Buenos Aires, Argentina

* Co-corresponding Authors

Address correspondence to:

Malcolm J Low, MD, PhD, Department of Molecular & Integrative Physiology, University of

Michigan Medical School, North Campus Research Complex, Bldg 25, Rm.25-3680, 2800

Plymouth Rd., Ann Arbor, MI 48109-2800.   Email: mjlow@umich.edu

Marcelo Rubinstein, PhD, Instituto de Investigaciones en Ingeniería Genética y Biología

Molecular, Consejo Nacional de Investigaciones Científicas y Técnicas, 1428 Buenos Aires,

Argentina. Email: mrubins@dna.uba.ar

Hui Yu, PhD, Department of Molecular & Integrative Physiology, University of Michigan Medical

School, North Campus Research Complex, Bldg 25, Rm.25-3634, 2800 Plymouth Rd., Ann

Arbor, MI 48109-2800. Email: huiyuz@umich.edu

Abstract

Proopiomelanocortin (POMC) neurons of the hypothalamic arcuate nucleus are essential to regulate food intake and energy balance. However, the ontogenetic transcriptional programs that specify the identity and functioning of these neurons are poorly understood. Here, we use scRNAseq to define the transcriptomes characterizing POMC progenitors in the developing hypothalamus and their transition into mature neurons. Our data show that *Pomc*-expressing neurons originate from two independent developmental pathways expressing distinct combinations of transcription factors. The predominant cluster, featured by high levels of *Pomc* and *Prdm12* transcripts represents the precursors of canonical arcuate POMC neurons. Additional clusters of progenitors expressing lower levels of *Pomc* mature into different neuronal phenotypes characterized by distinct combinations of transcription factors, neuropeptides, processing enzymes, cell surface and nuclear receptors. We conclude that the genetic programs specifying the identity and differentiation of arcuate POMC neurons are diverse and generate a heterogeneous repertoire of neuronal phenotypes early in development.

Introduction

Proopiomelanocortin (POMC)-expressing neurons located in the arcuate nucleus of the
hypothalamus play an essential role in the regulation of food intake by maintaining a
melanocortin-dependent anorexigenic tone. The critical importance of central melanocortins,
specifically in arcuate neurons, is apparent in mice lacking *Pomc* that develop hyperphagia and
early-onset extreme obesity[1]. This phenotype is closely mirrored in humans carrying biallelic
null-mutations in *POMC*[2]. In addition, arcuate POMC neurons release the *Pomc*-encoded opioid
peptide *β*-endorphin, which is critical in mediating normal food intake and stress-induced
analgesia[3].

Postmitotic POMC precursors acquire their phenotypic identity in the developing
hypothalamus days earlier than any other peptidergic neurons born in the presumptive arcuate
nucleus (eg. NPY/AGRP, GHRH, SST, KISSPEPTIN). *Pomc* mRNA is initially observed in the
tuberal portion of the prospective mouse hypothalamus of E10.5 embryos, following the
concurrent advent of the essential transcription factors (TFs) ISL1[4], NKX2-1[5] and PRDM12[6]. In
fact, the early ablation of either *Isl1*, *Nkx2-1* or *Prdm12* disrupts the onset of hypothalamic *Pomc*
expression in conditional mutant mouse embryos[4-6]. Given that the number of arcuate neurons
coexpressing *Nkx2-1*, *Isl1* and *Prdm12* greatly exceeds that of POMC neurons in this area, it is
expected that additional and still unknown TFs integrate the complete genetic program that
specifies the early identity of POMC neurons.

Other molecular markers of arcuate POMC neurons such as transporters, receptors,
channels and co-transmitters exhibit great variability indicating that a heterogeneous pool of
diverse subpopulations of these neurons are involved in multiple circuits and functions. In fact,
previous studies have suggested that subsets of arcuate *Pomc*-expressing neurons may act as
precursors of terminally differentiated neurons of alternative neuropeptidergic phenotypes such
as NPY/AGRP and KISSPEPTIN/NEUROKININ-B/DYNORPHIN (KNDY) neurons in the adult

hypothalamus[7, 8]. Initial attempts to characterize the transcriptome of adult POMC neurons at the

single-cell level confirmed the heterogeneous nature of arcuate POMC neurons and revealed the

existence of distinct clusters including those co-expressing the aforementioned neuropeptide

genes[9-12]. However, the molecular signatures of the precursors of the various hypothalamic *Pomc*-

expressing lineages and their transition into mature neurons within the postnatal Arc are still

lacking.

Here, we track the origin and maturation of hypothalamic POMC precursors and the

diverse transcriptional programs leading to alternative cellular fates in the developing and early

postnatal hypothalamus by performing single-cell RNA-seq transcriptomics of fluorescently

labeled cells taken from *Pomc-TdDiscomaRed-Sv40PolyA* (*Pomc-TdDsRed*) transgenic mice at

different embryonic (E11.5, E13.5, E15.5 and E17.5) and early postnatal (P5 and P12) ages. Our

results show that POMC precursors arise from two different transcriptional routes. Some of these

precursors further develop into mature POMC neurons while others follow diverse

neuropeptidergic trajectories indicating that a heterogeneous population of *Pomc*-expressing

neurons is established early in the developing arcuate nucleus**.**

84 Results

Cluster analysis of gene expression in Pomc positive hypothalamic cells integrated across

six development ages

We captured the single cell transcriptomes of hypothalamic *Pomc*-expressing cells at four key

embryonic days (E11.5, E13.5, E15.5 and E17.5) and two early postnatal days (P5 and P12)

when the maturing neurons already develop axons and innervate key target nuclei distal to the

Arc[13-15]. Medial basal hypothalami from mice known to express the reporter transgene *Pomc*-

TdDsRed selectively in POMC cells[17,28] were pooled from multiple embryos or postnatal pups at

the indicated ages. Dissociated single cell suspensions were sorted for DsRed fluorescence and

used for library preparation with the 10X genomics platform (Figure 1A). After a rigorous screening

procedure to remove substandard scRNA sequencing data (see Methods), we used a

combination of computational platforms to analyze the transcriptomes of 13,962 cells that had

Pomc counts of ≥ 1 unique molecular identifiers (UMI).

A Seurat analysis of all cells integrated across the six-time points revealed 11 distinct cell

clusters projected onto a UMAP plot (Figure 1B). The same data are represented in a heat map

corresponding to the top 30 expressed genes per cluster (Supplementary Figure 1). Clusters 1-

8 (10,828 cells) were all highly enriched for a panel of three neuronal marker gene transcripts.

The embedded table in Figure 1B lists the percentages of neurons of each cluster at each

developmental age. The other three clusters (3,134 cells) were enriched in transcripts

characteristic of either microglia, vascular and leptomeningeal cells (VLMC) or astrocytes (Figure

1C). GO ontology analyses for biological processes further confirmed the neuronal or non-

neuronal identities of the 11 individual clusters (data not shown).

By definition, all cells in the neuronal and non-neuronal clusters contained *Pomc*

transcripts (UMI ≥ 1), however there were significant differences in the average expression levels

per cluster, ranging from 1.3 to 33.8 [exp(*log Normalized* counts) -1)]. Nomenclature of the 8

neuronal clusters is based on a combination of relatively high (>20), medium or low (<5) *Pomc*

levels and high expression of one differential key feature gene as follows: no. 1, *Pomc^{high}/Prdm12*;

no. 2 *Pomc^{med}/Ebf1*; no. 3, *Pomc^{med}/Nr4a2*; no. 4, *Pomc^{low}/Otp*; no. 5, *Pomc^{low}Tac2*; no. 6,

Pomc^{low}/Dlx5; no. 7, *Pomc^{low}/Prdm13*; and no. 8, *Pomc^{low}/Pitx2*. The cluster distribution and

relative expression of those feature genes are represented by individual UMAP plots (Figure 1D).

Average expression levels of all genes, including *Pomc*, in the 11 clusters and complete lists of

all feature genes that define the individual clusters are presented in Supplemental Tables 1 and

2, respectively.

In order to validate our experimental approach of isolating *Pomc*-expressing cells from the

developing hypothalamus by FACS for the transgenic surrogate marker *Pomc-TdDsRed*, we

performed an independent unsupervised cluster analysis of the transcriptomes from the 5,724

cells that had *DsRed* counts of ≥ 1 UMI (Supplemental Figure 2A-C and Supplemental Table 3).

A projection analysis (Supplemental Figure 2D-F) of the differentially expressed feature genes

defined 9 DsRed clusters (Supplemental Table 4) which closely corresponded to 9 of the 11

clusters based on *Pomc* counts of ≥ 1 UMI (Supplemental Table 2). The two exceptions were the

smallest clusters 7. *Pomc^{low}/Prdm13* and Astrocytes (Supplemental Figure 2D-F). Furthermore,

the percentages of cells at each developmental age from the corresponding neuronal clusters

were remarkably similar in the *Pomc* UMI ≥ 1 data set compared to the *DsRed* UMI ≥ 1 data set

(compare Figure 1B to Supplemental Figure 3A). The different population of DsRed fluorescent

sorted cells showing UMI ≥ 1 for *Pomc* (13,962) or *DsRed* (5,724) may be due to differential half-

lives of the fluorescent protein DSRED used during cell sorting and its transcript used for

barcoding the RNAseq libraries.

Temporal gene expression analysis of the eight neuronal clusters at individual

developmental stages

Next, we analyzed temporal gene expression patterns for each of the 8 neuronal clusters at the

6 distinct developmental ages (Supplemental Table 5). The progressive changes in feature genes

across time are responsible for the observed within cluster heterogeneity on the heat map in

Supplemental Figure 1.

Cluster 1. $Pomc^{high}/Prdm12$ cells were the most abundant among the eight neuronal clusters and consistently had the highest levels of *Pomc* expression at all six developmental ages (Figure 2A). We consider them to be the direct precursors of canonical POMC neurons in the adult hypothalamus (Figure 2). Other feature genes included a set of four TFs known to be critical for the initiation and/or maintenance of Arc *Pomc* expression, *Isl1, Nkx2-1, Prdm12* and $Tbx3^{4-6,}$ [16] (Figure 2B). They were expressed in similar developmental patterns to each other, with the exception of *Tbx3* that was significantly expressed only from day E15.5. Notably, there was a lack of the pituitary-specific *Tbx19* TF, confirming that no pituitary tissue was included in the hypothalamic dissections. Two homologues of the *Drosophila sine oculus* homeobox gene, *Six3* and *Six6,* which have not been described previously in the context of Arc POMC neuron development, were also expressed robustly at all six time points (Figure 2B). Co-expression of their mRNAs with *Pomc* by RNAScope at ages E12.5 and E16.5 confirmed the scRNAseq analysis, although the two TFs were expressed in additional hypothalamic domains and the pituitary gland (Figure 2C, D). *Vim1, Sox1, 2, 3 and 14,* and *Myc*, whose expression is typically associated with neuronal precursors or immature neurons, tended to have higher expression at the earlier embryonic ages (Figure 2B). The nuclear receptor gene *Nr5a1*, also known as steroidogenic factor 1 (*Sf1*), was a key feature gene expressed almost exclusively at ages E11.5 and E13.5 (Figure 2B), but only minimally expressed at later embryonic time points when NR5A1 is critically involved in the development of the hypothalamic ventromedial nucleus (VMN)[17]. This differential spatio-temporal presence of NR5A1 within or outside POMC neurons was confirmed by immunofluorescence on hypothalamic sections taken from E11.5 and E17.5 mouse embryos (Figure 2E). Interestingly, *Nr5a1, Nr0b1* (*Dax1*) and *Shh*, transcripts were also present in Cluster 1 at the two earliest ages, and are factors known to interact with the *Wnt/*β-catenin signaling pathway in early cell fate determination in the adrenal and pituitary glands[18-20].

Cluster 2. $Pomc^{med}/Ebf1$ neurons made up the second highest abundance cluster of POMC neurons, particularly at E11.5 and E13.5. After these time points, the fraction of all neurons

constituting cluster 2 dropped by approximately two thirds from ages E15.5 through P12. In

contrast, the relative percentage of neurons in clusters 3-8 increased over time with peaks at

different embryonic ages (Figure 1B). Cluster 2 neurons, on average, had medium levels of *Pomc*

transcripts that were divided distinctly from their lowest at ages E11.5-E13.5 to highest at later

embryonic and postnatal ages (Figure 3A and B; Supplemental Tables 5). Major cluster 2 feature

markers included members of the non-basic helix-loop-helix early B-cell TF family EBF

(Supplemental Table 2). *Ebf1* is typically co-expressed with either *Ebf2* or *Ebf3* and pairs of these

factors were shared not only in cluster 2, but also in clusters 3 and 8 (Figure 4B and F). Dual

RNAScope *in situ* hybridization showed overlap of *Ebf1* and *Pomc* mRNAs in hypothalamic

neurons at an early embryonic stage (E13.5), whereas at later embryonic stages (E16.5 and

E17.5), POMC neurons had barely detectable *Ebf1* mRNAs (Figure 3C). A similar expression

pattern was found with another cluster 2 featured gene, *Pou4f1 (POU Class 4 Homeobox 1)*

(Figure 3B). Dual RNAScope *in situ* hybridization confirmed the overlap of *Pou4f1* and *Pomc*

mRNAs at E13.5, but not at E17.5 (Figure 3D). These results, together, suggest that *Pomc* mRNA

levels peak only after expression of *Ebf1* and *Pou4f1* decrease. Two other prominent feature

genes in cluster 2 were *Crabp1* and *Crabp2*, which encode cellular retinoic acid binding proteins.

Visual examination of the UMAP plots in Figure 1D suggested that cluster 2 was

composed of two compartments, one with high and one with low *Pomc* abundance. Unsupervised

reclustering of the differentially expressed genes from integrated cluster 2 confirmed the existence

of two subclusters (Figure 3E). Subcluster 2-1 contained two-thirds of the neurons, which were

POMC^{high} and present only from ages E15.5-P12. Subcluster 2-2 contained the other one-third of

cluster 2 neurons, which in contrast were *Pomc*^{low} and detected at ages E11.5 and E13.5.

Consistently, known activating TFs for *Pomc* expression such as *Nkx2-1*, *Isl1* and *Tbx3* were

limited to subcluster 2-1 while the *Ebf* and *Crabp* families were characteristic of subcluster 2-2

(Supplemental Table 6 and Supplemental Table 7). Conspicuously absent from both subclusters

was *Prdm12* mRNA.

Cluster 3. $Pomc^{med}/Nr4a2$ neurons displayed high expression of its major feature gene *Nr4a2* (Nuclear Receptor Subfamily 4 Group A Member 2) (Figure 4A-4C). *Nr4a2* mRNA was also expressed at relatively high levels in clusters 2 and 8. Originally known as *Nurr1*, *Nr4a2* has been implicated in the regulation of *Pomc* expression in the pituitary corticotropic cell line AtT20[21]. Furthermore, *Nr4a2* is a pioneer transcriptional regulator important for the differentiation and maintenance of meso-diencephalic dopaminergic (mdDA) neurons during development[22]. It is crucial for expression of a set of genes including *Slc6a3* (plasma membrane dopamine transporter, DAT), *Slc18a2* (synaptic vesicular monoamine transporter, VMAT), *Th* (Tyrosine hydroxylase), *Foxa1* and *Foxa2* (forkhead family TFs) and *Drd2* (dopamine receptor 2) that together define the dopaminergic neuronal phenotype[23-26]. An additional key factor involved in the generation and maintenance of mdDA neurons, *Lmx1a* (Lim- and homeodomain factor 1a)[23, 27], was also a key feature gene of clusters 3 and 8 (Figure 4B and F) suggesting that POMC and DA neurons share at least part of their genetic differentiation programs.

Similar to cluster 2, reclustering of the differentially expressed genes from cluster 3 revealed two subclusters (Figure 4D). Subcluster 3-1 constituted 72% of the main cluster 3 and was characterized by $Pomc^{low}$ neurons with strong *Nr4a2* expression at all developmental ages. Feature genes for subcluster 3-1 also included *Lmx1a, Slc6a3, Slc18a2*, *Foxa2* and *Th,* all feature genes for developmental subclusters E13.5-10 and E15.5-11 derived from cluster 3 (Supplemental Figure 6) indicating that a fraction of cluster 3 neurons were dopaminergic and existed transiently during a three-day developmental window (Supplemental Tables 2, 5 and 9). In contrast, subcluster 3-2 neurons were $POMC^{high}$ together with characteristic expression of the known *Pomc* transcriptional activator genes and were present only at ages P5 and P12. The dissociation of high *Pomc* expression with low *Nr4a2* expression in subcluster 3-2 suggests that the latter is not necessary for hypothalamic *Pomc* expression, unlike its role in pituitary corticotrophs[28].

Cluster 8. *Pomc^low^/Pitx2* is more closely related to clusters 2 and 3 than to any of the

others (Figure 4E and F). In addition to *Pitx2,* major feature genes were *Barhl1, Foxa1, Foxp2,*

Lmx1a/b, and *Tac1* (Supplemental Table 2). Intriguingly, virtually all of the top cluster 8 defining

genes (Supplemental Table 2) are identical to those that characterize both the mesencephalic

dopamine neurons and glutamatergic, non-dopaminergic, subthalamic nucleus neurons[29]. Dual

RNAscope in situ hybridization for *Pitx2* and *Pomc* showed limited overlap at E12.5 (Figure 4G)*.*

Cluster 8 is the only cluster with the neuropeptide *Cck* as a feature gene at developmental ages

P5 and P12 (Supplemental Figure 8).

Cluster 4. *Pomc^low^/Otp* cells uniquely expressed *Otp* at all developmental ages (Figure 5A

and B). *Pomc* expression was moderate at E11.5 and then dropped to consistently lower levels

through age P12 despite continued expression of *Isl1, Nkx2-1* and *Tbx3.* Interestingly, expression

of the other transcriptional modulator important for *Pomc* expression, *Prdm12*, was detected only

at E11.5. Other characteristic features of Cluster 4 were increasing gradients of *Agrp, Npy* and

Calcr expression over the course of development. Minimal co-expression of *Pomc* and *Npy* was

confirmed by FISH at age E15.5 (Figure 5C). *Sst* was highly expressed in this cluster at ages

E11.5, E13.5, P5 and P12 with a temporary drop off at E15.5 and E17.5. Altogether, these

patterns of gene expression indicate that a subpopulation of *Pomc* neuronal progenitors

differentiates into the canonical AGRP/NPY neurons located in the adult Arc as suggested

previously by studies based on transgenic lineage tracing[7]. A map of gene expression in individual

cells within the Cluster 4 UMAP plot demonstrated that *Npy* was expressed in virtually all *Agrp*

neurons, but few *Sst* neurons. *Sst* and *Agrp* neurons were essentially separate cell populations

(Figure 5D). The GABAergic marker *Slc32a1* encoding the vesicular inhibitory amino acid

transporter was expressed at the second highest level in cluster 4 relative to the other clusters

(Supplemental Table 1), consistent with the GABAergic phenotype of mature AGRP/NPY neurons

(Supplementary Figure5).

Cluster 5. *Pomc*^{low}*/Tac2* represents a distinct neuronal population from Cluster 1 in its

patterns of TFs associated with the development of mature POMC neurons (Figure 6A and B). In

particular, *Isl1* was highly expressed only at E11.5 and *Prdm12* at E11.5 and E13.5 in cluster 5.

Starting from E15.5 through P12 there was a gradual onset of expression for a set of genes

including *Tac2, Ar, Esr1, Tacr3, Prlr, Pdyn,* and *Kiss1/Gm28040* isoforms. Confirmations of co-

expression for *Pomc* with either *Tac2, Tacr3* and *Sox14* or TDTOMATO with ESR1 at several

embryonic ages are shown in Figure 6C, D, E and F, respectively. Interestingly, *Sox14* has been

reported to be necessary for expression of *Kiss1*[30]. Taken together, this ensemble of genes is

characteristic of adult Arc KNDY neurons that play an essential role in the control of reproductive

physiology. A previous publication based on transgene lineage tracing also concluded that a

significant subpopulation of KNDY neurons is derived from POMC progenitor cells[8].

A prominent and unique characteristic of **cluster 6.** *Pomc*$^{low/Dlx5}$ neurons was the

expression of members of the *Dlx* family of homeobox TFs (Supplementary Figure 4A and 4B).

The six family members are usually expressed in 3 linked pairs of isoforms: 1 and 2, 3 and 4 or 5

and 6. Cluster 6 was enriched in *Dlx1/2,* and *Dlx5/6* together with *Dlxos1*. The *Dlx1/2* pair is known

to activate expression of *Ghrh* in the mouse Arc[31]. However, there were no transcripts for *Ghrh*

expressed in cluster 6 or any of the other neuronal clusters suggesting that *Dlx1/2* are not

sufficient to induce *Ghrh* expression. Moreover, it has been recently shown that *Dlx1/2* binding to

Otp regulatory elements inhibits OTP production, subsequently reducing downstream *Agrp*

expression[31]. Cluster 6 had the highest expression of the opioid prepronociceptin (*Pnoc*)

compared to the other integrated clusters, but low levels of preprodynorphin (*Pdyn*), similar to

cluster 5 KNDY neurons.

Cluster 6 had the highest *Gad1*, *Gad2* and *Slc32a1* (*Vgat,* vesicular inhibitory amino acid

transporter) expression density of all clusters with virtually no *Slc17a6* (*Vglut2,* vesicular

glutamate transporter 2) expression, suggesting that these differentiating neurons were

exclusively GABAergic. A comparison of GABAergic and glutamatergic markers among the eight

neuronal clusters is shown in Supplementary Figure 5. Cluster 4 neurons were also uniformly

GABAergic. Cluster 1 neurons contained a mixture of glutamatergic and GABAergic markers with

the exception of undetectable *Slc32a1.* However, they did express low levels of *Slc18a2* (*Vmat2,*

vesicular monoamine transporter 2), which possibly functions as an alternative vesicular GABA

transporter in some dopamine neurons[32]. This combination of mixed glutamatergic and

GABAergic features in cluster 1 is characteristic of postnatal POMC neurons[33, 34] and was also

present in cluster 2. . The remaining neuronal clusters 3, 5, 7 and 8 were all uniformly

glutamatergic.

Cluster 7. *Pomc^low/Prdm13* contained the smallest number of cells among the eight

neuronal clusters, and the cells were primarily limited to developmental ages E13.5 to E17.5

(Figure 1B). Feature genes included *Prdm13, Nr5a1, Adcypap1, Cnr1, Fam19a1, Rbp1* and *Pdyn*

(Supplementary Figure 4D, E and F). *Prdm13* was reported recently[35] to also be highly expressed

in E15.5 POMC cells.

We further analyzed all of the neurons and nonneuronal cells by reclustering them based

on their individual developmental ages (Supplemental Figure 6) rather than for their gene

expression profiles integrated across all ages. Average gene expression levels for each of these

developmental subclusters are listed in Supplemental Table 8 and the feature genes defining the

developmental subclusters are listed in Supplemental Table 9. Based on these data, it was then

possible to trace the temporal continuity between developmental subclusters from age E11.5 to

P12 relative to the original integrated cell cluster identities (Supplemental Figure 7). These data

are important because they connected the dynamic nature of gene expression profiles at each

embryonic stage leading to the eventual transcriptomes present in the postnatal time points.

Comparison of transcriptional profiles for neuropeptides, neuropeptide processing

enzymes, neuroendocrine secretory proteins and GPCRs in the eight neuronal POMC

clusters at each developmental age

We identified 19 unique neuropeptide prohormone genes that were clearly differentially expressed

features in at least one of the neuronal clusters at one or more developmental ages (Supplemental

Figure 8, Supplemental Table 5). In addition to the neuropeptide genes already mentioned as

characteristic features of certain clusters, additional neuropeptides were featured in other clusters

primarily at postnatal ages P5 and P12. Notable examples are *Gal, Tac1, Adcyap1, Cartpt* and

Pthlh. Unlike the mosaic of neuropeptide gene expression profiles across clusters and

developmental ages, the neuroendocrine secretory proteins characteristic of dense core granules

were expressed ubiquitously in all eight neuronal clusters with similar gradients across all

developmental ages (Supplemental Figure 8). These included members of the chromogranin

(*Chga* and *Chgb*) and secretogranin (*Scg2, Scg3* and *Scg5*) families. Among the prohormone

processing enzymes, *Pcsk1* was prominently expressed only at P5 and P12 in most neuronal

clusters while *Pcsk2* was expressed at both embryonic and postnatal ages in the same clusters

as *Pcsk1*. *Pam* was expressed in the same cluster/developmental age pattern as *Pcsk1*. Only

Cpe was expressed in virtually all neurons at all developmental ages.

Split plots of GPCR gene expression revealed a wide range of cluster- and age-specific

patterns (Supplemental Figures 9A and 9B). *Cnr1* (Cannabinoid receptor 1) was an important

feature primarily of clusters 7 and the two interrelated clusters 3 and 8. The two metabotropic

GABA receptors *Gabbr1* and *Gabbr2* were strongly expressed in all eight neuronal clusters but

differed in their developmental gradients. *Gabbr1* was expressed at all developmental ages while

Gabbr2 transcripts were largely present only in more mature neurons at postnatal ages P5 and

P12. The opioid receptor *Oprl1* that is selectively activated by nociception/orphanin FQ[36] stood

out for its consistently strong expression in all eight neuronal clusters, particularly at postnatal

days P5 and P12.

The prokineticin receptor 1 (*Prokr1*) was selectively expressed together with the GPCR

accessory protein gene *Mrap2* in cluster 1 *Pomc^high^/Prdm12* neurons. Although MRAP2 was

initially identified as an activating modulator of melanocortin receptor signaling, consequently

reducing food intake, it was subsequently shown to be promiscuous in its interaction with

additional GPCRs[37]. Unlike its action on MC4R signaling, MRAP2 inhibits PROKR1 signaling to

promote food intake in mice independently of its interaction with MC4R[38]. Therefore, the co-

expression of *Prokr1* and *Mrap2* in anorexigenic POMC neurons suggests an additional

mechanism for PROKR1's regulation of energy homeostasis by modulating the release of

melanocortins from POMC neurons. Finally, the high expression of *Npy2r* (neuropeptide Y

receptor 2) in both clusters 1 and 4 matches well with its expression in adult anorexigenic POMC

neurons and orexigenic AGRP/NPY neurons, respectively.

Developmental relationships of gene expression patterns in the eight neuronal POMC

clusters at four embryonic ages

We altered the color coding of the original UMAP plot of the neuronal clusters shown in Figure 1B

to visualize the developmental age of each neuron superimposed on the integrated cluster

identities (Figure 7A). An alternative depiction of the same data isolated the individual

developmental ages into separate UMAP plots with color coding based on the cluster identities

(Figure 7B). Although it is possible to infer the developmental relationships among the

transcriptional profiles of each neuronal cluster at selected embryonic and postnatal ages as

depicted in Figure 7A and B, dimensionality reduction using a diffusion maps algorithm is an

alternative method to visualize the developmental relationships in three-dimensional Euclidean

space. This analysis for all neurons with *Pomc* UMIs ≥1 color-coded for the four embryonic ages

is shown in Figure 7C. The image depicts a triangular beveled star polygon with rays 1 and 2

derived primarily from distinct groups of neurons at ages E11.5 and E13.5, a center vertex

containing individual neurons from the intermediate ages E13.5 and E15.5 and a third ray with

neurons from ages E15.5 and E17.5. The same data are alternatively color-coded for each

neuron's cluster identity (Figure 7D) showing that ray 1 is primarily derived from cluster 2, ray 2

from cluster 1, the center vertex from a convergence of all clusters 1-8 and ray 3 from clusters 1,

2, 4, 5 and 6.

We then overlaid the geometric location of each neuron with the expression levels of

selected feature genes from each of the eight clusters (Figure 7E). Consistent with the diffusion

maps analyses, the more granular data for individual genes and neurons illustrates the essentially

identical patterns of high *Pomc* expression with the two originally discovered cognate

transactivating factors *Isl1* and *Nkx2-1* across embryonic development (Rays 2 and 3). However,

Isl1 was also expressed in numerous *Pomc^{low}* neurons at E11.5 and E13.5 (Ray 1). *Prdm12* and

Tbx3 exhibited complementary temporal patterns of high expression at early (Ray 2) versus late

(Ray 3) embryonic ages, respectively. The combined developmental and geometric patterns of

Six3, Six6 and *Sox14* expression were all very similar to both *Pomc* and *Nkx2-1*, suggesting that

all four of these TF genes might be necessary to define the cell fate of a subpopulation of

hypothalamic neuronal progenitors into mature differentiated POMC neurons.

In contrast to those genes with expression patterns that defined the continuum of neuronal

identity from Ray 2 to Ray 3, another set of genes including *Ebf1, Pou4f1* and *Nr4a2* was similarly

highly expressed only at the two earliest embryonic ages in neurons constituting Ray 1. The

expression of *Pitx2,* which characterizes cluster 8 and a portion of cluster 3, was high in Ray 1 as

it merges into the center vertex and was mostly derived from age E13.5. Finally, the temporal and

geometric patterns of high expression levels for *Otp, Prdm13, Tac2* and *Dlx5* were distinct from

each other and the aforementioned genes, consistent with their ontogeny into the distinct

differentiated cell fates of cluster 4 AGRP/NPY/GABA, cluster 5 KNDY and cluster 6 GABA/PNOC

neurons, respectively.

Projection of phenotypically distinct cell clusters identified by developmental

transcriptomic profiles to clusters in the adult Arc

We next asked how the transcriptomes of developing POMC neuron progenitors were related to

those of neurons in the adult Arc at age P60. The raw single cell sequencing data from Campbell,

et al.[9] was transformed, filtered and then combined with our data using the standard data sets

integration method from the Seurat package as detailed in the section Materials and Methods.

The combined data were scaled and the top 50 principal components were selected for

unsupervised clustering. Each of the eight *Pomc* neuronal clusters defined from our scRNAseq

data and the 34 neuronal clusters from the adult study of Campbell *et al.* (2017)[9] were then

mapped onto the combined UMAP plots to predict the cell projections between the two cluster

sets (Supplemental Figure 10). A summary of the cluster projections is shown in Figure 8.

66% of cells in the *Pomc^{high}/Prdm12* neuronal cluster projected to cluster n14 *Pomc/Ttr*

from the adult Arc and another 15% projected to n15 *Pomc/Anxa2* neurons. Although, neither of

the adult feature genes *Ttr* or *Anxa2* was expressed significantly in any of the integrated

developmental clusters, other key feature genes such as *Pomc, Nkx2-1, Tbx3, Tox2, Six3, Six6,*

Sox2, and Dlk1 do correlate. All together, these data indicate the continued phenotypic evolution

of POMC neurons between ages P12 to P60.

93% of *Pomc^{low}/Tac2* cells projected to the adult cluster n20 *Kiss1/Tac2*, providing further

evidence that at least a subpopulation of embryonic POMC progenitors are further differentiated

into mature KNDY neurons. Similarly, there was a 67% correspondence of *Pomc^{low}/Otp* neurons

with cluster n13 *Agrp/GM8773* neurons, while another 27% of the former projected to n23

Sst/Unc13c neurons. These two pairings are consistent with the developmental subclustering of

Pomc^low^/Otp neurons into those that co-express *Agrp* and *Npy* and those that highly express *Sst*

but neither of the other two neuropeptides (Figure 5D). 72% of *Pomc^low^/Dlx5* neurons projected

to n22 *Tmem215* neurons. *Tmem215* was identified as a feature gene in combination with *Dlx5*

in subcluster P12-12 from our data set (Supplemental Figure 6 and Supplemental Table 9).

78% of cluster 7 *Pomc^low^/Prdm13* neurons projected to the adult neuronal cluster n29

Nr5a1/Adcyap1. In the adult hypothalamus, expression of these two genes is characteristic of the

ventromedial hypothalamic nucleus instead of the Arc. Expression of the trio of *Prdm13*, *Nr5a1*

and *Adcyap1* genes was a significant defining feature unique to cluster 7 *Pomc^low^/Prdm13*

neurons from our developmental data set, although *Nr5a1* and *Adcyap1* individually were also

feature genes in additional clusters (Supplemental Table 2). Interestingly, the other 11% of

Pomc^low^/Prdm13 neurons projected to cluster n20 *Kiss/Tac2*. Cluster 5 KNDY neurons had

significant *Tac1* and *Sox14* expression (Figure 6 and Supplemental Table 2), which may suggest

why cluster 7 also projected to the adult n20 *Kiss/Tac2* cluster.

Integrated cluster 2 *Pomc^med^/Ebf1* neurons projected primarily to n34 unassigned2 cells,

but 18% projected to n21 *Pomc/Glipr1* neurons and 8% to n14 *Pomc/Ttr* neurons. However, when

cluster 2 was reclustered into two distinct components, subcluster 2-1 clearly projected to both

clusters n14 *Pomc/Ttr* and n15 *Pomc/Anxa2* neurons, exactly the same as integrated

developmental cluster no. 1 *Pomc^high^/Prdm12* canonical POMC neurons. Similarly, integrated

cluster 3 *Pomc^med^/Nr4a2* neurons projected mostly to n33 unassigned1 cells but 16% of the

neurons projected to n32 *Slc17a6/Trhr* neurons. VGLUT2 encoded by *Slc17a6* was most

prominently expressed in cluster 3 *Pomc^med^/Nr4a2* neurons compared to all the other integrated

clusters.

Discussion

In this study, we sought to track the genetic programs that give rise to *Pomc*-expressing neurons in the developing hypothalamus and follow their progression into terminally differentiated POMC neurons and alternative phenotypic destinies. To this end, we performed a single-cell RNA-seq study specifically of *Pomc*-expressing cells present in developing mouse hypothalami at four embryonic time points (E11.5, E13.5, E15.5, E17.5) and two early postnatal days (P5 and P12). These developmental stages were selected based on the ontogeny of *Pomc* expression in the presumptive hypothalamus[39], peak of neurogenesis of POMC neurons[4, 5], subsequent developmental maturation[7, 8] and terminal differentiation[16]. Unlike previous studies using fluorescently-labeled POMC neurons, we validated our cluster profile of sorted cells by an independent unsupervised transcriptome cluster analysis of hypothalamic cells expressing *Pomc*-driven *TdDsRed* (UMIs ≥ 1). This analysis yielded 9 clusters that closely corresponded to 9 of the 11 clusters based on *Pomc* UMIs.

Our study revealed the existence of a major group of POMC neurons (Cluster 1. *Pomc*$^{high/}$*Prdm12*) that showed by far the highest levels of *Pomc* UMIs relative to all other neuronal clusters at all ages. We believe these developing cells constitute the direct precursors of canonical POMC neurons in the adult hypothalamus. In addition, cluster 1 neurons were the most abundant compared to the other clusters at all six developmental ages (Fig. 1B). Cluster 1 was characterized by a unique combinatorial set of feature TFs that includes *Isl1*, *Nkx2-1* and *Prdm12*. In the last few years, our laboratories demonstrated that these three TFs are present in neurons that start expressing *Pomc* in the developing arcuate nucleus at E10.5, and their expression continues at later developmental ages and adulthood in hypothalamic POMC neurons. Indeed, we showed that ISL1, NKX2-1 and PRDM12 are absolutely necessary to specify the identity of POMC neurons[4-6]. Moreover, we also demonstrated that adult mice lacking any of these three TFs exclusively from POMC cells display greatly reduced levels of hypothalamic *Pomc* mRNA together with increased adiposity and body weight. However, our results also indicated that the

sole presence of ISL1, NKX2-1 and PRDM12 at E10.5-E11.5 is insufficient to determine the early

identity of hypothalamic POMC neurons, suggesting that the combinatorial set of TFs necessary

for neuronal-specific *Pomc* expression has additional early components, yet to be discovered. In

this regard, our scRNA-seq study revealed several novel candidates in cluster 1 cells including

Six3, *Six6, Sox1/2/3, Sox14* and other TFs (Figure 2). Future molecular and functional genetic

studies will be needed to assess their potential contribution to Arc *Pomc* expression and the

functioning of POMC neurons.

Six3 and *Six6* are two evolutionarily related TF genes known to play critical roles in the

development of the eyes, forebrain and pituitary[40-42]. The expression patterns of *Six3* and *Six6*

highly overlap during early embryogenesis but segregate into different territories during late

embryonic development, indicating possible distinct postnatal functions[43]. Indeed, *Six3* knockout

mice die at birth whereas mice lacking *Six6* exhibit decreased numbers of GNRH neurons and

impaired fertility[44]. Although *Six3* and *Six6* are abundantly expressed in Arc POMC neurons

across all examined developmental stages (Fig. 2), their participation in the control of *Pomc*

expression and the patterning and function of POMC neurons requires further investigation.

Regarding Sox genes, a family of TFs known to regulate cell fate by transactivating cell-specific

genes, *in silico* analysis of conserved SOX protein binding motifs suggests the potential binding

of SOX proteins including SOX1, SOX2, SOX3, and SOX14 to <u>*Pomc*</u> neuronal enhancers nPE1

and nPE2[35]. All four of these *Sox* genes were highly enriched in *Pomc*^{high}*/Prdm12* cluster 1 (Figure

2 and Supplementary Table 2) as well as in *Pomc*^{low}*/Tac2* cluster 5 and *Pomc*^{low}*/Prdm13* cluster

7. Furthermore, mice lacking *Sox14* exhibit a great reduction in KNDY neurons[30].

A particular feature gene present in cluster 1 was *Tbx3*, a T-box TF that has been recently

reported to play a key role in Arc *Pomc* expression[16]. Unlike *Isl1*, *Nkx2.1* and *Prdm12*, which are

all strictly required at the onset of *Pomc* expression (E10.5), *Tbx3* expression commences several

days later (E15.5) in the Arc and, therefore, apparently does not participate in the early

specification of POMC neurons. However, *Tbx3* plays an important role in the functional

maturation of these neurons as a transcriptional booster of arcuate *Pomc* expression. It was

recently demonstrated in mice specifically lacking *Tbx3* from POMC neurons that *Pomc*

expression was diminished leading to hyperphagia and obesity[16].

An unanticipated finding of our study is that Arc *Pomc*-expressing neurons originate from

two distinct genetic pathways. Neuronal precursors in cluster 1 characterized by the coexpression

of feature genes *Isl1*, *Nkx2-1*, *Prdm12*, *Six3*, *Six6*, and the *Sox* genes 1, 2, 3 and 14 (Figure 2)

express high *Pomc* levels and give rise to the major group of canonical POMC neurons. In

contrast, cluster 2 characterized by expression of the feature genes *Ebf1* and *Pou4f1*, the

absence of *Prdm12* and low levels of *Pomc,* differ essentially from cluster 1 POMC neurons at

developmental ages 11.5 through 15.5. Subsequently, about two thirds of cluster 2 precursors

mature into identifiable POMC neurons expressing more moderate expression levels of *Pomc*

and TFs such as *Isl1*, *Nkx2-1* and *Tbx3* compared to cluster 1 (Figure 3, subcluster 2-1). The

other descendants of Cluster 2 retain the expression of *Ebf* factors, *Crabp1*, *Crabp2* and *Nr4a2*

and express only very low *Pomc* levels (Figure 3, subcluster 2-2). A previous study identified a

novel neuronal subtype in the htypothalamus featured by *Crabp1* [10]. Both ISH and

immunostaining showed the restricted expression of *Crabp1* in the Arc. Analyzing the raw data

from that study, we found 11% of neurons co-expressed *Crabp1* and low-levels of *Pomc*.

Collectively, these data indicate that the $Pomc^{med}$/*Ebf1* cluster contains the early progenitor cells

that become adult *Crabp1* clusters in the Arc. Further studies are needed to understand the

functional significance of this non-canonical population of *Pomc*-expressing neurons that originate

from an alternative route.

The identification of $Pomc^{low}$/*Tac2* and $Pomc^{low}$/*Otp* clusters confirmed that some fertility-

regulating KNDY neurons[8] and orexigenic AGRP/NPY neurons[7] arise from *Pomc* progenitors.

Moreover, our data indicate that *Pomc* precursors may also give rise to four other less

characterized neuronal populations noted as *Pomc^med^/Nr4a2*, *Pomc^low^/Pitx2*, *Pomc^low^/Dlx5* and

Pomc^low^/Prdm13. The *Pomc^med^/Nr4a2* and *Pomc^low^/Pitx2* are related clusters sharing 20% of their

top 150 feature genes and both of them projected to the same two heterogeneous clusters n33

and n34 in adult Arc neurons (Figure 8). Interestingly, many of the top feature genes in cluster 8

Pomc^low^/Pitx2, including *Pitx2, Lmx1a, Nr4a2, Foxa1, Foxp2*, and *Barhl2* were reported previously

as marker genes for the supramammillary nucleus (SMN) [45]. This cluster also had the most

abundant *Cck* transcripts, further indicating that it may represent a group of cells from the SMN

considering the distribution of *Cck*-expressing neurons in the brain[46]. Notably, POMC cells are

undetectable in the SMN in adult mice, suggesting that early *Pomc* expression in these cells may

disappear during postnatal growth. This holds true when we compared our datasets to Lam *et.*

al's study profiling FACS sorted individual POMC-EGFP cells from adult hypothalamus[11]. None

of the clusters from their study were characterized by *Pitx2*, *Lmx1a* or *Foxa1* expression. Similarly,

the *Pomc^low^/Prdm13* cluster failed to match any clusters from Lam *et. al's* study[11], which is not

unexpected since this cluster had the most abundant *Nr5a1* transcripts and corresponded to the

n29 *Nr5a1/Adcyap1* (Figure 8) cluster[9], representing a cell population in the ventromedial

hypothalamus featured by *Nr5a1* (*Sf1*) expression. Immunohistochemistry showed that NR5A1 is

abundantly co-localized in *Pomc*-expressing cells at early embryonic days (E11.5-15.5) but not

later perinatal days (E17.5) (Figure 2E). Conversely, cluster 2 adult POMC cells (featured with

Fzd5, *Setd4* and *4732456N10Rik*) from Lam *et.al's* . study failed to match any of the clusters in

our study, indicating that this cluster may contain cells generated postnatally. The signature genes

for our *Pomc^low^/Dlx5* cluster including *Dlx1, Dlx2, Dlx5*, and *Dlx6* were highly enriched and specific

for cluster 1 in Lam *et. al's* study, indicating that the *Pomc^low^/Dlx5* cluster may become a subset

of adult POMC neurons. Other than the above four less defined clusters, the *Pomc^high^/Prdm12*

cluster is highly similar to cluster 4 in Lam *et. al's* study characterized by *Prdm12, S100a10,*

Cpne4 and *Calb1* expression, representing the classic POMC neurons in adult mice.

The *Pomc^med^/Nr4a2* cluster, particularly subcluster 3-1, is the second cluster enriched with

Pitx2, Lmx1a, and *Foxa1* expression and also has the most abundant *Nr4a2*, *Pitx3*, and *Th*

transcripts (Supplemental Table1). The early B cell factors (*Ebf1-3*) were also present in this

subgroup of cells in a temporally restricted expression pattern (Figure 4B). Since the *Ebfs* control

differentiation or migration of MbDA subpopulation in a temporal-dependent manner[47], it is

reasonable to infer that *Pomc^med^/Nr4a2* subcluster 3-1 (Figure 4) may define a subpopulation of

immature MbDA. Indeed, *Nr4a2* is expressed in 18.5% of cells in cluster 4 of adult POMC cells

from the Lam *et.al.* study[11].

It is worth mentioning that despite some extent of similarities between embryonic and adult

POMC cells, the majority of cells profiled in our study are distantly related to mature adult POMC

neurons. For example, most adult POMC cells express proprotein convertase PC1/3 (*Pcsk1*) and

leptin receptor (*Lepr*), whereas only 22% and 2.7% of cells from our datasets express the two

genes, respectively, indicating dramatic transcriptional signaling changes in POMC neurons

during postnatal maturation after age P12.

Concluding remarks

In summary, this study is the first to comprehensively characterize transcriptomes of POMC cells

during embryonic and early postnatal development. This dataset extends our understanding of

the diversification of early POMC cells and provides a valuable resource for further elucidating

the regulatory mechanism for *Pomc* expression and neuronal maturation. The identification of

new marker genes in each subpopulation as mentioned above and listed in Supplementary Table

2 will potentially lead to new directions for future functional studies.

Acknowledgements

We thank Dr. Graham L. Jones, Dr. Zoe Thompson and Charles Keane for their assistance in animal dissection, tissue sectioning and data analysis, Dr. Joshua Welch and Tongyu Liu for advice on data analysis, the Flow Cytometry Core, the Advanced Genomics Core, and the Microscopy, Imaging and Cellular Physiology Core (supported by NIH Grant: P30DK020572) and the University of Michigan Center for Gastrointestinal Research (supported by NIH grant: P30DK034933) at University of Michigan for data collection.

Author Contributions

H.Y., M.R., and M.J.L. jointly conceived the study, designed the experiments, analyzed data and drafted the manuscript. H.Y. conducted the experiments. All of the authors edited and approved the final manuscript.

Funding

This study was supported by NIH grant DK068400 (M.J.L. and M.R.)

Competing Interests

The authors declare no competing interests.

Accession Codes

554 All raw data have been deposited in the Gene Expression Omnibus under accession number
555 GSExxxx. No custom code was used in this study. The detailed analytical methods were
556 described in the methods.

Methods

Mice

All procedures were performed in accordance with the Institutional Animal Care and Use
Committee (IACUC) at the University of Michigan and followed the Public Health Service
guidelines for the humane care and use of experimental animals. Mice were housed in ventilated
cages under controlled temperature and photoperiod (12-h light/12-h dark cycle, lights on from
6:00 AM to 6:00 PM), with free access to tap water and laboratory chow (5L0D, LabDiet). Breeding
mice were fed with the breeder chow diet (5008, LabDiet). Transgenic mice expressing the
fluorescent protein Tdimer-Discosoma red (TdDsRed) in POMC neurons were generated
previously[17]. The vast majority of POMC-TdDsRed cells were validated as authentic POMC
neurons based on co-localization of ACTH immunostaining with the POMC-TdDsRed fluorophore
[17].

Generation of single cell (sc) suspension for scRNAseq

Adult male POMC-TdDsRed/+ mice were bred with female POMC-TdDsRed/+ mice and the day
of copulation plug detection was counted as embryonic day 0.5 (E0.5). Embryonic hypothalami
were dissected at E11.5, E13.5, E15.5, and E17.5. Postnatal hypothalami were isolated from
pups at postnatal day 5 (P5) and P12. In each dissection, hypothalami from at least 6 pups were
pooled together to acquire enough cells for fluorescence-activated cell sorting (FACS). Pooled
tissue samples were digested in papain solution supplemented with DNAse for 20 mins (E11.5
and E13.5), 30 mins (E15.5), 40 mins (E17.5) or 1hr (P5, and P12) with gentle agitation based on

a published protocol[14]. The isolated cells were stained with DAPI as an indicator for cell viability

and sorted by FACS for the transgenic red fluorophore (Supplemental Figure 11). The collected

cells were counted and subjected to scRNA-seq library preparation and sequencing (10x

Genomics).

Data processing

A total of 37,053 single cells (E11.5: 5,329, E13.5: 7,051, E15.5: 5,148, E17.5: 2,488, P5: 7,551

and P12: 9,486) from six different developmental stages were processed using the 10X Genomics

Chromium system. The libraries were sequenced on Illumina HiSeq 4000 and NovaSeq platforms.

We obtained a total of over 2.6 billion reads with an average of 70,7896 reads per cell. Over 85%

of reads mapped confidently to the mouse genome across all six developmental stages. Raw

reads were processed with Cell Ranger (version 2.2 and version 3.0). Seurat package (version

3.1.5)[48] was used for downstream analysis. Since the number of genes and UMI counts varied

across developmental stagewe applied different criteria at each stage to filter out possible

doublets and low quality cells. Specifically, at E11.5, we removed outlier cells that had UMI counts

< 5,000 or >60,000 and gene counts <1,000 or >8,000 (determined by the visualization of UMI

counts and gene count distributions). At E13.5, we removed outlier cells that had UMI counts

<2,500 or >30,000 and gene counts >1,500. At E15.5, we removed outlier cells that had UMI

counts >40,000 and gene counts <1,000 or > 6,000. At E17.5, we removed outlier cells that had

UMI counts >25,000 and gene counts <1.000 or > 5,000. At P5, we removed outlier cells that had

UMI counts <1,500 or >60,000 and gene counts <1,500 or >8,000. At P12, we removed outlier

cells that had UMI counts<2,000 or >60,000and gene counts <1,500 or > 8000. Moreover, cells

with high proportions of mitochondrial genes (>10%) or hemoglobin genes (>10%) were filtered

out. Finally, we removed all the cells without any *Pomc* transcript UMI counts. A total of 13,953

cells passed the criteria (E11.5: 1,498, E13.5: 1,796, E15.5: 2,078, E17.5: 1,139, P5: 3,909 and

P12: 3,533) for downstream analysis.

The gene expression level for each cell was normalized by the total expression, multiplied

by a scaling factor of 10,000 followed by log-transformation[48]. The top 2,000 most variable genes

were selected from each age and 60 integration anchors were used to combine all the datasets

together. After scaling and centering the integrated dataset, we performed principal components

(PC) analysis on the data matrix and the top 50 PCs were selected based on JackStraw and

Elbow plots from Seurat Package[19] for data visualization using Uniform Manifold Approximation

and Projection (UMAP) technique. We next constructed a Shared Nearest Neighbor (SNN) graph

by setting an expected number of neighbors to 50. In order to cluster the cells, a SNN modularity

optimization technique within a function FindClusters was used to group cells together with a

resolution parameter 0.1. The identity for each cluster was assigned based on the level of *Pomc*

transcripts and prior knowledge of marker genes. The similar analysis was applied to define

TdDsRed cell clusters (Supplementary Figure 2 and Supplementary Figure 3). In addition to the

above criteria applied to remove unwanted cells, we chose cells with at least one *TdDsRed*

transcript UMI count and used the top 2,000 highly variable genes for principal components

analysis. The top 30 PCs were selected for unsupervised clustering with a resolution of 0.3.

Unsupervised cell clustering from each developmental age (Supplementary Figure 6) was

conducting using the following PC numbers (E11.5: 25; E13.5:24, E15.5: 26, E17.5: 19, P5: 40,

P12: 34) with their corresponding resolutions (E11.5: 0.2; E13.5: 0.6, E15.5: 0.5, E17.5: 0.5, P5:

0.5, P12: 0.5. The UMAP plots, violin plots, feature plots and dot plots were all generated in the

Seurat Package. The average expression levels for each gene within each developmental

subcluster (Supplemental Table 8) were obtained from Seurat and calculated by the average of

[*exp(logNormalized counts) -1*]. The differentially expressed genes in each identified cluster was

identified by the comparison of gene expression levels in a specific cluster to all the other clusters.

Cluster identity comparisons

We took the following steps to compare cluster identities from developmental POMC cells to 34

adult ARC neuronal clusters[9] (Figure 8 and Supplementary Figure 10). 1) We obtained the adult

ARC single cell sequencing data from https://hemberg-

lab.github.io/scRNA.seq.datasets/mouse/brain[49] and transformed the dataset from

SingleCellExperiment format to Seurat object format. 2) We normalized adult ARC data and

acquired the top 2000 most variable gene for further analysis. 3) The standard datasets

integration method from Seurat package was performed with 50 integration anchors and used to

combine the two datasets. 4) The integrated data were scaled and the top 50 PCs were selected

for unsupervised clustering implemented in Seurat. 5) We chose 50 transfer anchors to predict

the developmental POMC cells' identities based on adult ARC cell clusters. A similar approach

was applied to compare cluster identities between POMC cells and DTDSRED cells

(Supplementary Figure 2) with the top 50 anchors for data integration and 50 anchors to transfer

cell identities from POMC cells to DSRED cells. We also applied this approach to conduct age-

pairwise comparisons (Supplementary Figure 6) with default setting from Seurat V3.1.5 package.

Cell lineage construction

Cell lineage was constructed using the Destiny (version 2.14.0) package, which implements the

formulation of diffusion maps[50]. Diffusion maps are a spectral method for non-linear dimension

reduction, which is especially suitable for analyzing single-cell gene expression data from different

time-courses. We removed VLMC, glial cells and astrocytes (cluster 9, 10 and 11 in Figure1) and

kept only neurons at embryonic stages (E11.5, E13.5, E15.5 and E17.5), resulting in a total of

6,005 cells for this analysis. The log-transformed normalized data from Seurat data slot and the

corresponding annotation information were imported to construct an expression matrix. The

diffusion maps were constructed under the default setting for the Gaussian kernel width sigma (σ)

and 300 nearest neighbors. The first two Diffusion Components (DCs) were used to visualize the

results. The 3D plots were produced using rgl package (version 0.100.50) with the top three DCs.

Tissue collection and Immunofluorescence staining

Depending on developmental stage, embryos at E10.5 – E17.5 were fixed in 4%

paraformaldehyde in phosphate buffer solution (PBS) at 4 °C for a various period of time.

Specifically, embryos at E10.5 to E12.5 were fixed for 1h; heads from embryos at E15.5-E16.6

and E17.5 were fixed for 2 h and 4 h respectively. Tissues were stabilized in 10% sucrose/10%

gelatin in PBS at 37 °C for 30 min prior to embedding in O.C.T. compound as described

previously[9]. Embryonic brains were sectioned sagittally on a cryostat (Leica CM1950) at 20 μm

thickness. For dual immunofluorescence, after washing excess O.C.T. compound with PBS, a

heat-induced antigen retrieval process was performed using citrate buffer (10 mM anhydrous citric

acid and 0.05% Tween-20, pH 6.0) at 80 °C for 30 min followed by two washes in PBS. Sections

were then blocked with 3% normal goat serum and 0.1% Triton X-100 for 1 hour and incubated

with primary antibodies at room temperature overnight. After PBS washes, sections were

incubated with goat secondary antibodies for 2 hours at room temperature. Nuclei were stained

with DAPI (1 mg/l) for 10 min and the slides washed five times with PBS before mounting with

ProLong™ Gold Antifade Mountant (ThermoFisher). Sections were imaged on a Nikon 90i

fluorescence microscope with NIS-Elements software. Information on the sources and dilution of

antibodies are listed in Supplementary Table 10.

***In situ* hybridization**

Fluorescence *in situ* hybridization (FISH) was performed using RNAscope® Multiplex Fluorescent

V2 Assay (Advanced Cell Diagnostics) according to the manufacturer's instructions with slight

modifications. Specifically, after washing with PBS, sections were post-fixed with 4%

paraformaldehyde for 20 minutes, and dehydrated in a series of ethanol solutions (50%, 70% and

100%). After drying, dehydrated sections were incubated in RNAscope® Hydrogen Peroxide

solution for 10 mins followed by the protease treatment using RNAscope® Protease IV for 20 mins.

Sections were then hybridized to target RNA probes for 2 hours in the HybEZ™ II hybridization

oven. Hybridized probes were amplified using a cascade of signal amplification solutions (AMP

1-3) followed by standard signal developing protocol as described in the manufacturer's brochure.

Detailed information on RNA probes and dilutions are listed in Supplementary Table10. Confocal

images were obtained using a Nikon Instruments A1 Confocal Laser Microscope with NIS-

Elements software.

Figure 1. Single cell sequencing of $Pomc^{DsRed}$ cells from developing mouse hypothalami at E11.5,

E13.5, E15.5, E17.5, P5 and P12 reveals eight distinct neuronal clusters and three non-neuronal

clusters. (A) Schematic diagram showing the overall experimental procedure. (B) UMAP (Uniform

Manifold Approximation and Projection) plot showing the distribution of major cell clusters and

their corresponding cell% at each developmental stage. (C) Violin plots showing the expression

of three signature genes and *Pomc* for each cluster. (D) UMAP plots showing the enrichment of

feature genes in each cluster. Genes representing each cluster were colored and highlighted to

show the cluster-specific enrichment. The color intensity corresponds to the normalized gene

expression from Seurat analysis, where gene expression for each cell was normalized by the total

expression, multiplied by a scale factor of 10,000, and then log-transformed. VLMC, vascular and

leptomeningeal cell.

Figure 2. Temporal gene expression patterns of the *Pomc^high^/Prdm12* cluster. (A) UMAP plot

highlights cells from the *Pomc^high^/Prdm12* cluster. (B) Violin plots showing the expression of

signature genes at each developmental stage. Fluorescence *in situ* hybridization showing the co-

localization of *Pomc* (green) and *Six6* (red) (C), and the co-localization of *Pomc* (green) and *Six3*

(red) (D) at indicated developmental stages. (E) Immunofluorescence showing the co-localization

TDTOMATO (red) NR5A1 (green) at the indicated developmental stages. Insets are magnified

views of the indicated boxes. Arrows indicate co-expressing neurons in the merged panels. Scale

bar: 50 µm. Image orientation: left, posterior(P); right, anterior(A). RP: Rathke's pouch; Pit:

pituitary gland. Arc: arcuate nucleus; VMH: ventromedial hypothalamus.

Figure 3. Temporal gene expression patterns of the *Pomc^med^/Ebf1* cluster. (A) UMAP plot

highlights cells from *Pomc^med^/Ebf1* cluster. (B) Violin plots showing the expression of signature

genes at each developmental stage. (C) Fluorescence *in situ* hybridization showing the co-

localization of *Pomc* (green) and *Ebf1* (red). (D) Fluorescence *in situ* hybridization showing the

co-localization of *Pomc* (green) and *Pou4f1* (red). (E) Re-clustering cells from the *Pomc^med^/Ebf1*

cluster reveals two sub-clusters denoted as 2-1 *Pomc^high^* and 2-2 *Pomc^low^*. Violin plots showing

the signature gene expression in 2-1 *Pomc^high^* and 2-2 *Pomc^low^* clusters. Insets are magnified

views of the indicated boxes. Arrows indicate co-expressing neurons in the merged panels. Scale

bar: 50 μm. Image orientation: left, posterior(P); right, anterior(A). Pit: pituitary gland. Mes:

mesenchyme.

Figure 4. Temporal gene expression patterns of the *Pomc^med^/Nr4a2* and *Pomc^low^/Pitx2* clusters.

(A) UMAP plot highlights cells from the *Pomc^med^/Nr4a2* cluster. (B) Violin plots showing the

expression of signature genes in the *Pomc^med^/Nr4a2* cluster at each developmental stage. (C)

Fluorescence *in situ* hybridization showing the co-localization of *Pomc* (green) and *Nr4a2* (red).

(D) Re-clustering cells from the *Pomc^med^/Nr4a2* cluster reveals two sub-clusters denoted as 3-1

*Pomc^low^* and 3-2 *Pomc^high^* clusters. Violin plot showing the signature gene expression in 3-1

*Pomc^low^* and 3-2 *Pomc^high^* clusters. (E) UMAP plot highlights cells from the *Pomc^low^/Pitx2* cluster.

(F) Violin plots showing the expression of signature genes at each developmental stage in the

Pomc^low^/Pitx2 cluster. (G) Fluorescence *in situ* hybridization showing the co-localization of *Pomc*

(green) and *Pitx2* (red). Scale bar: 50 μm. Insets are magnified views of the indicated boxes.

Arrows indicate co-expressing neurons in the merged panels. Scale bar: 50 μm. Image

orientation: left, posterior (P); right, anterior (A). RP: rathke's pouch; Pit: pituitary gland.

Figure 5. Temporal gene expression patterns of the *Pomc^low^/Otp* cluster. (A) UMAP plot highlights

cells from the *Pomc^low^/Otp* cluster. (B) Violin plots showing the expression of signature genes at

each developmental stage. (C) Fluorescence *in situ* hybridization showing the co-localization of

Pomc (green) and *Npy* (red) at E15.5. Insets are magnified views of the indicated boxes. Arrows

indicate co-expressing neurons in the merged panels. (D) UMAP plots showing cells in the

Pomc^low^/Otp cluster expressing *Sst*, *Agrp*, *Npy*, *Sst*/*Agrp*, or *Sst*/ *Npy*. Cells expressing *Sst* and

cells expressing both *Agrp*/*Npy* are from two distinct cell populations. Scale bar: 50 μm. Image

orientation: left, posterior; right, anterior; Pit: pituitary gland

Figure 6. Temporal gene expression patterns of the *Pomc^low^/Tac2* cluster. (A) UMAP plot

highlights cells from the *Pomc^low^/Tac2* cluster. (B) Violin plots showing the expression of signature

genes at each developmental stage. Fluorescence *in situ* hybridization showing the co-

localization of *Pomc* (green) and *Tac2* (red) (C), and the co-localization of *Pomc* (green) and

Tacr3 (red) (D) at the indicated developmental stages. (E) Fluorescence *in situ* hybridization

showing the co-localization of *Pomc* (green) and *Sox14* (red). (F) Immunofluorescence showing

the co-localization of TDTOMATO (red) and ESR1 (green) at the indicated developmental stages.

Insets are magnified views of the indicated boxes. Arrows indicate co-expressing neurons in the

merged panels. Scale bar: 50 μm. Image orientation: left, posterior; right, anterior; Pit: pituitary

gland.

Figure 7. Diffusion maps analysis illustrating the developmental trajectories of hypothalamic

neuronal clusters. (A) UMAP plot showing the distribution of cells in each cluster from their

corresponding developmental stages. (B) UMAP plots showing the splitting of neuronal clusters

by developmental stages. Clusters were dash-dot circled to show the emerging / evolving

progress through development. (C-D) Diffusion map showing the cell lineages of POMC neurons

during embryogenesis. Cells are colored on the basis of their developmental stage (C) or based

on their previously defined clusters (D). (E) Diffusion maps showing the expression of selected

genesaccording to previous characterization of the feature genes representing each cluster. The

color intensity corresponds to the normalized gene expression from Seurat analysis.

Figure 8. Projection of embryonic POMC[DsRed] cell subtypes to adult arcuate nucleus neuronal

clusters. Projected cell clusters from the *Campbell et al.* study [14] were labeled with the same color

as the POMC[DsRed] cells clusters. The adult ARC neuronal clusters highlighted in bold font indicate

a higher percentage of cells projected to this cluster from the corresponding developmental

clusters. The detailed cell embedding and percentage of cells projecting to adult Arc neuronal

clusters are shown in Supplemental Figure 10.

Supplemental Data

Figure S1

Heat map showing the expression of the top 30 marker genes defining each cluster from Figure 1. M: microglia; V: vascular and leptomeningeal cells; A: astrocytes.

Figure S2

Analysis of POMC cells with *DsRed* transcripts reveals 9 distinct cell clusters. (A) UMAP plot showing the distribution of major cell clusters. (B) Violin plots showing signature gene expression in each cell type. Three feature genes were chosen to represent each cell type. (C) UMAP plots showing the enrichment of feature genes in each cluster. Genes representative of each cluster were colored and highlighted to show the cluster-specific enrichment. The color intensity corresponds to the normalized gene expression from Seurat analysis. (D) Identification of similar cell clusters between POMC cells with *DsRed* transcripts and POMC cell clusters. Red: POMC cell clusters are from Figure 1. Blue: POMC cells with *DsRed* transcripts from Supplemental Figure 2A. (E) Venn diagram showing the number of cells expressing *Pomc* transcripts, *DsRed* transcripts or both *Pomc* and *DsRed* transcripts. Pie chart showing the percentage of POMC cells expressing *DsRed* transcripts at each developmental stage. (F) Vertical bar-chart showing the percentage of cells with *DsRed* transcripts projected to corresponding POMC cell clusters. VLMC, vascular and leptomeningeal cell.

Figure S3

Overview of cell clusters from POMC cells with *DsRed* transcripts. (A) UMAP plot showing the distribution of major cell clusters and their corresponding cell% from each developmental stage. (B) UMAP plots showing the number of cells at each developmental stage. (C) UMAP plots showing the splitting of clusters by developmental stages.

Figure S4

Temporal gene expression patterns of the *Pomc^low^/Dlx5* and *Pomc^low^/Prdm13* clusters. (A) UMAP

plot highlights cells from *Pomc^low^/Dlx5* cluster. (B) Violin plots showing the expression of signature

genes at each developmental stage in the *Pomc^low^/Dlx5* cluster. (C) Fluorescence *in situ*

hybridization showing the co-localization of *Pomc* (green) and *Dlx5* (red). (D) UMAP plot highlights

cells from the *Pomc^low^/Prdm13* cluster. (E) Violin plots showing the expression of signature genes

at each developmental stage in the *Pomc^low^/Prdm13* cluster. (F) Fluorescence *in situ* hybridization

showing the co-localization of *Pomc* (green) and *Prdm13* (red). Insets are magnified views of the

indicated boxes. Arrows indicate co-expressing neurons in the merged panels. Scale bar: 50 μm.

Image orientation: left, posterior; right, anterior; Pit: pituitary gland.

Figure S5

Distribution of GABAergic and glutamatergic neurons in POMC^DsRed^ cell clusters (A) UMAPs plot

showing the distribution of cells with *Gad1* or *Vglut2* gene expression. (B) Violin plots showing

expression of the GABAergic or glutamatergic marker genes in each neuronal cluster.

Figure S6

UMAP plots showing the distribution of major cell clusters at each developmental stage. (A)

E11.5, (B) E13.5, (C) E15.5, (D) E17.5, (E) P5 and (F) P12. The developmental subclusters

derived from each primary cluster are colored to corresponding cells in the UMAP plots.

Figure S7

POMC^DsRed^ developmental cell subclusters shown in Supplemental Figure 6 are projected to the

original integrated cell clusters from Figure 1. The projections were based on the Seurat analysis

and manual comparisons of feature genes in each subcluster (Supplemental Table 2 and

supplemental Table5). The clusters from individual stages leading to the same final integrated

clusters are labeled with the same color. Question marks indicate uncertainties of individual

projections.

Figure S8

Overview of neuropeptide, secretory granule and processing enzyme gene expression patterns

across clusters and developmental ages. The size of dots indicates the percentage of cells

expressing the specific transcripts. The color intensity corresponds to the levels of normalized

gene expression.

Figure S9 A and B

Overview of G-protein coupled receptor (GPCR) gene expression patterns across clusters and

developmental ages. The size of dots indicates the percentage of cells expressing the specific

transcripts. The color intensity corresponds to the levels of normalized gene expression. The list

of genes was acquired from https://www.guidetopharmacology.org/targets.jsp and Gene

Ontology (GO:0005576).

Figure S10

Identification of similar clusters between the neuronal POMC[DsRed] developmental clusters and the

neuronal clusters derived from adult arcuate nucleus[9]. (A) UMAP plots showing the integration of

POMC[DsRed] clusters and adult arcuate nucleus neuronal clusters. (B) The percentage of

POMC[DsRed] cells projecting to corresponding adult arcuate nucleus neuronal clusters.

Figure S11

Fluorescence-activated cell sorting (FACS) strategy for collecting the single POMC-TdDsRed

cells used for scRNAseq.

Table S1 (corresponding to Figure 1)

Average gene expression in each defined POMC[DsRed] clusters. Average gene expression was

obtained from Seurat analysis, which was calculated by the average of *exp(logNormalized counts)*

-1 in each gene within each cluster.

Table S2 (corresponding to Figure 1)

Differentially expressed genes in each identified POMC[DsRed] cluster. For each cluster, the fold-

change values for each gene was calculated by the comparison of expression level in a specific

cluster to the rest of other clusters. The data was sorted by average_logFC > 0.25. pct.1: the

percentage of cells expressed the gene in the defined cluster; pct.2: the percentage of cells

expressed the gene in other clusters; p-val: p-value; p_val_adj: adjusted p-value. Wilcox test was

performed to identify differentially expressed genes in each cluster.

Table S3 (corresponding to Supplemental Figure S2)

Average gene expression in each defined DsRed clusters. Average gene expression was

obtained from Seurat analysis, which was calculated by the average of *exp(logNormalized counts)*

-1 in each gene within each cluster.

Table S4 (corresponding to Supplemental Figure S2)

Differentially expressed genes in each identified DsRed cluster. For each cluster, the fold-change

values for each gene was calculated by the comparison of expression level in a specific cluster

to the rest of other clusters. The data was sorted by average_logFC > 0.25. pct.1: the percentage

of cells expressed the gene in the defined cluster; pct.2: the percentage of cells expressed the

gene in other clusters; p-val: p-value; p_val_adj: adjusted p-value. Wilcox test was performed to

identify differentially expressed genes in each cluster.

Table S5 (corresponding to Figures 2 –6)

Average gene expression at individual developmental ages for each integrated cluster. Average

gene expression was obtained from Seurat analysis, which was calculated by the average of

*exp(logNormalized counts) -1*for each gene within each cluster.

Table S6 (corresponding to Figures 3 and 4)

Average gene expression in subclusters 2-1, 2-2, 3-1 and 3-2. Average gene expression was

obtained from Seurat analysis, which was calculated by the average of *exp(logNormalized counts)*

-1 in each gene within each cluster.

Table S7 (corresponding to Figures 3 and 4)

Differentially expressed genes in subclusters 2-1, 2-2, 3-1 and 3-2 derived from integrated

clusters 2 and 3, respectively. The data were sorted by average_logFC > 0.25. pct.1: the

percentage of cells expressed the gene in the defined cluster; pct.2: the percentage of cells

expressed the gene in other clusters; p-val: p-value; p_val_adj: adjusted p-value. Wilcox test was

performed to identify differentially expressed genes in each cluster.

Table S8 (corresponding to Supplemental Figure 6)

Average gene expression in each defined cluster from the individual developmental ages.

Average gene expression was obtained from Seurat analysis, which was calculated by the

average of [*exp(logNormalized counts) -1*] for each gene within each cluster.

Table S9 (corresponding to Supplemental Figure 6)

Differentially expressed genes in each identified cluster from the individual developmental ages.

For each cluster, the fold-change values for each gene was calculated by the comparison of

expression level in a specific cluster to the rest of other clusters. The data was sorted by

average_logFC > 0.25. pct.1: the percentage of cells expressed the gene in the defined cluster;

pct.2: the percentage of cells expressed the gene in other clusters; p-val: p-value; p_val_adj:

adjusted p-value. Wilcox test was performed to identify differentially expressed genes in each

cluster.

Table S10

Table of key resources.

References

1. Bumaschny, V.F.*, et al.* Obesity-programmed mice are rescued by early genetic intervention. *J Clin Invest* **122**, 4203-4212 (2012).
2. Krude, H.*, et al.* Severe early-onset obesity, adrenal insufficiency and red hair pigmentation caused by POMC mutations in humans. *Nat Genet* **19**, 155-157 (1998).
3. Parikh, D.*, et al.* Stress-induced analgesia and endogenous opioid peptides: the importance of stress duration. *Eur J Pharmacol* **650**, 563-567 (2011).
4. Nasif, S.*, et al.* Islet 1 specifies the identity of hypothalamic melanocortin neurons and is critical for normal food intake and adiposity in adulthood. *Proc Natl Acad Sci U S A* **112**, E1861-1870 (2015).
5. Orquera, D.P.*, et al.* The Homeodomain Transcription Factor NKX2.1 Is Essential for the Early Specification of Melanocortin Neuron Identity and Activates Pomc Expression in the Developing Hypothalamus. *J Neurosci* **39**, 4023-4035 (2019).
6. Hael, C.E., Rojo, D., Orquera, D.P., Low, M.J. & Rubinstein, M. The transcriptional regulator PRDM12 is critical for Pomc expression in the mouse hypothalamus and controlling food intake, adiposity, and body weight. *Mol Metab* **34**, 43-53 (2020).
7. Padilla, S.L., Carmody, J.S. & Zeltser, L.M. Pomc-expressing progenitors give rise to antagonistic neuronal populations in hypothalamic feeding circuits. *Nat Med* **16**, 403-405 (2010).
8. Sanz, E.*, et al.* Fertility-regulating Kiss1 neurons arise from hypothalamic POMC-expressing progenitors. *J Neurosci* **35**, 5549-5556 (2015).
9. Campbell, J.N.*, et al.* A molecular census of arcuate hypothalamus and median eminence cell types. *Nat Neurosci* **20**, 484-496 (2017).
10. Chen, R., Wu, X., Jiang, L. & Zhang, Y. Single-Cell RNA-Seq Reveals Hypothalamic Cell Diversity. *Cell Rep* **18**, 3227-3241 (2017).
11. Lam, B.Y.H.*, et al.* Heterogeneity of hypothalamic pro-opiomelanocortin-expressing neurons revealed by single-cell RNA sequencing. *Mol Metab* **6**, 383-392 (2017).
12. Huisman, C.*, et al.* Author Correction: Single cell transcriptome analysis of developing arcuate nucleus neurons uncovers their key developmental regulators. *Nat Commun* **11**, 1359 (2020).
13. Bouret, S.G., Draper, S.J. & Simerly, R.B. Formation of projection pathways from the arcuate nucleus of the hypothalamus to hypothalamic regions implicated in the neural control of feeding behavior in mice. *J Neurosci* **24**, 2797-2805 (2004).
14. Bouret, S.G., Draper, S.J. & Simerly, R.B. Trophic action of leptin on hypothalamic neurons that regulate feeding. *Science* **304**, 108-110 (2004).
15. Bouret, S.G., Bates, S.H., Chen, S., Myers, M.G., Jr. & Simerly, R.B. Distinct roles for specific leptin receptor signals in the development of hypothalamic feeding circuits. *J Neurosci* **32**, 1244-1252 (2012).
16. Quarta, C.*, et al.* Functional identity of hypothalamic melanocortin neurons depends on Tbx3. *Nat Metab* **1**, 222-235 (2019).
17. Ikeda, Y., Luo, X., Abbud, R., Nilson, J.H. & Parker, K.L. The nuclear receptor steroidogenic factor 1 is essential for the formation of the ventromedial hypothalamic nucleus. *Mol Endocrinol* **9**, 478-486 (1995).
18. Luo, X., Ikeda, Y. & Parker, K.L. A cell-specific nuclear receptor is essential for adrenal and gonadal development and sexual differentiation. *Cell* **77**, 481-490 (1994).
19. Meeks, J.J., Weiss, J. & Jameson, J.L. Dax1 is required for testis determination. *Nat Genet* **34**, 32-33 (2003).
20. Mizusaki, H.*, et al.* Dax-1 (dosage-sensitive sex reversal-adrenal hypoplasia congenita critical region on the X chromosome, gene 1) gene transcription is regulated by wnt4 in the female developing gonad. *Mol Endocrinol* **17**, 507-519 (2003).
21. Kovalovsky, D.*, et al.* Activation and induction of NUR77/NURR1 in corticotrophs by CRH/cAMP: involvement of calcium, protein kinase A, and MAPK pathways. *Mol Endocrinol* **16**, 1638-1651 (2002).

22. Perlmann, T. & Wallen-Mackenzie, A. Nurr1, an orphan nuclear receptor with essential functions
in developing dopamine cells. *Cell Tissue Res* **318**, 45-52 (2004).
23. Hong, S.*, et al.* Functional roles of Nurr1, Pitx3, and Lmx1a in neurogenesis and phenotype
specification of dopamine neurons during in vitro differentiation of embryonic stem cells. *Stem Cells Dev*
23, 477-487 (2014).
24. Jankovic, J., Chen, S. & Le, W.D. The role of Nurr1 in the development of dopaminergic neurons
and Parkinson's disease. *Prog Neurobiol* **77**, 128-138 (2005).
25. Saucedo-Cardenas, O.*, et al.* Nurr1 is essential for the induction of the dopaminergic phenotype
and the survival of ventral mesencephalic late dopaminergic precursor neurons. *Proc Natl Acad Sci U S A*
95, 4013-4018 (1998).
26. Smits, S.M., Ponnio, T., Conneely, O.M., Burbach, J.P. & Smidt, M.P. Involvement of Nurr1 in
specifying the neurotransmitter identity of ventral midbrain dopaminergic neurons. *Eur J Neurosci* **18**,
1731-1738 (2003).
27. Andersson, E.*, et al.* Identification of intrinsic determinants of midbrain dopamine neurons. *Cell*
124, 393-405 (2006).
28. Murphy, E.P. & Conneely, O.M. Neuroendocrine regulation of the hypothalamic pituitary adrenal
axis by the nurr1/nur77 subfamily of nuclear receptors. *Mol Endocrinol* **11**, 39-47 (1997).
29. Kee, N.*, et al.* Single-Cell Analysis Reveals a Close Relationship between Differentiating Dopamine
and Subthalamic Nucleus Neuronal Lineages. *Cell Stem Cell* **20**, 29-40 (2017).
30. Huisman, C.*, et al.* Single cell transcriptome analysis of developing arcuate nucleus neurons
uncovers their key developmental regulators. *Nat Commun* **10**, 3696 (2019).
31. Lee, B.*, et al.* Dlx1/2 and Otp coordinate the production of hypothalamic GHRH- and AgRP-
neurons. *Nat Commun* **9**, 2026 (2018).
32. German, C.L., Baladi, M.G., McFadden, L.M., Hanson, G.R. & Fleckenstein, A.E. Regulation of the
Dopamine and Vesicular Monoamine Transporters: Pharmacological Targets and Implications for Disease.
Pharmacol Rev **67**, 1005-1024 (2015).
33. Wittmann, G., Hrabovszky, E. & Lechan, R.M. Distinct glutamatergic and GABAergic subsets of
hypothalamic pro-opiomelanocortin neurons revealed by in situ hybridization in male rats and mice. *J*
Comp Neurol **521**, 3287-3302 (2013).
34. Jones, G.L.*, et al.* Selective Restoration of Pomc Expression in Glutamatergic POMC Neurons:
Evidence for a Dynamic Hypothalamic Neurotransmitter Network. *eNeuro* **6** (2019).
35. Chen, X.*, et al.* Comparative Transcriptomic Analyses of Developing Melanocortin Neurons Reveal
New Regulators for the Anorexigenic Neuron Identity. *J Neurosci* **40**, 3165-3177 (2020).
36. Toll, L., Bruchas, M.R., Calo, G., Cox, B.M. & Zaveri, N.T. Nociceptin/Orphanin FQ Receptor
Structure, Signaling, Ligands, Functions, and Interactions with Opioid Systems. *Pharmacol Rev* **68**, 419-457
(2016).
37. Srisai, D.*, et al.* MRAP2 regulates ghrelin receptor signaling and hunger sensing. *Nat Commun* **8**,
713 (2017).
38. Chaly, A.L., Srisai, D., Gardner, E.E. & Sebag, J.A. The Melanocortin Receptor Accessory Protein 2
promotes food intake through inhibition of the Prokineticin Receptor-1. *Elife* **5** (2016).
39. Japon, M.A., Rubinstein, M. & Low, M.J. In situ hybridization analysis of anterior pituitary hormone
gene expression during fetal mouse development. *J Histochem Cytochem* **42**, 1117-1125 (1994).
40. Liu, W. & Cvekl, A. Six3 in a small population of progenitors at E8.5 is required for neuroretinal
specification via regulating cell signaling and survival in mice. *Dev Biol* **428**, 164-175 (2017).
41. Liu, W., Lagutin, O., Swindell, E., Jamrich, M. & Oliver, G. Neuroretina specification in mouse
embryos requires Six3-mediated suppression of Wnt8b in the anterior neural plate. *J Clin Invest* **120**, 3568-
3577 (2010).
42. Suh, C.S.*, et al.* Autoregulatory binding sites in the zebrafish six3a promoter region define a new
recognition sequence for Six3 proteins. *FEBS J* **277**, 1761-1775 (2010).

43.   Geng, X*., et al.* Haploinsufficiency of Six3 fails to activate Sonic hedgehog expression in the ventral
forebrain and causes holoprosencephaly. *Dev Cell* **15**, 236-247 (2008).
44.   Larder, R., Clark, D.D., Miller, N.L. & Mellon, P.L. Hypothalamic dysregulation and infertility in mice
lacking the homeodomain protein Six6. *J Neurosci* **31**, 426-438 (2011).
45.   Kim, D.W*., et al.* The cellular and molecular landscape of hypothalamic patterning and
differentiation from embryonic to late postnatal development. *Nat Commun* **11**, 4360 (2020).
46.   Burgunder, J.M. & Young, W.S., 3rd. Neurons containing cholecystokinin mRNA in the mammillary
region: ontogeny and adult distribution in the rat. *Cell Mol Neurobiol* **9**, 281-294 (1989).
47.   Yin, M*., et al.* Ventral mesencephalon-enriched genes that regulate the development of
dopaminergic neurons in vivo. *J Neurosci* **29**, 5170-5182 (2009).
48.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic
data across different conditions, technologies, and species. *Nat Biotechnol* **36**, 411-420 (2018).
49.   Kiselev, V.Y., Yiu, A. & Hemberg, M. scmap: projection of single-cell RNA-seq data across data sets.
Nat Methods **15**, 359-362 (2018).
50.   Angerer, P*., et al.* destiny: diffusion maps for large-scale single-cell data in R. *Bioinformatics* **32**,
1241-1243 (2016).

1002

Figure 1A

Timeline of hypothalami collection

Cell Isolation · **FACS** · **Library Prep** · **Sequencing** · **Analysis**

E11.5 E13.5 E15.5 E17.5 P5 P12

B

Microglia

Astrocytes

VLMC

Percentage of Total Neurons

	E11.5	E13.5	E15.5	E17.5	P5	P12
1. $Pomc^{high}/Prdm12$	50.8%	40.6%	34.7%	33.3%	44.1%	47.2%
2. $Pomc^{med}/Ebf1$	43.4%	31.1%	11.8%	13.7%	13.8%	14.4%
3. $Pomc^{med}/Nr4a2$	0.5%	3.4%	5.9%	1.1%	5.6%	9.1%
4. $Pomc^{low}/Otp$	2.4%	6.0%	20.7%	20.9%	2.8%	4.2%
5. $Pomc^{low}/Tac2$	1.2%	3.3%	10.7%	24.6%	29.1%	16.0%
6. $Pomc^{low}/Dlx5$	0.5%	1.0%	6.2%	4.9%	3.9%	5.7%
7. $Pomc^{low}/Prdm13$	0.1%	1.4%	3.3%	1.1%	0.2%	0.4%
8. $Pomc^{low}/Pitx2$	1.0%	13.2%	6.8%	0.4%	0.6%	2.9%

C

D

Figure 3 A

Figure 4 A

Figure 5

Figure 6

Figure 8

Developmental POMC⁺ Cells		Adult ARC Neurons
1.Pomc^high^/Prdm12	⋙	**n14. *Pomc/Ttr*** n15. *Pomc/Anxa2*
2.Pomc^med^/Ebf1--1	⋙	**n14. *Pomc/Ttr*** **n15. *Pomc/Anxa2***
2.Pomc^med^/Ebf1--2	⋙	**n34. unassigned2** n03. *Th/Sst*
3.Pomc^med^/Nr4a2--1	⋙	**n33. unassigned1** n34. unassigned2
3.Pomc^med^/Nr4a2--2	⋙	**n14. Pomc/Ttr** n15. Pomc/Anxa2
4.Pomc^low^/Otp	⋙	**n13. *Agrp/Gm8773*** n23. *Sst/Unc13c*
5.Pomc^low^/Tac2	⋙	**n20. *Kiss1/Tac2***
6.Pomc^low^/Dlx5	⋙	**n22. *Tmem215*** n34. unassigned2
7.Pomc^low^/Prdm13	⋙	**n29. *Nr5a1/Adcyap1*** n20. *Kiss1/Tac2*
8.Pomc^low^/Pitx2	⋙	**n34. unassigned2** n32. *Slc17a6/Trhr*