The cyclic dinucleotide 2'3'-cGAMP induces a broad anti-bacterial and anti-viral response in the sea anemone *Nematostella vectensis*

Shally R. Margolis[1], Peter A. Dietzen[1], Beth M. Hayes[2], Stephen C. Wilson[1,3], Brenna C. Remick[1], Seemay Chou[2,4], Russell E. Vance[1,5,6*]

[1] Division of Immunology and Pathogenesis, Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720 USA
[2] Department of Biochemistry and Biophysics, University of California, San Francisco, San Francisco, CA 94158, USA.
[3] Current Address: Bristol Myers Squibb, 200 Cambridgepark Dr., Cambridge, MA 02140
[4] Chan Zuckerberg Biohub, San Francisco, CA
[5] Cancer Research Laboratory and the Immunotherapeutics and Vaccine Research Initiative, University of California, Berkeley, CA 94720
[6] Howard Hughes Medical Institute, University of California, Berkeley, CA 94720
* correspondence: rvance@berkeley.edu

Author Contributions: S.R.M., P.A.D., B.M.H., S.C.W., S.C. and R.E.V. designed research; S.R.M., P.A.D., B.M.H., S.C.W., and B.C.R. performed research; S.R.M., P.A.D., and B.M.H. analyzed data; S.R.M. and R.E.V. wrote the paper; and P.A.D., B.M.H., S.C.W, and S.C. provided critical review.
Competing Interest Statement: R.E.V. consults for Ventus Therapeutics and Tempest Therapeutics. The other authors declare no competing interests.
Preprint Servers: https://doi.org/10.1101/2021.05.13.443009.
Available under CC-BY 4.0.
Classification: Biological Sciences; Immunology and Inflammation
Keywords: STING, Cnidaria, *Nematostella vectensis*, NF-κB, innate immunity

Abstract

In mammals, cyclic dinucleotides (CDNs) bind and activate STING to initiate an anti-viral type I interferon response. CDNs and STING originated in bacteria and are present in most animals. By contrast, interferons are believed to have emerged in vertebrates; thus, the function of CDN signaling in invertebrates is unclear. Here, we use a CDN, 2'3'-cGAMP, to activate immune responses in a model cnidarian invertebrate, the starlet sea anemone *Nematostella vectensis*. Using RNA-Seq, we found that 2'3'-cGAMP induces robust transcription of both anti-viral and anti-bacterial genes in *N. vectensis*. Many of the anti-viral genes induced by 2'3'-cGAMP are homologs of vertebrate interferon stimulated genes, implying that the interferon response predates the evolution of interferons. Knockdown experiments identified a role for NF-κB in specifically inducing anti-bacterial genes downstream of 2'3'-cGAMP. Some of these putative anti-bacterial genes were also found to be induced during *Pseudomonas aeruginosa* infection. We characterized the protein product of one of the putative anti-bacterial genes, the *N. vectensis* homolog of Dae4, and found that it has conserved anti-bacterial activity. This work suggests that a broad anti-bacterial and anti-viral transcriptional response is an evolutionarily ancestral output of 2'3'-cGAMP signaling in animals.

Significance statement

Cyclic dinucleotides are signaling molecules that originated in bacteria and were subsequently acquired and co-opted by animals for immune signaling. The major cyclic dinucleotide signaling pathway in mammals results in the production of anti-viral molecules called interferons. Invertebrates such as sea anemones lack interferons, and thus it was unclear whether cyclic dinucleotide signaling would play a role in immunity in these animals. Here we report that in the anemone *Nematostella vectensis*, cyclic dinucleotides activate both anti-viral and anti-bacterial immune responses, and do so through a conserved pathway. These results provide insights into the evolutionary origins of innate immunity, and suggest a broader ancestral role for cyclic dinucleotide signaling that evolved toward more specialized anti-viral functions in mammals.

Introduction

The innate immune system is an evolutionarily ancient system that detects pathogens and initiates their elimination. In mammals, the cGAS-STING pathway is critical for sensing and responding to intracellular DNA, which is particularly important for innate responses to DNA viruses (1, 2). The sensor protein in this pathway, cyclic-GMP-AMP synthase (cGAS), is an enzyme that binds directly to cytosolic DNA and produces 2'3'-cGAMP, a cyclic dinucleotide (CDN) second messenger that binds and activates STING (3-8). Active STING uses its C-terminal tail (CTT) to recruit TBK1, which then phosphorylates and activates the transcription factor IRF3 to induce the expression of type I interferons (IFNs) (9-12). Type I IFNs are secreted cytokines that signal via JAK-STAT signaling to induce transcription of hundreds of anti-viral genes known as interferon-stimulated genes (ISGs) (13, 14). STING also activates NF-κB, MAP kinase (15), STAT6 (16), and autophagy-like pathways (17-20), as well as senescence (21) and cell death (22-26), although the mechanism of activation of these pathways, and their importance during infection, are less well understood.

Type I IFNs are thought to be a relatively recent evolutionary innovation, with identifiable interferon genes found only in vertebrates (27). In contrast, STING and cGAS are conserved in the genomes of most animals and some unicellular choanoflagellates. Remarkably, CDN to STING signaling seems to have originated in bacteria, where it may be important in bacteriophage defense (28, 29). Studies on the function of STING in animals that lack type I IFN have been mostly limited to insects, where STING seems to be protective during viral (30-33), bacterial (34), and microsporidial (35) infection. In insects, STING may promote defense through activation of autophagy (32, 35) and/or induction of NF-κB-dependent defense genes (30, 31, 34). Most ISGs are lacking from insect genomes and are not induced by CDN-STING signaling (31). In addition, the biochemical mechanisms of STING activation and signaling in insects remain poorly understood.

Biochemically, perhaps the best-characterized invertebrate STING is that of the starlet sea anemone, *Nematostella vectensis,* a member of one of the oldest animal phyla (Cnidaria). *N. vectensis* encodes a surprisingly complex genome that harbors many gene families found in vertebrates but absent in other invertebrates such as *Drosophila* (36). *N. vectensis* STING (nvSTING) and human STING adopt remarkably similar conformations when bound to 2'3'-cGAMP, and nvSTING binds to this ligand with high affinity ($K_d < 1nM$) (37). The *N. vectensis* genome also encodes a cGAS enzyme that produces 2'3'-cGAMP in mammalian cell culture (17). In vertebrates, STING requires its extended CTT to initiate transcriptional responses (38, 39); however, nvSTING lacks an extended CTT and thus its signaling mechanism and potential for inducing transcriptional responses is unclear. Based on experiments with nvSTING in mammalian cell lines, CTT-independent induction of autophagy has been proposed as the 'ancestral' function of STING (17), but the endogenous function of STING in *N. vectensis* has never been described.

Despite the genomic identification of many predicted innate immune genes (40, 41), few have been functionally characterized in *N. vectensis*. The sole *N. vectensis* Toll-like receptor (TLR) is reported to bind flagellin and activate NF-κB in human cell lines, and is expressed in cnidocytes, the stinging cells that define cnidarians (42). *N. vectensis* NF-κB (nvNF-κB) binds to conserved κB sites, is inhibited by *N. vectensis* IκB (43), and seems to be required for the development of cnidocytes (44). However, no activators of endogenous nvNF-κB have yet been identified. Recent work probing the putative anti-viral immune response in *N. vectensis* found that double-stranded RNA (dsRNA) injection into *N. vectensis* embryos leads to transcriptional induction of genes involved

in the RNAi pathway as well as genes with homology to ISGs (45). This response is
partially dependent on a RIG-I-like receptor, indicating deep conservation of anti-viral
immunity (45). However, no anti-viral or anti-bacterial effectors from *N. vectensis* have
been functionally tested.
Here, we characterize the response of *N. vectensis* to 2'3'-cGAMP stimulation.
Similar to the response of vertebrates to 2'3'-cGAMP, we find robust transcriptional
induction of putative anti-viral genes with homology to vertebrate ISGs. In addition, we
observed induction of numerous anti-bacterial genes that are not induced during the
vertebrate response to 2'3'-cGAMP. Although we were unable to show that the
response to 2'3'-cGAMP is nvSTING dependent, we did find a selective requirement for
nvNF-κB in the induction of some of the anti-bacterial genes. Many of these genes are
also induced during *Pseudomonas aeruginosa* infection, suggesting a functional role in
anti-bacterial immunity. We selected and characterized the anti-bacterial activity of one
2'3'-cGAMP-induced *Nematostella* gene product, domesticated amidase effector
(nvDae4), a peptidoglycan cleaving enzyme that we found can kill Gram-positive
bacteria. This work demonstrates an evolutionarily ancient role for 2'3'-cGAMP in the
transcriptional induction of both anti-viral and anti-bacterial immunity.
126
Results
Transcriptional response to 2'3'-cGAMP in Nematostella vectensis
To assess the *in vivo* role of 2'3'-cGAMP signaling in *Nematostella vectensis*, we
treated 2-week-old polyps with 2'3'-cGAMP for 24 hours and performed RNA-Seq (Fig.
1A, Fig. S1). Thousands of genes were induced by 2'3'-cGAMP, many of which are
homologs of genes known to function in mammalian immunity. Despite the lack of
associated gene ontology (GO) terms for many of the differentially regulated genes,
unbiased GO term analysis revealed significant enrichment of immune-related terms
(Fig. 1B). We also treated animals with 3'3'-linked cyclic dinucleotides, which are
thought to be produced exclusively by bacteria, and which also bind to nvSTING *in vitro*,
albeit at lower affinity (37). Both 3'3'-cGAMP and cyclic-di-AMP treatment also induced
a smaller number of genes, although all of these genes were induced more strongly by
2'3'-cGAMP (Fig. S1). Interestingly, cyclic-di-GMP treatment led to almost no
transcriptional induction, despite having relatively high affinity for nvSTING *in vitro*. This
discrepancy may be due to differences in cell permeability among different CDNs, as
the ligands were added extracellularly. In order to be able to perform subsequent
genetic experiments, we confirmed that the immune gene induction downstream of by
2'3'-cGAMP also occurred in embryos (which are amenable to microinjection of
shRNAs). Quantitative reverse transcription PCR (qRT-PCR) on 48-hour-embryos
treated for 4 hours with a lower dose of 2'3'-cGAMP revealed that many immune genes
were also induced at this early developmental stage after a much shorter treatment (Fig.
1C).
Several interesting classes of genes were found to be upregulated in response to
2'3'-cGAMP. For example, several genes involved in the RNAi pathway were induced,
including homologs of Argonaute (AGO2), Dicer, and RNA-dependent RNA polymerase
(Rdrp1). In addition, many genes that are considered ISGs in mammals were also
induced in *N. vectensis*, including Viperin, RNase L, 2'-5'-oligoadenylate synthase
(OAS), interferon regulatory factors (IRFs), guanylate-binding proteins (GBPs), and the
putative pattern recognition receptors RIG-I-like receptor a (RLRa) and RLRb. These
results suggest a conserved role for 2'3'-cGAMP signaling in anti-viral immunity and ISG
induction, despite an apparent lack of conservation of type I interferons in *N. vectensis*.
Interestingly, we also found that many putative anti-bacterial genes were upregulated in

response to 2'3'-cGAMP, including homologs of LPS-binding protein (LBP), lysozyme,
perforin-2, Dae4, and mucins. These results indicate that 2'3'-cGAMP stimulation leads
to a broad immune response in *N. vectensis*.
To determine whether 2'3'-cGAMP signaled via nvSTING to induce these genes,
we injected shRNAs targeting nvSTING into 1-cell embryos and treated with 2'3'-
cGAMP 48 hours later. We extracted RNA and performed RNA-Seq on these samples,
and surprisingly, while nvSTING transcripts were reduced by ~50%, there was no
significant impact on 2'3'-cGAMP-induced gene expression (Fig. S2A, S2B). These
negative results were recapitulated in numerous independent qRT-PCR and Nanostring
experiments using 9 different shRNAs (3 shown in Fig. S2C). There are several
possible explanations for the failure to observe a requirement for nvSTING in 2'3'-
cGAMP signaling : (1) a 2-fold reduction in STING transcript levels may not result in a
reduction in STING protein levels if the protein is very stable; (2) even if STING protein
levels are reduced 2-fold, the reduction may not affect STING signaling due to threshold
effects; or (3) nvSTING may not be required for signaling downstream of 2'3'-cGAMP
due the presence of a redundant 2'3'-cGAMP sensor in *N. vectensis*. We generated an
anti-nvSTING antibody to validate knockdown efficiency at the protein level, but this
reagent did not appear to specifically detect nvSTING in anemone lysates. We also
tested whether an nvSTING translation-blocking morpholino could inhibit induction of
genes in response to 2'3'-cGAMP, but this also had no effect (Fig. S2D). Lastly, we
made multiple attempts to generate nvSTING mutant animals using CRISPR, using
multiple different guide RNAs, but the inefficiency of CRISPR in this organism and
issues with mosaicism prevented the generation of nvSTING null animals. We
previously solved the crystal structure of nvSTING bound to 2'3'-cGAMP and showed
that binding occurs with high affinity (Kd < 1nM) and in a similar mode as compared to
vertebrate STING (37). In addition, we found that when expressed in mammalian cells,
nvSTING forms puncta only in the presence of 2'3'-cGAMP, indicating some functional
change induced by this ligand (Fig. S2E). Thus, we hypothesize that 2'3'-cGAMP
signals via nvSTING, but technical issues and possible redundancy with additional
sensors prevent formal experimental evidence for this hypothesis.
189
The N. vectensis NF-κB homolog plays a role in the 2'3'-cGAMP response
We next tested the role of conserved transcription factors that are known to
function downstream of STING mammals in the *N. vectensis* response to 2'3'-cGAMP.
Interestingly, many of these transcription factors are themselves transcriptionally
induced by 2'3'-cGAMP in *N. vectensis* (Fig. 1A). In mammals, the transcription factors
IRF3 and IRF7 induce type I IFN downstream of STING activation. While the specific
function of these IRFs in interferon induction are thought to have arisen in vertebrates,
other IRF family members, with conserved DNA binding residues, are present in *N.*
vectensis (Fig. S3). We microinjected 1-cell embryos with short hairpin RNAs (shRNAs)
targeting each of the 5 nvIRFs or a GFP control, treated with 2'3'-cGAMP, and
assessed gene expression by qRT-PCR and/or Nanostring. Knockdown of IRF
transcripts by 40-60% did not measurably impact gene induction by 2'3'-cGAMP (Fig.
S4). We similarly tested the role of the single *N. vectensis* STAT gene, as mammalian
STATs both induce anti-viral genes downstream of Type I IFN signaling, and may even
be directly activated by STING (16). Similar to the nvIRFs, we did not observe a
significant loss of gene induction by 2'3'-cGAMP in nvSTAT knockdown embryos by
both RNA-Seq and Nanostring (Fig. S4). There are several explanations for these
findings: (1) sufficient IRF or STAT protein may remain in knockdown animals to
transduce the signal, either due to low efficiency of the knockdowns, or to protein

stability; (2) the IRFs may act redundantly with each other, and therefore no effect will
be seen in single knockdown experiments; or (3) nvIRFs and nvSTAT may not play a
role in the response to 2'3'-cGAMP.
NF-κB is also known to act downstream of mammalian STING, and appears to
be functionally conserved in *N. vectensis* (43). We found that NF-κB signaling
components are transcriptionally induced by 2'3'-cGAMP (Fig. 1A). To test the role of
nvNF-κB in the 2'3'-cGAMP response, we microinjected embryos with shRNAs targeting
nvNF-κB, treated with 2'3'-cGAMP, and performed RNA-Seq (Fig. 2A). 241 genes were
transcribed at significantly lower levels in the nvNF-κB knockdown embryos, and of
these, 98 were genes induced by 2'3'-cGAMP. Of these genes, 40 are uncharacterized,
and no GO terms were significantly enriched (data not shown). Of the induced genes
that were annotated in NCBI, we noticed many were homologs of anti-bacterial proteins,
including homologs of perforin-2/Mpeg-1, LPS-binding protein (LBP), linear gramicidin
synthase, and mucins. We confirmed that 2'3'-cGAMP-mediated induction of these
putative anti-bacterial genes was NF-κB-dependent by performing qRT-PCR and
Nanostring (Fig. 2B; Fig. S5A). Of note, the induction of nvLysozyme was not nvNF-κB
dependent (both by RNA-Seq and qRT-PCR; Fig. S5B), indicating either the existence
of another pathway for anti-bacterial gene induction, or that our knockdown experiment
was not able to affect expression of all nvNF-κB dependent genes. In addition, all of the
putative anti-viral genes we examined appeared to be induced independent of nvNF-κB
(Fig. S5).
We performed BLAST searches of unannotated 2'3'-cGAMP-induced, nvNF-κB-
dependent genes and identified several other genes with predicted anti-bacterial
activity, including two homologs of bacterial *tae4* genes, and a putative guanylate
binding protein (GBP) (*N. vectensis* LOC5515806, hereafter nvGBP-806). The Tae4
homologs had been previously identified and will be referred to as nvDae4 proteins
(discussed further below; (46)). To confirm the identity of nvGBP-806 as a true GBP
homolog, we performed phylogenetic analysis. We identified four conserved *N.
vectensis* proteins harboring an N-terminal GBP GTPase domain with conserved GBP-
specific motifs, including nvGBP-806 (Fig. S6). All of the nvGBP homologs cluster with
vertebrate IFN-inducible GBPs and are themselves induced by 2'3'-cGAMP. Finally, we
identified several unannotated nvNF-κB dependent, 2'3'-cGAMP-induced genes that
appeared to be cnidarian-specific with no identifiable homologs in other animal phyla
(Table S1).
To test directly whether nvNF-κB is activated in *N. vectensis* upon 2'3'-cGAMP
treatment, we treated polyps with cGAMP and performed immunostaining for nvNF-κB
(Fig. 2C). Inactive NF-κB is localized to the cytosol, and we observed sparse, cytosolic
staining of ectodermal cells in untreated animals, as has been previously reported (43).
In contrast, in 2'3'-cGAMP treated animals, we found many more nvNF-κB-positive
cells, and in almost all of these, nvNF-κB was found in the nucleus. We performed
automated quantification of nuclear nvNF-κB staining, and found that ~3-20% of nuclei
captured in our images were positive for nvNF-κB (Fig. 2D). In sum, 2'3'-cGAMP leads
to nvNF-κB nuclear localization, and nvNF-κB appears to be required for expression of
many putative anti-bacterial, but not anti-viral, genes. Our results demonstrate the first
NF-kB agonist in *N. vectensis*, and indicate a conserved immune function for NF-kB in
this organism.
255
Gene induction during Pseudomonas aeruginosa challenge
In order to test whether the putative anti-bacterial, NF-κB-dependent genes are
induced during bacterial infection, we infected *N. vectensis* with *Pseudomonas*

aeruginosa, a pathogenic Gram-negative bacterium. *P. aeruginosa* can infect a range of
hosts, including plants, mammals, and hydra (47, 48), though infections of *N. vectensis*
have not previously been reported. Infection of *N. vectensis* polyps with the *P.*
aeruginosa strain PA14 led to polyp death in a dose and temperature dependent
manner (Fig. 3A). 48 hours after infection, we isolated RNA from infected polyps and
assayed gene expression. Interestingly, nvSTING expression was induced during PA14
infection (Fig. 3B), and many of the putative anti-bacterial genes we identified as 2'3'-
cGAMP-induced were also induced during infection (Fig. 3C), although this expression
was not sufficient to protect from death. In addition, some putative anti-viral genes were
also induced in some animals (Fig. S5B), perhaps reflecting a broader immune
response in *Nematostella*. Importantly, the PA14 genome is not known to encode any
proteins that produce CDNs other than c-di-GMP. Since c-di-GMP was not sufficient to
robustly activate gene expression in *N. vectensis*, we believe that it is likely that the
response to PA14 is independent of bacterial CDNs, although we cannot rule out an
effect from PA14-produced c-di-GMP. In addition, we have no reason to believe that
PA14 in activating these genes via nv-cGAS, as we do not know the activator of this
enzyme. Nevertheless, taken together, these results indicate that the putative anti-
bacterial genes we identified as induced by 2'3'-cGAMP are also induced after bacterial
challenge.
278
nvDae4 is a peptidoglycan-cleaving enzyme with anti-bacterial activity
We decided to investigate directly whether any of the genes induced by both 2'3'-
cGAMP and bacterial infection are in fact anti-bacterial. Type VI secretion amidase
effector (Tae) proteins are bacterial enzymes that are injected into neighboring cells to
cleave peptidoglycan, an essential component of bacterial cell walls, leading to rapid
cell death (49). While the *tae* genes originated in bacteria, they have been horizontally
acquired multiple times in evolution by eukaryotes, and at least one of these so-called
"domesticated amidase effectors" (Daes) also has bactericidal activity (46, 50). The *N.*
vectensis genome has two *tae4* homologs, both of which were upregulated by 2'3'-
cGAMP in an nvNF-κB-dependent manner. However, only one of the *N. vectensis* Dae
proteins is predicted to encode a conserved catalytic cysteine (46) required for
peptidoglycan hydrolysis. Therefore, we focused our efforts on this homolog, which we
call nvDae4 (GI: 5507694). We first tested whether nvDae4 has conserved bactericidal
properties by expressing nvDae4 in *E. coli* either with or without a periplasm-targeting
signal sequence and measuring bacterial growth (assessed by OD_{600}) over time (Fig.
4A). *E. coli* are Gram-negative bacteria and thus have peptidoglycan
compartmentalized within the periplasmic space. Consistent with the predicted
peptidoglycan-cleaving function of nvDae4, only periplasmic wild-type (WT) but not
catalytic mutant (C63A) nvDae4 expression led to bacterial lysis. In order to test directly
whether nvDae4 cleaves peptidoglycan, we produced recombinant protein in insect
cells. Since nvDae4 encodes a secretion signal, recombinant nvDae4 was secreted by
the insect cells and purified from the cell supernatant. Purified nvDae4 protein was
incubated with purified peptidoglycan from either *E. coli* or *Staphylococcus epidermis*.
Analysis by high performance liquid chromatography (HPLC) showed that nvDae4
cleaves both Gram-negative (Fig. 4B) and Gram-positive (Fig. S7) derived
peptidoglycan. Finally, we wondered whether nvDae4 could directly kill Gram-positive
bacteria, as these bacteria contain a peptidoglycan cell wall that is not protected by an
outer membrane and is therefore accessible to extracellular factors. We treated *B.*
subtilis with recombinant nvDae4 and found that bacteria treated with WT but not C63A
nvDae4 protein were killed (Fig. 4C) in a dose-dependent manner (Fig. 4D). Overall

these results show that the 2'3'-cGAMP-induced protein nvDae4 is a peptidoglycan-
cleaving enzyme with the capacity to kill bacteria.
311
Discussion
In this study, we identified hundreds of *N. vectensis* genes that are induced by
the STING ligand 2'3'-cGAMP. Despite over 600 million years of divergence and the
absence of interferons, *N. vectensis* responds to 2'3'-cGAMP similarly to mammals by
inducing a variety of anti-viral genes. Similarly, Lewandowska et al. (45) recently
reported that *N. vectensis* responds to the synthetic double-stranded RNA poly(I:C), a
viral mimic and pathogen-associated molecular pattern (PAMP). In *N. vectensis*,
poly(I:C) induced both RNAi pathway components and genes traditionally thought of as
vertebrate ISGs. Our combined findings indicate that the pathways linking PAMP
detection to ISG expression existed prior to the vertebrate innovation of type I IFNs.
Interestingly, some invertebrate species have protein-based anti-viral signaling
pathways that perform similar functions to type I IFNs in vertebrates. For example,
mosquito cells secrete the peptide Vago upon viral infection, which signals through the
JAK-STAT pathway to activate anti-viral immunity (51). Additionally, the oyster
Crassostrea gigas is thought to have an IFN-like system, but no secreted proteins have
yet been identified in this organism (52). *N. vectensis* may also encode an undiscovered
IFN-like protein; at a minimum, *N. vectensis* encodes several IRF-like genes (Fig. S3).
One attractive hypothesis is that these IRFs are important for the anti-viral response of
N. vectensis; however, we were unable to see any impact of single knockdown
experiments on the induction of genes by 2'3'-cGAMP, though this may be explained by
redundancy or technical limitations of our knockdown approach. Nevertheless, an
important conclusion of our work is that induction of a broad transcriptional program is
an ancestral function of 2'3'-cGAMP signaling, similar to what has been seen in
Drosophila (31) and choanoflagellates (53). This ancestral transcriptional response
complements an additional autophagy response to 2'3'-cGAMP that was previously
reported to be induced by nvSTING in mammalian cells (17), and has now also been
shown to be induced by 2'3'-cGAMP and STING in choanoflagellates (53).
We found that in addition to an anti-viral response, *N. vectensis* responds to 2'3'-
cGAMP by inducing a variety of anti-bacterial genes, including lysozyme, Dae4,
perforin-2-like, LPB, and GBPs. With the exception of GBPs, which have dual anti-viral
and anti-bacterial activity, these anti-bacterial genes are not induced by 2'3'-cGAMP in
vertebrates; thus, the anti-bacterial response appears to be a unique feature of 2'3'-
cGAMP signaling in *N. vectensis*, and it will be interesting to see whether this proves
true in other invertebrates, or in additional cell types or contexts in vertebrates. Indeed,
a recent study found that during oral *L. monocytogenes* infection of mice, a STING-
dependent and IFN-independent response helps clear bacteria. Several of the anti-
bacterial genes are also induced by poly(I:C) in *Nematostella* (45), and we found that at
least one putative anti-viral gene was also induced during *P. aeruginosa* infection,
perhaps indicating a broader anti-pathogen response to PAMPs in *N. vectensis*.
Interestingly, we found that nvNF-κB was specifically required for the induction of many
of the anti-bacterial genes. This suggests that nvNF-κB activation downstream of 2'3'-
cGAMP signaling may have been present in the most recent common ancestor of
cnidarian and mammals, and confirms a role for nvNF-κB in *N. vectensis* immunity.
Consistent with this speculation, *Drosophila* STING also appears to activate NF-κB (30,
31, 34).
To further establish that 2'3'-cGAMP induces proteins with anti-bacterial activity, we
functionally characterized one 2'3'-cGAMP-induced, nvNF-κB-dependent protein,

nvDae4. We found that nvDae4 is a peptidoglycan-cleaving enzyme with direct bactericidal activity against Gram-positive bacteria. Many of the 2'3'-cGAMP-induced NF-kB dependent genes are not recognizable homologs of proteins of known function; thus, they represent good candidates for the discovery of novel anti-bacterial genes in *N. vectensis.*

Using shRNAs to knockdown nvSTING failed to confirm an essential role for nvSTING in the response to 2'3'-cGAMP. However, our previous biochemical and structural studies showed nvSTING binds 2'3'-cGAMP with high affinity ($K_d < 1nM$) and in a very similar manner as vertebrate STING (37). STING is essential for the response to 2'3'-cGAMP in diverse organisms, including vertebrates, choanoflagellates (53), and insects (31). In addition, nvSTING is highly induced by 2'3'-cGAMP. So despite our negative results, we favor the idea that nvSTING is at least partially responsible for the response of *N. vectensis* to 2'3'-cGAMP. It is possible that *N. vectensis* encodes a redundant 2'3'-cGAMP sensor, but such a sensor would have had to evolve specifically in Cnidarians, or be lost independently from choanoflagellates, insects and vertebrates. It is likely that technical limitations of performing shRNA knockdowns in *N. vectensis* accounts for our inability to observe a role for nvSTING in the response to 2'3'-cGAMP, though we cannot exclude the possibility that *N. vectensis* utilizes a distinct 2'3'-cGAMP-sensing pathway.

If indeed 2'3'-cGAMP is signaling via nvSTING, this presents several mechanistic questions. First, in mammals, all known transcriptional responses downstream of STING, including those requiring NF-κB activation, require the CTT (38, 39), leading to the question of how invertebrate STING proteins, which lack a discrete CTT, can activate this pathway. Also, nvNF-κB knockdown did not impact the vast majority of 2'3'-cGAMP-induced genes, which may imply the existence of other signaling pathways downstream of nvSTING. How these unidentified pathways become activated is another interesting question and one that could also shed light on mammalian STING signaling. Finally, mammalian STING can also be activated by direct binding to bacterial 3'3'-linked CDNs (54), and nvSTING also binds to these ligands, albeit with lower affinity (37). We found that treatment of *N. vectensis* with these ligands also led to induction of many of the same genes, likely through the same pathway. This perhaps indicates a role for the nvSTING pathway in bacterial sensing, though our preliminary attempts to observe an impact of 2'3'-cGAMP-induced gene expression on bacterial colonization of *N. vectensis* were unsuccessful. Further development of a bacterial infection model for *N. vectensis* will be required to study the anti-bacterial response of this organism *in vivo*.

A crucial remaining question is what activates nvcGAS to produce 2'3'-cGAMP. Double-stranded DNA did not seem to activate this protein *in vitro* (37), but this could be due to the absence of cofactors. This protein is also constitutively active when transfected into mammalian cells, but this could be due to overexpression. Unlike human cGAS, nvcGAS does not have any clear DNA-binding domains, although this does not necessarily exclude DNA as a possible ligand. The *Vibrio* cGAS-like enzyme DncV is regulated by folate-like molecules (55), so there is a diverse range of possible nvcGAS activators. Understanding the role of CDN sensing pathways in diverse organisms can shed light on the mechanisms of evolution of viral and bacterial sensing, and on unique ways divergent organisms have evolved to respond to pathogens.

Methods

Nematostella vectensis culture and spawning

N. vectensis adults were a gift from Mark Q. Martindale (University of Florida) and were cultured and spawned as previously described (56). Briefly, animals were kept in 1/3x seawater (12ppt salinity) in the dark at 17°C and fed freshly hatched *Artemia* (Carolina Biological Supply Company) weekly. Spawning was induced every two weeks by placing animals at 23°C under bright light for 8 hours, followed by 2 hours in the dark, and then finally moved to the light where they were monitored for spawning. Egg masses were de-jellied in 4% L-Cysteine (pH 7-7.4) in 1/3x sea water for 10-15 minutes and washed 3 times with 1/3x sea water. Water containing sperm was added to the washed eggs and these were either used immediately for microinjection or allowed to develop at room temperature.

CDN treatment

For the RNA sequencing experiment on polyps (Fig. 1 and Fig. S1), ~4 week old polyps were treated in duplicate in a bath of 500μM c-di-AMP, c-di-GMP, 2'3'-cGAMP, or 3'3'-cGAMP (all InvivoGen) in 1/3x sea water for 24 hours. For remaining cGAMP treatment experiments, 50-100 48-hour old embryos were treated with 100μM 2'3'-cGAMP (InvivoGen) in 1/3x sea water for 4 hours.

RNA sequencing

For the initial CDN treatment experiment using polyps, total RNA was extracted using Qiagen RNeasy Mini kits according to the manufacturer's protocol. Libraries were prepared by the Functional Genomics Laboratory at UC Berkeley using WaferGen PrepX library prep kits with oligo dT beads for mRNA enrichment according to the manufacturer's protocol, and 50 nt single-end sequencing was carried out on the HiSeq4000 (Illumina) by the Vincent J.Coates Genomics Sequencing Laboratory. For all other RNA sequencing experiments on 48 hour embryos, RNA was extracted using Trizol (Thermo Fisher Scientific) according to the manufacturer's protocol. Libraries were prepared by the Functional Genomics Laboratory at UC Berkeley as follows: oligo dT beads from the KAPA mRNA Capture Kit (KK8581) were used for mRNA enrichment; fragmentation, adapter ligation and cDNA synthesis were performed using the KAPA RNA HyperPrep kit (KK8540). Libraries were pooled evenly by molarity and sequenced by the Vincent J.Coates Genomics Sequencing Laboratory on a NovaSeq6000 150PE S4 flowcell (Illumina), generating 25M read pairs per sample. Read quality was assessed using FastQC. Reads were mapped to the *N. vectensis* transcriptome (NCBI: GCF_000209225.1) using kallisto and differential expression was analyzed in R with DESeq2. Differential expression was deemed significant with a log_2 fold change greater than 1 and an adjusted p-value less than 0.05. GO term analysis was performed using goseq with GO annotations from https://figshare.com/articles/dataset/Nematostella_vectensis_transcriptome_and_gene_models_v2_0/807696. The EnhancedVolcano package (https://github.com/kevinblighe/EnhancedVolcano) was used to generate volcano plots. Heatmaps are based on regularized log-transformed normalized counts and Z-scores are scaled by row. All RNA-Seq results can be found in Supplementary Dataset 1. Raw sequencing reads and normalized gene counts can be found at the NCBI GEO under accession GSE175984.

Quantitative Real-Time PCR (qRT-PCR)

Embryos and polyps were lysed in TRIzol (Invitrogen) and RNA was extracted according to the manufacturer's protocol. 500ng of RNA was treated with RQ1 RNase-

free DNase (Promega) for and reverse transcribed with Superscript III (Invitrogen). Quantitative PCR was performed using SYBR Green (Thermo Fisher Scientific) with 0.8 μM of forward and reverse primers on a QuantStudio 5 Real-Time PCR System (Applied Biosystems) with the following cycling conditions: 50°C 2 min; 95°C 10 min; [95°C 15 sec, 60°C 1 min] x 40; 95°C 15 sec; 60°C 1 min; melt curve: step 0.075 °C/s to 95°C. Fold changes in expression levels were normalized to actin and calculated using the $2^{-\Delta\Delta Ct}$ method. Student's t-tests were performed on ΔCt values. All primer sequences used in this study can be found in Supplementary Dataset 2.

shRNA microinjection

Short hairpin RNAs for microinjection were prepared by *in vitro* transcription as previously described (57). Briefly, unique 19 nucleotide targeting motifs were identified and used to create oligonucleotides with the following sequence: T7 promoter—19nt motif—linker—antisense 19nt motif–TT. RNA secondary structure was visualized using mfold (http://www.unafold.org/mfold/applications/rna-folding-form.php) to ensure a single RNA conformation. Both sense and anti-sense oligonucleotides were synthesized and mixed to a final concentration of 25μM, heated to 98°C for 5 minutes and cooled to 24°C before use as template for *in vitro* transcription using the Ampliscribe T7-Flash Transcription Kit (Lucigen). Reactions were allowed to proceed overnight, followed by a 15 minute treatment with DNase and subsequent purification with Direct-zol™ RNA MiniPrep Plus (Zymo Research). All shRNAs used in this study can be found in Supplementary Dataset 2.

Microinjections of one-cell embryos were carried out as previously described (58). shRNAs were diluted to a concentration of 500-900 ng/μl (ideal concentrations were determined experimentally) in RNase-free water with fluorescent dextran for visualization. Injected embryos were monitored for gross normal development at room temperature and used for experiments 48 hours later unless otherwise indicated. Knockdowns for each gene were performed using at least two different shRNAs and phenotypes were confirmed in at least 3 independent experiments.

Immunohistochemistry, imaging, and quantification

Polyps treated for 4 hours with 100μM 2'3'-cGAMP were stained for nvNF-κB as previously described (59). Briefly, polyps were fixed in 4% paraformaldehyde in 1/3x sea water overnight at 4°C with rocking, and subsequently washed 3 times with wash buffer (1× PBS, 0.2% Triton X-100). Antigen retrieval was performed by placing anemones in 95°C 5% urea for 5 minutes and allowing them to cool to room temperature before washing 3 times in wash buffer. Samples were blocked overnight at 4°C in blocking buffer (1× PBS, 5% normal goat serum, 1% bovine serum albumin, 0.2% Triton X-100). Samples were stained with anti-nvNF-κB (1:100; gift of Thomas Gilmore, Boston University) in blocking buffer for 90 minutes at room temperature and washed 4 times in wash buffer. Samples were then incubated in FITC-anti-Rabbit IgG (1:160; F9887, Sigma-Aldrich) in blocking buffer for 90 minutes at 37°C. Finally, samples were washed in wash buffer, stained with 1 μg/mL of DAPI for 10 minutes, washed again, and mounted in Vectashield HardSet Mounting medium and imaged on a Zeiss LSM 710 AxioObserver. Imaris 9.2 (Bitplane) was used to create 3D surfaces based on DAPI expression, and surface statistics were exported and analyzed in FlowJo (BD) to quantify nuclear nvNF-κB expression as previously described (60).

Bacterial infection

Single colonies of *Pseudomonas aeruginosa* strain PA14-GFP were cultured
overnight in LB with 50μg/ml of carbenicillin, centrifuged for 5 minutes at 3000 x *g,*
resuspended to an OD_{600} of 0.1, 0.01, or 0.001 in 1/3x sea water and used to infect
polyps, which were kept at room temperature or 28°C as indicated. Inputs were plated
to calculate CFU/ml. Polyp survival was monitored daily. For expression analysis,
polyps were homogenized in Trizol and RNA extraction and qPCR were performed as
indicated above.
515
Protein purification
nvDae4 lacking its signal peptide was cloned from cDNA. These fragments were
then cloned into the pAcGP67-A baculovirus transfer vector for secreted, His-tagged
protein expression. The plasmid for expressing mutant nvDae4 (C63A) was made from
the pAcGP67-A-nvDae4 plasmid using Q5 site-directed mutagenesis (NEB) according
to the manufactures protocol. Plasmids were transfected into Sf9 insect cells ($2x10^6$
cells/ml in 2ml) using Cellfectin II Reagent (Gibco) along with BestBac 2.0 v-cath/chiA
Deleted Linearized Baculovirus DNA (Expression Systems) for 6 hours, after which
media was replaced and cells were left for 1 week at 25°C. Supernatants were
harvested and 50 μL were used to infect $7x10^6$ Sf9 cells in 10ml of media for 1 week at
25°C. Supernatants containing secondary virus were harvested, tested, and used to
infect High Five cells (2 L at $1.5x10^6$ cells/ml) for 72 hours at 25°C with shaking.
Supernatants containing protein were harvested by centrifuging for 15 min at 600 x *g* at
4°C and subsequently passing through a 0.45 μm filter to remove all cells. Supernatants
were buffered to 1x HBS (20mM HEPES pH 7.2, 150mM NaCl), mixed with 2 mL of Ni-
NTA agarose were per liter, and rotated at 4°C for 2 hours. Ni-NTA resins with bound
protein were collected on a column by gravity-flow and washed with 30x column volume
of wash buffer (20mM HEPES, 1M NaCl, 30mM imidazole, 10% glycerol). Protein was
eluted in 1 mL fractions using 1xHBS supplemented with 200mM imidazole. Buffer was
exchanged to 1xHBS+ 2mM DTT using Econo-Pac10DG Desalting Prepacked Gravity
Flow Columns (Bio-rad) according to the manufacturers protocol, and proteins were
concentrated using 10kDa concentrators (Millipore).
538
Bacterial killing assays
For expression in *E. coli*, nvDae4 WT and C63A lacking the endogenous signal
sequence were cloned into the pET28a vector for inducible cytosolic expression, or the
pET22b vector for inducible periplasmic expression. *E. coli* (BL21 DE3 strain) were
freshly transformed with the vectors and grown overnight in LB with 50 μg/mL
carbenicillin shaking at 37°C. Overnight cultures were backdiluted in LB to the same
OD_{600} and grown to log phase before induction with 0.25mM IPTG. Plates were kept
shaking at 37°C and OD_{600} was read every 5 minutes for 3 hours.
Bacillus subtilis-GFP (derivative of strain 168; BGSC accession #1A1139) was
grown to log-phase in LB with 100 μg/mL spectinomycin, centrifuged, resuspended in
0.5xHBS, and incubated alone or with nvDae4 WT or C63A at indicated concentrations
for 2-3 hours at 37°C. Serial dilutions were plated on LB agar with 100 μg/mL
spectinomycin to determine CFU.
552
Peptidoglycan cleavage assays
Peptidoglycan (PG) was purified and analyzed as previously described (50).
Briefly, *Escherichia coli* BW11325 (from Carol Gross, UCSF) and *Staphylococcus*
epidermis BCM060 (from Tiffany Scharschmidt, UCSF) were grown to an OD600 of 0.6,
harvested by centrifugation, and boiled in SDS (4 % final concentration) for 4 hours with

stirring. After washing in purified water to remove SDS, the peptidoglycan was treated
with Pronase E for 2 h at 60°C (0.1 mg/ml final concentration in 10 mM Tris-HCl pH7.2
and 0.06 % NaCl; pre-activated for 2 h at 60 °C). Pronase E was heat inactivated at
100°C for 10 min and washed with sterile filtered water (5 x 20 min at 21k × *g*). PG from
Gram-positive bacteria (*Se*) was also treated with 48% HF at 37°C for 48 h to remove
teichoic acids, followed by washes with sterile filtered water. nvDae4 enzyme (WT and
C63A) was added (1-10 µM in 10 mM Tris-HCl pH 7.2 and 0.06% NaCl) and incubated
O/N at 37°C. Enzymes were heat inactivated at 100°C for 10 min. Mutanolysin (Sigma
M9901, final concentration 20 µg/ml) was added to the purified peptidoglycan and
incubated overnight at 37 °C. The peptidoglycan fragments were reduced, acidified,
analyzed via HPLC (0.5 ml/min flow rate, 55°C with Hypersil ODS C18 HPLC column,
Thermo Scientific, catalog number: 30103-254630).
570
571 ***Supplemental methods***
572
Phylogenetic analysis
Protein sequences containing domains of interest were downloaded from NCBI,
with the exception of nvGBP6 and nvGBP7, for which RNA-seq data showed additional
nucleotide usage relative to the reference sequence (all sequences can be found in
Supplementary Dataset 2). These were aligned using on phylogeny.fr using MUSCLE
and manipulated in Geneious. For the GBP and IRF alignments, only the domain of
interest was used for phylogenetic analysis. Maximum likelihood phylogenetic trees
were generated with PhyML using 100 bootstrap replicates. Alignments shown were
made in Geneious using Clustal Omega.
582
Nanostring gene expression analysis
A custom codeset targeting 36 genes of interest and 4 housekeeping genes for
normalization (Supplementary Dataset 2) was designed by NanoString Technologies
(Seattle, WA) for use in nCounter XT CodeSet Gene Expression Assays run on an
nCounter SPRINT Profiler (NanoString Technologies). RNA was isolated as it was for
qRT-PCR experiments, and hybridized to probes according to the manufacturer's
protocol using 50ng/µL of total RNA. Quality control, data normalization, and
visualization was performed in nSolver 4.0 analysis software (NanoString Technologies)
according to the manufacturer's protocol.
592
Mammalian cell immunofluorescence and confocal microscopy
Glass coverslips were seeded with 293T cells and grown to ~50% confluency.
Cells were transfected for 24-48 hours with a total of 1.25 µg of DNA and 3 µL
Lipofectamine 2000. Each well contained the following: pcDNA4-STING (10 ng),
pEGFP-LC3 (5 ng) and either empty vector or pcDNA4 with the indicated cyclic-
dinucleotide synthase (1,235 ng). Cells were washed once in PBS, fixed for 15 minutes
in 4% paraformaldehyde, washed once in PBS, and permeabilized for 5 minutes in
0.5% saponin in PBS. Cells were then washed once in PBS, and treated with 0.1%
sodium borohydride/0.1% saponin/PBS for 5 – 10 minutes in order to consume any
remaining paraformaldehyde. Cells were then washed 3 times in PBS, and blocked with
blocking buffer (1% BSA/0.1% saponin/PBS) for 45 minutes. Cells were then incubated
HA antibody (1:200 dilution, Sigma 11867423001 rat IgG from Roche) in blocking buffer
for one hour and washed 3 times in PBS. Each well was then incubated with 0.1%
saponin/PBS and secondary antibody (1:500 dilution, Jackson ImmunoResearch, Cy3
affinipure donkey anti-rat IgG, 712-165-153) for 45 minutes. Finally, cells were washed

3 times in PBS, mounted using VectaShield with DAPI, and dried overnight. Images
were acquired using a Zeiss LSM 780 NLO AxioExaminer.
610
Acknowledgements

We are especially grateful to Mark Q. Martindale and Miguel Salinas-Saavedra
for *N. vectensis* animals and training. We also thank Matt Gibson for sharing shRNA
protocols, and Thomas Gilmore for the anti-nvNF-κB antibody. We thank members of
the Vance and Barton labs for discussions, and Arielle Woznica for comments on the
manuscript. This work used the Functional Genomic Laboratory and Vincent J. Coates
Genomics Sequencing Laboratory at UC Berkeley, supported by NIH S10 OD018174
Instrumentation Grant. Confocal imaging experiments were conducted at the CRL
Molecular Imaging Center supported by the Gordon and Betty Moore Foundation; we
would like to thank Holly Aaron and Feather Ives for training and assistance. REV is an
HHMI Investigator and is supported by NIH grants AI0663302, AI075039, and
AI155634. SRM is supported by the National Science Foundation Graduate Research
Fellowship under Grant Numbers DGE 1106400 and DGE 1752814. B.H. and S.C. were
supported by funding from the NIH (R01AI132851 to S.C.), the Chan Zuckerberg
Biohub, and the Sangvhi-Agarwal Innovation Award. Any opinions, findings, and
conclusions or recommendations expressed in this material are those of the author(s)
and do not necessarily reflect the views of the National Science Foundation, HHMI, or
the National Institutes of Health.

References

1. J. Ahn, G. N. Barber, STING signaling and host defense against microbial infection. *Exp Mol Med* **51**, 1-10 (2019).
2. A. Ablasser, Z. J. Chen, cGAS in action: Expanding roles in immunity and inflammation. *Science* **363** (2019).
3. L. Sun, J. Wu, F. Du, X. Chen, Z. J. Chen, Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science* **339**, 786-791 (2013).
4. P. Gao *et al.*, Cyclic [G(2',5')pA(3',5')p] is the metazoan second messenger produced by DNA-activated cyclic GMP-AMP synthase. *Cell* **153**, 1094-1107 (2013).
5. A. Ablasser *et al.*, cGAS produces a 2'-5'-linked cyclic dinucleotide second messenger that activates STING. *Nature* **498**, 380-384 (2013).
6. E. J. Diner *et al.*, The innate immune DNA sensor cGAS produces a noncanonical cyclic dinucleotide that activates human STING. *Cell reports* **3**, 1355-1361 (2013).
7. X. Zhang *et al.*, Cyclic GMP-AMP containing mixed phosphodiester linkages is an endogenous high-affinity ligand for STING. *Molecular cell* **51**, 226-235 (2013).
8. H. Ishikawa, G. N. Barber, STING is an endoplasmic reticulum adaptor that facilitates innate immune signalling. *Nature* **455**, 674-678 (2008).
9. Y. Tanaka, Z. J. Chen, STING specifies IRF3 phosphorylation by TBK1 in the cytosolic DNA signaling pathway. *Science signaling* **5**, ra20 (2012).
10. S. Liu *et al.*, Phosphorylation of innate immune adaptor proteins MAVS, STING, and TRIF induces IRF3 activation. *Science* **347**, aaa2630 (2015).
11. C. Zhang *et al.*, Structural basis of STING binding with and phosphorylation by TBK1. *Nature* **567**, 394-398 (2019).
12. B. Zhao *et al.*, A conserved PLPLRT/SD motif of STING mediates the recruitment and activation of TBK1. *Nature* **569**, 718-722 (2019).
13. D. B. Stetson, R. Medzhitov, Type I interferons in host defense. *Immunity* **25**, 373-381 (2006).
14. J. W. Schoggins, Interferon-Stimulated Genes: What Do They All Do? *Annu Rev Virol* **6**, 567-584 (2019).
15. T. Abe, G. N. Barber, Cytosolic-DNA-mediated, STING-dependent proinflammatory gene induction necessitates canonical NF-kappaB activation through TBK1. *Journal of virology* **88**, 5328-5341 (2014).
16. H. Chen *et al.*, Activation of STAT6 by STING is critical for antiviral innate immunity. *Cell* **147**, 436-446 (2011).
17. X. Gui *et al.*, Autophagy induction via STING trafficking is a primordial function of the cGAS pathway. *Nature* **567**, 262-266 (2019).
18. T. D. Fischer, C. Wang, B. S. Padman, M. Lazarou, R. J. Youle, STING induces LC3B lipidation onto single-membrane vesicles via the V-ATPase and ATG16L1-WD40 domain. *J Cell Biol* **219** (2020).
19. R. O. Watson, P. S. Manzanillo, J. S. Cox, Extracellular M. tuberculosis DNA targets bacteria for autophagy by activating the host DNA-sensing pathway. *Cell* **150**, 803-815 (2012).

20.   R. O. Watson *et al.*, The Cytosolic Sensor cGAS Detects Mycobacterium
tuberculosis DNA to Induce Type I Interferons and Activate Autophagy. *Cell host*
& microbe **17**, 811-819 (2015).
21.   S. Gluck, A. Ablasser, Innate immunosensing of DNA in cellular senescence.
Current opinion in immunology **56**, 31-36 (2019).
22.   M. M. Gaidt *et al.*, The DNA Inflammasome in Human Myeloid Cells Is Initiated
by a STING-Cell Death Program Upstream of NLRP3. *Cell* **171**, 1110-1124
e1118 (2017).
23.   A. Sze *et al.*, Host restriction factor SAMHD1 limits human T cell leukemia virus
type 1 infection of monocytes via STING-mediated apoptosis. *Cell host &*
microbe **14**, 422-434 (2013).
24.   S. R. Paludan, L. S. Reinert, V. Hornung, DNA-stimulated cell death: implications
for host defence, inflammatory diseases and cancer. *Nature reviews.*
Immunology **19**, 141-153 (2019).
25.   M. F. Gulen *et al.*, Signalling strength determines proapoptotic functions of
STING. *Nature communications* **8**, 427 (2017).
26.   J. Wu *et al.*, STING-mediated disruption of calcium homeostasis chronically
activates ER stress and primes T cell death. *The Journal of experimental*
medicine **216**, 867-883 (2019).
27.   S. R. Margolis, S. C. Wilson, R. E. Vance, Evolutionary Origins of cGAS-STING
Signaling. *Trends Immunol* **38**, 733-743 (2017).
28.   D. Cohen *et al.*, Cyclic GMP-AMP signalling protects bacteria against viral
infection. *Nature* **574**, 691-695 (2019).
29.   B. R. Morehouse *et al.*, STING cyclic dinucleotide sensing originated in bacteria.
Nature **586**, 429-433 (2020).
30.   A. Goto *et al.*, The Kinase IKKbeta Regulates a STING- and NF-kappaB-
Dependent Antiviral Response Pathway in Drosophila. *Immunity* **49**, 225-234
e224 (2018).
31.   H. Cai *et al.*, 2'3'-cGAMP triggers a STING- and NF-kappaB-dependent broad
antiviral response in Drosophila. *Science signaling* **13** (2020).
32.   Y. Liu *et al.*, Inflammation-Induced, STING-Dependent Autophagy Restricts Zika
Virus Infection in the Drosophila Brain. *Cell host & microbe* **24**, 57-68 e53 (2018).
33.   X. Hua *et al.*, Stimulator of interferon genes (STING) provides insect antiviral
immunity by promoting Dredd caspase-mediated NF-kappaB activation. *J Biol*
Chem **293**, 11878-11890 (2018).
34.   M. Martin, A. Hiroyasu, R. M. Guzman, S. A. Roberts, A. G. Goodman, Analysis
of Drosophila STING Reveals an Evolutionarily Conserved Antimicrobial
Function. *Cell reports* **23**, 3537-3550 e3536 (2018).
35.   X. Hua, W. Xu, S. Ma, Q. Xia, STING-dependent autophagy suppresses Nosema
bombycis infection in silkworms, Bombyx mori. *Developmental and comparative*
immunology **115**, 103862 (2021).
36.   N. H. Putnam *et al.*, Sea anemone genome reveals ancestral eumetazoan gene
repertoire and genomic organization. *Science* **317**, 86-94 (2007).
37.   P. J. Kranzusch *et al.*, Ancient Origin of cGAS-STING Reveals Mechanism of
Universal 2',3' cGAMP Signaling. *Molecular cell* **59**, 891-903 (2015).
38.   L. H. Yamashiro *et al.*, Interferon-independent STING signaling promotes
resistance to HSV-1 in vivo. *Nature communications* **11**, 3382 (2020).

39. S. Yum, M. Li, Y. Fang, Z. J. Chen, TBK1 recruitment to STING activates both
IRF3 and NF-kappaB that mediate immune defense against tumors and viral
infections. *Proceedings of the National Academy of Sciences of the United*
States of America **118** (2021).
40. D. J. Miller *et al.*, The innate immune repertoire in cnidaria--ancestral complexity
and stochastic gene loss. *Genome Biol* **8**, R59 (2007).
41. A. M. Reitzel, J. C. Sullivan, N. Traylor-Knowles, J. R. Finnerty, Genomic survey
of candidate stress-response genes in the estuarine anemone Nematostella
vectensis. *Biol Bull* **214**, 233-254 (2008).
42. J. J. Brennan *et al.*, Sea anemone model has a single Toll-like receptor that can
function in pathogen detection, NF-kappaB signal transduction, and
development. *Proceedings of the National Academy of Sciences of the United*
States of America **114**, E10122-E10131 (2017).
43. F. S. Wolenski *et al.*, Characterization of the core elements of the NF-kappaB
signaling pathway of the sea anemone Nematostella vectensis. *Molecular and*
cellular biology **31**, 1076-1087 (2011).
44. F. S. Wolenski, C. A. Bradham, J. R. Finnerty, T. D. Gilmore, NF-kappaB is
required for cnidocyte development in the sea anemone Nematostella vectensis.
Developmental biology **373**, 205-215 (2013).
45. M. Lewandowska, T. Sharoni, Y. Admoni, R. Aharoni, Y. Moran, Functional
characterization of the cnidarian antiviral immune response reveals ancestral
complexity. *Molecular biology and evolution* 10.1093/molbev/msab197 (2021).
46. S. Chou *et al.*, Transferred interbacterial antagonism genes augment eukaryotic
innate immune function. *Nature* **518**, 98-101 (2015).
47. L. G. Rahme *et al.*, Plants and animals share functionally common bacterial
virulence factors. *Proceedings of the National Academy of Sciences of the United*
States of America **97**, 8815-8821 (2000).
48. S. Franzenburg *et al.*, MyD88-deficient Hydra reveal an ancient function of TLR
signaling in sensing bacterial colonizers. *Proceedings of the National Academy of*
Sciences of the United States of America **109**, 19374-19379 (2012).
49. A. B. Russell *et al.*, Type VI secretion delivers bacteriolytic effectors to target
cells. *Nature* **475**, 343-347 (2011).
50. B. M. Hayes *et al.*, Ticks Resist Skin Commensals with Immune Factor of
Bacterial Origin. *Cell* **183**, 1562-1571 e1512 (2020).
51. P. N. Paradkar, L. Trinidad, R. Voysey, J. B. Duchemin, P. J. Walker, Secreted
Vago restricts West Nile virus infection in Culex mosquito cells by activating the
Jak-STAT pathway. *Proceedings of the National Academy of Sciences of the*
United States of America **109**, 18915-18920 (2012).
52. T. J. Green, P. Speck, Antiviral Defense and Innate Immune Memory in the
Oyster. *Viruses* **10** (2018).
53. A. Woznica *et al.*, STING mediates immune responses in a unicellular
choanoflagellate. *bioRxiv* https://doi.org/10.1101/2021.05.13.443778 (2021).
54. D. L. Burdette *et al.*, STING is a direct innate immune sensor of cyclic di-GMP.
Nature **478**, 515-518 (2011).
55. D. Zhu *et al.*, Structural biochemistry of a Vibrio cholerae dinucleotide cyclase
reveals cyclase activity regulation by folates. *Molecular cell* **55**, 931-937 (2014).

56.   D. J. Stefanik, L. E. Friedman, J. R. Finnerty, Collecting, rearing, spawning and
769   inducing regeneration of the starlet sea anemone, Nematostella vectensis. *Nat*
770   *Protoc* **8**, 916-923 (2013).
771   57.   A. Karabulut, S. He, C. Y. Chen, S. A. McKinney, M. C. Gibson, Electroporation
772   of short hairpin RNAs for rapid and efficient gene knockdown in the starlet sea
773   anemone, Nematostella vectensis. *Developmental biology* **448**, 7-15 (2019).
774   58.   M. J. Layden, E. Rottinger, F. S. Wolenski, T. D. Gilmore, M. Q. Martindale,
775   Microinjection of mRNA or morpholinos for reverse genetic analysis in the starlet
776   sea anemone, Nematostella vectensis. *Nat Protoc* **8**, 924-934 (2013).
777   59.   F. S. Wolenski, M. J. Layden, M. Q. Martindale, T. D. Gilmore, J. R. Finnerty,
778   Characterizing the spatiotemporal expression of RNAs and proteins in the starlet
779   sea anemone, Nematostella vectensis. *Nat Protoc* **8**, 900-915 (2013).
780   60.   D. I. Kotov, T. Pengo, J. S. Mitchell, M. J. Gastinger, M. K. Jenkins, Chrysalis: A
781   New Method for High-Throughput Histo-Cytometry Analysis of Images and
782   Movies. *Journal of immunology* **202**, 300-308 (2019).
783

Figure Legends

Figure 1: 2'3'-cGAMP induces many putative immune genes in *Nematostella vectensis*

    A) Volcano plot showing differential gene expression (DE) in *N. vectensis* polyps untreated vs. treated with 2'3'-cGAMP for 24 hours. A positive fold-change indicates higher expression in polyps treated with 2'3'-cGAMP. Genes of interest with homologs known to be involved in immunity in other organisms are labeled.

    B) Breakdown of DE genes into categories based on known GO terms. Gene set enrichment analysis shows a clear enrichment of GO terms associated with immunity.

    C) qRT-PCR measuring genes of interest in 48-hour-old *N. vectensis* embryos untreated or treated with 2'3'-cGAMP for 4 hours. Fold changes were calculated relative to untreated as $2^{-\Delta\Delta Ct}$ and each point represents one biological replicate. Unpaired t test performed on ΔΔCt before log transformation. *p ≤ 0.05; **p ≤ 0.01; ***p ≤ 0.001; ****p ≤ 0.0001.

Figure 2: The induction of many anti-bacterial genes by 2'3'-cGAMP is nvNF-κB dependent

    A) Heatmap showing all genes that are significantly (padj < 0.05, $\log_2 FC < -1$) downregulated in 2'3'-cGAMP -treated embryos microinjected with NF-κB shRNA vs. GFP shRNA. Genes with predicted anti-bacterial function are labeled

    B) qRT-PCR of anti-bacterial genes in nvNF-kB shRNA or control GFP shRNA treated samples after induction by 2'3'-cGAMP. Fold change was calculated relative to untreated, GFP shRNA injected as $2^{-\Delta\Delta Ct}$ and each point represents one biological replicate. Unpaired t test performed on ΔΔCt before log transformation. *p ≤ 0.05; **p ≤ 0.01.

    C) Whole mount immunofluorescence of polyps stained with anti-nvNF-κB antiserum. Right two panels are enlargements of the boxed regions indicated in the left two panels.

    D) Quantification of cells with nuclear localization of nvNF-kB after treatment with cGAMP (representative images shown in C). Each point represents a single polyp, in which at least 1500 cells were analyzed. Statistical analysis was performed by unpaired t test; *p = 0.0481.

Figure 3: *Pseudomonas aeruginosa* infection induces putative anti-bacterial genes

    A) Survival curves of *N. vectensis* polyps infected with *P. aeruginosa* at indicated dose and temperature.

B+C) qRT-PCR of nvSTING (B) or putative anti-bacterial genes (C) assayed at 48 hours post *Pa* infection (2×10^7 CFU/ml). Each point represents one biological replicate; unpaired t test performed on ΔΔCt before log transformation. **p ≤ 0.01; ***p ≤ 0.001; ****p ≤ 0.0001.

Figure 4: A 2'3'-cGAMP induced, nvNF-κB-dependent protein has anti-bacterial activity

A) Growth of *E. coli* expressing either periplasmic (Peri-) or cytosolic (Cyto-) nvDae4 (WT or C63A) induced with 250µM IPTG. Error bars +/- SD; n=3. Unpaired t test; **p = 0.0063.

B) Partial HPLC chromatograms of *E. coli* peptidoglycan sacculi after overnight incubation with buffer only (no enzyme), or 1 µM nvDae4 WT or C63A enzyme.

C) *Bacillus subtilis* CFU after 2 hour incubation with buffer alone, nvDae4 WT or catalytic mutant C63A (25 µM). Error bars +/- SEM; n=3. Unpaired t test performed on log-tranformed values; **p ≤ 0.01.

D) Dose dependent killing of *B. subtilis* by WT nvDae4 enzyme (same assay as in C). Error bars +/- SD; n=2 per concentration.

Figure S1: Treatment with other CDNs leads to some gene induction

A) Heatmap showing differentially expressed genes in response to c-di-AMP, c-di-GMP, and 3'3'-cGAMP. Almost all of these are also significantly induced by 2'3'-cGAMP.

B-D) Volcano plots of differential gene expression in *N. vectensis* polyps untreated vs. treated with cyclic-di-AMP (B), cyclic-di-GMP (C) and 3'3'-cGAMP (D) for 24 hours.

Figure S2: nvSTING knockdown does not impact the induction of genes by 2'3'-cGAMP

A) Volcano plot showing differential gene expression in 48 hour embryos treated with 2'3'-cGAMP that were injected with GFP shRNA or nvSTING shRNA. Positive fold-change indicates higher expression in GFP shRNA injected embryos.

B) Clustered heatmap showing the expression of the top 1000 varied genes by RNA-Seq between embryos injected with either GFP or nvSTING shRNA and either untreated or treated with 2'3'-cGAMP.

C) Fold change of nvSTING, nvOAS, and nvLBP assayed by Nanostring from experiments using 3 different shRNAs to knock down nvSTING expression.

D) qRT-qPCR measuring genes of interest in 48-hour-old embryos injected with a control (ctrl) or nvSTING translation-inhibiting morpholino (MO) and treated with 2'3'-cGAMP. Fold changes were calculated as $2^{-\Delta\Delta Ct}$ and each point represents one biological replicate. Unpaired t test performed on ΔΔCt before log transformation; no significant differences.

E) Immunofluorescence images of 293T cells transfected with plasmids encoding nvSTING-HA, LC3-GFP, and either empty vector, human cGAS or *V. cholera* DncV. Human cGAS is activated by the transfected DNA to produce 2'3'-cGAMP, and DncV, which produces 3'3'-cGAMP, is constitutively active in 293T cells.

Figure S3: Phylogenetic study of *N. vectensis* IRFs

A) Phylogenetic tree of all human and *N. vectensis* IRF proteins. 3 nvIRFs cluster with members of the human IRF1 supergroup, while the other 2 nvIRFs cluster with the IRF4 supergroup.

B) Full protein alignment of sequences in A). The DNA-binding domain is highly conserved between all *N. vectensis* and human IRFs. Among *N. vectensis* paralogs, only nvIRF5 contains an IAD1 domain.

877   C) Alignment of all nvIRF DNA-binding domains with conserved tryptophan pentad
878   outlined in red
879
Figure S4: Knockdowns of nvIRFs or nvSTAT have no effect on 2'3'-cGAMP-
induced gene expression
A) Fold changes in gene expression as determined by Nanostring in embryos
microinjected with shRNAs targeting EGFP, nvIRF1, or nvIRF5 either untreated
or treated with 2'3'-cGAMP.
B) Fold changes in gene expression as determined by qRT-PCR in samples
microinjected with shRNAs targeting EGFP, nvIRF2, nvIRF-67, or nvIRF4 either
untreated or treated with 2'3'-cGAMP. Note that IRF2 is not induced by cGAMP
and was mostly undetected in all samples; therefore it is possible that the
knockdowns were unsuccessful.
C) Volcano plot showing differential gene expression as determined by RNA-Seq in
48 hour embryos treated with 2'3'-cGAMP that were injected with GFP shRNA or
nvSTAT shRNA. Positive fold-change indicates higher expression in GFP shRNA
injected embryos. The GFP shRNA samples here are the same as those shown
in Figure 2A.
D) Fold changes in gene expression as determined by Nanostring in embryos
microinjected with shRNAs targeting EGFP or nvSTAT either untreated or treated
with 2'3'-cGAMP.
898
Figure S5: Anti-viral gene induction is not dependent on nvNF-κB
A) Fold changes in gene expression as determined by Nanostring in embryos
microinjected with shRNAs targeting EGFP or nvNF-κB, either untreated or
treated with 2'3'-cGAMP. The GFP shRNA samples here are the same as those
shown in Figure S4. GBP-806 expression included to show anti-bacterial gene
induction is lower in these samples.
B) Fold changes in nvLysozyme expression as determined by qRT-PCR in embryos
microinjected with shRNAs targeting EGFP or nvNF-κB, either untreated or
treated with 2'3'-cGAMP.
A+B) No significant differences in gene expression are observed between any
cGAMP treated samples.
C) qRT-PCR of putative anti-viral genes assayed at 48 hours post *Pa* infection
($2x10^7$ CFU/ml). Each point represents one biological replicate; unpaired t test
performed on ΔΔCt before log transformation. **p $\leq$ 0.01.
913
Figure S6: Phylogenetic study of *N. vectensis* GBPs
A) Phylogenetic tree of mammalian GTPases and putative *N. vectensis* GBPs made
with the full protein sequences. Branches with mammalian interferon-induced
GTPases and cGAMP-induced *N. vectensis* GBPs are colored red; these tend to
cluster together. Domain structures of *N. vectensis* GBPs are displayed.
B) Alignment of the GTPase domains of all *N. vectensis* GBPs and select
mammalian GTPase. Conserved GBP and atlastin specific residues are
highlighted in red.
922

923 **Figure S7: nvDae4 cleaves peptidoglycan from Gram-positive bacteria.** Partial
924 HPLC chromatograms of *Staphylococcus epidermis* peptidoglycan sacculi products
925 resulting from incubation with buffer only (no enzyme), or 1 µM nvDae4 WT or C63A
926 enzyme.

Figure 1: 2'3'-cGAMP induces many putative immune genes in *Nematostella vectensis*

A) Volcano plot showing differential gene expression (DE) in *N. vectensis* polyps untreated vs. treated with 2'3'-cGAMP for 24 hours. A positive fold-change indicates higher expression in polyps treated with 2'3'-cGAMP. Genes of interest with homologs known to be involved in immunity in other organisms are labeled.

B) Breakdown of DE genes into categories based on known GO terms. Gene set enrichment analysis shows a clear enrichment of GO terms associated with immunity.

C) qRT-PCR measuring genes of interest in 48-hour-old *N. vectensis* embryos untreated or treated with 2'3'-cGAMP for 4 hours. Fold changes were calculated relative to untreated as $2^{-\Delta\Delta Ct}$ and each point represents one biological replicate. Unpaired t test performed on ΔΔCt before log transformation. *p ≤ 0.05; **p ≤ 0.01; ***p ≤ 0.001; ****p ≤ 0.0001.

Figure 2: The induction of many anti-bacterial genes by 2'3'-cGAMP is nvNF-κB dependent

A) Heatmap showing all genes that are significantly (padj < 0.05, log$_2$FC<-1) downregulated in 2'3'-cGAMP -treated embryos microinjected with NF-κB shRNA vs. GFP shRNA. Genes with predicted anti-bacterial function are labeled

B) qRT-PCR of anti-bacterial genes in nvNF-kB shRNA or control GFP shRNA treated samples after induction by 2'3'-cGAMP. Fold change was calculated relative to untreated, GFP shRNA injected as 2$^{-\Delta\Delta Ct}$ and each point represents one biological replicate. Unpaired t test performed on ΔΔCt before log transformation. *p ≤ 0.05; **p ≤ 0.01.

C) Whole mount immunofluorescence of polyps stained with anti-nvNF-κB antiserum. Right two panels are enlargements of the boxed regions indicated in the left two panels.

D) Quantification of cells with nuclear localization of nvNF-kB after treatment with cGAMP (representative images shown in C). Each point represents a single polyp, in which at least 1500 cells were analyzed. Statistical analysis was performed by unpaired t test; *p = 0.0481.

Figure 3: *Pseudomonas aeruginosa* infection induces putative anti-bacterial genes

 A) Survival curves of *N. vectensis* polyps infected with *P. aeruginosa* at indicated dose and temperature.

B+C) qRT-PCR of nvSTING (B) or putative anti-bacterial genes (C) assayed at 48 hours post *Pa* infection (2x10^7 CFU/ml). Each point represents one biological replicate; unpaired t test performed on ΔΔCt before log transformation. **p ≤ 0.01; ***p ≤ 0.001; ****p ≤ 0.0001.

Figure 4: A 2'3'-cGAMP induced, nvNF-κB-dependent protein has anti-bacterial activity

A) Growth of *E. coli* expressing either periplasmic (Peri-) or cytosolic (Cyto-) nvDae4 (WT or C63A) induced with 250μM IPTG. Error bars +/- SD; n=3. Unpaired t test; **p = 0.0063.

B) Partial HPLC chromatograms of *E. coli* peptidoglycan sacculi after overnight incubation with buffer only (no enzyme), or 1 μM nvDae4 WT or C63A enzyme.

C) *Bacillus subtilis* CFU after 2 hour incubation with buffer alone, nvDae4 WT or catalytic mutant C63 (25 μM). Error bars +/- SEM; n=3. Unpaired t test performed on log-tranformed values; **p ≤ 0.01.

D) Dose dependent killing of *B. subtilis* by WT nvDae4 enzyme (same assay as in C). Error bars +/- SD; n=2 per concentration.

Figure S1: Treatment with other CDNs leads to some gene induction

A) Heatmap showing differentially expressed genes in response to c-di-AMP, c-di-GMP, and 3'3'-cGAMP. Almost all of these are also significantly induced by 2'3'-cGAMP.

B-D) Volcano plots of differential gene expression in *N. vectensis* polyps untreated vs. treated with cyclic-di-AMP (B), cyclic-di-GMP (C) and 3'3'-cGAMP (D) for 24 hours.

Figure S2: nvSTING knockdown does not impact the induction of genes by 2'3'-cGAMP

A) Volcano plot showing differential gene expression in 48 hour embryos treated with 2'3'-cGAMP that were injected with GFP shRNA or nvSTING shRNA. Positive fold-change indicates higher expression in GFP shRNA injected embryos.

B) Clustered heatmap showing the expression of the top 1000 varied genes by RNA-Seq between embryos injected with either GFP or nvSTING shRNA and either untreated or treated with 2'3'-cGAMP.

C) Fold change of nvSTING, nvOAS, and nvLBP assayed by Nanostring from experiments using 3 different shRNAs to knock down nvSTING expression.

D) qRT-qPCR measuring genes of interest in 48-hour-old embryos injected with a control (ctrl) or nvSTING translation-inhibiting morpholino (MO) and treated with 2'3'-cGAMP. Fold changes were calculated as $2^{-\Delta\Delta Ct}$ and each point represents one biological replicate. Unpaired t test performed on ΔΔCt before log transformation; no significant differences.

E) Immunofluorescence images of 293T cells transfected with plasmids encoding nvSTING-HA, LC3-GFP, and either empty vector, human cGAS or *V. cholera* DncV. Human cGAS is activated by the transfected DNA to produce 2'3'-cGAMP, and DncV, which produces 3'3'-cGAMP, is constitutively active in 293T cells.

Figure S3: Phylogenetic study of *N. vectensis* IRFs

A) Phylogenetic tree of all human and *N. vectensis* IRF proteins. 3 nvIRFs cluster with members of the human IRF1 supergroup, while the other 2 nvIRFs cluster with the IRF4 supergroup.

B) Full protein alignment of sequences in A). The DNA-binding domain is highly conserved between all *N. vectensis* and human IRFs. Among *N. vectensis* paralogs, only nvIRF5 contains an IAD1 domain.

C) Alignment of all nvIRF DNA-binding domains with conserved tryptophan pentad outlined in red

Figure S4: Knockdowns of nvIRFs or nvSTAT have no effect on 2'3'-cGAMP-induced gene expression

A) Fold changes in gene expression as determined by Nanostring in embryos microinjected with shRNAs targeting EGFP, nvIRF1, or nvIRF5 either untreated or treated with 2'3'-cGAMP.

B) Fold changes in gene expression as determined by qRT-PCR in samples microinjected with shRNAs targeting EGFP, nvIRF2, nvIRF-67, or nvIRF4 either untreated or treated with 2'3'-cGAMP. Note that IRF2 is not induced by cGAMP and was mostly undetected in all samples; therefore it is possible that the knockdowns were unsuccessful.

C) Volcano plot showing differential gene expression as determined by RNA-Seq in 48 hour embryos treated with 2'3'-cGAMP that were injected with GFP shRNA or nvSTAT shRNA. Positive fold-change indicates higher expression in GFP shRNA injected embryos. The GFP shRNA samples here are the same as those shown in Figure 2A.

D) Fold changes in gene expression as determined by Nanostring in embryos microinjected with shRNAs targeting EGFP or nvSTAT either untreated or treated with 2'3'-cGAMP.

Figure S5: Anti-viral gene induction is not dependent on nvNF-κB

A) Fold changes in gene expression as determined by Nanostring in embryos microinjected with shRNAs targeting EGFP or nvNF-κB, either untreated or treated with 2'3'-cGAMP. The GFP shRNA samples here are the same as those shown in Figure S4. GBP-806 expression included to show anti-bacterial gene induction is lower in these samples.

B) Fold changes in nvLysozyme expression as determined by qRT-PCR in embryos microinjected with shRNAs targeting EGFP or nvNF-κB, either untreated or treated with 2'3'-cGAMP.

A+B) No significant differences in gene expression are observed between any cGAMP treated samples.

C) qRT-PCR of putative anti-viral genes assayed at 48 hours post *Pa* infection ($2x10^7$ CFU/ml). Each point represents one biological replicate; unpaired t test performed on ΔΔCt before log transformation. **$p \leq 0.01$.

Figure S6: Phylogenetic study of *N. vectensis* GBPs

A) Phylogenetic tree of mammalian GTPases and putative *N. vectensis* GBPs made with the full protein sequences. Branches with mammalian interferon-induced GTPases and cGAMP-induced *N. vectensis* GBPs are colored red; these tend to cluster together. Domain structures of *N. vectensis* GBPs are displayed.

B) Alignment of the GTPase domains of all *N. vectensis* GBPs and select mammalian GTPase. Conserved GBP and atlastin specific residues are highlighted in red.

Figure S7: nvDae4 cleaves peptidoglycan from Gram-positive bacteria. Partial HPLC chromatograms of *Staphylococcus epidermis* peptidoglycan sacculi products resulting from incubation with buffer only (no enzyme), or 1 µM nvDae4 WT or C63A enzyme.

Table S1: Cnidarian-specific genes that are induced by 2'3'-cGAMP in an nvNF-κB-dependent manner

NCBI Gene ID	Domains	Only in Cnidaria?	Only in Anthozoa?	Homolog found in immune cells in *Stylophora pistillata (1)*?
5501851	none	yes	yes	no
5504224	MDN1 super family	yes	yes	no
5516219	none	yes	yes	no
5518710	none	yes	yes	no
116603205	none	yes	yes	No homolog in *Stylophora pistillata*
116603727	none	yes	yes	No homolog in *Stylophora pistillata*
116604070	none	yes; except Bacillus spore coat proteins	yes; except Bacillus spore coat proteins	No homolog in *Stylophora pistillata*
116604505	none	yes	yes	No homolog in *Stylophora pistillata*
116612667	none	yes	yes	no
116613998	none	yes	yes	no
116616875	none	yes	yes; similar to 116620239	No homolog in *Stylophora pistillata*
116619128	none	yes	no	yes
116620239	none	yes	yes; similar to 116616875	No homolog in *Stylophora pistillata*

Dataset 1: DESeq2 results for all RNA-Seq experiments
Dataset 2: All primers, oligos, and probes used in this study

SI References

1.   S. Levy *et al.*, A stony coral cell atlas illuminates the molecular and cellular basis of coral symbiosis, calcification, and immunity. *Cell* 10.1016/j.cell.2021.04.005 (2021).