Localization and RNAi-driven inhibition of a *Brugia malayi* encoded

Interleukin-5 Receptor Binding protein

Rojelio Mejia[1], Sasisekhar Bennuru[1], Yelena Oksov[2] Sara Lustigman[2],

Gnanasekar Munirathinam[3,] Ramaswamy Kalyanasundaram[3], Thomas B.

Nutman[1*]

[1]Laboratory of Parasitic Diseases, National Institute of Allergy and Infectious

Diseases, Bethesda, Maryland, USA. [2]Lindsay F. Kimball Research Institute,

New York Blood Center, New York, NY, and [3]Department of Biomedical

Sciences, College of Medicine, University of Illinois, Rockford, IL, USA.

Running title: *Brugia malayi* RNAi in L3

Corresponding author: Thomas B. Nutman tnutman@niaid.nih.gov

Keywords: *Brugia malayi*, RNAi, Host-parasite defense, Interleukin 5

Abstract

A molecule termed BmIL5Rbp (aka Bm8757) was identified from *Brugia malayi*

filarial worms and found to competitively inhibit human IL-5 binding to its human

receptor.

After the expression and purification of a recombinant BmIL5Rbp and generation

of BmIL5Rbp-specific rabbit antibody, we localized the molecule on *B. malayi*

worms through immunohistochemistry and immunoelectron microscopy. RNA

interference was used to inhibit BmIL5Rbp mRNA and protein production.

BmIL5Rbp was shown to localize to the cuticle of *Brugia malayi* and to be

released in their excretory/secretory products. RNAi inhibited BmIL5Rbp mRNA

production by 33% and reduced the surface protein expression by ~50% and

suppressed the release of BmIL5Rbp in the excretory/secretory products.

RNAi has been used successfully to knock down the mRNA and protein

expression of BmIL5Rbp in the early larval stages of *B. malayi* and provided a

proof-of-principle for the local inhibition of the human IL5 receptor. These findings

provide evidence that a parasite encoded IL5R antagonist could be utilized

therapeutically.

Introduction

Filarial parasites are vector-borne nematodes that are responsible for >200

million infections, the most pathogenic of which are caused by *Wuchereria*

bancrofti, Brugia malayi (two of the causative agents of lymphatic filariasis),

Onchocerca volvulus (the causative agent of onchocerciasis or "river blindness")

and *Loa loa* (the causative agent of loiasis or "the Africa eyeworm"). Infection

occurs when third-stage larvae (L3) are introduced into the skin, after which they

migrate through the dermis into either deep or superficial lymphatics, where they

continue their development. With the identification of innate lymphoid cells (ILC)

(1) and tissue-resident eosinophils in the skin (2) and the fat associated

subcutaneous tissue, and the fact that the filarial L3s can evade these crucial

innate cell populations to establish infection suggests that there must be a

common mechanism used by these skin-transiting parasites to establish infection

in humans. This lack of response is believed to be due to the parasite's evasive

immune mechanisms, especially the third-stage larvae (3).

Several helminth-encoded cytokine and chemokine mimics have been

identified, largely through *in silico* analyses of genomic information. These

include migration inhibitory factor (4-6) and TGF-β (7-9). A number of additional

studies have used other methods to identify parasite-encoded molecules that are

either agonists or antagonists of host-derived cytokine. Indeed, we identified in

2004, using a *B. malayi* L3 phage display library (10) a parasite-produced

molecule called BmIL5Rbp (subsequent genome annotation Bm8757).

To understand the precise localization of this molecule in the *B. malayi* L3

and explore the feasibility of using L3s that fail to express this molecule for *in*

vivo assessment/functional capabilities in relevant animal models, we used

immunostaining and confocal microscopy to demonstrate the location and

expression of BmIL5Rb. Using the RNAi approach on adults and microfilariae of

B. malayi (11-14), we demonstrate its utility in L3s so that its role could be

elucidated at the host-parasite interface.

Results

BmIL5Rbp is located on the surface of the worm and in the excretory-

secretory products.

Using immunostaining to localize BmIL5Rbp on various stages of *B. malayi*, it

can be seen that BmIL5Rbp was found to be expressed on the surface of the

worm using electron microscopy (Figure 1) and laser confocal microscopy of the

adult male, female, and L3 (Figure 2). As seen, BmIL5Rbp (red) is only

demonstrated in the stacked images on the parasite surfaces. Interestingly, the

protein is found dispersed throughout the worm's cuticle. No BmIL5Rbp was

seen internal to the cuticle even when using fragmented or transected worms

(data not shown).

Silencing BmIL5Rbp mRNA in L3 worms

To undertake the silencing of the BmIL5Rbp in L3s, we soaked L3 worms for two

days in dsRNAi constructs for BmIL5Rbp or BmCPL (control) and examined the

mRNA expression of the BmIL5Rbp. As shown in Figure 3A, BmIL5Rbp mRNA

expression was diminished significantly compared to the BmCPL inhibited worms

(33.1% vs. 4.89%, p = 0.0007) (Figure 3A). Approximately 85% of worms

survived in both groups. Supernatants from live worms used in these RNAi

cultured showed that the BmIL5Rbp protein content from these silenced worms

showed levels of BmIL5Rbp (Figure 3B) that were 50% of the control worms (GM

205 pg/ml for the control siRNA construct and GM 105 pg for the siBmIL5Rbp

group (P=0.034).

Detection and quantification of surface and excretory/secretory BmIL5Rbp

There was a 54% decrease in surface protein expression between the two

groups of worms, as shown by the confocal images in Figure 4A. Quantification

of fluorescence at the surface of the worms (Imaris) showed a similar reduction

induced by siBmIL5Rbp (2.125 Intensity Sum/Voxels compared to 1.146 Intensity

Sum/Voxels (p = 0.0025).

Human IL-5 exhibited decrease binding to human IL-5 receptors with

increasing rBmIL5Rbp concentrations (Spearman r = -0.9977, p < 0.0001)

(Supplementary Figure 1). RNAi worms excrete less BmIL5Rbp to inhibit binding

of human IL5 to its receptor. Supernatant from worms (200 in each group) at two

days of soaking showed a decrease in inhibition of human IL-5 for the BmIL5Rbp

RNAi worms compared to the negative control worms by 23% (Figure 4B).

Results are presented in percentage of inhibition to control worms.

Discussion

Lymphatic filarial parasites are notorious for their ability to manipulate host

immune responses. In the present study, we show that one such mechanism

includes interfering with the function of host-derived IL-5. Here we characterize

and localize the BmIL5Rbp produced by the third stage infective larvae of *B.*

malayi, a secreted molecule that can bind to soluble human IL-5 receptor alpha

and block the binding of human IL-5 to this receptor. Thus, BmIL5Rbp, a protein

produced by *B. malayi,* appears to be a novel host-immunomodulatory molecule,

acting as an IL5 antagonist.

The ability of a nematode-derived protein to alter the human host immune

response is a unique evolutionary response that likely helps the worm to

establish a foothold. We describe a nematode-produced protein found on the

worm's surface (Figures 1, 2) and in its excretory/secretory product (Figures 3,

4).

B. malayi, along with all of the pathogenic filariae, induce a robust

eosinophil response, particularly in the acute phase of infection (19, 20) related

to the tissue migration L3 and L4 larvae of filarial nematodes (21-23). Our data

suggest that these parasites secrete a molecule that antagonizes the activity of

human IL-5, a cytokine found both early in infection and within days of

anthelmintic chemotherapy (24-26).

The localization of BmIL5Rbp to the worm's surface appears to be directly

on (and within) the cuticle. Its location gives it accessibility to directly tissue-

dwelling host cells bearing the IL5R (eosinophils, basophils, mast cells). It

suggests that there may be a physiologic role for this molecule at the host tissue

level. Since eosinophil worm killing is contact or localized dependent action (27),

then the immediate areas, including the tissue, around the worm's surface will

have higher concentrations of BmIL5Rbp, leading possibly to a loss of eosinophil

survival at the site of inflammation/tissue damage.

The use of RNAi to selectively inhibit BmIL5Rbp helps to provide proof of

principle that the worm produces a protein to inhibit the binding of human IL5 to

its receptor. The RNAi-treated worms produced specifically less BmIL5Rbp that

had less inhibition of human IL-5 to its human IL-5 receptor *in vitro*. With the

advent of CRISPR-CAS9 in genomic splicing (28), the ability to knock out this

gene now is a possibility (29-31). However, for the filarial nematodes, RNAi

remains the most useful method of gene silencing (11, 12, 14, 17, 32-34).

Conclusion

While this study shows that it is feasible to silence the BmIL5Rb expression, *in*

vivo experiments will be required to demonstrate the importance of this molecule

(13). Our study provides a framework for utilizing these parasite-encoded

cytokine antagonists for mechanistic and/or host-directed therapy studies that

should, in turn, provide new insights into specific host-parasite interactions.

Methods

Expression and purification of rBmIL5Rbp

The entire coding sequence (Supplementary Table 1) of the IL5Rbp was

amplified and inserted into entry clones containing an upstream Tev protease

cleavage site (ENLYFQG). Both entry clones were completely sequenced to

verify that they contained the correct sequence. Expression clones were

transferred to six different T7 promoter-based Destination vectors for *E. coli*

expression. Among the six, one termed 2110-X1-566 contained His6-MBP-tev-

I5RBP that was found to produce the most abundant soluble protein when

transformed into *E. coli* Rosetta (DE3). The soluble fraction of the MBP-BmIL5

was bound to the affinity column of the immobilized metal-ion affinity

chromatography (IMAC) and eluted at 200 mM imidazole (~200 mM). The eluted

material was incubated overnight with 5% TEV protease at 16^0 C. The

contaminating components of the protease reaction were removed in a

subsequent IMAC step. The final protein was ~ 95% pure as confirmed by SDS-

PAGE analysis and was stored at 1mg/ml in PBS, pH 7.5 at -80^0 C until used.

Recombinant BmIL5 (rBmIL5Rbp) was also expressed in Sf9 and Hi5 cells by

infection with baculoviruses. 2 x 10^6 insect cells from 48 post infection Hi5 cells

incubated at 21°C were resuspended in lysis buffer (20 mM Na phosphate pH

7.5, 500 mM NaCl, 5% glycerol, Complete protease inhibitor, Roche Diagnostics,

Rotkreuz, Switzerland) and sonicated until cells were > 99% lysed as determined

by microscopic examination. Soluble and insoluble fractions were separated by

centrifugation. Baculovirus-infected Hi5 cell paste was resuspended with

extraction buffer (20 mM HEPES buffer, pH7.3, 500 mM NaCl, 45 mM imidazole,

5 mM $MgCl_2$, 10% glycerol, 2 mm βME, complete protease inhibitor [Roche] at 2

tablets/50 ml of lysate) to a volume in milliliters equal to four times the wet weight

of the pellet in grams. The sample was sonicated to lyse the cells (verified by

microscopic examination), treated with 1.0 U benzonase/ml for 20 min on ice,

clarified by centrifugation at 111,000 x g for 50 min, filtered (0.45 µm PES

membrane), diluted five-fold with extraction buffer and applied to a 5 ml HisTrap

column (GE Healthcare, Chicago, IL) equilibrated with extraction buffer. The

column was washed to baseline, proteins eluted over a 20-column volume (CV)

gradient to 400 mM imidazole and fractions analyzed by SDS-PAGE. The pool

created from the IMAC fractions was dialyzed against binding buffer without

additional imidazole and treated with TEV protease (1% v/v with a lab stock) at

RT for 4 hours and then shifted to 16 °C overnight and analyzed by SDS-PAGE.

An additional IMAC step similar to the initial IMAC was used to purify the target

protein away from the contaminants of the protease digest. Fractions were

analyzed by SDS-PAGE.

Electron microscopy and immunogold staining

The native BmIL5Rbp was identified in thin sections of *B. malayi* L3 by immuno-

electron microscopy. The larvae were fixed with 4% paraformaldehyde/0.15%

glutaraldehyde in 0.1 m sodium cacodylate buffer for 1 hour and embedded in LR

White medium as previously described (15). Thin sections (70 nm) were then

blocked with 2% BSA and incubated with primary antibody (rabbit anti-rBmIL5) or

control serum (pre-immunization bleed of the same rabbit) overnight at 4°C.

Following washes in PBS containing 0.1% BSA and 0.01% Tween-20, grids were

incubated with gold-labeled goat anti-mouse IgG (Fc) at 1:20 dilution (Amersham

Pharmacia Biotech, Little Chalfont, United Kingdom) for 1 hour at room

temperature, washed, counterstained in 5% uranyl acetate solution and observed

using a Philips 410 electron microscope (Figure 1).

Immunostaining of L3 worms

L3 worms were fixed with a final concentration of 2% paraformaldehyde

overnight at 4°C. Worms were washed three times with blocking buffer: 1%

bovine serum albumin, 0.5% Triton X-100, 0.1% sodium azide, phosphate-

buffered saline (PBS). The worms were soaked in 10 mMol sodium citrate (in

blocking buffer) at 70°C for 60 minutes, washed three times in PBS, and soaked

in blocking buffer overnight at 4°C. The worms were washed three times in PBS

and then soaked in rabbit anti-BmIL5Rbp (1:10 in blocking buffer) for three days

at 4°C. After washing three times in blocking buffer for 30 minutes, the worms

were incubated with 1:500 dilution of goat anti-rabbit IgG Alexa Fluor 594

(Thermo Fisher Scientific, Waltham, MA) in blocking buffer for three days at 4°C.

Phalloidin Alexa Fluor 488 (Thermo Fisher Scientific) in methanol (200 units/mL)

was added and incubated overnight at 4°C. The worms were then washed three

times with moderate shaking at 4°C for three hours each. Individual worms were

collected and set with a mounting solution (Hardset VectaShield Vector H-1400,

Vector Laboratories, Burlingame, CA) overnight a 4°C. All worms were visualized

using a confocal microscope, Leica SP5 X-WLL (Mannheim, Germany). Dapi

stain was detected at 460 nm, Phalloidin at 520 nm, Alexa Fluor 594 at 620 nm

with excitation at 595 nm at 15% laser power, a smart gain of 838, offset -0.3%.

Synthesis of siRNA constructs for BmIL5Rbp

Primers and probes for the gene sequence of BmIL5Rbp were designed

using siRNA Target Designer-Version 1.6, Promega (Madison, WI) that used

proprietary algorithms to select the optimal primer/probe sequences.

Forward 5'AAA AAG GGA ATT AGT TGT TG3'

Reverse 5'ATA CTA ACA AAC TTC TGC AAA TT3'

Plasmid DNA for BmIL5Rbp was generated and transformed to

recombinant protein using TOP 10 *E. coli* competent cells (AMP resistant)

(Thermo Fisher Scientific, Waltham, Ma). RNA PCR was performed using T7, T3

primers. Single-stranded RNA was annealed per-protocol (Agilent Technologies,

Santa Clara, CA) with a final product of double-stranded RNA (dsRNA). DsRNA

was concentrated using microcentrifugation filters and the final product was

resuspended in PBS.

RNAi of *Brugia malayi* L3 worms

The L3 worms were cultured, as described previously (16). Double-

stranded BmIL5Rbp (dsBmIL5Rbp) RNA was introduced to L3 worms by

soaking, in which 20 L3 worms were soaked in 100 µL of dsBmIL5Rbp (500

µg/mL) in L3 media (16) for two days at 37°C. The media was removed, and 100

µL TRIzol® (Thermo Fisher Scientific) was added to the L3's and quickly frozen.

The L3 larvae went through three freeze-thaw cycles and homogenated for five

minutes at room temperature with Qiagen shredders (Qiagen, Venlo,

Netherlands), washing with 100 µl TRIzol® (Thermo Fisher Scientific) three times.

The macerated larvae were centrifuged at 12,000 g for one minute at 4°C

between each wash. A total of 20 µL chloroform was then added to the shredded

larvae and shaken vigorously for 15 seconds. The mixture was then centrifuged

for 15 minutes at 12,000 g at 4°C. The aqueous phase was retained, and equal

volumes of isopropanol were added. The aqueous phase mixture was

centrifuged at 16,000 g for 15 minutes at 4°C. The pellet was washed with 70%

ethanol then centrifuged at 16,000 g for 5 minutes at 4°C. The pellet was air-

dried for 20 minutes, then re-suspended with 10 µL TE buffer (Thermo Fisher

Scientific). Reverse transcription PCR was performed using standard techniques

to the final volume of 100 µL. An internal control siRNA for *Brugia malayi*

cathepsin L intron (BmCPL) (17) was created using the above protocol and was

used to show specific inhibition of BmIL5Rbp using a SYBR Green real-time PCR

(Applied Biosystems) per manufacturer specifications.

Primers for BmCPL (14).

Forward: 5' GAC AAA GAT TAC AAA CAG GGC3'

Reverse: 5' TGA TTG GGC AGT CGA AGT C3'

Real-time PCR quantification of BmIL5Rbp

Real-time PCR with final volumes of 20 μL, 1 μL of the probe (FAM-5'AAG

CAA GCA TTG ATT TCT3'-MGBNFQ), 2 μL of template, 10 μL Taqman™ fast

mix (Applied Biosystems, Foster City, CA), forward/reverse primers 1 μL each, 5

μl of dH$_2$O. Primers for BmIL5Rbp were selected upstream of the siRNA

selection using the following primers:

Forward: 5'AAA ATG ATG GAA GCA GCA GAA ACT G3'

Reverse: 5'TAC AGG AAT ACA TCA TGC TCA CAA GT3'

All reactions were performed on an ABI 7900HT Fast Real-Time PCR System

(Applied Biosystems). All primers and probes were purchased from Applied

Biosystems. The PCR conditions were 95°C for 1 min 1 x cycle; 95°C for 25 s,

55°C for 20 s, 72°C for 1 min 35x cycles; 72°C for 10 min 1x cycle.

For standard comparison, housekeeping control Bm-tub primers were used with

SYBR Green real-time PCR (Applied Biosystems) per manufacturer

specifications (14).

Primers for Bm-tub (17)

Forward: 5'ATA TGT GCC ACG AGC AGT C3'

Reverse: 5'CGG ATA CTC CTC ACG AAT TT3'

Penetration of dsBmIL5Rbp in live *Brugia malayi*

DsBmIL5Rbp was Cy3 labeled using the CyDye™Post-Labelling Reactive Dye

Pack (GE, Piscataway, NJ) per manufacturer's instructions. A total of 5 μL

dsBmIL5Rbp (2000 μg/mL) was added to 100 mMol sodium bicarbonate and

mixed with Cy3 in 20 μL DMSO stirred in the dark for 90 minutes at room

temperature. Then 15 µL of 4M hydroxylamine was added and mixed for 15

minutes in the dark at room temperature. Next, the mixture was precipitated with

5 µL of 5M sodium chloride for 60 minutes at 20°C. The solution was centrifuged

at 16,000 g for 15 minutes at 4°C. Then to the supernatant, 175 µL of 70% cold

ethanol was added, then centrifuged 16,000 g for 5 minutes at 4°C. Next, the

supernatant was removed, and the pellet air-dried for 10 minutes in the dark and

solubilized in 100 µL in L3 media (16). A total of 20 live L3 worms were soaked in

the dsBmIL5Rbp media overnight. The worms were then fixed and mounted at

the above methods. Also, 20 other worms were used in non-CY3 L3 media as a

control group

Quantifying surface BmIL5Rbp on L3 worms using confocal microscopy.

Using the above RNAi methods, approximately 60 worms were exposed to

BmIL5Rbp RNAi constructs. These were mounted along with appropriate control

worms. Using a laser confocal microscope, capturing several 0.5mm stack

images were obtained of entire worms. The intensity was calculated by using

sum/voxels in Imaris (South Windsor, CT). Voxels are the total number of pixels

at a specific wavelength.

Quantification of excretory/secretory BmIL5Rbp in L3 worms

BmIL5Rbp was measured using an immune dot blot. Briefly, nitrocellulose paper

(Schleicher and Schuell (Keene, NH). was trimmed to fit a 96-well manifold. The

paper was activated in methanol for three minutes and then washed for 1 min in

distilled water. The paper was dried, and 50 µL of rBmIL5Rbp (baculovirus

derived) in various dilutions, duplicates were placed in wells with 50 µL of

supernatants from RNAi and control worms. The plate was incubated at 37°C for

one hour. The plate was dried, and the membrane was placed in a blocking

buffer (5% milk, 5% rabbit serum, 5% goat serum) overnight at 4°C with gently

tilting action. The membranes were incubated at room temperature in primary

antibody Rabbit anti-BmIL5Rbp 1:100 dilution in 5% milk solution. The

membrane was washed three times with 0.5% Tween 20 (Sigma-Aldrich, St.

Louis, MO) in PBS solution for 5 min each cycle. Then a secondary antibody

Goat anti-Rabbit Horseradish peroxidase (HRP) (Thermo Fisher Scientific)

1:1000 dilution in 5% milk was added for a 1-hour incubation at room

temperature. The membrane was washed three times with 0.5% Tween and PBS

solution for 5 minutes each cycle. For the ELISA developing step, a SuperSignal

West Pico Chemiluminescent Substrate kit (Thermo Fisher Scientific) was used

on the membrane as per the manufacturer's instructions. Luminescence was

detected using Kodak X-OMAT film (Rochester, NY) and exposed for 10 minutes,

then quantitated using BmIL5Rbp standard curves.

Measurement of decrease inhibition of human IL5 to its receptor *in vitro*

using RNAi worms

A total of 50 µL/well of recombinant human IL5-Receptor alpha (R&D Systems,

Minneapolis, MN) final concentration of (4 µg/mL) was plated using coating buffer

(18) on Immulon 4 plates (Thermo Fischer Scientific) overnight at 4°C. The plates

were then washed with (0.2% Tween 20 in PBS) 6 times. A total of 100 µL of

ELISA blocking buffer (0.05 Tween 20, 5% bovine serum albumin, PBS) was

added for 1 hour at 37°C, with subsequent washing as above. Afterward, 50 µL

of E/S product from 200 worms in each group (BmCPL Internal control and

BmIL5Rbp RNAi) was added and incubated for 1 hour at 37°C. The plates were

washed again as above. Subsequently, 25 µL of biotinylated recombinant human

IL5 (rhIL5) protein (50 µg/mL) (R&D Systems) was added at 1:50 in PBS.

Biotinylated rhIL5 was produced using the Biotin-X-NHS system (Merck,

Darmstadt, Germany). A total of 25 µL of rhIL5 (100 µg/mL) was immediately

added for a final concentration of 50 µg/mL and incubated at 37°C for one hour.

The plate was washed as above, and 50 µL of streptavidin (Thermo Fisher

Scientific) was added and incubated at 37°C for one hour. The plate was washed

for a final time as above. The plate was developed with pNPP phosphatase

substrate (Sigma, St. Louis, MO) per the manufacturer's instructions for 30 min at

room temperature. Plates were read at 405 nm in an ELISA reader (SpectraMax

Plus; Molecular Devices, Sunnyvale, CA). Similar methods were used with

increasing concentrations of rBmIL5Rbp inhibiting hIL-5 to human IL-5 receptor

(Supplementary Figure 1). In determining the optimal concentration for

Biotinylated rhIL5, a separate set of experiments was done with 2-fold dilutions

using the above methods. Results showed that a 1:100 final concentration of

biotinylated rhIL5 was the optimal concentration (data not shown).

Statistics and calculations

All statistical analysis was performed using Prism version 5.0d (GraphPad, La

Jolla, CA). Comparisons were analyzed using the Mann-Whitney test, and P

values < 0.05 were considered significant. RNAi results were shown as the

percent reduction compared to a standard housekeeping control gene (Bm-tub)

as previously described (17). In summary, the control group value was set as the

maximum amount with the RNAi-treated worms reported as a relative reduction

compared to the internal control group (BmCPL). Calculation of intensity

sum/voxels was completed with Imaris Software (Zurich, Switzerland).

References

1. Artis D, Spits H. 2015. The biology of innate lymphoid cells. Nature 517:293-
301.
2. Cheng LE, Sullivan BM, Retana LE, Allen CD, Liang HE, Locksley RM. 2015.
IgE-activated basophils regulate eosinophil tissue entry by modulating
endothelial function. J Exp Med 212:513-24.
3. Cotton RN, McDonald-Fleming R, Boyd A, Spates K, Nutman TB, Tolouei
Semnani R. 2015. Brugia malayi infective larvae fail to activate Langerhans
cells and dermal dendritic cells in human skin. Parasite Immunol 37:79-91.
4. Wang Y, Lu M, Wang S, Ehsan M, Yan R, Song X, Xu L, Li X. 2017.
Characterization of a secreted macrophage migration inhibitory factor
homologue of the parasitic nematode Haemonchus Contortus acting at the
parasite-host cell interface. Oncotarget 8:40052-40064.
5. Filbey KJ, Varyani F, Harcus Y, Hewitson JP, Smyth DJ, McSorley HJ, Ivens A,
Nylen S, Rottenberg M, Loser S, Maizels RM. 2019. Macrophage Migration
Inhibitory Factor (MIF) Is Essential for Type 2 Effector Cell Immunity to an
Intestinal Helminth Parasite. Front Immunol 10:2375.
6. Chauhan N, Sharma R, Hoti SL. 2015. Identification and biochemical
characterization of macrophage migration inhibitory factor-2 (MIF-2)
homologue of human lymphatic filarial parasite, Wuchereria bancrofti. Acta
Trop 142:71-8.
7. Gomez-Escobar N, Gregory WF, Maizels RM. 2000. Identification of tgh-2, a
filarial nematode homolog of Caenorhabditis elegans daf-7 and human
transforming growth factor beta, expressed in microfilarial and adult stages
of Brugia malayi. Infect Immun 68:6402-10.

8.   White MPJ, McManus CM, Maizels RM. 2020. Regulatory T-cells in helminth
infection: induction, function and therapeutic potential. Immunology
160:248-260.
9.   Johnston CJC, Smyth DJ, Kodali RB, White MPJ, Harcus Y, Filbey KJ, Hewitson
JP, Hinck CS, Ivens A, Kemter AM, Kildemoes AO, Le Bihan T, Soares DC,
Anderton SM, Brenn T, Wigmore SJ, Woodcock HV, Chambers RC, Hinck AP,
McSorley HJ, Maizels RM. 2017. A structurally distinct TGF-beta mimic from
an intestinal helminth parasite potently induces regulatory T cells. Nat
Commun 8:1741.
10.   Gnanasekar M, Rao KV, He YX, Mishra PK, Nutman TB, Kaliraj P, Ramaswamy
K. 2004. Novel phage display-based subtractive screening to identify vaccine
candidates of Brugia malayi. Infect Immun 72:4707-15.
11.   Singh M, Singh PK, Misra-Bhattacharya S. 2012. RNAi mediated silencing of
ATPase RNA helicase gene in adult filarial parasite Brugia malayi impairs in
vitro microfilaria release and adult parasite viability. J Biotechnol 157:351-8.
12.   Landmann F, Foster JM, Slatko BE, Sullivan W. 2012. Efficient in vitro RNA
interference and immunofluorescence-based phenotype analysis in a human
parasitic nematode, Brugia malayi. Parasit Vectors 5:16.
13.   Kushwaha S, Singh PK, Shahab M, Pathak M, Bhattacharya SM. 2012. In vitro
silencing of Brugia malayi trehalose-6-phosphate phosphatase impairs
embryogenesis and in vivo development of infective larvae in jirds. PLoS Negl
Trop Dis 6:e1770.
14.   Aboobaker AA, Blaxter ML. 2003. Use of RNA interference to investigate gene
function in the human filarial nematode parasite Brugia malayi. Mol Biochem
Parasitol 129:41-51.
15.   Zheng HJ, Tao ZH, Zhang XH, Piessens WF. 1989. [Further study on in vitro
culture of third-stage larvae of Wuchereria bancrofti and Brugia malayi].
Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi 7:276-9.
16.   Bennuru S, Semnani R, Meng Z, Ribeiro JM, Veenstra TD, Nutman TB. 2009.
Brugia malayi excreted/secreted proteins at the host/parasite interface:
stage- and gender-specific proteomic profiling. PLoS Negl Trop Dis 3:e410.
17.   Ford L, Zhang J, Liu J, Hashmi S, Fuhrman JA, Oksov Y, Lustigman S. 2009.
Functional analysis of the cathepsin-like cysteine protease genes in adult
Brugia malayi using RNA interference. PLoS Negl Trop Dis 3:e377.
18.   Ravi V, King TP, Andersen JF, Nutman TB, Neva FA. 2005. Strongyloides
stercoralis recombinant NIE antigen shares epitope with recombinant Ves v
5 and Pol a 5 allergens of insects. Am J Trop Med Hyg 72:549-53.
19.   Gebreselassie NG, Moorhead AR, Fabre V, Gagliardo LF, Lee NA, Lee JJ,
Appleton JA. 2012. Eosinophils preserve parasitic nematode larvae by
regulating local immunity. J Immunol 188:417-25.
20.   Specht S, Saeftel M, Arndt M, Endl E, Dubben B, Lee NA, Lee JJ, Hoerauf A.
2006. Lack of eosinophil peroxidase or major basic protein impairs defense
against murine filarial infection. Infect Immun 74:5236-43.
21.   Patnode ML, Bando JK, Krummel MF, Locksley RM, Rosen SD. 2014.
Leukotriene B4 amplifies eosinophil accumulation in response to nematodes.
J Exp Med 211:1281-8.

22.   Faccioli LH, Mokwa VF, Silva CL, Rocha GM, Araujo JI, Nahori MA, Vargaftig
BB. 1996. IL-5 drives eosinophils from bone marrow to blood and tissues in a
guinea-pig model of visceral larva migrans syndrome. Mediators Inflamm
5:24-31.
23.   Chitkara RK, Krishna G. 2006. Parasitic pulmonary eosinophilia. Semin Respir
Crit Care Med 27:171-84.
24.   Limaye AP, Abrams JS, Silver JE, Ottesen EA, Nutman TB. 1990. Regulation of
parasite-induced eosinophilia: selectively increased interleukin 5 production
in helminth-infected patients. J Exp Med 172:399-402.
25.   Cooper PJ, Chico ME, Sandoval C, Nutman TB. 2004. Atopic phenotype is an
important determinant of immunoglobulin E-mediated inflammation and
expression of T helper cell type 2 cytokines to ascaris antigens in children
exposed to ascariasis. J Infect Dis 190:1338-46.
26.   Klion AD, Law MA, Noel P, Kim YJ, Haverty TP, Nutman TB. 2004. Safety and
efficacy of the monoclonal anti-interleukin-5 antibody SCH55700 in the
treatment of patients with hypereosinophilic syndrome. Blood 103:2939-41.
27.   Cadman ET, Thysse KA, Bearder S, Cheung AY, Johnston AC, Lee JJ, Lawrence
RA. 2014. Eosinophils are important for protection, immunoregulation and
pathology during infection with nematode microfilariae. PLoS Pathog
10:e1003988.
28.   Ghorbal M, Gorman M, Macpherson CR, Martins RM, Scherf A, Lopez-Rubio JJ.
2014. Genome editing in the human malaria parasite Plasmodium falciparum
using the CRISPR-Cas9 system. Nat Biotechnol 32:819-21.
29.   You H, Mayer JU, Johnston RL, Sivakumaran H, Ranasinghe S, Rivera V,
Kondrashova O, Koufariotis LT, Du X, Driguez P, French JD, Waddell N, Duke
MG, Ittiprasert W, Mann VH, Brindley PJ, Jones MK, McManus DP. 2021.
CRISPR/Cas9-mediated genome editing of Schistosoma mansoni
acetylcholinesterase. FASEB J 35:e21205.
30.   Shao H, Li X, Lok JB. 2017. Heritable genetic transformation of Strongyloides
stercoralis by microinjection of plasmid DNA constructs into the male
germline. Int J Parasitol 47:511-515.
31.   Liu C, Grote A, Ghedin E, Unnasch TR. 2020. CRISPR-mediated Transfection of
Brugia malayi. PLoS Negl Trop Dis 14:e0008627.
32.   Song C, Gallup JM, Day TA, Bartholomay LC, Kimber MJ. 2010. Development of
an in vivo RNAi protocol to investigate gene function in the filarial nematode,
Brugia malayi. PLoS Pathog 6:e1001239.
33.   Misra S, Gupta J, Misra-Bhattacharya S. 2017. RNA interference mediated
knockdown of Brugia malayi UDP-Galactopyranose mutase severely affects
parasite viability, embryogenesis and in vivo development of infective larvae.
Parasit Vectors 10:34.
34.   Iqbal S, Fosu-Nyarko J, Jones MG. 2016. Genomes of parasitic nematodes
(Meloidogyne hapla, Meloidogyne incognita, Ascaris suum and Brugia
malayi) have a reduced complement of small RNA interference pathway
genes: knockdown can reduce host infectivity of M. incognita. Funct Integr
Genomics 16:441-57.
478

Figure Legends

Figure 1. Evaluation of *Brugia malayi* L3 larvae sections under a transmission

electron microscope showed that BmIL5Rbp is predominantly concentrated on

the surface of the parasite (arrows), suggesting that BmIL5Rbp is expressed on

the parasite's surface. The micrograph (A) shows worms coated with pre-

immunized rabbit sera against BmIL5Rbp compared to the micrograph (B, C)

using post-immunized rabbit sera identifying antibody localization on the worm's

cuticle. Micrograph (C) is at higher magnification with Bar 500 nm.

Figure 2. Immunohistochemical staining localization with post-immunization

rabbit sera to BmIL5Rbp and laser confocal microscopy on *Brugia malayi*.

Worms were stained for BmIL5Rbp (red), nuclei (blue), and actin (green). Adult

male worms (A, B), adult female worms (C, D), and infectious stage L3 worms

(E, F) all express BmIL5Rbp on the surface of the cuticle. No BmIL5Rbp was

seen internally in the worms, even dissected specimens. Control worms (B, D, F)

had minimal red staining with pre-immunized rabbit sera to BmIL5Rbp.

Figure 3. (A) BmIL5Rbp mRNA expression in live *Brugia malayi* L3 larvae is

inhibited using RNAi. RNAi-specific BmIL5Rbp dsRNA had a significant percent

reduction to the internal control BmCPL dsRNA (33.1% versus 4.9% respectively,

$p = 0.007$). A total of 450 worms were processed with an 85% survival rate in

both groups. (B) RNAi inhibits BmIL5Rbp protein expression in

excretory/secretory products of live *Brugia malayi* L3 larvae compared to control

worms (3.4 versus 8.2 pg/µL respectively, p = 0.034).

Figure 4. (A) RNAi can effectively reduce BmIL5Rbp on the surface of *Brugia*

malayi L3 larvae. Intensity sum is the calculation of all red-stained BmIL5Rbp

normalize to the total number of pixels (voxels) (Intensity Sum/Voxels) of all

stained components of the worm. Pre-RNAi worms showed significant BmIL5Rbp

staining to post-RNAi worms (2.125 Intensity Sum/Voxels compared to 1.146

Intensity Sum/Voxel respectively, p = 0.0025). (B) The supernatant (sups) from

RNAi *Brugia malayi* L3 larvae contain less BmIL5Rbp in their excretory/secretory

product compared to internal controls and thus had less inhibition of human IL-5

to its human receptor (63.5% versus 85.1% respectively, p = 0.005).

Figure 1.

Pre-Immunization sera

Post Immunization sera

Figure 2.

Figure 3.

Figure 4.