**STING mediates immune responses in the closest living relatives of animals**

Arielle Woznica[1*], Ashwani Kumar[2], Carolyn R. Sturge[1], Chao Xing[2], Nicole King[3], Julie K. Pfeiffer[1*]

[1] Department of Microbiology, University of Texas Southwestern Medical Center, Dallas TX 75390, USA
[2] McDermott Center Bioinformatics Lab, University of Texas Southwestern Medical Center, Dallas TX, USA
[3] Howard Hughes Medical Institute, and Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA 94720, USA.

*Correspondence: Arielle.Woznica@UTSouthwestern.edu and Julie.Pfeiffer@UTSouthwestern.edu

**Abstract**

Animals have evolved unique repertoires of innate immune genes and pathways that provide their first line of defense against pathogens. To reconstruct the ancestry of animal innate immunity, we have developed the choanoflagellate *Monosiga brevicollis*, one of the closest living relatives of animals, as a model for studying mechanisms underlying pathogen recognition and immune response. We found that *M. brevicollis* is killed by exposure to *Pseudomonas aeruginosa* bacteria. Moreover, *M. brevicollis* expresses STING, which, in animals, activates innate immune pathways in response to cyclic dinucleotides during pathogen sensing. *M. brevicollis* STING increases the susceptibility of *M. brevicollis* to *P. aeruginosa*-induced cell death and is required for responding to the cyclic dinucleotide 2'3' cGAMP. Furthermore, similar to animals, autophagic signaling in *M. brevicollis* is induced by 2'3' cGAMP in a STING-dependent manner. This study provides evidence for a pre-animal role for STING in antibacterial immunity and establishes *M. brevicollis* as a model system for the study of immune responses.

**Introduction**

Innate immunity is the first line of defense against pathogens for all animals, in which it is crucial for distinguishing between self and non-self, recognizing and responding to pathogens, and repairing cellular damage. Some mechanisms of animal immunity have likely been present since the last common eukaryotic ancestor, including RNAi, production of antimicrobial peptides, and the production of nitric oxide[1,2]. However, many gene families that play critical roles in animal innate immune responses are unique to animals[3].

Comparing animals with their closest relatives, the choanoflagellates, can provide unique insights into the ancestry of animal immunity and reveal other key features of the first animal, the 'Urmetazoan'[4–6]. Choanoflagellates are microbial eukaryotes that live in diverse aquatic environments and survive by capturing and phagocytosing environmental bacteria[7] using their "collar complex," an apical flagellum surrounded by actin-filled microvilli (Figure 1A)[7,8]. Several innate immune pathway genes once considered to be animal-specific are present in choanoflagellates, including cGAS and STING, both of which are crucial for innate responses to cytosolic DNA in animals (Figure 1- Supplement 1)[3,9,10]. Although the phylogenetic distribution of these gene families reveals that they first evolved before animal origins, their functions in choanoflagellates and their contributions to the early evolution of animal innate immunity are unknown.

STING (stimulator of interferon genes) is a signaling protein that activates innate immune responses to cytosolic DNA during bacterial or viral infection[11,12]. Although STING homologs are conserved in diverse invertebrate and vertebrate animals (reviewed in Margolis et al. 2017)[9,13,14], mechanisms of STING activation are best understood in mammals. In mammals, STING is activated by binding 2'3' cGAMP, an endogenous cyclic dinucleotide produced by the sensor cGAS (cyclic GMP-AMP synthase) upon detecting cytosolic DNA[15–20.]. In addition, cyclic dinucleotides produced by bacteria can also activate STING[17,21]. Importantly, STING domain-containing systems are present in bacteria where they may contribute to anti-phage defense[22,23], raising the possibility that eukaryotic STING-like proteins were acquired from lateral gene transfer[24]. Comparative genomics suggests that STING domains arose at least three independent times in eukaryotes, including once in the stem lineage leading to Choanozoa, the clade containing animals and choanoflagellates[24].

Choanoflagellates have already served as powerful models for studying the origin of animal multicellularity and cell differentiation[10,25–30] and are ideally positioned to yield insights into the evolution of animal immune pathways. Therefore, we sought to establish the choanoflagellate *Monosiga brevicollis* as a model for studying pathogen recognition and immune responses. Here, we report that *Pseudomonas aeruginosa* bacteria are pathogenic for *M. brevicollis.* Through our study of interactions between *P.*

*aeruginosa* and *M. brevicollis*, we determined that STING functions in the
choanoflagellate antibacterial response. In addition, we demonstrate that STING is
necessary for mediating responses to the STING agonist 2'3' cGAMP *in vivo*, and that
2'3' cGAMP induces STING-dependent autophagic signaling.  Our results demonstrate
that key features of STING-mediated immune responses are conserved in *M.*
*brevicollis*, thereby expanding our understanding of the pre-metazoan ancestry of
STING signaling.
88
89  **Results**
90
91  ***P. aeruginosa* has pathogenic effects on *M. brevicollis***
92
One impediment to studying immune responses in choanoflagellates has been
the lack of known choanoflagellate pathogens. While bacteria are obligate prey and can
regulate mating, multicellular development, and cell contractility in choanoflagellates, to
our knowledge no bacteria with pathogenic effects have been described [25,28,29,31–33]. For
this study, we focused on the choanoflagellate *Monosiga brevicollis,* which has a
sequenced genome[4] and grows robustly under laboratory conditions in co-culture with
*Flavobacterium* prey bacteria[5]. To identify potential pathogens of choanoflagellates*,* we
screened select bacteria – including environmental isolates and known animal
pathogens and commensals (Table 1) – to test whether any of these induced *M.*
*brevicollis* behavioral changes or reduced cell survival.
After co-culturing *M. brevicollis* with bacteria for 24 hours, only the
gammaproteobacterium *Pseudomonas aeruginosa,* a ubiquitous environmental
bacterium and opportunistic pathogen of diverse eukaryotes[34–37], altered the behavior
and growth dynamics of *M. brevicollis*. Under standard laboratory conditions, *M.*
*brevicollis* is a highly motile flagellate and swims up in the water column (Video 1,
Supplementary file 1). However, after 12-14 hours in the presence of *P. aeruginosa*
strains PAO1 and PA14, a large proportion of *M. brevicollis* cells settled to the bottom of
the culture dish (Video 1, Supplementary file 1 ). Immunofluorescence staining revealed
that *M. brevicollis* cells exposed to *P. aeruginosa* had truncated flagella compared to
cells exposed to *E. coli* or other bacteria that did not induce cell settling (Figure 1B). To
determine the effects of *P. aeruginosa* on cell viability, we added *P. aeruginosa* strain
PAO1 or control gammaproteobacteria to *M. brevicollis* and monitored cell density over
the course of 72 hours (Figure 1C). While *M. brevicollis* continued to proliferate in the
presence of control gammaproteobacteria, exposure to *P. aeruginosa* PAO1 resulted in
cell death.
Choanoflagellates prey upon bacteria and ingest them through phagocytosis[7,8].
However, many bacterial pathogens, including *P. aeruginosa*, have evolved strategies
to prevent or resist phagocytosis by eukaryotic cells[38,39]. Therefore, we examined

whether phagocytosis of *P. aeruginosa* is required to induce cell death. To track
phagocytosis, we added GFP-expressing *E. coli* DH5α (Figure 1D) or *P. aeruginosa*
PAO1 (Figure 1E) to *M. brevicollis* and monitored the cultures by live imaging. After one
hour, while 92% of *M. brevicollis* cells incubated with *E. coli*–GFP had GFP+ food
vacuoles, only 3% of cells incubated with PAO1-GFP had GFP+ food vacuoles (Figure
1F). *M. brevicollis* also robustly phagocytosed GFP-expressing *V. parahaemolyticus*
and *C. jejuni* (Figure 1F).
Next, to determine if *P. aeruginosa* broadly disrupts *M. brevicollis* phagocytosis,
which could induce cell death through starvation, we incubated *M. brevicollis* with GFP-
expressing PAO1 or GFP-expressing *E. coli* for one hour, and then added 0.2 mm
fluorescent beads for an additional 30 minutes as an independent measure of
phagocytic activity. The fraction of *M. brevicollis* cells with internalized 0.2 mm beads
was similar in cultures incubated with *E. coli* DH5α and PAO1 (Figure 1G). Moreover,
exposure to *P. aeruginosa* did not inhibit phagocytic uptake of *E. coli* (Figure 1H)*. These*
results suggest that exposure to *P. aeruginosa* does not broadly inhibit phagocytosis.
The above results suggested that the pathogenic effects of *P. aeruginosa* are
induced by factors secreted by extracellular bacteria. In addition, diverse secreted
bacterial molecules have been previously shown to influence choanoflagellate cell
biology[25,28,31]. Therefore, we next investigated the effects of secreted *P. aeruginosa*
molecules on *M. brevicollis* viability. Exposure of *M. brevicollis* to conditioned medium
from *P. aeruginosa* PAO1 or diverse non-pathogenic gammaproteobacteria revealed
that PAO1 conditioned medium is sufficient to restrict growth and induce cell death
(Figure 1I). Similar to live bacteria, exposure to *P. aeruginosa* conditioned medium led
to reduced motility and truncated flagella in *M. brevicollis* after approximately 8-10
hours.
Because numerous *P. aeruginosa* secreted virulence factors have been
characterized[34,40], we screened a battery of isogenic PAO1 strains with deletions in
known virulence genes to determine if any of these factors contribute to the pathogenic
effects on *M. brevicollis* (Table 2). All strains tested induced similar levels of *M.*
*brevicollis* cell death as the parental PAO1 strain, suggesting that none of the deleted
virulence genes alone are essential for inducing cytotoxicity in *M. brevicollis*. The
bioactivity in the conditioned media was also found to be heat, protease, and nuclease
resistant, indicating that the virulence factors are unlikely to be proteins or nucleic acids
(Table 3). In addition, we found that subjecting the conditioned media to methanol
extraction followed by liquid chromatography-mass spectrometry resulted in specific
fractions that recapitulated the bioactivity of the conditioned media. Although further
detailed chemical analysis is required to determine the molecular nature of these
factors, these data indicate that secreted *P. aeruginosa* small molecules are sufficient
for inducing cell death in *M. brevicollis.*

Finally, we investigated whether transient exposure to *P. aeruginosa* is sufficient
to induce cell death in *M. brevicollis* (Figure 1J). Because choanoflagellates likely
experience fluctuations in the chemical composition of aquatic microenvironments on
various timescales, we exposed stationary-phase *M. brevicollis* to *P. aeruginosa*
conditioned media for increasing durations, and then assessed survival relative to
*Flavobacterium*-treated controls after 24 hours. While *M. brevicollis* cell death was not
observed after treatment with *P. aeruginosa* conditioned media for 4 hours or less,
exposures lasting for 6 hours or longer reduced *M. brevicollis* survival (Figure 1J).
These results suggest that cell death pathways are not induced immediately in
response *P. aeruginosa* virulence factors, but are activated after longer exposures to *P.*
*aeruginosa.*

**Upregulation of *M. brevicollis* STING in response to *P. aeruginosa***

To identify potential genetic pathways activated by *M. brevicollis* in response to
pathogenic bacteria*,* we performed RNA-seq on *M. brevicollis* exposed to conditioned
medium from either *P. aeruginosa* or *Flavobacterium sp*., the non-pathogenic bacterial
strain used as a food source (Table 1).  We found that 674 genes were up-regulated
and 232 genes were downregulated two-fold or greater (FDR≤$10^{-4}$) upon *P. aeruginosa*
exposure compared to cells exposed to *Flavobacterium* (Figure 2A). The up-regulated
genes were enriched in biological processes including response to stress, endocytosis,
microtubule-based movement, mitochondrial fission, and carbohydrate metabolism.
Genes down-regulated in response to *P. aeruginosa* were enriched in biological
processes including RNA modification and metabolism (Figure 2 - figure supplement
1A). We also found that the transcription of several genes encoding proteins that
function in animal antibacterial innate immunity was upregulated in response to *P.*
*aeruginosa*, including C-type lectin, glutathione peroxidase, and STING (Figure 2A,B).
Using an antibody we raised against the C-terminal portion of *M. brevicollis* STING
(Figure 2 - figure supplement 1C) we found that STING protein levels are also elevated
in response to *P. aeruginosa* (Figure 2C). Given the importance of STING in animal
immunity and its upregulation in response to *P. aeruginosa*, we pursued its functional
relevance in the *M. brevicollis* pathogen response.

**The cyclic dinucleotide 2'3' cGAMP induces elevated expression of STING in *M.***
***brevicollis***

The predicted domain architecture of *M. brevicollis* STING consists of four
transmembrane domains followed by a cytosolic STING domain (Fig. 3A, Figure 2 -
figure supplement 1D), and likely matches the structure of the ancestral animal STING
protein. Vertebrate STING proteins contain a C-terminal tail (CTT; Fig. 3A, Figure 2 -

figure supplement 1D) that is required for the induction of interferons[41–43], and for the
activation of other downstream responses, including NFkB[44] and autophagy[45–47]
pathways. Both the STING CTT and interferons evolved in vertebrates, and it is
currently unclear how choanoflagellate and invertebrate STING proteins mediate
downstream immune responses[13,48]. However, the conservation of putative cyclic
dinucleotide-binding residues in *M. brevicollis* STING (Fig. 3B) led us to hypothesize
that STING signaling may be induced by cyclic dinucleotides. In addition, because *M.*
*brevicollis* has a cGAS-like enzyme (Figure 1- Supplement 1A), it is possible that *M.*
*brevicollis* produces an endogenous cyclic dinucleotide similar to mammalian 2'3'
cGAMP[9,12,49].
To identify potential STING inducers[49,50], we treated *M. brevicollis* with purified
cyclic dinucleotides, including mammalian cGAMP (2'3' cGAMP) and bacterial cyclic
dinucleotides (3'3' c-di-GMP, 3'3' c-di-AMP, 3'3' cGAMP). We first performed dose-
response curves to determine if the different cyclic dinucleotides affect the viability of *M.*
*brevicollis* (Fig. 3C). Interestingly, we found that exposure to 2'3' cGAMP induced cell
death in a dose-dependent manner. In contrast, exposure to 3'3' cGAMP, c-di-GMP,
and c-di-AMP did not alter *M. brevicollis* survival. Transcriptional profiling of *M.*
*brevicollis* revealed a robust transcriptional response to 2'3' cGAMP after three hours
(Figure 3 - figure supplement 1A).  Moreover, transcriptional profiling of *M. brevicollis*
exposed to 2'3' cGAMP or 3'3' cGAMP for three hours revealed that *STING* mRNA
levels increase in response to 2'3' cGAMP, but remain unchanged in response to 3'3
cGAMP (Figure 3 - figure supplement 1A-C). Therefore, we next treated *M. brevicollis*
with the cyclic dinucleotides for five hours, and measured STING protein levels by
immunoblot (Fig. 3D). Treatment with 2'3' cGAMP, but not the bacterially-produced
cyclic dinucleotides, led to elevated levels of STING protein compared to unstimulated
cells. A time course of 2'3' cGAMP treatment revealed that STING protein levels
increase as early as three hours after exposure to the cyclic dinucleotide and remain
elevated for at least 7 hours, approximately one cell cycle (Fig. 3E). While we also
observed sustained upregulation of STING in the presence of *P. aeruginosa*, this is
markedly different from what has been described in mammals, wherein STING
activation results in its translocation to lysosomes and degradation[47]. In addition,
immunostaining for STING in fixed *M. brevicollis* revealed that the number and intensity
of STING puncta increases after exposure to 2'3' cGAMP (Figure 3 - figure supplement
1E,F), although the localization of STING was difficult to assess by immunostaining due
to a lack of available subcellular markers. These data suggest that *M. brevicollis* STING
responds to 2'3' cGAMP, and that this cyclic dinucleotide can be used to further
characterize the role of STING in *M. brevicollis*.
237
**Transfection reveals that STING localizes to the *M. brevicollis* endoplasmic**
**reticulum**

A key barrier to investigating gene function in *M. brevicollis* has been the
absence of transfection and reverse genetics. We found that the transfection protocol
recently developed for the choanoflagellate *Salpingoeca rosetta*[50] was not effective in
*M. brevicollis*, but by implementing a number of alterations to optimize reagents and
conditions (see Methods) we were able to achieve both reproducible transfection and
establishment of stably transformed cell lines in *M. brevicollis*.

To investigate the subcellular localization of STING, we established a robust
transfection protocol for *M. brevicollis* that would allow the expression of fluorescently-
labeled STING along with fluorescent subcellular markers for different organelles.
We observed that STING-mTFP protein localized to tubule-like structures around the
nucleus (Figure 5A) similar to what was observed by immunostaining with an antibody
to STING (Figure 3 - figure supplement 1E,F). We then co-transfected STING-mTFP
alongside fluorescent reporters marking the endoplasmic reticulum (ER) or mitochondria
(Figure 5B,C) and performed live-cell imaging. STING-mTFP co-localized with a
fluorescent marker highlighting the ER (Figure 5B). Thus, as in mammalian cells[15,51],
STING localizes to regions of the ER in *M. brevicollis*.

**Genetic disruption of STING reveals its role in responding to 2'3' cGAMP and *P.***
***aeruginosa***

Disrupting the *STING* locus using CRISPR/Cas9-mediated genome editing
(Figure 5A) enabled us to investigate the function of STING.  To overcome low gene
editing efficiencies in *M. brevicollis,* we based our gene editing strategy on a protocol
recently developed for *S. rosetta* that simultaneously edits a gene of interest and
confers cycloheximide resistance[52]. By selecting for cycloheximide resistance and then
performing clonal isolation, we were able to isolate a clonal cell line that has a deletion
within the *STING* locus that introduces premature stop codons (Figure 5 - figure
supplement 1A). We were unable to detect STING protein in *STING*⁻ cells by
immunoblot (Figure 5B). Wild type and *STING*⁻ cells have similar growth kinetics (Figure
5 - figure supplement 1B), suggesting that STING is not required for cell viability under
standard laboratory conditions. In addition, overexpression of STING-mTFP did not
affect *M. brevicollis* viability.

To investigate the connection between 2'3' cGAMP and STING signaling in *M.*
*brevicollis*, we exposed *STING*⁻ cells to increasing concentrations of 2'3' cGAMP. In
contrast to wild type *M. brevicollis*, *STING*⁻ cells are resistant to 2'3' cGAMP-induced
cell death (Figure 5C). The 2'3' cGAMP resistance phenotype could be partially
reversed by stably expressing STING within the *STING*⁻ mutant background (Figure
5D). In addition, *STING*⁻ cells fail to induce a strong transcriptional response to 2'3'
cGAMP compared to wild type cells (Figure 5E, Figure 3 - figure supplement 1A, Figure

5 - figure supplement 1C,D). While 371 genes are differentially expressed in wild type
cells after exposure to 2'3' cGAMP for three hours, only 28 genes are differentially
expressed in *STING*⁻ cells (FC ≥3; FDR≤$10^{-4}$). Thus, 2'3' cGAMP induces a STING-
dependent transcriptional response in *M. brevicollis*.

Interestingly, of the 22  choanoflagellate species with sequenced
transcriptomes[3,29], only *M. brevicollis* and *Salpingoeca macrocollata*, express homologs
of both STING and cGAS (Figure 1- Supplement 1A, Figure 5F, Figure 5 - figure
supplement 1E). Therefore, we were curious whether other choanoflagellate species
are able to respond to 2'3' cGAMP in the absence of a putative STING protein. We
exposed four other choanoflagellate species (*Salpingoeca infusionum*, *S. macrocollata,*
*S. rosetta*, and *Salpingoeca punica*) to increasing 2'3' cGAMP concentrations*,* and
quantified survival after 24 hours (Figure 5G). Of these additional species, only *S.*
*macrocollata* had impaired survival in the presence of 2'3' cGAMP. Thus, it is possible
that STING also responds to 2'3' cGAMP in *S. macrocollata.*

We next asked whether *STING*⁻ cells have altered responses to other immune
agonists. Although *M. brevicollis* is continuously co-cultured with feeding bacteria, we
observed that treatment with high concentrations of *E. coli* lipopolysaccharides induces
cell death (Figure 5H). As LPS is not known to activate STING signaling, we treated wild
type and *STING*⁻ cells with LPS to probe the specificity of STING-mediated immune
responses in *M. brevicollis*.  The survival responses of wild type and *STING*⁻ cells to
LPS were indistinguishable (Figure 5H), suggesting that there are separable pathways
for responding to 2'3' cGAMP and LPS. We also examined the survival of *STING*⁻ cells
exposed to *P. aeruginosa* conditioned medium (Figure 5I,J). In growth curve
experiments, *P. aeruginosa* hindered the growth rate and stationary phase cell density
of *STING*⁻ cells compared to *Flavobacterium* (Figure 5I,J)*.* However, *STING*⁻ cells were
still able to divide in the presence of *P. aeruginosa,* whereas wild type cell growth was
completely restricted (Figure 5I,J). These results indicate that wild type cells are more
susceptible to *P. aeruginosa* than *STING*⁻ cells, although it is unclear how STING
contributes to *P. aeruginosa*-induced growth restriction and cell death.

309
310

**2'3' cGAMP-induces autophagic signaling via STING**

One downstream consequence of STING signaling in animals is the initiation of
autophagy[21,45,46,49,53]. Based on viral infection studies in *D. melanogaster*[53] and
experiments expressing invertebrate STING in mammalian cells[45],  it has been
proposed that the induction of autophagy may be an interferon-independent ancestral
function of STING.  Although *M. brevicollis* lacks many effectors required for immune
responses downstream of STING in animals (including TBK1 and NF-kB; Figure S1A),

autophagy machinery is well conserved in *M. brevicollis*. Therefore, we asked if one
outcome of 2'3' cGAMP exposure in *M. brevicollis* is the induction of autophagy.
The evolutionarily conserved protein Atg8/LC3 is a ubiquitin-like protein that can
be used to monitor autophagy[54,55]. During autophagosome formation, unmodified Atg8,
called Atg8-I, is conjugated to phosphatidylethanolamine. Lipidated Atg8, called Atg8-II,
remains associated with growing autophagosomes. As such, two indicators of
autophagy are elevated Atg8-II levels relative to Atg8-I and increased formation of
Atg8+ autophagosome puncta. Because antibodies are not available to detect
endogenous *M. brevicollis* autophagy markers or cargo receptors, we generated wild
type and *STING*[−] cell lines stably expressing mCherry-Atg8. Stable expression of
mCherry-Atg8 under the control of the constitutive pEFL promoter did not alter the
relative susceptibilities of these cell lines to 2'3'cGAMP (Figure 6 - figure supplement
1A). By immunoblot, mCherry-Atg8-II can be distinguished from mCherry-Atg8-I based
on its enhanced gel mobility. When we exposed both cell lines to 2'3' cGAMP for three
hours, we observed increased levels of Atg8-II relative to Atg8-I by immunoblot in wild
type, but not *STING*[−] cells (Figure 6A). These results suggest that treatment with 2'3'
cGAMP induces autophagic signaling in a STING-dependent manner; however, making
this conclusion requires evidence of autophagy induction through inhibitor studies. To
confirm autophagy induction, we treated cells with chloroquine, a lysosomotropic agent
which inhibits autophagy by blocking endosomal acidification, thereby preventing
amphisome formation and Atg8-II turnover[54]. Exposing wild type cells pretreated with
chloroquine to 2'3' cGAMP for three hours resulted in increased levels of Atg8-II relative
to Atg8-I, suggesting that 2'3' cGAMP treatment indeed induces the autophagic
pathway (Figure 6B, Figure 6 - figure supplement 1B). In cells pretreated with
chloroquine, STING levels did not markedly increase after exposure to 2'3' cGAMP
(Figure 6 - figure supplement 1B), raising the possibility that the autophagic pathway is
important for regulating STING protein levels. We next examined whether 2'3' cGAMP
induces Atg8+ puncta formation by treating wild type and *STING*[−] cells with 2'3' cGAMP
for three hours, and observing mCherry foci by microscopy (Figure 6C-F). Quantifying
images revealed that Atg8+ puncta accumulate after 2'3' cGAMP treatment in wild type,
but not *STING*[−] cells (Figure 6G). Overall, these results suggest that *M. brevicollis*
responds to 2'3' cGAMP through STING-dependent induction of the autophagy
pathway.
Finally, we asked whether STING-mediated autophagic pathway induction affects
survival after exposure to 2'3' cGAMP (Figure 6H). To determine if inhibiting autophagy
impacts 2'3' cGAMP-induced cell death, we examined the survival responses of wild
type and *STING*[−] *M. brevicollis* to 2'3' cGAMP after pretreatment with lysosomotropic
agents chloroquine or NH$_4$Cl. Chloroquine or NH$_4$Cl pretreatment rescued 2'3' cGAMP-
induced cell death in wild type *M. brevicollis* (Figure 6H), whereas the survival of
*STING*[−] cells, which are already resistant to 2'3' cGAMP-induced cell death, was not

affected. Therefore, we hypothesize that 2'3' cGAMP induces cell death in *M. brevicollis*
by overstimulating STING-mediated autophagic signaling.
361
362    **Discussion**
363
Investigating choanoflagellate immune responses has the potential to inform the
ancestry of animal immune pathways. In this study, we screened diverse bacteria to
identify a choanoflagellate pathogen, and determined that *M. brevicollis* is killed by
sustained exposure to *P. aeruginosa* bacteria. We found that STING, a crucial
component of animal innate responses to cytosolic DNA, is upregulated in *M. brevicollis*
after exposure to *P. aeruginosa* or the STING ligand 2'3' cGAMP. The application of
newly-developed transgenic and genetic tools for *M. brevicollis* revealed that, similar to
mammalian STING, *M. brevicollis* STING localizes to perinuclear endoplasmic reticulum
regions. In addition, STING mediates responses to *P. aeruginosa* bacteria, and is
required for inducing transcriptional changes and autophagic signaling in response to
2'3' cGAMP. These data reveal that STING plays conserved roles in choanoflagellate
immune responses, and provide insight into the evolution of STING signaling on the
animal stem lineage.
The discovery that *M. brevicollis* STING mediates immune responses raises a
number of interesting questions about the full extent of its physiological roles in
choanoflagellates. For example, while our results demonstrate that *M. brevicollis* STING
responds to exogenous 2'3' cGAMP, the endogenous triggers of STING activation in *M.*
*brevicollis* remain to be determined. *M. brevicollis* has a putative cGAS homolog,
suggesting that STING may respond to an endogenously produced cyclic dinucleotide
similar to 2'3' cGAMP. Determining the enzymatic activities of *M. brevicollis* cGAS and
identifying the endogenous trigger of *M. brevicollis* STING will be critical steps towards
elucidating mechanisms of STING activation. Although cGAS and STING are rare
among sequenced choanoflagellate species, both species with STING homologs, *M.*
*brevicollis* and *S. macrocollata*, also harbor a cGAS homolog (Figure 1- Supplement
1A), suggesting the presence of an intact choanoflagellate cGAS-STING pathway.
Our results suggest that *M. brevicollis* has distinct responses to 2'3' cGAMP
versus 3'3'-linked cyclic dinucleotides produced by bacteria (Figure 3C,D, Figure 3 -
figure supplement 1). In contrast to 2'3' cGAMP, bacterial cyclic dinucleotides (3'3'
cGAMP, c-di-AMP, c-di-GMP) do not induce cell death in *M. brevicollis*. However, 3'3'
cGAMP induces a robust transcriptional response in *M. brevicollis,* indicating that
STING, or a different cyclic dinucleotide receptor[57], responds to these bacterial
molecules. One hypothesis is that *M. brevicollis* STING, similar to animal STING
proteins[14], may have different binding affinities for 2'3' and 3'3'-linked cyclic
dinucleotides. It is also possible that bacterial cyclic dinucleotides activate additional
pathways that influence survival in *M. brevicollis*. As bacterivores, choanoflagellates

likely benefit from a fine-tuned response to bacterial cyclic dinucleotides that enables
them to interpret higher and lower concentrations in their environment. Elucidating
mechanisms of STING activation in *M. brevicollis* could help reveal how STING proteins
in animals evolved to respond to both bacterially-produced and endogenous cyclic
dinucleotides.
While it is clear that 2'3' cGAMP stimulates STING-dependent transcriptional
responses and autophagic signaling in *M. brevicollis* (Figure 5, Figure 6), the signaling
pathways downstream of STING in choanoflagellates are unknown. Much of what is
known about STING signaling comes from mammals and involves the extended CTT
domain of STING, which *M. brevicollis* lacks, and immune genes that are restricted to
vertebrates. Two pathways downstream of STING activation that are conserved in
invertebrates, and as such are proposed ancestral functions of STING, are autophagy
and NF-kB signaling. Here, we observed that STING is required for induction of the
autophagy pathway in response to 2'3' cGAMP in *M. brevicollis*, indicating that the role
of STING in regulating autophagy predates animal origins. While exposure to 2'3'
cGAMP leads to NF-kB activation in the sea anemone *N. vectensis*[58] and in insects[59–62],
neither *M. brevicollis* nor *S. macrocollata*, the two choanoflagellate species with STING,
possess a NF-kB homolog[3].  Nonetheless, 2'3' cGAMP activates an extensive
transcriptional response downstream of STING in *M. brevicollis*, although the specific
signaling pathways remain to be identified. This does not negate the hypothesis that
STING signaling led to NF-kB activation in the Urmetazoan, but strongly suggests that
additional pathways exist downstream of STING activation in choanoflagellates, and
potentially in animals.
Choanoflagellates forage on diverse environmental bacteria for sustenance, yet
how they recognize and respond to pathogens is a mystery. Our finding that *P.*
*aeruginosa* has pathogenic effects on *M. brevicollis* (Figure 1) provides a much-needed
framework for uncovering mechanisms of pathogen recognition and antibacterial
immunity in choanoflagellates. Profiling the host transcriptional response to *P.*
*aeruginosa* has allowed us to identify choanoflagellate genes that may be involved in
recognizing (C-type lectins) and combating (polysaccharide lyases, antimicrobial
peptides) bacteria; yet, it has also revealed the immense complexity of this interaction,
with more than 600 *M. brevicollis* genes differentially expressed in response *P.*
*aeruginosa*. Thus, identifying specific *P. aeruginosa* virulence factors will be critical for
understanding why *P. aeruginosa* – but not other bacteria – have pathogenic effects on
*M. brevicollis,* and facilitate the characterization of mechanisms underlying
choanoflagellate pathogen responses.
With the establishment of molecular genetic techniques in choanoflagellates --
first for *S. rosetta*[50,52], and here for *M. brevicollis* -- we now have the opportunity to
explore the functions of candidate immune genes. Identifying additional
choanoflagellate pathogens, particularly viral pathogens, will also be key to delineating

immune response pathways. Finally, as choanoflagellates are at least as genetically
diverse as animals[3],  expanding studies of immune responses to diverse
choanoflagellate species will be essential for reconstructing the evolution of immune
pathways in animals.
443
444

445 <u>Acknowledgements</u>
446

We thank Bill Jackson, Tera Levin, Shally Margolis, Russell Vance, and Nan Yan
for helpful advice and/or comments on the manuscript, and David Greenberg, David
Hendrixson, Andrew Koh, Kim Orth, Russell Vance, and Sebastian Winter for bacterial
strains. We thank Neal Alto for use of the widefield microscope, Monika Sigg for use of
a reporter construct, and Mya Breitbart for sea water samples. We are grateful for David
Booth's invaluable advice on developing transfection and gene editing protocols.
This work was funded by a Pew Innovation Fund award (JKP and NK), a HHMI
Hanna Gray Fellows award (AW), a HHMI Faculty Scholar award (JKP), and a
Burroughs Wellcome Fund Investigators in the Pathogenesis of Infectious Diseases
(JKP). We acknowledge the assistance of the UT Southwestern Live Cell Imaging
Facility, a Shared Resource of the Harold C. Simmons Cancer Center, supported in part
by an NCI Cancer Center Support Grant, 1P30 CA142543-01.

459
460

461 <u>Materials and Methods</u>
462
463

**Key Resources Table**

| Reagent type (species) or resource | Designation | Source or reference | Identifiers | Additional information |
|---|---|---|---|---|
| Strain, strain background (*M. brevicollis*) | *M. brevicollis* | ATCC PRA-258 | PMID: 18276888 | |
| Genetic reagent, (*M. brevicollis*) | *M. brevicollis STING⁻* | This study | | *STING⁻* knockout strain ; cell line maintained |

| | | | | |
|---|---|---|---|---|
| | | | | by A. Woznica |
| Transfected construct (*M. brevicollis*) | pEFL5'-pac-P2A-STING-mTFP-3'act | This study | | Construct to express *M, brevicollis* STING fused to mTFP; can be obtained from A. Woznica |
| Transfected construct (*M. brevicollis*) | pEFL5'-pac-P2A-mCherry-Atg8--3'act | This study | | Construct to express mCherry fused to *M. brevicollis* Atg8; can be obtained from A. Woznica |
| Strain, strain background(*Flavobacterium*) | *Flavobacterium* sp. | This study | | Isolated from MX1 (ATCC PRA-258) culture; can be obtained from A. Woznica |
| Strain, strain background(*Pseudomonas aeruginosa*) | PAO1 | ATCC 15692 | PMID: 13961373 | |
| *P. aeruginosa*, transgenic strain | PAO1-GFP | ATCC 15692GFP | PMID: 9361441 | |
| Antibody | anti-choano STING (rabbit polyclonal) | This study | | Generated by Pacific Immunology; dilution 1:200 for IF, 1:2000 dilution for |

| | | | | WB; can be obtained from A. Woznica |
|---|---|---|---|---|
| antibody | Anti-mCherry 16D7 (rat monoclonal) | Invitrogen | Cat# M11217 | 1:2000 dilution for WB |
| antibody | Anti-human Tubulin E7 (Mouse monoclonal) | Developmental Studies Hybridoma Bank | Cat# AB_2315513 | 1:200 dilution for IF |
| antibody | Alpha-human tubulin (Mouse monoclonal) | Sigma Aldrich | Cat # T64074 | 1:7000 dilution for WB |
| Chemical compound, drug | 2'3' cGAMP | Cayman Chemical | Cat# 19887 | |
| Chemical compound, drug | 3'3' cGAMP | Cayman chemical | Cat# 17966 | |
| Sequence-based reagent | *STING556* gRNA | This study | Guide RNA | TTTCGGGATT CAGATGTGG G |
| Sequenced-based reagent | *STING* locus PCR primers | This study | PCR primers | F: 5' ATG ATG GTT AAT CTC TCT GAT CTT TCA CAT C 3' R: 5' TTA TGG CAT CGC ATA CTG GTC C 3' |

| Commercial assay, kit | SG Cell Line 4D-NucleofectorT M X Kit S | Lonza, | Cat# V4XC-3032 | |
|---|---|---|---|---|

## Culturing choanoflagellates

All strains of *M. brevicollis* were co-cultured with *Flavobacterium sp.* bacteria[4]
(American Type Culture Collection [ATCC], Manassas, VA; Cat. No. PRA-258) in a
seawater based media enriched with glycerol, yeast extract, peptone and cereal grass
(details in Media Recipes). Cells were grown either at room temperature, or at 16ºC in a
wine cooler (Koldfront). All *M. brevicollis* cell lines were verified by 18S sequencing and
RNA-seq. Choanoflagellate cell lines *S. rosetta*, *S. macrocollata*, *S. punica*, and *S.*
*infusionum* were verified by 18S sequencing.

## Bacterial effects on *M. brevicollis*

### *Isolating environmental bacteria*

Environmental bacterial species were isolated from water samples from Woods
Hole, MA, St. Petersburg, FL, and Dallas, TX. Water samples were streaked onto Sea
Water Complete media or LB plates, and grown at 30º C or 37º C. After isolating
individual colonies, partial 16S sequencing using 16S universal primers (27F: 5′-
AGAGTTTGATCCTGGCTCAG-3′, 1492R: 5′-TACGGYTACCTTGTTACGACTT-3′) was
used to determine the identity of the bacterial isolates.

### *Screening for pathogenic effects*

*M. brevicollis* was grown for 30 h, and feeding bacteria were reduced through
one round of centrifugation and resuspension in artificial seawater (ASW). Cells were
counted on a hemocytometer and diluted to $5 \times 10^6$ cells/mL in High Nutrient Medium,
and plated into 24-well plates.

For each bacterium, a single colony was inoculated into LB and grown shaking
overnight at either 30º C (environmental isolates) or 37º C (mouse isolates). Bacterial
cells were pelleted by centrifugation for 5 minutes at 4000 x *g,* and resuspended in
artificial seawater (ASW) to an OD~1.

Each bacterial species was added to *M. brevicollis* culture at two concentrations
(10mL/mL and 50 mL/mL) in duplicate. *M. brevicollis* was then monitored at regular
intervals for changes in behavior and growth.

### *Growth curves in the presence of bacteria*

All bacteria were grown shaking at 30º C in Sea Water Complete media or LB (to
optical density of 0.8). For each bacterial strain, CFU plating was used to estimate the
number of bacterial cells/ mL under these growth conditions. To prepare bacterial
conditioned media, bacterial cells were pelleted by centrifugation for 10 minutes at 4000
x *g,* and supernatant was passed through a 0.22mm sterilizing filter.

*M. brevicollis* was grown for 30 h, and bacteria were washed away through two
consecutive rounds of centrifugation and resuspension in artificial seawater (ASW).
Cells were counted on a hemocytometer and diluted to $1.0 \times 10^6$ cells/mL (growth curves
with live bacteria) or $1.5 \times 10^5$ cells/mL (growth curves with conditioned medium) in High
Nutrient Medium. To test the effects of live bacteria, $1.5 \times 10^6$ bacterial cells were added
per 1 mL of *M. brevicollis* culture. To test the effects of bacterial conditioned media, 50
ml of bacterial conditioned media was added per 1 mL of *M. brevicollis* culture. For each
growth curve biological replicate, cells were plated into 24-well plates, and two wells
were counted per time point as technical replicates. At least three biological replicates
are represented in each graph.

***Bacterial internalization***
Fluorescent *E. coli* and *P. aeruginosa* were grown shaking at 30º C in LB to an
optical density of $OD_{600}=0.8$. Fluorescent *C. jejuni* was grown from freezer stocks in
microaerobic conditions on Mueller-Hinton agar. For each bacterial strain, CFU plating
was used to estimate the number of bacterial cells/ mL under these growth conditions.
*M. brevicollis* was grown for 30 h, and feeding bacteria were washed away with
one round of centrifugation and resuspension in artificial seawater (ASW). Cells were
counted and diluted to $1.5 \times 10^5$ cells/mL in ASW. $1.5 \times 10^7$ bacterial cells were added to 2
mL *M. brevicollis* culture (MOI=50), and co-incubated at room temperature for 1 hr with
gentle mixing at regular intervals to avoid settling. To quantify bead internalization, *M.*
*brevicollis* was co-incubated with bacteria for 1 hour (as described above), at which
point ~$1 \times 10^{10}$ beads (0.2mm diameter, resuspended in 1% BSA to prevent clumping)
were added to the conical for an additional 30 minutes.
Prior to imaging, 200mL aliquots were transferred to 8-well glass bottom
chambers (Ibidi Cat. No 80827). Live imaging was performed on a Zeiss Axio Observer
widefield microscope using a 63x objective. Images were processed and analyzed using
Fiji[63].

***P. aeruginosa deletion mutants***
*P. aeruginosa* deletion strains were acquired from the Seattle PAO1 transposon
mutant library (NIH P30 DK089507). Strains RP436 and RP576 (PMID 15731071) were
acquired from Russell Vance, The effects of both live bacteria and bacterial conditioned
medium were tested for all acquired strains at a range of PFU/mL (live bacteria) or
percent volume (conditioned medium).
534

<u>Immune agonist dose-response curves</u>
*M. brevicollis* was grown to late-log phase, and feeding bacteria were reduced
through one round of centrifugation and resuspension in artificial seawater (ASW). Cells
were counted on a hemocytometer and diluted to $1.0 \times 10^6$ cells/mL (growth curves with
live bacteria) in High Nutrient Medium, and aliquoted into 96-well (100µL/well) or 24-well
(1mL/well) plates. Immune agonists were added at indicated concentrations in technical
duplicate, and cells were counted again after 24 hours. % survival is a calculation of:

542 [mean experimental (cells/mL) / mean control (cells/mL)]. Each dose-response curve is
543 representative of at least three biological replicates.
544
545 <u>RNA-seq</u>
546 ***Growth of choanoflagellate cultures***
547   After thawing new cultures, growth curves were conducted to determine the
548 seeding density and time required to harvest cells at late-log phase growth. To grow
549 large numbers of cells for RNA-seq, cells were seeded one to two days prior to the
550 experiment in either 3-layer flasks (Falcon; Corning, Oneonta, NY, USA; Cat. No. 14-
551 826-95) or 75 cm$^2$ flasks (Falcon; Corning, Oneonta, NY, USA; Cat. No. 13-680-65),
552 and grown at room temperature. Bacteria were washed away from choanoflagellate
553 cells through two rounds of centrifugation and resuspension in artificial seawater (ASW).
554 To count the cell density, cells were diluted 100-fold in 200 µl of ASW, and fixed with 1
555 µL of 16% paraformaldehyde. Cells were counted on a hemocytometer, and the
556 remaining cells were diluted to a final concentration of $4 \times 10^6$ choanoflagellate cells/mL.
557 The resuspended cells were divided into 2.5 mL aliquots and plated in 6-well plates
558 prior to treatment. After treatment, cells were transferred to a 15 mL conical and
559 pelleted by centrifugation at 2400 x *g* for 5 min, flash frozen with liquid nitrogen, and
560 stored at -80°C.
561 ***RNA isolation***
562   Total RNA was isolated from cell pellets with the RNAqueous kit (Ambion,
563 Thermo Fisher Scientific). Double the amount of lysis buffer was used to increase RNA
564 yield and decrease degradation, and RNA was eluted in minimal volumes in each of the
565 two elution steps (40 µL and 15 µL). RNA was precipitated in LiCl to remove
566 contaminating genomic DNA. Total RNA concentration and quality was evaluated using
567 the Agilent Bioanalyzer 2100 system and RNA Nano Chip kit (Cat No. 5067-1511).
568 ***Library preparation, sequencing, and analysis***
569   Libraries were prepared and sequenced by the UTSW Genomics Sequencing
570 Core. RNA libraries were generated with the Illumina TruSeq® Stranded mRNA Library
571 prep kit (Cat No. 20020594), using a starting total RNA input of 2-3 µg. To remove
572 contaminating bacterial RNA, samples were first poly-A selected using oligo-dT
573 attached magnetic beads. Following purification, the mRNA was fragmented at 94°C for
574 4 minutes, and cleaved RNA fragments were synthesized into cDNA. After an end
575 repair step, UMI adapters (synthesized by IDT) were ligated to the cDNA, and the
576 products were twice purified using AMPure XP beads before amplification.
577 Library quantity was measured using the Quant-iT™ PicoGreen dsDNA Assay kit by
578 Invitrogen (Cat No. P7589) and a PerkinElmer Victor X3, 2030 Multilabel Reader.
579 Library quality was verified on an Agilent 2100 Bioanalyzer instrument using Agilent
580 High sensitivity DNA kit (Cat No. 5067-4626) or DNA 1000 kit (Cat No. 5067-
581 1504). Libraries were pooled, and sequenced in different batches on either the Illumina

NextSeq 550 system with SE-75 workflow, or the Illumina NovaSeq 6000 system with
S4 flowcell and XP PE-100 workflow, generating 25-40 million reads per sample. Reads
were checked for quality using fastqc (v0.11.2) and fastq_screen (v0.4.4), and trimmed
using fastq-mcf (ea-utils, v1.1.2-806). Trimmed fastq files were mapped to the *Monosiga*
*brevicollis* reference genome (NCBI:txid81824) using TopHat[64] (v2.0.12). Duplicates
were marked using picard-tools (v2.10.10).  Read counts were generated using
featureCounts[65], and differential expression analysis was performed using edgeR[66].
Statistical cutoffs of FDR≤$10^{-4}$ were used to identify significant differentially expressed
genes. GO enrichment analysis of differentially expressed genes was performed using
DAVID (https://david.ncifcrf.gov/).
592
RT-qPCR
RNA was isolated as described above, and cDNA was synthesized from total
RNA with the High Capacity cDNA Reverse Transcription kit (Applied Biosystems;
Thermo Fisher. Real-time PCR was performed with iTaq Universal SYBR Green
Supermix (Biorad; Cat No. 1725121) or SYBR Green PCR master mix (Applied
Biosystems; Cat No. 4309155) in either an 7500 Fast Real-Time PCR System (Applied
Biosystems), or a QuantStudio 3 Real-Time PCR System (Applied Biosystems). Ct
values were converted into relative gene expression using the ΔΔCt method (Livak and
Schmittgen, 2001) and the internal control gene RPL15 (MONBRDRAFT_38309).
602
Immunoblotting
*M. brevicollis* was harvested by centrifugation at 5,000 x *g* for 5 min at 4°C, and
resuspended in 100 µL lysis buffer (50mM Tris, pH 7.4, 150 mM NaCl, 1 mM EDTA, 1
mM ethyleneglycoltetraacetic acid [EGTA], 0.5% sodium deoxycholate, 1% NP-40)
containing protease inhibitor cocktail (Roche) for 10 min at 4°C. The crude lysate was
clarified by centrifugation at 10,000 x *g* for 10 min at 4°C, and denatured in Laemmli
buffer before SDS-PAGE. Proteins were transferred to an Immobilon-P PVDF
membrane (Millipore), and blocked for two hours in PBST (1x PBS containing 5% nonfat
dry milk and 0.05% Tween-20). Membranes were incubated with primary antibodies
diluted in PBST overnight at 4°C and washed extensively in PBST. Membranes were
incubated with secondary antibodies for 1 hour at room temperature, washed
extensively in PBST, and developed using Immobilon Western Chemiluminescent HRP
Substrate (Millipore Sigma). Source data files for all western blots are provided as
Source Data File 1.
***STING antibody production***
The anti-mbreSTING antibody was generated by Pacific Immunology. Rabbits
were immunized with a KLH-conjugated peptide corresponding to residues 320-338 of
*M. brevicollis* protein EDQ90889.1 (Cys-KNRSEVLKKMRAEDQYAMP), and serum was
affinity purified against the peptide to reduce cross-reactivity and validated using
immunoblotting.

623
## Immunofluorescence Staining and Imaging

Depending on the cell density of the starting culture, between 0.2-1 mL of cells were concentrated by centrifugation for 5 min at 2500 × *g*. The cells were resuspended in 200 µl of artificial seawater and applied to poly-L-lysine–coated coverslips (Corning Life Sciences; Cat. No.354085) placed at the bottom of each well of a 24-well cell culture dish. After the cells were allowed to settle on the coverslip for 30 min, 150 µl of the cell solution was gently removed from the side of the dish. All of the subsequent washes and incubations during the staining procedure were performed by adding and removing 200 µl of the indicated buffer.

Cells were fixed in two stages. First, 200 µl cold 6% acetone diluted in 4X PBS was added for 5 min at room temperature. Next, 200 µl cold 8% paraformaldehyde diluted in 4X PBS was added (yielding a final concentration of 4% paraformaldehyde), and the fixative mixture was incubated for 15 min at room temperature. After fixation, the coverslip was gently washed three times with 200 µl 4X PBS.

Cells were permeabilized by incubating in permeabilization buffer (4X PBS; 3% [wt/vol] bovine serum albumin (BSA)-fraction V; 0.2% [vol/vol] Triton X-100) for 30 min. After removing permeabilization buffer, the coverslip was incubated in primary antibody for 1 hour at room temperature, and then washed three times in 4X PBS. The coverslip was then incubated with secondary antibody for 1 hour at room temperature, and then washed twice in 4X PBS. The coverslip was next incubated in 4 U/ml Phalloidin (Thermo Fisher Scientific) for 30 min at room temperature, washed once in 4X PBS. Lastly, the coverslip was incubated in 10 µg/ml Hoechst 33342 (Thermo Fisher Scientific) for 5 min at room temperature, and then washed once with 4X PBS.

To prepare a slide for mounting, 10 µl of Pro-Long Gold (Thermo Fisher Scientific) was added to a slide. The coverslip was gently removed from the well with forceps, excess buffer was blotted from the side with a piece of filter paper, and the coverslip was gently placed on the drop of Pro-Long diamond. The mounting media cured overnight before visualization.

Images were acquired on either: (1) a Zeiss LSM 880 Airyscan confocal microscope with a 63x objective by frame scanning in the superresolution mode (images processed using the automated Airyscan algorithm (Zeiss)), or (2) a Nikon CSU-W1 SoRa spinning disk confocal microscope with a 60x objective in SR mode (images processed using Imaris).

## Live-Cell Imaging

Cells transfected with fluorescent reporter plasmid were prepared for microscopy by transferring 200 µl of cells to a glass-bottom dish or glass-bottom 8-well chamber (Ibidi). Confocal microscopy was performed on a Zeiss Axio Observer LSM 880 with an Fast Airyscan detector and a 63x/NA1.40 Plan-Apochromatic oil immersion objective

(Carl Zeiss AG, Oberkochen, Germany). Confocal stacks were acquired by frame

scanning in superresolution mode, and images were processed using the automated

Airyscan algorithm (Zeiss).

Transfection of *M. brevicollis*

**Cell Culture.** One day prior to transfection, 60 ml of High Nutrient Medium was

inoculated with *M. brevicollis* to a final concentration of 10000 cells/ml. The culture was

split in two, and grown in two 75 cm$^2$ flasks at room temperature, approximately 22°C

(Falcon; Corning, Oneonta, NY, USA; Cat. No. 13-680-65).

**Cell Washing.** After 24 hours of growth, bacteria were washed away from *M.*

*brevicollis* cells through three consecutive rounds of centrifugation and resuspension in

artificial seawater (ASW). The culture flasks were combined and vigorously shaken for

30 s, and then transferred to 50-ml conical tubes and spun for 5 min at 2000 × *g* and

22°C. The supernatant was removed with a serological pipette, and residual media were

removed with a fine-tip transfer pipette. The cell pellets were resuspended in a single

conical tube in a total volume of 50 ml of ASW, vigorously shaken for 30 s, and then

centrifuged for 5 min at 2050 × *g*. The supernatant was removed as before. In a final

washing step, the cell pellet was resuspended in 50 mL ASW, shaken vigorously, and

centrifuged for 5 min at 2100 × *g*. After the supernatant was removed, the cells were

resuspended in a total volume of 400 µl of ASW. To count the cell density, cells were

diluted 100-fold in 200 µl of ASW, and fixed with 1 µl  of 16% paraformaldehyde. Cells

were counted on a hemocytometer, and the remaining cells were diluted to a final

concentration of 5 × 10$^7$ choanoflagellate cells/ml. The resuspended cells were divided

into 100-µl aliquots with 5 × 10$^6$ cells per aliquot to immediately prime cells in the next

step.

**Cell Priming.** Each aliquot of *M. brevicollis* cells was incubated in priming buffer (40

mM HEPES-KOH, pH 7.5; 55 mM lithium citrate; 50 mM L-cysteine; 10% [wt/vol] PEG

8000; and 2 µM papain) to remove the extracellular material coating the cell. The 100-µl

aliquots, which contained 5 × 10$^6$ cells, were centrifuged for 5 min at 1700 × *g*. The

supernatant was removed, and cells were resuspended in 100 µl of priming buffer and

then incubated for 35 min at room temperature. Priming was quenched by adding 4 µl of

50-mg/ml bovine serum albumin-fraction V (Thermo Fisher Scientific, Waltham, MA;

Cat. No. BP1600-100) and then centrifuged for 5 min at 1250 ×*g* and 22°C with the

centrifuge brake set to a "soft" setting. The supernatant was removed with a fine-tip

micropipette, and the cells were resuspended in 25 µl of SG Buffer (Lonza).

**Nucleofection.** Each transfection reaction was prepared by adding 2 µl of "primed"

cells resuspended in SG buffer (Lonza) to a mixture of: 16 µl of SG buffer, 2 µl of 20

µg/µl pUC19, 1 µl of 250 mM ATP (pH 7.5), 1 µl of 100 mg/ml sodium heparin, and ≤7 µl

of reporter DNA (volume is dependent on the number of constructs transfected). Each

transfection reaction was transferred to one well in 16-well nucleofection strip (Lonza;

Cat. No. V4XC-2032). The nucleofection strip was placed in the X-unit (Lonza; Cat. No.
AAF-1002F) connected to a Nucleofector 4D core unit (Lonza; Cat. No. AAF-1002B),
and the EO100 pulse was applied to each well.
*Recovery.* 100 µl of cold recovery buffer (10 mM HEPES-KOH, pH 7.5; 0.9 M
sorbitol; 8% [wt/vol] PEG 8000) was added to the cells immediately after pulsation. After
5 minutes, the whole volume of the transfection reaction plus the recovery buffer was
transferred to 2 ml of Low Nutrient Medium in a 12-well plate. The cells were grown for
24–48 hours before being assayed for luminescence or fluorescence.
*Puromycin Selection.* To generate stably transfected *M. brevicollis* cell lines,
puromycin was added to cells 24 hours after transfection at a final concentration of 300
µg/mL. Cells were monitored over the course of 7-21 days, and fresh High Nutrient
Media + 300 µg/mL puromycin was added to the cells as needed.
715
Genome editing
For a more detailed description of gRNA and repair oligonucleotide design, refer to
Booth et al. 2018[50].
*Design and preparation of gRNAs* First, crRNAs were designed by using the
extended recognition motif 5'-HNNGRSGGH-3' (in which the PAM is underlined, N
stands for any base, R stands for purine, S stands for G or C, and H stands for any
base except G) to search for targets in cDNA sequences[67]. Next, we confirmed that the
RNA sequence did not span exon-exon junctions by aligning the sequence to genomic
DNA.
Functional gRNAs were prepared by annealing synthetic crRNA with a synthetic
tracrRNA[52]. To prepare a functional gRNA complex from synthetic RNAs, crRNA and
tracrRNA (Integrated DNA Technologies [IDT], Coralville, IA, USA) were resuspended
to a final concentration of 200 µM in duplex buffer (30 mM HEPES-KOH, pH 7.5; 100
mM potassium acetate; IDT, Cat. No. 11-0103-01). Equal volumes of crRNA and
tracrRNA stocks were mixed together, incubated at 95°C for 5 min in an aluminum
block, and then the entire aluminum block was placed at room temp to slowly cool the
RNA to 25°C. The RNA was stored at -20°C
*Design and preparation of repair oligonucleotides* Repair oligonucleotides for
generating knockouts were designed by copying the sequence 50 bases upstream and
downstream of the SpCas9 cleavage site. A knockout sequence
(5'TTTATTTAATTAAATAAA-3') was inserted at the cleavage site[52].
Dried oligonucleotides (IDT) were resuspended to a concentration of 250 µM in a
buffer of 10 mM HEPES-KOH, pH 7.5, incubated at 55°C for 1 hour, and mixed well by
pipetting up and down. The oligonucleotides were stored at -20°C.
*Delivery of gene editing cargoes with nucleofection*
The method for delivering *Sp*Cas9 RNPs and DNA repair templates into *M.*
*brevicollis* is as follows:

**Cell Culture.** One day prior to transfection, 60 ml of High Nutrient Medium was
inoculated to a final concentration of *M. brevicollis* at 10000 cells/ml. The culture was
split in two, and grown in two 75 cm$^2$ flasks at room temperature, approximately 22°C
(Falcon; Corning, Oneonta, NY, USA; Cat. No. 13-680-65).
**Assembly of Cas9/gRNA RNP.** Before starting transfections, the *Sp*Cas9 RNP was
assembled. For one reaction, 2 µl of 20 µM *Sp*Cas9 (NEB, Cat. No. M0646M) was
placed in the bottom of a 0.25 ml PCR tube, and then 2 µl of 100 µM gRNA was slowly
pipetted up and down with *Sp*Cas9 to gently mix the solutions. The mixed solution was
incubated at room temperature for 1 hour, and then placed on ice.
**Thaw DNA oligonucleotides.** Before using oligonucleotides in nucleofections, the
oligonucleotides were incubated at 55°C for 1 hour.
**Cell Washing.** After 24 hours of growth, bacteria were washed away from *M.
brevicollis* cells through three consecutive rounds of centrifugation and resuspension in
artificial seawater (ASW). The culture flasks were combined and vigorously shaken for
30 s, and then transferred to 50-ml conical tubes and spun for 5 min at 2000 × *g* and
22°C. The supernatant was removed with a serological pipette, and residual media were
removed with a fine-tip transfer pipette. The cell pellets were resuspended in a single
conical tube in a total volume of 50 ml of ASW, vigorously shaken for 30 s, and then
centrifuged for 5 min at 2050 × *g*. The supernatant was removed as before. In a final
washing step, the cell pellet was resuspended in 50 mL ASW, shaken vigorously, and
centrifuged for 5 min at 2100 × *g*. After the supernatant was removed, the cells were
resuspended in a total volume of 400 µl of ASW. To count the cell density, cells were
diluted 100-fold in 200 µl of ASW, and fixed with 1 µl of 16% paraformaldehyde. Cells
were counted on a hemocytometer, and the remaining cells were diluted to a final
concentration of 5 × 10$^7$ choanoflagellate cells/ml. The resuspended cells were divided
into 100-µl aliquots with 5 × 10$^6$ cells per aliquot to immediately prime cells in the next
step.
**Cell Priming.** Each aliquot of *M. brevicollis* cells was incubated in priming buffer (40
mM HEPES-KOH, pH 7.5; 50 mM lithium citrate; 50 mM ʟ-cysteine; 15% [wt/vol] PEG
8000; and 2 µM papain) to remove the extracellular material coating the cell. The 100-µl
aliquots, which contained 5 × 10$^6$ cells, were centrifuged for 5 min at 1700 × *g* and at
room temperature. The supernatant was removed, and cells were resuspended in 100
µl of priming buffer and then incubated for 35 min. Priming was quenched by adding 10
µl of 50-mg/ml bovine serum albumin-fraction V (Thermo Fisher Scientific, Waltham,
MA; Cat. No. BP1600-100). Cells were then centrifuged for 5 min at 1250 ×*g* and 22°C
with the centrifuge brake set to a "soft" setting. The supernatant was removed with a
fine-tip micropipette, and the cells were resuspended in 25 µl of SG Buffer (Lonza).
**Nucleofection.** Each nucleofection reaction was prepared by adding 16 µl of cold
SG Buffer to 4 µl of the *Sp*Cas9 RNP that was assembled as described above. For
reactions that used two different guide RNAs, each gRNA was assembled with *Sp*Cas9

separately and 4 µl of each RNP solution were combined at this step. 2 µl of the repair
oligonucleotide template was added to the *Sp*Cas9 RNP diluted in SG buffer. Finally, 2
µl of primed cells were added to the solution with Cas9 RNP and the repair template.
The nucleofection reaction was placed in one well of a 16-well nucleofection strip
(Lonza; Cat. No. V4XC-2032). The nucleofection strip was placed in the X-unit (Lonza;
Cat. No. AAF-1002F) connected to a Nucleofector 4D core unit (Lonza; Cat. No. AAF-
1002B), and the EO100 pulse was applied to each well.
***Recovery.*** 100 µl of cold recovery buffer (10 mM HEPES-KOH, pH 7.5; 0.9 M
sorbitol; 8% [wt/vol] PEG 8000) was added to the cells immediately after pulsation. After
5 minutes, the whole volume of the transfection reaction plus the recovery buffer was
transferred to 1 ml of High Nutrient Medium in a 12-well plate.
***Cycloheximide Selection in M. brevicollis.*** One day after transfection, 10 µl of 10
µg/ml cycloheximide was added per 1 mL culture of transfected cells. The cells were
incubated with cycloheximide for 5 days prior to clonal isolation and genotyping.
***Genotyping***. Cells were harvested for genotyping by spinning 0.5ml of cells at
4000*g* and 22°C for 5 min. The supernatant was removed and DNA was isolated either
by Base-Tris extraction [in which the cell pellet was resuspended in 20uL base solution
(25mM NaOH, 2mM EDTA), boiled at 100°C for 20 min, cooled at 4°C for 5 min, and
neutralized with 20uL Tris solution (40mM Tris-HCl, pH 7.5)], or by DNAzol Direct [in
which the cell pellet was resuspended in 50uL and incubated at room temperature for
30 min (Molecular Research Center, Inc. [MRC, Inc.], Cincinnati, OH; Cat. No. DN131)].
3 µl of the DNA solution was added to a 25 µl PCR reaction (DreamTaq Green PCR
Master Mix, Thermo Fisher Scientific Cat No K1082) and amplified with 34 rounds of
thermal cycling.
807
808 <u>Data Availability</u>
809 Source data files for all western blots are provided in Source Data File 1. Raw
810 sequencing reads and normalized gene counts for all RNA-seq experiments have been
811 deposited at the NCBI GEO under accession GSE174340.
812
813
814

  **Figure Legends**

**Figure 1. *P. aeruginosa* has pathogenic effects on *M. brevicollis***

**(A)** Immunofluorescence illuminates the diagnostic cellular architecture of *M. brevicollis*,

including an apical flagellum (f) made of microtubules, surrounded by an actin-filled

microvilli feeding collar (co). Staining for tubulin (green) also highlights cortical

microtubules that run along the periphery of the cell body, and staining for F-actin

(magenta) highlights basal filopodia (fp). DNA staining (blue) highlights the nucleus (n).

**(B)** *M. brevicollis* exhibits truncated flagella after exposure to *P. aeruginosa. M.*

*brevicollis* were exposed to *E. coli* or *P. aeruginosa* for 24 hours, and then fixed and

immunostained. Arrows point to flagella. Green: anti-tubulin antibody (flagella and cell

body), magenta: phalloidin (collar), blue: Hoechst (bacterial and choanoflagellate

nuclei). Scale bars represent 10$\mu$m. Flagellar length was quantified using Fiji, and

statistical analysis (unpaired t-tests) was performed in GraphPad software. **(C)**

Exposure to *P. aeruginosa*, but not other Gammaproteobacteria, results in *M. brevicollis*

cell death. Bacteria were added to *M. brevicollis* culture at an MOI of 1.5 (at Hours=0),

and *M. brevicollis* cell density was quantified at indicated time points. Data represent

mean +/- SD for three biological replicates. Statistical analysis (multiple unpaired t-tests)

was performed in GraphPad software; *p*-values shown are from comparisons between

*Flavobacterium* and *P. aeruginosa.* **(D-F)** *M. brevicollis* does not ingest *P. aeruginosa*

bacteria. **(D,E)** *M. brevicollis* were fed either fluorescent *E. coli* (D) or *P. aeruginosa* (E)

for one hour, and then visualized by DIC (D,E, left) and green fluorescence (D, E, right).

Fluorescent food vacuoles were observed in choanoflagellates fed *E. coli,* but not *P.*

*aeruginosa.* **(F)** *M. brevicollis* was exposed to GFP-expressing *E. coli*, *V.*

*parahaemolyticus*, *C. jejuni*, or *P. aeruginosa* (MOI=50) for one hour, and then imaged

by DIC and green fluorescence to quantify number of cells with internalized bacteria.

Choanoflagellate cells with ≥1 GFP+ food vacuole were scored as GFP+, and cells

without any GFP+ food vacuoles were scored as GFP–. Data represent cells quantified

over three biological replicates. **(G,H)** *P. aeruginosa* does not broadly inhibit *M.*

*brevicollis* phagocytosis. **(G)** Internalization of 0.2$\mu$m fluorescent beads was used to

quantify phagocytic activity after exposure to *E. coli* or *P. aeruginosa* bacteria. Although

cells did not phagocytose *P. aeruginosa,* cells exposed to *E. coli* and *P. aeruginosa* had

similar phagocytic uptake of beads. Data represent n=600 cells from three biological

replicates. Statistical analyses (multiple unpaired t-tests) were performed in GraphPad

software. **(H)** Exposure to *P. aeruginosa* does not inhibit phagocytic uptake of *E. coli.*

Internalization of fluorescent *E. coli* or *P. aeruginosa* bacteria was quantified after

exposure to unlabeled *P. aeruginosa* (PAO1 strain)*.* Data represent n=200 cells from

two biological replicates. Statistical analysis (unpaired t-test) was performed in

GraphPad software. **(I)** Secreted *P. aeruginosa* molecules are sufficient to induce *M.*

*brevicollis* cell death. 5% (vol/vol) bacterial conditioned medium was added to *M.*

*brevicollis* culture (at Hours=0), and *M. brevicollis* cell density was quantified at
indicated time points. Data represent mean +/- SD for three biological replicates.
Statistical analysis (multiple unpaired t-tests) was performed in GraphPad software, and
*p*-values shown are from comparisons between *Flavobacterium* and *P. aeruginosa.* **(J)**
Sustained exposure to secreted *P. aeruginosa* molecules is required to induce *M.*
*brevicollis* cell death. *P. aeruginosa* or *Flavobacterium* conditioned medium (5% vol/vol)
was added to stationary-phase *M. brevicollis* cultures. After indicated times, cultures
were washed and resuspended in fresh media. *M. brevicollis* cell density was quantified
after 24 hours. The % survival is a measure of the cell density of *P. aeruginosa*-treated
cells relative to *Flavobacterium*-treated controls. Data represent mean +/- SD for three
biological replicates.
866
867
868
869   **Figure 1 – figure supplement 1. Presence of animal innate immune genes in**
870   **choanoflagellates (A)** The transcriptomes of 21 choanoflagellate species[3] were
871   searched for genes that play key roles in animal innate immune responses. Evidence
872   for gene presence was based on sequence homology in a BLAST-based approach and
873   conserved domain architectures, as described in Richter et al., 2018.
874
875   **Video 1, Supplementary file 1. *P. aeruginosa* influences *M. brevicollis* motility.**
876   Movies depicting *M. brevicollis* cultures after exposure to *E. coli* or *P. aeruginosa*
877   bacteria for 16 hours. In the absence of pathogenic bacteria, like *E. coli*, M. *brevicollis* is
878   a highly motile flagellate and swims up in the water column. However, co-culturing *M.*
879   *brevicollis* with *P. aeruginosa* results in reduced motility and cell settling.
880
**Figure 2. STING is upregulated in *M. brevicollis* after exposure to *P. aeruginosa***
**(A,B)** *STING* transcript levels increase in response to *P. aeruginosa.* **(A)** Volcano plot
displaying genes differentially expressed between *M. brevicollis* exposed to *P.*
*aeruginosa* PAO1 and *Flavobacterium* (control) conditioned medium for three hours.
Differentially expressed genes are depicted by blue (674 upregulated genes) and yellow
(232 downregulated genes) dots (fold change≥2; FDR≤1e$^{-4}$). Select genes that are
upregulated or may function in innate immunity are labeled. RNA-seq libraries were
prepared from four biological replicates. **(B)** After a three-hour treatment, *STING* mRNA
levels (determined by RNA-seq) increase 1.42 fold in cells exposed to *Flavobacterium*
conditioned medium and 5.54 fold in cells exposed to *P. aeruginosa* conditioned
medium, compared to untreated controls. **(C)** STING protein levels increase after
exposure to *P. aeruginosa*. STING levels were examined by immunoblotting at indicated
timepoints after exposure to *Flavobacterium* or *P. aeruginosa* conditioned medium (5%
vol/vol). Tubulin is shown as loading control, and intensity of STING protein bands were

quantified relative to tubulin.  Statistical analysis (one-way ANOVA, Dunnett's multiple
comparison) was performed in GraphPad software, and *p*-values shown are calculated
using 0 hour timepoint as the control group.
898
899
900
**Figure 2 – figure supplement 2. *M. brevicollis* response to *P. aeruginosa,* and**
**STING antibody validation and protein alignment. (A)** Gene ontology enrichment
analysis of genes identified as differentially expressed (fold change≥2; FDR≤1e$^{-4}$) after
exposure to *P. aeruginosa*. Due to lack of annotation, >40% of the differentially
expressed genes were not included in the enrichment analysis. **(B)** qRT-PCR validation
of *STING* mRNA after exposure to *Flavobacterium* or *P. aeruginosa* conditioned media
for three hours, compared to vehicle control. Error bars represent SD. Statistical
analysis (t-test) was performed in GraphPad software  **(C)** To validate the *M. brevicollis*
STING antibody, cell lysates from *M. brevicollis* were immunoblotted alongside cell
lysates from *S. rosetta,* a closely-related choanoflagellate species that does not have a
STING homolog. A band at 36kD, the predicted size of *M. brevicollis* STING, is
detectable in *M. brevicollis* lysate but not *S. rosetta* lysate. Arrow indicates STING band.
Non-specific bands are likely due to co-cultured feeding bacteria. Tubulin is shown as
loading control. **(D)** Protein sequence alignment (generated by Clustal Omega multiple
sequence alignment) of *M. brevicollis* and animal STING proteins, colored by similarity.
*M. brevicollis* STING and human STING are 19.1% identical and 36.6% similar at the
amino acid level.
918
919
920
**Figure 3. 2'3' cGAMP, but not bacterially-produced cyclic dinucleotides, induces**
**elevated levels of STING**
**(A)** Schematic of choanoflagellate (*M. brevicollis*), sea anemone (*N. vectensis*), insect
(*D. melanogaster*) and mammalian (*M. musculus* and *H. sapiens*) STING proteins.
Transmembrane (TM) domains are depicted in gray, STING cyclic dinucleotide binding
domain (CDN) in purple, and C-terminal tail domain (CTT) in blue. **(B)** Partial protein
sequence alignment (generated by Clustal Omega multiple sequence alignment) of *M.*
*brevicollis* and animal STING proteins, colored by similarity. *M. brevicollis* STING and
human STING are 19.1% identical and 36.6% similar at the amino acid level. Key cyclic
dinucleotide-interacting residues from human STING structure are indicated by circles.
**(C)** Dose-response curves of *M. brevicollis* exposed to cyclic dinucleotides for 24 hours
reveal that treatment with 2'3'cGAMP, but not 3'3' cGAMP, c-di-AMP, or c-di-GMP,
leads to *M. brevicollis* cell death in a dose-dependent manner. Data represent mean +/-
SD for at least three biological replicates. **(D)** STING protein levels increase after

exposure to 2'3'cGAMP, but not bacterially-produced cyclic dinucleotides. *M. brevicollis*
STING levels were examined by immunoblotting 5 hours after exposure to 2'3'cGAMP
(100 μM), 3'3'cGAMP (200 μM), c-di-GMP (200 μM), or c-di-AMP (200 μM). Tubulin is
shown as loading control, and intensity of STING protein bands were quantified relative
to tubulin. Shown is a representative blot from three biological replicates. Statistical
analysis (one-way ANOVA, Dunnett's multiple comparison) was performed in GraphPad
software. **(E)** STING protein levels increase and remain elevated after exposure to 100
μM 2'3'cGAMP. Tubulin is shown as loading control, and data are representative of
three biological replicates. Statistical analysis (one-way ANOVA, Dunnett's multiple
comparison) was performed in GraphPad software, and *p*-values shown are calculated
using 0 hour timepoint as control group.

**Figure 3 – figure supplement 1. *M. brevicollis* has distinct responses to 2'3'**
**cGAMP and 3'3' cGAMP**
**(A,B)** Volcano plots displaying RNA-seq differential expression analysis of *M. brevicollis*
treated with (A) 100 μM 2'3'cGAMP or (B) 200 μM 3'3'cGAMP for 3 hours, relative to an
untreated control. Genes with a fold change ≥2 and false discovery rate ≤10e$^{-4}$ are
depicted by black dots. STING is highlighted in red. RNA-seq libraries were prepared
from three (2'3' cGAMP) or two (3'3' cGAMP) biological replicates. **(C)** *M. brevicollis*
*STING* mRNA levels increase in response to 2'3'cGAMP. (Left) RNA-seq fold change of
*STING* mRNA after exposure to 100 μM 2'3'cGAMP or 200 μM 3'3'cGAMP for three
hours, compared to vehicle control. (Right) qRT-PCR fold change of *STING* mRNA after
exposure to 100 μM 2'3'cGAMP or 200 μM 3'3'cGAMP for three hours, compared to
vehicle control.  **(D)** Venn diagram comparing the overlap of genes identified as
differentially expressed after treatment with 2'3'cGAMP, 3'3'cGAMP, and *P. aeruginosa*
(DEG cutoff: fold change ≥ 3, false discovery rate ≤ 10e$^{-4}$). **(E,F)** Representative
immunostained *M. brevicollis* demonstrating 2'3'cGAMP stimulates the formation of
STING puncta at perinuclear regions. *M. brevicollis* was left untreated (E), or exposed to
100 μM 2'3'cGAMP (F) for 5 hours. Cells were fixed and STING levels and localization
were probed using an anti-STING antibody. **(E',F')** Exposure to 2'3'cGAMP results in
increased numbers of STING puncta compared to untreated controls. **(E'',F'')** Z-slice
images of the plane containing the nucleus 'n' show that STING puncta localize to
perinuclear regions. Green: anti-tubulin antibody (flagella and cell body), magenta: anti-
STING antibody, blue: Hoechst. Scale bar represents 2 μm.

**Figure 4. Transfection reveals STING localization to perinuclear and endoplasmic reticulum regions**

**(A)** STING-mTFP localizes to tubule-like structures around the nucleus in cells stably expressing STING-mTFP. Green: anti-tubulin antibody (flagella and cell body), magenta: anti-STING antibody, blue: Hoechst. Scale bar represents 2 µm. **(B,C)** Fluorescent markers and live cell imaging reveal that STING is localized to the endoplasmic reticulum (ER). Cells were co-transfected with STING-mTFP and an mCherry fusion protein that localizes either to the endoplasmic reticulum (B; mCherry-HDEL) or mitochondria (C; Cox4-mCherry )[4]. Cells were recovered in the presence of *Flavobacterium* feeding bacteria for 28 hours after co-transfection, and then live cells were visualized with super-resolution microscopy. Each panel shows Z-slice images of a single representative cell. In confocal Z-slice images, cells are oriented with the apical flagella pointing up, and the nucleus is marked by 'n' when clearly included in the plane of focus. STING colocalized with the ER marker (B''), but not the mitochondrial marker (C''). Scale bar represents 2 µm.

**Figure 5. STING mediates responses to 2'3'cGAMP and *P. aeruginosa***

**(A)** The genotypes of wild type and genome-edited *STING⁻* strains at the *STING* locus. **(B)** STING protein is not detectable by immunoblot in *STING⁻* cells. Shown is a representative blot from three biological replicates. **(C,D)** STING is necessary for 2'3'cGAMP-induced cell death. **(C)** Wild type and *STING⁻* strains were treated with increasing concentrations of 2'3'cGAMP, and survival was quantified after 24 hours. In contrast to wild type cells, 2'3'cGAMP does not induce cell death in *STING⁻* cells. Data represent mean +/- SD for four biological replicates. **(D)** Wild type and *STING⁻* cells were transfected with STING-mTFP, and treated with puromycin to generate stable clonal strains. Stable expression of STING-mTFP in *STING⁻* cells partially rescued the phenotype of 2'3'cGAMP-induced cell death. Data represent mean +/- SD for three biological replicates. Statistical analysis (multiple unpaired t-tests) was performed in GraphPad software. **(E)** Wild type and *STING⁻* strains have distinct transcriptional responses to 2'3' cGAMP. Differential expression analysis was performed on wild type and *STING⁻* cells treated with 100µM 2'3'cGAMP or a vehicle control for three hours. A heatmap comparing the $\log_2$ fold change of genes identified as differentially expressed (FC ≥2; FDR≤10⁻⁴) in wild type cells after 2'3' cGAMP treatment, to their $\log_2$ fold change in *STING⁻* cells after 2'3' cGAMP treatment. RNA-seq libraries were prepared from two biological replicates. **(F)** Presence of STING in the transcriptomes of diverse choanoflagellate species. Data from Richter et al. 2018[3]. **(G)** Effects of 2'3'cGAMP on different choanoflagellate species. Choanoflagellates were grown to late-log phase, and treated with increasing concentrations of 2'3'cGAMP. Survival was quantified after 24

hours. 2'3'cGAMP only affected the survival of *M. brevicollis* and *S. macrocollata*, the

two sequenced choanoflagellate species with a STING homolog. Data represent mean

+/- SD for three biological replicates. **(H)** Wild type and *STING⁻* cells have similar

survival responses to LPS, suggesting that STING is not required for mediating a

response to LPS. Wild type and *STING⁻* strains were treated with increasing

concentrations of *E. coli* LPS, and survival was quantified after 24 hours. Data represent

mean +/- SD for four biological replicates. Statistical analysis (multiple unpaired t-tests)

was performed in GraphPad software. **(I,J)** STING renders *M. brevicollis* more

susceptible to *P. aeruginosa*-induced growth inhibition. **(I)** Wild type and *STING⁻* cells

were exposed to control *Flavobacterium* or *P. aeruginosa* conditioned medium (5%

vol/vol), and cell densities were quantified at indicated time points. Data represent mean

+/- SD for three biological replicates. **(J)** Percent survival calculated from growth curves

in (I). Statistical analysis (multiple unpaired t-tests) was performed in GraphPad

software*.*

1028

1029

1030

**Figure 5- figure supplement 1. Characterizing *STING⁻* *M. brevicollis***

**(A)** Sanger sequences of the consensus genotype at the site of gene editing in wild type

and *STING⁻* cells. *STING⁻* cells have a 7 base-pair deletion that leads to premature stop

codons. **(B)** Growth curves of wild type and *STING⁻* cells indicate that both strains have

similar growth dynamics. Statistical analysis (multiple unpaired t-tests) was performed in

GraphPad software. **(C)** Volcano plot displaying RNA-seq differential expression

analysis of *STING⁻* cells treated with 100 µM 2'3'cGAMP for 3 hours, relative to an

untreated control. Genes with a fold change ≥2 and false discovery rate ≤10e$^{-4}$ are

depicted by black dots. RNA-seq libraries were prepared from two biological replicates.

**(D)** Venn diagram comparing the overlap of genes identified as differentially expressed

(FC ≥3; FDR≤10$^{-4}$) after treatment with 2'3'cGAMP in wild type and *STING⁻* cells. **(E)**

Protein sequence alignment (generated by Clustal Omega multiple sequence

alignment) of STING proteins from choanoflagellates *S. macrocollata* and *M. brevicollis*

and animals, colored by similarity.

1045

1046

1047

**Figure 6. STING mediates 2'3'cGAMP-induced autophagic pathway**

**(A)** 2'3'cGAMP-induced Atg8 lipidation requires STING. WT and *STING⁻* cells stably

expressing mCherry-Atg8 were treated with a vehicle control or 100 µM 2'3'cGAMP for

3 hours, followed by immunoblotting. The band intensity of Atg8-I (unmodified Atg8) and

Atg8-II (lipidated Atg8) were quantified for each sample. Relative levels of Atg8

lipidation were assessed by dividing the band intensities of Atg8-II/Atg8-I. Tubulin is

shown as loading control. Immunoblot is representative of three biological replicates.
**(B)** 2'3'cGAMP induces Atg8 lipidation in chloroquine-treated wild type cells. WT cells
stably expressing mCherry-Atg8 were first incubated with 40 mM chloroquine for 6
hours, and then treated with a vehicle control or 100 μM 2'3'cGAMP for 3 hours in the
presence of chloroquine, followed by immunoblotting. For each sample, relative levels
of Atg8 lipidation were assessed by dividing the band intensities of Atg8-II/Atg8-I.
Tubulin is shown as loading control. Immunoblot is representative of three biological
replicates. For a representative immunoblot and quantification of Atg8-II/Atg8-I levels in
chloroquine-treated *STING⁻* cells, refer to Figure 6 - figure supplement 1B.  **(C-G)**
STING is required for 2'3'cGAMP-induced autophagosome formation. **(C-F)** WT and
*STING⁻* cells stably expressing mCherry-Atg8 were treated with a vehicle control or 100
μM 2'3'cGAMP for 3 hours, and then fixed and immunostained. **(C,D)** Representative
confocal images of wild type cells show that Atg8 puncta accumulate after 2'3'cGAMP
treatment. Magenta: mCherry-Atg8; Green: anti-tubulin antibody (flagella and cell body).
**(E,F)** Representative confocal images of *STING⁻* cells show that Atg8 remains evenly
distributed in the cytoplasm after 2'3'cGAMP treatment. **(G)** The number of Atg8
puncta/cell was quantified for WT and *STING⁻* cells treated with a vehicle control or
2'3'cGAMP for three hours. Data represent cells quantified from two biological replicates
(n=150 cells per treatment group). Statistical analyses (unpaired two-tailed t-tests) were
performed in GraphPad software. **(H)** Treatment with lysosomotropic agents that inhibit
autophagy rescue 2'3' cGAMP-induced cell death in wild type cells. WT and *STING⁻*
cells were pre-treated with 40 mM chloroquine, 10mM $NH_4Cl$, or a vehicle control for 6
hours. Cells were then exposed to either 100 μM 2'3'cGAMP or a vehicle control for 18
hours before quantifying survival. Data represent mean +/- SD for three biological
replicates. Statistical analyses (multiple unpaired t-tests) were performed in GraphPad
software.
1080
1081
**Figure 6 – figure supplement 1. STING mediates 2'3'cGAMP-induced autophagic**
**signaling**
**(A)** Overexpression of mCherry-Atg8 does not alter the susceptibility of wild type and
*STING⁻* strains to 2'3'cGAMP. Wild type and *STING⁻* strains stably expressing mCherry-
Atg8 were treated with increasing concentrations of 2'3'cGAMP, and survival was
quantified after 24 hours. Data represent mean +/- SD for two biological replicates. **(B)**
2'3'cGAMP does not induce increased Atg8 lipidation in chloroquine-treated *STING⁻*
cells. *STING⁻* cells stably expressing mCherry-Atg8 were incubated with 40 mM
chloroquine for 6 hours, and then treated with a vehicle control or 100 μM 2'3'cGAMP
for 3 hours in the presence of chloroquine, followed by immunoblotting. For each
sample, relative levels of Atg8 lipidation were assessed by dividing the band intensities
of Atg8-II/Atg8-I. Tubulin is shown as loading control. Immunoblot is representative of

three biological replicates. Statistical analyses (unpaired two-tailed t-tests) were
performed in GraphPad software.
1096
1097    **Source data file 1.** Uncropped western blots for: Figure 2-  figure supplement 1C,
1098    Figure 2C, Figure 3D, Figure 3E, Figure 5B, Figure 6A, Figure 6B, Figure 6-
1099    supplement1B

## References

1. Shabalina, S. A. & Koonin, E. V. Origins and evolution of eukaryotic RNA interference. *Trends Ecol Evol* **23**, 578–587 (2008).

2. Richter, D. J. & Levin, T. C. The origin and evolution of cell-intrinsic antibacterial defenses in eukaryotes. *Curr Opin Genet Dev* **58–59**, 111–122 (2019).

3. Richter, D. J., Fozouni, P., Eisen, M. B. & King, N. Gene family innovation, conservation and loss on the animal stem lineage. *Elife* **7**, 946 (2018).

4. King, N. *et al.* The genome of the choanoflagellate Monosiga brevicollis and the origin of metazoans. *Nature* **451**, 783 788 (2008).

5. Brunet, T. & King, N. The Origin of Animal Multicellularity and Cell Differentiation. *Dev Cell* **43**, 124 140 (2017).

6. Richter, D. J. & King, N. The genomic and cellular foundations of animal origins. *Annu Rev Genet* **47**, 509 537 (2013).

7. Leadbeater, B. S. C. *The Choanoflagellates: Evolution, Ecology, and Biology*. (Cambridge University Press, 2015).

8. Dayel, M. J. & King, N. Prey capture and phagocytosis in the choanoflagellate Salpingoeca rosetta. *Plos One* **9**, e95577 (2014).

9. Wu, X. *et al.* Molecular evolutionary and structural analysis of the cytosolic DNA sensor cGAS and STING. *Nucleic Acids Res* **42**, 8243–8257 (2014).

10. Levin, T. C., Greaney, A. J., Wetzel, L. & King, N. The Rosetteless gene controls development in the choanoflagellate S. rosetta. *Elife* **3**, (2014).

11. Ablasser, A. & Chen, Z. J. cGAS in action: Expanding roles in immunity and inflammation. *Science* **363**, eaat8657 (2019).

12. Ahn, J. & Barber, G. N. STING signaling and host defense against microbial infection. *Exp Mol Medicine* **51**, 1–10 (2019).

13. Margolis, S. R., Wilson, S. C. & Vance, R. E. Evolutionary Origins of cGAS-STING Signaling. *Trends Immunol* **38**, 733–743 (2017).

14. Kranzusch, P. J. *et al.* Ancient Origin of cGAS-STING Reveals Mechanism of Universal 2′,3′ cGAMP Signaling. *Mol Cell* **59**, 891 903 (2015).

15. Ishikawa, H. & Barber, G. N. STING is an endoplasmic reticulum adaptor that facilitates innate immune signalling. *Nature* **455**, 674–678 (2008).

16. Sun, L., Wu, J., Du, F., Chen, X. & Chen, Z. J. Cyclic GMP-AMP Synthase Is a Cytosolic DNA Sensor That Activates the Type I Interferon Pathway. *Science* **339**, 786–791 (2013).

17. Burdette, D. L. *et al.* STING is a direct innate immune sensor of cyclic di-GMP. *Nature* **478**, 515–518 (2011).

18. Diner, E. J. *et al.* The Innate Immune DNA Sensor cGAS Produces a Noncanonical Cyclic Dinucleotide that Activates Human STING. *Cell Reports* **3**, 1355–1361 (2013).

19. Gao, P. *et al.* Cyclic [G(2′,5′)pA(3′,5′)p] Is the Metazoan Second Messenger Produced by DNA-Activated Cyclic GMP-AMP Synthase. *Cell* **153**, 1094–1107 (2013).

20. Ablasser, A. *et al.* cGAS produces a 2′-5′-linked cyclic dinucleotide second messenger that activates STING. *Nature* **498**, 380–384 (2013).

21. Moretti, J. *et al.* STING Senses Microbial Viability to Orchestrate Stress-Mediated Autophagy of the Endoplasmic Reticulum. *Cell* **171**, 809-823.e13 (2017).

22. Cohen, D. *et al.* Cyclic GMP–AMP signalling protects bacteria against viral infection. *Nature* **574**, 691–695 (2019).

23. Morehouse, B. R. *et al.* STING cyclic dinucleotide sensing originated in bacteria. *Nature* 1–8 (2020) doi:10.1038/s41586-020-2719-5.

24. Burroughs, A. M. & Aravind, L. Identification of Uncharacterized Components of Prokaryotic Immune Systems and Their Diverse Eukaryotic Reformulations. *J Bacteriol* **202**, (2020).

25. Alegado, R. A. *et al.* A bacterial sulfonolipid triggers multicellular development in the closest living relatives of animals. *Elife* **1**, e00013 (2012).

26. Brunet, T. *et al.* A flagellate-to-amoeboid switch in the closest living relatives of animals. *Elife* **10**, e61037 (2021).

27. Dayel, M. J. *et al.* Cell differentiation and morphogenesis in the colony-forming choanoflagellate Salpingoeca rosetta. *Dev Biol* **357**, 73 82 (2011).

28. Woznica, A., Gerdt, J. P., Hulett, R. E., Clardy, J. & King, N. Mating in the Closest Living Relatives of Animals Is Induced by a Bacterial Chondroitinase. *Cell* **170**, 1175 1183.e11 (2017).

29. Brunet, T. *et al.* Light-regulated collective contractility in a multicellular choanoflagellate. *Science* **366**, 326–334 (2019).

30. Laundon, D., Larson, B. T., McDonald, K., King, N. & Burkhardt, P. The architecture of cell differentiation in choanoflagellates and sponge choanocytes. *Plos Biol* **17**, e3000226 (2019).

31. Woznica, A. *et al.* Bacterial lipids activate, synergize, and inhibit a developmental switch in choanoflagellates. *Proc National Acad Sci* **113**, 7894 7899 (2016).

32. Ireland, E. V., Woznica, A. & King, N. Synergistic Cues from Diverse Bacteria Enhance Multicellular Development in a Choanoflagellate. *Appl Environ Microb* **86**, (2020).

33. Hake, K. *et al.* Colonial choanoflagellate isolated from Mono Lake harbors a microbiome. doi:10.1101/2021.03.30.437421.

34. Moradali, M. F., Ghods, S. & Rehm, B. H. A. Pseudomonas aeruginosa Lifestyle: A Paradigm for Adaptation, Survival, and Persistence. *Front Cell Infect Mi* **7**, 39 (2017).

35. Mahajan-Miklos, S., Tan, M.-W., Rahme, L. G. & Ausubel, F. M. Molecular Mechanisms of Bacterial Virulence Elucidated Using a Pseudomonas aeruginosa– Caenorhabditis elegans Pathogenesis Model. *Cell* **96**, 47–56 (1999).

36. Pukatzki, S., Kessin, R. H. & Mekalanos, J. J. The human pathogen Pseudomonas aeruginosa utilizes conserved virulence pathways to infect the social amoeba Dictyostelium discoideum. *Proc National Acad Sci* **99**, 3159–3164 (2002).

37. Rahme, L. G. *et al.* Use of model plant hosts to identify Pseudomonas aeruginosa virulence factors. *Proc National Acad Sci* **94**, 13245–13250 (1997).

38. Uribe-Querol, E. & Rosales, C. Control of Phagocytosis by Microbial Pathogens. *Front Immunol* **8**, 1368 (2017).

39. Yoon, S. *et al.* Pseudomonas syringae evades phagocytosis by animal cells via type III effector-mediated regulation of actin filament plasticity. *Environ Microbiol* **20**, 3980–3991 (2018).

40. Klockgether, J. & Tümmler, B. Recent advances in understanding Pseudomonas aeruginosa as a pathogen. *F1000research* **6**, 1261 (2017).

41. Tanaka, Y. & Chen, Z. J. STING Specifies IRF3 Phosphorylation by TBK1 in the Cytosolic DNA Signaling Pathway. *Sci Signal* **5**, ra20–ra20 (2012).

42. Liu, S. *et al.* Phosphorylation of innate immune adaptor proteins MAVS, STING, and TRIF induces IRF3 activation. *Science* **347**, aaa2630 (2015).

43. Zhang, C. *et al.* Structural basis of STING binding with and phosphorylation by TBK1. *Nature* **567**, 394–398 (2019).

44. Abe, T. & Barber, G. N. Cytosolic-DNA-Mediated, STING-Dependent Proinflammatory Gene Induction Necessitates Canonical NF-κB Activation through TBK1. *J Virol* **88**, 5328–5341 (2014).

45. Gui, X. *et al.* Autophagy induction via STING trafficking is a primordial function of the cGAS pathway. *Nature* **567**, 262–266 (2019).

46. Yamashiro, L. H. *et al.* Interferon-independent STING signaling promotes resistance to HSV-1 in vivo. *Nat Commun* **11**, 3382 (2020).

47. Prabakaran, T. *et al.* Attenuation of cGAS-STING signaling is mediated by a p62/SQSTM1-dependent autophagy pathway activated by TBK1. *Embo J* **37**, (2018).

48. Mann, C. C. de O. *et al.* Modular Architecture of the STING C-Terminal Tail Allows Interferon and NF-κB Signaling Adaptation. *Cell Reports* **27**, 1165-1175.e5 (2019).

49. Watson, R. O. *et al.* The Cytosolic Sensor cGAS Detects Mycobacterium tuberculosis DNA to Induce Type I Interferons and Activate Autophagy. *Cell Host Microbe* **17**, 811–819 (2015).

50. Booth, D. S., Szmidt-Middleton, H. & King, N. Choanoflagellate transfection illuminates their cell biology and the ancestry of animal septins. *Mol Biol Cell* **29**, mbcE18080514 (2018).

51. Dobbs, N. *et al.* STING Activation by Translocation from the ER Is Associated with Infection and Autoinflammatory Disease. *Cell Host Microbe* **18**, 157–168 (2015).

52. Booth, D. S. & King, N. Genome editing enables reverse genetics of multicellular development in the choanoflagellate Salpingoeca rosetta. *Elife* **9**, e56193 (2020).

53. Liu, Y. *et al.* Inflammation-Induced, STING-Dependent Autophagy Restricts Zika Virus Infection in the Drosophila Brain. *Cell Host Microbe* **24**, 57-68.e3 (2018).

54. Klionsky, D. J. *et al.* Guidelines for the use and interpretation of assays for monitoring autophagy (4th edition) 1. *Autophagy* **17**, 1–382 (2021).

55. Tanida, I., Minematsu-Ikeguchi, N., Ueno, T. & Kominami, E. Lysosomal Turnover, but Not a Cellular Level, of Endogenous LC3 is a Marker for Autophagy. *Autophagy* **1**, 84–91 (2005).

56. Pokatayev, V. *et al.* Homeostatic regulation of STING protein at the resting state by stabilizer TOLLIP. *Nat Immunol* **21**, 158–167 (2020).

57. McFarland, A. P. *et al.* Sensing of Bacterial Cyclic Dinucleotides by the Oxidoreductase RECON Promotes NF-κB Activation and Shapes a Proinflammatory Antibacterial State. *Immunity* **46**, 433–445 (2017).

58. Margolis, S. R. *et al.* The STING ligand 2′3′-cGAMP induces an NF-κB-dependent anti-bacterial innate immune response in the starlet sea anemone Nematostella vectensis. (2021) doi:10.1101/2021.05.13.443009.

59. Cai, H. *et al.* 2′3′-cGAMP triggers a STING- and NF-κB–dependent broad antiviral response in Drosophila. *Sci Signal* **13**, eabc4537 (2020).

60. Martin, M., Hiroyasu, A., Guzman, R. M., Roberts, S. A. & Goodman, A. G. Analysis of Drosophila STING Reveals an Evolutionarily Conserved Antimicrobial Function. *Cell Reports* **23**, 3537-3550.e6 (2018).

61. Goto, A. *et al.* The Kinase IKKβ Regulates a STING- and NF-κB-Dependent Antiviral Response Pathway in Drosophila. *Immunity* **49**, 225-234.e4 (2018).

62. Hua, X. *et al.* Stimulator of interferon genes (STING) provides insect antiviral immunity by promoting Dredd caspase–mediated NF-κB activation. *J Biol Chem* **293**, 11878–11890 (2018).

63. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676–682 (2012).

64. Kim, D. *et al.* TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol* **14**, R36 (2013).

65. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

66. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

67. Peng, D., Kurup, S. P., Yao, P. Y., Minning, T. A. & Tarleton, R. L. CRISPR-Cas9-Mediated Single-Gene and Gene Family Disruption in Trypanosoma cruzi. *Mbio* **6**, e02097-14 (2015).

**Table 1. Bacteria screened for pathogenic effects**

| Bacterium | Pathogenic effects | Reference or details | Source |
|---|---|---|---|
| *Aeromonas hyrophila* | – | Environmental isolate | This study |
| *Bacillus aquimaris* | – | Environmental isolate | This study |
| *Bacillus badius* | – | Mouse isolate | Julie Pfeiffer |
| *Bacillus cereus* | – | Environmental isolate | This study |
| *Bacillus indicus* | – | Environmental isolate | This study |
| *Bacillus marisflavi* | – | Environmental isolate | This study |
| *Bacillus pumilus* | – | Mouse isolate | Julie Pfeiffer |
| *Bacillus safensis* | – | Mouse isolate | Julie Pfeiffer |
| *Bacillus subtilus* | – | ATCC 6633 | Julie Pfeiffer |
| *Bacteroides acidifaciens* | – | Mouse isolate | Julie Pfeiffer |
| *Burkholderia multivorans* | – | ATCC 17616 | David Greenberg |
| *Campylobacter jejuni* GFP | – | DRH3123 | David Hendrixson |
| *Deinococcus* sp. | – | Environmental isolate | This study |
| *Enterococcus cloacae* | – | Mouse isolate | Julie Pfeiffer |
| *Enterococcus faecium* | – | Mouse isolate | Julie Pfeiffer |
| *Escherichia coli* BW25113 | – | Datsenko and Wanner, 2000 | David Greenberg |
| *Escherichia coli DH5a* GFP | – | | David Hendrixson |
| *Escherichia coli* ECC-1470 | – | Leimbach et al., 2015 | Julie Pfeiffer |
| *Escherichia coli* K12 | – | ATCC 10798 | Julie Pfeiffer |
| *Flavobacterium sp.* | – | King et al., 2008 | Isolated from ATCC PRA-258 |
| *Lactobacillus johnsonii* | – | Mouse isolate | Julie Pfeiffer |
| *Pseudoalteromonas sp.* | – | Environmental isolate | This study |
| *Pseudomonas aeruginosa* PA-14 | + | Rahm et al., 1995 | Andrew Koh |
| *Pseudomonas aeruginosa* PAO1 | + | ATCC 15692 | David Greenberg |
| *Pseudomonas aeruginosa* PAO1-GFP | + | Bloemberg et al., 1997 | David Greenberg |
| *Pseudomonas granadensis* | – | Environmental isolate | This study |
| *Staphylococcus aureus* | – | ATCC 23235 | Julie Pfeiffer |
| *Staphylococcus* sp. | – | Mouse isolate | Julie Pfeiffer |
| *Vibrio alginolyticus* | – | Environmental isolate | Kim Orth |
| *Vibrio furnissii* | – | Environmental isolate | This study |
| *Vibrio parahaemolyticus* | – | Environmental isolate | This study |
| *Vibrio parahaemolyticus* RimD-GFP | – | Ritchie et al., 2012 | Kim Orth |
| *Vibrio ruber* | – | Environmental isolate | This study |
| *Vibrio sp.* | – | Environmental isolate | This study |

**Table 2. *P. aeruginosa* deletion strains**

| Strain name | Gene | Putative ORF function | Effects on *M. brevicollis* | |
|---|---|---|---|---|
| | | | **Truncated Flagellum/ Settling** | **Cell Death** |
| MPAO1 | | parent to library stain | + | + |
| PW5035 | pvdE | pyoverdine biosynthesis protein PvdE | + | + |
| PW5034 | pvdE | pyoverdine biosynthesis protein PvdE | + | + |
| PW1059 | exoT | exoenzyme T | + | + |
| PW3078 | toxA | exotoxin A precursor | + | + |
| PW3079 | toxA | exotoxin A precursor | + | + |
| PW4736 | exoY | adenylate cyclase ExoY | + | + |
| PW4737 | exoY | adenylate cyclase ExoY | + | + |
| PW6886 | rhlA | rhamnosyltransferase chain A | + | + |
| PW6887 | rhlA | rhamnosyltransferase chain A | + | + |
| PW7478 | exoS | exoenzyme S | + | + |
| PW7479 | exoS | exoenzyme S | + | + |
| PW7303 | lasB | elastase LasB | + | + |
| PW7302 | lasB | elastase LasB | + | + |
| PW3252 | aprA | alkaline metalloproteinase precursor | + | + |
| PW3253 | aprA | alkaline metalloproteinase precursor | + | + |
| PW4282 | lasA | LasA protease precursor | + | + |
| PW4283 | lasA | LasA protease precursor | + | + |
| RP436 | popB | T3SS translocase | + | + |
| RP576 | exoS, exoT, exoY | T3SS effector molecules | + | + |

**Table 3. *M. brevicollis* response to *P. aeruginosa* factors**

| Treatment | Cell Death | Interpretation |
| --- | --- | --- |
| Live *P. aeruginosa* | ++ | |
| *P. aeruginosa* conditioned media (CM) | ++ | Factor(s) are secreted by *P. aeruginosa* |
| *P. aeruginosa* outer membrane vesicles | – | Factor(s) are not present in outer membrane vesicles |
| CM, boiled 20 min | ++ | Factor(s) are not heat labile |
| CM + proteinase K, followed by 80C for 30 min | ++ | Factor(s) are not proteins |
| CM + DNAse and RNAse | ++ | Factor(s) are not nucleic acids |
| CM MeOH extraction | ++ | Factor(s) are organic compounds |

Figure 1

**A**

Figure 1 - figure supplement 1

Figure 2

**A**

Condensin complex
Chromosome condensation
Glutathione peroxidase activity
Membrane depolarization during action potential
Alpha-galactosidase activity
Glycoside catabolic process
Response to stress
Cilium movement
Mitochondrial fission
Intracellular membrane-bounded organelle
ATPase activity
DNA-directed DNA polymerase activity
Microtubule-based movement
Hydrolase activity
Carbohydrate metabolic process
Oxidoreductase activity
Regulation of metaphase/anaphase transition
RNA modification
Intra-S DNA damage checkpoint
Heme binding
Zinc ion binding

Upregulated genes
Downregulated genes

Fold enrichment

**B**

$STING$ $\log_2$ fold-change (qRT-PCR)

$p=1.5*10^{-3}$

Flavobacterium
P. aeruginosa

**C**

*M. brevicollis*  *S. rosetta*

kDa

STING

Tubulin
55

**D**

```
M. brevicollis   1 MM----VNLSDLSHLSQRGWAQVFVTLA-------------------AAAISIF   31
D. melanogaster  1 MAI--ASNVVEAGNAVRAEKGRKYFYFRK--------------MIGDYIDTSIRIV   40
N. vectensis     1 MRRAEENNGFGTI-PKRRNQHTPFYASIGMIVVIVAFTSYHITSYGDDRNRAMRQY   56
M. musculus      1 ---MPYSNLHPAI-PRPRGHRSKYVALI-FLVASL----MILWVAKDPPNHTLKYL   47
H. sapiens       1 ---MPHSSLHPSI-PCPRGHGAQKAALV-LLSACL-----VTLWGLGPPEHTLRYL   47

M. brevicollis  32 YFSFSPFEVAAGICASIAAAGAVPLLFDAVHYLIAFCISAPDARP------PLRTVW   82
D. melanogaster 41 --------------ATVFLADLLLRLYRCVV-EYGSNGRYYLPEDRLW   73
N. vectensis    57 SFTFSLA---------------YLAFLVGELLRRCCLFAEEYRHIETRY--NGSLK   95
M. musculus     48 ----ALH---------------LASHELGLLLKNLCCLAEELCHVQSRY--QGSYW   82
H. sapiens      48 ----VLH---------------LASLQLGLLLLNGVCSLAEELRHIHSRY--RGSYW   82

M. brevicollis  83 TKTRLQRWSGLSILTFIVLACGLY---FASPSP-------------------112
D. melanogaster 74 IILRR--SCTYNNRSIYLIVGFLLVAFFRISVT-GNY-----RNVMPTTLFLFQMP  121
N. vectensis    96 KAIQTTFSFGHNNV-LFVASLLFFVVFVASNDPNGSSSVIQGNSTAEPHT--EMRQT  149
M. musculus     83 VRACLGCPIHCMAMILLSSYFYFL-QNTADI---------------Y--------  115
H. sapiens      83 RTVRACLGCPLRRGALLLLSIYFYSLPNAVGP--------------P--------  116

M. brevicollis 113 MSASNLFAVFALSLCNSVFSAMVAPRCGRLEVVD-LATHPNTVLFRDFGAASAMSYW  168
D. melanogaster 122 LYWI--WSFT-----------DMDQSTLSYSHWIRDSHGLDYAAGMASNYF  159
N. vectensis   150 SGWQGLWGQFIIS---ALLTPLVVHLLGLRELSKVEESQLNEKENKNVADGLAWSYY  203
M. musculus    116 LSWM--FGL-------LVLYKSLSMLGLQSLTPAEVSAVCEEKKLNVAHGLAWSYY  163
H. sapiens     117 ITWM--LAL-------LGLSQALNILLGLKGLAPAEISAVCEKGNFNVAHGLAWSYY  164

M. brevicollis 169 HGYLQHIAGDRAA--------LEERFRDSDVGGMRVPRKLYILVPQDCEVNASDNKY  217
D. melanogaster 160 HGYLKLSLPERKDDGLKHRLAVYEDKN N----VTFGIKRLVLIPDEMFVNGVLESH  212
N. vectensis   204 FGYLKFVLPELEK-----QIEKTSKFRS-----KEKFVKKMFILIPSNCFWDDKIPGS  251
M. musculus    164 IGYLRLILPGLQA-----RIRMFNQLHN-NMLSGAGSRRLYILFPLDCGVPDNLSVV  214
H. sapiens     165 IGYLRLILPELQA-----RIRTYNQHYN-NLLRGAVSQRLYILLPLDCGVPDNLSMA  215

M. brevicollis 218 E----GLKATEHYISPKPITIGGVVDREMGKHTLYTPSESAETASVAFAMELASPLN  270
D. melanogaster 213 LLD-----KAEPLETQFINRAGVY-RPF-KHDVYRMNKKVNGRTYYFAVEGATPMI  261
N. vectensis   252 DYDPDQNRITFEGNTEPLEKTRGGVFLRHY-KHSVYEILKDGE-NEPWFCIMEYATPLL  306
M. musculus    215 DP----NIRFRDMLPQONIDRAGIKNRVY-SNSVYEILENG-QPAGVCILEYATPLQ  265
H. sapiens     216 DP----NIRFLDKLPQOTGDRAGIKDRVY-SNSIYELLENG-QRAGTCVLEYATPLQ  266

M. brevicollis 271 TMKNALKDAAEGLLEL------QSEAFYLTLRGILKQEGVLDQGNDEGMIKLV-WGK  320
D. melanogaster 262 SFFDATYSNLSGTWQMQELKREIWIKFYKHLKELITTWPET-----RDLVELIIYNS  313
N. vectensis   307 TLYDMSVAQPGELSR-EERDAQVVVFLRKLQDILEGDRAC-----QGKVELVTFSP  356
M. musculus    266 TLFAMSQDAKAGFSR-EDRLEQAKLFCRTLEEILEDVPES-----RNNCRLIVYQE  315
H. sapiens     267 TLFAMSQYSQAGFSR-EDRLEQAKLFCRTLEDILADAPES-----QNNCRLIAYQE  316

M. brevicollis 321 N---------RSEVLKKMFAEDQYAMP-------------------------338
D. melanogaster 314 HDSKGNLVDVGELLVAHMQNKTKTIDEISN-----------------------343
N. vectensis   357 DRDLA------DVMLRKLKDSE---LEIGG-------------------------377
M. musculus    316 PTDGNS-FSLSQEVLRHIRQEEKEEVTMNAPMTSVAPPPSVLSQEPRLLISGMDQPL  371
H. sapiens     317 PADDSS-FSLSQEVLRHLRQEEKEEVTVGSLKTSAVPSTSTMSQEPELLISGMEKPL  372

M. brevicollis     -------
D. melanogaster    -------
N. vectensis       -------
M. musculus    372 PLRTDLI  378
H. sapiens     373 PLRTDFS  379
```

Figure 2 - figure supplement 1

Figure 3

**A** 2'3'cGAMP

**B** 3'3'cGAMP

**C**

**D**

Z-projection | STING | Z-Slice

Control

2'3'cGAMP

Figure 3 - figure supplement 1

Figure 4

Figure 5

Figure 5 - figure supplement 1

Figure 6

Figure 6 - figure supplement 1