Microtubule inhibitors identified through non-biased screening

enhance DNA transfection efficiency by delaying p62-dependent

ubiquitin recruitment

Megumi Tsuchiya[1#], Hidesato Ogawa[1#*], Kento Watanabe[1], Takako Koujin[2], Chie

Mori[2], Kazuto Nunomura[3], Bangzhong Lin[3], Akiyoshi Tani[3], Yasushi Hiraoka[1,2], and

Tokuko Haraguchi[1,2*]

[1]Graduate School of Frontier Biosciences, Osaka University, 1-3 Yamadaoka, Suita

565-0871, Japan

[2]Advanced ICT Research Institute Kobe, National Institute of Information and

Communications Technology, 588-2 Iwaoka, Iwaoka-cho, Nishi-ku, Kobe 651-2492,

Japan

[3]Graduate School of Pharmaceutical Science, Osaka University, 1-6 Yamadaoka, Suita

565-0871, Japan

# These authors contributed equally to this manuscript.

*Correspondence should be addressed to:

Hidesato Ogawa, Tel: +81 668794621; Fax: +81 668794622

E-mail address: hidesato@fbs.osaka-u.ac.jp

Tokuko Haraguchi, Tel: +81 668794621; Fax: +81 668794622

E-mail address: haraguchi@fbs.osaka-u.ac.jp

Short title: MT drugs enhance transfection efficiency

Abstract

Ectopic gene expression is an indispensable tool in biology and medicine, but is often limited by the low efficiency of DNA transfection. We previously reported that depletion of the autophagy receptor p62/SQSTM1 enhances DNA transfection efficiency by preventing the degradation of transfected DNA. Therefore, p62 is a potential target for drugs to increase transfection efficiency. To identify such drugs, a non-biased high-throughput screening was applied to over 4,000 compounds from the Osaka University compound library, and their p62-dependency was evaluated. The top-scoring drugs were mostly microtubule inhibitors, such as colchicine and vinblastine, and all of them showed positive effects only in the presence of p62. To understand the p62-dependent mechanisms, the time required for p62-dependent ubiquitination, which is required for autophagosome formation, was examined using polystyrene beads that were introduced into cells as materials that mimicked transfected DNA. Microtubule inhibitors caused a delay in ubiquitination. Furthermore, the level of phosphorylated p62 at S405 was markedly decreased in the drug-treated cells. These results suggest that microtubule inhibitors inhibit p62-dependent autophagosome formation. Our findings demonstrate for the first time that microtubule inhibitors suppress p62 activation as a mechanism for increasing DNA transfection efficiency and provide solutions to increase efficiency.

Keywords: high-throughput screening; transfection efficiency; autophagy; p62; phosphorylation; ubiquitination; gene delivery

Introduction

Gene delivery is one of the most important steps in gene therapy and genetic

modification in basic science. Gene therapy has great potential in clinical medicine, and

this concept has become well established in therapeutic approaches. In basic science,

DNA transfection is a powerful tool that enables the study of gene functions and their

products in cells.

Endocytosis is a crucial pathway for the regulation of cellular uptake of

plasmid DNA (Rejman *et al.* 2005; Khalil *et al.* 2006; Midoux *et al.* 2008). Importantly,

endocytosis can promote the induction of selective autophagy, also called xenophagy

(Chen *et al.* 2014). Generally, autophagy is a cytosolic bulk degradation pathway for the

recycling of biomolecules through nonspecific degradation of proteins and organelles

under nutrient starvation conditions. In contrast, selective autophagy plays an important

defensive role against cellular infection by pathogens, as part of a starvation-

independent autophagic defense system (Nakatogawa *et al.* 2009; Mizushima *et al.*

2011; Galluzzi *et al.* 2017). The conjugation of ubiquitin (Ub) to target pathogens is an

important initial process in selective autophagy. Ubiquitination assists in the recruitment

of autophagy receptor proteins, including p62/sequestosome-1 (p62/SQSTM1), against

pathogens or transfected DNA (Dupont *et al.* 2010; Roberts *et al.* 2013; Chen *et al.*

2014; Alomairi *et al.* 2015; Tsuchiya *et al.* 2016). Therefore, suppression of the

autophagy pathway can be a target for drugs to increase transfection efficiency.

We have previously reported that the depletion of the p62/SQSTM1 protein

(hereafter designated p62) greatly increases the efficiency of DNA transfection in

cultured cells (Tsuchiya *et al.* 2016). To monitor the behavior of the transfected DNA,

we developed an experimental system using DNA-conjugated beads that mimic the

transfected DNA. In this system, DNA-conjugated beads together with pHrodo, a

fluorescent marker for endosome rupture, were incorporated into cells, and their

intracellular dynamics were analyzed in a live cell (Tsuchiya *et al.* 2016). Using this

system, we demonstrated that the transfected DNA was incorporated into cells through

endocytosis, released from the endosomes into the cytosol, and entrapped via autophagy

in a p62-dependent manner. Furthermore, we demonstrated that the recruitment of Ub

around the transfected material was significantly delayed in p62 gene-knockout murine

embryonic fibroblast (p62KO-MEF) cells compared with that in normal MEF cells

(Tsuchiya *et al.* 2018). Additionally, p62 phosphorylation at S405 (equivalent to human

S403) is a crucial step for the recruitment of Ub to the target site of transfected materials

that mimic ectopic DNA (Tsuchiya *et al.* 2016). Hence, p62 phosphorylation at S405 is

an essential step for transfection-induced selective autophagy (Tsuchiya *et al.* 2018). As

p62 plays an important role in gene delivery through initial ubiquitination of the

transfected DNA, inhibition of p62 may increase transfection efficiency. The aim of this

study was to identify a small chemical compound that blocks the initial step of p62-

dependent selective autophagy to enhance DNA transfection efficiency in mammalian

cells.

Results

High-throughput screening for drugs enhancing transfection efficiency

To identify potential compounds that can enhance transfection efficiency, high-

throughput screening based on a luciferase assay was performed on MEF cells using an

automated workstation (Fig. 1A). The MEF cells were seeded in 384-well plates and

incubated for 18 h with each compound from the Osaka University compound library

(4,400 compounds; Tachibana *et al.* 2018) at a final concentration of 10 µM. The

negative control was dimethyl sulfoxide (DMSO) at a final concentration of 1%. The

cells were transfected with the pCMV-Luc plasmid (encoding luciferase driven by the

cytomegalovirus immediate early (CMV-IE) promoter) for 28 h using Lipofectamine

2000. The cells were subjected to a cell viability assay, followed by a luciferase assay

(Fig. 1A). Luciferase activity, as expressed from the transfected plasmid, was measured,

and transfection efficiency was evaluated based on luciferase activity normalized to cell

viability. Among the 4,400 compounds tested, 160 were removed from the list for

selection as valid compounds because of severe defects in cell morphology. The

transfection efficiency of the cells treated with each of the remaining 4,240 compounds

is plotted in Fig. 1B. In this first screening, out of the 4,240 tested compounds, we

identified 87 compounds that increased luciferase activity compared with that in the

negative control (approximately 2.1% positive hit rate). The cutoff value used for

selection was the mean value of the DMSO control + 4 × standard deviation (SD) (mean

0.347, SD = 0.242).

110   The activities of these 87 compounds were further analyzed via a second

screening. To achieve this, MEF cells were cultured in 96-well plates and transfected

with the pCMV-Luc plasmid for 28 h using Lipofectamine 2000 in the presence of each

of these compounds at a concentration of 1 µM (Fig. 2A). Transfection efficiency was

determined by a luciferase assay, in which luciferase activity expressed from the

transfected plasmid was measured and normalized to cell viability, as described in

Experimental procedures. The transfection efficiency values for the 87 compounds are

listed in Table S1. Notably, the top 9 compounds with high transfection efficiencies (>

20-fold that of the negative control) were all microtubule inhibitors (Fig. 2B; Table S1).

As microtubule inhibitors were found to increase transfection efficiency, we

also tested nocodazole, an extensively used microtubule inhibitor, which was not

included in the Osaka University compound library. Transfection efficiency was

determined in the presence or absence of 1 μM nocodazole in 96-well plates (Fig. S1).

The results show that nocodazole has a transfection-enhancing activity at 1 μM, similar

to colchicine (Fig. S1).

We further examined whether microtubule inhibitors are effective against

various transfection reagents with different mechanisms. Effectene (a non-liposomal

lipid reagent) and Viofectin (a non-lipid high-charge cationic polymer reagent) were

examined in addition to the previously used Lipofectamine 2000 (a cationic lipid

reagent). In the presence of 300 nM colchicine, the transfection efficiencies in all these

transfection reagents increased, although the degree of effectiveness varied (Fig. S2).

The increase in transfection efficiency in the presence of 300 nM colchicine was

approximately 2-, 6-, and 29-fold in Effectene, Viofectin, and Lipofectamine 2000,

respectively (Fig. S2), suggesting that colchicine is effective with various non-viral

transfection reagents.

Microtubule inhibitors enhance gene transfection efficiency

To further evaluate the effects of microtubule inhibitors on DNA transfection efficiency,

we selected two commonly used microtubule inhibitors, colchicine and vinblastine.

These were ranked second and third, respectively, in the 2nd screening (Fig. 2A). First,

we examined the dose-dependency of these inhibitors in MEF cells (Fig. 2C). Cells

were treated with colchicine or vinblastine at various concentrations and transfected

with the pCMV-Luc DNA plasmid. Then, their luciferase activity was measured to

determine transfection efficiency. Transfection efficiency increased in a dose-dependent

manner in the MEF cells (Fig. 2C). Statistical analysis showed that the EC50 (half-

maximal effective concentration) of colchicine and vinblastine was 239.1 and 26.29 nM,

respectively, in the MEF cells (Fig. 2C). Furthermore, we evaluated transfection

efficiency by observing GFP expression in MEF cells. The number of GFP-positive

cells was greater in colchicine- and vinblastine-treated cells than in DMSO-treated

control cells, and cell viability was approximately 70% in both drug-treated cells

compared with that in the control under these conditions (Fig. 2D). These results

suggest that treatment with colchicine and vinblastine can enhance transfection

efficiency.

As microtubule depolymerization activates the transcription factor NF-κB to

induce NF-κB-dependent gene expression (Rosette & Karin, 1995) and the CMV

promoter in the pCMV-Luc plasmid possesses an NF-κB binding site (Rosette & Karin,

1995; Wang & MacDonald, 2004), treatment with microtubule inhibitors may induce

luciferase gene expression through NF-κB-mediated CMV promoter activation. To

eliminate this possibility, we tested promoter activity in a cell line (MEF-LUC cells)

with a CMV-driven luciferase plasmid integrated into the genome. In these cells, we

measured the levels of luciferase activity in the presence or absence of colchicine or

vinblastine. Treatment with colchicine for 27 h increased luciferase gene expression

only marginally (approximately 1.2-fold), which did not explain the considerably

greater increase (approximately 40-fold) in the transfected DNA plasmids (Fig. S3A;

compare with Fig. 2C). Treatment with vinblastine also yielded a similar result, with

almost no increase in luciferase expression (Fig. S3B; compare with Fig. 2C). These

results indicate that the increase in transfection efficiency in the presence of microtubule

inhibitors does not reflect NF-κB-dependent promoter activation.

Transfection enhancement occurs in the presence of autophagy receptor p62

As p62 acts as an inhibitory factor for DNA transfection by recruiting LC3

(microtubule-associated protein 1A/1B-light chain 3; also called ATG8), a marker

protein for autophagosomes, to transfected DNA (Tsuchiya *et al.* 2016), the

transfection-enhancing activity of the tested compounds may be deduced by the

invalidation of the p62-dependent autophagic pathway. Based on this hypothesis, we

examined the activity of the 87 compounds at a concentration of 1 μM in p62KO-MEF

cells under the same conditions as those in Fig. 2A (Fig. 3A). All compounds exhibited

little or no transfection-enhancing activity in p62KO-MEF cells (Fig. 3A). As the

microtubule inhibitors (compounds 1–9 in Fig. 2B) seemed to show a slight increase in

transfection enhancing activity (Fig. 3A), we further tested colchicine and vinblastine

for transfection-enhancing activity at various concentrations from 15.6 to 2000 nM (Fig.

3B, C) and found that these two microtubule inhibitors did not show transfection-

enhancing activity at any of the tested concentrations (Fig. 3B, C). This result suggests

that microtubule inhibitors function in a p62-dependent autophagic pathway.

p62-dependent ubiquitination is delayed by microtubule inhibitors

To understand the molecular mechanisms of the p62-dependent enhancement of

transfection efficiency by colchicine and vinblastine, we employed an experimental

method using polystyrene beads that had been developed to monitor the behavior of the

transfected DNA (Kobayashi *et al.* 2010) (Fig. 4A). In this method, the beads are

incorporated into cells with transfection reagents via endocytosis and enter the cytosol

after rupture of the endosomal membrane. The beads that appeared in the cytosol were

targeted for autophagy, similar to the transfected DNA (Kobayashi *et al.* 2010;

Kobayashi *et al.* 2015) (Fig. 4A). For monitoring in living cells, the beads were pre-

conjugated with pHrodo dye, which emits fluorescence under acidic pH conditions,

such as those in the acidic endosome but not in the cytosol. Therefore, this dye serves as

a marker of endosome membrane rupture, as previously described (Kobayashi *et al.*

2010). The pHrodo-conjugated beads were incorporated into MEF cells expressing

GFP-fused Ub protein (GFP-Ub MEF cells) (Tsuchiya *et al.* 2018), and the assembly of

GFP-Ub around the beads was observed in a living cell using time-lapse fluorescence

microscopy (Fig. 4B). In the control DMSO-treated cells, the time for Ub recruitment to

the beads was approximately 3–4 min (Fig. 4B, middle panel) after the disappearance of

pHrodo fluorescence (Fig. 4B, upper panel). In contrast, the time for GFP-signal

accumulation was 9–10 min in 500 nM colchicine-treated MEF cells (Fig. 4C, middle

panel), which was longer than that in the control cells (Fig. 4B). Statistical analysis was

performed to determine the timing of GFP-signal accumulation around the beads after

the loss of pHrodo signals in cells expressing GFP-Ub with or without the inhibitor

(Fig. 4D). In the DMSO-treated cells, the time for Ub recruitment to the beads was ~4

min (median) after the disappearance of pHrodo fluorescence (mean and SD, 4.167 ±

1.88 min, n = 24 beads; lane 1 in Fig. 4D). Moreover, the timing of GFP-signal

accumulation was ~6 min (median) in the 100 nM colchicine-treated MEF cells (mean

and SD: 8.286 ± 8.36 min, n = 28 beads: lane 2 in Fig. 4D), ~11 min (median) in the

500 nM colchicine-treated MEF cells (mean and SD: 11.32 ± 11.30 min, n = 22 beads:

lane 3 in Fig. 4D), and ~6 min (median) in the vinblastine-treated MEF cells (mean and

SD: 6.48 ± 3.81 min, $n = 31$ beads; lane 4 in Fig. 4D). Additionally, the GFP-Ub signals

did not accumulate within 60 min in colchicine- or vinblastine-treated GFP-Ub MEF

cells (these beads were not counted and were not included in the n; upper column in Fig.

4D). These results show that GFP-Ub accumulation around the beads was significantly

delayed in colchicine- or vinblastine-treated MEF cells. Colchicine and vinblastine have

different effects on microtubule depolymerization; colchicine forms diffusely

distributed tubulins/microtubules, whereas vinblastine forms the paracrystal structure of

the microtubules (Haraguchi *et al.* 1997). Nonetheless, both have the same function of

increasing DNA transfection efficiency, suggesting that normal microtubule function or

structure is important for Ub recruitment to the target site.

Active form of p62 is suppressed by microtubule inhibitors

It has been reported that the phosphorylated form of p62 at amino acid residue S405

(p62 S405) is required for Ub recruitment during selective autophagy (Tsuchiya *et al.*

2018). Therefore, Ub recruitment can be delayed by a decrease in the level of

phosphorylated p62 at S405. To test this hypothesis, we performed Western blot

analysis to evaluate the total p62 protein levels and p62 S405 phosphorylation levels

(Fig. 5). The p62 S405 phosphorylation levels were very low before DNA transfection

("no transfection" in Fig. 5). However, after transfection ("DNA transfection 24 h"),

these levels greatly increased in DMSO-treated MEF cells, although the p62 levels

remained unchanged. This suggests that DNA transfection induced an increase in p62

S405 phosphorylation. However, in MEF cells treated with colchicine or vinblastine,

the level of p62 S405 phosphorylation decreased. This suggests that microtubule

inhibitors inhibit Ub recruitment by decreasing the level of phosphorylated p62 at S405.

Discussion

Microtubule structure/function and transfection efficiency

In this study, we used a non-biased, high-throughput screening approach to identify

small chemical compounds that increase DNA transfection efficiency and identified

microtubule inhibitors as the top 9 compounds. Therefore, inhibition of the microtubule

structure or function seems to trigger an increase in transfection efficiency. This finding

is consistent with previous reports that microtubule depolymerizing reagents,

nocodazole, increased transfection efficiency in CV-1 cells (Lindberg *et al.* 2001) and

colchicine and vinblastine increased transfection efficiency in cultured vascular smooth

muscle cells (Wang & MacDonald 2004). Additionally, microtubule-polymerizing

reagents, such as paclitaxel, also increase the transfection efficiency in COS-7 and A549

cells (Hasegawa *et al.* 2001; Nair *et al.* 2002). The polymerizing agent docetaxel, a

chemical compound closely related to paclitaxel, was included in our screening results

for the top 9 compounds (Fig. 2B, Rank 7). Moreover, tubulin deacetylation inhibitors,

such as HDAC6 inhibitors, also increase transfection efficiency in A549 cells, PC3

cells, and mesenchymal stem cells (Vaughan *et al.* 2008; Barua & Rege 2010; Ho *et al.*

2017). This is likely because acetylated tubulin can stabilize microtubule structures,

which in turn implies that microtubule structure stabilization can also increase gene

transfection efficiency. Taken together, these findings strongly suggest that inhibition of

the intact structure or dynamic nature of microtubules is important for increasing the

efficiency of gene transfection.

Microtubule inhibitors affect selective autophagy pathways

Our results showed that treatment of cells with colchicine or vinblastine increased

transfection efficiency in a p62-dependent manner. However, it is difficult to attribute

these observations to a particular cause. Microtubules are major contributors to the

trafficking of several components in the endosomal/lysosomal pathway; hence, it is

logical to propose that these inhibitors may block the autophagy pathway. The role of

microtubules in autophagosome formation appears to be different based on the culture

medium conditions, e.g., vegetatively growing medium (basal) or starvation medium

(inducible) conditions. Several studies have used microtubule inhibitors under basal

conditions to show that microtubules do not participate in autophagosome formation

(Reunanen *et al.* 1988; Aplin *et al.* 1992; Kochl *et al.* 2006). However, under inducible

conditions, disassembling the microtubules with these inhibitors prevented

autophagosome formation, suggesting that microtubules play a crucial role in this

process (Kochl *et al.* 2006; Geeraert *et al.* 2010). Previous studies have shown that

transfected DNA also induces the selective autophagy pathway (Roberts *et al.* 2013;

Chen *et al.* 2014; Tsuchiya *et al.* 2016); hence, microtubule inhibitors may affect

transfection-induced autophagosome formation. The detailed mechanisms of the

inhibition of autophagosome formation following microtubule inhibitor treatment are

not clear; however, several autophagy factors are associated with microtubules,

including ATG8 (LC3), ATG1 (ULK1), ATG6 (Beclin1), ATG18 (WIPI1), and p62

(Mackeh *et al.* 2013). LC3 has long been thought to be involved in the regulation of

microtubule assembly and disassembly (Mann & Hammarback 1994; Deretic 2008).

Furthermore, ATG18-positive pre-autophagosomal structures can move along

microtubules, and this movement is extremely sensitive to microtubule inhibitor

treatment (Polson *et al.* 2010). These data suggest that microtubules contribute to the

sequestration, recruitment, and movement of autophagy factors for the formation of

inducible autophagosomes. Therefore, microtubule inhibitors may block the inducible

autophagy pathway, resulting in a decrease in the likelihood of DNA degradation and

increase in transfection efficiency.

Microtubule inhibitors decrease the level of phosphorylated p62

The ubiquitination of endosome membrane proteins surrounding exogenous material,

such as transfected DNA, is the initial step in inducible selective autophagy (Randow

2011; Wileman 2013; Tsuchiya *et al.* 2016). In our study, treatment with colchicine or

vinblastine delayed the recruitment of Ub proteins. This delay in recruitment is also

caused by the depletion of p62 or mutation of the p62 phosphorylation site at S405

(human S403) (Tsuchiya *et al.* 2018). As p62 has been reported as a microtubule-

associated factor (Geeraert *et al.* 2010), treatment with a microtubule inhibitor may

affect p62-mediated Ub recruitment. Specifically, our results showed that p62 S405

phosphorylation was significantly impaired by treatment with colchicine or vinblastine

after transfection. p62 S405 is phosphorylated by kinases such as CK2 and TBK1

(Matsumoto *et al.* 2011; Matsumoto *et al.* 2015). These kinases are also associated with

tubulin (Faust *et al.* 1999; Lim *et al.* 2004; Pillai *et al.* 2015; Yang *et al.* 2018). After

transfection, the recruitment of these kinases may be affected by treatment with

microtubule inhibitors. Further studies are necessary to elucidate the detailed

mechanisms by which these microtubule inhibitors affect the phosphorylation levels of

p62.

308   In this study, a non-biased high-throughput screening method revealed that

microtubule inhibitors are the most effective reagents for increasing transfection

efficiency among the drugs present in the Osaka University compound library, and

revealed for the first time, the mechanism by which microtubule inhibitors increase

transfection efficiency; that is, microtubule inhibitors suppress p62 phosphorylation,

which is required for autophagosome formation to degrade transfected DNA. Our

findings provide solutions to increase transfection efficiency.

Experimental procedures

Plasmids

pCMV-Luc plasmid encoding the luciferase gene *luc2*, used as a luciferase expression

vector in this study, was purchased from Promega (E1310; pGL4.50 [*luc2*/CMV/Hygro]

vector; Promega, Madison, WI, USA). The GFP expression plasmid pCMX-AFAP was

prepared as previously described (Ogawa *et al.* 2004).

To create cells stably expressing luciferase (see the next section), we constructed

the PB-CMV-LUC-Zeo vector. We first constructed the PB-EF1-MCS-IRES-Zeo

vector. To insert the zeocin resistance gene DNA sequence, two DNA fragments (#1

and #2) were amplified as follows: the cloning site with Kozak sequence fragment #1

was amplified from the PB-EF1-MCS-IRES-Neo vector (PB533A-2; System

Biosciences, Palo Alto, CA, USA) using PCR and the following primers: 5′-

CTGAAGGATGCCCAGAAGGTACCCCATTGT-3′ and 5′-

TCCGGACGCCATGGTTGTGG-3′. The zeocin coding fragment #2 was amplified

from the pcDNA3.1/Zeo (+) vector (V86020; Thermo Fisher Scientific, Yokohama,

Japan) using PCR and the following primers: 5′-

ACAACCATGGCGTCCGGAATGGCCAAGTTGACCAGTGCCGTTCC-3′ and 5′-

TCCAGAGGTTGATTGTCGACTCAGTCCTGCTCCTCGGCCACGAA-3′.

Following digestion with KpnI and SalI, fragments #1 and #2 were inserted into the PB-

EF1-MCS-IRES-Neo vector using the In-Fusion HD Cloning Kit (639648; Takara Bio

Inc. Kusatsu, Japan). This resulted in the PB-EF1-MCS-IRES-Zeo vector. Next, the

human cytomegalovirus immediate early enhancer and promoter (CMV-IE) DNA

sequence (#3) was amplified from the pGL4.50 [*luc2*/CMV/Hygro] vector using PCR

and the following primers: 5′-

GGGGATACGGGGAAAAGGCCTCGTTACATAACTTACGGTAAATG-3′ and 5′-

GAATTCGCTAGCTCTAGAAGCTCTGCTTATATAGACCTCCCACC-3′. The

luciferase DNA sequence (#4) was amplified from the pGL4.50 [*luc2*/CMV/Hygro]

vector using PCR and the following primers: 5′-

TCTAGAGCTAGCGAATTCATGGAAGATGCCAAAAACATTAAGAA-3′ and 5′-

CGATTTAAATTCGAATTCTTACACGGCGATCTTGCCGCCCTTCT-3′. Following

digestion with EcoRI and StuI, fragments #3 and #4 were inserted into the PB-EF1-

MCS-IRES-Zeo vector using the In-Fusion HD Cloning Kit. This resulted in the PB-

CMV-LUC-Zeo vector.

Cell strains

p62KO-MEF cells (p62$^{-/-}$ cells) and their parental MEF cells were kindly provided by

Dr. Tetsuro Ishii (University of Tsukuba) (Komatsu *et al.* 2007). MEF cells stably

expressing GFP-Ub were generated as previously described (Tsuchiya *et al.* 2018). To

obtain MEF cells or p62KO-MEF cells stably expressing luciferase, MEF cells or

p62KO-MEF cells were transfected with the PB-CMV-LUC-Zeo plasmid (generated as

described in the previous section) and cultured in the presence of 100 μg/mL Zeocin

(R25501, Thermo Fisher Scientific), and then, single clones (MEF-LUC cells) were

selected. Each stable clone was examined for luciferase protein expression using a

luciferase assay.

Cell culture

All cell lines were maintained in culture medium (Dulbecco's modified Eagle medium

(DMEM) (D6429; Sigma-Aldrich, St. Louis, MO, USA) supplemented with 10% fetal

bovine serum) in the presence of 5% CO_2 at 37°C.

High-throughput screening for enhancer compounds

The MEF cells were seeded at a density of 0.8×10^3 cells per well (384-well microplate,

781091; Greiner Bio-One, Tokyo, Japan) using a Multidrop COMBI (Thermo Fisher

Scientific) and cultured for 6 h in culture medium. The cells were pretreated for 18 h

with 1% DMSO (negative control) or the screening compounds (10 μM each) using a

Fluent780® Automation Workstation (Tecan Japan, Kawasaki, Japan) with a 96-

channel head adapter and Tecan sterile tips (30048824; Tecan Japan). The pretreated

cells in each well were transfected with 25 ng of pCMV-Luc plasmid using

Lipofectamine 2000 (Thermo Fisher Scientific) for 28 h in the presence of each

compound. Cell viability was measured with the RealTime-Glo™ MT Cell Viability

Assay (E9713; Promega) using the GloMax® Discover Microplate Reader (Promega).

Luciferase activity was measured with the ONE-Glo™ Luciferase Assay System

(E6120; Promega) using the GloMax® Discover Microplate Reader according to the

manufacturer's protocol. The values of luciferase activity were normalized to the cell

number obtained from cell viability measurements.

Transfection efficiency measurements

Transfection efficiency was determined based on the luciferase activity expressed from

the transfected plasmid. Cells were seeded at a density of 0.45×10^4 cells per well (96-

well assay plate; 3603 Corning, Corning, NY, USA) and cultured overnight in culture

medium. The cells were pretreated with 1% DMSO (negative control) or the screening

compounds (0.1–10 μM each) for 30 min. The cells were then transfected with 100 ng

of pCMV-Luc plasmid using Lipofectamine 2000 for 28 h in the presence of each

compound. Cell viability was measured with the RealTime-Glo™ MT Cell Viability

Assay using the GloMax® 96 Microplate Luminometer (Promega). Luciferase activity

was measured with the ONE-Glo™ Luciferase Assay System using the GloMax® 96

Microplate Luminometer according to the manufacturer's protocol. The values of

luciferase activity were normalized to the cell number obtained from cell viability

measurements. Mean EC50 values and SDs were determined from three independent

experiments. Transfection efficiencies were determined by luciferase activity

normalized to cell viability.

Viofectin (VFT1001; Viogene, Nacalai tesque, Kyoto, Japan) and Effectene

(301425; Qiagen, Tokyo, Japan) were also used as transfection reagents. For Viofectin,

the same transfection procedure as that described above for Lipofectamine 2000 was

used, except that the reagents were different. For Effectene, some procedures were

modified as follows: initial cell density was 1×10^4 cells per well and the cells were

transfected for 4 h and further incubated for another 20 h in culture medium without

transfection reagents.

GFP expression

MEF cells were seeded at a density of 1.5×10^5 cells per well (6-well assay plate; 3335

Corning, Corning, NY, USA) and cultured overnight in culture medium. The cells were

pretreated with 1% DMSO (negative control), 250 nM cholchicine, or 50 nM

vinblastine for 30 min. The cells were then transfected with 2 μg of pCMV-Luc plasmid

using Lipofectamine 2000 for 24 h in the presence of each drug. The cells were

observed using an objective lens (N Plan 5x/0.12 PH 0) on a Leica DM IL LED

fluorescence microscope (Leica Microsystems, Wetzlar, Germany).

Preparation of pHrodo-conjugated beads

pHrodo-conjugated beads were prepared as previously described (Kobayashi *et al.*

2010). Briefly, Dynabeads M-270 Streptavidin (DB65306; Thermo Fisher Scientific)

were washed three times with phosphate buffered saline (PBS) and resuspended in 100

mM sodium bicarbonate buffer (pH 8.5) to an appropriate concentration (typically a

1:10 or 1:20 dilution). pHrodo-succinimidyl ester (P36600; Thermo Fisher Scientific)

was then added to the bead suspension and incubated in sodium bicarbonate buffer for 1

h at room temperature (about 26°C). After the conjugation reaction, the beads were

washed with sodium bicarbonate buffer and suspended in PBS.

Incorporation of beads into living cells

One day before incorporating the beads, GFP-Ub MEF cells were seeded onto 35 mm

glass-bottom culture dishes (P35G-1.5-10-C; MatTek, Ashland, MA, USA) at a density

of 1.5×10^5 cells/dish in culture medium. The cells were pretreated with each

microtubule inhibitor (100 nM colchicine, 500 nM colchicine, or 100 nM vinblastine) or

DMSO as a control for 30 min at 37°C in a CO_2 incubator before bead incorporation.

The transfection reagent Effectene (301425; Qiagen, Tokyo, Japan) was used to

incorporate the beads. Transfection-reagent-coated beads were prepared by mixing

pHrodo-conjugated beads with the Effectene transfection reagent according to the

manufacturer's protocol, except that a bead suspension was used instead of the DNA

solution. The resulting bead mixture (~10 μL) was mixed with 90 μL of the culture

medium containing each microtubule inhibitor and added to the cells by replacing the

medium. After incubation for 1 h at 37°C in a CO_2 incubator, the cells were washed

twice with fresh growth medium to remove unattached beads and then further incubated

with the medium containing each microtubule inhibitor or control DMSO for the time

indicated in each experiment.

Time-lapse imaging

Cells were treated with 100 ng/mL Hoechst 33342 (B2261; Sigma-Aldrich) for 15 min

to stain chromosomes, as previously described (Haraguchi *et al.* 1997), except that the

medium containing each microtubule inhibitor or control DMSO was used. After

washing with the culture medium, the cells were cultured in fresh culture medium

(phenol red-free DMEM containing 10% fetal bovine serum) containing each

microtubule inhibitor or control DMSO. Time-lapse observations were performed using

an oil-immersion objective lens (UApo40/NA1.35; Olympus, Tokyo, Japan) on a

DeltaVision microscope system (GE Healthcare Life Sciences Japan, Tokyo, Japan)

placed in a temperature-controlled room (37°C), as previously described (Haraguchi *et*

al. 1997). Images were obtained every minute for approximately 60 min.

The timing of GFP-Ub fluorescence signal accumulation around the beads was

determined as previously described (Tsuchiya *et al.* 2018). Briefly, the fluorescence

intensity of the region (18 pixels square) surrounding the beads was measured using the

FUJI software suite (IMAGE J; National Institutes of Health, Bethesda, MD, USA), and

the background fluorescence intensity of the region with no beads in the same cell was

subtracted.

Western blot analysis

MEF cells were seeded at a density of 1.5×10^5 cells per well (6-well assay plate; 3335

Corning, Corning, NY, USA) and cultured overnight in culture medium. The cells were

pretreated with 1% DMSO (negative control), 500 nM colchicine, or 100 nM

vinblastine for 30 min. The cells were then transfected with 2 μg of pCMX-AFAP

plasmid using Lipofectamine 2000 for 24 h in the presence of each drug. Western blot

analysis was performed as previously described (Tsuchiya *et al.* 2018). Briefly, cell

lysates were prepared in a lysis buffer [50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1 mM

EDTA, 1% Triton X-100, 1 x Phosphatase Inhibitor Cocktail Solution Ⅱ (160-24371;

FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan) and 1 x protease inhibitor

cocktail (Nacalai tesque Inc., Kyoto, Japan)]. The lysates were subjected to

electrophoresis on NuPAGE 4% to 12% Bis-Tris gels (NP0321; Thermo Fisher

Scientific). Proteins were transferred to polyvinylidene fluoride membranes and probed

using anti-p62 (SQSTM1) (PM045; MBL, Nagoya, Japan), anti-Phospho-

SQSTM1/p62(Ser403) (D8D6T; Cell Signaling Technology, Danvers, MA, USA), and

anti-glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (14C10; Cell Signaling

Technology) antibodies, and secondary antibody conjugated to horseradish peroxidase

(NA9340V; GE Healthcare Life Sciences). Protein bands were stained with

ImmunoStar Zeta (295-72404; FUJIFILM Wako Pure Chemical Corporation) and

detected by chemiluminescence using a ChemiDoc MP imaging system (Bio-Rad,

Tokyo, Japan).

Statistical analysis

The *p*-values were obtained by performing Kruskal-Wallis tests using GraphPad Prism

8 software (GraphPad Software, Inc., La Jolla, CA, USA).

Acknowledgments

This work is dedicated to the late Professor Tomoko Ogawa. We are grateful to Dr. Eiji

Warabi and Dr. Tetsuro Ishii (Tsukuba University) for providing p62KO-MEF cells and

the corresponding wild-type MEF cells. We thank J. Iacona, Ph.D., from Edanz Group

(https://en-author-services.edanzgroup.com/ac) for English-editing a draft of this

manuscript. This study was supported by The JSPS Kakenhi Grant Numbers

JP19K06488 to MT, JP19KK0218, JP20H05322 and JP20K07029 to HO, JP18H05533

and JP20H00454 to YH, and JP17K19505 and JP18H05528 to TH. This research was

supported by Platform Project for Supporting Drug Discovery and Life Science

Research (Basis for Supporting Innovative Drug Discovery and Life Science Research

(BINDS)) from AMED under Grant Number JP20am0101084.

Author contributions

MT, HO, KW, TK, CM, KN, BL and AT performed the experiments. MT, HO, AT,

YH, and TH designed the experiments. All authors analyzed and discussed the data, and

MT, HO, YH, and TH wrote the manuscript.

Conflict of interest

All authors declare that: (i) no support, financial or otherwise, has been received from

any organization that may have an interest in the submitted work; and (ii) there are no

other relationships or activities that could appear to have influenced the submitted work.

References

Alomairi, J., Bonacci, T., Ghigo, E. & Soubeyran, P. (2015) Alterations of host cell
ubiquitination machinery by pathogenic bacteria. *Frontiers in cellular and*
infection microbiology **5**, 17. https://doi.org/10.3389/fcimb.2015.00017

Aplin, A., Jasionowski, T., Tuttle, D.L., Lenk, S.E. & Dunn, W.A., Jr. (1992)
Cytoskeletal elements are required for the formation and maturation of autophagic
vacuoles. *J Cell Physiol* **152**, 458-466. https://doi.org/10.1002/jcp.1041520304

Barua, S. & Rege, K. (2010) The influence of mediators of intracellular trafficking on
transgene expression efficacy of polymer-plasmid DNA complexes. *Biomaterials*
31, 5894-5902. https://doi.org/10.1016/j.biomaterials.2010.04.007

Chen, X., Khambu, B., Zhang, H., Gao, W., Li, M., Chen, X., Yoshimori, T. & Yin,
X.M. (2014) Autophagy induced by calcium phosphate precipitates targets
damaged endosomes. *Journal of Biological Chemistry* **289**, 11162-11174.
https://doi.org/10.1074/jbc.m113.531855

Deretic, V. (2008) Autophagosome and phagosome. *Methods Mol Biol* **445**, 1-10.
https://doi.org/10.1007/978-1-59745-157-4_1

Dupont, N., Temime-Smaali, N. & Lafont, F. (2010) How ubiquitination and autophagy
participate in the regulation of the cell response to bacterial infection. *Biology of*
the Cell **102**, 621-634. https://doi.org/10.1042/BC20100101

Faust, M., Schuster, N. & Montenarh, M. (1999) Specific binding of protein kinase CK2
catalytic subunits to tubulin. *FEBS letters* **462**, 51-56.
https://doi.org/10.1016/S0014-5793(99)01492-1

Galluzzi, L., Baehrecke, E.H., Ballabio, A. *et al.* (2017) Molecular definitions of
autophagy and related processes. *The EMBO journal* **36**, 1811-1836.
https://doi.org/10.15252/embj.201796697

Geeraert, C., Ratier, A., Pfisterer, S.G., Perdiz, D., Cantaloube, I., Rouault, A.,
Pattingre, S., Proikas-Cezanne, T., Codogno, P. & Pous, C. (2010) Starvation-
induced hyperacetylation of tubulin is required for the stimulation of autophagy by
nutrient deprivation. *J Biol Chem* **285**, 24184-24194.
https://doi.org/10.1074/jbc.M109.091553

Haraguchi, T., Kaneda, T. & Hiraoka, Y. (1997) Dynamics of chromosomes and

microtubules visualized by multiple-wavelength fluorescence imaging in living

mammalian cells: effects of mitotic inhibitors on cell cycle progression. *Genes to*

Cells **2**, 369-380. https://doi.org/10.1046/j.1365-2443.1997.1280326.x

Hasegawa, S., Hirashima, N. & Nakanishi, M. (2001) Microtubule involvement in the

intracellular dynamics for gene transfection mediated by cationic liposomes. *Gene*

Ther **8**, 1669-1673. https://doi.org/10.1038/sj.gt.3301573.

Ho, Y.K., Zhou, L.H., Tam, K.C. & Too, H.P. (2017) Enhanced non-viral gene delivery

by coordinated endosomal release and inhibition of beta-tubulin deactylase.

Nucleic Acids Research **45**, e38. https://doi.org/10.1093/nar/gkw1143

Khalil, I.A., Kogure, K., Akita, H. & Harashima, H. (2006) Uptake pathways and

subsequent intracellular trafficking in nonviral gene delivery. *Pharmacol Rev* **58**,

32-45. https://doi.org/10.1124/pr.58.1.8

Kobayashi, S., Kojidani, T., Osakada, H., Yamamoto, A., Yoshimori, T., Hiraoka, Y. &

Haraguchi, T. (2010) Artificial induction of autophagy around polystyrene beads in

nonphagocytic cells. *Autophagy* **6**, 36-45. https://doi.org/10.4161/auto.6.1.10324

Kobayashi, S., Koujin, T., Kojidani, T., Osakada, H., Mori, C., Hiraoka, Y. &

Haraguchi, T. (2015) BAF is a cytosolic DNA sensor that leads to exogenous DNA

avoiding autophagy. *Proceedings of the National Academy of Sciences of the*

United States of America **112**, 7027-7032.

https://doi.org/10.1073/pnas.1501235112

Kochl, R., Hu, X.W., Chan, E.Y. & Tooze, S.A. (2006) Microtubules facilitate

autophagosome formation and fusion of autophagosomes with endosomes. *Traffic*

(Copenhagen, Denmark) **7**, 129-145. https://doi.org/10.1111/j.1600-

0854.2005.00368.x

Komatsu, M., Waguri, S., Koike, M. *et al.* (2007) Homeostatic levels of p62 control

cytoplasmic inclusion body formation in autophagy-deficient mice. *Cell* **131**, 1149-

1163. https://doi.org/10.1016/j.cell.2007.10.035

Lim, A.C., Tiu, S.Y., Li, Q. & Qi, R.Z. (2004) Direct regulation of microtubule

dynamics by protein kinase CK2. *J Biol Chem* **279**, 4433-4439.

https://doi.org/10.1074/jbc.M310563200

Lindberg, J., Fernandez, M.A., Ropp, J.D. & Hamm-Alvarez, S.F. (2001) Nocodazole
treatment of CV-1 cells enhances nuclear/perinuclear accumulation of lipid-DNA
complexes and increases gene expression. *Pharmaceutical Research* **18**, 246-249.
https://doi.org/10.1023/a:1011001022570

Mackeh, R., Perdiz, D., Lorin, S., Codogno, P. & Pous, C. (2013) Autophagy and
microtubules - new story, old players. *Journal of Sell Science* **126**, 1071-1080.
https://doi.org/10.1242/jcs.115626

Mann, S.S. & Hammarback, J.A. (1994) Molecular characterization of light chain 3. A
microtubule binding subunit of MAP1A and MAP1B. *J Biol Chem* **269**, 11492-
11497. https://doi.org/10.1016/S0021-9258(19)78150-2

Matsumoto, G., Shimogori, T., Hattori, N. & Nukina, N. (2015) TBK1 controls
autophagosomal engulfment of polyubiquitinated mitochondria through
p62/SQSTM1 phosphorylation. *Human Molecular Genetics* **24**, 4429-4442.
https://doi.org/10.1093/hmg/ddv179

Matsumoto, G., Wada, K., Okuno, M., Kurosawa, M. & Nukina, N. (2011) Serine 403
phosphorylation of p62/SQSTM1 regulates selective autophagic clearance of
ubiquitinated proteins. *Molecular Cell* **44**, 279-289.
https://doi.org/10.1016/j.molcel.2011.07.039

Midoux, P., Breuzard, G., Gomez, J.P. & Pichon, C. (2008) Polymer-based gene
delivery: a current review on the uptake and intracellular trafficking of polyplexes.
Curr Gene Ther **8**, 335-352. https://doi.org/10.2174/156652308786071014

Mizushima, N., Yoshimori, T. & Ohsumi, Y. (2011) The role of Atg proteins in
autophagosome formation. *Annu Rev Cell Dev Biol* **27**, 107-132.
https://doi.org/10.1146/annurev-cellbio-092910-154005

Nair, R.R., Rodgers, J.R. & Schwarz, L.A. (2002) Enhancement of transgene expression
by combining glucocorticoids and anti-mitotic agents during transient transfection
using DNA-cationic liposomes. *Mol Ther* **5**, 455-462.
https://doi.org/10.1006/mthe.2002.0567

Nakatogawa, H., Suzuki, K., Kamada, Y. & Ohsumi, Y. (2009) Dynamics and diversity
in autophagy mechanisms: lessons from yeast. *Nature Reviews. Molecular Cell*
Biology **10**, 458-467. https://doi.org/10.1038/nrm2708

Ogawa, H., Yu, R. T., Haraguchi, T., Hiraoka, Y., Nakatani, Y., Morohashi, K., &
Umesono, K. (2004). Nuclear structure-associated TIF2 recruits glucocorticoid
receptor and its target DNA. *Biochem Biophys Res Commun* **320**, 218–225.
https://doi.org/10.1016/j.bbrc.2004.05.161

Pillai, S., Nguyen, J., Johnson, J., Haura, E., Coppola, D. & Chellappan, S. (2015) Tank
binding kinase 1 is a centrosome-associated kinase necessary for microtubule
dynamics and mitosis. *Nature Communications* **6**, 10072.
https://doi.org/10.1038/ncomms10072

Polson, H.E., de Lartigue, J., Rigden, D.J., Reedijk, M., Urbe, S., Clague, M.J. &
Tooze, S.A. (2010) Mammalian Atg18 (WIPI2) localizes to omegasome-anchored
phagophores and positively regulates LC3 lipidation. *Autophagy* **6**, 506-522.
https://doi.org/10.4161/auto.6.4.11863

Randow, F. (2011) How cells deploy ubiquitin and autophagy to defend their cytosol
from bacterial invasion. *Autophagy* **7**, 304-309
https://doi.org/10.4161/auto.7.3.14539

Rejman, J., Bragonzi, A. & Conese, M. (2005) Role of clathrin- and caveolae-mediated
endocytosis in gene transfer mediated by lipo- and polyplexes. *Mol Ther* **12**, 468-
474. https://doi.org/10.1016/j.ymthe.2005.03.038

Reunanen, H., Marttinen, M. & Hirsimaki, P. (1988) Effects of griseofulvin and
nocodazole on the accumulation of autophagic vacuoles in Ehrlich ascites tumor
cells. *Exp Mol Pathol* **48**, 97-102. https://doi.org/10.1016/0014-4800(88)90048-2

Roberts, R., Al-Jamal, W.T., Whelband, M., Thomas, P., Jefferson, M., van den
Bossche, J., Powell, P.P., Kostarelos, K. & Wileman, T. (2013) Autophagy and
formation of tubulovesicular autophagosomes provide a barrier against nonviral
gene delivery. *Autophagy* **9**, 667-682. https://doi.org/10.4161/auto.23877

Rosette, C., & Karin, M. (1995). Cytoskeletal control of gene expression:
depolymerization of microtubules activates NF-kappa B. *The Journal of cell*
biology, **128**, 1111–1119. https://doi.org/10.1083/jcb.128.6.1111

Tachibana, K., Yuzuriha, T., Tabata, R. *et al.* (2018) Discovery of peroxisome
proliferator-activated receptor alpha (PPARalpha) activators with a ligand-
screening system using a human PPARalpha-expressing cell line. *J Biol Chem* **293**,
10333-10343. https://doi.org/10.1074/jbc.ra118.002077

Tsuchiya, M., Ogawa, H., Koujin, T., Kobayashi, S., Mori, C., Hiraoka, Y. &
Haraguchi, T. (2016) Depletion of autophagy receptor p62/SQSTM1 enhances the
efficiency of gene delivery in mammalian cells. *FEBS letters* **590**, 2671-2680.
https://doi.org/10.1002/1873-3468.12262
Tsuchiya, M., Ogawa, H., Koujin, T., Mori, C., Osakada, H., Kobayashi, S., Hiraoka, Y.
& Haraguchi, T. (2018) p62/SQSTM1 promotes rapid ubiquitin conjugation to
target proteins after endosome rupture during xenophagy. *FEBS Open Bio* **8**, 470-
480. https://doi.org/10.1002/2211-5463.12385
Vaughan, E.E., Geiger, R.C., Miller, A.M., Loh-Marley, P.L., Suzuki, T., Miyata, N. &
Dean, D.A. (2008) Microtubule acetylation through HDAC6 inhibition results in
increased transfection efficiency. *Mol Ther* **16**, 1841-1847.
https://doi.org/10.1038/mt.2008.190
Wang, L. & MacDonald, R.C. (2004) Effects of microtubule-depolymerizing agents on
the transfection of cultured vascular smooth muscle cells: enhanced expression
with free drug and especially with drug-gene lipoplexes. *Mol Ther* **9**, 729-737.
https://doi.org/10.1016/j.ymthe.2004.02.009
Wileman, T. (2013) Autophagy as a defence against intracellular pathogens. *Essays in*
Biochemistry **55**, 153-163. https://doi.org/10.1042/bse0550153
Yang, H., Mao, W., Rodriguez-Aguayo, C., Mangala, L.S., Bartholomeusz, G., Iles,
L.R., Jennings, N.B., Ahmed, A.A., Sood, A.K., Lopez-Berestein, G., Lu, Z. &
Bast, R.C., Jr. (2018) Paclitaxel Sensitivity of Ovarian Cancer Can be Enhanced by
Knocking Down Pairs of Kinases that Regulate MAP4 Phosphorylation and
Microtubule Stability. *Clin Cancer Res* **24**, 5072-5084.
https://doi.org/10.1158/1078-0432.ccr-18-0504
654

Figure legends

Fig. 1. High-throughput screening for drugs enhancing transfection efficiency

(A) Schematic diagram of the process of high-throughput screening using automated workstations. Murine embryonic fibroblast (MEF) cells were treated with each compound and then transfected with the pCMV-Luc plasmid in the presence of each compound. Cell viability and the luciferase activity from the transfected plasmid were measured. Transfection efficiency was evaluated based on luciferase activity normalized to cell viability. (B) High-throughput screening was performed for 4,400 compounds from the Osaka University compound library as described. Values of transfection efficiency were plotted for the selected 4,240 compounds. Dimethyl sulfoxide (DMSO) was used as the negative control. The red broken line represents the mean $+ 4 \times$ SD of all negative control assay points.

Fig. 2. Second screening of compounds selected through high-throughput screening

(A) The top 87 compounds identified in the primary screen were further examined. MEF cells were transfected with the pCMV-Luc plasmid for 28 h in the presence of 1 μM of each compound. DMSO was used as the negative control. Transfection efficiency was determined based on luciferase activity normalized to the cell viability. Bar graphs show fold changes in transfection efficiency relative to the value from the control DMSO-treated cells in descending order. The second and third columns represent data for colchicine and vinblastine as indicated. (B) The top 9 potential compounds for enhancement of transfection efficiency. (C) MEF cells were transfected with the

pCMV-Luc plasmid for 28 h in the presence of colchicine (left graph) or vinblastine

(right graph) at the indicated concentrations. Transfection efficiency was determined as

described above. Filled circles and vertical lines indicate the mean and SD, respectively,

of at least three independent experiments. (D) Control DMSO-treated MEF cells (a, d),

250 nM colchicine-treated MEF cells (b, e), and 50 nM vinblastine-treated MEF cells

(c, f) were examined using fluorescence microscopy 24 h after transfection with a GFP-

expressing plasmid pCMX-AFAP (upper panels; GFP); the lower panels represent the

corresponding bright-field (BF) images of the upper panels. Scale bar = 500 μm.

Fig. 3. Colchicine- and vinblastine-treated MEF cells do not enhance transfection

efficiency in the absence of p62

(A) The top 87 compounds were examined in p62KO-MEF cells. The p62KO-MEF

cells were transfected with the pCMV-Luc plasmid for 28 h in the presence of 1 μM of

each compound, and transfection efficiency was determined as described in the legend

to Fig. 2A. The fold changes in transfection efficiency relative to the value from the

control DMSO-treated cells are shown in the same order as in Fig. 2A. The two

columns on the right are reproduced from Fig. 2A for comparison. (B, C) The p62KO-

MEF cells were transfected for 28 h with the pCMV-Luc plasmid in the presence of the

indicated concentrations of colchicine (B) or vinblastine (C). Each value of the

transfection efficiency is indicated as the mean ± SD of at least three independent

experimental results.

Fig 4. Colchicine- and vinblastine-treated MEF cells affect the timing of

ubiquitination

(A) Schematic diagram of the experimental system using beads incorporated into cells

to monitor the behavior of transfected materials. (B, C) Time-lapse images of pHrodo

and GFP-Ub fluorescence around a single pHrodo-conjugated bead in MEF cells.

Images were obtained every minute for approximately 60 min. The panels show

representative images of pHrodo and GFP-Ub fluorescence. (B) GFP-Ub MEF cells

treated with DMSO as a control. Arrows indicate the timing of appearance of GFP-Ub

signals around the beads. (C) GFP-Ub MEF cells treated with 500 nM colchicine.

Arrows indicate the timing of appearance of GFP-Ub signals around the beads. Scale

bar = 2 μm. (D) Statistical analysis was performed for the time taken for GFP-signal

accumulation around the beads after the loss of pHrodo signals in the GFP-Ub MEF

cells. Lane 1: control treated with DMSO. Lane 2: 100 nM colchicine-treated MEF

cells. Lane 3: 500 nM colchicine-treated MEF cells. Lane 4: 100 nM vinblastine-treated

MEF cells. The median values were 4 min for GFP-Ub ($n = 24$ beads), 6 min for 100

nM colchicine ($n = 28$ beads), 6 min for 500 nM colchicine ($n = 22$ beads), and 6 min

for 100 nM vinblastine ($n = 31$ beads). Three independent experiments were performed

for each lane, and the total bead number is indicated as n. Statistical differences ($p <$

0.0001) were determined using the Kruskal–Wallis test. Error bars indicate 95%

confidence intervals.

Fig. 5. Colchicine- and vinblastine-treated MEF cells decrease the level of

phosphorylated p62 at S405

Western blot analysis was performed for total p62 and S405-phosphorylated p62 in

MEF cells. Cells were pretreated with 500 nM colchicine or 100 nM vinblastine for 30

min, and then transfected with the pCMX-AFAP plasmid in the presence of 500 nM

colchicine or 100 nM vinblastine. DMSO was used as the control. Samples were

subjected to Western blot analysis with no transfection or with transfection for 24 h.

GAPDH was used as the loading control.

Fig. 1

A

B

Fig. 2

A

B

No	chemical name	target	function	PubChem CID
1	PODOFILOX	microtubule	antineoplastic, antimitotic agent	10607
2	COLCHICINE	microtubule	antimitotic, antigout agent	6167
3	VINBLASTINE SULFATE	microtubule	antineoplastic, spindle poison	241902
4	DEOXYSAPPANONE B 7,4'-DIMETHYL ETHER	microtubule		4026888
5	VINCRISTINE SULFATE	microtubule	antineoplastic	249332
6	PICROPODOPHYLLIN ACETATE	microtubule / IGF-1R	antineoplastic	233299
7	DOCETAXEL	microtubule	antineoplastic	148124
8	FENBENDAZOLE	microtubule	anthelmintic, antinematodal drug	3334
9	FLUBENDAZOLE	microtubule	anthelmitic, antinematodal drug	35802

C

D

Fig. 3

Fig. 4

A

colchicine or vinblastine

GFP-Ub

pHrodo-conjugated beads

acidic endosome
pHrodo-conjugated beads

raptured acidic endosome
GFP-Ub

GFP-Ubwt MEF cell

B GFP-Ub MEF, DMSO

[min] -2 0 2 4 6 8 10 12 14 16

2 μm

pHrodo

GFP-Ub

merge

C GFP-Ub MEF, colchicine 500 nM

[min] -2 0 2 4 6 8 10 12 14 16

2 μm

pHrodo

GFP-Ub

merge

D

over 60 min   1   16   3   10

GFP-Ub accumulation time [min]

p < 0.0001
p < 0.0001
p < 0.0001

inhibitor treatment

1 DMSO
2 100 nM colchicine
3 500 nM colchicine
4 100 nM vinblastine

Fig. 5

Supporting Information

Microtubule inhibitors identified through non-biased screening enhance DNA transfection efficiency by delaying p62-dependent ubiquitin recruitment

Megumi Tsuchiya[1#], Hidesato Ogawa[1#]*, Kento Watanabe[1], Takako Koujin[2], Chie Mori[2], Kazuto Nunomura[3], Bangzhong Lin[3], Akiyoshi Tani[3], Yasushi Hiraoka[1,2], and Tokuko Haraguchi[1,2]*

List of contents:

Table S1. Identification of 87 potential compounds for enhancement of transfection efficiency.

Rank	Structure	Name	Formula	Fold transfection efficiency in MEF	Fold cell viability in MEF	Fold transfection efficiency in p62KO-MEF	Fold cell viability in p62KO-MEF
1		PODOFILOX	C22 H22 O8	297.24	64%	4.46	281%
2		COLCHICINE	C22 H25 N O6	294.20	95%	4.94	258%
3		VINBLASTINE SULFATE	C46 H58 N4 O9 . H2 O4 S	263.07	68%	6.90	318%
4		DEOXYSAPPANONE B 7,4'-DIMETHYL ETHER	C18 H18 O5	54.43	96%	4.72	95%
5		VINCRISTINE SULFATE	C46 H56 N4 O10 . H2 O4 S	46.77	40%	0.66	47%
6		PICROPODOPHYLLIN ACETATE	C24 H24 O9	35.46	34%	2.19	46%
7		DOCETAXEL	C43 H53 N O14 . 3 H2 O	33.78	72%	9.43	316%
8		FENBENDAZOLE	C15 H13 N3 O2 S	27.14	63%	7.83	337%
9		FLUBENDAZOLE	C17 H13 F N2 O3	21.35	71%	4.53	357%
10			C18 H19 Cl F N O2	8.77	59%	1.09	80%
11			C18 H17 N O2 S2	6.80	74%	1.80	68%
12			C15 H26 N4 O5 S2	6.19	89%	1.70	91%
13		OXIBENDAZOLE	C12 H15 N3 O3	4.64	45%	0.85	49%
14		DEOXYSAPPANONE B 7,3'-DIMETHYL ETHER ACETATE	C20 H20 O6	4.50	82%	2.07	85%
15		PHENETHYL CAFFEATE (CAPE)	C17 H16 O4	4.44	72%	1.63	89%
16		MEBENDAZOLE	C16 H13 N3 O3	3.01	95%	1.57	365%
17		AZACITIDINE	C8 H12 N4 O5	2.92	59%	0.93	67%
18			C15 H10 Br F N2 O2	2.84	96%	0.53	109%
19			C15 H18 N4 O3	2.52	97%	0.62	105%
20		PIROCTONE OLAMINE	C14 H23 N O2 . C2 H7 N O	2.04	40%	0.78	42%
21			C18 H16 N2	1.96	94%	0.38	102%

22			C25 H21 N3 O5 S	**1.91**	**113%**	**0.76**	**101%**
23			C15 H29 N3 O2	**1.82**	**113%**	**1.52**	**116%**
24		ISOOSAJIN	C25 H24 O5	**1.59**	**62%**	**1.50**	**76%**
25		ALBENDAZOLE	C12 H15 N3 O2 S	**1.50**	**92%**	**0.62**	**222%**
26		PICROPODOPHYLLIN	C22 H22 O8	**1.49**	**93%**	**0.32**	**110%**
27		DEGUELIN(-)	C23 H22 O6	**1.44**	**36%**	**1.51**	**33%**
28		EPOXYGEDUNIN	C28 H34 O8	**1.37**	**77%**	**2.66**	**83%**
29		NISOLDIPINE	C20 H24 N2 O6	**1.32**	**91%**	**1.05**	**59%**
30		METHYLPREDNISOLO NE SODIUM SUCCINATE	C26 H33 O8 . Na	**1.30**	**81%**	**1.24**	**65%**
31		METERGOLINE	C25 H29 N3 O2	**1.29**	**61%**	**0.93**	**54%**
32		THIOGUANINE	C5 H5 N5 S	**1.28**	**111%**	**0.45**	**106%**
33			C11 H15 Cl N2	**1.26**	**76%**	**1.02**	**88%**
34		TOLONIUM CHLORIDE	C15 H16 N3 S . Cl	**1.24**	**67%**	**0.49**	**45%**
35		MANIDIPINE HYDROCHLORIDE	C35 H38 N4 O6	**1.22**	**70%**	**0.90**	**51%**
36		PREGNENOLONE SUCCINATE	C25 H36 O5	**1.20**	**78%**	**0.85**	**73%**
37			C12 H13 Br N4 O S	**1.15**	**103%**	**0.56**	**102%**
38		MOMETASONE FUROATE	C27 H30 Cl2 O6	**1.14**	**80%**	**1.26**	**54%**
39		ESTRAMUSTINE	C23 H31 Cl2 N O3	**1.08**	**71%**	**0.79**	**77%**
40		PHENOLPHTHALEIN	C20 H14 O4	**1.07**	**76%**	**0.86**	**70%**
41		BROXYQUINOLINE	C9 H5 Br2 N O	**1.07**	**62%**	**0.63**	**58%**
42			C19 H30 Cl N O2	**1.06**	**101%**	**1.04**	**113%**
43		PHENOTHIAZINE	C12 H9 N S	**1.05**	**44%**	**0.86**	**64%**
44		PEUCENIN	C15 H16 O4	**1.05**	**54%**	**1.45**	**31%**

#		Name	Formula				
45		GANGALEOIDIN	C18 H14 Cl2 O7	1.04	78%	1.71	71%
46		7,3'-DIMETHOXYFLAVONE	C17 H14 O4	1.01	84%	1.43	96%
47		3-ACETYLGEDUNOL	C30 H40 O8	1.00	102%	1.37	96%
48		GARCINOLIC ACID	C38 H46 O9	0.99	62%	0.88	54%
49			C12 H12 N6 S2	0.97	109%	0.94	126%
50		SULBENTINE	C17 H18 N2 O S	0.92	72%	0.63	73%
51			C16 H16 Cl N3 O S	0.92	85%	0.85	95%
52		ISRADIPINE	C19 H21 N3 O5	0.90	91%	0.80	71%
53		DIHYDROGEDUNIN	C28 H36 O7	0.90	37%	1.23	69%
54		NITRENDIPINE	C18 H20 N2 O6	0.88	72%	0.82	72%
55		GEDUNIN	C28 H34 O7	0.87	54%	1.23	80%
56		PYRONARIDINE TETRAPHOSPHATE	C29 H32 Cl N5 O2 . 4 H3 O4 P	0.87	79%	0.83	66%
57		BENAZEPRIL HYDROCHLORIDE	C24 H28 N2 O5 . Cl H	0.86	76%	1.06	60%
58		PYRVINIUM PAMOATE	C26 H28 N3 . C23 H15 O6	0.85	79%	1.46	147%
59			C12 H12 N4 S	0.83	101%	0.70	105%
60			C16 H13 N3 O S2	0.82	129%	0.80	137%
61		SODIUM TETRADECYL SULFATE	C14 H29 O4 S . Na	0.81	76%	0.83	69%
62		DIENESTROL	C18 H18 O2	0.81	92%	0.61	219%
63		AMSACRINE	C22 H20 N2 O3 S	0.81	38%	0.18	35%
64		BISMUTH SUBSALICYLATE	C7 H5 Bi O4	0.78	84%	0.84	48%
65			C20 H16 N2 O3 S	0.78	94%	0.86	104%
66		AMINOETHOXYDIPHENYLBORANE	C14 H16 B N O	0.76	94%	0.88	96%
67			C11 H9 N O2	0.75	97%	0.69	106%

68		OXFENDAZOLE	C15 H13 N3 O3 S	0.73	77%	0.80	78%
69		CHLOROXINE	C9 H5 Cl2 N O	0.71	87%	0.72	80%
70		DIETHYLSTILBESTROL	C18 H20 O2	0.66	105%	0.45	204%
71		ESTRADIOL ACETATE	C20 H26 O3	0.60	28%	1.21	49%
72			C23 H26 O3	0.55	104%	0.81	97%
73			C22 H17 N3 O3	0.44	110%	0.86	99%
74			C15 H18 N4 O3 S3	0.41	14%	7.11	37%
75			C14 H14 N4 O S3	0.07	203%	2.33	265%
76			C15 H10 F3 N3 S	0.07	257%	2.35	270%
77			C15 H17 N3 O3	0.07	199%	2.67	260%
78			C16 H10 Br Cl2 N5	0.06	203%	2.62	257%
79			C20 H23 N O4 S	0.06	203%	1.80	265%
80			C17 H10 Cl N O4	0.06	206%	2.52	262%
81			C17 H19 Br O3	0.06	222%	2.36	268%
82			C15 H13 Cl N2 O4 S	0.05	164%	1.75	155%
83			C15 H14 Cl N3 O	0.04	171%	1.52	163%
84			C16 H13 Cl N4 O S2 . Cl H	0.04	186%	1.59	162%
85			C15 H13 Br N2 O2 S	0.03	152%	1.58	150%
86			C14 H18 N6 O2 S	0.03	182%	1.26	190%
87			C16 H18 N2 O2	0.03	203%	1.27	188%

Comparison of 87 compounds derived from the Osaka University compound library, showing their chemical structure, name, formula, transfection efficiency and cell viability in MEF cells and p62KO-MEF cells.

Fig. S1. Nocodazole increases gene transfection efficiency.

MEF cells were transfected with the pCMV-Luc plasmid for 28 h in the presence of 1 μM colchicine (left) and 1 μM nocodazole (right). DMSO was used as the negative control. Transfection efficiency was determined based on luciferase activity normalized to the cell viability. Bar graphs show the fold changes in transfection efficiency relative to the value from the control DMSO-treated cells. Each value is indicated as the mean ± SD of at least threeindependent experiments.

Fig. S2. Colchicine increases transfection efficiency for various transfection reagents.

MEF cells were transfected with the pCMV-Luc plasmid in the presence of 300 nM colchicine by using Effectene, Viofectin, or Lipofectamine 2000. DMSO was used as the negative control. Details of transfection procedures for each transfection reagent are described in the Experimental procedures. Bar graphs show the fold changes in transfection efficiency relative to the value from the control DMSO-treated cells. (mean ± SD of at least three independent experiments).

Fig. S3. Colchicine and vinblastine treatments do not affect the promoter activity of the luciferase gene.

(A, B) MEF cells stably expressing luciferase (MEF-LUC) were treated with the indicated concentrations of colchicine (A) or vinblastine (B). After incubation for 27 h, the luciferase activity of the cells was measured and normalized to cell viability. The graphs show the fold changes in luciferase activity relative to the value from the control DMSO-treated cells (indicated as mean ± SD of at least three independent experimental results).