Using ephaptic coupling to estimate the synaptic cleft resistivity of the calyx of Held synapse.

Short title: **Electronic model of the synaptic cleft.**

Martijn C. Sierksma & J. Gerard G. Borst

Department of Neuroscience, Erasmus MC, University Medical Center Rotterdam, 3000 CA, Rotterdam,
The Netherlands.

Correspondence to g.borst@erasmusmc.nl

A conference paper including parts of this paper was presented at Proceedings of the 23rd International
Congress on Acoustics (2019, doi: 10.18154/RWTH-CONV-240008)

Keywords: Prespike; Cleft conductance; Giant synapse; Cleft potential; Auditory brainstem; phasor.

Abstract

At synapses, the pre- and postsynaptic cell get so close that currents entering the cleft do not flow

exclusively along its conductance, g_{cl}. A prominent example is found in the calyx of Held synapse in the

medial nucleus of the trapezoid body, where the presynaptic action potential can be recorded in the

postsynaptic cell in the form of a prespike. Here, we developed a theoretical framework for ephaptic

coupling via the synaptic cleft. We found that the capacitive component of the prespike recorded in

voltage clamp is closely approximated by the second time derivative of the presynaptic action potential.

Its size scales with the fourth power of the radius of the synapse, explaining why intracellularly recorded

prespikes are uncommon in the CNS. We show that presynaptic calcium currents can contribute to the

prespike and that their contribution is closely approximated by the scaled first derivative of these currents.

We confirmed these predictions in juvenile rat brainstem slices, and used the presynaptic calcium currents

to obtain an estimate for g_{cl} of ~1 μS. We demonstrate that for a typical synapse geometry, g_{cl} is scale-

invariant and only defined by extracellular resistivity, which was ~75 Ωcm, and by cleft height. During

development the calyx of Held develops fenestrations. These fenestrations effectively minimize the cleft

potentials generated by the adult action potential, which would otherwise interfere with calcium channel

opening. We thus provide a quantitative account of the dissipation of currents by the synaptic cleft, which

can be readily extrapolated to conventional, bouton-like synapses.

Author Summary

At chemical synapses two neurons are separated by a cleft, which is very narrow. As a result, the

concentration of released neurotransmitter can rapidly peak, which is essential for fast synaptic

transmission. At the same time, the currents that flow across the membranes that face the synaptic cleft are

also substantial, and a fraction of these currents will enter the other cell. We made an electronic model of

the synaptic cleft, which indicated that if the size of the membrane currents that flow across both

membranes are known, our model can be used to infer the electrical resistance of the synaptic cleft. We

tested several of the predictions of our model by comparing the membrane currents during a presynaptic

action potential in a giant terminal, the calyx of Held, with the much smaller prespike evoked by these

currents in the postsynaptic cell. We estimate the cleft resistance to be about 1 MΩ, which means that the

changes in the cleft potential due to the membrane currents can become large enough to have an impact on

voltage-dependent calcium channels controlling neurotransmitter release.

Introduction

When sir Sherrington introduced the term 'synapsis' [1], it was still unknown whether or not nerve impulses reach their target through direct electrical coupling [2, 3]. Together with other evidence, it was the visualization by electron microscopy of the synaptic cleft, the narrow gap between an axon and its target, which put the debate to rest in favor of transmitter-mediated neurotransmission [4-6]. At about the same time, Eccles and Jaeger (7) outlined the electrical circuitry of the synapse, which included a cleft leak conductance (g_{cl}) linking synaptic currents with the interstitial space. Since then, the role of the synaptic cleft for synaptic transmission has gradually become more important. Its small volume not only allows a rapid buildup of released neurotransmitter, but protons [8, 9], and potassium ions [10-12] may accumulate as well, whereas calcium ions may become partially depleted [13, 14]. Interestingly, during transmission a cleft potential may arise that may alter the dwelling time of mobile, charged particles within the cleft [15, 16], the distribution of ligand-gated ion channels [17], or the kinetics of voltage-gated channels in the cleft-facing membrane [18]. These features all stem from the small size of the synaptic cleft, which thus forms a barrier for particle movements [19].

Except for some specialized synapses, the contribution of g_{cl} to synaptic transmission has largely been ignored or deemed irrelevant, but its importance will of course depend on its magnitude and on the size of the currents that flow within the cleft. Eccles and Jaeger (7) already indicated that the maximal cleft current will be limited by this conductance. g_{cl} decreases as the path length l_{path} to the extracellular space increases ($g_{cl} \sim A/(R_{ex} \, l_{path})$), where R_{ex} is extracellular resistivity in Ω cm and A the area through which the current flows (see also Savtchenko, Antropov (20)). Due to their much longer path length, it has been argued that giant synapses have a lower g_{cl} than conventional synapses [21]. Remarkably, estimates of g_{cl} of very different synapses vary over four orders of magnitude (range 1.4 nS to 40 μS) without a clear relation to synapse size [7, 11, 18, 20, 22-30]. Despite the remarkable progress in our understanding of the synaptic cleft, we need a better theoretical understanding of its conductive properties, which is ideally confirmed by experimental observations.

One of the largest synapses in the mammalian brain is the calyx of Held synapse, which spans about 25

µm [31, 32]. Its large size matches its strength, with the maximal glutamatergic conductance exceeding

100 nS, and its fast presynaptic AP allowing firing rates >300 Hz [33]. Large currents, related to the

excitatory postsynaptic current (EPSC) and capacitive currents, need to flow within the cleft through g_{cl},

and under these conditions g_{cl} may limit synaptic transmission. A characteristic feature of this synapse is

that the presynaptic AP can be detected in the postsynaptic recording (Figure 1). This hallmark of the

calyx of Held synapse has been called the prespike [34]. It is probably due to ephaptic coupling of the

presynaptic AP via the synaptic cleft [30, 35], but the precise relation between the presynaptic AP and the

prespike has not been studied. Here, we use the prespike to infer some key electrical properties of this

giant synapse. We provide a theoretical description of ephaptic coupling by defining the electrical

properties of the synaptic cleft, which is validated through dual patch-clamp recordings of the calyx of

Held and its postsynaptic target.

Results

The prespike is the ephaptically coupled calyx of Held AP recorded in its target neuron, a principal neuron

in the medial nucleus of the trapezoid body (MNTB) [34]. Some current clamp (CC) and voltage clamp

(VC) prespike examples are shown in Figure 1A-B. Figure 1C shows that the amplitudes of the CC and

the VC prespikes recorded in the same cell are correlated. Figure 1D shows examples of the CC prespikes

recorded during *in vivo* whole-cell recordings in young rats, whose shape resemble those recorded in the

simultaneous pre- and postsynaptic slice recordings (Figure 1A-B, 2A).

Figure 1 – Examples of prespikes. (A-B) Two examples of prespikes in simultaneous pre- and postsynaptic

patch-clamp recordings. From top to bottom: calyx of Held AP, which was elicited by afferent stimulation, CC

prespike, associated VC prespike. Recordings were made at room temperature in brainstem slices from 7-10

days old Wistar rats [36]. Horizontal scale bar: 0.5 ms. Vertical (from top to bottom): 50 mV, 2 mV, 1 nA. (C)

Relation between amplitude of the positive peak of the prespike measured in CC and the amplitude of the first

negative peak of the prespike measured in VC in the same cell. Slope of the fitted line is -4.6 mV/nA. (D). Six

examples of CC prespikes from *in vivo* recordings of a target neuron of the calyx of Held in 3-5 days old Wistar

rats [37]. During this period the calyx of Held expands over the postsynaptic cell. Note the small upward

deflection immediately preceding the EPSP in the bottom three prespikes. Scale bar indicates 0.5 mV and 0.5

ms.

Figure 2. An electrical circuit for the prespike. (A) Drawing of the calyx of Held synapse. The calyx of Held

terminal ('pre') covers a large part of the soma of its target neuron ('post'). In between is the synaptic cleft (not

drawn to scale), a small space where membrane currents will flow during a presynaptic AP. (B) Electrical

circuit to reproduce the prespike. The capacitance facing the cleft (c_{cl}) is assumed to be equal for the calyx and

its target. Presynaptic membrane currents can leave the synaptic cleft via the cleft conductance (g_{cl}) or through

the postsynaptic cell. It is assumed that the conductance from cleft to ground is relatively large compared to the

cleft conductance [38, 39].

Phasor analysis of an electrical circuit for the prespike

The prespike is generated by currents that flow from the calyx to the principal neuron in the

MNTB. While the current flow through the cleft depends on the synapse geometry, which we will

consider later in this article, we can already get a good understanding by analyzing a prespike-

generating, electrical circuit [30] with the following components (Figure 2B): (1) the terminal's

release face, consisting of voltage-gated conductances (not shown), a capacitor (c_{cl}), and a

conductance (g_{pre}); (2) the synaptic cleft, which is separated from the reference by a leak

conductance (g_{cl}), and (3) a passive neuron, which partly faces the synaptic cleft. This part consists

of a capacitor (c_{cl}) and a leak conductance (g_{syn}). The remainder of the postsynaptic cell consists of

a capacitor ($c_{post}-c_{cl}$) and a leak conductance ($g_{post}-g_{syn}$). As the prespike is small, we ignore the

postsynaptic voltage-dependent conductances. We also did not include a direct resistive connection

between calyx and target since there is no physiological evidence for gap junctions at the calyx of

Held synapse [40], and capacitive currents are likely to dominate the electrical coupling: the

susceptance at 1 kHz across an area of 1000 μm^2 is 63 nS ($2\pi f C_m A$, where f is the frequency

component, C_m is the specific capacitance, and A the membrane area), which is much larger than

the postsynaptic resting conductance of ~4 nS.

We use phasors to describe analytically how currents flow and voltages change within this circuit in

response to the presynaptic AP. In our notation, capitals indicate phasors, e.g. V, I; lowercases

indicate real-valued responses, v, i; j indicates the imaginary number $\sqrt{-1}$. To come to an analytical

description of the prespike, we first describe the relation between the presynaptic AP and the cleft

potential. The cleft potential can be approximated as the presynaptic potential multiplied by the

voltage divider formed by the presynaptic capacitance Y_2 and the parallel combination of cleft

conductance Y_3 and postsynaptic capacitance: $V_{cl} = V_{pre}(-jY_2)\left(Y_3 - j(Y_2 + Y_5)\right)^{-1}$ (Appendix

A1.1). This notation can be transformed to an angular notation, which shows how amplitudes are

scaled by the modulus and shifted in time by the phase shift: $V_{cl} = V_{pre}\, 2\pi f \tau_{cl}\,((4\pi f \tau_{cl})^2 +$

$1)^{-0.5} \angle atan2(-1, 4\pi f \tau_{cl})$ (Appendix A1.2 and A1.3), where the cleft time constant τ_{cl} is defined

as $\tau_{cl} = c_{cl}/g_{cl}$. These phasor relations can be simplified when we assume that g_{cl} is substantially

higher than the cleft susceptance ($g_{cl} \gg 2\pi f c_{cl}$). The cleft potential then becomes $V_{cl} =$

$V_{pre} 2\pi f \tau_{cl} \angle atan2(-1, 4\pi f \tau_{cl})$ (Appendix A1.6). We thus observe a prominent contribution of the

frequency component (f), indicating that faster voltage and current changes will be more readily

transmitted. We see that v_{cl} is a high-pass filtered v_{pre}, and that it approaches the first derivative of

v_{pre} (as the phase shift approaches -π/2). This is to be expected as the current that enters the

synaptic cleft is mainly the capacitive current $c_{cl}\, dv_{pre}/dt$. Using the waveform of a presynaptic AP,

we can calculate the cleft potential (Figure 3A-B), which, indeed, resembles the biphasic capacitive

current.

These changes in v_{cl} will generate a postsynaptic current, the VC prespike, $I_{post} = -jY_5V_{cl}$, which

yields $I_{post} = -V_{pre}\,g_{cl}(2\pi f\tau_{cl})^2((4\pi f\tau_{cl})^2 + 1)^{-0.5}\,\angle atan2(-4\pi f\tau_{cl},\,-1)$ (Appendix A2.1-2.5;

we use the convention that an inward current is negative). This current underlies the CC prespike,

which can be described as $V_{post} = -I_{post}\,(Y_4 - jY_5 + Y_6 - jY_7)^{-1}$, which gives $V_{post} =$

$V_{pre}(2\pi f c_{cl})^2\left(\left[g_{post}{}^2 + \left(2\pi f c_{post}\right)^2\right]\left[g_{cl}{}^2 + (4\pi f c_{cl})^2\right]\right)^{-0.5}$

$\angle atan2\left(-\left(2\pi f c_{post}g_{cl} + 4\pi f c_{cl}g_{post}\right), -g_{post}g_{cl} + (2\pi f)^2c_{post}c_{cl}\right)$ (Appendix A3.1-3.7). When

we again assume that g_{cl} is substantially higher than the cleft susceptance, the VC prespike becomes

$I_{post} = -V_{pre}\,(2\pi f\tau_{cl})^2g_{cl}\,\angle atan2(-4\pi f\tau_{cl}, -1)$ (Appendix A2.6); and when we also assume for

the CC prespike that the postsynaptic susceptance is substantially higher than the postsynaptic

conductance ($2\pi f c_{post} \gg g_{post}$), we obtain $V_{post} = V_{pre}\,2\pi f\tau_{cl}\;c_{cl}c_{post}^{-1}$

$\angle atan2\left(-2\pi f\tau_{post}, ((2\pi f)^2\tau_{cl}\tau_{post} - 1)\right)$ (Appendix A3.8 and A3.9). The VC prespike therefore

reflects the second derivative of v_{pre} scaled by g_{cl} as the phase shift approaches $-\pi$ (Figure 3C).

Lastly, the CC prespike is the first derivative of v_{pre} scaled by the relative postsynaptic cleft

capacitance (Figure 3D).

Figure 3 – Transformation of the presynaptic AP by phasors yields prespike-like responses. (A) An AP

waveform used in voltage-clamp experiments to elicit presynaptic calcium currents. (B-D) The capacitive

response in the cleft potential (B), in the capacitive prespike currents (C), and in the capacitive prespike

potentials in response to the presynaptic AP (D). (E) The calcium current can be simulated as a Gaussian

function. (F) The calcium prespike in response to the calcium current. (G-H) The VC prespike (G) and CC

prespike (H) in response to both the capacitive current and the calcium current. G is the combined currents in C

and F. The parameter values were: c_{cl} = 10 pF, g_{cl} = 1 μS, c_{post} = 30 pF, g_{post} = 5 nS, q_{ca} = 1 pC, and τ_{Ca} = 0.21

ms. The calcium current peak was aligned with a membrane potential of +30 mV relative to the resting

potential on the downward slope of the AP waveform.

It is more challenging to obtain an analytical description for the impact of the voltage-gated

conductances on the prespike, as these conductances depend on the transmembrane potential at the

cleft-facing membrane, and therefore are not expected to remain constant in time. However, if the

amplitude and time course of an ionic current (i_{ion}) are known, its impact on the prespike currents

can be described as $I_{post} = I_{ion}\,(-jY_5)(Y_3 - j(Y_2 + Y_5))^{-1}$, which simplifies to $I_{post} =$

$I_{ion} 2\pi f \tau_{cl} \angle atan2(-1, 4\pi f \tau_{cl})$ (Appendix A4.1). The calcium current of the calyx of Held closely

follows a Gaussian function [41], which has the advantage that its Fourier transform is another

Gaussian function. We describe the calcium current with a charge transfer q_{ca} centered at its peak

as $i_{ca} = i_{peak}\exp(-0.5(t/\tau_{Ca})^2)$ (Appendix 4.2; Figure 3E; τ_{Ca} corresponds to the standard

deviation of the Gaussian function). Its Fourier transform is $I_{ca} = q_{ca}\exp(-2(\pi\tau_{Ca}f)^2)$ (Appendix

4.3). We multiply I_{ca} with the transfer function and perform an inverse Fourier transform: $i_{post}(t) =$

$-\tau_{cl}\,i_{peak}\tau_{Ca}^{-2}\,t\,\exp(-0.5(t/\tau_{Ca})^2)$, which is $-\tau_{cl}\,di_{Ca}/dt$ (Appendix A4.4-A4.6). This shows a

biphasic response with a positive and a negative peak (Figure 2F) at $\pm\,\tau_{Ca}$ with an amplitude of

$\tau_{cl}(\tau_{Ca}\sqrt{e})^{-1}\,i_{peak}$. We therefore define a simple scaling factor $\tau_{cl}/(\tau_{Ca}\sqrt{e})$ that (with a phase shift)

relates the presynaptic calcium current amplitude to a biphasic, postsynaptic current. This biphasic

response is only the calcium component of the VC prespike (the calcium prespike), and it needs to

be combined with the capacitive currents generated by the presynaptic AP (the capacitive prespike)

to obtain the VC prespike (Figure 3G) and subsequently the CC prespike (Figure 3H). We observe

that the first part of the VC prespike, including its first negative peak, is mainly due to the

capacitive prespike, while the last part with the second negative peak is mostly due to the calcium

prespike.

Experimental validation of the model

Based on our phasor analysis, we predict that the VC prespike resembles the second derivative of

the presynaptic AP. To test this prediction, we revisited published simultaneous whole-cell

presynaptic current clamp recordings with postsynaptic voltage clamp recordings [36, 42]. In these

recordings, the afferent axon of the calyx of Held was stimulated with an electrode to trigger a

presynaptic AP, and a prespike followed by an EPSC was recorded postsynaptically (Figure 4A).

The prespike generally matched the scaled inverted second derivative of the AP (Figure 4A-B),

taking into account that it may have been filtered by the series resistance. Figure 4B shows that in

these recordings the time course of the second derivative of the AP indeed provides a better match

for the prespike than its first derivative (n = 13 paired recordings). This is in agreement with the

idea that the prespike is generated by two capacitors in-series.

Figure 4 – Relation between the calyceal AP and the prespike. (A) *Left column, from top to bottom*: AP

recorded in CC from the calyx of Held, its inverted first and second derivatives, and postsynaptic voltage clamp

recording showing the accompanying prespike. Insets show the same presynaptic AP (top) and the postsynaptic

EPSC (bottom) elicited by the calyceal stimulation on a longer time scale. Vertical scale bars (from top to

bottom panels): 80 mV, 0.8 kV/s, 8 MV/s^2, 0.2 nA. Horizontal: 0.5 ms. (B) Relation between the delay between

the first negative and positive peak of the prespike and the first two peaks in the first (blue) and second (black)

derivative of the presynaptic AP. Dashed line is the identity line. (C) *Left column, from top to bottom*:

Presynaptic calcium current (after P/5 subtraction), its first derivative, postsynaptic currents (black, active) and

passive prespike (blue, P/5-scaled), difference between active and passive prespike (black minus blue), which

corresponds to the calcium prespike. Vertical scale bars: 2 nA, 15 nA/ms, 0.15 nA, 0.1 nA. Horizontal: 0.5 ms.

(D) Relation between the first derivative of the presynaptic calcium current shown in the bottom panel of C and

the calcium prespike. Maximum correlation was obtained when delaying the first derivative by 40 μs. Blue line

shows regression line with a slope of 5.6 μs.

A second prediction of our phasor analysis is that presynaptic calcium currents that arise from the

cleft will have an impact on the shape of the prespike. We turned to simultaneous whole-cell pre-

and postsynaptic voltage clamp recordings of the calyx of Held synapse that were originally

reported in Borst and Sakmann (43), which were re-analyzed to evaluate the presence or absence of

a calcium prespike. In these experiments the presynaptic compartment was voltage clamped with an

action potential waveform (APW) command while the presynaptic calcium currents were

pharmacologically isolated (Figure 4C). To isolate the postsynaptic current related to the calcium

currents, we averaged the postsynaptic responses to the P/5 AP waveform and subtracted the scaled

passive prespike from the 'active' prespike to obtain the calcium prespike. The remaining currents

indeed showed the biphasic response predicted by the phasor analysis (Appendix A4; cf. second and

bottom panel of Figure 4C).

From the phasor analysis we know that when the presynaptic calcium current and the VC prespike

are simultaneously recorded, τ_{cl} can be calculated by comparing di_{Ca}/dt and the calcium prespike.

To estimate τ_{cl}, we first analyzed the presynaptic calcium current that was evoked by the APW. It

had a peak amplitude of -1.9 ± 0.6 nA and a full-width at half-maximum (FWHM) of 0.51 ± 0.05

ms (n = 5 paired recordings). Next, we obtained the values for the calcium prespike. Its peak-to-

peak amplitude was 80 ± 60 pA and its peak-to-peak delay was 0.54 ± 0.11 ms. This peak-to-peak

delay should be about twice τ_{Ca}, which was thus slightly higher than the 0.21 ms reported by Borst

and Sakmann (44). Its time course closely matched the model predictions (cf. Figure 4C and Figure

3F). As predicted by the phasor analysis, the amplitude of the calcium current was linearly related

to the calcium prespike peak (r = 0.98, p = 0.003); the calcium prespike peak was 1.9 ± 0.7% of the

calcium current amplitude. Above, we derived that this percentage should be $\tau_{cl}\,\tau_{Ca}^{-1}e^{-0.5}$, yielding

an estimated τ_{cl} of 9 ± 5 μs. As an alternative approach, we plot the first derivative of the

presynaptic calcium current against the calcium prespike. Its maximal slope (allowing for a slight

time shift) provided a τ_{cl} of 6 ± 2 μs (Figure 4D). The two methods thus gave comparable values for

the cleft time constant.

The ratio of cleft capacitance c_{cl} and cleft leak conductance g_{cl} constitutes τ_{cl}. If we assume that c_{cl}

is 10 pF, the estimated g_{cl} becomes about 0.5-1 μS. From the phasor analysis we know that the

capacitive VC prespike coming from the presynaptic AP and the calcium prespike do not scale the

same way to c_{cl} and g_{cl} (*cf.* Appendix A2.1 with A4.1). This difference allows us to further define

these two parameters. We define the relation between the postsynaptic calcium component and the

presynaptic calcium current as $I_{post}/I_{Ca} = 2\pi f \tau_{cl} = R_{Ca}$. If we substitute $2\pi f \tau_{cl}$ in equation A2.6 with

R_{Ca}, the equation for the capacitive prespike becomes $I_{post} = V_{pre} R_{Ca}^2 g_{cl}$, and the Fourier transform

will yield the same scaling factor of $(R_{Ca})^2 g_{cl}$. The first peak of the VC prespike had an amplitude

of -120 ± 110 pA, which corresponds to the capacitive currents. We can then calculate a g_{cl} of 2.6 ±

0.8 μS and a c_{cl} of 24 ± 16 pF. (In this calculation we assume that the time course of the calcium

current and the presynaptic AP have a very similar frequency composition.)

For τ_{cl} we assumed that the calcium current entirely originated from the membrane facing the cleft.

These estimates should be considered as a lower bound as some of the current may not arise from

the cleft-facing membrane. Immunolabeling for calcium channels identified a fraction of the N-type

and R-type calcium channels to be in the non-cleft-facing membrane [45]. If we assume that only

two-third of the calcium current originated from the cleft, our estimate of τ_{cl} increases to 13.5 ± 7.8

μs. This estimate could be considered as an upper bound, as less than 50% of the calcium current

was mediated by N-type and R-type channels, and the other, P/Q-type, calcium channels were

mostly associated with the release face [45]. When we assume that 2/3 of the total presynaptic

calcium current originated from the cleft, we calculate a g_{cl} of 1.2 ± 0.4 μS and a c_{cl} of 16 ± 11 pF.

With our limited data set we determine that g_{cl} ranges between 1-2 μS.

Cleft currents

For the calyx of Held synapse to be strong and fast, it needs to rapidly conduct large currents. We focus on the generation of the prespike, but a similar analysis can be made for the feedback effect of postsynaptic currents on the calyx of Held [30]. The calyceal EPSC is very fast and can easily be 10 nA [36, 46-49]. This EPSC should flow not only through the postsynaptic glutamate receptor channels, but also through g_{cl} [7]. For the synapse to be able to conduct an EPSC of 10 nA at a driving force of 100 mV, the cleft leak conductance must be at least >0.1 μS. This is consistent with the observed range for g_{cl}.

Besides the capacitive currents and the EPSC, voltage-gated channels facing the cleft may also contribute to the cleft currents. We therefore explored the potential contribution of Na^+ and K^+ currents to the prespike. We estimated their density based on literature values, and assumed a homogeneous distribution within the terminal. As voltage-gated channels in the cleft-facing membrane sense v_{pre}-v_{cl} instead of v_{pre}, their gating depends on v_{cl}. This feedback effect of v_{cl} could not be incorporated in the phasor calculations. We instead simulated the consequences of v_{cl} on voltage-gated currents by numerically solving ordinary differential equations with different values for g_{cl}. Figure 5 illustrates that the presence of these channels at the release face may have a large impact on the shape and size of the prespike. Due to the differences in their gating properties, the timing of each current during the presynaptic AP is quite different (Figure 5A), and the same holds for their impact on v_{cl} and on the prespike (Figure 5B). The prespike examples (Figure 1) did not resemble the prespike of the homogeneous model, which included Na^+, K^+ and Ca^{2+} currents at the cleft-facing membrane, suggesting that some voltage-gated ion channels are excluded (at least functionally) from the release face. In particular, the voltage-gated sodium conductance generates a current that partially counteracts the first prespike peak. These very fast changes in the prespike during the rising phase of the presynaptic AP would point at a contribution by sodium channels. As

these were not observed, we conclude that there is no substantial contribution of sodium channels at

the release face to the prespike.

Figure 5 – Impact on the prespike of voltage-gated ion channels located at the release face. (A) Timing of

different voltage-gated currents at the release face during the AP. Note that the amplitudes of each of the

currents depend on cleft potential, thus creating a mutual dependency. Vertical scale bar: 60 mV, 2.4 nA, 1 nA,

1 nA, 3 nA. Horizontal: 0.5 ms. (B) The cleft potential (top), the VC prespike (middle), and the CC prespike

(bottom) under different conductance density conditions. Homogen.: homogeneous density; no VGSCs: no

sodium conductance at the release face; only VGCCs: calcium and capacitive currents at the release face; no

channels: the capacitive-only model. Scale bars: 4 mV, 0.2 nA, 2 mV.; 0.5 ms, except for the homogeneous

model where the vertical scale bar corresponds to 1.6 nA.

Potassium channels do not start to open until around the peak of the presynaptic AP. They therefore do not

impact the first part of the VC/CC prespike. The potassium current contributes to a more positive synaptic

cleft potential, which would be opposite to the effect of the calcium current. If the potassium current at the

cleft were significantly larger than the other currents, we would expect a fast inward current in the VC

prespike after the first positive peak, and the CC prespike would become a single positive deflection.

However, CC prespikes are typically biphasic (Figure 1), therefore the contribution of the potassium

current is unlikely to outweigh the contribution of the other currents.

Lastly, we revisit the impact of calcium channels on the prespike. The observation of a calcium prespike

already demonstrated the presence of these channels at the release face. The calcium current occurs during

the repolarization phase of the presynaptic AP [44], and therefore contributes little to the first two peaks of

the VC or to the first peak of the CC prespike. Its contribution is easily obscured by noise or by the EPSC,

which starts rapidly after the calcium-induced glutamate release. Nevertheless, it may generate a small,

upward inflection in the CC prespike and such an inflection can sometimes be observed (Figure 1D).

These simulations show that the prespike waveform may contain information about the presence of

voltage-dependent ion channels at the presynaptic membrane facing the cleft. This means that a

comprehensive experimental study of the impact of specific channel blockers on the prespike

waveforms in experiments similar to Figure 3 may be informative about their presence at the

release face.

Voltage-dependent kinetics of cleft-facing calcium channels

During development the presynaptic AP narrows from a half width of 0.5 to 0.2 ms [46]. The

briefer AP necessitates a much larger capacitive current, which, all else being equal, would generate

a larger change in v_{cl}, which may impact the calcium channel kinetics. We again simulated the

impact of v_{cl} on the calcium current by solving ordinary differential equations for different values

for g_{cl} (Figure 6). For a slow AP, a g_{cl} of 0.67 μS and a c_{cl} of 10 pF, v_{cl} first rises to a maximum of

+3.9 mV, and then decreases to a minimum of -4.2 mV. While v_{cl} has little effect on the current

amplitude (Figure 6B), it generates a delay of 45 μs in the onset of the calcium current (Figure 6C).

In contrast, under the same conditions we observe for a fast AP that v_{cl} rises to a maximum of +16.4

mV, followed by a decrease to a minimum of -8.2 mV. As a consequence, the peak calcium current

was delayed by 21 µs and reduced by 19% (-2.0 nA vs -1.7 nA). These effects became even more

pronounced at lower g_{cl}. The observed delay corresponded to the delay in reaching the $v_{pre} - v_{cleft}$

maximum compared to the v_{pre} maximum; the reduction in calcium current was due to a smaller

effective AP, which reduced the maximal open probability of the calcium channels. We conclude

that developmental acceleration of the calyceal AP will increase the capacitive current to a level

that potentially could delay and impede the presynaptic calcium current substantially if g_{cl} were <1

µS.

Figure 6 – Potential impact of the cleft potential on presynaptic calcium currents. (A) Simulation of the

impact of a slow and a fast AP (top; FWHM: left 1.0, right 0.2 ms) on the cleft potential (second row), the

transmembrane potential sensed by the calcium channels in the cleft (third row) and the presynaptic calcium

current (bottom) at three leak resistances (0 MΩ, 1 MΩ and 5 MΩ). Scale bar: 228 mV, 103 mV, 57 mV, 1 nA.

(B) Relation between peak calcium currents and leak resistance for the two APs. Driving force slightly

increases (not shown). (C) The change in delay between the calcium peak current and the AP for the two APs

(solid line). The dashed line shows the delay after correcting for the decrease in the calcium current shown in

B. Calcium conductance density was adjusted to 0.2 nS µm^{-2} and 2.8 nS µm^{-2} for the slow and fast AP,

respectively, in order to elicit a 2 nA-current at 0 MΩ cleft leak resistance. The apposition area was 1000 µm^2.

Other conductances were set to 0 nS.

Fenestration of the calyx of Held reduces cleft potentials.

So far we modeled the cleft as a single compartment with capacitive and conductive properties. However,

it is expected that the voltage profile within the cleft will depend on synapse geometry [20]. We therefore

need an analytical description of the voltage profile within the cleft. Previous work considered the impact

of cleft geometry on the EPSC, a current that runs radially through the cleft and enters at a centered region

containing the glutamate receptors [20]. Our situation is different as we consider a homogeneously

distributed current: the capacitive currents arise at every membrane patch facing the cleft. For an

analytical description we start by assuming a radially-symmetric synapse with a radius r and a cleft height

h. Within this synapse we only consider capacitive currents from the cleft-facing membrane (Figure 7A).

As only a small fraction of these currents enters the postsynaptic cell, we simply assume that all currents

run radially to the synaptic membranes. The electric field at location x relative to the cleft center will be

$dv_{cl}/dx = x R_{ex} C_m (2h)^{-1} dv_{pre}/dt$ (Appendix B1.3) where R_{ex} is the extracellular cleft resistivity and

C_m is the specific membrane capacitance. The electric field grows linearly with the distance from the

center. The cleft potential is equal to $\int_x^r dv_{cl}\, dx$, which gives $v_{cl} = (r^2 - x^2) R_{ex} C_m (4h)^{-1} dv_{pre}/dt$

(Appendix B1.4). The size of the cleft potential thus depends on r^2, indicating that for larger terminals the

cleft potential may become increasingly important (Figure 7B). The voltage profile also shows that not all

voltage-gated channels in the cleft will be affected, but only the channels located towards the cleft center.

The temporal changes in the cleft potential will generate a capacitive current at the postsynaptic side. To

obtain the VC prespike, we derive $i_{post} = r^4 \pi R_{ex} C_m^2 (8h)^{-1} d^2 v_{pre}/dt^2$ (Appendix B2.2). The VC

prespike therefore grows with the fourth power of the radius, which explains why prespikes are only

observed in giant synapses such as the calyx of Held synapse. With our geometrical relation for the VC

prespike we can relate g_{cl} to extracellular resistivity: $g_{cl} = 8\pi h (R_{ex})^{-1}$ (Appendix B3). This relation

leads to an estimate for the intra-cleft resistivity of 75 Ω cm for an h of 30 nm [50] and a g_{cl} of 1 μS.

Considering that there may be extended extracellular spaces within the synaptic cleft [51], and taking into

account the variation in g_{cl} (1-2 μS), we estimate an R_{ex} in the range of 50-100 Ω cm.

Figure 7. Synapse geometry determines the profile of the cleft potential. (A) Drawing of a radially-

symmetric synapse with cleft height h and radius r. The capacitive currents will generate a cleft potential v(x)

at location x. (B) Cleft voltage profile for synapses with different radii (color-coded) at the first peak in the

capacitive current. The horizontal axis depicts the location from the center (0) relative to the edge (1.0). In (B),

R_{ex} is 100 Ω cm, C_m is 1 μF cm^{-2}, and h is 30 nm.

Giant synapses typically fenestrate during development [52-55]. We initially assumed that fenestration

minimizes v_{cl} by reducing g_{cl} through reducing the path length to the extracellular fluid (r). However, we

analytically derived that for a radially-symmetric synapse, g_{cl} does not dependent on its radius. The relation

between the conductance and the resistivity is $g_{cl} \sim A \, (r \, R_{ex})^{-1}$, where A is the area through which the current

escapes. As A of a radially-symmetric cleft is $2\pi rh$, when r is doubled, A is also doubled, leaving the

conductance independent of synapse size. The cup-shape of the young calyx makes the radially-symmetric

synapse a reasonable approximation. This also means that our estimate of the resistivity of the synaptic cleft is

relevant for smaller synapses with a comparable radially-symmetric shape.

Following a period of fenestration, the calyx of Held becomes, as its name implies, a synapse with

extended fingers [31, 56]. However, for finger-like synapses with a sheet-like or cylindrical cleft, the

geometrical relations are slightly different. For these synapses we assume that the currents only run

transversally, *i.e.* perpendicular to the longitudinal axis. The cleft potential then becomes $v(x) =$

$(r^2 - x^2)(2h)^{-1}R_{ex}C_m dv_{pre}/dt$ (Appendix B4.4), where $2r$ equals the width of the sheet (or r the length

of the cylinder). As the escape route for the current is more restricted than for a radially-symmetric

synapse, its voltage gradient is steeper. However, the diameter of a calyceal finger will be much less than

the diameter of the juvenile calyx, effectively reducing the cleft potential. For a calyceal finger of 4 μm

diameter [31, 57] the cleft potential at the center of the finger-like synapse would be only ~2.5% of the

cleft potential in the cup-shape calyceal synapse (equation B5.2). As a consequence, the VC prespike

would become ~3.3% assuming that the total contact surface is conserved (equation B5.4). The cleft

potential and VC prespike with more complex synapse morphology may be investigated using finite-

element modeling, but this is beyond the scope of this study.

Fenestration will effectively reduce the amplitude of the cleft potentials, and therefore we expect that the

prespike in the adult calyx of Held synapse should be significantly smaller than the juvenile prespike. To

study the adult prespike, we revisited *in vivo* whole-cell current-clamp recordings of MNTB principal

neurons from wild-type C57BL/6J mice previously reported in Wang, van Woerden (58) or in Lorteije,

Rusu (59). CC prespikes consisted of a small positive deflection, preceding the start of the rapid rising

phase of the EPSP by about 0.3 ms. In 11 of 12 recordings, the amplitude of the positive deflection was

<0.2 mV. In contrast to the juvenile recordings (Figure 1), the positive peak was not followed by a

negative peak. An example is shown in Figure 8. The good correlation between the size of CC and VC

prespikes (Figure 1C) in combination with the absence of a large decrease in the input resistance of adult

principal neurons [58, 59] suggests that the VC prespike can be expected to also be relatively small. Our

anecdotal evidence from *in vivo* whole-cell recordings in adult mice therefore confirms that the prespike is

smaller than the juvenile prespike and that it possibly has a different shape (*cf.* Figure 1 and 8).

Figure 8 – Adult prespike. (A) Black traces: Spontaneous, sub- and suprathreshold events recorded during a

whole-cell *in vivo* recording of a principal neuron in the MNTB from a P52 C57BL/6J mouse [58]. Events were

aligned on the start of their EPSP and (<1 mV) differences in the resting membrane potential between events

were subtracted for the display. Minimum interval between events was 20 ms. Red trace: average of >1000

events. Resting potential was -64 mV. Arrow points at the small prespike. Vertical scale bar: 10 mV. Horizontal:

1 ms. (B) As (A), showing closeup of the prespike (arrow). Vertical scale bar: 0.2 mV. Horizontal: 0.2 ms.

Discussion

Here, we studied how electrical currents that are needed to depolarize a presynaptic terminal or trigger transmitter release are dissipated in the synaptic cleft. We derived equations for the ephaptic coupling between the presynaptic terminal and the postsynaptic cell. We found that the capacitive component of the prespike is closely approximated by the second time derivative of the presynaptic action potential. Its size scales with the fourth power of the radius of the synapse, explaining why intracellularly recorded prespikes are uncommon in the CNS. We showed that presynaptic calcium currents can contribute to the prespike and that their contribution is closely approximated by the scaled first derivative of these currents. We used their contribution to infer an estimate of the cleft conductance, which was ~1 µS for the juvenile calyx of Held synapse. We showed that the cleft conductance of a radially-symmetric synapse does not depend on its radius, but is entirely defined by its height and the extracellular resistivity. This led to an estimate of the cleft resistivity that is at most 50% larger than of the interstitial fluid, which means that the cleft conductance is large enough to allow for the large EPSCs of the calyx of Held, but small enough to have a potential impact on the opening of presynaptic voltage-dependent calcium channels. Finally, we showed that developmental fenestration of the calyx of Held constitutes a very effective adaptation to minimize synaptic cleft potentials.

Comparison of cleft electrical properties across synapses

Using the calcium component of the prespike, we estimated a cleft conductance of about 1 µS. This value is subject to some uncertainty. For the calculation we assumed that most, but not all, presynaptic calcium channels are facing the synaptic cleft. The juvenile calyx contains N- and R-channels, which are not located near the release sites based on immunostaining and their lower efficacy at triggering release, which is in contrast to the P/Q-type calcium channels [45]. Our assumption that only two-third of the

calcium current originated from the cleft is therefore reasonable, considering that the N- and R-channels

make up about half of the calcium currents of the juvenile calyx.

Even though we calculated the AP-related capacitive currents that run through the cleft-facing membrane,

we did not record the AP itself at the cleft-facing membrane. The AP passively invades the terminal [60],

and therefore may be slowed down and have a reduced amplitude at the cleft-facing membrane. We

consider it unlikely that this filtering is significant for two reasons. First, passive filtering from one end to

the other end of the calyx reduces the AP peak by only 6 mV with a delay of <0.2 ms [61]. Second, the

axonal AP is able to invade the fenestrated endings of the mature calyx [62], which should have a

significantly higher impedance than the cup-shaped calyx. Our calculation of the presynaptic capacitive

currents as the rate of change of the recorded calyceal AP relative to the cleft potential is therefore likely

to be accurate.

Interestingly, with similar values for cleft height and extracellular resistivity other studies came to a wide

range of estimated values for the cleft conductance, both at least an order of magnitude smaller than 1 µS

[11, 26, 28-30, 63], but in some cases also considerably larger than 1 µS [7, 64]. In some of the synapses,

an unusually large and narrow path to the interstitial fluid was responsible for the low estimates. Examples

are mossy fiber boutons penetrated by a thin (<1 µm) spinule [28], the unusually long calyx-type

vestibular synapse [11], or the cone-to-horizontal cell synapse, for which a large part of the horizontal cell

is located within an invagination of the cone photoreceptors [65]. While cleft resistivity should be

independent of morphology, the cleft conductance for membrane currents at the deep end of these

synapses would be significantly less than for the homogeneously distributed current addressed here. In

these morphologically complex synapses, slow removal of ions or neurotransmitter may serve a function,

for example in the case of vestibular hair cells, where potassium ions accumulate [10, 11], or in the cone

photoreceptor synapse, where prolonged effects of glutamate are important, and where postsynaptic

currents are typically relatively small [66]. At the other end of the range lies the high estimate for the cleft

conductance of the neuromuscular junction, to which the postjunctional folds were suggested to contribute

[7]. The unique morphological properties of these synapses have therefore made a large contribution to the

large range of estimated values for the cleft conductance.

Our calculations indicated that for homogeneously-distributed currents the associated cleft conductance is

fully determined by the extracellular resistivity, the cleft height and the morphological type of the synapse,

but not by the size of the synapse. The cleft height is constrained within narrow boundaries because of the

opposing requirements of low volume for high neurotransmitter concentrations and high conductance to

allow large synaptic currents [16]. The cleft height typically ranges across synapses between 10-30 nm,

but there is some uncertainty regarding the impact of chemical fixation. Assuming a cleft height of 30 nm,

we estimated a resistivity of 75 Ω cm based on the prespike, only about 30% larger than of cerebrospinal

fluid or Ringer solution [16, 67]. While we are not aware of other estimates of the extracellular resistivity

within the synaptic cleft, our estimate is similar to the intracellular resistivity of neurons [68-70] and the

periaxonal resistivity of myelinated axons [70]. The extracellular resistivity can also be estimated based on

the diffusional mobility of small molecules. Our estimate agrees with those made with an imaging method

in hippocampal mossy fiber synapses, which also estimated that small molecules move on average about

50% slower than in the Ringer solution [71]. Estimates of neurotransmitter diffusion constants within the

synaptic cleft range from 1.5 to 4 times smaller than in free solution [72-74], but for neurotransmitters

diffusion within the synaptic cleft may be slowed down by neurotransmitter-binding proteins. We

therefore suggest that a cleft resistivity of ~75 Ωcm may apply to other synapses as well. Given the range

of cleft heights and the estimated cleft resistivity, the cleft conductance can be inferred for a distributed

current using our analytical relations (B3.2). It should be not so different from the calyx of Held

considering that cleft height is conserved across synapses, while the geometry of the juvenile calyx

translates well to bouton synapses by simple downscaling. Cleft conductances for circular synapses would

then range between 0.3-1.0 μS.

The prespike: what is it good for?

The extracellular AP of the calyx of Held can be observed in extracellular recordings at the target neuron

[75], and the characteristic shape of the complex waveform consisting of the presynaptic AP, the EPSP

and the postsynaptic AP has facilitated identification of *in vivo* single unit recordings at the MNTB [59].

Importantly, it was shown that the calyceal AP can be picked up in postsynaptic whole-cell recordings as

well [34]. In postsynaptic voltage clamp recordings the prespike showed a resemblance to a high-pass

filtered version of the presynaptic AP [36], but we showed here that it is closer to the second derivative of

the AP. At the related endbulb of Held synapse, the presynaptic AP is also readily picked up in

extracellular recordings [76]. However, no prespikes were reported during postsynaptic whole-cell

recordings [77] possibly due to the smaller size of the endbulb compared to the calyx of Held. In the

visual system, prespikes can be seen in a subset of lateral geniculate neurons [78]. However, in many

synapses in which the presynaptic AP is readily observed in extracellular recordings, it is not reported in

intracellular postsynaptic recordings [18, 79-83], attesting to the efficiency of the synaptic cleft in

dissipating these currents. Interestingly, especially in the hippocampus, neurons can display so-called

spikelets with an amplitude of mV in intracellular recordings. They have a biphasic shape resembling the

calyceal prespike recorded in current-clamp. These spikelets are thought to result from ephaptic,

capacitive coupling with one or more nearby neurons [84]. The equations we derived here should be

applicable to these spikelets as well: we predict them to resemble the second derivative of the AP in the

nearby neurons [85], and to critically depend on contact area, but also on proximity and on the resistivity

of the extracellular fluid.

We believe that the major reason for the lack of prespikes at most synapses is that the prespike amplitude

is proportional to the 4^{th} power of the synapse radius (Appendix B2). During the postnatal development of

the MNTB, it can therefore be argued that the absence of postsynaptic recordings with multiple prespikes

cannot be used as firm evidence against the presence of multiple calyces on these cells. We previously

found anecdotal evidence for the presence of multiple large inputs during early postnatal development

[86]. In a small fraction of the recordings multiple large inputs were observed, one with and one without a

prespike. It might be that the one without the prespike had a smaller surface area or a higher cleft height,

going below the radar. This highly nonlinear dependence on apposition area may thus provide an

explanation for the discrepancy between anatomical evidence favoring multiple large inputs during

development [87] and the lack of recordings with multiple prespikes.

Divide and conquer

Virtually all synapses have in common that neurotransmission is initiated by a presynaptic action potential

that results in capacitive cleft currents. The size of these currents depends on the amplitude and speed of

the presynaptic AP. The properties of the presynaptic APs have mostly been studied at giant synapses,

where they tend to be large and fast [46, 88-93]. These giant synapses are typically relay synapses that are

specialized for rapid signaling, but recent measurements at more conventional bouton synapses in the

neocortex show that their presynaptic action potentials are large and fast as well [94]. These capacitive

currents, together with the presynaptic ion channel currents, will lead to a change in the cleft potential

whose time course will closely follow the time course of the presynaptic membrane currents, and whose

amplitude will critically depend on the cleft resistivity. Because of this cleft potential, the presynaptic ion

channels facing the cleft will sense a reduced action potential. We showed that this should result in a less

effective and delayed opening of the presynaptic calcium channels facing the synaptic cleft. Since rapid

and large changes in the voltage sensed by the presynaptic ion channels serve to rapidly turn transmitter

release on and off again, large cleft potentials seem undesirable, even though they may serve other, more

beneficial functions such as influencing the neurotransmitter mobility within the cleft [15, 16], or rapidly

inhibiting postsynaptic action potential generation [18, 95].

Above, we argued that many radial-like synapses should have a cleft conductance similar to what was

measured here. An important difference between small and large synapses, however, is that the total

current will be much larger in the large synapse, as the capacitive current scales with the presynaptic

apposition area. A large synapse thus has to dissipate this much larger current over a cleft conductance

that is similar to the cleft conductance of a small synapse. There are three obvious different neural

strategies to mitigate significant cleft potentials. Firstly, a strategy may be widenings of the synaptic cleft

in between synaptic junctions. These intrasynaptic space enlargements [4], also called channels of

extracellular spaces [96, 97] or subsynaptic cavities [98], have been found at many synapses, including the

calyx of Held synapse, and may increase cleft conductance. Interestingly, a knockout of brevican, a

component of the extracellular matrix of perineuronal nets, leads to a reduction of the size of the

subsynaptic cavities at the calyx of Held synapse and an increased synaptic delay, even though the relation

with a change in cleft conductance remains to be established [98]. The second strategy is to divide the

cleft membrane currents over many postsynaptic junctions. For example, in the case of the squid giant

synapse, the postsynaptic axon has numerous processes contacting the presynaptic axon, each with a width

that is only ~1 μm [99]. At each of these processes the cleft potential will be small, thus effectively

mitigating its impact. The last strategy is to fenestrate presynaptically. Most large synapses fenestrate

presynaptically during development, leading to extended extracellular spaces [97]. Well-known examples

include the neuromuscular junction, where the terminal arbor assumes a pretzel shape [100], the calyx of

Held, which develops from a cup shape to the calyx shape [52, 53], the endbulb of Held synapse [54] or

the avian ciliary ganglion [55]. In Appendix B5 we quantified the impact this will have on the size of the

cleft potential change and on the prespike. Previously, other roles for these adaptations have been

emphasized, such as speeding up of transmitter clearance or minimization of crosstalk between

postsynaptic densities, both for the calyx of Held synapse [52, 101, 102] and at other synapses [103, 104].

The digitated adult morphology of the calyx of Held is accompanied by an increase in the number of

puncta adhaerentia [102] and the formation of swellings connected via necks to the fingers [51, 105]. In

the adult calyx of Held, a large fraction of the release takes place at these swellings, which should

facilitate the dissipation of capacitive currents by the cleft, and reduce the impact of the cleft conductance

on the postsynaptic currents.

In conclusion, we show here that the prespike of the calyx of Held synapse can be used to infer critical

properties of its synaptic cleft; apart from the cleft conductance, it may also provide information about the

presence or absence of ion channels at the cleft, or the effectiveness of morphological changes in reducing

the size of cleft potentials during development. We thus provide a quantitative description for how the

synaptic cleft, despite its narrow, limited space, can sustain the large and fast currents that are at the heart

of chemical neurotransmission.

Materials & Methods

Analytical solutions for the transfer functions were derived using phasors. Analytical solutions for the

Fourier transformation were checked using Wolfram Alpha. The derivations are outlined in Appendix A.

Phasor calculations were done in Igor Pro 6.37 (WaveMetrics Inc., Lake Oswego, Oregon) using built-in

functions: cmplx, FFT, IFFT and MatrixOP. The AP template shown in Figure 3A was based on an AP

template previously used in voltage-clamp experiments [42] that was smoothed to eliminate the high-

frequency oscillations that can arise after inverse Fourier transformation.

For the experimental procedures and recording conditions for the dual recordings of the calcium current

and the postsynaptic currents, we refer the reader to the original paper [43]. *In vivo* whole-cell current-

clamp recordings of developing MNTB principal neurons with afferent stimulation were previously

reported in Sierksma, Slotman (37) (n = 6). *In vivo* whole-cell current-clamp recordings of MNTB

principal neurons from adult C57BL/6J mice were previously reported in Wang, van Woerden (58) (n = 7,

only wild-type animals) or in Lorteije, Rusu (59) (n=5). We also refer to these papers for the respective

ethics statements.

For numerical simulations with voltage-gated channels the following two differential equations were

defined:

$$584 \qquad c_{post} \frac{dv_{post}}{dt} = c_{cl} \frac{dv_{cl}}{dt} - g_{post}(v_{post} - v_{rest})$$

$$585 \qquad 2c_{cl} \frac{dv_{cl}}{dt} = c_{cl} \left(\frac{dv_{pre}}{dt} + \frac{dv_{post}}{dt} \right) - g_{cl}v_{cl} + \sum g_{max}p_{open}(v_{pre} - v_{cl} - v_{channel})$$

Where g_{max} represents the maximal conductance of a voltage-gated channel, p_{open} the open probability of

the voltage-gated channels, which is calculated using a multiple-state model (see below), and $v_{channel}$ the

reversal potential of the voltage-gated channel. The reversal potentials were 50 mV, -90 mV, 40 mV and -

70 mV for the sodium conductance, potassium conductance, calcium conductance, and the resting

membrane potential of the postsynaptic neuron, respectively. v_{pre} was defined by an AP waveform.

Voltage-gated conductances were modeled with multiple states following Yang, Wang (106) except for

the calcium conductance. In the models C indicates a closed state, O the open state and I the inactivated

state. The sodium channel followed a five states-model, the high-threshold and low-threshold potassium

channel were modeled as a six-states channel, and the calcium channel as a two-state channel for which

the open probability was squared:

Five states-model:
$$C_1 \underset{\beta}{\overset{2\alpha}{\rightleftarrows}} C_2 \underset{2\beta}{\overset{\alpha}{\rightleftarrows}} C_3 \underset{\delta}{\overset{\gamma}{\rightleftarrows}} O \underset{\epsilon}{\overset{\theta}{\rightleftarrows}} I$$

597   Six states-model:
$$C_1 \underset{\beta}{\overset{4\alpha}{\rightleftarrows}} C_2 \underset{2\beta}{\overset{3\alpha}{\rightleftarrows}} C_3 \underset{3\beta}{\overset{2\alpha}{\rightleftarrows}} C_4 \underset{4\beta}{\overset{\alpha}{\rightleftarrows}} C_5 \underset{\delta}{\overset{\gamma}{\rightleftarrows}} O$$

598   Two states-model:
$$C \underset{\beta}{\overset{\alpha}{\rightleftarrows}} O$$

599   The rate constants indicated by the Greek letters in the state models were calculated as

600
$$k = a\, e^{b(v_{pre} - v_{cl})}$$

where k represents a particular rate constant, and *a* and *b* are channel-specific parameters listed in Table 1.

To take into account a possible effect of the cleft potential on the channel kinetics, the rate constant was

calculated using the voltage difference between the presynaptic potential and the cleft potential.

The ordinary differential equations were numerically solved in Igor using IntegrateODE with default

settings.

Statistics are reported as mean and standard deviation. Pearson's correlation coefficient was calculated to

quantify correlations.

Table 1. Rate constants for the voltage-gated channels			
		a (ms^{-1})	**b (mV^{-1})**
Sodium conductance	**α**	48.9	1/128.4
	β	1.46	-1/8.9
	γ	392.8	1/36.0
	δ	0.73	-1/15.5
	ε	7.8	1/35.9
	θ	0.01	-1/18
High-threshold potassium conductance	**α**	1.097	1/57.404
	β	0.794	-1/79.264
	γ	33.750	0
	δ	74.360	0
Low-threshold potassium conductance	**α**	1.204	1/37.574
	β	0.360	-1/230
	γ	245.488	0
	δ	132.566	0
Calcium conductance	**α**	1.78	1/23.3
	β	0.14	-1/15
Parameters were taken from Yang, Wang (106), except for the calcium conductance [44].			

609   **Acknowledgements**

610   We thank Dr. Henrique von Gersdorff for helpful discussions on the developmental changes in the

611   prespike.

References

1.   Foster M. A text-book of physiology: Lea Brothers & Company; 1895.

2.   Bennett MR. The early history of the synapse: from Plato to Sherrington. Brain Research Bulletin.
1999;50(2):95-118. doi: 10.1016/S0361-9230(99)00094-5.

3.   Sotelo C. The history of the synapse. The Anatomical Record. 2020;303(5):1252-79. doi:
10.1002/ar.24392.

4.   Palay SL. Synapses in the central nervous system. The Journal of Cell Biology. 1956;2(4):193-202.
doi: 10.1083/jcb.2.4.193.

5.   De Robertis E. Submicroscopic changes of the synapse after nerve section in the acoustic
ganglion of the guinea pig; an electron microscope study. The Journal of Biophysical and Biochemical
Cytology. 1956;2(5):503-12. doi: 10.1083/jcb.2.5.503.

6.   Harris KM, Weinberg RJ. Ultrastructure of synapses in the mammalian brain. Cold Spring Harbor
perspectives in biology. 2012;4(5):a005587. doi: 10.1101/cshperspect.a005587.

7.   Eccles JC, Jaeger JC. The relationship between the mode of operation and the dimensions of the
junctional regions at synapses and motor end-organs. Proceedings of the Royal Society of London Series
B-Biological Sciences. 1958;148(930):38-56. doi: 10.1098/rspb.1958.0003.

8.   Cho S, von Gersdorff H. Proton-mediated block of Ca^{2+} channels during multivesicular release
regulates short-term plasticity at an auditory hair cell synapse. The Journal of Neuroscience.
2014;34(48):15877-87. doi: 10.1523/jneurosci.2304-14.2014.

9.   DeVries SH. Exocytosed protons feedback to suppress the Ca^{2+} current in mammalian cone
photoreceptors. Neuron. 2001;32(6):1107-17. doi: 10.1016/S0896-6273(01)00535-9.

10.   Contini D, Price SD, Art JJ. Accumulation of K^+ in the synaptic cleft modulates activity by
influencing both vestibular hair cell and calyx afferent in the turtle. The Journal of Physiology.
2017;595(3):777-803. doi: 10.1113/JP273060.

11.   Goldberg JM. Theoretical analysis of intercellular communication between the vestibular type I
hair cell and its calyx ending. Journal of Neurophysiology. 1996;76(3):1942-57. doi:
10.1152/jn.1996.76.3.1942. PubMed PMID: 8890305.

12.   Shih PY, Savtchenko LP, Kamasawa N, Dembitskaya Y, McHugh TJ, Rusakov DA, et al. Retrograde
synaptic signaling mediated by K^+ efflux through postsynaptic NMDA receptors. Cell Rep. 2013;5(4):941-
51. doi: 10.1016/j.celrep.2013.10.026. PubMed PMID: 24268779.

13.   Borst JGG, Sakmann B. Depletion of calcium in the synaptic cleft of a calyx-type synapse in the
rat brainstem. Journal of Physiology. 1999;521:123-33. doi: 10.1111/j.1469-7793.1999.00123.x.

14.   Stanley EF. Presynaptic calcium channels and the depletion of synaptic cleft calcium ions. J
Neurophysiol. 2000;83(1):477-82. doi: 10.1152/jn.2000.83.1.477. PubMed PMID: 10634889.

15.   Sylantyev S, Savtchenko LP, Niu Y-P, Ivanov AI, Jensen TP, Kullmann DM, et al. Electric fields due
to synaptic currents sharpen excitatory transmission. Science. 2008;319(5871):1845-9. doi:
10.1126/science.1154330.

16.   Savtchenko LP, Rusakov DA. The optimal height of the synaptic cleft. Proceedings of the National
Academy of Sciences. 2007;104(6):1823. doi: 10.1073/pnas.0606636104.

17.   Savtchenko LP, Poo MM, Rusakov DA. Electrodiffusion phenomena in neuroscience: a neglected
companion. Nat Rev Neurosci. 2017;18(10):598-612. doi: 10.1038/nrn.2017.101. PubMed PMID:
28924257.

18.   Blot A, Barbour B. Ultra-rapid axon-axon ephaptic inhibition of cerebellar Purkinje cells by the
pinceau. Nat Neurosci. 2014;17(2):289-95. Epub 2014/01/15. doi: 10.1038/nn.3624. PubMed PMID:
24413696.

19.   Rusakov DA, Savtchenko LP, Zheng K, Henley JM. Shaping the synaptic signal: molecular mobility
inside and outside the cleft. Trends in Neurosciences. 2011;34(7):359-69. doi:
10.1016/j.tins.2011.03.002.
20.   Savtchenko LP, Antropov SN, Korogod SM. Effect of voltage drop within the synaptic cleft on the
current and voltage generated at a single synapse. Biophysical Journal. 2000;78(3):1119-25. doi:
10.1016/S0006-3495(00)76670-7.
21.   Berretta N, Rossokhin AV, Kasyanov AM, Sokolov MV, Cherubini E, Voronin LL. Postsynaptic
hyperpolarization increases the strength of AMPA-mediated synaptic transmission at large synapses
between mossy fibers and CA3 pyramidal cells. Neuropharmacology. 2000;39(12):2288-301. doi:
10.1016/s0028-3908(00)00076-9. PubMed PMID: 10974312.
22.   Faber DS, Funch PG. Differential properties of orthodromic and antidromic impulse propagation
across the Mauthner cell initial segment. Brain Research. 1980;190(1):255-60. doi: 10.1016/0006-
8993(80)91176-2.
23.   del Castillo J, Katz B. Localization of active spots within the neuromuscular junction of the frog.
The Journal of Physiology. 1956;132(3):630-49. doi: 10.1113/jphysiol.1956.sp005554.
24.   Martin AR, Pilar G. An analysis of electrical coupling at synapses in the avian ciliary ganglion. J
Physiol. 1964;171:454-75. doi: 10.1113/jphysiol.1964.sp007390. PubMed PMID: 14193934; PubMed
Central PMCID: PMCPMC1368844.
25.   Dmitriev AV, Mangel SC. Electrical feedback in the cone pedicle: A computational analysis.
Journal of Neurophysiology. 2006;95(3):1419-27. doi: 10.1152/jn.00098.2005. PubMed PMID: 16319220.
26.   Fahrenfort I, Steijaert M, Sjoerdsma T, Vickers E, Ripps H, van Asselt J, et al. Hemichannel-
mediated and pH-based feedback from horizontal cells to cones in the vertebrate retina. PLoS One.
2009;4(6):e6090. doi: 10.1371/journal.pone.0006090. PubMed PMID: 19564917; PubMed Central
PMCID: PMCPMC2699542.
27.   Kamermans M, Fahrenfort I, Schultz K, Janssen-Bienhold U, Sjoerdsma T, Weiler R. Hemichannel-
mediated inhibition in the outer retina. Science. 2001;292(5519):1178-80. doi: 10.1126/science.1060101.
PubMed PMID: 11349152.
28.   Kasyanov AM, Maximov VV, Byzov AL, Berretta N, Sokolov MV, Gasparini S, et al. Differences in
amplitude-voltage relations between minimal and composite mossy fibre responses of rat CA3
hippocampal neurons support the existence of intrasynaptic ephaptic feedback in large synapses.
Neuroscience. 2000;101(2):323-36. doi: 10.1016/S0306-4522(00)00366-3. PubMed PMID: 11074156.
29.   Byzov AL, Shura-Bura TM. Electrical feedback mechanism in the processing of signals in the outer
plexiform layer of the retina. Vision Res. 1986;26(1):33-44. doi: 10.1016/0042-6989(86)90069-6. PubMed
PMID: 3012877.
30.   Savtchenko LP. Bilateral processing in chemical synapses with electrical 'ephaptic' feedback: a
theoretical model. Math Biosci. 2007;207(1):113-37. doi: 10.1016/j.mbs.2006.09.016. PubMed PMID:
17112549.
31.   Morest DK. The growth of synaptic endings in the mammalian brain: a study of the calyces of the
trapezoid body. Zeitschrift für Anatomie und Entwicklungsgeschichte. 1968;127(3):201-20. doi:
10.1007/BF00526129.
32.   Rollenhagen A, Lübke JHR. The morphology of excitatory central synapses: from structure to
function. Cell Tissue Res. 2006;326(2):221-37. doi: 10.1007/s00441-006-0288-z. PubMed PMID:
16932936.
33.   Borst JGG, Soria van Hoeve J. The calyx of Held synapse: from model synapse to auditory relay.
Annual Review of Physiology. 2012;74:199-224. Epub 2011/11/01. doi: 10.1146/annurev-physiol-
020911-153236. PubMed PMID: 22035348.

34.   Forsythe ID. Direct patch recording from identified presynaptic terminals mediating
glutamatergic EPSCs in the rat CNS, *in vitro*. J Physiol. 1994;479(3):381-7. doi:
10.1113/jphysiol.1994.sp020303.
35.   von Gersdorff H, Borst JGG. Short-term plasticity at the calyx of Held. Nature Reviews
Neuroscience. 2002;3:53-64.
36.   Borst JGG, Helmchen F, Sakmann B. Pre- and postsynaptic whole-cell recordings in the medial
nucleus of the trapezoid body of the rat. J Physiol. 1995;489(Pt 3):825-40. doi:
10.1113/jphysiol.1995.sp021095.
37.   Sierksma MC, Slotman JA, Houtsmuller AB, Borst JGG. Structure–function relation of the
developing calyx of Held synapse *in vivo*. The Journal of Physiology. 2020;598(20):4603-19. doi:
10.1113/JP279976.
38.   Korogod N, Petersen CCH, Knott GW. Ultrastructural analysis of adult mouse neocortex
comparing aldehyde perfusion with cryo fixation. Elife. 2015;4. doi: 10.7554/eLife.05793. PubMed PMID:
26259873; PubMed Central PMCID: PMCPMC4530226.
39.   Syková E, Nicholson C. Diffusion in brain extracellular space. Physiol Rev. 2008;88(4):1277-340.
doi: 10.1152/physrev.00027.2007. PubMed PMID: 18923183.
40.   Rubio ME, Nagy JI. Connexin36 expression in major centers of the auditory system in the CNS of
mouse and rat: Evidence for neurons forming purely electrical synapses and morphologically mixed
synapses. Neuroscience. 2015;303:604-29. Epub 2015/07/19. doi: 10.1016/j.neuroscience.2015.07.026.
PubMed PMID: 26188286; PubMed Central PMCID: PMC4576740.
41.   Meinrenken CJ, Borst JGG, Sakmann B. Calcium secretion coupling at calyx of Held governed by
nonuniform channel-vesicle topography. J Neurosci. 2002;22(5):1648-67. doi: 10.1523/JNEUROSCI.22-05-
01648.2002. PubMed PMID: 11880495.
42.   Borst JGG, Sakmann B. Calcium influx and transmitter release in a fast CNS synapse. Nature.
1996;383(6599):431-4. doi: 10.1038/383431a0.
43.   Borst JGG, Sakmann B. Effect of changes in action potential shape on calcium currents and
transmitter release in a calyx-type synapse of the rat auditory brainstem. Philos Trans R Soc Lond B Biol
Sci. 1999;354(1381):347-55. doi: 10.1098/rstb.1999.0386. PubMed PMID: 10212483.
44.   Borst JGG, Sakmann B. Calcium current during a single action potential in a large presynaptic
terminal of the rat brainstem. J Physiol. 1998;506:143-57. doi: 10.1111/j.1469-7793.1998.143bx.x.
45.   Wu L-G, Westenbroek RE, Borst JGG, Catterall WE, Sakmann B. Calcium channel types with
distinct presynaptic localization couple differentially to transmitter release in single calyx-type synapses.
Journal of Neuroscience. 1999;19:726-36. doi: 10.1523/JNEUROSCI.19-02-00726.1999
46.   Taschenberger H, von Gersdorff H. Fine-tuning an auditory synapse for speed and fidelity:
developmental changes in presynaptic waveform, EPSC kinetics, and synaptic plasticity. J Neurosci.
2000;20(24):9162-73. doi: 10.1523/JNEUROSCI.20-24-09162.2000 PubMed PMID: 11124994.
47.   Barnes-Davies M, Forsythe ID. Pre- and postsynaptic glutamate receptors at a giant excitatory
synapse in rat auditory brainstem slices. Journal of Physiology. 1995;488(2):387-406. doi:
10.1113/jphysiol.1995.sp020974.
48.   Forsythe ID, Barnes-Davies M. The binaural auditory pathway: excitatory amino acid receptors
mediate dual timecourse excitatory postsynaptic currents in the rat medial nucleus of the trapezoid
body. Proceedings of the Royal Society B. 1993;251(1331):151-7. doi: 10.1098/rspb.1993.0022.
49.   Joshi I, Wang L-Y. Developmental profiles of glutamate receptors and synaptic transmission at a
single synapse in the mouse auditory brainstem. J Physiol. 2002;540(Pt 3):861-73. doi:
10.1113/jphysiol.2001.013506. PubMed PMID: 11986375.
50.   Sätzler K, Söhl LF, Bollmann JH, Borst JGG, Frotscher M, Sakmann B, et al. Three-dimensional
reconstruction of a calyx of Held and its postsynaptic principal neuron in the medial nucleus of the
trapezoid body. J Neurosci. 2002;22(24):10567-79. PubMed PMID: 12486149.

51.   Rowland KC, Irby NK, Spirou GA. Specialized synapse-associated structures within the calyx of
Held. J Neurosci. 2000;20(24):9135-44. doi: 10.1523/JNEUROSCI.20-24-09135.2000. PubMed PMID:
11124991.
52.   Ford MC, Grothe B, Klug A. Fenestration of the calyx of Held occurs sequentially along the
tonotopic axis, is influenced by afferent activity, and facilitates glutamate clearance. J Comp Neurol.
2009;514(1):92-106. PubMed PMID: 19260071.
53.   Kandler K, Friauf E. Pre- and postnatal development of efferent connections of the cochlear
nucleus in the rat. Journal of Comparative Neurology. 1993;328(2):161-84. doi: 10.1002/cne.903280202.
54.   Ryugo DK, Montey KL, Wright AL, Bennett ML, Pongstaporn T. Postnatal development of a large
auditory nerve terminal: the endbulb of Held in cats. Hear Res. 2006;216-217:100-15. Epub 2006/02/25.
doi: 10.1016/j.heares.2006.01.007. PubMed PMID: 16497457.
55.   Hess A, Pilar G, Weakly JN. Correlation between transmission and structure in avian ciliary
ganglion synapses. J Physiol. 1969;202(2):339-54. doi: 10.1113/jphysiol.1969.sp008815. PubMed PMID:
4306543; PubMed Central PMCID: PMCPMC1351486.
56.   Baydyuk M, Xu J, Wu LG. The calyx of Held in the auditory system: Structure, function, and
development. Hear Res. 2016;338:22-31. Epub 2016/03/29. doi: 10.1016/j.heares.2016.03.009. PubMed
PMID: 27018297; PubMed Central PMCID: PMC4967386.
57.   Spirou GA, Chirila FV, von Gersdorff H, Manis PB. Heterogeneous Ca^{2+} influx along the adult calyx
of Held: a structural and computational study. Neuroscience. 2008;154(1):171-85. PubMed PMID:
18485608.
58.   Wang T, van Woerden GM, Elgersma Y, Borst JGG. Enhanced transmission at the calyx of Held
synapse in a mouse model for Angelman syndrome. Front Cell Neurosci. 2017;11:418. Epub 2018/01/23.
doi: 10.3389/fncel.2017.00418. PubMed PMID: 29354033; PubMed Central PMCID: PMC5758499.
59.   Lorteije JAM, Rusu SI, Kushmerick C, Borst JGG. Reliability and precision of the mouse calyx of
Held synapse. J Neurosci. 2009;29(44):13770-84. doi: 10.1523/JNEUROSCI.3285-09.2009. PubMed PMID:
19889989.
60.   Leão RM, Kushmerick C, Pinaud R, Renden R, Li G-L, Taschenberger H, et al. Presynaptic Na$^+$
channels: locus, development, and recovery from inactivation at a high-fidelity synapse. J Neurosci.
2005;25(14):3724-38. doi: 10.1523/JNEUROSCI.3983-04.2005. PubMed PMID: 15814803.
61.   Yang Y-M, Wang L-Y. Amplitude and kinetics of action potential-evoked Ca^{2+} current and its
efficacy in triggering transmitter release at the developing calyx of Held synapse. J Neurosci.
2006;26(21):5698-708. doi: 10.1523/JNEUROSCI.4889-05.2006 PubMed PMID: 16723526.
62.   Fekete A, Nakamura Y, Yang YM, Herlitze S, Mark MD, DiGregorio DA, et al. Underpinning
heterogeneity in synaptic transmission by presynaptic ensembles of distinct morphological modules. Nat
Commun. 2019;10(1):826. Epub 2019/02/20. doi: 10.1038/s41467-019-08452-2. PubMed PMID:
30778063; PubMed Central PMCID: PMC6379440.
63.   Martin AR, Pilar G. Transmission through the ciliary ganglion of the chick. Journal of Physiology.
1963;168:464-75.
64.   Funch PG, Faber DS. Action-potential propagation and orthodromic impulse initiation in
Mauthner axon. Journal of Neurophysiology. 1982;47(6):1214-31. doi: 10.1152/jn.1982.47.6.1214
65.   Vandenbranden CA, Verweij J, Kamermans M, Muller LJ, Ruijter JM, Vrensen GF, et al. Clearance
of neurotransmitter from the cone synaptic cleft in goldfish retina. Vision Research. 1996;36(24):3859-
74. doi: 10.1016/s0042-6989(96)00134-4.
66.   DeVries SH, Li W, Saszik S. Parallel processing in two transmitter microenvironments at the cone
photoreceptor synapse. Neuron. 2006;50(5):735-48. doi: 10.1016/j.neuron.2006.04.034. PubMed PMID:
16731512.

67.   Baumann SB, Wozny DR, Kelly SK, Meno FM. The electrical conductivity of human cerebrospinal
fluid at body temperature. IEEE Trans Biomed Eng. 1997;44(3):220-3. doi: 10.1109/10.554770. PubMed
PMID: 9216137.
68.   Roth A, Häusser M. Compartmental models of rat cerebellar Purkinje cells based on
simultaneous somatic and dendritic patch-clamp recordings. J Physiol. 2001;535(Pt 2):445-72. doi:
10.1111/j.1469-7793.2001.00445.x. PubMed PMID: 11533136.
69.   Golding NL, Mickus TJ, Katz Y, Kath WL, Spruston N. Factors mediating powerful voltage
attenuation along CA1 pyramidal neuron dendrites. J Physiol. 2005;568(Pt 1):69-82. doi:
10.1113/jphysiol.2005.086793. PubMed PMID: 16002454; PubMed Central PMCID: PMCPMC1474764.
70.   Cohen CCH, Popovic MA, Klooster J, Weil MT, Mobius W, Nave KA, et al. Saltatory conduction
along myelinated axons involves a periaxonal nanocircuit. Cell. 2020;180(2):311-22 e15. doi:
10.1016/j.cell.2019.11.039. PubMed PMID: 31883793; PubMed Central PMCID: PMCPMC6978798.
71.   Zheng K, Jensen TP, Savtchenko LP, Levitt JA, Suhling K, Rusakov DA. Nanoscale diffusion in the
synaptic cleft and beyond measured with time-resolved fluorescence anisotropy imaging. Scientific
Reports. 2017;7(1):42022. doi: 10.1038/srep42022.
72.   Land BR, Harris WV, Salpeter EE, Salpeter MM. Diffusion and binding constants for acetylcholine
derived from the falling phase of miniature endplate currents. Proc Natl Acad Sci U S A. 1984;81(5):1594-
8. doi: 10.1073/pnas.81.5.1594. PubMed PMID: 6584895; PubMed Central PMCID: PMCPMC344884.
73.   Faber DS, Funch PG, Korn H. Evidence that receptors mediating central synaptic potentials
extend beyond the postsynaptic density. Proc Natl Acad Sci U S A. 1985;82(10):3504-8. doi:
10.1073/pnas.82.10.3504. PubMed PMID: 2582417; PubMed Central PMCID: PMCPMC397805.
74.   Nielsen TA, DiGregorio DA, Silver RA. Modulation of glutamate mobility reveals the mechanism
underlying slow-rising AMPAR EPSCs and the diffusion coefficient in the synaptic cleft. Neuron.
2004;42(5):757-71. doi: 10.1016/j.neuron.2004.04.003. PubMed PMID: 15182716.
75.   Guinan JJ, Jr., Li RY-S. Signal processing in brainstem auditory neurons which receive giant
endings (calyces of Held) in the medial nucleus of the trapezoid body of the cat. Hear Res. 1990;49(1-
3):321-34. doi: 10.1016/0378-5955(90)90111-2.
76.   Pfeiffer RR. Anteroventral cochlear nucleus: wave forms of extracellularly recorded spike
potentials. Science. 1966;154(749):667-8. doi: 10.1126/science.154.3749.667 PubMed PMID: 5923782.
77.   Typlt M, Haustein MD, Dietz B, Steinert JR, Witte M, Englitz B, et al. Presynaptic and postsynaptic
origin of multicomponent extracellular waveforms at the endbulb of Held-spherical bushy cell synapse.
Eur J Neurosci. 2010;31(9):1574-81. Epub 2010/06/09. doi: 10.1111/j.1460-9568.2010.07188.x. PubMed
PMID: 20525070.
78.   Eysel UT. Quantitative studies of intracellular postsynaptic potentials in the lateral geniculate
nucleus of the cat with respect to optic tract stimulus response latencies. Exp Brain Res. 1976;25:469-86.
doi: 10.1007/bf00239782. PubMed PMID: 182514.
79.   Dudel J. Effect of inhibition on the presynaptic nerve terminal in the neuromuscular junction of
the crayfish. Nature. 1962;193:587-8. doi: 10.1038/193587b0. PubMed PMID: 13888196.
80.   Takeuchi A, Takeuchi N. Electrical changes in pre- and postsynaptic axons of the giant synapse of
Loligo. J Gen Physiol. 1962;45:1181-93. doi: 10.1085/jgp.45.6.1181. PubMed PMID: 13919241; PubMed
Central PMCID: PMCPMC2195243.
81.   Katz B, Miledi R. Propagation of electric activity in motor nerve terminals. Proceedings of the
Royal Society B. 1965;161:453-82. doi: 10.1098/rspb.1965.0015.
82.   Mathy A, Ho SSN, Davie JT, Duguid IC, Clark BA, Häusser M. Encoding of oscillations by axonal
bursts in inferior olive neurons. Neuron. 2009;62(3):388-99. Epub 2009/05/19. doi:
10.1016/j.neuron.2009.03.023. PubMed PMID: 19447094; PubMed Central PMCID: PMC2777250.

83.   Maruta J, Hensbroek RA, Simpson JI. Intraburst and interburst signaling by climbing fibers. J
Neurosci. 2007;27(42):11263-70. Epub 2007/10/19. doi: 10.1523/JNEUROSCI.2559-07.2007. PubMed
PMID: 17942720.
84.   Michalikova M, Remme MWH, Schmitz D, Schreiber S, Kempter R. Spikelets in pyramidal
neurons: generating mechanisms, distinguishing properties, and functional implications. Rev Neurosci.
2019;31(1):101-19. doi: 10.1515/revneuro-2019-0044. PubMed PMID: 31437125.
85.   Vigmond EJ, Perez Velazquez JL, Valiante TA, Bardakjian BL, Carlen PL. Mechanisms of electrical
coupling between pyramidal cells. J Neurophysiol. 1997;78(6):3107-16. doi: 10.1152/jn.1997.78.6.3107.
PubMed PMID: 9405530.
86.   Sierksma MC, Tedja MS, Borst JGG. *In vivo* matching of postsynaptic excitability with
spontaneous synaptic inputs during formation of the rat calyx of Held synapse. J Physiol.
2016;595(1):207-31. Epub 2016/07/19. doi: 10.1113/JP272780. PubMed PMID: 27426483.
87.   Holcomb PS, Hoffpauir BK, Hoyson MC, Jackson DR, Deerinck TJ, Marrs GS, et al. Synaptic inputs
compete during rapid formation of the calyx of Held: a new model system for neural development. J
Neurosci. 2013;33(32):12954-69. Epub 2013/08/09. doi: 10.1523/JNEUROSCI.1087-13.2013. PubMed
PMID: 23926251; PubMed Central PMCID: PMC3735879.
88.   Ritzau-Jost A, Delvendahl I, Rings A, Byczkowicz N, Harada H, Shigemoto R, et al. Ultrafast action
potentials mediate kilohertz signaling at a central synapse. Neuron. 2014;84(1):152-63. doi:
10.1016/j.neuron.2014.08.036.
89.   Sierksma MC, Borst JGG. Resistance to action potential depression of a rat axon terminal in vivo.
Proc Natl Acad Sci U S A. 2017;114(16):4249-54. Epub 2017/04/05. doi: 10.1073/pnas.1619433114.
PubMed PMID: 28373550; PubMed Central PMCID: PMC5402426.
90.   Lin KH, Oleskevich S, Taschenberger H. Presynaptic Ca^{2+} influx and vesicle exocytosis at the
mouse endbulb of Held: a comparison of two auditory nerve terminals. J Physiol. 2011;589(17):4301-20.
doi: 10.1113/jphysiol.2011.209189. PubMed PMID: 21746778; PubMed Central PMCID:
PMCPMC3180584.
91.   Rancz EA, Ishikawa T, Duguid I, Chadderton P, Mahon S, Häusser M. High-fidelity transmission of
sensory information by single cerebellar mossy fibre boutons. Nature. 2007;450(7173):1245-8. doi:
10.1038/nature05995. PubMed PMID: 18097412.
92.   Geiger JR, Jonas P. Dynamic control of presynaptic Ca^{2+} inflow by fast-inactivating K^+ channels in
hippocampal mossy fiber boutons. Neuron. 2000;28(3):927-39. PubMed PMID: 11163277.
93.   Bischofberger J, Geiger JRP, Jonas P. Timing and efficacy of Ca^{2+} channel activation in
hippocampal mossy fiber boutons. J Neurosci. 2002;22(24):10593-602. Epub 2002/12/18. doi:
10.1523/JNEUROSCI.22-24-10593.2002 PubMed PMID: 12486151.
94.   Ritzau-Jost A, Tsintsadze T, Krueger M, Ader J, Bechmann I, Eilers J, et al. Large, stable spikes
exhibit differential broadening in excitatory and inhibitory neocortical boutons. Cell Rep.
2021;34(2):108612. doi: 10.1016/j.celrep.2020.108612. PubMed PMID: 33440142; PubMed Central
PMCID: PMCPMC7809622.
95.   Furukawa T, Furshpan EJ. Two inhibitory mechanisms in the Mauthner neurons of goldfish. J
Neurophysiol. 1963;26:140-76. doi: 10.1152/jn.1963.26.1.140. PubMed PMID: 13945908.
96.   Tatsuoka H, Reese TS. New structural features of synapses in the anteroventral cochlear nucleus
prepared by direct freezing and freeze-substitution. Journal of Comparative Neurology. 1989;290(3):343-
57. doi: 10.1002/cne.902900304.
97.   Nakajima Y. Fine structure of the medial nucleus of the trapezoid body of the bat with special
reference to two types of synaptic endings. J Cell Biol. 1971;50(1):121-34. doi: 10.1083/jcb.50.1.121
98.   Blosa M, Sonntag M, Jäger C, Weigel S, Seeger J, Frischknecht R, et al. The extracellular matrix
molecule brevican is an integral component of the machinery mediating fast synaptic transmission at the

calyx of Held. J Physiol. 2015;593(19):4341-60. Epub 2015/08/01. doi: 10.1113/jp270849. PubMed PMID:
26223835.
99.   Martin R, Miledi R. The form and dimensions of the giant synapse of squids. Philosophical
Transactions of the Royal Society B. 1986;312(1155):355-77. doi: 10.1098/rstb.1986.0013.
100.   Sanes JR, Lichtman JW. Development of the vertebrate neuromuscular junction. Annu Rev
Neurosci. 1999;22:389-442. Epub 1999/04/15. doi: 10.1146/annurev.neuro.22.1.389. PubMed PMID:
10202544.
101.   Renden R, Taschenberger H, Puente N, Rusakov DA, Duvoisin R, Wang L-Y, et al. Glutamate
transporter studies reveal the pruning of metabotropic glutamate receptors and absence of AMPA
receptor desensitization at mature calyx of Held synapses. J Neurosci. 2005;25(37):8482-97. doi:
10.1523/JNEUROSCI.1848-05.2005. PubMed PMID: 16162930.
102.   Taschenberger H, Leão RM, Rowland KC, Spirou GA, von Gersdorff H. Optimizing synaptic
architecture and efficiency for high-frequency transmission. Neuron. 2002;36(6):1127-43. doi:
10.1016/S0896-6273(02)01137-6. PubMed PMID: 12495627.
103.   Graydon CW, Cho S, Diamond JS, Kachar B, von Gersdorff H, Grimes WN. Specialized
postsynaptic morphology enhances neurotransmitter dilution and high-frequency signaling at an
auditory synapse. J Neurosci. 2014;34(24):8358-72. doi: 10.1523/JNEUROSCI.4493-13.2014. PubMed
PMID: 24920639; PubMed Central PMCID: PMCPMC4051984.
104.   Cathala L, Holderith NB, Nusser Z, DiGregorio DA, Cull-Candy SG. Changes in synaptic structure
underlie the developmental speeding of AMPA receptor-mediated EPSCs. Nat Neurosci.
2005;8(10):1310-8. doi: 10.1038/nn1534. PubMed PMID: 16172604.
105.   Wimmer VC, Horstmann H, Groh A, Kuner T. Donut-like topology of synaptic vesicles with a
central cluster of mitochondria wrapped into membrane protrusions: a novel structure-function module
of the adult calyx of Held. J Neurosci. 2006;26(1):109-16. doi: 10.1523/JNEUROSCI.3268-05.2006
PubMed PMID: 16399677.
106.   Yang YM, Wang W, Fedchyshyn MJ, Zhou Z, Ding J, Wang LY. Enhancing the fidelity of
neurotransmission by activity-dependent facilitation of presynaptic potassium currents. Nat Commun.
2014;5:4564. Epub 2014/08/01. doi: 10.1038/ncomms5564. PubMed PMID: 25078759.
107.   Borst JGG, Helmchen F. Calcium influx during an action potential. In: Conn PM, editor. Methods
in Enzymology. Ion Channels. 293: Academic Press; 1998. p. 352-71.

Appendix A. Phasors

For the circuit shown in Figure 2, we assume the presence of a voltage oscillator at the calyx. For

simplicity, we also assume that the admittances Y_1 and Y_4 of the resistive membrane components are

small compared to the admittances Y_2 and Y_5 of the capacitive components and can therefore both be

neglected. However, following the logic of the solutions below, it should be straightforward to derive the

solution for a situation that also includes either or both resistive components.

Question A1: How does the cleft potential depend upon the presynaptic membrane potential?

The admittances surrounding the cleft form a voltage divider. We assume that the impact of the prespike

at the postsynaptic cell is very small ($Y_5 \gg Y_6$ or Y_7), so we can neglect the contribution of Y_6 and Y_7:

930
$$\frac{V_{cleft}}{V_{pre}} = \frac{-jY_2}{Y_3 - j(Y_2 + Y_5)} = \frac{-jY_2}{Y_3 - j(Y_2 + Y_5)} \frac{Y_3 + j(Y_2 + Y_5)}{Y_3 + j(Y_2 + Y_5)}$$

931
$$= \frac{Y_2}{Y_3{}^2 + (Y_2 + Y_5)^2} \left((Y_2 + Y_5) - jY_3\right) \qquad (A1.1)$$

932 This notation can be converted to the modulus:

933
$$\left|\frac{V_{cleft}}{V_{pre}}\right| = \frac{Y_2}{Y_3{}^2 + (Y_2 + Y_5)^2} \sqrt{(Y_2 + Y_5)^2 + Y_3{}^2} = \frac{Y_2}{\sqrt{(Y_2 + Y_5)^2 + Y_3{}^2}} \qquad (A1.2)$$

934 And a phase shift:

935
$$\varphi_f = atan2(-Y_3, (Y_2 + Y_5)) \qquad (A1.3)$$

936 If we assume that the pre- and the postsynaptic capacitance facing the cleft are identical ($Y_2 = Y_5$):

937
$$\left|\frac{V_{cleft}}{V_{pre}}\right| = \frac{2\pi f c_{cl}}{\sqrt{(4\pi f c_{cl})^2 + g_{cl}{}^2}} \qquad (A1.4)$$

938
$$\varphi_f = atan2(-g_3, 4\pi f c_{cl}) \qquad (A1.5)$$

939   This can be simplified if $(4\pi f c_{cl})^2 \ll g_3^2$, which gives:

940
$$\left|\frac{V_{cleft}}{V_{pre}}\right| \leq \frac{2\pi f c_{cl}}{g_{cl}} = 2\pi f \tau_{cl} \qquad (A1.6)$$

941   The phase shift would approach $-\pi/2$, which reflects the first derivative of v_{pre}.

943   ***Question A2: how does the VC prespike depend on the passive synaptic parameters?***

944   The current that flows into the synaptic cleft from the terminal will either leak away via the cleft

945   conductance or enter into the postsynaptic cell. The current into the postsynaptic cell (i_{post}) can be

946   described by:

947
$$I_{post} = -jY_5 V_{cleft} = -jY_5 \frac{-jY_2}{Y_3 - j(Y_2 + Y_5)} V_{pre} \qquad (A2.1)$$

948
$$\frac{I_{post}}{V_{pre}} = \frac{-Y_2 Y_5}{Y_3 - j(Y_2 + Y_5)} = \frac{Y_2 Y_5}{Y_3^2 + (Y_2 + Y_5)^2}\left(-Y_3 - j(Y_2 + Y_5)\right) \qquad (A2.2)$$

949   We then get:

951
$$\left|\frac{I_{post}}{V_{pre}}\right| = Y_2 Y_5 \frac{\sqrt{Y_3^2 + (Y_2 + Y_5)^2}}{Y_3^2 + (Y_2 + Y_5)^2} = \frac{Y_2 Y_5}{\sqrt{Y_3^2 + (Y_2 + Y_5)^2}} \qquad (A2.3)$$

950   If we assume that the pre- and postsynaptic cleft capacitance are the same ($Y_2 = Y_5$):

952
$$\left|\frac{I_{post}}{V_{pre}}\right| = \frac{(2\pi f c_{cl})^2}{\sqrt{(4\pi f c_{cl})^2 + g_{cl}^2}} \qquad (A2.4)$$

953
$$\varphi_f = atan2(-(Y_2 + Y_5), -Y_3) = atan2(-4\pi f c_{cl}, -g_{cl}) \qquad (A2.5)$$

954   Again, we may approximate the modulus by simplifying the equation to:

955
$$\left|\frac{I_{post}}{V_{pre}}\right| \leq \frac{(2\pi f c_{cl})^2}{g_{cl}} = g_{cl}(2\pi f \tau_{cl})^2 \qquad (A2.6)$$

Question A3: what is the equation for the CC prespike?

The postsynaptic voltage response can be described by i_{post} divided by the combinations of the different

routes to the reference:

$$V_{post} = \frac{-I_{post}}{Y_{to\ the\ cleft} + Y_6 - jY_7} = \frac{-I_{post}}{(Y_3^{-1} + (Y_4 - jY_5)^{-1})^{-1} + Y_6 - jY_7} \qquad (A3.1)$$

Here, we already assumed that none of the currents will reenter the calyx. As the cleft leak conductance is

likely to be much higher than the cleft-facing admittance of the postsynaptic membrane ($Y_3^{-1} \ll$

$(Y_4 - jY_5)^{-1}$), we get:

$$V_{post} = \frac{-I_{post}}{Y_4 - jY_5 + Y_6 - jY_7}$$

$$= \frac{-jY_5 V_{cleft}}{Y_4 - jY_5 + Y_6 - jY_7}$$

$$= \frac{1}{Y_4 - jY_5 + Y_6 - jY_7} \frac{-Y_2 Y_5}{Y_3 - j(Y_2 + Y_5)} V_{pre} \qquad (A3.2)$$

$$\frac{V_{post}}{V_{pre}} = \frac{-Y_2 Y_5}{(Y_4 + Y_6 - j(Y_5 + Y_7))(Y_3 - j(Y_2 + Y_5))} \qquad (A3.3)$$

To simplify the derivation, we combine $Y_4 + Y_6$ to be the total postsynaptic conductance g_{post}, and we

combine $Y_5 + Y_7$ to be the total postsynaptic susceptance $2\pi f c_{post}$:

$$\frac{V_{post}}{V_{pre}} = \frac{-Y_2 Y_5}{(g_{post} - j2\pi f c_{post})(Y_3 - j(Y_2 + Y_5))}$$

$$= \frac{Y_2 Y_5}{\left(g_{post}^2 + (2\pi f c_{post})^2\right)(Y_3^2 + (Y_2 + Y_5)^2)} \left(-g_{post}Y_3 + 2\pi f c_{post}(Y_2 + Y_5)\right.$$

$$\left. - j\left(2\pi f c_{post}Y_3 + g_{post}(Y_2 + Y_5)\right)\right) \qquad (A3.4)$$

From this we can deduce the modulus:

$$\left|\frac{V_{post}}{V_{pre}}\right| = \frac{Y_2 Y_5 \sqrt{\left(2\pi f c_{post}(Y_2 + Y_5) - g_{post}Y_3\right)^2 + \left(2\pi f c_{post}Y_3 + g_{post}(Y_2 + Y_5)\right)^2}}{\left(g_{post}{}^2 + \left(2\pi f c_{post}\right)^2\right)(Y_3^2 + (Y_2 + Y_5)^2)}$$

$$= \frac{Y_2 Y_5}{\sqrt{\left(g_{post}{}^2 + \left(2\pi f c_{post}\right)^2\right)(Y_3^2 + (Y_2 + Y_5)^2)}} \quad \text{(A3.5)}$$

974   Assuming again that the susceptance towards the synaptic cleft ($Y_2 = Y_5$) is the same:

$$\left|\frac{V_{post}}{V_{pre}}\right| = \frac{(2\pi f c_{cl})^2}{\sqrt{\left(g_{post}{}^2 + \left(2\pi f c_{post}\right)^2\right)(g_{cl}^2 + (4\pi f c_{cl})^2)}} \quad \text{(A3.6)}$$

$$\varphi_f = atan2\left(-2\pi f c_{post}g_{cl} - 4\pi f c_{cl}g_{post} , -g_{post}g_{cl} + (2\pi f)^2 c_{post}c_{cl}\right) \quad \text{(A3.7)}$$

979   This can be further simplified by assuming that $g_{cl} \gg 4\pi f c_{cl}$ and $2\pi f c_{post} \gg g_{post}$:

$$\left|\frac{V_{post}}{V_{pre}}\right| = \frac{(2\pi f c_{cl})^2}{2\pi f c_{post}g_{cl}} = \frac{2\pi f c_{cl}^2}{c_{post}g_{cl}} = 2\pi f \tau_{cl} \frac{c_{cl}}{c_{post}} \quad \text{(A3.8)}$$

$$\varphi_f = atan2\left(-2\pi f c_{post}g_{cl} , -g_{post}g_{cl} + (2\pi f)^2 c_{post}c_{cl}\right)$$

$$= atan2\left(-2\pi f \tau_{post}, (2\pi f)^2 \tau_{cl}\tau_{post} - 1\right) \quad \text{(A3.9)}$$

984   ***Question A4: what is the impact of a current injected into the cleft on the VC prespike?***

985   This Appendix deals with the question how i_{post} relates to i_{ion}, which is relevant for how the calcium

986   current changes the prespike. The current i_{ion} that runs through the presynaptic cleft-facing membrane can

987   leak away through the cleft leak conductance or via the postsynaptic cell. How these currents distribute

988   depend on the relative admittances:

$$\frac{I_{post}}{I_{ion}} = \frac{-jY_5}{Y_3 + Y_{ion} - j(Y_2 + Y_5)} \quad \text{(A4.1)}$$

which becomes equation A1.1 when we assume that $g_{ion} \ll g_{cl}$ and $Y_2 = Y_5$.

We know that the calcium current during a presynaptic AP approaches a Gaussian function with a peak

amplitude of 1.0-2.0 nA, a standard deviation τ_{Ca} of 0.21 ms and a calcium charge q_{Ca} of ~1 pC [44, 107].

The Gaussian function has the advantage that its Fourier transform is exact, allowing us to derive an exact

solution for the impact of the calcium current on the postsynaptic prespike.

The Gaussian function to describe the calcium current over time is:

996
$$i_{ca} = \frac{q_{Ca}}{\tau_{Ca}\sqrt{2\pi}} e^{-\frac{t^2}{2\tau_{Ca}^2}} \quad or \quad i_{max} e^{-\frac{t^2}{2\tau_{Ca}^2}} \qquad (A4.2)$$

997 Its Fourier transform is:

999
$$I_{Ca} = \int_{-\infty}^{\infty} i_{ca}\, e^{j2\pi ft} dt = \int_{-\infty}^{\infty} \frac{q_{Ca}}{\tau_{Ca}\sqrt{2\pi}} e^{-\frac{t^2}{2\tau_{Ca}^2}} e^{j2\pi ft} dt = \frac{q_{Ca}}{\tau_{Ca}\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{\frac{-1}{2\tau_{Ca}^2}(t^2 - j4\pi\tau_{Ca}^2 ft)} dt$$

1000
$$= \frac{q_{Ca}}{\tau_{Ca}\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{\frac{-1}{2\tau_{Ca}^2}(t^2 - 2t(2\pi j\tau_{Ca}^2 f) + (2\pi j\tau_{Ca}^2 f)^2 - (2\pi j\tau_{Ca}^2 f)^2)} dt$$

1001
$$= \frac{q_{Ca}}{\tau_{Ca}\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{\frac{-1}{2\tau_{Ca}^2}(t - 2\pi j\tau_{Ca}^2 f)^2} dt\; e^{\frac{-1}{2\tau_{Ca}^2}(-2\pi j\tau_{Ca}^2 f)^2}$$

1002
$$= \frac{q_{Ca}}{\tau_{Ca}\sqrt{2\pi}}\; \tau_{Ca}\sqrt{2\pi}\; e^{-2(\pi\tau_{Ca}f)^2}$$

998
$$= q_{Ca}\, e^{-2(\pi\tau_{Ca}f)^2} \quad or \quad i_{max}\, \tau_{Ca}\sqrt{2\pi}\, e^{-2(\pi\tau_{Ca}f)^2} \qquad (A4.3)$$

1003 To obtain the calcium current component of the prespike, we use equation A1.1 and assume that $g_{cl} \gg$

1004 $4\pi f c_{cl}$:

1005
$$I_{post} = -j2\pi f \tau_{cl} I_{ca} = -j2\pi f \tau_{cl}\, q_{Ca}\, e^{-2(\pi\tau_{Ca}f)^2} \qquad (A4.4)$$

1006 We perform the inverse Fourier transform by using that the Fourier transform of di_{Ca}/dt is equal to $j2\pi f I_{Ca}$:

1007
$$i_{post} = \int_{-\infty}^{\infty} I_{post}\, e^{-j2\pi ft} df = \int_{-\infty}^{\infty} -j2\pi f \tau_{cl} I_{ca}\, e^{-j2\pi ft} df = -\tau_{cl} \int_{-\infty}^{\infty} j2\pi f I_{ca}\, e^{-j2\pi ft} df$$

1008
$$= -\tau_{cl} \frac{di_{ca}}{dt} \qquad (A4.5)$$

1009
$$-\tau_{cl}\frac{di_{ca}}{dt} = -\frac{\tau_{cl}q_{Ca}}{\tau_{Ca}{}^3\sqrt{2\pi}}\, t\, e^{-\frac{t^2}{2\tau_{Ca}{}^2}} \quad or \quad -\frac{\tau_{cl}i_{max}}{\tau_{Ca}{}^2}\, t\, e^{-\frac{t^2}{2\tau_{Ca}{}^2}} \qquad \text{(A4.6)}$$

1010 This function will show two peaks at $t = \pm\tau$ with an amplitude of:

1011
$$i_{post}(\pm\tau) = \pm\frac{\tau_{cl}}{\tau_{Ca}{}^2\sqrt{2\pi e}}\, q_{Ca} = \pm\frac{\tau_{cl}}{\tau_{Ca}\sqrt{e}}\, i_{Ca,max} \qquad \text{(A4.7)}$$

1012 We thus define a simple scaling factor $\tau_{cl}(\tau_{Ca}\sqrt{e})^{-1}$ that relates amplitude of the presynaptic calcium current

1013 to the amplitude of the calcium prespike.

1014

Appendix B. Geometry

Question B1: What is the voltage profile in the synaptic cleft for a simplified synapse?

We assume a radially-symmetrical synapse with radius r and cleft height h. The capacitive current that

enters the synaptic cleft equals:

$i_{cap} = c_m v'_{pre} = C_m \pi r^2 v'_{pre}$ (B1.1)

with v'_{pre} indicating the first temporal derivative of the presynaptic AP. We assume that the current is

homogeneously distributed over the cleft-facing membrane and that all currents within the cleft run

radially towards the interstitial fluid. The current density $k(x)$ (typically denoted J, but we use k instead to

avoid confusion with our imaginary j) can be defined as a quasi-static function of the radial location x,

where 0 indicates the center and r the synapse edge. $k(x)$ will be equal to the current that runs through the

cleft-facing membrane from the center to radius x divided by the surface through which it escapes, $A_{esc}(x)$.

This surface is the surface of an open cylinder with radius x and height h:

1027
$$k(x) = \frac{i(x)}{A_{esc}(x)} = \frac{\pi x^2 C_m v'_{pre}}{2\pi x h} = \frac{C_m v'_{pre}}{2h}x \qquad \text{(B1.2)}$$

1028   Based on $k(x)$, the electric field can be defined using a homogeneous extracellular cleft resistivity R_{ex},

1029   where the negative sign reflects the convention that current runs down the voltage gradient:

$$1030 \qquad -\frac{dv(x)}{dx} = R_{ex}k(x) = \frac{R_{ex}C_m v'_{pre}}{2h}x \qquad \text{(B1.3)}$$

1031   The voltage profile relative to the cleft edge is the integral of equation B1.3 from x to r:

$$1032 \qquad v(x) = \int_x^r \left(-\frac{R_{ex}C_m v'_{pre}}{2h}x\right)dx = -\frac{R_{ex}C_m v'_{pre}}{2h}\int_x^r (x)dx = \frac{R_{ex}C_m v'_{pre}}{4h}(r^2 - x^2) \qquad \text{(B1.4)}$$

1033   We can also rewrite equation B1.4 as a relative distance to the cleft edge:

$$1034 \qquad v(x/r) = \frac{r^2 R_{ex}C_m v'_{pre}}{4h}(1 - (x/r)^2) \qquad \text{(B1.5)}$$

The voltage profile will apply as long as all current runs radially to the extracellular matrix. When this is

not the case -for instance, when currents enter into the postsynaptic cell or return through the presynaptic

cleft-facing membrane- the current density will be reduced, and the voltage gradient will be reduced as

well.

Question B2: How much current will be generated at the postsynaptic membrane?

We assume that the currents that run through the postsynaptic membrane are purely capacitive:

$$1041 \qquad i_{post} = \int_{x=0}^r i_{cap}(x)dx = \int_{x=0}^r (C_m A(x)v'(x))dx \qquad \text{(B2.1)}$$

1042   Here, the current through an annulus of membrane (at x, with thickness dx) will depend on the radial

1043   location as the surface of the annulus equals $2\pi x dx$ and the cleft potential is also defined by the radial

1044   location (equation B1.4). Only v'_{pre} will be affected by the time derivative, which will become the second

1045   time derivative (v''_{pre}) . The equation then becomes:

$$1046 \qquad i_{post} = C_m \int_{x=0}^r \left(2\pi x \frac{R_{ex}C_m v''_{pre}}{4h}(r^2 - x^2)\right)dx$$

1047
$$= \frac{\pi R_{ex} C_m^2 v''_{pre}}{2h} \int_{x=0}^{r} (x\,(r^2 - x^2))dx$$

1048
$$= \frac{\pi R_{ex} C_m^2 v''_{pre}}{2h} \left[\frac{1}{2}r^2 x^2 - \frac{1}{4}x^4\right]_{x=0}^{r}$$

1049
$$= \frac{\pi R_{ex} C_m^2 v''_{pre}}{8h} r^4 \qquad \text{(B2.2)}$$

1050

1051 *Question B3: How does the cleft resistivity relate to the cleft conductance?*

1052 With our phasor analysis we defined that (A2.6 after Fourier transformation):

1053
$$i_{post} = \frac{(c_{cl})^2}{g_{cl}} v''_{pre} \qquad \text{(B3.1)}$$

1054 By combining equation B2.2 and B3.1 we find that g_{cl} relates to the cleft resistivity:

1055
$$g_{cl} = \frac{8\pi h}{R_{ex}} \qquad \text{(B3.2)}$$

1056 Combining equations B3.1 and B3.2 yields an alternative way to calculate the extracellular resistivity:

1057
$$R_{ex} = \frac{i_{post}}{c_{cl}^2} \frac{8\pi h}{v''_{pre}} \qquad \text{(B3.3)}$$

1058

Question B4. What are the relations for a finger-like synapse?

We assume a sheet-like synapse with a cleft height h, a constant width of $2r$, and a length L that is closed

at the longitudinal ends. Any current that enters the synaptic cleft cannot escape along the longitudinal

axis, leaving only the transversal route to the interstitial space. Examples of such synapses are finger-like

synapses, such as the neuromuscular junction or the endbulb of Held synapse for which we ignore the

longitudinal endings, or deeply-invaginating, cylindrical synapses (spinules) with a circular longitudinal

axis (*i.e.* a rolled-up sheet forming a cylinder). Because of the constant width, the voltage profile will be

identical along the longitudinal axis, varying only along the synapse width. The capacitive current will be

equal to $i_{cap}(x) = C_m v'_{pre} dr dl$ (B4.1).

The current density k at each location will be:

1069
$$k(x) = \frac{i(x)}{A_{esc}(x)} = \frac{C_m v'_{pre} x dl}{h dl} = \frac{C_m v'_{pre}}{h} x \qquad (B4.2)$$

1070   The voltage gradient then is:

1071
$$-\frac{dv(x)}{dx} = R_{ex} k(x) = \frac{R_{ex} C_m v'_{pre}}{h} x \qquad (B4.3)$$

1072   And the voltage at x is:

1073
$$v(x) = -\frac{R_{ex} C_m v'_{pre}}{h} \int_x^r (x) dx = \frac{R_{ex} C_m v'_{pre}}{2h} (r^2 - x^2) \qquad (B4.4)$$

1074   The changes in the cleft potential will generate a postsynaptic, capacitive current. To calculate the total

1075   current from the finger-like synapse, we sum the current along the two dimensions:

1076
$$i_{fin} = \int_{l=0}^{L} 2 \int_{x=0}^{r} i_{cap}(x) dx \, dl = \int_{l=0}^{L} 2 \int_{x=0}^{r} C_m A v'(x) dx \, dl$$

1077
$$= \int_{l=0}^{L} 2 \int_{x=0}^{r} C_m \frac{R_{ex} C_m v''_{pre}}{2h} (r^2 - x^2) dx \, dl = \frac{R_{ex} C_m^2 v''_{pre}}{h} \int_{l=0}^{L} \int_{x=0}^{r} (r^2 - x^2) dx \, dl$$

1078
$$= \frac{R_{ex} C_m^2 v''_{pre}}{h} \int_{l=0}^{L} \left[x r^2 - \frac{1}{3} x^3 \right]_{x=0}^{r} dl = \frac{2 R_{ex} C_m^2 v''_{pre} r^3}{3h} \int_{l=0}^{L} dl$$

1079
$$= \frac{2 R_{ex} C_m^2 v''_{pre} r^3 L}{3h} \qquad (B4.5)$$

For the sheet-like synapse, x=0 indicates the cleft center and currents can run in two, transversal

directions. We therefore multiply the capacitive current by 2 at the start of the derivation of B4.5. For a

spinule synapse, no current can escape from the 'deep' end of the cylinder (at x=0) and therefore the

currents run in a single, transversal direction. The right-hand side of B4.5 should therefore be halved for

spinule-like synapses or other deeply-invaginating synapses, neglecting any contribution from the cup-

shaped cleft at the deep end. The characteristic difference between the radial-like or sheet-like synapse is

whether the escape area along the longitudinal axis increases or stays constant, respectively. Following the

logic outlined in B1, B2 and B4, one can calculate the cleft potentials and postsynaptic currents for other

synaptic geometries.

1089

1090   **Question B5. How does fenestration change the cleft potential and the VC prespike?**

1091   To assess the impact of fenestration on the cleft potential and the VC prespike, we compare a radially-

1092   symmetric synapse and a synapse with n equally-sized fingers. The relative cleft potential can be

1093   calculated using equation B1.5 for the radially-symmetric synapse (v_{sym}) and equation B4.4 for the finger-

1094   like synapse (v_{fin}), while assuming that the cleft potential of each finger is independent from each other:

1095
$$\frac{v_{fin}}{v_{sym}} = \frac{r_{fin}^2 R_{ex} C_m v'_{pre}(1 - x^2/r_{fin}^2)}{2h} \frac{4h}{r_{sym}^2 R_{ex} C_m v'_{pre}(1 - x^2/r_{sym}^2)}$$

1096
$$= \frac{2r_{fin}^2(1 - x^2/r_{fin}^2)}{r_{sym}^2(1 - x^2/r_{sym}^2)} \qquad \text{(B5.1)}$$

1097   This gives at the center (x=0) a ratio of:

1098
$$\frac{v_{fin}}{v_{sym}} = 2(r_{fin}/r_{sym})^2 \qquad \text{(B5.2)}$$

1099   To compare the VC prespike of a radially-symmetric synapse and a fenestrated synapse, we assume that

1100   the currents from each finger are synchronized and that they sum linearly. We use equation B2.2 and

1101   equation B4.5:

1102
$$\frac{n\,i_{fin}}{i_{sym}} = \frac{2nR_{ex}C_m^2 v''_{pre}r_{fin}^3 L}{3h} \frac{8h}{\pi R_{ex}C_m^2 v''_{pre}r_{sym}^4} = \frac{16nLr_{fin}^3}{3\pi r_{sym}^4} \qquad \text{(B5.3)}$$

1103   We assume that the total contact surface is conserved, so the surface of the fenestrated synapse (n $2r_{fin}$ L)

1104   and the radially-symmetric synapse ($\pi r_{sym}{}^2$) should be equal. Therefore L = $\pi r_{sym}{}^2$(n $2r_{fin}$)$^{-1}$, which we insert

1105   into equation B5.3:

$$\frac{n\, i_{fin}}{i_{sym}} = \frac{16nr_{fin}^3\ \pi r_{sym}^2}{3\pi r_{sym}^4\ n\ 2r_{fin}} = \frac{8}{3}\left(r_{fin}/r_{sym}\right)^2 \qquad (B5.4)$$

1107