Transcriptional landscape of human microglia reveals robust gene expression signatures that implicates age, sex and *APOE*-related immunometabolic pathway perturbations

Tulsi Patel Ph.D[1], Troy P. Carnwath[1], Xue Wang Ph.D[2], Mariet Allen Ph.D[1], Sarah J. Lincoln[1], Laura J. Lewis-Tuffin Ph.D[3], Zachary S. Quicksall[2], Shu Lin[1], Frederick Q. Tutor-New[1], Charlotte C.G. Ho[1], Yuhao Min[1], Kimberly G. Malphrus[1], Thuy T. Nguyen[1], Elizabeth Martin[5], Cesar A. Garcia[5], Rawan M. Alkharboosh[5,6,7], Sanjeet Grewal M.D[5], Kaisorn Chaichana M.D[5], Robert Wharen M.D[5], Hugo Guerrero-Cazares M.D Ph.D[5], Alfredo Quinones-Hinojosa M.D[5], Nilüfer Ertekin-Taner M.D Ph.D[1,4,#]

Author Affiliations:

1) Department of Neuroscience, Mayo Clinic, Jacksonville, FL 32224 USA

2) Department of Quantitative Health Sciences, Mayo Clinic, Jacksonville, FL 32224 USA

3) Department of Cancer, Mayo Clinic, Jacksonville, FL 32224 USA

4)   Department of Neurology, Mayo Clinic, Jacksonville, FL 32224 USA

5)   Department of Neurosurgery, Mayo Clinic, Jacksonville, FL 32224 USA

6)   Neuroscience Graduate Program, Mayo Clinic Graduate School of Biomedical Sciences,

Mayo Clinic, Rochester, MN 55905, USA.

7)   Regenerative Sciences Training Program, Center for Regenerative Medicine, Mayo Clinic,

Rochester, MN 55905, USA

Corresponding Author

Corresponding Author Contact Information:

Departments of Neurology and Neuroscience, Mayo Clinic, 4500 San Pablo Road, Birdsall 3,

Jacksonville, FL 32224. Email: taner.nilufer@mayo.edu, Phone: 904-953-7103, Fax: 904-953-

7353.

Number of Figures: 4

Number of Supplementary Tables: 20

Number of Supplementary Figures: 10

Word Count: 5,609 (excludes Methods word count of 2,183)

References: 91

Abbreviations:

AD: Alzheimer's disease

APOE: Apolipoprotein E

BM: Brodmann's area

BSA: Bovine serum albumin

CERAD: Consortium to Establish a Registry for Alzheimer's Disease

CNS: Central nervous system

CQN: Conditional quantile normalization

DAM: Disease-associated microglia

DPBS: Dulbecco's phosphate buffered saline

FACS: Fluorescence-activated cell sorting

FPKM: Fragments per kilobase of transcript per million mapped reads

GEM: Gel bead-in emulsion

GO: Gene ontology

MACS: Magnetic-activated cell sorting

ME: Module eigengenes

MM: Module membership

PBS: Phosphate buffered saline

PC: Principal component

PCA: Principal component analysis

PFA: Paraformaldehyde

QC: Quality control

RNAseq: RNA sequencing

ROSMAP: Rush University Religious Order Study-Memory and Aging Project

scRNAseq: Single cell RNA sequencing

snRNAseq: Single nuclei RNA sequencing

UMI: Unique molecular identifier

WGCNA: Weighted gene co-expression network analysis

Abstract:

Microglia have fundamental roles in health and disease, however effects of age, sex and genetic factors on human microglia have not been fully explored. We applied bulk and single cell approaches to comprehensively characterize human microglia transcriptomes and their associations with age, sex and *APOE*. We identified a novel microglial signature, characterized its expression in bulk tissue and single cell microglia transcriptomes. We discovered microglial co-expression network modules associated with age, sex and *APOE*-ε4 that are enriched for lipid and carbohydrate metabolism genes. Integrated analyses of modules with single cell transcriptomes revealed significant overlap between age-associated module genes and both pro-inflammatory and disease-associated microglial clusters. These modules and clusters harbor known neurodegenerative disease genes including *APOE, PLCG2* and *BIN1*. Meta-analyses with published bulk and single cell microglial datasets further supported our findings. Thus, these data represent a well-characterized human microglial transcriptome resource; and highlight age, sex and *APOE*-related microglial immunometabolism perturbations with potential relevance in neurodegeneration.

Introduction

Microglia are the resident macrophages of the central nervous system (CNS), responsible for clearance of cellular debris and pathological protein aggregates. In the healthy brain they exist in a resting state and can be induced to a reactive state in response to changes in the CNS microenvironment, such as inflammation and neuronal damage[1]. They are fundamental to maintaining brain homeostasis during development, aging and disease, therefore microglial dysfunction could ultimately lead to neurodegeneration[2]. Microglia are integral to the pathophysiology of neurodegenerative diseases, including Alzheimer's disease (AD) and multiple sclerosis, with chronic inflammation implicated as a contributing factor[3-5].

Fresh human brain tissue studies are imperative to the characterization of the microglial transcriptome in health and disease; however, accessibility is limited. Although single nuclei studies using frozen tissue provide an easier alternative, recent studies have demonstrated limitations in detecting substantial populations of less abundant cell types[6,7]. Additionally, it was recently reported that many microglial activation genes are expressed in the cytosol and therefore are likely to be missed by single nuclei RNA sequencing (snRNAseq)[8]. Recent single cell studies aiming to characterize microglial gene expression using fresh tissue have highlighted the heterogeneity in microglial phenotypes[9-11]. This has revealed that phenotypic changes are not binary but rather a spectrum of states in which microglia can simultaneously co-exist during transition from resting to more reactive states. Additionally, these different subsets could have specialized functions in brain homeostasis and dysfunction. Thus, it is increasingly important to characterize these heterogeneous subpopulations to understand their roles in health and disease. This could also help facilitate the design of novel therapeutic approaches to target specific subpopulations of cells and modulate their activity[2].

Utilizing fresh human tissue from neurosurgeries allows us to study the mechanisms of microglial function in living cells. By necessity, this tissue is typically obtained from surgical resection of tumor or epilepsy-affected regions. While there are studies which investigated the effect of glioblastoma (GBM)[12] or temporal lobe epilepsy[13,14] on myeloid cells within or

surrounding the disease tissue, studies which leverage 'normal' tissue from these resections

to explore microglial transcriptome are rare[9,11,15]. Despite careful surgical resection to obtain

'normal' tissue, there may still be microglial transcriptional changes that occur as a result of

nearby disease tissue. Combined analyses of data from multiple available studies can

enhance sample size, rigor and reveal common microglial transcriptional signatures robust to

differences in tissue source or technique. However, such joint analyses are lacking from the

literature. In this study we leveraged the microglial expression data from our neurosurgical

samples and that made available by others[9,16-18] and discovered robust transcriptional

changes associated with age, sex and *APOE*.

Microglial expression has been shown to be affected by aging[17,18], however few studies have

investigated the effects of sex and genetic factors on human microglia. Sex differences in

microglia have been previously reported in mice, with females being predisposed to harboring

more activated microglia than males[19-21]. *APOE*, a lipoprotein of which the ε4 allele (*APOE*-

ε4) is a major risk factor for AD and also implicated in other neurodegenerative diseases[22], is

upregulated in disease-associated microglia (DAM) in mice and humans, but downregulated

in astrocyte and oligodendrocyte subpopulations[4,6,23,24]. In microglia and neurons, *APOE*

interacts with LDL receptors to facilitate endocytosis of cholesterol and phospholipids and

modulate lipid homeostasis in the brain[25]. Such studies provide growing support for cell type-

specific functions of *APOE*, however, its effects on microglia remain to be fully elucidated.

Thereby identifying age, sex and *APOE*-associated pathways in microglia will provide greater

insight into the functions of specific microglial subsets in relation to these risk factors. Inter-

individual variability and diversity in functional states makes targeting specific microglial

subsets in disease challenging for modulating these cells[2]. Identifying the mechanisms

regulating microglial homeostasis and activation can allow us to manipulate these cells for

therapeutic purposes.

In this study, we leveraged both bulk and single cell approaches to provide a comprehensive

characterization of the adult human microglial transcriptome. We obtained fresh intraoperative

neurosurgical brain tissue and isolated an enriched population of microglial cells to investigate

transcriptional changes associated with age, sex and *APOE*-ε4 in bulk microglia and further

explored these in single microglial cells. Our findings support age-, sex- and *APOE*-related

microglial transcriptome changes involving lipid and carbohydrate metabolic pathways and

implicate microglial immunometabolism perturbations relevant to neurodegenerative

diseases.

Results

To uncover microglial transcriptional profiles and their associations with age, sex and *APOE,* we performed microglial cell-type specific and single cell RNA sequencing (scRNAseq) studies in fresh human brain tissue. We obtained neurosurgical tissue unaffected by the primary disease process from 19 human donors (**Supplementary Figure S1**). Microglia were isolated by CD11b⁺ microbead selection followed by FACS sorting of cells expressing the CD11b⁺/CD45^{intermediate} microglial signature to acquire a more purified population. These samples underwent bulk microglia RNAseq, with subsets of these and additional samples also undergoing 10x scRNAseq (n=5) and bulk tissue RNAseq (n=9) (**Figure 1A**; **Supplementary Table S1**). Validation of sorted microglia using qPCR showed the expected *CD11b⁺*/*CD45*^{intermediate}/*P2RY12*⁺ microglial signature[2] with no expression of other cell type markers, indicating that we isolated a highly enriched microglial population (**Supplementary Figure S2**).

Identification of a core human microglial transcriptional signature

To define a core human microglial signature, we calculated $\log_2$ fold change and q-values of differential expression for each gene between bulk microglia RNAseq data in our study and bulk brain RNAseq data from 7 AMP-AD datasets provided by Mayo Clinic[26], Mount Sinai Brain Bank[27] and Rush University Religious Orders Study and Memory and Aging Project (ROS-MAP)[28] representing 6 brain regions from 515 human samples. Using a cutoff of 4-fold greater expression in our bulk microglia and a q-value threshold of 0.05, we identified 1,971 genes (**Supplementary Table S2-4**). These genes were expressed at significantly greater levels in our bulk microglial transcriptome data in comparison to each of the bulk brain transcriptome datasets. Therefore, we considered these 1,971 genes as the core microglial signature in our dataset. This signature comprises several known marker genes, with 12.7% of the genes being BRETIGEA[29] microglial genes, suggesting that it also likely harbors novel microglial markers of interest (**Supplementary Table S5**). GO enrichment using MSigDB showed that this signature was enriched for genes involved in immune-related and

inflammatory response pathways as would be expected, and leukocyte mediated immunity

(**Figure 1B**).

To determine the ability of bulk brain tissue data to capture microglial genes, we assessed the

expression levels of our microglial signature genes in each of the 7 AMP-AD bulk brain

RNAseq datasets. Of the 1,971 microglial signature genes in our study, 37-47% were captured

in these bulk brain datasets (**Supplementary Figure S3A-B**). Our microglial signature genes

comprised 3.6-4.5% of the expressed bulk brain transcriptome, consistent with prior

estimations[6,30]. We next compared bulk microglia RNAseq transcript levels to that obtained

from bulk tissue RNAseq of neurosurgical fresh brain tissue samples. Bulk fresh brain tissue

does not capture all microglial marker genes, as demonstrated by the low correlation between

bulk tissue and bulk microglia data (R=0.46) (**Supplementary Figure S3C, S4,**

Supplementary Table S6). This reiterates the need for complementary single cell type data

to deconvolute cell type specific expression. We provide the list of microglial signature genes

that are also expressed at high levels in bulk brain tissue data (**Supplementary Table S7**),

which can serve as a validated resource for microglial signature gene markers in bulk RNAseq

datasets.

To determine how the microglial signature in this study compared to previously published

signatures, we performed hypergeometric tests of overrepresentation with Galatro, et al. [17],

Gosselin, et al. [15] and Olah, et al. [18] studies which had signatures consisting of 881-1296

genes and also with Srinivasan et al.[16] with a smaller signature of 66 genes. Significant overlap

was observed across all datasets, except with Srinivasan[16] et al., with 350 genes common to

ours and the three larger signatures (**Figure 1C-D, Supplementary Table S4, S8**). This

comprised several established microglial marker genes, including *P2RY12*, *TMEM119* and

CX3CR1. The most significant overlap was shared with Gosselin, et al. [15] signature [OR=19.6

(17.0-Inf) p=3.8E-261], where 49.7% of their genes were also present in our signature, and

22% of ours in their signature. Gosselin, et al. [15] samples were also obtained from

neurosurgical tissue resections like our cohort; and are unlike Galatro, et al. [17] and Olah, et al.

[18] samples that were harvested during autopsy. We repeated these analyses focusing on the

subset of our samples with matching ages to the comparison groups, where applicable, and

still observed significant overlaps in signature (**Supplementary Table S9**), suggesting these

findings are robust. Although there appears to be a common set of microglial genes consistent

across signatures, each also harbors many unique genes, which could be due to study or

individual specific differences.

Transcriptional profiling of microglia discovers co-expression networks and implicates

lipid and carbohydrate metabolism pathways associated with age, sex and *APOE*

We generated gene co-expression networks using WGCNA[31] to reduce number of tests and

increase power to detect genetic associations with age, sex and *APOE*. We identified 7

modules with significant associations (**Figure 2; Supplementary Figure S5; Supplementary**

Table S10). Modules ME14 and ME34 associated with age, however, in opposite directions.

ME14 was enriched for genes involved in the lipid localization pathway that were upregulated

with age (R=0.50, p=0.03) (**Figure 2A-C)**. ME34, enriched for DNA endoreduplication genes,

had negative association with both age (R=-0.55, p=0.01) and *APOE*-ε4 (R=-0.50, p=0.03),

indicating that microglial transcripts involved in this pathway are downregulated with aging

and in *APOE*-ε4 carriers (**Figure 2A**). Several other modules also associated with *APOE*-ε4,

in either direction. The only module associated with sex was ME26, which was downregulated

in females (R=-0.54, p=0.02), and enriched for genes involved in cholesterol absorption and

lipid digestion. This module also had the most significant association with *APOE*, in the

positive direction with presence of *APOE*-ε4 (R=0.66, p=0.002) (**Figure 2A,B,E**). Of the *APOE*

associated modules, ME23 had the second most significant association (R=-0.61, p=0.006)

and was enriched for carbohydrate metabolism genes (**Figure 2A,B,D**). Given recent

discoveries in microglial immunometabolism[32-35], we focused on ME14, ME23 and ME26 that

are enriched for lipid and carbohydrate metabolism genes.

ME14 co-expression network (**Figure 2C**) hub genes *NPC2*, *MSR1* and *PLAU* are also

microglial signature genes in our study and known to be involved in microglial functions[36-40,41].

Several disease-associated microglial (DAM) markers are also present in this network, including *CD9*, *ARAP2* and *MYO1E*[4,42,43] that are increased with aging, implicating activated microglial lipid localization pathways in aging (**Figure 2F**). Several genes in this module were also previously linked to neurodegeneration, including *MYO1E*[44,45], *CTSL*[46] and *UNC5B*[47,48]. Due to the nature of the neurosurgical tissue, we also compared expression levels of the key module genes between tumor and epilepsy samples (**Supplementary Figure S6**). Out of 18 comparisons we only observed significant differences due to diagnosis in *NPC2, PLAU, APOC1* and *IKBKE*; therefore, it is important to note that some of these associations may be confounded due to the disease state.

Our microglial signature (**Supplementary Tables S2-S4**) had significant overrepresentation of the age-associated ME14 genes (**Supplementary Table S10**) (OR=1.55 [95% CI=1.23-INF], p=0.001), highlighting age-related increases in microglial signature genes. Galatro, et al. [17] and Olah, et al. [18] also reported age-related microglial signatures. Comparison of ME14 genes revealed significant overlap with Olah, et al. [18] (OR=1.34 [95% CI=1.05-INF] p=0.03), but not with Galatro, et al. [17] microglial aging signature genes (OR=1.09 [95% CI=0.81-INF] p=0.33).

ME26 cholesterol metabolism pathway genes exhibited reduced expression in males and were elevated in *APOE*-ε4 carriers (**Figure 2A,B**). This module contains known microglial genes *LDLR*, *CD36* and *CRIP1* (**Figure 2E,F**). Assessment of individual ME26 network genes revealed *C17orf49*, *RP11-589P10.7* and *MIR497HG* to be the only microglial signature genes in this network to be associated with both sex and *APOE* (**Figure 2E**). Other microglial signature genes in ME26 associated with only sex or only *APOE*, suggesting that these traits may have independent effects on expression of some microglial genes. Several *APOE*-associated genes in ME26 were previously implicated in AD, including *CASP7*[49,50] and *LDLR*[51,52] (**Figure 2F**).

248   Carbohydrate metabolism gene enriched module ME23 is downregulated in *APOE*-ε4 carriers

249   (**Figure 2A,B,D**). AD risk genes *BIN1*[53] and *PLCG2*[54] are present in this network, which have

250   both been implicated in microglial dysfunction in neurodegeneration (**Figure 2D**).

Single cell transcriptome reveals specific subtypes of microglia

To uncover distinct microglial subtypes, a subset of sorted microglial samples from

neurosurgical brain tissue underwent single cell expression profiling. We obtained 26,558 cells

from 5 unique individuals, including one individual who underwent epilepsy surgery and had

samples from two brain regions (**Supplementary Table S1**). Analysis of the scRNAseq data

from these samples revealed 13 distinct cell clusters which were annotated using established

neuronal and glial marker genes from the literature[4,6,9-11,43,55,56] (**Figure 3A, Supplementary**

Table S11). Myeloid markers (*AIF1*, *PTPRC*, *C1QA*) were detected in all clusters except

cluster 12 which expressed oligodendrocyte markers (*PLP1*, *MBP*, *MOBP*). Cluster 9

expressed macrophage-specific markers (*VCAN*, *FCN1*, *CRIP1*, *S100A8*). These two clusters

comprised <3% of all cells, indicating that our sorted samples represent a very pure microglial

population. Each myeloid cluster had cellular contributions from all samples, albeit with some

variability in their proportions, likely due to intrinsic differences between individuals (**Figure**

3B, Supplementary Table S12). For these samples, the most marked difference was

observed for macrophages (cluster 9) and homeostatic microglia (cluster 2), which had greater

contributions from the mesiotemporal and anterior temporal regions, respectively. This could

be due to the proximity of the mesiotemporal sample to the disease-affected region.

We characterized the microglial clusters by their expression of established microglial subtype

markers (**Figure 3C, Supplementary Figure S7**) and their most significant marker genes

(**Supplementary Figure S8**). Homeostatic (*TMEM119*, *P2RY12*, *CX3CR1*)[10,11,43,56], pro-

inflammatory (*CCL2*, *CCL4*, *EGR2*)[10,11] and DAM markers (*APOE*, *C1QA*, *C1QB*)[4,9,11,23] were

observed in clusters 2, 1/6 and 10, respectively. Cluster marker genes are defined as those

expressed in at least 70% of the cells in the cluster with log fold change > 0.6 and q < 0.05 in

comparison to all other clusters. Expression levels of the top marker genes per cluster are

shown (**Figure 3C; Supplementary Figure S8; Supplementary Table S13**). Most of these

markers are distinct to a single cluster, although some clusters appeared to have similarities

in their marker expressions. To define the proximity of their transcriptional profiles, we

performed hierarchical clustering of the microglial clusters (**Figure 3D**). We determined that

the homeostatic microglia cluster 2 was transcriptionally closest to clusters 7 and 11, which

may represent subtypes of homeostatic microglia. Clusters 1 and 6 both expressed

chemokines *CCL2* and *CCL4* representative of pro-inflammatory microglia, however cluster 6

was more closely related to DAM, whereas cluster 1 represented a more distinct microglial

signature. Cluster 6 highly expressed interferon-related marker *IFITM3* and *ISG15*, also

observed in a cluster by Olah et al (2020)[9], which they defined as an interferon response-

enriched subset. These findings highlight different transcriptional profiles for the two pro-

inflammatory microglial clusters that may represent distinct activated microglia subtypes.

Cluster 3 highly expressed heat shock protein *HSPA1A*, an immediate early gene[57] reportedly

involved in antigen processing[58], response to stress and injury and exhibiting decreased gene

expression in multiple sclerosis patients[59,60]. Several were upregulated in this cluster,

suggesting that this may represent cells that underwent dissociation-induced stress[11]. Several

of the clusters did not express well known existing cell type markers. Clusters 5/8 and FOS[+]

0/4 were transcriptionally closest to one another (**Figure 3D**). Cluster 5 has distinct expression

of immunoreactive marker *CD163*, which was not observed in other subsets except

macrophages. Several *HLA* genes are also highly expressed in this cluster, suggesting that

these may be immunoreactive microglia[61]. Cluster 8 marker FTL has recently been used to

characterize iron-accumulating microglia[62]. Our findings highlight transcriptional profiles for

known microglial clusters, describe the transcriptional proximity of these clusters and suggest

that less well-defined clusters could potentially represent novel or intermediate transcriptional

states of microglia. Our microglial signature was significantly enriched in more clusters

expressing more activated markers (**Supplementary Table S14**), implicating this as the

dominant expression profile within our samples. However, there is also enrichment of

homeostatic cluster 2, demonstrating that we have not only captured activated cells as might

be expected due to the nature of the tissue, but also homeostatic microglia subtypes.

To determine whether the bulk microglial co-expression networks (**Figure 2A,C-E,**

Supplementary Figure S5) were representative of microglial subtypes, we performed

enrichment analyses of the module genes within the myeloid clusters with sufficient cell

numbers (**Figure 3E; full enrichment statistics provided in Supplementary Table S15**).

Age-associated co-expression network ME14, implicated in lipid metabolism, was significantly

enriched in pro-inflammatory (cluster 6) and DAM (cluster 10) clusters. Genes within module

28, which was significantly upregulated with *APOE*-ε4, had statistically significant enrichment

in all clusters except cluster 7. There was no statistically significant enrichment for any of the

other microglial modules that had significant age, sex or *APOE* associations, suggesting that

these factors may have ubiquitous effects on most microglial subtypes. Some of the remaining

microglial   co-expression   networks   had   distinct   patterns   of   cluster   enrichment

(**Supplementary Figure S9**), suggesting that some but not all networks could be

representative of distinct microglial subtypes.

Meta-analyses with published datasets support age, sex and *APOE* associations

We identified three bulk microglial[16-18] and one single cell microglial[9] transcriptome study with

available data that could be analyzed with our own. Each bulk microglia transcriptome dataset

was individually processed through the same MAPR-seq pipeline[63] for quality control to

minimize variability due to data processing. Meta-analysis of WGCNA results was performed

by coercing the external dataset co-expression networks onto our existing co-expression

networks. The forest plot in **Figure 4A** highlights the individual and combined associations for

networks of interest across the datasets where age, sex or *APOE* information was available

(**Supplementary Table S16**). For module ME14 associations with age, Galatro et al (2017)

exhibited a similar direction of effect (R=0.30, P=0.06), whereas Srinivasan et al (2020) did

not show any association (R=-0.003, P=0.99), likely due to a higher median age of individuals

in that study. Meta-analyses of these two datasets with ours showed association of ME14 with

age (R=0.25, P=0.05). ME26 showed lower levels in males in all but Olah et al[18]. When meta-

analyzed, sex-associated ME26 genes were significantly down-regulated in males (R=-0.30,

P=0.007), supporting our finding. This module was also significantly associated with *APOE* e4

carriers in the Srinivasan dataset (R=0.42, P=0.04), as in ours (R=0.57, P=0.01) with

significant meta-analysis association (R=0.49, P=0.001). Likewise, Srinivasan data was

inversely correlated with *APOE* for ME23 (R=-0.38, P=0.07), as is our data (R=-0.58,

P=0.009), with significant meta-analysis results (R=-0.0.47, P=0.0.002). The consistent

direction of association between most datasets and ours is also evident when hub genes from

the modules are plotted by the analyzed variable (**Supplementary Figure S10**). These

consistent results across multiple datasets provide support for our findings.

We subsequently integrated microglial single cell data from Olah et al (2020)[6] (n=9, 15,819

cells) with our samples (n=6, 26,558 cells) to compare these single cell datasets. There were

11 integrated-clusters when both datasets were analyzed together. The UMAP in **Figure 4B**

is split by study and shows relatively even contributions to each cluster from both datasets,

also observed in the stacked bar plot in **Figure 4C,** demonstrating slightly greater proportions

from our dataset commensurate with our greater cell numbers. One exception is the

integrated-cluster 10 that has a greater number of cells from Olah et al. This cluster has similar

expression patterns to our original FOS+ cluster 0. Hypergeometric tests of enrichment

showed significant overlap between many of the original and integrated clusters, showcasing

the high congruence between our data and Olah et al.[9] (**Supplementary Table S17**). To

further identify where our individual cells clustered when combined with the Olah dataset, we

overlaid the cell IDs from the original clusters to the integrated dataset and calculated the

overlap (**Supplementary Table S18**). Greatest cellular overlaps were observed between the

original clusters and integrated-clusters with the same subtype features. Many integrated-

clusters had the identities observed in the original cluster, as evidenced by the dot plot (**Figure**

4D) which shows some of the same top marker genes depicted in **Figure 3C**. We again

observed enrichment of our microglial signature in several integrated single cell clusters, many

of which highly expressed markers of activated microglia (**Supplementary Table S19**).

Finally, we compared the microglial subtype marker genes from a published single cell[11] and

single nucleus[6] study for many of the clusters from our single cell data (**Supplementary Table**

S20), with greater enrichment for like-subtypes from the former dataset and ours. Collectively,

these findings demonstrate the robustness of the microglial subtypes we identified and provide

detailed characterization of these in both our and integrated single cell datasets.

Discussion

Given their critical functions in maintaining homeostasis in the central nervous system (CNS) in health and their multifaceted roles during neurological diseases[2,3], understanding the biology of microglia and characterizing microglial subtypes is essential. Large scale studies in bulk brain tissue[26-28] have been instrumental in establishing transcriptional profiles in health and neurodegenerative diseases. Although these studies yielded information on brain expression signatures and uncovered perturbed pathways and molecules implicated in Alzheimer's disease and other neurological disorders[64-67], they are limited in their ability to provide cell-type specific transcriptional outcomes, especially for less abundant CNS cells such as microglia[30]. Analytic deconvolution approaches began to leverage these bulk tissue transcriptome datasets to estimate cell-type specific expression profiles[29,30], but the accuracy of these methods relies on the availability of high-quality single cell-type datasets. Such microglia-specific transcriptome datasets are gradually emerging[9,15,17,18], although the numbers of unique samples assessed remain limited given the arduous nature of collecting fresh human brain tissue. Additionally, comparative assessment of bulk brain vs. single cell-type bulk microglia vs. single-cell microglia studies are still rare[9,16,68]. To our knowledge there are no studies that evaluate human microglial transcriptome using all three approaches, as in our study. Further, investigations on effects of genetic and other factors on microglial transcriptional signatures in humans is likewise sparse, with the exception of age-related effects assessed in a few studies[15,17,18]. Finally, unlike in bulk tissue studies[29,64-67], microglia-specific co-expression networks, their molecular signatures and functional implications have not been evaluated.

In this study, we sought to overcome these knowledge gaps by characterizing the transcriptome of sorted bulk and single-cell microglial populations isolated from fresh human brain tissue. We identified a robust microglial signature comprising 1,971 genes enriched for immune-related functions. These signature genes were selected due to their consistently higher expression levels in our sorted bulk microglial transcriptome in comparison to 7 different

bulk brain tissue datasets from 6 different regions[26-28]. We also compared sorted bulk microglia

to bulk fresh brain tissue and identified transcripts that are expressed in both. The microglial

signature genes that are also reliably detected in bulk brain tissue represent a validated list of

microglial markers that can be utilized in bulk brain tissue transcriptome analytic deconvolution

studies[29,30].

Our microglial signature significantly overlapped with other signatures from bulk microglia

previously reported by Galatro, et al. [17], Gosselin, et al. [15] and Olah et al.[18], implicating a core

set of genes consistently expressed in this cell type. However, there were additional genes

unique to each signature, likely to be driven by factors such as patient demographics or study

differences. Galatro, et al. [17] and Olah et al.[18] both also reported age-related microglial

expression signatures. We found significant overlap of our age-associated microglial gene

expression module ME14 genes with the latter, which was also enriched for our microglial

signature. This indicates that bulk microglial profiles can effectively capture genes affected by

aging in microglia.

We leveraged the co-expression network structure of sorted bulk microglia to further explore

whether microglial subsets were associated with age, sex or *APOE*-ε4. To our knowledge sex-

differences in microglial transcriptome were previously studied only in mice[19-21], however

APOE genotype-specific microglial interactions with amyloid plaques have been previously

observed in mice[20,69] and humans[70]. We identified two network modules associated with age,

one with sex and six with *APOE*-ε4. We observed that two modules, ME14 that is positively

associated with increased age; and ME26 that is positively associated with both *APOE*-ε4 and

female sex, were both enriched for lipid metabolism biological terms[32-34]. Module ME14

included genes involved in lipid localization and storage pathways (*PLIN2, IL6, LPL, MSR1,*

ENPP1, PPARG, PTPN2, SOAT1, IKBKE) and ME26 had lipid digestion/cholesterol transport

pathway genes (*CD36, LDLR*). Both modules harbored known microglial genes (*LDLR, CD36,*

CRIP1, NPC2, *MSR1, PLAU)* and those that are included in our microglial signature (*PLIN2,*

IL6, MSR1, SOAT1, IKBKE, NPC2, PLAU).

Comparing the sorted bulk microglial network modules to scRNAseq microglial clusters, we

determined that ME14 genes were significantly over-represented in pro-inflammatory cluster

6 and disease-associated microglia (DAM) cluster 10. In our study, DAM cluster 10 included

APOE, APOC1, ASAH1 and *CTSD*. Of these *APOE*[22,33,71], *APOC1* and *ASAH1*[72] are involved

in lipid metabolism and neurodegenerative diseases. *APOE*[4,5,23], *APOC1*[23] and *CTSD*[4] were

also signature genes in mouse models of neurodegenerative diseases[4,5] or aging[23]. Our pro-

inflammatory cluster 6 also included genes associated with mice microglial neurodegenerative

(*FTH1*[4]) or aging signatures (*CCL4*[23]), as well as *IFITM3*[34], *GOLGA4*[34], previously shown to

be upregulated in aging lipid droplet accumulating microglia[34]. Our findings that integrate

human sorted bulk RNAseq and scRNAseq data, support a model where aging human

microglia transition to a pro-inflammatory and disease-associated transcriptional profile which

is also associated with perturbations in lipid metabolism in these cells.

There is increasing evidence that tightly controlled lipid metabolism is essential to the

functions of microglia during development and homeostatic functions of adulthood and may

be disrupted in aging and disease[32,33]. The complex interactions between microglial lipid

metabolism and its cellular functions rely on lipid sensing by microglial receptors such as CD36

and TREM2 and uptake of lipids, including LDL and APOE[32,33]. These interactions are

necessary for microglia to become activated and perform functions including phagocytosis of

myelin[73] and misfolded proteins like amyloid ß[74], cytokine release, migration and

proliferation[32,35]. Studies primarily focused on *in vitro* and animal models suggest disruption of

the microglial immunometabolism and assumption of a pro-inflammatory phenotype with

aging[23,34,75,76] and diseases including multiple sclerosis (MS) and Alzheimer's disease[4,5,77].

Interestingly, microglial lipid droplet accumulation has been demonstrated under all these

conditions[32-34,73] and lipid droplet accumulating microglia in aging mice were shown to have a

unique transcriptional state[34]. Our findings in sorted cells from fresh human brain tissue

provide transcriptional evidence for immunometabolism changes and pro-inflammatory

phenotype with microglial aging, thereby contributing essential complementary data from

humans for this cell type.

Besides module ME14, we determined that ME26 is also enriched for lipid metabolism genes.

ME26 module expression is higher in both *APOE*-ε4 and female sex, however we note that in

our sorted bulk microglia RNAseq samples, there were no male *APOE*-ε4 carriers. Therefore,

the distinct influence of sex and *APOE* on the expression of this module remains to be

established. *APOE*-ε4, a major risk factor for Alzheimer's disease, has the lowest lipid binding

efficiency compared with other *APOE* isoforms[32]. Increased cholesterol accumulation has

been reported in both iPSC-driven astrocytes from *APOE*-ε4 carriers[78] and in Apoe-deficient

microglia[73]. These findings collectively support a role for *APOE*-ε4 associated microglial

transcriptional changes and disrupted cholesterol metabolism. Using our sorted microglia

RNAseq data, we identified five additional modules that associate with *APOE*-ε4, one in a

positive direction (ME28) and four negatively (ME4, ME23, ME34, ME36). Of these, module

ME23 had the second most significant *APOE*-ε4 association after ME26. Interestingly, ME23

was enriched for carbohydrate metabolism biological processes, which are also tightly

regulated in microglia[35]. Module ME23 harbors known AD risk genes *BIN1* and *PLCG2*, where

the latter is a microglial gene that modulates signaling through *TREM2*[79] and also a hub gene

in this module. ME23 genes *BIN1, JUN* and *TGFBR2* were found to be reduced in a mouse

microglial neurodegenerative phenotype gene signature[5]. These findings further demonstrate

the consistency of our human microglial data with that from mouse models and supports

perturbed microglial immunometabolism as a potential pathogenic mechanism in

neurodegeneration. Importantly, using available transcriptome data from three other bulk

microglia datasets of Galatro et al (2017)[17], Olah et al (2018)[18] and Srinivasan et al (2020)[16],

we performed meta-analyses with our data thereby providing results from a collective sample

size of 93, 83 and 44 fresh tissue samples for respectively, sex, age and *APOE* associations.

We observed that the associations in our data replicated in most of the other available datasets

and the meta-analyses improved the significance of most associations. These results

represent the largest collective bulk microglial associations from human fresh brain tissue to

our knowledge and provide further support for our findings.

In addition to analyzing gene expression modules from sorted bulk microglia, we also identified

microglial clusters from sorted microglial scRNAseq data. To our knowledge, there are only

two prior publications of scRNAseq characterizations on human microglia[9,10]. Masuda et al.[10]

analyzed 1,602 microglia isolated from 5 control and 5 MS patient brains, compared their

findings to those from mice demonstrating clusters that are common and others that are

species-specific. Olah et al. assessed 16,242 microglia from 17 individuals and characterized

subclusters of microglia from patients with mild cognitive impairment, AD and epilepsy[9]. Our

scRNAseq dataset is from 5 unique individuals comprising 26,558 cells, 99.98% of which have

myeloid markers. We identified microglial clusters that share characteristics of those

previously reported in mice[4] and humans[9,70], such as DAM. We also uncovered clusters that

had not been previously characterized in the literature, including cluster 7, exhibiting high

expression levels of astrocytic marker gene *SLC1A3*. Microglial expression of *SLC1A3* was

previously shown to occur in mice and humans especially in disease states[80-82]. We also

leveraged these scRNAseq data to further characterize the sorted bulk microglial expression

modules. Integration of our samples with Olah et al.[9] allowed us to explore microglial subtypes

at single cell resolution with increased power. We found that even in this larger dataset, our

microglial signature was similarly enriched in the more activated subtype clusters,

demonstrating the robustness of this signature across datasets. The homogeneity in clustering

highlighted shared expression patterns between cells from both datasets, providing further

support for the putative microglial subtypes we identified in our current study. Hence our

microglial scRNAseq data contribute further to the emerging single cell landscape of this cell

type.

Activated microglia are a hallmark in both brain cancers and temporal lobe epilepsy in

response to the inflammation[12-14,83,84]. We acknowledge that the transcriptome data generated

in this study is sourced from grossly normal tissue obtained during resection of brain tumor or

epilepsy regions. Despite careful resection to exclude disease tissue, it is possible that the

grossly normal tissue transcriptome may still be influenced by nearby pathology. Therefore,

we cannot rule out the possibility that the activated microglial subtypes we identify may be due

to the nature of the tissue. Indeed, sex-specific expression differences have been observed in

glioma-activated microglia[84]. Darmanis et al[12] investigated the effect of glioblastoma tumors

on CNS cell types and surrounding tissue, and found that peri-tumor myeloid populations were

primarily pro-inflammatory microglia compared to macrophages within the tumor core[12]. In

temporal lobe epilepsy, two distinct microglial phenotypes have been identified with microglia

present in sclerotic areas with few neurons expressing markers of activation, including anti-

inflammatory cytokine IL10[13,14]. The other phenotype occurs transiently following a seizure,

with secretion of interleukins CXCL8 and IL1B mediated by the NLRP3 inflammasome[13].

Nonetheless, we did not observe significant differences in most genes we highlighted here

based on tumor vs. epilepsy diagnosis (**Supplementary Figure S6**). There is also a large

cluster of homeostatic microglia detected in our samples, indicating that not all cells are

activated. Further, the consistency of our findings with other single cell[9] or bulk microglia[16-18]

data supports the notion that the microglial signatures and associations we identified in our

study are applicable to microglia biology in general rather than being disease-specific.

We recognize that our study has several limitations, primarily owing to the difficulty in obtaining

high quality neurosurgical brain tissue, which leads to limited sample size and variability in

tissue, diagnoses and patient demographics. Even though we have utilized grossly normal

tissue surgically separated from disease tissue, the samples are from epilepsy and various

brain tumor patients representing multiple diagnoses. We isolated microglia using an

approach which should minimize activation however we cannot definitively rule out stress-

induced transcriptomic changes during isolation. Despite these caveats, we could identify

microglial co-expression modules and subclusters with multiple features that are consistent

with prior publications from model systems[4,5,23,34]. Our scRNAseq clusters have contributions

from both tumor and epilepsy samples, suggesting that our findings are unlikely to be driven

by any one diagnoses. Furthermore, there are few published studies using fresh brain tissue

to study microglia and thus our integrated single cell data showing a homogenous cluster of

microglia highlights the robustness of the methodology. Finally, our joint and meta-analysis

using all available datasets enhance the rigor of and support our conclusions.

In summary, our study on sorted bulk microglia RNAseq and scRNAseq from fresh brain tissue

yield several key findings. We identify a microglial gene signature from sorted bulk microglia,

characterize its expression in bulk brain RNAseq across 7 datasets comprising 6 regions, in

bulk fresh brain RNAseq and in microglial scRNAseq subtype clusters. This signature provides

a well-characterized resource which can be utilized in analytic deconvolution studies of bulk

transcriptome data[29,30]. We uncovered microglial gene expression modules associated with

age, sex and/or *APOE*-ε4. Modules with age and *APOE*-ε4 associated transcriptional changes

implicate microglial lipid and carbohydrate metabolism perturbations and microglial activation.

Microglial scRNAseq data highlight the transcriptional complexity of this cell type, reveal both

known and novel cell types and demonstrate utility of this data in characterizing sorted bulk

RNAseq   data.   These   findings   provide   support   for   the   emerging   microglial

immunometabolism[32,35] pathway as a plausible therapeutic target in aging-related disorders;

and provide a well-characterized human transcriptome resource for the research community

on this cell type with central roles in health and disease[1].

Methods

Patient Samples

Fresh human brain tissue was obtained from patients undergoing neurosurgical procedures for epilepsy or tumor resection. Tissues determined to be grossly unaffected by the primary disease process were utilized for the present study **(Supplementary Figure 1)**. Patient samples were transported from the operating room to the laboratory in 1X DPBS (Thermofisher; 14287080) for processing within 1-2 hours of resection. Human tissue was collected with informed consent prior to surgery and all procedures were approved by the Mayo Clinic Institutional Review Board and are HIPAA compliant.

Tissue Dissociation

Tissue was dissected to remove necrotic tissue, white matter and excess vascular tissue, to retain only cortical grey matter. The remaining tissue was cut into sagittal slices and weighed before being processed using the Adult Brain Dissociation Kit (Miltenyi; 130-107-677) as per the manufacturer's protocol. Debris removal (Miltenyi; 130-109-398) and red blood cell lysis (Miltenyi; 130-094-183) were also performed. All procedures were carried out on ice. The resulting homogenate was filtered through a 70μm filter before proceeding.

Magnetic-Activated Cell Sorting (MACS)

The cell suspension was first enriched for $CD11b^+$ cells by incubating with anti-CD11b microbeads (Miltenyi; 130-049-601 clone M1/70) for 15 minutes according to manufacturer's recommendation. This was then washed with PB buffer (0.5% BSA, 1X PBS Ca^{2+}/Mg^{2+} free pH 7.4) and filtered through a 70μm cell strainer before being applied to a large separation column (Miltenyi; 130-042-401) in a QuadroMACS separator magnet (Miltenyi; 130-090-976). The $CD11b^+$ fraction was collected and resuspended in sterile filtered FACS buffer (1X PBS Ca^{2+}/Mg^{2+} free, 0.5% BSA, 2% FBS, 3mM EDTA) for antibody staining.

Fluorescence-Activated Cell Sorting (FACS)

MACS sorted CD11b$^+$ cells subsequently underwent FACS sorting to further purify the

microglial population. The cell suspension was incubated in Human TruStain FcX blocking

solution (1:20, Biolegend; 422302) at room temperature for 10 minutes. Subsequently, cells

were stained with anti-CD11b PE/Cy7 (1:100, Biolegend; 101206, M1/70) and anti-CD45

Alexa Fluor 647 (1:100, Biolegend; 304056, HI30) antibodies for 30 minutes on ice. Following

two washes with FACS buffer, SYTOX Green viability dye (1:1000, ThermoFisher; S7020)

was added for an additional 20 minutes. Single cell suspensions were filtered through a 40μm

cell strainer (Falcon; 352235) before sorting on a BD FACS Aria II (BD Biosciences).

CD11b$^+$/CD45^{intermediate}/SYTOX green$^-$ cells were sorted directly into FACS buffer.

Independently CD11b and CD45 are not microglial specific markers, however gating cells with

a CD11b+/CD45^{intermediate} signature allowed us to differentiate microglia from macrophages

which are expected to be CD45^{high}. An example of our FACS gating strategy is provided in

Supplementary Figure S2A.

RNA Isolation and Sequencing

RNA from sorted microglial cells was isolated using the miRNeasy Serum/Plasma Kit

(QIAGEN; 217184) and quantified on the Agilent BioAnalyzer 2100. cDNA libraries were

generated using SMARTSeq2 v4 and Nextera Low Input Library Prep Kit. Samples were

multiplexed and sequenced on the Illumina HiSeq 4000.

RNA from frozen bulk tissue was isolated using Trizol and chloroform, followed by DNase and

clean up using the RNeasy Kit (QIAGEN; 74106). Libraries were generated using the TruSeq

Stranded mRNA Library Prep Kit. Samples were multiplexed and sequenced on the Illumina

HiSeq 4000. Base-calling of all sequence data was performed using Illumina's RTA v2.7.7.

10X Single Cell 3' v3 Library Preparation of Sorted Microglia

Viability of MACS plus FACS sorted cells was assessed by Trypan blue (Gibco; 15250061)

exclusion and cell density was determined using a hemocytometer prior to adjustment to target

4000-5000 cells. Cells were loaded onto a 10X Chromium chip and run on the GemCode

Single Cell Instrument (10X Genomics) to generate single cell gel beads-in-emulsion (GEMs).

Single cell RNA-seq libraries were prepared using the Chromium Single Cell 3' Gel Bead and

Library Kit v2 and v3 (10X Genomics; 120237) and the Chromium i7 Multiplex Kit (10X

Genomics; 120262) according to the manufacturer's instructions. Quality of cDNA libraries

was determined using a BioAnalyzer 2100 DNA High Sensitivity assay (Agilent; 5067-4626)

prior to sequencing one per lane on an Illumina HiSeq 4000.

Validation with Quantitative Real-Time PCR

Total RNA was extracted from sorted cells using the miRNeasy Serum/Plasma Kit (QIAGEN;

217184). Concentration and quality were assessed using the Agilent BioAnalyzer RNA 6000

Pico Kit (Agilent; 5067-1514). RNA was normalized to 0.5ng/µl for cDNA synthesis using the

SuperScript IV VILO Master Mix (ThermoFisher; 11756050). TaqMan PreAmp Master Mix

(ThermoFisher; 4391128) was used to pre-amplify the cDNA, followed by TaqMan Universal

PCR Master Mix (ThermoFisher; 4304437) with the following gene expression probes: *MOG,*

AQP4, THY1, PTPRC, ITGAM, P2RY12, PECAM1, CD34, GAPDH (ThermoFisher;

Hs01555268_m1, Hs00242342_m1, Hs00174816_m1, Hs04189704_m1, Hs00355885_m1,

Hs00224470_m1, Hs01065279_m1, Hs02576480_m1, Hs99999905_m1). RT-qPCR was

performed on a QuantStudio 7 Flex Real-Time PCR System (ThermoFisher) using a relative

standard curve to quantify gene expression.

Validation with Immunocytochemistry

Cultured cells were fixed with 4% paraformaldehyde (PFA) overnight at 4°C and blocked with

blocking solution (10% BSA, 5% normal goat serum and 0.1% Triton-X). Fixed cells were

stained with anti-TMEM119 (1:100, Biolegend; 853302) extracellular primary antibody with

Goat anti-mouse IgG secondary antibody conjugated to Alexa-488 (1:100, Abcam; ab150113).

Nuclei were stained with 1µg/ml DAPI (1:1000, ThermoFisher; 62248) before mounting with

AquaPoly Mount (Poly Sciences, 18606-20). Images were acquired with a Zeiss LSM880

Confocal microscope using a Plan-Apochromat 20x magnification and 0.8 objective at 1024

by 1024 pixels with a 0.5 microsecond pixel dwell time.

Data Analysis

Bulk Microglia RNA-seq Processing

The MAPR-Seq pipeline[63] was used to align reads to human reference genome hg38 using STAR[85] and count reads using featureCounts[86]. FastQC was used for quality control (QC) of raw sequence reads, and RSeQC was used for QC of mapped reads. Quality measures were examined including base calling quality, GC content, mapping statistics and sex check to ensure consistency between the recorded and inferred sex from expression of chromosome Y genes. Raw read counts were normalized using Conditional Quantile Normalization (CQN) to generate $\log_2$ scaled expression values via the Bioconductor package cqn, accounting for sequencing depth, gene length and GC content. Normalized CQN expression values were assessed using Principal components analysis (PCA) to identify and remove outliers, defined as greater than 4 standard deviations from the mean of the first two principal components. In addition, RPKM (reads per kilo bases per million) values were calculated.

Identification of a Core Microglial Signature from Bulk Microglia Data

To define a core microglial signature, we compared our bulk microglia data to cognitively normal control samples from the AMP-AD bulk tissue transcriptome data from 7 different datasets representing 6 brain regions (Synapse ID: syn2580853); Mayo Clinic[26] (cerebellum and superior temporal gyrus), Mount Sinai Brain Bank[27] BM10 (frontal pole), BM22 (superior temporal gyrus), BM36 (parahippocampal gyrus), BM44 (inferior frontal gyrus) and Rush University Religious Order Study-Memory and Aging Project (ROS-MAP)[28] (dorsolateral prefrontal cortex). Raw gene counts and metadata (see Acknowledgements) were obtained from the AMP-AD RNAseq Harmonization study which had performed alignment and processing of all datasets and brain regions through a consensus pipeline[87]. Samples were removed that had inconsistent sex between that indicated in metadata and that inferred from RNAseq expression; a RIN < 5; were identified as gene expression outliers based on principal component analysis (PCA) (> 4 standard deviation (SD) from mean PC1 or PC2), or missing metadata. In addition, duplicates (lowest read count sample removed) and those with rRNA

(>5%) were removed from the MSBB datasets. Furthermore, samples not meeting

neuropathological criteria as Alzheimer's disease (AD)[88] or control were excluded. To

generate the microglial expression signature, only control samples from the AMP-AD datasets

were included. Raw read counts were normalized using Conditional Quantile Normalization

(CQN). Log_2 fold change and q-values between each bulk tissue brain region and the bulk

microglia profiles were calculated for each gene via linear regression using log_2(RPKM)

without correction for covariates. Genes were filtered using a cutoff of 4-fold greater

expression in bulk microglia compared to each bulk tissue region and q < 0.05. Genes that

passed these criteria and were significant in comparisons with all 7 bulk brain datasets

determined the microglial signature. These signature genes were assessed for GO term

enrichment with biological pathways using MSigDB. REViGO[89] tree plots were generated in R

using GO terms obtained from MSigDB.

Weighted Gene Co-Expression Network Analysis

The CQN normalized expression values from bulk microglia were input to R WGCNA[31]

package v1.69. This analysis included 14,149 expressed genes, i.e. median(CQN) > 2.

Modules were identified, their eigengenes were calculated and merged if correlation of

eigengenes > 0.7. Genes in the 40 modules identified were tested for GO term enrichment via

WGCNA. Module membership (MM) for each gene was calculated as the correlation between

expression of each gene and its module eigengene. Genes with MM $\geq$ 0.7 are considered the

hub genes for the network. Gene co-expression network plots were generated in Cytoscape

v3.8 (http://www.cytoscape.org/). Each module eigengene was tested for association with age,

sex and *APOE* ε4 carrier status independently using Pearson correlation. Co-expression

network genes were annotated if they were significantly associated (p < 0.05) with the tested

trait.

Over-Representation and Correlation Analyses

Hypergeometric testing was performed in R to determine the enrichment of a select set of

genes in previously reported signatures, bulk tissue expressed genes, WGCNA modules or

10X single cell clusters. Correlation between bulk tissue and bulk microglial normalized CQN

data was calculated using Spearman's rank correlation. Concordant and discordantly

correlated genes were determined using the upper and lower quartiles from each dataset.

Single Cell Data Analysis

For single cell RNA samples, 10X Genomics Cell Ranger Single Cell Software Suite v3.1.0[90]

was used to demultiplex raw base call files generated from the sequencer into FASTQ files.

Raw reads were aligned to human genome build GRCh38. Reads aligned to gene transcript

locus were counted to generate raw UMI counts per gene per barcode for each sample. The

raw UMI matrices were filtered to only keep barcodes with > 500 UMIs and those that were

classified as cells by Cell Ranger's cell calling algorithm.

Quality control, normalization, clustering and marker gene identification were performed with

Seurat v3[91], followed by annotation of clusters using established cell type markers. We kept

1) barcodes with > 10% of UMI mapped to mitochondrial genome; 2) barcodes with < 400 or

> 8000 detected genes; 3) barcodes with < 500 or > 46,425 mapped UMIs; 4) genes that are

detected in < 5 cells. These thresholds were determined by UMI or gene distribution to identify

undetectable genes and outlier barcodes that may encode background, damaged or multiple

cells. UMI counts of remaining cells and genes were normalized using NormalizeData

function, which gave natural log transformed expression adjusted for total UMI counts in each

cell. The top 2000 genes whose normalized expression varied the most across cells were

identified through FindVariableFeatures function with default parameters. Using those genes,

cells from 6 samples were integrated using functions FindIntegrationAnchors and

IntegrateData with default parameters. Principal components (PCs) of the integrated and

scaled data were computed; and the first 31 PCs, which accounted for > 95% variance, were

used in clustering cells. Cell clustering was performed using FindNeighbors and FindClusters

with default parameters. Marker genes were identified in each cluster using FindMarkers in

Seurat. Marker genes on one cluster must 1) be present in > 20% cells in the cluster; 2) the

log(fold change) between expression in the cluster and other clusters must be > 0.25; 3) the

rank sum test p-value (Bonferroni-adjusted) between cells in the cluster and cells in other

clusters < 0.05.

Meta-Analysis of Bulk Microglia RNA-seq Datasets

For these analyses we identified and obtained RNAseq data from three bulk microglial

datasets that were available for download[16-18] (**Supplementary Table S16**). We obtained sets

of co-expression genes i.e. WGCNA modules from our own data for those modules the

eigengenes of which were significantly correlated with age/sex/*APOE*. To determine whether

these correlations replicated in the external datasets, we downloaded raw reads in FASTQ

format for the three external bulk microglia datasets listed in **Supplementary Table S16**.

Reads were mapped to human genome build hg38 and were counted and normalized in the

same fashion as described previously. For each WGCNA module, we identified the central

genes (i.e. genes whose Pearson correlation with module eigengene > 0.75). Using these

central genes, module eigengenes were calculated in all datasets – ours and three external

ones. We correlated module eigengenes with traits. Finally, we performed meta-analysis to

combine correlations from multiple datasets using metacor function (random effect model) in

R meta package.

Joint Analysis of Single Cell RNA-seq Datasets

Raw single cell RNA-seq data from Olah et al (2020)[9] was downloaded through Synapse

(syn21438358) and processed through Cell Ranger. Quality control was performed as

described prior to integration with our samples using Seurat v4.0.4. Integration was performed

to combine the datasets by individual rather than sample. A total of 9 samples and 15,819

cells were retained from the Olah data[9]. Cells were annotated to include dataset of origin.

Hypergeometric tests were performed to determine the enrichment of our microglial signature

within each integrated cluster. Additionally, we also looked at overlap between integrated

cluster genes and marker genes from our original single cell dataset. To identify where our

original cells localized within the integrated dataset, we mapped the original cell IDs to the

integrated clusters and calculated percentage overlap.

Acknowledgements

The authors thank the patients and their families for their participation, without whom these studies would not have been possible.

This study was supported by NIH funding U01 AG046193, RF1 AG051504, R01 AG051504 to NET.

We thank our colleagues in the neurosurgery team Christopher Louie, Karim ReFaey and Ivan Segura Duran. We thank our colleagues at the Mayo Clinic Genome Analysis Core (GAC) for their collaboration, particular gratitude to Bruce Eckloff and Julie Lau. We also acknowledge the AMP-AD RNAseq reprocessing team, in particular Dr. Kirsten Dang, Dr. Thanneer Perumal and Dr. Ben Logsdon at Sage Bionetworks.

AMP-AD RNASeq datasets: This study is a cross-consortia project using RNAseq data generated through grants U01AG046152, U01AG046170, and U01AG046139. **For the Mayo RNAseq study:** The results published here are in whole or in part based on data obtained from the AD Knowledge Portal (**https://adknowledgeportal.synapse.org/**). Study data were provided by the following sources: The Mayo Clinic Alzheimers Disease Genetic Studies, led by Dr. Nilüfer Ertekin-Taner and Dr. Steven G. Younkin, Mayo Clinic, Jacksonville, FL using samples from the Mayo Clinic Study of Aging, the Mayo Clinic Alzheimers Disease Research Center, and the Mayo Clinic Brain Bank. Data collection was supported through funding by NIA grants P50 AG016574, R01 AG032990, U01 AG046139, R01 AG018023, U01 AG006576, U01 AG006786, R01 AG025711, R01 AG017216, R01 AG003949, NINDS grant R01 NS080820, CurePSP Foundation, and support from Mayo Foundation. Study data includes samples collected through the Sun Health Research Institute Brain and Body Donation Program of Sun City, Arizona. The Brain and Body Donation Program is supported by the National Institute of Neurological Disorders and Stroke (U24 NS072026 National Brain and Tissue Resource for Parkinson's Disease and Related Disorders), the National Institute on Aging (P30 AG19610 Arizona Alzheimer's Disease Core Center), the Arizona Department

of Health Services (contract 211002, Arizona Alzheimer's Research Center), the Arizona

Biomedical Research Commission (contracts 4001, 0011, 05-901 and 1001 to the Arizona

Parkinson's Disease Consortium) and the Michael J. Fox Foundation for Parkinson's

Research. **For the ROSMAP study**: The results published here are in whole or in part based

on data obtained from the AD Knowledge Portal (**https://adknowledgeportal.synapse.org**).

Study data were provided by the Rush Alzheimer's Disease Center, Rush University Medical

Center, Chicago. Data collection was supported through funding by NIA grants P30AG10161

(ROS), R01AG15819 (ROSMAP; genomics and RNAseq), R01AG17917 (MAP),

R01AG30146, R01AG36042 (5hC methylation, ATACseq), RC2AG036547 (H3K9Ac),

R01AG36836 (RNAseq), R01AG48015 (monocyte RNAseq) RF1AG57473 (single nucleus

RNAseq), U01AG32984 (genomic and whole exome sequencing), U01AG46152 (ROSMAP

AMP-AD, targeted proteomics), U01AG46161(TMT proteomics), U01AG61356 (whole

genome sequencing, targeted proteomics, ROSMAP AMP-AD), the Illinois Department of

Public Health (ROSMAP), and the Translational Genomics Research Institute (genomic).

Additional phenotypic data can be requested at www.radc.rush.edu. **For the MSBB study:**

The results published here are in whole or in part based on data obtained from the AD

Knowledge Portal (**https://adknowledgeportal.synapse.org/**). These data were generated

from postmortem brain tissue collected through the Mount Sinai VA Medical Center Brain Bank

and were provided by Dr. Eric Schadt from Mount Sinai School of Medicine.

Data Sharing Statement

The data in this manuscript are available via the AD Knowledge Portal

(**https://adknowledgeportal.synapse.org**). The AD Knowledge Portal is a platform for

accessing data, analyses and tools generated by the Accelerating Medicines Partnership

(AMP-AD) Target Discovery Program and other National Institute on Aging (NIA)-supported

programs to enable open-science practices and accelerate translational learning. The data,

analyses and tools are shared early in the research cycle without a publication embargo on

secondary use. Data is available for general research use according to the following

requirements for data access and data attribution

(**https://adknowledgeportal.synapse.org/DataAccess/Instructions**).

Dataset	Data Type	Description	SynapseID	DoD
Mayo RNAseq TCX	Metadata	Individual human and RNAseq	syn5550404	na
Mayo RNAseq CER	Metadata	Individual human and RNAseq	syn5550404	na
Mayo RNAseq TCX	RNASeq Expression	Consensus processed RNASeq raw counts	syn8690799	10/2/2019
Mayo RNAseq CER	RNASeq Expression	Consensus processed RNASeq raw counts	syn8690904	10/2/2019
ROSMAP	Metadata	ID Key	syn3382527	10/2/2019
ROSMAP	Metadata	Individual human	syn3191087	10/2/2019
ROSMAP	Metadata	Assay RNAseq	syn21088596	1/2/2020
ROSMAP	RNASeq Expression	Consensus processed RNASeq raw counts	syn8691134	10/2/2019
MSBB	Metadata	Individual human	syn6101474	11/22/2019
MSBB	Metadata	Assay RNAseq	syn6100548	10/2/2019
MSBB	RNASeq Expression	Consensus processed RNASeq raw counts	syn8691099	10/2/2019

787 **Data from AD knowledge portal utilized in this study.** DoD = Date of download, "na" indicates data
788 that was generated by study authors and shared within the AD knowledge portal. Data were obtained
789 from the RNAseq Harmonization Study in the AD knowledge portal (DOI:
790 https://doi.org/10.7303/syn9702085).

Author Contributions

TP and NET wrote the manuscript; NET and MA designed the study; TP, XW and ZQ

performed data analysis; JC consulted on statistical methods; TP, TPC, XW, YM, RMA

generated tables and figures; EM, CAG, SG, KC, RW, HGC and AQH provided neurosurgical

tissue samples; TP, TPC, LJLT, SJL, SL, FQTN, CCGH, KGM, and TN performed

experimental procedures from blood and tissue samples. All authors read the manuscript and

provided input and consultation. NET oversaw the study and provided direction, funding and

resources.

Competing Financial Interests

801 None

References

1. Masuda, T., Sankowski, R., Staszewski, O. & Prinz, M. Microglia heterogeneity in the single-cell era. *Cell Reports* **30**, 1271-1281 (2020).
2. Li, Q. & Barres, B.A. Microglia and macrophages in brain homeostasis and disease. *Nat Rev Immunol* **18**, 225-242 (2018).
3. Hickman, S., Izzy, S., Sen, P., Morsett, L. & El Khoury, J. Microglia in neurodegeneration. *Nat Neurosci* **21**, 1359-1369 (2018).
4. Keren-Shaul, H.*, et al.* A Unique Microglia Type Associated with Restricting Development of Alzheimer's Disease. *Cell* **169**, 1276-1290 e1217 (2017).
5. Krasemann, S.*, et al.* The TREM2-APOE pathway drives the transcriptional phenotype of dysfunctional microglia in neurodegenerative diseases. *Immunity* **47**, 566-581 e569 (2017).
6. Mathys, H.*, et al.* Single-cell transcriptomic analysis of Alzheimer's disease. *Nature* **570**, 332-337 (2019).
7. Del-Aguila, J.L.*, et al.* A single-nuclei RNA sequencing study of Mendelian and sporadic AD in the human brain. *Alzheimer's Research & Therapy* **11**, 71 (2019).
8. Thrupp, N.*, et al.* Single-nucleus RNA-seq is not suitable for detection of microglial activation genes in humans. *Cell Reports* **32**, 108189 (2020).
9. Olah, M.*, et al.* Single cell RNA sequencing of human microglia uncovers a subset associated with Alzheimer's disease. *Nat Commun* **11**, 6129 (2020).
10. Masuda, T.*, et al.* Spatial and temporal heterogeneity of mouse and human microglia at single-cell resolution. *Nature* **566**, 388-392 (2019).
11. Sankowski, R.*, et al.* Mapping microglia states in the human brain through the integration of high-dimensional techniques. *Nat Neurosci* **22**, 2098-2110 (2019).
12. Darmanis, S.*, et al.* Single-Cell RNA-Seq Analysis of Infiltrating Neoplastic Cells at the Migrating Front of Human Glioblastoma. *Cell Rep* **21**, 1399-1410 (2017).
13. Morin-Brureau, M.*, et al.* Microglial phenotypes in the human epileptic temporal lobe. *Brain* **141**, 3343-3360 (2018).
14. Kinoshita, S. & Koyama, R. Pro- and anti-epileptic roles of microglia. *Neural Regeneration Research* **16**, 1369-1371 (2021).
15. Gosselin, D.*, et al.* An environment-dependent transcriptional network specifies human microglia identity. *Science* **356**(2017).
16. Srinivasan, K.*, et al.* Alzheimer's Patient Microglia Exhibit Enhanced Aging and Unique Transcriptional Activation. *Cell Rep* **31**, 107843 (2020).
17. Galatro, T.F.*, et al.* Transcriptomic analysis of purified human cortical microglia reveals age-associated changes. *Nat Neurosci* **20**, 1162-1171 (2017).
18. Olah, M.*, et al.* A transcriptomic atlas of aged human microglia. *Nature Communications* **9**, 539 (2018).
19. Frigerio, C.*, et al.* The major risk factors for Alzheimer's Disease: Age, sex, and genes modulate the microglia response to amyloid-beta plaques. *Cell Reports* **27**, 1293-1306 e1296 (2019).
20. Stephen, T.L.*, et al.* APOE genotype and sex affect microglial interactions with plaques in Alzheimer's disease mice. *Acta Neuropathol Commun* **7**, 82 (2019).
21. Nelson, L.H., Warden, S. & Lenz, K.M. Sex differences in microglial phagocytosis in the neonatal hippocampus. *Brain, Behavior, and Immunity* **64**, 11-22 (2017).

22. Yamazaki, Y., Zhao, N., Caulfield, T.R., Liu, C.C. & Bu, G. Apolipoprotein E and Alzheimer
disease: pathobiology and targeting strategies. *Nat Rev Neurol* **15**, 501-518 (2019).

23. Hammond, T.R*., et al.* Single-cell RNA sequencing of microglia throughout the mouse
lifespan and in the injured brain reveals complex cell-state changes. *Immunity* **50**, 253-
271 e256 (2019).

24. Grubman, A*., et al.* A single-cell atlas of entorhinal cortex from individuals with
Alzheimer's disease reveals cell-type-specific gene expression regulation. *Nat
Neurosci* **22**, 2087-2097 (2019).

25. Gamache, J., Yun, Y. & Chiba-Falek, O. Sex-dependent effect of APOE on Alzheimer's
disease and other age-related neurodegenerative disorders. *Disease Models &
Mechanisms* **13**, dmm045211 (2020).

26. Allen, M*., et al.* Human whole genome genotype and transcriptome data for
Alzheimer's and other neurodegenerative diseases. *Sci Data* **3**, 160089 (2016).

27. Wang, M.*, et al.* The Mount Sinai cohort of large-scale genomic, transcriptomic and
proteomic data in Alzheimer's disease. *Sci Data* **5**, 180185 (2018).

28. De Jager, P.L.*, et al.* A multi-omic atlas of the human frontal cortex for aging and
Alzheimer's disease research. *Sci Data* **5**, 180142 (2018).

29. McKenzie, A.T.*, et al.* Brain Cell Type Specific Gene Expression and Co-expression
Network Architectures. *Sci Rep* **8**, 8868 (2018).

30. Wang, X.*, et al.* Deciphering cellular transcriptional alterations in Alzheimer's disease
brains. *Mol Neurodegener* **15**, 38 (2020).

31. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics* **9**, 559 (2008).

32. Chausse, B., Kakimoto, P.A. & Kann, O. Microglia and lipids: how metabolism controls
brain innate immunity. *Semin Cell Dev Biol* (2020).

33. Loving, B.A. & Bruce, K.D. Lipid and Lipoprotein Metabolism in Microglia. *Front Physiol*
11, 393 (2020).

34. Marschallinger, J*., et al.* Lipid-droplet-accumulating microglia represent a
dysfunctional and proinflammatory state in the aging brain. *Nat Neurosci* (2020).

35. Bernier, L.P., York, E.M. & MacVicar, B.A. Immunometabolism in the Brain: How
Metabolism Shapes Microglial Function. *Trends Neurosci* **43**, 854-869 (2020).

36. El Khoury, J., Hickman, S.E., Thomas, C.A., Loike, J.D. & Silverstein, S.C. Microglia,
scavenger receptors, and the pathogenesis of Alzheimer's disease. *Neurobiol Aging*
19, S81-84 (1998).

37. DePaula-Silva, A.B.*, et al.* Differential transcriptional profiles identify microglial- and
macrophage-specific gene markers expressed during virus-induced
neuroinflammation. *J Neuroinflammation* **16**, 152-152 (2019).

38. Butovsky, O. & Weiner, H.L. Microglial signatures and their role in health and disease.
Nature Reviews Neuroscience **19**, 622-635 (2018).

39. Colombo, A*., et al.* Loss of NPC1 enhances phagocytic uptake and impairs lipid
trafficking in microglia. *Nature Communications* **12**, 1158 (2021).

40. Mehra, A., Ali, C., Parcq, J., Vivien, D. & Docagne, F. The plasminogen activation system
in neuroinflammation. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of
Disease* **1862**, 395-402 (2016).

41. Cunningham, O*., et al.* Microglia and the urokinase plasminogen activator
receptor/uPA system in innate brain inflammation. *Glia* **57**, 1802-1814 (2009).

42. Sobue, A.*, et al.* Microglial gene signature reveals loss of homeostatic microglia
associated with neurodegeneration of Alzheimer's disease. *Acta Neuropathologica*
Communications **9**, 1 (2021).
43. Rangaraju, S.*, et al.* Identification and therapeutic modulation of a pro-inflammatory
subset of disease-associated-microglia in Alzheimer's disease. *Mol Neurodegener* **13**,
24 (2018).
44. Rangaraju, S.*, et al.* Quantitative proteomics of acutely-isolated mouse microglia
identifies novel immune Alzheimer's disease-related proteins. *Mol Neurodegener* **13**,
34 (2018).
45. Gerrits, E.*, et al.* Distinct amyloid-β and tau-associated microglia profiles in
Alzheimer's disease. *Acta Neuropathologica* **141**, 681-696 (2021).
46. Cermak, S.*, et al.* Loss of Cathepsin B and L Leads to Lysosomal Dysfunction, NPC-Like
Cholesterol Sequestration and Accumulation of the Key Alzheimer's Proteins. *PLOS*
ONE **11**, e0167428 (2016).
47. Xu, H., Han, Y., Liu, B. & Li, R. Unc-5 homolog B (UNC5B) is one of the key downstream
targets of N-α-Acetyltransferase 10 (Naa10). *Scientific Reports* **6**, 38508 (2016).
48. Ahn, E.H., Kang, S.S., Qi, Q., Liu, X. & Ye, K. Netrin1 deficiency activates MST1 via
UNC5B receptor, promoting dopaminergic apoptosis in Parkinson's disease.
Proceedings of the National Academy of Sciences **117**, 24503-24513 (2020).
49. Zhang, X.*, et al.* A rare missense variant of CASP7 is associated with familial late-onset
Alzheimer's disease. *Alzheimers Dement* **15**, 441-452 (2019).
50. Ayers, K.L.*, et al.* A loss of function variant in CASP7 protects against Alzheimer's
disease in homozygous APOE ε4 allele carriers. *BMC Genomics* **17 Suppl 2**, 445 (2016).
51. Katsouri, L. & Georgopoulos, S. Lack of LDL receptor enhances amyloid deposition and
decreases glial response in an Alzheimer's disease mouse model. *PLoS One* **6**, e21880
(2011).
52. Lämsä, R.*, et al.* Genetic study evaluating LDLR polymorphisms and Alzheimer's
disease. *Neurobiol Aging* **29**, 848-855 (2008).
53. Crotti, A.*, et al.* BIN1 favors the spreading of Tau via extracellular vesicles. *Scientific*
Reports **9**, 9477 (2019).
54. Sims, R.*, et al.* Rare coding variants in PLCG2, ABI3, and TREM2 implicate microglial-
mediated innate immunity in Alzheimer's disease. *Nature Genetics* **49**, 1373-1384
(2017).
55. Darmanis, S.*, et al.* A survey of human brain transcriptome diversity at the single cell
level. *Proceedings of the National Academy of Sciences* **112**, 7285 (2015).
56. Zhou, Y.*, et al.* Human and mouse single-nucleus transcriptomics reveal TREM2-
dependent and TREM2-independent cellular responses in Alzheimer's disease. *Nat*
Med **26**, 131-142 (2020).
57. Schmunk, G.*, et al.* Human microglia upregulate cytokine signatures and accelerate
maturation of neural networks. *bioRxiv*, 2020.2003.2024.006874 (2020).
58. Aung, L.L.*, et al.* Multiple sclerosis-linked and interferon-beta-regulated gene
expression in plasmacytoid dendritic cells. *J Neuroimmunol* **250**, 99-105 (2012).
59. Satoh, J.*, et al.* Microarray analysis identifies an aberrant expression of apoptosis and
DNA damage-regulatory genes in multiple sclerosis. *Neurobiol Dis* **18**, 537-550 (2005).
60. Gandhi, K.S.*, et al.* The multiple sclerosis whole blood mRNA transcriptome and
genetic associations indicate dysregulation of specific T cell pathways in pathogenesis.
Hum Mol Genet **19**, 2134-2143 (2010).

61. Hendrickx, D.A.E., van Eden, C.G., Schuurman, K.G., Hamann, J. & Huitinga, I. Staining of HLA-DR, Iba1 and CD68 in human microglia reveals partially overlapping expression depending on cellular morphology and pathology. *Journal of Neuroimmunology* **309**, 12-22 (2017).

62. Kenkhuis, B.*, et al.* Iron loading is a prominent feature of activated microglia in Alzheimer's disease patients. *Acta Neuropathol Commun* **9**, 27 (2021).

63. Kalari, K.R.*, et al.* MAP-RSeq: Mayo Analysis Pipeline for RNA sequencing. *BMC Bioinformatics* **15**, 224 (2014).

64. Allen, M.*, et al.* Divergent brain gene expression patterns associate with distinct cell-specific tau neuropathology traits in progressive supranuclear palsy. *Acta Neuropathol* **136**, 709-727 (2018).

65. Allen, M.*, et al.* Conserved brain myelination networks are altered in Alzheimer's and other neurodegenerative diseases. *Alzheimers Dement* **14**, 352-366 (2018).

66. Mostafavi, S.*, et al.* A molecular network of the aging human brain provides insights into the pathology and cognitive decline of Alzheimer's disease. *Nat Neurosci* **21**, 811-819 (2018).

67. Neff, R.A.*, et al.* Molecular subtyping of Alzheimer's disease using RNA sequencing data reveals novel mechanisms and targets. *Sci Adv* **7**(2021).

68. Alsema, A.M.*, et al.* Profiling Microglia From Alzheimer's Disease Donors and Non-demented Elderly in Acute Human Postmortem Cortical Tissue. *Frontiers in Molecular Neuroscience* **13**(2020).

69. Shi, Y.*, et al.* Microglia drive APOE-dependent neurodegeneration in a tauopathy mouse model. *Journal of Experimental Medicine* **216**, 2546-2561 (2019).

70. Nguyen, A.T.*, et al.* APOE and TREM2 regulate amyloid-responsive microglia in Alzheimer's disease. *Acta Neuropathologica* **140**, 477-493 (2020).

71. Leduc, V., Jasmin-Belanger, S. & Poirier, J. APOE and cholesterol homeostasis in Alzheimer's disease. *Trends Mol Med* **16**, 469-477 (2010).

72. Paciotti, S., Albi, E., Parnetti, L. & Beccari, T. Lysosomal Ceramide Metabolism Disorders: Implications in Parkinson's Disease. *J Clin Med* **9**(2020).

73. Nugent, A.A.*, et al.* TREM2 Regulates Microglial Cholesterol Metabolism upon Chronic Phagocytic Challenge. *Neuron* **105**, 837-854 e839 (2020).

74. Yeh, F.L., Wang, Y., Tom, I., Gonzalez, L.C. & Sheng, M. TREM2 Binds to Apolipoproteins, Including APOE and CLU/APOJ, and Thereby Facilitates Uptake of Amyloid-Beta by Microglia. *Neuron* **91**, 328-340 (2016).

75. Norden, D.M. & Godbout, J.P. Review: microglia of the aged brain: primed to be activated and resistant to regulation. *Neuropathol Appl Neurobiol* **39**, 19-34 (2013).

76. Koellhoffer, E.C., McCullough, L.D. & Ritzel, R.M. Old maids: Aging and its impact on microglia function. *Int J Mol Sci* **18**(2017).

77. Ulland, T.K.*, et al.* TREM2 Maintains Microglial Metabolic Fitness in Alzheimer's Disease. *Cell* **170**, 649-663 e613 (2017).

78. Lin, Y.T.*, et al.* APOE4 Causes Widespread Molecular and Cellular Alterations Associated with Alzheimer's Disease Phenotypes in Human iPSC-Derived Brain Cell Types. *Neuron* **98**, 1141-1154 e1147 (2018).

79. Andreone, B.J.*, et al.* Alzheimer's-associated PLCgamma2 is a signaling node required for both TREM2 function and the inflammatory response in human microglia. *Nat Neurosci* **23**, 927-938 (2020).

80. Grassivaro, F.*, et al.* Convergence between Microglia and Peripheral Macrophages
Phenotype during Development and Neuroinflammation. *The Journal of Neuroscience*
40, 784-795 (2020).
81. Chrétien, F.*, et al.* Expression of Excitatory Amino Acid Transporter-1 (EAAT-1) in Brain
Macrophages and Microglia of Patients with Prion Diseases. *Journal of*
Neuropathology & Experimental Neurology **63**, 1058-1071 (2004).
82. Wilhelmsson, U.*, et al.* Injury Leads to the Appearance of Cells with Characteristics of
Both Microglia and Astrocytes in Mouse and Human Brain. *Cerebral Cortex* **27**, 3360-
3377 (2017).
83. Kaminska, B., Ochocka, N. & Segit, P. Single-cell omics in dissecting immune
microenvironment of malignant gliomas: Challenges and perspectives. *Cells* **10**(2021).
84. Ochocka, N.*, et al.* Single-cell RNA sequencing reveals functional heterogeneity of
glioma-associated brain macrophages. *Nature Communications* **12**, 1151 (2021).
85. Dobin, A.*, et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).
86. Liao, Y., Smyth, G.K. & Shi, W. featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).
87. Wan, Y.W.*, et al.* Meta-Analysis of the Alzheimer's Disease Human Brain
Transcriptome and Functional Dissection in Mouse Models. *Cell Rep* **32**, 107908
(2020).
88. McKhann, G.*, et al.* Clinical diagnosis of Alzheimer's disease: report of the NINCDS-
ADRDA Work Group under the auspices of Department of Health and Human Services
Task Force on Alzheimer's Disease. *Neurology* **34**, 939-944 (1984).
89. Supek, F., Bosnjak, M., Skunca, N. & Smuc, T. REVIGO summarizes and visualizes long
lists of gene ontology terms. *PLoS One* **6**, e21800 (2011).
90. Zheng, G.X.Y.*, et al.* Massively parallel digital transcriptional profiling of single cells.
Nature Communications **8**, 14049 (2017).
91. Stuart, T.*, et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-
1902.e1821 (2019).
1016
1017

 Main Figures

Figure 1. Characterization of our core human microglial signature. (A) Schematic illustrating our experimental approach for isolating microglial populations from fresh brain tissue and data analyses. [Created with BioRender.com] (B) MSigDB GO terms enriched in our microglial signature genes and top 25 genes for each. (C) Venn diagram showing number of overlapping genes between our microglial signature and those previously reported from Galatro et al (2017), Gosselin et al (2017) and Olah et al (2018). (D) Hypergeometric tests of overrepresentation showing overlap with the published signatures.

1019

1020

Figure 2. Age, sex and *APOE* ε4 pathway correlations in bulk microglia. (a) Heatmap showing correlation of age, sex and *APOE* ε4 status with WGCNA module eigengenes (MEs) significantly associated (p<0.05) with traits, with top GO terms listed for each module. (b) Module eigengenes stratified by age or *APOE* ε4 status. (c) Module M14 gene co-expression network, with genes of interest highlighted according to the key. Genes upregulated with age shown in red triangle (▲). Bar plot of top 10 significant GO terms (p<0.05) for this module. (d) Module 23 gene co-expression network, with genes downregulated in *APOE* ε4 carriers shown in blue arrow (▼). Bar plot of top 10 significant GO terms (p<0.05) for this module. (e) Module 26 gene co-expression network, with genes upregulated in *APOE* ε4 carriers shown in orange triangle (▲). Bar plot of top 10 significant GO terms (p<0.05) for this module. (f) Violin plots showing expression of key genes in modules, stratified by age or *APOE*. * p < 0.05; ** p < 0.01; *** p < 0.001.

Figure 3. Single cell microglial data. (A) UMAP of clustered cells annotated with putative subtypes using cell type markers from the literature. (B) Stacked bar plot showing the distribution of cells across the clusters. (C) Dot plot showing the expression of key significant module genes across clusters. (D) Hierarchical clustering to highlight relationships between clusters. (E) Hypergeometric distribution of enrichment between module genes and clusters, showing number of overlapping genes. * Represents module genes that were significantly enriched in the cluster (p < 0.05).

1021

Figure 4. Meta-analysis with published datasets. (A) Forest plots of module eigengene correlations across individual bulk microglial datasets. Results from meta-analyses of relevant samples are shown in bold. (B) UMAPs of Olah et al's (2020) (left) and our (right) single cell data, split by dataset. Samples were integrated prior to clustering and downstream analyses. Cells from each dataset contribute to all clusters. (C) Stacked bar plot showing the distribution of cells across each of the clusters. (D) Dot plot showing the expression of top cluster marker genes across all clusters.

1022