Title:

An ultrasensitive method for analysis of viral spike N-glycoforms

Authors:

Sabyasachi Baboo*, Jolene K Diedrich, Salvador Martínez-Bartolomé, Xiaoning Wang, Torben

Schiffner, Bettina Groschel, William R Schief, James C Paulson & John R Yates III*

Affiliations:

Department of Molecular Medicine, The Scripps Research Institute, La Jolla, CA, USA

Sabyasachi Baboo, Jolene K Diedrich, Salvador Martínez-Bartolomé, Xiaoning Wang, James C

Paulson and John R Yates III

IAVI Neutralizing Antibody Center, The Scripps Research Institute, La Jolla, CA, USA

Torben Schiffner, Bettina Groschel, William R Schief

Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla,

CA, USA

Torben Schiffner, Bettina Groschel, William R Schief, James C Paulson

The Ragon Institute of Massachusetts General Hospital, Massachusetts Institute of

Technology and Harvard, Cambridge, MA, USA

Torben Schiffner, William R Schief

Present address:

Institut für Wirkstoffentwicklung, Universität Leipzig, Leipzig, Germany

Torben Schiffner

Corresponding authors:

Correspondence to Sabyasachi Baboo (sbaboo@scripps.edu) or John R Yates III

(jyates@scripp.sedu)

Keywords:

proteomics, mass spectrometry, Proteinase K, N-glycan, heterogeneity, HIV, Env, viral spike

Words:

1,462 (abstract/70 + main)

Abstract:

Viruses can evade the host immune system by displaying numerous glycans on their surface

"spike-proteins" that cover immune epitopes. We have developed an ultrasensitive "single pot"

method to assess glycan occupancy and the extent of glycan processing from high-mannose to

complex forms at each N-glycosylation site. Though aimed at characterizing glycosylation of

viral spike proteins as potential vaccines, this method is applicable for analysis of site-specific

glycosylation of any glycoprotein.

Main:

Viral spike-proteins initiate virus entry into host cells and are the primary targets of vaccine

design. Spike-proteins are often heavily N-glycosylated which help to shield the protein from the

host immune response[1]. These glycans add complexity to the production and characterization of

recombinant protein-based vaccines[2]. This is particularly a concern for characterization of the

envelope spike-protein (*Env*) of the Human Immunodeficiency Virus (HIV) comprising a trimer

with each monomer containing 26-30 unique N-linked glycosylation sites (NGS) as defined by

the sequon NX[S|T], where X is any amino acid except P[3]. To address this analytical challenge

several mass spectrometry-based strategies using multiple proteases[4-6] have been implemented to

create sufficient numbers of peptides unique to each glycosylation site[3, 7, 8]. In these strategies,

individual aliquots are digested with each protease and analyzed separately by liquid

chromatography-mass spectrometry (LC-MS/MS), or pooled and analyzed together. To broadly

characterize the nature of the glycosylation at each NGS, we had previously introduced the use

of endoglycosidases that create residual mass signatures[3]. This helped us to determine the degree

of glycan occupancy, and the degree of glycan processing – from the high mannose form that is

initially attached to the protein, which may mature into the complex form when mannose

residues are replaced by "terminal" monosaccharide sequences. To achieve coverage for all

NGS, we had combined several digestions performed with different proteases and achieved

>95% sequence coverage. Here we show that it is possible to replace these multiple proteolytic

digestions with a single Proteinase K (PK) digestion, and moreover through careful choice of

volatile buffers we have developed an improved "single pot" strategy with significantly

increased sensitivity.

PK is a broadly specific serine-protease that has previously been exploited for its potential to

generate overlapping peptides and high sequence coverage[9]. The redundancy afforded by

overlapping sequences significantly increases confidence in identifications, especially when

covalent modifications are present[4]. However, because proteinase K is an aggressive protease it

is necessary to attenuate its proteolytic activity to obtain high sequence coverage of proteins.

Attenuation of PK can be achieved using suboptimal reaction conditions[9] (to reduce the rate of

enzyme activity) and limited reaction time. Using a mildly acidic, chaotrope-free solution to

attenuate the activity of PK, we were able to achieve >95% sequence coverage of candidate viral

spike proteins and identify all NGS. The proteomic strategy of our new method is conceptually

similar to our previous approach[3], but it is significantly faster and more sensitive. These

improvements result from three key changes to the strategy: [1] using only mass spectrometry-

compatible constituents, samples are processed in a single solution except the final step of

PNGase F deglycosylation; [2] reaction volumes are kept to a minimum (5-8 µl) to increase the

rate of reaction to limit sample loss on surfaces and minimize freeze-drying time; and [3] the use

of PK provides faster digestion and excellent sequence and NGS coverage with less starting

material. Our new method reduces sample preparation time from 3 days to 1 day, reduces LC-

MS/MS run times by 9-24 fold, and can achieve 50-180 times higher sensitivity than existing

methods[3, 7, 8]. In addition, we have developed the data analysis tool GlycoMSQuant which further

reduces analysis times and simplifies data analysis.

We tested our approach on BG505 SOSIP.664 MD39[10], a stabilized native-like HIV *Env* trimer

being developed for an HIV vaccination strategy[11] targeting germline precursors of broadly

neutralizing antibodies (bNAbs) that are impacted by N-glycans. As in our previous approach[3],

the glycosylated peptides generated by PK digestion were sequentially deglycosylated; first with

Endo H to remove high-mannose and hybrid N-glycans, and then with PNGase F, which

removes all remaining N-glycans. The resulting residual masses on asparagine (N) in NGS is

+203 Da at sites occupied by high-mannose/hybrid N-glycans or +3 Da at sites occupied by

complex N-glycans when PNGase F deglycosylation is carried out in the presence of $H_2^{18}O$

(differentiating these sites from any deamidated Ns). Unoccupied NGS results in no (+0 Da)

residual mass on N. Using this new method (***Fig. 1a***), we achieved >99% amino acid sequence

coverage and identified all theoretically possible 27 NGSs from a single LC-MS/MS run of 0.5

µg of peptides generated from a starting material of 5 µg purified protein (***Fig. 1b*** and

Supplementary Figures 1a,b). We used semi-quantitative label-free analysis based on precursor

peak areas to calculate the proportion of N-glycan occupancy (unoccupied: complex: high-

mannose/hybrid N-glycans) for each NGS. We reanalyzed the N-glycan microheterogeneity

pattern on BG505 SOSIP.664[12] HIV *Env* trimer from data obtained using our previous approach[3]

and compared it with results using the method we describe here (***Supplementary Figures 2***). The

results with improved method were highly comparable to those using our original approach, in

spite of being processed differently and the samples being prepared at different times in different

laboratories.

Initial results demonstrated that the new approach is at least 18 times more sensitive than our

previous approach[3] even though it uses a simpler and shorter workflow. To evaluate the limit of

sensitivity of our method, we processed progressively decreasing amounts of starting material,

ranging from 1 µg to 5 ng. We observed that a single LC-MS/MS run with 1 µg of starting

material was enough to cover >95% of the amino acid sequence and all NGS (***Fig. 2a***), which is

90 times more sensitive than our previous approach[3]. Major differences in microheterogeneity at

each NGS were generally observed when we started with <100 ng material (***Supplementary***

Figure 1c). This is likely due to low sampling as evidenced by a decrease in amino acid

sequence and NGS coverage (*Fig. 2a*), as well as the absolute number of identified peptides

representing each NGS (*Supplementary Figure 1d*).

The improved method is agnostic to mass spectrometry platform (*Fig. 2b*). A timsTOF Pro mass-

spectrometer coupled to an Evosep One HPLC (timsTOF/Evosep)[13] was used to achieve >99%

sequence coverage and identification of all NGS using a single LC-MS/MS run with 0.5 µg of

starting material and an 88-minute LC gradient (*Fig. 1c* and *Supplementary Figures 3a,b*).

Thus, the sensitivity of our method on this platform was 180 times higher than our previous

approach[3].

N-glycan heterogeneity reflects the immunogenicity of the viral spike-protein and is critical for

designing vaccines[14]. The reproducibility of N-glycan heterogeneity patterns obtained with our

new method suggests that this is a robust procedure. Variability in sequence coverage are not

observed until the limits of detection are reached on an LC-MS/MS platform. Although our

results within the same LC-MS platform were reproducible (except some variations when using

different proteases, *Supplementary Figure 5*), we may infer the effects of sampling differences

when comparing two different LC-MS platforms (QE-HFX *vs.* timsTOF/Evosep), as in case of

N156, N160, N197, N386, N392 (relative peptide abundance is persistently low) and N88, N295,

N301, N332, N355, N406, N411 (possible skewing of timsTOF/Evosep identification against

N+203 peptides when peptide sampling per NGS decreases due to less starting material) (*Fig. 2b*

and *Supplementary Figures 1c,d* and *3c,d*). When enough sampling per NGS is achieved, these

variations are diminished (*Figs. 1b,c*).

138 We attribute improvements in our method to efficient sample handling strategies. We observed

139 reduced sequence coverage if the sample was from the digestion of a small amount of starting

140 material rather than an equal aliquot from a larger sample (***Supplementary Figure 3e***). We infer

141 that the sensitivity differences are not occurring during LC-MS/MS, but that sample is being lost

142 to the reaction-tube surface (during reaction and lyophilization) and the proportion of loss is

143 more pronounced when we start with less material. The kinetics of the enzyme-substrate reaction

144 may also account for sensitivity differences since a more "crowded" reactant environment (low

145 reaction volumes) is expected to result in better reaction kinetics[15].

147 The simplicity and high reproducibility of this procedure will allow for high-throughput analyses

148 of viral spike-proteins and for any glycoprotein whether produced recombinantly or purified

149 from natural sources. Results were highly comparable in the two LC-MS/MS platforms we used.

150 A single analysis using a QE-HFX/nLC platform can determine the complete N-glycan

151 heterogeneity pattern from 1 microgram of purified viral spike-protein. The timsTOF/Evosep

152 platform was observed to be more sensitive than QE-HFX/nLC across all NGS, although limited

153 sampling may not allow us to confidently infer N-glycan microheterogeneity at all NGS. We

154 view the high sensitivity of this robust method to be an important step to analysis of the

155 glycosylation of more complex samples such as whole virus or virus in infected blood[16].

157 **Methods:**

159 Expression and purification of HIV *Env* trimers

BG505 SOSIP.664[12] and BG505 SOSIP.664 MD39[10] *Env* trimers were expressed and purified

essentially as described previously[10]. Briefly, sequences with codons optimized for expression in

human cells were synthesized and cloned into pHLSec between *AgeI*/*KpnI* by Genscript. The

constructs were co-transfected with Furin-encoding plasmid, using polyethylenimine in Freestyle

293F cells cultured in 293 FreeStyle media (Thermo Fisher Scientific). Where indicated, 15 μM

sterile-filtered Kifunensine (Cayman Chemical) was added after transfection. After 6-7 days,

supernatant was collected after passing through 0.22 μm filter (Nalgene), and the C-terminally

His-tagged trimers were purified using a HisTrap affinity column (Cytiva) with a linear elution

gradient from 20-500 mM imidazole, followed by a Superdex 200 Increase SEC column (Cytiva)

in Tris-buffered saline/TBS (20 mM Tris, 100 mM NaCl, pH 7.5). The oligomeric state and

purity of trimer was verified using size exclusion chromatography coupled with multi-angle light

scattering (SEC-MALS; DAWN HELEOS II/ Optilab T-rEX, Wyatt Technology).

Proteinase K treatment and deglycosylation

HIV Env trimer was exchanged to water using Microcon Ultracel-10 centrifugal device

(Millipore Sigma). Trimer was reduced with 5 mM tris(2-carboxyethyl)phosphine hydrochloride

(TCEP-HCl, Thermo Scientific) and alkylated with 10 mM 2-chloroacetamide (Sigma Aldrich)

in 100 mM ammonium acetate for 20 min at room temperature (RT, 24°C). Initial protein-level

deglycosylation was performed using 250 U of Endo H (New England Biolabs) for up to 5 μg

trimer, for 1 h at 37°C (pH 5.5-6.0). Trimer was digested with 1:25 Proteinase K (Sigma

Aldrich) for 4 h at 37°C (pH 5.5-6.0). PK was denatured by incubating at 90°C for 15 min, then

cooled to RT. Peptides were deglycosylated again with 250 U Endo H for 1 h at 37°C (pH 5.5-

6.0), then frozen at –80°C and lyophilized. 100 U PNGase F (New England Biolabs) was

lyophilized (for up to 5 µg trimer), resuspended in 100 mM ammonium bicarbonate prepared in

$H_2^{18}O$ (97% [18]O, Sigma-Aldrich), and added to the lyophilized peptides. The resulting 5-8 µl

reaction solutions (except PNGase F reaction, in 5-20 µl) were then incubated for 1 h at 37°C

(pH 8.0-8.5) in 0.2 ml PCR tubes on a thermocycler with heated lid.

Validating efficiency of glycosidases

BG505 SOSIP.664 HIV *Env* trimer glycosylated with only high-mannose N-glycans (purified

from cells treated with Kifunensine[17], which inhibits processing of high-mannose N-glycans to

complex N-glycans during protein maturation), was sequentially treated with Endo H, followed

by PNGase F. After treatment with both enzymes, 99.2% of identified peptides were N+203;

100% of peptides identified with only PNGase F treatment were N+3 (***Supplementary Figures***

4a,b; proportions do not consider unoccupied NGS because they remained similar in both

experiments – 8-10%). We realize the possibility that glycosidase PNGase F may occasionally

cleave the remnant GlcNAc (N-Acetylglucosamine)[18] post-Endo H processing of high-

mannose/hybrid N-glycans and thus, convert the mass modification characteristic of high-

mannose/hybrid (+203) to complex (+3) N-glycans, which would affect our analyses. However,

we have not observed any significant evidence of this possibility in our results, though it may

explain why we observe a few peptides with +3 mass modified NGS in Kifunensine treated

samples (***Supplementary Figure 4a***).

Trypsin proteolysis

The Proteinase K/deglycosylation method described above was followed, except PK was

replaced with trypsin and reactions were incubated overnight at 37°C. Trypsin generated a lower

total number of peptides than PK, but we obtained >95% sequence coverage, including 26 of 27

NGS (*Supplementary Figure 5a*). Variations in N-glycan microheterogeneity at certain NGS

may be explained by low sampling at these sites (N386, N392) or difference in cleavage-

specificity between PK and trypsin (N88, N611) (*Supplementary Figures 5b,c*).

<u>LC-MS/MS</u>

Q Exactive HF-X with EASY-nLC 1200

Samples were analyzed on an Q Exactive HF-X mass spectrometer (Thermo). Samples were

injected directly onto a 25 cm, 100 μm ID column packed with BEH 1.7 μm C18 resin (Waters).

Samples were separated at a flow rate of 300 nL/min on an EASY-nLC 1200 (Thermo). Buffers

A and B were 0.1% formic acid in 5% and 80% acetonitrile, respectively. The following

gradient was used: 1–25% B over 160 min, an increase to 40% B over 40 min, an increase to

90% B over another 10 min and 30 min at 90% B for a total run time of 240 min. Column was

re-equilibrated with solution A prior to the injection of sample. Peptides were eluted from the tip

of the column and nanosprayed directly into the mass spectrometer by application of 2.8 kV at

the back of the column. The mass spectrometer was operated in a data dependent mode. Full

MS1 scans were collected in the Orbitrap at 120,000 resolution. The ten most abundant ions per

scan were selected for HCD MS/MS at 25 NCE. Dynamic exclusion was enabled with exclusion

duration of 10 s and singly charged ions were excluded.

timsTOF Pro with Evosep One

Samples were loaded onto EvoTips following manufacturer protocol. The samples were run on

an Evosep One (Evosep) coupled to a timsTOF Pro (Bruker Daltonics). Samples were separated

on a 15 cm × 150 μm ID column with BEH 1.7 μm C18 beads (Waters) and integrated tip pulled

in-house using either the 30 SPD or 15 SPD methods. Mobile phases A and B were 0.1% formic

acid in water and 0.1% formic acid in acetonitrile, respectively. MS data was acquired in PASEF

mode with 1 MS1 survey TIMS-MS and 10 PASEF MS/MS scans acquired per 1.1 s acquisition

cycle. Ion accumulation and ramp time in the dual TIMS analyzer was set to 100 ms each and we

analyzed the ion mobility range from $1/K_0 = 0.6$ Vs cm^{-2} to 1.6 Vs cm^{-2}. Precursor ions for

MS/MS analysis were isolated with a 2 Th window for m/z < 700 and 3 Th for m/z >700 with a

total m/z range of 100-1700. The collision energy was lowered linearly as a function of

increasing mobility starting from 59 eV at $1/K_0 = 1.6$ VS cm^{-2} to 20 eV at $1/K_0 = 0.6$ Vs cm^{-2}.

Singly charged precursor ions were excluded with a polygon filter, precursors for MS/MS were

picked at an intensity threshold of 2,500, target value of 20,000 and with an active exclusion of

24 s.

<u>Data Processing</u>

Protein and peptide identification were done with Integrated Proteomics Pipeline (IP2, Bruker

Scientific LLC). Tandem mass spectra were extracted from raw files using RawConverter[19]

(timstofCoverter for timsTOF Pro data) and searched with ProLuCID[20] against a database

comprising UniProt reviewed (Swiss-Prot) proteome for *Homo sapiens* (UP000005640), UniProt

amino acid sequences for Endo H (P04067), PNGase F (Q9XBM8), and Proteinase K (P06873),

amino acid sequences for BG505 SOSIP.664[12] and BG505 SOSIP.664 MD39[10] (including a

preceding secretory signal sequence and followed by 6xHis-tag), and a list of general protein

contaminants. The search space included no cleavage-specificity (all fully tryptic and semi-

tryptic peptide candidates for trypsin treatment). Carbamidomethylation (+57.02146 C) was

considered a static modification. Deamidation in presence of $H_2^{18}O$ (+2.988261 N), GlcNAc

(+203.079373 N), oxidation (+15.994915 M) and N-terminal pyroglutamate formation (−

17.026549 Q) were considered differential modifications. Data was searched with 50 ppm

precursor ion tolerance and 50 ppm fragment ion tolerance. Identified proteins were filtered

using DTASelect2[21] and utilizing a target-decoy database search strategy to limit the false

discovery rate to 1%, at the spectrum level[22]. A minimum of 1 peptide per protein and no tryptic

end (or 1 tryptic end when treated with trypsin) per peptide were required and precursor delta

mass cut-off was fixed at 10 ppm for data acquired with Q Exactive HF-X or 20 ppm for data

acquired with timsTOF Pro. Statistical models for peptide mass modification (modstat) were

applied (trypstat was additionally applied for trypsin-treated samples). Census2[23] label-free

analysis was performed based on the precursor peak area, with a 10 ppm precursor mass

tolerance and 0.1 min retention time tolerance. "Match between runs" was used to find missing

peptides between runs for Q Exactive HF-X data (for timsTOF Pro data, reconstructed-MS1

based chromatograms combining isotope peaks for all triggered precursor ions were pre-

generated, and then chromatograms were assigned to identified peptides for quantitative analysis,

without retrieving missing peptides).

Data Analysis using GlycoMSQuant

Our new tool GlycoMSQuant v.1.4.1 (https://github.com/proteomicsyates/GlycoMSQuant) was

implemented to automate the analysis and to visualize the results. GlycoMSQuant summed

precursor peak areas across replicates, discarded peptides without NGS, discarded misidentified

peptides when N-glycan remnant-mass modifications were localized to non-NGS asparagines

and corrected/fixed N-glycan mislocalization where appropriate. The results were aligned to

NGS in *Env* of HXB2[24] HIV-1 variant.

Precursor peak area was calculated by Census2[23] from extracted-ion chromatogram (XIC) for

each peptide in each replicate. For each NGS (NX[S|T], where X is any amino acid except P), the

"N-glycosylation state" represented by proportions for unoccupied (+0, u), complex (+2.988261,

c) and, high-mannose/hybrid (+203.079373, h) N-glycans was calculated as follows.

The sum of the precursor peak areas $S_{g,pepz}$ was calculated as:

$$S_{g,pepz} = \sum xic_{pepz}$$

where N-glycosylated peptides with the same sequences and charge were grouped together

(*pepz*), g is the N-glycosylation state $\in G$ (u, c, h), and *xic* is the precursor peak area.

For each group (*pepz*), the abundance proportion $\%_{g,pepz}$ of each N-glycosylation state $g \in G$ was

calculated as:

$$\%_{g,pepz} = \frac{S_{g,pepz}}{\sum_{i \in G} S_{i,pepz}}$$

Finally, as each NGS may be covered by multiple groups (*pepz*), the proportion of each N-

glycosylation state g for a particular NGS (*ngs*) is calculated as the mean of all proportions

$\%_{g,pepz}$ of all groups (*pepz*) covering this NGS:

$$\%_{ngs,g} = \frac{1}{n_{ngs}} \sum \%_{g,pepz}$$

where n_{ngs} is the number of groups (*pepz*) covering a particular NGS.

The standard error of mean of the proportion of each N-glycosylation state $g \in G$ for a particular

NGS ($SEM_{ngs,g}$) was calculated as:

$$SEM_{ngs,g} = \frac{s_{ngs,g}}{\sqrt{n_{ngs}}}$$

where, $s_{ngs,g}$ is the standard deviation of $\%_{g,pepz}$ from all groups (*pepz*) covering a particular NGS.

Pairwise statistical comparisons of experiments (*a* and *b*) were performed for each $g \in G$ at each

NGS using proportion values $\%_{g,pepz}$ of groups (*pepz*) sharing the NGS, applying the Mann-

Whitney U test[25]. Testing $\%_{g,pepz,a}$ *vs.* $\%_{g,pepz,b}$ individually for *u*, *c* and *h* at each NGS, we

calculated *p*-values that were subjected to multiple hypothesis correction using the Benjamini-

Hochberg (BH) method[26]. If the corrected *p*-value was <0.05 (for *u*, *c* or *h* at any NGS), then the

difference was considered statistically significant.

For ***Supplementary Figure 2***, published data from our previous approach[3] was reanalyzed using

the data analysis workflow described here. Briefly, the data analyzed is from 3 replicates of 3

conditions, each separately analyzed by LC-MS/MS, with total starting material of 90 µg protein,

and Census2[23] label-free analysis was performed simultaneously on all 9 experiments without

"match between runs", and the results analyzed by GlycoMSQuant. This was compared with

data obtained from a single LC-MS/MS run (QE-HFX/nLC) with 0.5 µg of peptides generated

from a starting material of 5 µg purified protein.

Acknowledgements:

We thank Bruker Daltonics for providing access to timsTOF Pro MS. We thank Bruker

Scientific LLC and Robin S Park for making IP2 accessible for data analysis. We thank Titus

Jung in J.R.Y. lab for IT support. We thank members of the J.R.Y. lab for discussion and Claire

Delahunty for critically perusing the manuscript. We thank Saman Eskandarzadeh, Michael

Kubitz, and Erik Georgeson in W.R.S. lab for assistance in protein production. This work was

supported by the grants P41GM103533 (NIH) and UM1AI100663, UM1AI144462,

R01/R56AI113867 (NIH/NIAID). We thank Bill and Melinda Gates Foundation (BMGF)

Collaboration for AIDS Vaccine Discovery (CAVD) funding to IAVI Neutralizing Antibody

Center (NAC) and Ragon Institute post-doctoral fellowship to T.S.

Author Contributions:

326   S.B. conceived the method. S.B., J.K.D. and X.W. designed the experiments. S.B. prepared the

327   samples and analyzed data. J.K.D. performed LC-MS/MS. S.M.B. created the GlycoMSQuant

328   tool. T.S. and B.G. expressed and purified HIV *Env* trimers. W.R.S., J.C.P. and J.R.Y.

329   supervised the project. S.B. wrote the paper with contribution from all authors.

Competing Interests:

332   The authors declare no competing interests.

Data Availability:

335   Mass spectrometry data has been deposited in MassIVE-KB repository and is also accessible

336   through ProteomeXchange Consortium with identifiers MSV000087414 and PXD025990,

337   respectively.

Code Availability:

GlycoMSQuant source code is freely available at

https://github.com/proteomicsyates/GlycoMSQuant under a permissive Apache License 2.0.

References:

1.   Rudd, P.M. & Dwek, R.A. Glycosylation: heterogeneity and the 3D structure of proteins.

Crit Rev Biochem Mol Biol **32**, 1-100 (1997).

2.   Burton, D.R. & Hangartner, L. Broadly Neutralizing Antibodies to HIV and Their Role in

Vaccine Design. *Annu Rev Immunol* **34**, 635-659 (2016).

3.   Cao, L. et al. Global site-specific N-glycosylation analysis of HIV envelope glycoprotein.

Nat Commun **8**, 14954 (2017).

4.   MacCoss, M.J. et al. Shotgun identification of protein modifications from protein

complexes and lens tissue. *Proc Natl Acad Sci U S A* **99**, 7900-7905 (2002).

5.   Zhang, Y., Fonslow, B.R., Shan, B., Baek, M.C. & Yates, J.R., 3rd Protein analysis by

shotgun/bottom-up proteomics. *Chem Rev* **113**, 2343-2394 (2013).

6.   Tsiatsiani, L. & Heck, A.J. Proteomics beyond trypsin. *Febs j* **282**, 2612-2626 (2015).

7.   Behrens, A.J. et al. Molecular Architecture of the Cleavage-Dependent Mannose Patch

on a Soluble HIV-1 Envelope Glycoprotein Trimer. *J Virol* **91** (2017).

8.   Go, E.P. et al. Glycosylation Benchmark Profile for HIV-1 Envelope Glycoprotein

Production Based on Eleven Env Trimers. *J Virol* **91** (2017).

9.   Wu, C.C., MacCoss, M.J., Howell, K.E. & Yates, J.R., 3rd A method for the

comprehensive proteomic analysis of membrane proteins. *Nat Biotechnol* **21**, 532-538

(2003).

10.   Steichen, J.M. et al. HIV Vaccine Design to Target Germline Precursors of Glycan-

Dependent Broadly Neutralizing Antibodies. *Immunity* **45**, 483-496 (2016).

11.   Steichen, J.M. et al. A generalized HIV vaccine design strategy for priming of broadly

366   neutralizing antibody responses. *Science* **366** (2019).

367   12.   Sanders, R.W. et al. A next-generation cleaved, soluble HIV-1 Env trimer, BG505

368   SOSIP.664 gp140, expresses multiple epitopes for broadly neutralizing but not non-

369   neutralizing antibodies. *PLoS Pathog* **9**, e1003618 (2013).

370   13.   Meier, F. et al. Online Parallel Accumulation-Serial Fragmentation (PASEF) with a

371   Novel Trapped Ion Mobility Mass Spectrometer. *Mol Cell Proteomics* **17**, 2534-2545

372   (2018).

373   14.   Seabright, G.E., Doores, K.J., Burton, D.R. & Crispin, M. Protein and Glycan Mimicry in

374   HIV Vaccine Design. *J Mol Biol* **431**, 2223-2247 (2019).

375   15.   Zhou, H.X., Rivas, G. & Minton, A.P. Macromolecular crowding and confinement:

376   biochemical, biophysical, and potential physiological consequences. *Annu Rev Biophys*

377   **37**, 375-397 (2008).

378   16.   Bar-On, Y.M., Flamholz, A., Phillips, R. & Milo, R. SARS-CoV-2 (COVID-19) by the

379   numbers. *Elife* **9** (2020).

380   17.   Scanlan, C.N. et al. Inhibition of mammalian glycan biosynthesis produces non-self

381   antigens for a broadly neutralising, HIV-1 specific antibody. *J Mol Biol* **372**, 16-22

382   (2007).

383   18.   Fan, J.Q. & Lee, Y.C. Detailed studies on substrate structure requirements of

384   glycoamidases A and F. *J Biol Chem* **272**, 27058-27064 (1997).

385   19.   He, L., Diedrich, J., Chu, Y.Y. & Yates, J.R., 3rd Extracting Accurate Precursor

386   Information for Tandem Mass Spectra by RawConverter. *Anal Chem* **87**, 11361-11367

387   (2015).

20.   Xu, T. et al. ProLuCID: An improved SEQUEST-like algorithm with enhanced

sensitivity and specificity. *J Proteomics* **129**, 16-24 (2015).

21.   Tabb, D.L., McDonald, W.H. & Yates, J.R., 3rd DTASelect and Contrast: tools for

assembling and comparing protein identifications from shotgun proteomics. *J Proteome*

Res **1**, 21-26 (2002).

22.   Peng, J., Elias, J.E., Thoreen, C.C., Licklider, L.J. & Gygi, S.P. Evaluation of

multidimensional chromatography coupled with tandem mass spectrometry (LC/LC-

MS/MS) for large-scale protein analysis: the yeast proteome. *J Proteome Res* **2**, 43-50

(2003).

23.   Park, S.K., Venable, J.D., Xu, T. & Yates, J.R., 3rd A quantitative analysis software tool

for mass spectrometry-based proteomics. *Nat Methods* **5**, 319-322 (2008).

24.   Zhang, M. et al. Tracking global patterns of N-linked glycosylation site variation in

highly variable viral glycoproteins: HIV, SIV, and HCV envelopes and influenza

hemagglutinin. *Glycobiology* **14**, 1229-1246 (2004).

25.   Mann, H.B. & Whitney, D.R. On a Test of Whether one of Two Random Variables is

Stochastically Larger than the Other. *The Annals of Mathematical Statistics* **18**, 50-60

(1947).

26.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B*

(Methodological) **57**, 289-300 (1995).

Figures

Figure 2 | Factors affecting limit of sensitivity. (a) Decreasing trend of sequence and NGS coverage observed as starting material is diluted 200-fold (1 µg to 5 ng, QE-HFX/nLC) or 100-fold (0.5 µg to 5 ng, timsTOF/Evosep), using triplicates for each amount of starting material. The limit of sensitivity is revealed. **(b)** Relative abundance of peptides identified per NGS across the dilution series for the 2 LC-MS/MS platforms used. This comparison reveals a non-uniform digestion pattern that may be attributed to steric hindrance offered by the glycoprotein or characteristic behavior of individual peptides in LC-MS/MS. Error bars represent mean±SEM. Values for sequence and NGS coverage are mean