Gastrin producing syngeneic mesenchymal stem cells protect non-obese diabetic mice from type 1 diabetes

Marie-Claude Gaudreau[1*], Radhika R. Gudi[1*], Gongbo Li[2], Benjamin M. Johnson, and Chenthamarakshan Vasu[1,2,3]

[1]Department of Microbiology and Immunology, College of Medicine, Medical University of South Carolina, Charleston, SC-29425

[2]Department of Surgery, University of Illinois at Chicago, Chicago, IL-60612

Running title: MSC therapy for type 1 diabetes

[3]Address Correspondence: Chenthamarakshan Vasu
Medical University of South Carolina
Microbiology and Immunology
173 Ashley Avenue, MSC 509, BSB208E
Charleston, SC-29425
Phone: 843-792-1032, Fax: 843-792-9588
Email: vasu@musc.edu

Key words: Mesenchymal stem cells; Type 1 diabetes, Gastrin, Autoimmunity, insulitis, Immune modulation

*Equal contribution

Abstract:

Progressive destruction of pancreatic islet β-cells by immune cells is the primary feature of type 1 diabetes (T1D) and therapies that can restore the functional β-cell mass are needed to alleviate disease progression. Here, we report the use of mesenchymal stromal/stem cells (MSCs) for the production and delivery of Gastrin, a peptide-hormone which is produced by intestinal cells and fetal islets and can increase β-Cell mass, to promote protection from T1D. A single injection of syngeneic MSCs that were engineered to express Gastrin (Gastrin-MSCs) caused a profound delay in hyperglycemia in non-obese diabetic (NOD) mice compared to engineered control-MSCs. Similar treatment of early-hyperglycemic mice caused restoration of euglycemia for a significant duration, and these therapeutic effects were associated with protection of, and/or increase in, insulin producing islets and less severe insulitis. While the overall immune cell phenotype was not affected profoundly, pancreatic lymph node cells from Gastrin-MSC treated mice, upon ex vivo challenge with self-antigen, showed a Th2 and Th17 bias, and diminished the diabetogenic property in NOD-*Rag1* deficient mice suggesting a disease protective immune modulation upon Gastrin-MSC treatment. Overall, this study shows the potential of production and delivery of Gastrin in vivo, by MSCs, in protecting insulin producing β-cells and ameliorating the disease progression in T1D.

Introduction

Progressive destruction of pancreatic islet β-cells by immune cells, which ultimately results in the inability of pancreas to produce sufficient amounts of insulin, is the key event responsible for T1D[1,2]. The ideal goals of clinical intervention strategies for T1D are: 1) to prevent or arrest the onset and progression of autoimmunity before hyperglycemia is detected in at-risk individuals and 2) to restore insulin producing β-cell mass and glyco-metabolic and immune homeostasis after clinical diagnosis in T1D patients. Although immuno- and cellular- therapies are widely being tested for their efficacy in preventing/delaying T1D in pre-clinical models and clinical trials[3-10], there is an urgent need for safe and effective approaches to prevent the disease in at-risk subjects and restore insulin independence in T1D patients.

Recently, there has been a major focus on cellular therapy to generate β-cells for T1D treatment[11-13]. In addition, the immune-modulatory properties of some types of stem cells[14-17] can be exploited to suppress autoimmunity. Mesenchymal stromal/stem cells (MSCs), which can be isolated from a number of sources including the bone marrow (BM) and adipose tissue, are adult stem cells with a potential for differentiation into various tissues and they have the capacity for self-renewal and differentiation with a broad tissue distribution, and the ability to repair tissue damage[18-22]. Natural ability of MSCs to migrate in the body has been described[23-25]. Of particular interest for tissue repair, i.v. injection of MSCs results in their specific migration to a site of injury or inflammation[26-29]. Notably, most MSC based approaches for T1D diabetes therapy have been focused on their multi-potency to become insulin-producing cells (IPCs) or immune modulatory properties[30-35]. However, a major caveat is that MSCs from pre-clinical models as well as T1D patients show gross defects in many of these properties and lack the ability to promote protection from T1D[34,36], raising the concern that unaltered autologous MSCs may not have therapeutic value. Studies in NOD mice have shown that repeated injections with allogeneic or xenogeneic MSCs, but not syngeneic MSCs, promote protection from T1D[15,34,37].

One potential solution for overcoming the inherent defects of autologous/syngeneic MSCs is by engineering them to express desired factors that can protect/preserve β-cell function. Here, we examined the potential of syngeneic MSCs that are engineered to express Gastrin, a peptide hormone that is produced by intestinal cells as well as fetal islet hormone and has the ability to

stimulate β-Cell neogenesis and increase islet mass[38-41], to prevent T1D incidence and reverse the early-stage hyperglycemia in NOD mice. We show that, while non-engineered and control engineered MSCs had no substantial impact on T1D incidence in NOD mice, a single injection of Gastrin-MSCs 1) at pre-diabetic stage resulted in prevention of hyperglycemia and 2) at early-diabetic stages reversed hyperglycemia, for significant durations. We also show that Gastrin-MSC recipients had profoundly higher frequencies of insulin-producing islets compared to control-MSC recipients, suggesting the ability of Gastrin-MSCs to preserve/suppress the loss of functional β-cell mass. While Gastrin-MSC treatment had no profound impact on global Treg, IL10+ and IFNγ+ T cell frequencies, PnLN cells from these mice showed significantly altered cytokine responses upon challenge with β-cell antigen ex vivo. Overall, this study shows the potential of using autologous MSCs for in vivo production and delivery of Gastrin in protecting and/or increasing insulin producing β-cell mass to prevent and/or suppress the progression of T1D.

Materials and methods

Mice

Wild-type NOD/ShiLtJ (NOD) and NOD-*Rag1-/-* mice, originally purchased from the Jackson laboratory (Maine, USA), were bred in our facility. All animal studies were approved by the animal care and use committee of MUSC. To detect hyperglycemia in mice, glucose levels in the blood collected from tail vein were determined at timely intervals using the Ascensia Micro-fill blood glucose test strips and an Ascensia Contour blood glucose meter (Bayer, USA). Mice with glucose level of >250 mg/dl for two consecutive testing (at weekly time points) were considered diabetic/hyperglycemic. Untreated mice with glucose levels <100 mg/dl, 140-250 mg/dl, >450 mg/dl were considered pre-diabetic, early-hyperglycemic and overt-hyperglycemic respectively. In studies using mice at various stages of hyperglycemia, "high" glucose reading by Contour glucose meter for 2 consecutive testing were considered the endpoint.

Peptide antigens and other reagents

Immunodominant β-cell antigen peptides [viz., 1. Insulin B$_{(9-23)}$, 2. GAD65$_{(206-220)}$, 3. GAD65$_{(524-543)}$, 4. IA-2beta$_{(755-777)}$ and 5. IGRP$_{(123-145)}$ were custom synthesized (Genescript Inc) and used in this study. These peptides were pooled at an equal molar ratio and used as β-cell-Ag peptide

cocktail as described in our earlier studies[42-46]. PMA, ionomycin, Brefeldin A, ELISA and unlabeled and fluorochrome labeled antibodies, and other key reagents were purchased from Sigma-Aldrich, BD Biosciences, eBioscience, Invitrogen, Millipore, Miltenyi Biotec, StemCell Technologies, R&D Systems, Biolegend, and Santa-Cruz Biotechnology. Magnetic bead based 26-plex suspension bead array cytokine kits were purchased from Invitrogen. Suspension bead array plates were read using FlexMap3D instrument. Gastrin EIA kit was purchased from Sigma-Millipore. Flow cytometry data was acquired using FACS Calibur, FACS Verse or CyAn-ADP instruments and analyzed by using Flowjo application.

Cloning and Lentiviral production

The third-generation replication-incompetent lentiviral cDNA cloning vector (pCDH1-copGFP) and packaging plasmid were purchased from SBI and used as described in our previous report[47]. For cloning Gastrin cDNA, pCDH1-copGFP vector was digested using NheI (downstream of CMV promoter) and NotI restriction enzymes, and the restriction digestion fragment was replaced with PCR amplified and NheI and NotI digested mouse Gastrin cDNA. mRNA isolated from mouse stomach tissue was used to generate the Gastrin cDNA. For control (GFP) and Gastrin lentiviral production, HEK293T packaging cells were transfected with above control and Gastrin-cDNA vectors along with packaging vectors by calcium phosphate method. Occasionally, GPRG cells[48], provided by National Gene Vector Biorepository, were used as virus packaging cells. Virus-containing media were collected after 36 and 72 h, pooled, and centrifuged at 3500 RPM for 15 min and the supernatants were subjected to 0.22-µm filtration, and concentrated at least 25-fold by PEG precipitation. The infectious titers/transduction unit (TU) of the virus were determined by using fresh HEK293T cells.

Culture, lentiviral transduction and characterization of bone marrow (BM) MSCs

BM cells were isolated from the femurs and tibiae, red blood cells were removed by hypotonic lysis, and washed in complete DMEM/F12 medium containing 10% FBS, antibiotics/antimycotics, L-glutamine, sodium pyruvate, non-essential amino acids. The cells were seeded in cell culture treated dishes at $2x10^6$ cell/ml. Non-adherent cells were removed after 48h and 72h of cultures, trypsin-treated adherent cells were split every 3rd day starting day

5, and 13-16 day old cultures were used for lentiviral transduction and phenotypic characterization. For transducing the MSCs to obtain >80% efficiency, cells were seeded along with a final concentration of at least 5×10^8 TU/ml of virus in the presence of polybrene (8 μg/ml) and protamine sulfate (10 μg/ml) for up to 24 h, washed and cultured in fresh medium for 48h, split at 1:2 ratio and cultured for an additional 48h. Transduction efficiency and cell surface marker expression profile were determined after 72h of transduction by microscopy and/or flow cytometry. Gastrin secretion was determined in post-transduction supernatants collected at 72h time-points by using Gastrin EIA kit from Millipore-Sigma.

Treatment of NOD mice with engineered MSCs (eMSCs)

Ten-to-12-week-old pre-diabetic and 12-25-week-old early-diabetic, diabetic, overt-diabetic mice were injected i.v. with non-engineered MSCs, control (GFP)-MSCs or Gastrin-MSCs (approx. 1×10^6 cells/mouse). These mice were monitored for hyperglycemia by testing for blood glucose levels every week. Cohorts of mice were euthanized at different time-points to determine the degree of Gastrin production in the pancreas, insulitis, immune cell phenotype and auto-antigen specific response.

Adoptive T cell transfer experiment

Fresh PnLN cells from control- and Gastrin-MSC treated mice were transferred into 6-week-old NOD-*Rag1-/-* mice (i.v.; 1×10^6 cells/mouse) and tested for blood glucose levels every week.

Histochemical and immunofluorescence analysis of pancreatic tissues

Pancreata were fixed in 10% formaldehyde, 5-μm paraffin sections were made, and stained with hematoxylin and eosin (H&E) to assess insulitis. Stained sections were analyzed using a 0-4 insulitis severity grading system as described in our earlier studies[45,49,50]. In some experiments, pancreatic sections were stained using anti-insulin and/or anti-glucagon antibodies, followed by Alexa fluor 488- and 568- linked secondary antibodies and DAPI, and scored for insulitis based on DAPI-positive cells in islet areas and insulin positivity. In some experiments, insulin positive islets among total islet structures in different groups were compared.

Statistical analysis

Mean, SD, and statistical significance (*p-value*) were calculated using GraphPad Prism, Microsoft Excel, and/or online statistical applications. Unpaired *t*-test or Mann-Whitney test was

employed, unless specified, for values from *in vitro* and *ex vivo* assays. Fisher's exact test was used for comparing the total number of severely infiltrated islets (grades ≥3) relative to total number of islets with low or no infiltration (grades ≤2) in test vs. control groups. Log-rank analysis (http://bioinf.wehi.edu.au/software/russell/logrank/) was performed to compare T1D incidence (hyperglycemia) of the test group with that of respective control group. A *P* value of ≤0.05 was considered statistically significant.

Results

Engineering of MSCs to express Gastrin and phenotypic characterization of MSCs

For generating Gastrin-expressing MSCs, tissue culture plate-adherent BM cells from NOD mice were cultured for up to 16-days with splitting at least every 3rd day. Cells with MSC phenotype from 13-16-day old culture were used for various experiments. The cells were transduced with control- or Gastrin- lentivirus particles, both with GFP reporter, **(Supplemental Fig. 1)** and cultured for 72h before use in various assays. Transduction efficiency, indicated by GFP expression, was confirmed by fluorescence microscopy **(Fig. 1A)**. In some experiments, secreted levels of Gastrin in control- and Gastrin-lentivirus transduced/engineered MSC (eMSC) culture wells were examined using EIA kit **(Fig. 1B)**. Examination of cell surface expression of MSC related markers confirmed that control and Gastrin transduced cells maintained their MSC-phenotype after lentiviral transduction **(Fig. 1C)**. These results show that BM MSCs from NOD mice can be engineered effectively to express Gastrin, without affecting their phenotype.

Gastrin production is detected in pancreatic tissues of Gastrin-MSC treated mice

To examine if systemically injected eMSCs secrete Gastrin in the pancreatic microenvironment, early diabetic NOD mice (with blood glucose: 140-250 mg/dl) were injected i.v. with control- and Gastrin-MSCs (1x10^6 cells/mouse). Cohorts of mice were euthanized after 48h or 30-day time-points, and pancreatic tissues were digested using collagenase followed by trypsin. Washed cell suspension of each pancreas was cultured in 2ml complete DMEM/F12 medium for 48h and the spent media were tested for Gastrin levels by EIA. As shown in **Fig. 2**, profoundly higher levels of Gastrin were detected in the pancreatic suspension cultures of mice that received

Gastrin-MSCs compared to those of control-MSC recipients. Interestingly, Gastrin was detectable, albeit at much lower levels, in the pancreas even after 30-days of treatment. These results suggest that eMSCs reach the pancreatic microenvironment, persist, and secrete the factors of interest for a considerable duration.

Gastrin-MSC treatment at pre-diabetic stage results in diminished insulitis and prevention of T1D in NOD mice

To determine the effect of treatment with Gastrin-MSCs on insulitis and T1D incidence, pre-diabetic female NOD mice were injected once with non-engineered MSCs, control-MSCs or Gastrin-MSCs, and the degree of insulitis was determined after 30 days. H&E staining of pancreatic sections revealed that while mice that received non-engineered MSCs or control-MSCs did not show significant differences in the insulitis severity compared to untreated controls, mice that received Gastrin-MSCs had significantly less severe insulitis compared to control groups (**Fig. 3A**). To assess the impact of suppressed insulitis in Gastrin-MSC recipients on diabetes incidence, cohorts of treated mice were monitored for hyperglycemia for up to 25-weeks post-treatment. As shown in **Fig. 3B**, while 100% of Gastrin-MSC treated mice remained euglycemic for at least 10 weeks post-treatment, 90-100% of untreated, and non-engineered MSC and control-MSC treated mice turned hyperglycemic during this period. Moreover, about 50% Gastrin-MSC recipients remained euglycemic at the end of 25 weeks of monitoring. Importantly, diabetes onset was detected in the control groups of mice within 2-5 weeks of monitoring as compared to 12 weeks post-treatment in Gastrin-MSC treated mice. Overall, these results show that a single dose of Gastrin-MSCs can cause significant suppression of insulitis and prevention of hyperglycemia for a significant duration.

Gastrin-MSC treatment at pre-diabetic stage results in preservation of insulin producing islets

To determine the effect of treatment using Gastrin-MSCs on the abundance of functional islets, insulin positive islet frequencies of mice that received MSC treatment at pre-diabetic stage were determined 30 days after the treatment. Cryosections of pancreatic tissues of control-MSC and Gastrin-MSC treated mice were stained using anti-insulin antibody and the insulitis severity and total number of islet areas with and without insulin positive cells was assessed. Islet features

were then graded depending on immune cell infiltration (based on DAPI) and insulin staining. As shown in **Fig. 4**., while majority of the islets (>90%), including those with profound immune cell infiltration, in Gastrin-MSC treated mice were insulin positive, >60% pancreatic islets from control groups of mice were insulin negative. Interestingly, insulin positive as well as insulitis free islet frequencies of Gastrin-MSC treated mice appear to be considerably higher (>20%) than that of control-MSC recipients, which is <5%. This result suggests that prevention of T1D in Gastrin-MSC treated mice is associated with protection/preservation of overall insulin producing islet mass.

Treatment of NOD mice with Gastrin-MSCs at early-hyperglycemic and diabetic stages results in reversal of hyperglycemia

Next, we examined if blood glucose levels and insulin producing islet abundance in mice with early-hyperglycemia and established diabetes can be altered by treating with Gastrin-MSCs. Female NOD mice with blood glucose between 140-250 mg/dl (early-hyperglycemic) and >250 (diabetic) mg/dl were injected once with control-MSCs or Gastrin-MSCs and monitored for blood glucose levels. As shown in **Fig. 5A**, while all control-MSC treated early-hyperglycemic mice (n=6) progressed to diabetic stage, all 4 early-hyperglycemic mice that received Gastrin-MSCs reverted to euglycemic (blood glucose <100 mg/dl) within 2 weeks and remained euglycemic for at least 8 weeks post-treatment. In the case of diabetic mice, Gastrin-MSC treatment caused reversal of hyperglycemia in mice that received the treatment when the blood glucose levels were at about 400 mg/dl or less, but not in those at overt-hyperglycemic stage. On the other hand, control-MSC recipient diabetic mice failed to revert to euglycemic. Of note, a separate experiment using only overt diabetic mice (glucose levels: >500 mg/dl; n=6 mice/group) showed no therapeutic effect either with control-MSCs or Gastrin-MSCs (not shown).

Pancreatic tissues were collected, at 30-days post-treatment, from cohorts of mice that were treated with control-MSCs and Gastrin-MSCs at early-hyperglycemic stage to examine insulitis and islet function. Pancreatic tissues from naïve, early-hyperglycemic (at the time of euthanasia) control mice were also collected and processed. Tissue sections were stained using anti-insulin and anti-glucagon antibodies and examined for the frequencies of "functional islets". As shown

in **Fig. 5B**, about 80% of islet areas, with or without immune cell infiltration, of Gastrin-MSC recipient mice showed some degree of islet function, as indicated by insulin and glucagon positivity, compared to about 2% insulin positive islet remnants in control-MSC treated mice. Of note, functional islet areas were detectable only in mice that received treatment at the early diabetic stages, but not at overt-hyperglycemic stage (not shown). These results show that reversal of hyperglycemia is possible by Gastrin-MSC treatment, if initiated soon after detection of clinical stage disease.

Gastrin-MSC treated mice showed suppressed autoimmune response

While MSCs in general are thought to have immune modulatory properties[14-17], only the allogeneic or xenogeneic MSCs, but not the syngeneic MSCs showed immune modulatory properties and the ability to protect NOD mice from T1D upon repeated injections[14,15,34,37,51]. In fact, as shown in **Supplemental Fig. 2**, a single injection of syngeneic, non-engineered MSCs or control-eMSCs had no impact on T1D progression in NOD mice. On the other hand, a single injection of non-engineered allogeneic MSCs caused modest delay in T1D onset. Nevertheless, since T1D protection was achieved in NOD mice when treated with Gastrin-MSCs, we examined if the syngeneic control-MSC and Gastrin-MSC treatment recipient mice show differences in immune response. As observed in **Fig. 6A**, albeit modestly higher frequencies of IL10+ and IL17+ cells in Gastrin-MSC treated mice, differences in the frequencies of various T cell populations, including Foxp3+ cells, in the pancreatic LN of control- and Gastrin-MSC treated mice were not statistically significant. Similar observations were made when spleen cells were examined (not shown). To determine the pathogenic effect of immune cells, PnLN cells from control- and Gastrin-MSC treated mice were injected into NOD-*Rag1-/-* mice and monitored for hyperglycemia. As observed in **Fig. 6B**, recipients of PnLN cells from both control-MSC and Gastrin-MSC treated donors turned diabetic by 14 weeks of cell transfer. However, mice that received cells from Gastrin-MSC treated mice showed significantly delayed onset of hyperglycemia suggesting that the diabetogenic properties of immune cells are diminished in Gastrin-MSC treated mice. We, then, examined if self-antigen specific T cells are impacted by the treatment. PnLN cells from MSC-treated mice were cultured ex vivo in the presence of β-cell antigen peptide cocktail[42-46], and the profile of secreted cytokines was examined by employing suspension bead array. **Fig. 6C** shows that cells from Gastrin-MSC

treated mice, compared to that of control-MSC recipients, secreted significantly higher amounts of IL2, IL1β, IL4, IL5, IL13, IL17, IL22, GM-CSF and eotaxin, and lower amounts of IL6, MCP1 and MIP2 upon ex vivo activation using β-cell antigen. Interestingly, higher amounts of Th2 and Th17 response related cytokines (IL4, IL5, IL13, IL17 and IL22 particularly) were produced by immune cells from Gastrin-MSC recipients suggesting a skewed T cell response in these mice. Overall, these results show that while Gastrin-MSC treatment does not exert widespread immune modulation, self-antigen specific response and pathogenic properties of T cells in the pancreatic microenvironment are impacted as a consequence of this treatment, potentially contributing to protection from the disease.

Discussion

Gastrin, which is produced primarily by the stomach and duodenum cells, binds to cholecystokinin B receptor and stimulates the release of gastric acid from parietal cells[52,53]. This peptide hormone can also cause increased division and differentiation of cells in the acid-secreting mucosa [53-55]. Of interest, gastrin is also expressed in the pancreas at the embryonic stage as well as in adult pancreas under stress including hyperglycemia in type 2 diabetes (T2D) and cancer[38,39,56,57]. Transient high expression of Gastrin and its receptors in fetal pancreas, in a period of pronounced islet neogenesis[58,59], suggests its involvement in increasing islet and β-cell mass. In fact, multiple reports have shown that Gastrin can increase pancreatic β-cell mass[60-69]. Gastrin can also synergize with other agents such as GLP-1 and EGF and impact islet/β-cell function and regulate blood glucose[61,65,69-72]. The β-cell neogenesis inducing potential of Gastrin was demonstrated using pancreas specific transgenic mice as well as by Gastrin therapy[61,70]. While systemic use of Gastrin has been explored for enhancing β-cell mass [69,71,73], repeated high-dose systemic administration of this peptide hormone can be inefficient due to its short half-life. Further, its persistent high systemic levels can be unsafe due to global side effects including unregulated cell proliferation and malignancies. Hence, approaches for preferential delivery of Gastrin to, and its low-level continuous production in, the pancreatic microenvironment are needed to enhance β-cell mass safely and effectively. In this study, we demonstrate that the MSCs that are engineered to exogenously produce Gastrin could

be such delivery vehicles for protecting β-cell function to prevent, and potentially reverse, hyperglycemia in T1D.

It has been shown that MSCs can ameliorate tissue damage and improve function after lung injury, kidney disease, diabetes, myocardial infarction, liver injury, and neurological disorders[8,19,74,75]. Natural ability of MSCs to migrate in the body has also been described[23,24,76]. Of particular interest, for delivering desired protein factors to pancreas, i.v. injection of MSCs results in their specific migration to a site of injury or inflammation[23,25,77]. Therefore, the inherent property of MSCs to migrate to the site of inflammation makes them the ideal cell population for delivering protein factors to sites of inflammation. Engineering the MSCs exogenously and stably to express the protein factors of interest is expected to ensure prolonged delivery of factors such as Gastrin. Unlike other stem cells that are difficult to engineer with viral vectors without affecting their properties, MSCs can be readily transduced with most available viral vector systems to produce secreted, surface and cytoplasmic proteins of interest[78,79]. Importantly, MSCs have been engineered to either augment their own natural production of specific proteins or to enable them to express other therapeutic proteins for various clinical conditions[80-82]. Furthermore, eMSCs have been effectively used to enhance their ability to repair tissues including myocardium, CNS and liver in experimental models[83-85]. With respect to T1D therapy, eMSCs that express Pdx1 and other β-cell specific transcription factors have shown the ability to become insulin producing cells in vitro and in vivo[86-88]. While, how effective these MSC preparations are in differentiating into IPCs in an autoimmune microenvironment and if they can reverse established disease is largely unknown, these reports show the possibility that MSCs from T1D patients and pre-clinical models, despite their defective differentiation and immunomodulation properties[34,36], can be exploited as delivery vehicles for proteins of interest. Therefore, we postulated that if MSCs that are engineered to secrete Gastrin can, upon injection, reach and condition the pancreatic microenvironment, then these cells will be effective in increasing insulin producing cells and reversing hyperglycemia without being reliant on their multi-potent and immune modulatory properties. In fact, our observations that Gastrin production can be detectable, albeit at low levels, in the pancreatic tissue of Gastrin eMSC recipient NOD mice even after 30 days of injection, insulin producing islet frequencies are increased, diabetes can be prevented, and early hyperglycemia can be reversed do support this notion.

With respect to islet β-cell mass and insulin production in T1D, two key Gastrin associated effects have been reported. First, it has shown that Gastrin's mitogenic property not only triggers increased insulin production by residual β-cells, but it alone or along with other factors such as GLP-1, EGF, and TGFα also induces the proliferation of these cells[38-41,61,65,69-72]. Second, studies using pancreas specific Gastrin and TGFα transgenic mice, as well as that involving pancreatic ductal ligation, have shown that Gastrin can contribute to trans-differentiation of the exocrine acinar cells to duct-like cells as well as neogenesis of insulin producing islets from ductal cells[62,64,68,70]. Therefore, while the mechanism of Gastrin-MSC treatment induced protection of NOD mice from T1D in our study is not known, it is possible that higher insulin positive islets found in these mice as compared to control MSC-recipients are, at least in part, due to the direct effects on pancreatic tissues. Substantiating the likelihood of direct impact on pancreatic islets, we observed that many of the islet structures with severe insulitis in Gastrin-MSC treated mice, unlike in control MSC treated mice, were insulin positive. We also observed that the insulin positive islet frequencies of Gastrin-MSC recipients 30 days post-treatment was significantly higher than that of a cohort of pre-treatment stage mice.

Notably, syngeneic Gastrin-MSC treated NOD mice showed suppression of autoimmune and pro-inflammatory response to a certain degree. While previous report showing the need for repeated injections with syngeneic MSCs[34], and our observations that a single dose of syngeneic non-engineered or control-eMSCs had no impact on T1D incidence in NOD mice, suggest that the immune modulation observed in Gastrin-MSC recipients is treatment associated. A direct effect for Gastrin on immune cells is largely unknown. Our observations that immune function of Gastrin-MSC treated mice, compared to control-MSC treated mice, is skewed towards Th17 and Th2 types suggest that Gastrin does impact immune function, perhaps indirectly through causing changes in the pancreatic microenvironment. In this regard, our in vitro coculture assays using control and Gastrin-MSC and NOD spleen cells and purified immune cells showed comparable pro- and anti- inflammatory cytokine responses (not shown). It is possible that, as suggested in previous reports[89-92], hyperglycemia alone can impact various physiological functions including gut permeability and cause systemic pro-inflammatory response aggravating the disease complications. Hence, an explanation for the observed in vivo immune modulation in Gastrin-MSC recipients is that Gastrin induced enhanced β-cell function,

and the suppression of hyperglycemia, contributed to skewed immune cell function and diminishes pro-inflammatory and autoimmune responses.

Overall, our study suggests that use of autologous MSCs to deliver peptide hormone Gastrin in the pancreatic microenvironment can be an effective and safer approach for preventing T1D and reversing early-stage hyperglycemia. It is possible that repeated injections as well as combination treatment approaches involving other islet regeneration and immune modulatory factors could potentially enhance the efficacy of this approach for achieving permanent euglycemia, even in patients with clinical stage disease.

Conflict of Interest statement: The authors report no conflicts of interest.

Data availability: The datasets generated and/or analyzed during the current study are available from the corresponding author on reasonable request.

Resource availability: The cDNA vectors generated during the current study is available from the corresponding author on reasonable request.

Author contribution: M.G. researched and analyzed data and edited the manuscript, R.G. researched and analyzed data and edited the manuscript, G.L, researched data, B.J. researched data, and C.V. designed experiments, researched and analyzed data, and wrote/edited manuscript.

Acknowledgements: This work was supported by unrestricted research funds from MUSC and National Institutes of Health (NIH) grants R21AI069848, R21AI133798 and R01AI073858 to C.V. C.V. is the guarantor of this work and, as such, has full access to all the data in the study and takes responsibility for the integrity of the data and accuracy of the data analysis. The authors are thankful to Cell and Molecular Imaging, Pathology, Proteomics, immune monitoring and discovery, and flow cytometry cores of MUSC and UIC for the histology service, microscopy, FACS and multiplex assay instrumentation support.

References:

1. Lagani V, Koumakis L, Chiarugi F, et al. A systematic review of predictive risk models for diabetes complications based on large scale clinical studies. Journal of diabetes and its complications. 2013 Jul-Aug;27(4):407-13.
2. Best CH. Some thoughts on the etiology of human diabetes. Canadian Medical Association journal. 1962 Oct 6;87:731-4.
3. Giannopoulou EZ, Puff R, Beyerlein A, et al. Effect of a single autologous cord blood infusion on beta-cell and immune function in children with new onset type 1 diabetes: a non-randomized, controlled trial. Pediatr Diabetes. 2014 Mar;15(2):100-9.
4. Haller MJ, Wasserfall CH, Hulme MA, et al. Autologous umbilical cord blood infusion followed by oral docosahexaenoic acid and vitamin D supplementation for C-peptide preservation in children with Type 1 diabetes. Biology of blood and marrow transplantation : journal of the American Society for Blood and Marrow Transplantation. 2013 Jul;19(7):1126-9.
5. Seay HR, Putnam AL, Cserny J, et al. Expansion of Human Tregs from Cryopreserved Umbilical Cord Blood for GMP-Compliant Autologous Adoptive Cell Transfer Therapy. Mol Ther Methods Clin Dev. 2017 Mar 17;4:178-191.
6. Malmegrim KC, de Azevedo JT, Arruda LC, et al. Immunological Balance Is Associated with Clinical Outcome after Autologous Hematopoietic Stem Cell Transplantation in Type 1 Diabetes. Frontiers in immunology. 2017;8:167.
7. Diaz-de-Durana Y, Lau J, Knee D, et al. IL-2 immunotherapy reveals potential for innate beta cell regeneration in the non-obese diabetic mouse model of autoimmune diabetes. PloS one. 2013;8(10):e78483.
8. Manzoor F, Johnson MC, Li C, et al. beta-cell-specific IL-35 therapy suppresses ongoing autoimmune diabetes in NOD mice. Eur J Immunol. 2017 Jan;47(1):144-154.
9. Goudy KS, Tisch R. Immunotherapy for the prevention and treatment of type 1 diabetes. Int Rev Immunol. 2005 Sep-Dec;24(5-6):307-26.
10. Alhadj Ali M, Liu YF, Arif S, et al. Metabolic and immune effects of immunotherapy with proinsulin peptide in human new-onset type 1 diabetes. Sci Transl Med. 2017 Aug 9;9(402).
11. Cho J, D'Antuono M, Glicksman M, et al. A review of clinical trials: mesenchymal stem cell transplant therapy in type 1 and type 2 diabetes mellitus. Am J Stem Cells. 2018;7(4):82-93.
12. Sambathkumar R, Migliorini A, Nostro MC. Pluripotent Stem Cell-Derived Pancreatic Progenitors and beta-Like Cells for Type 1 Diabetes Treatment. Physiology (Bethesda). 2018 Nov 1;33(6):394-402.
13. Ma H, Wert KJ, Shvartsman D, et al. Establishment of human pluripotent stem cell-derived pancreatic beta-like cells in the mouse pancreas. Proceedings of the National Academy of Sciences of the United States of America. 2018 Apr 10;115(15):3924-3929.
14. Abdi R, Fiorina P, Adra CN, et al. Immunomodulation by mesenchymal stem cells: a potential therapeutic strategy for type 1 diabetes. Diabetes. 2008 Jul;57(7):1759-67.
15. Madec AM, Mallone R, Afonso G, et al. Mesenchymal stem cells protect NOD mice from diabetes by inducing regulatory T cells. Diabetologia. 2009 Jul;52(7):1391-9.
16. Glenn JD, Whartenby KA. Mesenchymal stem cells: Emerging mechanisms of immunomodulation and therapy. World journal of stem cells. 2014 Nov 26;6(5):526-39.

17.	Le Blanc K, Ringden O. Immunomodulation by mesenchymal stem cells and clinical experience. Journal of internal medicine. 2007 Nov;262(5):509-25.

18.	Prockop DJ. Marrow stromal cells as stem cells for nonhematopoietic tissues. Science. 1997 Apr 4;276(5309):71-4.

19.	Kucia M, Reca R, Jala VR, et al. Bone marrow as a home of heterogenous populations of nonhematopoietic stem cells. Leukemia. 2005 Jul;19(7):1118-27.

20.	Pittenger MF, Mackay AM, Beck SC, et al. Multilineage potential of adult human mesenchymal stem cells. Science. 1999 Apr 2;284(5411):143-7.

21.	Li M, Ikehara S. Bone-marrow-derived mesenchymal stem cells for organ repair. Stem cells international. 2013;2013:132642.

22.	Patel DM, Shah J, Srivastava AS. Therapeutic potential of mesenchymal stem cells in regenerative medicine. Stem cells international. 2013;2013:496218.

23.	Horwitz EM, Prockop DJ, Fitzpatrick LA, et al. Transplantability and therapeutic effects of bone marrow-derived mesenchymal cells in children with osteogenesis imperfecta. Nature medicine. 1999 Mar;5(3):309-13.

24.	Ji JF, He BP, Dheen ST, et al. Interactions of chemokines and chemokine receptors mediate the migration of mesenchymal stem cells to the impaired site in the brain after hypoglossal nerve injury. Stem cells. 2004;22(3):415-27.

25.	Sordi V, Malosio ML, Marchesi F, et al. Bone marrow mesenchymal stem cells express a restricted set of functionally active chemokine receptors capable of promoting migration to pancreatic islets. Blood. 2005 Jul 15;106(2):419-27.

26.	Wang B, Wu SM, Wang T, et al. Pre-treatment with bone marrow-derived mesenchymal stem cells inhibits systemic intravascular coagulation and attenuates organ dysfunction in lipopolysaccharide-induced disseminated intravascular coagulation rat model. Chinese medical journal. 2012 May;125(10):1753-9.

27.	Yu Y, Wu RX, Gao LN, et al. Stromal cell-derived factor-1-directed bone marrow mesenchymal stem cell migration in response to inflammatory and/or hypoxic stimuli. Cell adhesion & migration. 2016 Jul 3;10(4):342-59.

28.	Spaeth E, Klopp A, Dembinski J, et al. Inflammation and tumor microenvironments: defining the migratory itinerary of mesenchymal stem cells. Gene Ther. 2008 May;15(10):730-8.

29.	Kallmeyer K, Pepper MS. Homing properties of mesenchymal stromal cells. Expert opinion on biological therapy. 2015 Apr;15(4):477-9.

30.	Canibano-Hernandez A, Saenz Del Burgo L, Espona-Noguera A, et al. Hyaluronic Acid Promotes Differentiation of Mesenchymal Stem Cells from Different Sources toward Pancreatic Progenitors within Three-Dimensional Alginate Matrixes. Molecular pharmaceutics. 2019 Feb 4;16(2):834-845.

31.	Ojaghi M, Soleimanifar F, Kazemi A, et al. Electrospun poly-l-lactic acid/polyvinyl alcohol nanofibers improved insulin-producing cell differentiation potential of human adipose-derived mesenchymal stem cells. J Cell Biochem. 2018 Dec 11.

32.	Kharat A, Chandravanshi B, Gadre S, et al. IGF-1 and somatocrinin trigger islet differentiation in human amniotic membrane derived mesenchymal stem cells. Life Sci. 2019 Jan 1;216:287-294.

33.	Bassi EJ, Moraes-Vieira PM, Moreira-Sa CS, et al. Immune regulatory properties of allogeneic adipose-derived mesenchymal stem cells in the treatment of experimental autoimmune diabetes. Diabetes. 2012 Oct;61(10):2534-45.

34.   Fiorina P, Jurewicz M, Augello A, et al. Immunomodulatory function of bone marrow-derived mesenchymal stem cells in experimental autoimmune type 1 diabetes. Journal of immunology. 2009 Jul 15;183(2):993-1004.

35.   Jurewicz M, Yang S, Augello A, et al. Congenic mesenchymal stem cell therapy reverses hyperglycemia in experimental type 1 diabetes. Diabetes. 2010 Dec;59(12):3139-47.

36.   de Lima KA, de Oliveira GL, Yaochite JN, et al. Transcriptional profiling reveals intrinsic mRNA alterations in multipotent mesenchymal stromal cells isolated from bone marrow of newly-diagnosed type 1 diabetes patients. Stem cell research & therapy. 2016 Jul 12;7(1):92.

37.   Dang LT, Bui AN, Le-Thanh Nguyen C, et al. Intravenous Infusion of Human Adipose Tissue-Derived Mesenchymal Stem Cells to Treat Type 1 Diabetic Mellitus in Mice: An Evaluation of Grafted Cell Doses. Adv Exp Med Biol. 2018;1083:145-156.

38.   Suissa Y, Magenheim J, Stolovich-Rain M, et al. Gastrin: a distinct fate of neurogenin3 positive progenitor cells in the embryonic pancreas. PloS one. 2013;8(8):e70397.

39.   Dahan T, Ziv O, Horwitz E, et al. Pancreatic beta-Cells Express the Fetal Islet Hormone Gastrin in Rodent and Human Diabetes. Diabetes. 2017 Feb;66(2):426-436.

40.   Khan D, Vasu S, Moffett RC, et al. Expression of Gastrin Family Peptides in Pancreatic Islets and Their Role in beta-Cell Function and Survival. Pancreas. 2018 Feb;47(2):190-199.

41.   Rooman I, Lardon J, Flamez D, et al. Mitogenic effect of gastrin and expression of gastrin receptors in duct-like cells of rat pancreas. Gastroenterology. 2001 Oct;121(4):940-9.

42.   Perez N, Karumuthil-Melethil S, Li R, et al. Preferential costimulation by CD80 results in IL-10-dependent TGF-beta1(+) -adaptive regulatory T cell generation [Research Support, N.I.H., Extramural

Research Support, Non-U.S. Gov't]. Journal of immunology. 2008 May 15;180(10):6566-76.

43.   Li R, Perez N, Karumuthil-Melethil S, et al. Bone marrow is a preferential homing site for autoreactive T-cells in type 1 diabetes. Diabetes. 2007 Sep;56(9):2251-9.

44.   Karumuthil-Melethil S, Perez N, Li R, et al. Dendritic cell-directed CTLA-4 engagement during pancreatic beta cell antigen presentation delays type 1 diabetes. Journal of immunology. 2010 Jun 15;184(12):6695-708.

45.   Karumuthil-Melethil S, Sofi MH, Gudi R, et al. TLR2- and Dectin 1-associated innate immune response modulates T-cell response to pancreatic beta-cell antigen and prevents type 1 diabetes. Diabetes. 2015 Apr;64(4):1341-57.

46.   Karumuthil-Melethil S, Gudi R, Johnson BM, et al. Fungal beta-glucan, a Dectin-1 ligand, promotes protection from type 1 diabetes by inducing regulatory innate immune response. Journal of immunology. 2014 Oct 1;193(7):3308-21.

47.   Gudi RR, Karumuthil-Melethil S, Perez N, et al. Engineered Dendritic Cell-Directed Concurrent Activation of Multiple T cell Inhibitory Pathways Induces Robust Immune Tolerance. Sci Rep. 2019 Aug 19;9(1):12065.

48.   Throm RE, Ouma AA, Zhou S, et al. Efficient construction of producer cell lines for a SIN lentiviral vector for SCID-X1 gene therapy by concatemeric array transfection. Blood. 2009 May 21;113(21):5104-10.

49.   Sofi MH, Gudi R, Karumuthil-Melethil S, et al. pH of drinking water influences the composition of gut microbiome and type 1 diabetes incidence. Diabetes. 2014 Feb;63(2):632-44.

50.   Karumuthil-Melethil S, Perez N, Li R, et al. Induction of innate immune response through TLR2 and dectin 1 prevents type 1 diabetes. Journal of immunology. 2008 Dec 15;181(12):8323-34.

51.   Hu J, Wang Y, Wang F, et al. Effect and mechanisms of human Wharton's jelly-derived mesenchymal stem cells on type 1 diabetes in NOD model. Endocrine. 2015 Feb;48(1):124-34.

52.   Shulkes A, Baldwin GS. Biology of gut cholecystokinin and gastrin receptors. Clin Exp Pharmacol Physiol. 1997 Mar-Apr;24(3-4):209-16.

53.   Dockray GJ, Varro A, Dimaline R, et al. The gastrins: their production and biological activities. Annu Rev Physiol. 2001;63:119-39.

54.   Baldwin GS. The role of gastrin and cholecystokinin in normal and neoplastic gastrointestinal growth. Journal of gastroenterology and hepatology. 1995 Mar-Apr;10(2):215-32.

55.   Logan CJ, Connell AM. The effect of a synthetic gastrin-like pentapeptide (I.C.I. 50,123) on intestinal motility in man. Lancet. 1966 May 7;1(7445):996-9.

56.   Gittes GK, Rutter WJ, Debas HT. Initiation of gastrin expression during the development of the mouse pancreas. Am J Surg. 1993 Jan;165(1):23-5; discussion 25-6.

57.   Smith JP, Fonkoua LK, Moody TW. The Role of Gastrin and CCK Receptors in Pancreatic Cancer and other Malignancies. Int J Biol Sci. 2016;12(3):283-91.

58.   Larsson LI, Rehfeld JF, Sundler F, et al. Pancreatic gastrin in foetal and neonatal rats. Nature. 1976 Aug 12;262(5569):609-10.

59.   Brand SJ, Fuller PJ. Differential gastrin gene expression in rat gastrointestinal tract and pancreas during neonatal development. The Journal of biological chemistry. 1988 Apr 15;263(11):5341-7.

60.   Brand SJ, Tagerud S, Lambert P, et al. Pharmacological treatment of chronic diabetes by stimulating pancreatic beta-cell regeneration with systemic co-administration of EGF and gastrin. Pharmacology & toxicology. 2002 Dec;91(6):414-20.

61.   Suarez-Pinzon WL, Yan Y, Power R, et al. Combination therapy with epidermal growth factor and gastrin increases beta-cell mass and reverses hyperglycemia in diabetic NOD mice. Diabetes. 2005 Sep;54(9):2596-601.

62.   Rooman I, Lardon J, Bouwens L. Gastrin stimulates beta-cell neogenesis and increases islet mass from transdifferentiated but not from normal exocrine pancreas tissue. Diabetes. 2002 Mar;51(3):686-90.

63.   Song I, Patel O, Himpe E, et al. Beta Cell Mass Restoration in Alloxan-Diabetic Mice Treated with EGF and Gastrin. PloS one. 2015;10(10):e0140148.

64.   Tellez N, Montanya E. Gastrin induces ductal cell dedifferentiation and beta-cell neogenesis after 90% pancreatectomy. The Journal of endocrinology. 2014 Oct;223(1):67-78.

65.   Sasaki S, Miyatsuka T, Matsuoka TA, et al. Activation of GLP-1 and gastrin signalling induces in vivo reprogramming of pancreatic exocrine cells into beta cells in mice. Diabetologia. 2015 Nov;58(11):2582-91.

66.   Skarbaliene J, Secher T, Jelsing J, et al. The anti-diabetic effects of GLP-1-gastrin dual agonist ZP3022 in ZDF rats. Peptides. 2015 Jul;69:47-55.

67.   Tellez N, Vilaseca M, Marti Y, et al. beta-cell dedifferentiation, reduced duct cell plasticity and impaired beta-cell mass regeneration in middle-aged rats. Am J Physiol Endocrinol Metab. 2016 Jul 12:ajpendo 00502 2015.

68. Tellez N, Joanny G, Escoriza J, et al. Gastrin treatment stimulates beta-cell regeneration and improves glucose tolerance in 95% pancreatectomized rats. Endocrinology. 2011 Jul;152(7):2580-8.

69. Suarez-Pinzon WL, Lakey JR, Rabinovitch A. Combination therapy with glucagon-like peptide-1 and gastrin induces beta-cell neogenesis from pancreatic duct cells in human islets transplanted in immunodeficient diabetic mice. Cell transplantation. 2008;17(6):631-40.

70. Wang TC, Bonner-Weir S, Oates PS, et al. Pancreatic gastrin stimulates islet differentiation of transforming growth factor alpha-induced ductular precursor cells. The Journal of clinical investigation. 1993 Sep;92(3):1349-56.

71. Suarez-Pinzon WL, Power RF, Yan Y, et al. Combination therapy with glucagon-like peptide-1 and gastrin restores normoglycemia in diabetic NOD mice. Diabetes. 2008 Dec;57(12):3281-8.

72. Fosgerau K, Jessen L, Lind Tolborg J, et al. The novel GLP-1-gastrin dual agonist, ZP3022, increases beta-cell mass and prevents diabetes in db/db mice. Diabetes, obesity & metabolism. 2013 Jan;15(1):62-71.

73. Caballero F, Siniakowicz K, Hollister-Lock J, et al. Birth and death of human beta-cells in pancreases from cadaver donors, autopsies, surgical specimens, and islets transplanted into mice. Cell transplantation. 2014 Feb;23(2):139-51.

74. Krause DS. BM-derived stem cells for the treatment of nonhematopoietic diseases. Cytotherapy. 2002;4(6):503-6.

75. Prockop DJ. Marrow stromal cells as stem cells for continual renewal of nonhematopoietic tissues and as potential vectors for gene therapy. Journal of cellular biochemistry Supplement. 1998;30-31:284-5.

76. Ratajczak J, Wysoczynski M, Zuba-Surma E, et al. Adult murine bone marrow-derived very small embryonic-like stem cells differentiate into the hematopoietic lineage after coculture over OP9 stromal cells. Experimental hematology. 2011 Feb;39(2):225-37.

77. Barbash IM, Chouraqui P, Baron J, et al. Systemic delivery of bone marrow-derived mesenchymal stem cells to the infarcted myocardium: feasibility, cell migration, and body distribution. Circulation. 2003 Aug 19;108(7):863-8.

78. Meyerrose TE, De Ugarte DA, Hofling AA, et al. In vivo distribution of human adipose-derived mesenchymal stem cells in novel xenotransplantation models. Stem cells. 2007 Jan;25(1):220-7.

79. Zhang XY, La Russa VF, Bao L, et al. Lentiviral vectors for sustained transgene expression in human bone marrow-derived stromal cells. Molecular therapy : the journal of the American Society of Gene Therapy. 2002 May;5(5 Pt 1):555-65.

80. Reiser J, Zhang XY, Hemenway CS, et al. Potential of mesenchymal stem cells in gene therapy approaches for inherited and acquired diseases. Expert opinion on biological therapy. 2005 Dec;5(12):1571-84.

81. Bosch P, Stice SL. Adenoviral transduction of mesenchymal stem cells. Methods in molecular biology. 2007;407:265-74.

82. Fan L, Lin C, Zhuo S, et al. Transplantation with survivin-engineered mesenchymal stem cells results in better prognosis in a rat model of myocardial infarction. European journal of heart failure. 2009 Nov;11(11):1023-30.

83. Nomura T, Honmou O, Harada K, et al. I.V. infusion of brain-derived neurotrophic factor gene-modified human mesenchymal stem cells protects against injury in a cerebral ischemia model in adult rat. Neuroscience. 2005;136(1):161-9.

84. Kurozumi K, Nakamura K, Tamiya T, et al. BDNF gene-modified mesenchymal stem cells promote functional recovery and reduce infarct size in the rat middle cerebral artery occlusion model. Molecular therapy : the journal of the American Society of Gene Therapy. 2004 Feb;9(2):189-97.

85. Zhao MZ, Nonoguchi N, Ikeda N, et al. Novel therapeutic strategy for stroke in rats by bone marrow stromal cells and ex vivo HGF gene transfer with HSV-1 vector. Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism. 2006 Sep;26(9):1176-88.

86. Guo QS, Zhu MY, Wang L, et al. Combined transfection of the three transcriptional factors, PDX-1, NeuroD1, and MafA, causes differentiation of bone marrow mesenchymal stem cells into insulin-producing cells. Experimental diabetes research. 2012;2012:672013.

87. Lin HT, Kao CL, Lee KH, et al. Enhancement of insulin-producing cell differentiation from embryonic stem cells using pax4-nucleofection method. World journal of gastroenterology : WJG. 2007 Mar 21;13(11):1672-9.

88. Soyer J, Flasse L, Raffelsberger W, et al. Rfx6 is an Ngn3-dependent winged helix transcription factor required for pancreatic islet cell development. Development. 2010 Jan;137(2):203-12.

89. Thaiss CA, Levy M, Grosheva I, et al. Hyperglycemia drives intestinal barrier dysfunction and risk for enteric infection. Science. 2018 Mar 23;359(6382):1376-1383.

90. Nielsen TB, Pantapalangkoor P, Yan J, et al. Diabetes Exacerbates Infection via Hyperinflammation by Signaling through TLR4 and RAGE. MBio. 2017 Aug 22;8(4).

91. Gyurko R, Siqueira CC, Caldon N, et al. Chronic hyperglycemia predisposes to exaggerated inflammatory response and leukocyte dysfunction in Akita mice. Journal of immunology. 2006 Nov 15;177(10):7250-6.

92. Esposito K, Nappo F, Marfella R, et al. Inflammatory cytokine concentrations are acutely increased by hyperglycemia in humans: role of oxidative stress. Circulation. 2002 Oct 15;106(16):2067-72.

Figure Legends

Figure 1: Ectopic expression of Gastrin and phenotypic characterization of eMSCs. BM MSCs from NOD mice were transduced with control vector (GFP) or Gastrin lentiviral particles for 72h as described under materials and methods. **A)** Representative microscopy images of control-MSCs and Gastrin-MSCs are shown. **B)** Supernatants were subjected to competitive EIA and values of supernatants from three independently transduced MSC preparations are shown. *p*-value by paired *t*-test. **C)** eMSCs were stained for MSC markers (CD106, CD105, CD44, CD73, Sca-1 and CD29), and hematopoietic and myeloid cell markers (CD45 and CD11b) and analyzed by FACS. FACS plots and mean fluorescent intensity (MFI) values, representative of three independent assays, are shown.

Figure 2: Gastrin production by pancreatic cell suspension of Gastrin-MSC treated mice. Early diabetic female NOD mice (with blood glucose: 140-250 mg/dl) were left untreated (none), or injected with Control MSCs or Gastrin MSCs, i.v. (1×10^6 cells/mouse). Pancreatic tissues were harvested from cohorts of mice on day 2 or 30 post-treatment, single cell suspensions were cultured for 48h, and the supernatants were subjected to EIA to determine Gastrin levels. Values of 4 mice/group, each tested separately in triplicate, are shown. *p*-value by Mann-Whitney test.

Figure 3: Impact of Gastrin MSC treatment on T1D incidence and insulitis. Ten- to 12-week-old pre-diabetic female NOD mice were left untreated (none) or injected with Control-MSCs or Gastrin-MSCs as described for Fig. 2. A) Cohorts of mice (a total of 6 mice/group) were euthanized 30 days post-treatment, pancreatic sections were subjected to H&E staining and grading for insulitis severity. Examples of islets with different insulitis grades (left panel) and the percentage islets with different insulitis grades (right panel) are shown. Grades: 0 = no evidence of infiltration, 1 = peri-islet infiltration (<5%), 2= 5-25% islet infiltration, 3 = 25–50% islet infiltration, and 4 = >50% islet infiltration. At least 140 islets from intermittent sections were examined for each group. Statistical significance was assessed by Fisher's exact test comparing relative numbers of islets with insulitis grades ≤2 and ≥3 between groups. **B)** Cohorts of mice were monitored for hyperglycemia by testing for blood glucose levels every week. Mice that showed glucose >250 mg/dl for two consecutive weeks were considered diabetic, and cumulative results of multiple experiments, using 3-4 mice/group in each experiment (total of 14 to 15

mice/group), are shown. *p*-values of control-MSC vs Gastrin-MSC (0 to 15 week and 0 to 25 week) comparisons by log-rank test are shown.

Figure 4: Impact of Gastrin-MSC treatment on insulin positive islet frequencies. Ten- to 12-week-old pre-diabetic female NOD mice treated with control-MSCs or Gastrin-MSCs as described for Fig. 2, euthanized after 30 days post-treatment, and pancreatic tissue sections were stained using anti-insulin antibody (red) and the nuclear stain DAPI (blue) and examined for insulitis and insulin positive islets. **Left panel:** representative microscopy images indicating mostly insulin negative islet areas (red arrow) in control-MSC recipients and insulin positive islets (white arrow) in Gastrin-MSC treated mice are shown. **Right panel:** Islet features were graded 0-5 depending on immune cell infiltration (based on DAPI) and insulin staining (+ or -): 0+ (no infiltration/insulin+), 1+ (<5% infiltration/insulin+), 2+ (5-25% infiltration/insulin+), 3+ (25-50% infiltration/insulin+), 4+ (50-100% infiltration/insulin+), 4- (50-100% infiltration/insulin-), and 5- (only islet remnants left/insulin-), and percentage values of islets with specific grades among examined islet areas are shown. About 100 islets from multiple intermittent sections were examined for each group (total of 6 mice/group). *p*-value by Fisher's exact test comparing relative numbers of islets with insulitis grades ≤2+ and ≥3+ between control-MSC recipients vs Gastrin-MSC recipients.

Figure 5: Impact of Gastrin-MSC treatment at hyperglycemic stages on blood glucose levels and insulin producing islet frequencies. Hyperglycemic group of female NOD mice (which included early-hyperglycemic and diabetic mice) were treated with control-MSCs or Gastrin-MSCs as described for Fig. 2. **A)** Blood glucose levels were examined before and every week after the treatment, and glucose values of individual mice are shown. Dotted lines represent "diabetic" and "high (endpoint)" glucose levels. *p*-values by Fisher's exact test comparing the number of mice with diabetic glucose levels by week 5 post-treatment as well as by log-rank test comparing survival duration (euthanasia timing) of individual mice in each group. **B)** Cohorts of MSC treated early-hyperglycemic mice were euthanized 30 days post treatment, and pancreatic sections were subjected to insulin (green), glucagon (red) and DAPI (blue) staining and insulin positive islets and immune cell infiltration were scored as described for Fig. 4B.*Pancreatic tissues from a cohort of untreated mice that were at early-hyperglycemic stage at the time of euthanasia were included as "none" control group to assess the pre-treatment frequency of insulin positive islets. *p*-value by Fisher's exact test comparing relative numbers of islets with

insulitis grades ≤2+ and ≥3+ between control-MSC recipients vs Gastrin-MSC recipients and none group vs Gastrin-MSC recipients.

FIGURE 6: Impact of Gastrin-MSC treatment on immune cell phenotype and function. Prediabetic female NOD mice were treated with control-MSCs or Gastrin-MSCs as described for Fig. 2 and euthanized after 30 days to obtain PnLN cells. **A)** PnLN cells were stained for Foxp3, or for intracellular cytokines after brief stimulation using PMA and ionomycin in the presence of Brefeldin A, and analyzed by FACS. Representative FACS plots (left panel) and mean±SD values of 5 mice/group tested individually in duplicate (right panel) are shown. This experiment was repeated using similar number of mice. **B)** PnLN cells from at least 4 mice/group were pooled separately and injected i.v. into NOD-*Rag1-/-* mice ($1x10^6$ cells/mouse; 8 recipients/group) and examined for blood glucose levels every week to detect hyperglycemia. *p*-value by log-rank test. **C)** PnLN cells (4 mice/group) from individual mice were cultured overnight in the presence of β-cell Ag peptide cocktail for 48h and the supernatants were tested for cytokine levels by suspension bead array and the cytokine concentration values of individual mice were used for generating heatmap. *p*-value by Mann-Whitney test. This assay was repeated once using an additional 3 mice/group with comparable outcomes.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

p=0.007 (based on % diabetic mice at 5wk)
p=0.006 (based on euthanasia timing)

Fig. 6