Threonine phosphorylation regulates the molecular assembly and signaling of EGFR in cooperation with membrane lipids

Ryo Maeda[1], Hiroko Tamagaki-Asahina[2], Takeshi Sato[2], Masataka Yanagawa[1], and Yasushi Sako[1,3,*]

[1]Cellular Informatics Laboratory, RIKEN CPR, Wako, Saitama 351-0198, Japan; [2]Kyoto Pharmaceutical University, 5, Misasagi-cho, Yamashina, Kyoto, 607-8414, Japan; [3]CREST JST, 4-1-8, Honcho, Kawaguchi, 332-0012, Japan

*To whom correspondence should be addressed. E-mail: sako@riken.jp

Abstract

The cytoplasmic domain of the receptor tyrosine kinases (RTKs) plays roles as a phosphorylation enzyme and a protein scaffold but the regulation of these two functions is not fully understood. We here analyzed assembly of the transmembrane (TM)-juxtamembrane (JM) region of EGFR, one of the best studied species of RTKs, by combining single-pair FRET imaging and a nanodisc technique. The JM domain of EGFR contains a threonine residue that is phosphorylated after ligand association. We observed that the TM-JM peptides of EGFR form anionic lipid-induced dimers and cholesterol-induced oligomers. The two forms involve distinct molecular interactions, with a bias towards oligomer formation upon threonine phosphorylation. We further analyzed the functions of whole EGFR molecules, with or without a threonine to alanine substitution in the JM domain, in living cells. The results suggested an autoregulatory mechanism in which threonine phosphorylation of the JM domain causes a switch from kinase activation dimers to scaffolding oligomers.

Introduction

Epidermal growth factor receptor (EGFR) is an RTK responsible for cell proliferation and differentiation (1, 2) and consists of five domains; an extracellular domain that interacts with extracellular ligands, a single-pass transmembrane (TM) helix, a juxtamembrane (JM) domain, a cytoplasmic kinase domain, and a C-terminal tail domain for interaction with various cytoplasmic proteins (3, 4). Ligand association changes the conformation of EGFR in its extracellular domain (5) and induces formation of an asymmetric dimer of the intracellular kinase domains (6). This dimerization subsequently results in the phosphorylation of tyrosine residues on the tail domain and the recruitment of intracellular signal proteins such as GRB2 and PLCγ containing SH2 and/or PTB domains (7). Although the atomic structures of most of the EGFR domains excluding the tail domain have been elucidated individually (5, 6, 8-10), the overall architecture of this protein has not yet been revealed, leaving several unanswered questions about the molecular mechanisms underlying its functions. The correlation between the arrangement of EGFR molecules and their function is therefore still controversial, e.g., it has long been established that the dimerization of EGFR is necessary and sufficient for kinase activation (11), whereas several studies have reported the importance of higher-order oligomerization for EGFR-mediated signal transduction (12-14).

The TM helix and the JM domain (TM-JM) of EGFR play important roles in the conformational coupling of ligand binding to its activation and oligomerization (9, 11, 15). Previous NMR studies and molecular dynamics simulations have suggested that the TM domain forms an α-helix dimer that undergoes a change in the interaction between GXXXG motifs (9, 16, 17) following the ligand association with its extracellular domains (17, 18). This information regarding conformational changes in the TM dimer is then transmitted to the JM domain which comprises a JM-A (N-terminal half) region that can form an antiparallel helix dimer, and a JM-B (C-terminal half) region which makes intramolecular contact with the kinase domain (11). Both these JM regions contribute to the stable formation of an asymmetric kinase dimer conformation, which is crucial for kinase activation. The JM-A domain is rich in Lys and Arg residues, several of which are thought to interact with anionic lipid molecules of the plasma membrane and promote antiparallel dimer formation (19-21). In addition to the phospholipid species, cholesterol is a major component of the plasma membrane, mainly distributed as lipid rafts and caveolae, and has been implicated in the regulation of membrane fluidity and receptor function. Previous studies have shown that EGFR molecules are clustered in lipid rafts (14, 22), suggesting an interaction with cholesterols. Of note in this regard, it has been reported that the depletion of cholesterols induces various effects on EGFR signaling,

also this remains controversial (23-25). Another important factor in the regulation of
EGFR through the TM-JM is the phosphorylation of Thr654 at the JM-A domain.
Although Thr phosphorylation is known to be involved in EGFR deactivation, the precise
mechanism of this is still elusive (26).
In our present study, by combining single-pair FRET measurements and
nanodisc technology, we studied how the functions of anionic lipids, cholesterols, and
EGFR Thr654 phosphorylation (pT654) are orchestrated to achieve the regulation of
dimerization and/or oligomerization of EGFR. We previously reported that anionic lipids
cause the dimerization of JM domains, and that pT654 together with acidic lipids induces
the dissociation of the EGFR dimer (20). In this current study, we report that both the TM
and JM protomers of EGFR are positioned closer to each other in the presence of
cholesterols than in the EGFR dimers promoted by anionic lipids. Furthermore, we found
that TM-JM peptides were oligomerized in cholesterol containing membranes, which was
promoted by pT654. Finally, in living cells expressing whole EGFR molecules, we
observed differential functional roles of this crucial signaling factor that are dependent on
the pT654 levels. Our results suggest that the membrane cholesterol and pT654
cooperatively shift the EGFR function from the kinase dimer to the scaffold oligomer.
83

Results

Incorporation of TM-JM peptides into nanodiscs

Synthesized peptides of the TM-JM region of EGFR were prepared and labeled with a fluorophore Cy3 or Cy5 at the N-terminus (TM terminal region) or C-terminus (JM terminal region), respectively (Fig. 1a). The peptides were reconstituted into nanodisc structures with membrane scaffold proteins (MSPs) and lipid molecules (Fig. 1b, c). Mixtures of POPC (PC), POPS (PS), and cholesterol were used for reconstruction (Fig. 1d). The nanodiscs containing cholesterol showed two peaks following size exclusion gel chromatography, one of which had a smaller disc size relative to that without cholesterol (Fig. 1e). To avoid the effects of disc size, we collected and used nanodiscs involved in the first peak fraction which had a similar size without cholesterol. This fraction was composed of the same ratio of lipids as used in the preparation (Fig. 1f). Synthesized TM-JM peptides with pT654 were also reconstituted into nanodiscs. The nanodisc construction was examined under a transmission electron microscope (Fig. 1g). In total, 16 types of nanodiscs were applied to subsequent single-molecule measurements. Fractions of PS (inner leaflet) and cholesterol (inner and outer leaflets) mimicked those in the plasma membrane.

TM-TM interaction in the EGFR dimer

Nanodiscs containing Cy3 and Cy5-labeled peptides were immobilized onto glass surfaces and illuminated with a 532-nm laser for Cy3 excitation. A portion of the fluorescent spots contained Cy5 fluorescence derived from the occurrence of FRET (Fig. 2a, b). Based on the fluorescence intensity, we selected nanodiscs containing one Cy3- and one Cy5-labeled peptide and calculated the FRET efficiency, E_{FRET} (Fig. 2c).

We first examined the interactions between the N-terminal regions of the TM domains (Fig. 3). When Thr654 was not phosphorylated and the membrane contained only PC as lipid species, E_{FRET} distributed with a peak at a relatively high (> 0.90; Table I) value (Fig. 3a), indicating close proximity of the two TM domains. There may be additional stable structures between the TM domains, as suggested by the small peaks and shoulders in the E_{FRET} distribution. The addition of anionic lipid PS caused few effects, i.e., the TM dimers were maintained as the major structure (Fig. 3b). Peptides with pT654 slightly decreased the major peak positions of the E_{FRET} distributions in the PC or PC/PS membranes (Fig. 3e, f). The smooth distribution of the pT654 peptides suggested that pThr654 had homogenized possible substructures of the TM dimers of non-phosphorylated peptides. PS had a little effect on the TM-TM interactions regardless of the Thr654 phosphorylation. The presence of cholesterol in the membrane concentrated

the distributions to a high E_{FRET} (> 0.95) region (Fig. 3c, g, h), indicating that the N-terminal regions of TM-TM dimers were positioned in extremely close proximity. Although the peak position was similar in PC and PC/cholesterol membranes (Fig 3a, c; Table I), the distribution in PC/cholesterol membrane was more accumulated at the peak position. It also should be noted that the accumulation at a high E_{FRET} region was a remarkable observation for pT654 peptides. Thus, pT654 and the presence of membrane cholesterol decreased the distance between the two N-termini of the TM domains in cooperation. PS competed with cholesterol when Thr654 was non-phosphorylated (Fig. 3d).

JM-JM interaction in the EGFR dimer

To examine the effects of lipid species and pT654 on JM-JM interaction, E_{FRET} distributions were determined under the C-terminus labeling (Fig. 4). In PC membrane, E_{FRET} was broadly distributed with a peak around 0.81 (Fig. 4a; Table I). It is plausible that the JM-A dimers are fluctuating between minor dissociation and major association states. In the PC/PS membrane, the peak fraction was increased, indicating that PS stabilized the JM-A dimer conformation (Fig. 4b). pThr654 increased the low FRET fraction in the PC/PS membrane (Fig. 4f) but showed little effect in the PC membrane (Fig. 4e). These results confirmed the results of our previous study (20). Cholesterol moved the E_{FRET} peak between non-phosphorylated peptides to higher values (0.93–0.94) regardless of whether it was a PC or PC/PS membrane (Fig. 4c, d), i.e., cholesterol forced the C-termini of the JM-A domains to position closer.

The mixed effects of cholesterol and pT654 on the JM-JM interaction were further examined. In the PC membrane (Fig. 4g), cholesterol increased the high FRET population to a comparable level to those shown for non-phosphorylated peptides. Cholesterol in the PC/PS membrane (Fig. 4h) reversed the E_{FRET} distribution seen in the PC/PS membrane without cholesterol (Fig. 4f) to that observed in the PC membrane (Fig. 4e) i.e. minor low FRET and major high FRET states in the PC/PS/cholesterol membrane. However, it should be noted that the E_{FRET} values were not as large as those found in other conditions with cholesterol (Fig. 4c, d, g), i.e., the cholesterol effect on JM-JM interaction was partially diminished by the coexistence of PS and pThr654. Overall, our data showed that cholesterol increased the proximity between the C-terminus of JM domains in PC and PC/PS membranes, and that this effect overrode that of pThr654 in the PC/PS membrane.

Higher-order oligomerization of TM-JM peptides

We speculated that the accumulation of EGFR in lipid rafts, which has been reported in
previous studies, could be an effect of cholesterol in the raft membrane. We examined the
assembly of TM-JM peptides in the nanodiscs, collecting images of fluorescent spots
containing only Cy3-labeled peptides to avoid interference from the effects of FRET
occurring between Cy3 and Cy5. Figure 5 displays the fluorescence intensity histograms
of C-terminus-labeled TM-JM peptides in nanodiscs containing or not-containing
cholesterol. Cholesterol shifted the histograms toward higher intensities for the pT654
peptides (Fig. 5b, d), suggesting a cooperative effect of cholesterol and Thr
phosphorylation to induce higher-order assembly of the TM-JM peptides.

Assembly of TM regions

For analysis of the interactions between more than two TM or JM domains in the
assembled structures at the N-terminus, images of fluorescent spots containing two Cy3-
labeld peptides and one Cy5-labeled peptide were collected based on their 2-color
fluorescence trajectories (Fig. 6a). These nanodiscs showed a variety of Cy3 donor
fluorescence intensities before Cy5 photobleaching indicating that the three peptides
interacted diversely. We constructed maximum fluorescence intensity histograms in the
Cy3 donor channel before and after Cy5 acceptor photobleaching for inference of the
interactions between three TM domains (Fig. 6b–i). In all conditions other than non-
phosphorylated peptides in the PC/PS membrane, Cy3 distributions after Cy5
photobleaching (red) peaked at the fluorescence intensity of ~100 (in arbitrary units),
which was smaller than that observed for the C-terminal-labeled peptides (~150; Fig. 7b–
i). This result must have been caused by homo-FRET (self-quenching) between two N-
terminal labeled Cy3 peptides. Together with the very small intensity peaks prior to Cy5
photobleaching (blue), these distributions suggested that TM regions of the three peptides
(two of them were randomly labeled with Cy3) were oligomerized in very close proximity
to each other in the major configuration (trimer; Fig. 6j).

For non-phosphorylated peptides in the PC/PS membrane however (Fig. 6c), the
Cy3 intensity histogram after Cy5 photobleaching (red) had a peak intensity at ~150,
indicating that two Cy3-labeled peptides in the major population were positioned
separately. In addition, the low intensity shoulder in this distribution indicated the
presence of proximate dimers (and trimers). Taken together, these distributions suggested
that N-terminal regions of three non-phosphorylated peptides have a stronger tendency to
arrange as one dimer and one monomer in the PC/PS membrane than any other condition.
A similar dimer + monomer arrangement might be contained in the distributions under
other conditions as a minor fraction. Consistent with this suggestion, for the non-

phosphorylated peptides in PC/PS membrane before Cy5 photobleaching (Fig. 6c, blue),
a homo-FRET fraction (~100; with a low E_{FRET} to Cy5) was evident compared to other
conditions. It should be noted that the ability of PS to promote the dimer + monomer
arrangement was diminished for pT654 peptides (Fig. 6g). Cholesterol also reduced this
effect of PS even for non-phosphorylated peptides (Fig. 6e). As observed in the earlier
analysis of the TM-JM peptide dimer (Fig. 3), the effects of PS and cholesterol were
competitive.
199
Assembly of JM regions
We constructed Cy3 fluorescence intensity histograms of two Cy3 and one Cy5 peptide
with C-terminal-labeling in single nanodiscs in order to analyze the interactions between
three JM domains (Fig. 7). The distributions of the Cy3 florescence after Cy5
photobleaching (red) were similar under all conditions, exhibiting a single peak at ~150,
which was the fluorescence intensity of the two Cy3 molecules without strong
interactions to induce homo-FRET. Both the trimer and dimer + monomer arrangements
are possible if we assume that the three molecules in the trimer and two molecules in the
dimer are not so close that they will induce homo FRET (Fig. 7j).
Prior to Cy5 photobleaching (blue), the distribution peaks were observed in the
region of small Cy3 intensities indicating the proximity of both Cy3 molecules with Cy5
to induce high E_{FRET}, as observed between two molecules in a nanodisc (Fig. 4), i.e., the
formation of a JM trimer. An accumulation in the low intensity peak fraction was very
evident for non-phosphorylated peptides in the membranes containing cholesterol (Fig.
7d, e). On the other hand, fractions at the intensities similar to those observed after Cy5
photobleaching were significant for pT654 peptides in the membrane without cholesterol
(Fig. 7f, g). In general, pT654 peptides exhibited higher fluorescence intensity compared
to non-phosphorylated peptides in the corresponding membrane lipid compositions. One
possible explanation is that the fraction of high Cy3 intensity before Cy5 photobleaching
represents a Cy3 dimer in the dimer + monomer arrangement of three peptides (Fig. 7j).
Another possibility is that it was caused by an increased distance between three JM
domains in trimers, resulting from Thr phosphorylation to reduce E_{FRET} (Fig. 7j top
middle). These two arrangements could potentially coexist.
Considering the possible arrangement for the TM and JM regions of three TM-
JM peptides together (Figs. 6 and 7), we conclude that cholesterol induces the closely
proximate oligomerization of three JM domains of non-phosphorylated peptides whereas
PS preferentially causes a dimer + monomer arrangement, and the Thr phosphorylation
disrupts the JM dimer and facilitates oligomerization of peptides with separated JM

domains in the presence of cholesterols (Fig. 8). Because of the limitation of FRET
measurement, we could not distinguish whether the peptides in the oligomers directly
interacted or not.
231

Effect of Thr phosphorylation on the Tyr phosphorylation of EGFR

Our single-molecule structural analysis suggested that pT654 is a key regulator of the
molecular assembly of EGFR, which may affect its functions. We examined this
possibility in living cells. It is known that PKC activation under EGF signaling induces
pT654 in EGFR. This process has been thought to be a negative feedback pathway in the
EGFR system. We expressed a wild type (wt) or T654A mutant EGFR in CHO-K1 cells,
which have no intrinsic expression of EGFR. An increase in pT654 was observed for wt
EGFR after EGF stimulation, and treatment of these cells with phorbol-12-myristate 13-
acetate (PMA), a PKC activator, caused stronger phosphorylation of Thr654 regardless
of EGF stimulation (Fig. 9a). Application of a saturation amount (100 ng/ml) of EGF to
the culture medium induced phosphorylation of Tyr1068 (pY1068) of both the wt and
T654A mutant EGFR proteins (Fig. 9b). pY1068 is a major association site on EGFR for
the adaptor protein GRB2 and its levels after EGF stimulation were significantly
increased by the T654A mutation compared to wt, as expected from the negative effect of
pT654. Whereas, pretreatment with PMA decreased the pY1068 level in both the wt and
T654A mutant EGFR (Fig. 9c). The PMA-induced decrease in pY1068 for the wt protein
could be a negative effect of increased pT654, but the similar result with the T654A
mutant suggests that PMA has indirect effects independent of pT654.
250

Single-molecule imaging of the clustering and movement of EGFR

We expressed EGFR (wt and T654A) fused with GFP in CHO-K1 cells and, by using
single-molecule imaging, detected cluster size distributions and lateral diffusion
movements of EGFR molecules in the plasma membrane (Fig. 10a). Clustering of EGFR
was measured as the fluorescence intensity distribution of EGFR spots, and the lateral
diffusion movements were measured as the increase in the mean square displacement
(MSD) of the spots with time. Both measurements were performed before and after 10
min of EGF application to the medium. Application of EGF to the medium induced
clustering and immobilization of wt EGFR as we have reported previously(14, 27). The
distributions of EGFR cluster size suggest formation of oligomers containing up to more
than 10 molecules (Fig. 10b). The convex shapes of MSD curve with time indicate
subdiffusion of EGFR molecules (Fig. 10c).
Even in the absence of EGF, PMA treatment of cells increased fractions of

higher-order wt EGFR oligomers (Fig. 10b left), though diffusion movements were hardly
affected (Fig. 10c left). This oligomerization was not as strong as that induced by EGF in
the absence of PMA, and application of EGF to the PMA treated cells did not induce
further oligomerization at least up to 10 min but significantly decreased mobility.
Application of EGF induced strong oligomerization and immobilized wt EGFR in cells
without PMA treatment. For T654A mutant, PMA treatment hardly affected both
oligomerization (Fig. 10b right) and movements (Fig. 10c right) in the absence of EGF.
EGF induced strong oligomerization and immobilization of T654A mutant independent
of PMA treatment. The effect of PMA to induce EGFR oligomerization dependent on
Thr654 was consistent with pT654-induced oligomerization of TM-JM peptides in
nanodiscs.

In summary, single-molecule measurements suggest three states of EGFR
oligomerization depends on pT654 and EGF association (Fig. 10d). PMA treatment of wt
EGFR but not T654A mutant induced a medial level of oligomerization, which could be
stabilized by pT654. Strong oligomerization observed under the weak pT654 level in wt
EGFR and no pT654 in T654A mutant was caused by a distinct mechanism of pT654. On
the other hand, it is possible that immobilization of EGFR relates with its tyrosine
phosphorylation levels as we have observed previously (28). Immobilization in the
presence of PMA was more evident in T654A mutant in which pY1068 level was higher
than that in wt EGFR (Fig. 9), and PMA treatment decreased immobilization and pY1068
in wt EGFR.
285

Interaction of EGFR with GRB2

We finally measured the interaction of EGFR with GRB2 in living cells using a split
luciferase (NanoBiT) assay, in which the C-terminus of EGFR and the N-terminus of
GRB2 were conjugated with the large BiT (LgBiT) and the small BiT (SmBiT) of
NanoLuc luciferase (29), respectively. The association of EGFR and GRB2 promoted the
formation of active luciferase to produce chemiluminescence emission (Fig. 11a). From
the timecourses of the NanoBiT signal increases after EGF treatment of cells expressing
the wt or T654A mutant EGFR (Fig. 11b), the maximum intensities indicated a dose
dependent response to the EGF concentration in the medium (Fig. 11b, c). The maximum
intensity was significantly increased after pretreatment with PMA in cells expressing wt
EGFR, despite the fact that the pY1068 level was reduced after the PMA treatment (Fig.
9c). This effect of PMA was not observed to any extent with the T654A mutant, and the
GRB2 association was not increased by this mutation in the absence of PMA, even though
the pY1068 level after EGF stimulation was significantly increased in the mutant (Fig.

9c). The increase in GRB2 association observed for wt EGFR after PMA treatment was not likely to be an indirect effect of PMA because it was not observed for the T654A mutant, in which the pY1068 level was also affected by PMA. These results suggest that pT654 promotes the formation of a GRB2 recognition state for EGFR, and that the inhibition of Thr654 phosphorylation prevents a GRB2-EGFR association in spite of enhanced Tyr1068 phosphorylation.

Discussion

We have here studied the dimerization and oligomerization of EGFR molecules by reconstituting its TM-JM peptides into nanodiscs. As expected from the positively charged JM-A sequence and accumulation of EGFR in the raft membrane, PS and cholesterol affect the molecular assembly of the TM-JM peptides. Interestingly, these two lipid species each function in a specific fashion i.e. the E_{FRET} distributions between the two TM-JM peptides in the nanodiscs suggested that PS facilitates JM dimerization, while cholesterol induces closer positioning of both the TM and JM domains (Figs. 3 and 4). In addition, cholesterol promoted the oligomeric assembly of TM-JM peptides (Figs. 6 and 7). We herein propose schematic models for the formation of the EGFR TM-JM dimers and trimers under various conditions of lipid exposure and Thr654 phosphorylation (Fig. 8), in which PS and cholesterol exert competitive effects on dimerization and oligomerization, and pT654 disrupts the PS-induced JM dimer, thus promoting oligomerization of the peptides.

It should be noted that the E_{FRET} distribution was broad in every one of our observations in this present study, especially between JM domains, indicating multiple configurations coexisting under each condition. Non-phosphorylated peptide dimers showed a peak at around E_{FRET} > 0.90 and > 0.81 both for N- and C-terminus labeling, respectively. We can attribute this configuration to that suggested in a previous NMR study, in which two JM domains form an anti-parallel helix dimer (9). PS stabilized this configuration probably at the JM side of the non-phosphorylated peptide. Acidic lipids are known to interact with the positively charged JM-A domain (19, 30) whereas cholesterol induced more proximation of the two peptides at the both N- and C-termini in the major configuration (E_{FRET} 0.95 and 0.93), which must be distinct from the arrangement containing antiparallel JM helices (Fig. 8). If the TM-JM domains of whole EGFR dimer adopt similar configurations as suggested for the TM-JM peptides in the nanodiscs, the arrangement of two kinase domains indicates namely, the kinase activity would be affected by the lipid composition and by pT654 (Fig. 11d).

Cholesterol was found in our current analysis to increase the population of nanodiscs containing three TM-JM peptides with pT654 (Fig. 5). The fluorescence intensity distributions of the two Cy3 probes among the three peptides in the presence of cholesterol suggested that a close trimer was the major configuration (Figs. 6 and 7). The cholesterol-induced oligomerization of TM peptides with a short JM region (to T654) of EGFR in liposomes has been reported previously from NMR analysis (31). In that report however, pT654 in the peptide showed no obvious effect on the oligomerization in PC and PC/cholesterol liposomes (without any acidic lipids), consistent with our current and

previous results indicating interplay of acidic lipids and pT654. The induction of
oligomerization seems to be a general effect of cholesterol upon α-helix peptides in lipid
bilayers (32). Our result does not necessarily mean trimerization in the physiological
conditions but reflects a tendency of oligomerization. We picked up trimers in the analysis
to avoid complexity in the interactions between more than three molecules. While in the
PC/PS membrane without cholesterol (Figs. 6 and 7), the probability to adopt a one dimer
+ one monomer configuration seems to be increased for the non-phosphorylated peptides,
likely because peptides have difficulty forming trimers when containing the anti-parallel
helix JM dimer, the pT654 event appears to dissociate the JM dimer to help in the
formation of the close trimer especially in the presence of cholesterol. If this assumption
is correct, oligomer formation will be inhibitory for EGFR kinase activity. Our observed
increases in the pY1068 level in the T654A mutant of EGFR (Fig. 9) support this
possibility. Recently, we found that cholesterol depletion induces hyperphosphorylation
of Y1068 and Y1173 of EGFR upon EGF stimulation. This seems to be caused by an
increase in the pre-dimer fraction of EGFR. However, importantly, reactions downstream
of EGFR autophosphorylation, including the translocation of GRB2 to the cell surface
and phosphorylations of SHC, AKT, and ERK were diminished under the cholesterol
depletion. At the same time, clustering of EGFR over dimers was strongly inhibited (33).
These observations are consistent with the results of this study.

The antiparallel helix dimer of JM is thought to facilitate asymmetric interaction
between the kinase domains of EGFR, and hence its activation, in order to phosphorylate
tyrosine residues in the C-tail (11, 34). This tyrosine phosphorylation results in the
recruitment of PKC and other threonine kinases from the cytoplasm to the EGFR
molecules for the phosphorylation of Thr654, which is known to negatively regulate
EGFR signaling (26, 35). Our previous results suggested that the mechanism underlying
this negative effect of pT654 is the dissociation of JM dimers in the presence of acidic
lipids (20). At the same time, pT654 might induce the oligomerization of EGFR in the
presence of cholesterol. Supporting this possibility, our current single-molecule imaging
in living cells revealed the oligomerization of unliganded wt EGFR after PMA treatment,
which induced pT654 (Fig. 10). In the unliganded condition, the extracellular domains of
EGFR are thought to prevent the JM dimerization for kinase activation (9, 36). Clustering
of EGFR is also prevented largely in the unliganded condition. However, probably due to
the structural fluctuation, a small part of the unliganded EGFR are oligomerized (33).
pT654 (PMA) stabilized the unliganded oligomers possibly destructing the JM
antiparallel dimers.

We previously reported that oligomers of EGFR formed after cell stimulation

with EGF function as the major signal transduction sites for GRB2, where the dissociation
rate constant with GRB2 was reduced increasing the affinity between single-molecules
of EGFR and GRB2 (14). In addition, our current analyses found an increase in the wt
EGFR/GRB2 association following PMA treatment (Fig. 11). We speculate that after the
initial Tyr phosphorylation, successive Thr phosphorylation reduces Tyr kinase activity
of EGFR but enhances the formation of signal transduction oligomer in the medium
immobilized and oligomerized state of EGFR molecules (Fig. 10d). Further
immobilization and oligomerization were found in our current experiments to be induced
by EGF in the absence of PMA for wt EGFR and with or without PMA for the T654A
mutant. This process might include EGFR molecules accumulated into the clathrin coated
pits (37) and be independent of pT654. In the previous report (38), ERK activation was
reduced by the Thr654/669 of EGFR to Ala substitution suggesting only negative effects
of Thr654, however, oligomerization of EGFR was not observed in this report. Distinct
from wt EGFR, the T654A mutation in EGFR suppressed the GRB2 association after
EGF association in spite of the higher levels of Tyr phosphorylation at the binding site
compared to wt. Thus, even though pT654 is inhibitory for EGFR kinase activity, it
promotes signal transduction to the cytoplasmic protein, GRB2.

Based on our present results, we propose a model of EGFR signaling regulated
by membrane lipids and Thr654 phosphorylation (Fig. 11d). The signal transduction
mediated by EGFR is a complex multi-step process. Conformational changes in the
extracellular domain of EGFR upon ligand association allow JM domains to form acidic
lipid-facilitated anti-parallel JM helix dimers and asymmetric kinase domain dimers(11,
34). This is the activation process for EGFR kinase, in which acidic membrane lipids and
cholesterol play stimulative and inhibitory roles, respectively (20, 33). Tyrosine
phosphorylation in the kinase-active EGFR dimers recruits PKC from the cytoplasm (39).
The association of PLCγ to the EGFR phosphotyrosine for the degradation of PIP_2 is
involved in this process. PKC then phosphorylates Thr654 (40), which dissociates anti-
parallel JM dimers in the presence of remaining acidic lipids and supports the
oligomerization of EGFR in the presence of cholesterol, especially after the removal of
acidic lipids around the EGFR molecules. The cholesterol-induced oligomer of EGFR is
a major site of interaction with cytoplasmic proteins including GRB2 (14). Thus, a major
function of EGFR is shifted from a kinase for self-activation to a scaffold for signal
transduction. Thr654 phosphorylation is a key step underlying this role change of EGFR
and is not merely an inactivating mechanism. The degradation of PIP_2, a major anionic
lipid in the inner leaflet of the plasma membrane, may support this role change.
Importantly, both Thr654 phosphorylation and PIP_2 degradation are caused by the kinase

activation of EGFR. Hence, this represents an ingenious autoregulatory process involving
membrane proteins and lipids.
417

Materials and Methods

Materials

1-palmitoyl-2-oleoyl-*sn*-phosphatidylcholine (PC), 1-palmitoyl-2-oleoyl-*sn*-phosphatidylserine (PS), and cholesterol were purchased from Avanti Polar Lipids (Alabaster, AL) as chloroform solutions (PC and PS) or powders (cholesterol). Cy3-maleimide and Cy5-maleimide were purchased from GE Healthcare Life Sciences (Little Chalfont, UK). n-octyl-b-D-glucoside (OG) was purchased from Dojindo (Kumamoto, Japan). Monofunctional polyethylene glycol-succinimidyl valerate (s-PEG, 5000 mol wt) and biotinylated monofunctional polyethylene glycol-succinimidyl valerate (b-PEG, 5000 mol wt) were purchased from Laysan Bio (Arab, AL). Chinese hamster ovary K1 (CHO-K1) cells were provided from RIKEN BRC through the National Bio-Resource Project (MEXT, Tokyo, Japan).

Plasmid construction

Construction of the cDNA of full-length human EGFR (wt) fused with GFP was described previously (14). T654A mutant DNA was constructed using PrimeSTAR Max (Takara, Kusatsu, Japan) in the wt EGFR vector. The primer sequences were as follows: EGFR(T654A)-f: GAAGCGCGCGCTGCGGAGGCTGCTGC and EGFR(T654A)-r: CCGCAGCGCGCGCTTCCGAACGATGTG, respectively. For NanoBiT assays, full-length human EGFR (wt or T654A mutant) was fused with LgBiT at the C-terminus (wt or T654A EGFR-LgBiT), and GRB2 was fused with SmBiT at the N-terminus (GRB2-SmBiT) as follows. The LgBiT fragment amplified from pBiT1.1-C [TK/LgBiT] Vector (Promega) using KOD One PCR Master Mix (TOYOBO) was subcloned into the AgeI- and NotI-digested EGFP-N1 vector (Clontech), and subsequently full-length EGFR fragment was subcloned into the NheI- and HindIII-digested the LgBiT-inserted EGFP-N1 vector. The GRB2-SmBiT fragment was constructed using KOD One PCR Master Mix (TOYOBO) and subcloned into the AgeI- and SalI-digested EGFP-C2 vector (Clontech). The primer sequence of SmBiT was designed from pBiT2.1-N [TK/SmBiT] Vector (Promega).

Peptide synthesis and purification

Peptides corresponding to the TM-JM regions of EGFR (618-666) were synthesized by solid-phase methods with the sequence KIPSIATGMVGALLLLLVVALGIGLFM-RRRHIVRKRT$_{654}$LRRLLQERELVE-NH$_2$ (28). For the experiments with the C-terminal labeled EGFR peptide, peptides containing a cysteine at the C-terminus were synthesized. These synthetic peptides were purified by reverse-phase high-performance liquid

chromatography on a C4 column with a gradient of 1-propanol and acetonitrile (1:1) over
0.1% aqueous trifluoroacetic acid. To prepare the C-terminal labeled peptide, Cy3-
maleimide or Cy5-maleimide was introduced to the sulfide group on the cysteine at the
C-terminus of the TM-JM peptide by mixing the peptide and the fluorescence derivative
in dimethyl formamide under basic conditions. For experiments with the N-terminal
labeled peptide, Cy3-COOH or Cy5-COOH was reacted with an elongating peptide on
the resin in the presence of 1-[bis(dimethylamino)methylene]-1H-benzotriazolium 3-
oxide hexafluorophosphate (HBTU) and diisopropylethylamine (DIEA), which activate
the carboxyl group on the fluorophore derivative. For synthesis of Thr654 phosphorylated
peptides, phosphorylated threonine derivatives were utilized. The purity was confirmed
by reverse-phase high-performance liquid chromatography and matrix-assisted laser-
desorption/ionization time-of-flight mass spectroscopy analysis.
466

Nanodisc preparation
For nanodisc construction, fluorescent EGFR TM-JM peptides co-solubilized with lipids
and OG in hexafluoroisopropanol were first dried to form thin films. These peptide films
were then resolubilized in buffer A (0.5 M NaCl, 20 mM Tris/Cl, 0.5 mM EDTA)
containing 30 mM OG and 5 mM dithiothreitol (pH 7.5). His8-tagged MSP 1E3D1 (MSP)
was expressed in *E.* coli and purified as described previously (41). The concentration of
MSP was quantified based on the absorbance at 280 nm (29,910 $M^{-1}cm^{-1}$). Thin PC or PS
films were formed by evaporation of the solvent (chloroform) under a steam of nitrogen
gas and dried in vacuum. Cholesterol powders were first dissolved in chloroform, and a
thin film was formed as described above. PC, PS, and cholesterol were resuspended in
buffer A containing 0.4 M sodium cholate (pH 7.5) at a final concentration of 10 mM.
Cy3- and Cy5-labeled TM-JM peptides in buffer A were mixed in equal amounts and then
conjugated with MSP and phospholipid mixtures (PC, PC/PS, PC/cholesterol,
PC/PS/cholesterol) at a molar ratio of 1:1:120 μM (TM-JM/MSP/lipids). The mixture was
dialyzed against a buffer containing 0.5 M NaCl, 20 mM Tris/Cl, and 5 mM EDTA (pH
7.5) at 4°C to reconstitute the nanodiscs by removing the detergent. The aggregates and
liposomes were removed from the mixture by size-exclusion chromatography using a
Superdex 200 Increase column (GE Healthcare Life Sciences) and the peak fractions
containing nanodiscs of around 11 nm in diameter were collected.
486

Single-pair FRET (spFRET) measurements
Nanodisc samples were immobilized on the surface of a glass chamber as described
previously (20, 42, 43). Briefly, amine-modified glass surfaces were coated with 99% s-

PEG and 1% b-PEG. NeutrAvidin (Thermo Fisher Scientific、 Waltham, MA) was then

bound to the b-PEG. The nanodisc samples bound with biotinylated anti-His8-tag

antibody (MBL Life Science) were loaded into the glass chamber and allowed to bind to

the NeutrAvidin-coated glass surface, after which unbound nanodiscs were washed away.

To reduce the photobleaching rate of Cy3 and Cy5, the nanodisc-loaded chamber was

filled with dialysis buffer containing 2-mercaptoethanol at the final concentration of 0.5%

(w/v). The fluorescence of Cy3 and Cy5 was observed under a TIRF microscope based

on an inverted microscope (Ti2; Nikon) with a 60x oil-immersion objective (ApoTIRF

60x 1.49 NA; Nikon). The fluorescence activity of Cy3 was excited using a 532 nm laser

(Compass 315M-100). Dual-color imaging was carried out through a 4x relay lens by

using two EMCCD cameras (C9100-134, ImagEM; Hamamatsu Photonics, Hamamatsu,

Japan) with a 200x EM gain. Images of 512 x 512 pixels (67 nm/pixel) were recorded

with a temporal resolution of 100 ms/frame using MetaMorph (Molecular Devices, San

Jose, CA) or AIS (ZIDO, Toyonaka, Japan).

Analysis of FRET signals

The measurement of fluorescence intensities of single nanodiscs was performed using

ImageJ software, as described previously (44). The background noise was filtered out

using the Subtract Background function in ImageJ. Fluorescence intensities of Cy3 and

Cy5 in single nanodiscs were measured as averages from circles with a diameter of 12

pixels containing a fluorescence spot. The average intensity of the same sized circles in

which no spot was present was subtracted as the background. Along the fluorescence

trajectories of TM-JM-Cy3 and TM-JM-Cy5, the FRET efficiency, E_{FRET}, for each frame

(> 2000) was calculated from the fluorescence intensities in the donor I_D and FRET I_A

channels as

515
$$E_{FRET} = \frac{I_A - \beta I_D}{I_A + (\gamma - \beta)I_D},$$

where β and γ are coefficients for the compensation of fluorescence leakage from the

donor dye to the acceptor detector channel, and the difference in the detection efficiencies

of the dyes, respectively (45). Coefficients were calculated using the intensity time traces

as $\beta = 0.03$ and $\gamma = 0.4$, respectively. E_{FRET} distributions were all different in Kolmogorov-

Smirnov (KS) test with a rejection p < 0.006. The peak positions and their 95% percentile

section were estimated using bootstrap method from a discretized E_{FRET} distribution

(Table I), in which the bin width was set to 0.02 and the resampling number was 300. KS

test was done using "StatsKSTest" in Igor Pro 8.0 (WaveMetrix). Resampling for

bootstrap method was done using "sample" function in R.

Cell culture and transfection

CHO-K1 cells were maintained in HAM F12 medium supplemented with 10% fetal
bovine serum at 37°C under 5% CO_2. HEK293S cells were maintained in DMEM F12
medium supplemented with 10% fatal bovine serum at 37°C under 5% CO2. For western
blotting assays, DNA constructs of full-length wt and T654A EGFR (1 μg) were
transiently transfected into CHO-K1 cells using FuGENE HD Transfection Reagent
(Promega, Madison, WI). For single-molecule measurements, CHO-K1 cells were
transfected with either a wt or T654A EGFR-GFP gene (0.5 μg each) using Lipofectamine
3000 Reagent (Thermo Fisher Scientific). For NanoBiT assays, HEK293S cells were
transfected with a mixture of wt or T654A EGFR-LgBiT gene (1 μg each) and GBR2-
SmBiT (0.2 μg) using Lipofectamine 3000 Reagent in 60 mm dish.

PMA treatment and EGF stimulation

DNA constructs of full-length wt and T654A EGFR were transfected and cultured with
10% fetal bovine serum (FBS) on the day before each measurement. Cells were then
starved in modified Eagle's medium without FBS for 3 hours before the experiment.
Phorbol 12-myristate 13-acetate (PMA) was dissolved in DMSO and subsequently
diluted in PBS to a final concentration of 10 μM. For PMA pre-treatment, PMA solution
was added to the cell cultured medium at a final concentration of 100 nM and incubated
for 30 min at room temperature. For EGF stimulation, EGF (PeproTech, Cranbury, NJ)
dissolved in PBS was added to the cell cultured medium at a final concentration of 100
ng/mL (for western blotting assays and single-molecule measurements) or as a 0.001 to
100 nM dilution series (for the NanoBiT assay).

Western blotting analysis

In cells stimulated with EGF for 0, 5, 30 min at 37°C, threonine and tyrosine
phosphorylation of the wt and mutant T654A proteins was detected by western blotting
using rabbit anti-pT654 (ab75986; Abcam, Cambridge, UK) and rabbit anti-pY1086
antibody (#4407; Cell Signaling Technology, Danvers, MA), respectively. Rabbit anti-
EGFR antibody (#sc-03; Santa Cruz Biotechnology, Dallas, TX) was used to detect
protein expression. After being resolved by SDS-polyacrylamide gel electrophoresis
(PAGE), the electrophoresed proteins were transferred onto a polyvinylidene difluoride
(PVDF) membrane and incubated with each antibody (primary antibody) and then with a
horseradish peroxidase (HRP)-linked anti-rabbit IgG (secondary antibody; 7076, Cell

Signaling Technology). Immunoreactive proteins were detected with Amersham ECL
Prime Western Blotting Detection Reagent (GE Healthcare) using an ImageQuant LAS
500 device (GE Healthcare).
563
Single-molecule imaging in living cells
The methods for single-molecule measurement and analysis were described elsewhere
(Yanagawa and Sako, Methods in Mol Biol, in press; bioRxiv: doi:
10.1101/2020.06.08.141192). The single-molecule imaging of EGFR was performed at
the basal plasma membrane of the CHO-K1 cells at 25°C with the same microscopic
methods used for the spFRET measurements. The laser wavelength was 488 nm (Sapphire
488; Coherent, Santa Clara, CA) for the excitation of the GFP. Fluorescence images were
acquired every 50 ms using AIS software. The acquired multiple TIFF files were
processed by ImageJ software as follows: background subtraction was performed with a
rolling ball radius of 25 pixels, and two-frame averaging of the images was then
performed. Single-molecule tracking analysis was performed with AAS software (ZIDO).
All   subsequent   analyses   were   performed   using   smDynamicsAnalyzer
(https://github.com/masataka-yanagawa/IgorPro8-smDynamicsAnalyzer), an Igor Pro
8.0-based homemade program.
578
NanoBiT assay
HEK293S cells co-transfected with the plasmids of wt or T654A EGFR-LgBiT and
GRB2-SmBiT. Overnight after the transfection, cells were collected in 0.5 mM EDTA-
containing PBS, centrifuged, and suspended in 2 mL of HBSS containing 0.01 % bovine
serum albumin and 5 mM HEPES (pH 7.4) (assay buffer). The cell suspension was
dispensed in a 96-well white bottom plate at a volume of 80 μL per well and loaded with
20 μL of 25 μM Nano-Glo Vivazine Live Cell Substrates (Promega) diluted in the assay
buffer. After incubation for 2 hrs at room temperature in the assay buffer, cells were
pretreated with PMA or vehicle as described above. Basal luminescence was then
measured by using a microplate leader (SpectraMax L, Molecular Devices) with an
interval of 60 sec at room temperature. After 10 min, 20 μL of the EGF dilution series in
the assay buffer or the assay buffer (vehicle) were applied to each well using a benchtop
multi-pipetter (EDR-384SR, BioTec, Tokyo, Japan) under red dim light. Then,
luminescence was measured for 30 min with an interval of 60 sec. Each time-course of
luminescence counts was normalized with the luminescence counts of the vehicle-added
well. Dose-response curves were fitted with a Hill-equation to determine the maximum
intensity.

Acknowledgments

YS was supported by MEXT Japan with Grants-in-Aid for Scientific Research (19H05647) and by JST CREST (JPMJCR1912). We thank Hiromi Sato for technical assistance.

Author contributions

Conceptualization, R.M., Y.S.; methodology, R.M., T.S., M.Y., Y.S.; investigation, R.M., H. T., T. S., M.Y.; manuscript writing, R.M., T. S., Y.S. with feedback from all other coauthors; funding acquisition, Y.S.; supervision, Y.S.

Conflicts of interest

None.

References

1. M. A. Olayioye, R. M. Neve, H. A. Lane, N. E. Hynes, The ErbB signaling network:
receptor heterodimerization in development and cancer. *EMBO J.* **19**, 3159-3167
(2000).

2. M. K. Nyati, M. A. Morgan, F. Y. Feng, T. S. Lawrence, Integration of EGFR inhibitors
with radiochemotherapy. *Nat. Rev. Cancer* **6**, 876-885 (2006).

3. E. Kovacs, J. A. Zorn, Y. Huang, T. Barros, J. Kuriyan, A structural perspective on the
regulation of the epidermal growth factor receptor. *Annu. Rev. Biochem.* **84**, 739-764
(2015).

4. N. J. Bessman, M. A. Lemmon, Finding the missing links in EGFR. *Nat. Struct. Mol.*
Biol. **19**, 1-3 (2012).

5. H. Ogiso *et al.*, Crystal structure of the complex of human epidermal growth factor and
receptor extracellular domains. *Cell* **110**, 775-787 (2002).

6. X. Zhang, J. Gureasko, K. Shen, P. A. Cole, J. Kuriyan, An allosteric mechanism for
activation of the kinase domain of epidermal growth factor receptor. *Cell* **125**, 1137-
1149 (2006).

7. M. J. Wagner, M. M. Stacey, B. A. Liu, T. Pawson, Molecular mechanisms of SH2- and
PTB-domain-containing proteins in receptor tyrosine kinase signaling. *Cold Spring*
Harb Perspect Biol **5**, a008987 (2013).

8. K. M. Ferguson *et al.*, EGF activates its receptor by removing interactions that
autoinhibit ectodomain dimerization. *Mol. Cell* **11**, 507-517 (2003).

9. N. F. Endres *et al.*, Conformational coupling across the plasma membrane in activation
of the EGF receptor. *Cell* **152**, 543-556 (2013).

10. E. R. Wood *et al.*, A unique structure for epidermal growth factor receptor bound to
GW572016 (Lapatinib): relationships among protein conformation, inhibitor off-rate,
and receptor activity in tumor cells. *Cancer Res.* **64**, 6652-6659 (2004).

11. N. Jura *et al.*, Mechanism for activation of the EGF receptor catalytic domain by the
juxtamembrane segment. *Cell* **137**, 1293-1307 (2009).

12. S. E. Webb *et al.*, Single-molecule imaging and fluorescence lifetime imaging microscopy
show different structures for high- and low-affinity epidermal growth factor receptors in
A431 cells. *Biophys. J.* **94**, 803-819 (2008).

13. Y. Huang *et al.*, Molecular basis for multimerization in the activation of the epidermal
growth factor receptor. *Elife* **5** (2016).

14. M. Hiroshima *et al.*, Transient Acceleration of Epidermal Growth Factor Receptor
Dynamics Produces Higher-Order Signaling Clusters. *J. Mol. Biol.* **430**, 1386-1401
(2018).

15.   N. Jura, Y. Shan, X. Cao, D. E. Shaw, J. Kuriyan, Structural analysis of the catalytically
646   inactive kinase domain of the human EGF receptor 3. *Proc Natl Acad Sci U S A* **106**,
647   21608-21613 (2009).
648   16.   S. J. Fleishman, J. Schlessinger, N. Ben-Tal, A putative molecular-activation switch in
649   the transmembrane domain of erbB2. *Proc Natl Acad Sci U S A* **99**, 15937-15940
650   (2002).
651   17.   A. Arkhipov *et al.*, Architecture and membrane interactions of the EGF receptor. *Cell*
652   **152**, 557-569 (2013).
653   18.   M. Lelimousin, V. Limongelli, M. S. Sansom, Conformational Changes in the Epidermal
654   Growth Factor Receptor: Role of the Transmembrane Domain Investigated by Coarse-
655   Grained MetaDynamics Free Energy Calculations. *J. Am. Chem. Soc.* **138**, 10611-10622
656   (2016).
657   19.   C. Matsushita *et al.*, Transmembrane helix orientation influences membrane binding of
658   the intracellular juxtamembrane domain in Neu receptor peptides. *Proc Natl Acad Sci U*
659   *S A* **110**, 1646-1651 (2013).
660   20.   R. Maeda, T. Sato, K. Okamoto, M. Yanagawa, Y. Sako, Lipid-Protein Interplay in
661   Dimerization of Juxtamembrane Domains of Epidermal Growth Factor Receptor.
662   *Biophys. J.* **114**, 893-903 (2018).
663   21.   K. B. Abd Halim, H. Koldso, M. S. P. Sansom, Interactions of the EGFR juxtamembrane
664   domain with PIP2-containing lipid bilayers: Insights from multiscale molecular dynamics
665   simulations. *Biochim Biophys Acta* **1850**, 1017-1025 (2015).
666   22.   Y. Wang *et al.*, Regulation of EGFR nanocluster formation by ionic protein-lipid
667   interaction. *Cell Res.* **24**, 959-976 (2014).
668   23.   Y. Liu *et al.*, The involvement of lipid rafts in epidermal growth factor-induced
669   chemotaxis of breast cancer cells. *Mol. Membr. Biol.* **24**, 91-101 (2007).
670   24.   T. Furuchi, R. G. Anderson, Cholesterol depletion of caveolae causes hyperactivation of
671   extracellular signal-related kinase (ERK). *J. Biol. Chem.* **273**, 21099-21104 (1998).
672   25.   T. Ringerike, F. D. Blystad, F. O. Levy, I. H. Madshus, E. Stang, Cholesterol is
673   important in control of EGF receptor kinase activity but EGF receptors are not
674   concentrated in caveolae. *J. Cell Sci.* **115**, 1331-1340 (2002).
675   26.   K. A. Lund *et al.*, Phosphorylation of the Epidermal Growth-Factor Receptor at
676   Threonine 654 Inhibits Ligand-Induced Internalization and down-Regulation. *J. Biol.*
677   *Chem.* **265**, 20517-20523 (1990).
678   27.   M. Hiroshima, Y. Saeki, M. Okada-Hatakeyama, Y. Sako, Dynamically varying
679   interactions between heregulin and ErbB proteins detected by single-molecule analysis
680   in living cells. *Proc Natl Acad Sci U S A* **109**, 13984-13989 (2012).

28. M. Yasui, M. Hiroshima, J. Kozuka, Y. Sako, M. Ueda, Automated single-molecule
imaging in living cells. *Nat Commun* **9**, 3061 (2018).

29. A. S. Dixon *et al.*, NanoLuc Complementation Reporter Optimized for Accurate
Measurement of Protein Interactions in Cells. *ACS Chem Biol* **11**, 400-408 (2016).

30. T. Sato, P. Pallavi, U. Golebiewska, S. McLaughlin, S. O. Smith, Structure of the
membrane reconstituted transmembrane-juxtamembrane peptide EGFR(622-660) and
its interaction with Ca2+/calmodulin. *Biochemistry* **45**, 12704-12714 (2006).

31. D. H. Jones, K. R. Barber, C. W. Grant, The EGF receptor transmembrane domain: 2H
NMR study of peptide phosphorylation effects in a bilayer environment. *Biochemistry*
37, 7504-7508 (1998).

32. K. Matsuzaki, Why and how are peptide-lipid interactions utilized for self-defense?
Magainins and tachyplesins as archetypes. *Biochim Biophys Acta* **1462**, 1-10 (1999).

33. M. Hiroshima *et al.*, Membrane cholesterol interferes with tyrosine phosphorylation but
facilitates the clustering and signal transduction of EGFR. *bioRxiv*
10.1101/2021.08.28.457965 (2021).

34. M. Red Brewer *et al.*, The juxtamembrane region of the EGF receptor functions as an
activation domain. *Mol. Cell* **34**, 641-651 (2009).

35. X. Li, Y. Huang, J. Jiang, S. J. Frank, ERK-dependent threonine phosphorylation of EGF
receptor modulates receptor downregulation and signaling. *Cell. Signal.* **20**, 2145-2155
(2008).

36. H. J. Zhu, J. Iaria, S. Orchard, F. Walker, A. W. Burgess, Epidermal growth factor
receptor: association of extracellular domain negatively regulates intracellular kinase
activation in the absence of ligand. *Growth Factors* **21**, 15-30 (2003).

37. L. K. Goh, F. Huang, W. Kim, S. Gygi, A. Sorkin, Multiple mechanisms collectively
regulate clathrin-mediated endocytosis of the epidermal growth factor receptor. *J. Cell*
Biol. **189**, 871-883 (2010).

38. M. Kluba, Y. Engelborghs, J. Hofkens, H. Mizuno, Inhibition of Receptor Dimerization
as a Novel Negative Feedback Mechanism of EGFR Signaling. *PLoS One* **10**, e0139971
(2015).

39. K. Oda, Y. Matsuoka, A. Funahashi, H. Kitano, A comprehensive pathway map of
epidermal growth factor receptor signaling. *Mol. Syst. Biol.* **1**, 2005 0010 (2005).

40. S. Aifa *et al.*, Phosphorylation of Thr654 but not Thr669 within the juxtamembrane
domain of the EGF receptor inhibits calmodulin binding. *Biochem. Biophys. Res.*
Commun. **347**, 381-387 (2006).

41. K. Kojima, Y. Imamoto, R. Maeda, T. Yamashita, Y. Shichida, Rod visual pigment
optimizes active state to achieve efficient G protein activation as compared with cone

visual pigments. *J. Biol. Chem.* **289**, 5061-5073 (2014).

42.   M. Pirchi *et al.,* Single-molecule fluorescence spectroscopy maps the folding landscape

of a large protein. *Nat Commun* **2**, 493 (2011).

43.   R. Lamichhane *et al.,* Single-molecule view of basal activity and activation mechanisms

of the G protein-coupled receptor β_2AR. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 14254-

14259 (2015).

44.   R. Maeda *et al.,* Single-molecule observation of the ligand-induced population shift of

rhodopsin, a G-protein-coupled receptor. *Biophys. J.* **106**, 915-924 (2014).

45.   K. Okamoto, Y. Sako, Variational Bayes Analysis of a Photon-Based Hidden Markov

Model for Single-Molecule FRET Trajectories. *Biophys. J.* **103**, 1315-1324 (2012).

 Figures

Figure 1. Construction of nanodiscs containing fluorescent TM-JM peptides of EGFR.

(**a**) Amino acid sequence of the EGFR TM and JM-A domains. EX, extracellular domain;

TK, tyrosine kinase domain. (**b, c**) Schematic images of nanodiscs containing dimeric

TM-JM peptides fluorescently labeled at the N- (b) and C-terminus (c), respectively. (**d**)

Lipid compositions used in the preparation of each nanodisc sample. The fractional ratios

of PS and cholesterol mimic those in the mammalian plasma membranes. (**e**) Size-

exclusion chromatography used for the purification of nanodiscs containing cholesterol

(red) or not (black). The charge ratios of PC/PS/cholesterol are described in the upper

table. The fraction having a peak absorbance of around 12 elution volumes (mL) was

collected and used for the subsequent experiments. (**f**) A thin-layer chromatography of

the nanodisc fraction containing cholesterol (fraction 12 in (**e**)). (**g**) A negative stain

electron micrograph of fraction 12 in (**e**). Size distribution of the nanodiscs calculated

from the images was fitted with a Gaussian function, with a mean diameter of 11 ± 2 nm.

Figure 2. Single-pair FRET measurement of the EGFR TM-JM dimers in nanodiscs. (**a**) Fluorescence micrograph of nanodiscs illuminated with a green laser. Cy3 (green) and Cy5 (red) emissions were superimposed. The Cy5 emission was caused by FRET from Cy3. (**b**) Representative fluorescence trajectories of Cy3 (green) and Cy5 (red). Black allows indicate photobleaching points of Cy5. (**c**) FRET efficiency trajectories of the fluorescence trajectories in (**b**). The FRET efficiency, E_{FRET}, was calculated as described in the Materials and Methods section. Typical fluorescence and FRET trajectories between peptides labeled at the C-terminus are shown. Transitions to low FRET efficiency states suggested that dissociation of the JM dimer occurred occasionally. The Förster radius R_0 between Cy3 and Cy5 is 5.6 nm.

Figure 3. FRET efficiency (E_{FRET}) distributions in nanodiscs containing a single Cy3/Cy5-pair of N-terminal labeled peptides. Nanodiscs contained non-phosphorylated (**a–d**) and Thr654 phosphorylated (**e–h**) peptides at the indicated lipid conditions. Positions of the mode and its 5-95% percentile section are indicated by solid and dashed lines, respectively. See Table I for these values. In **(b–h)**, the distribution and the 95% percentile section shown in **(a)** are superimposed (red) for comparison.

Figure 4. E_{FRET} distributions in nanodiscs containing a single Cy3/Cy5-pair of C-terminal labeled peptides. Nanodiscs contained non-phosphorylated (**a–d**) and Thr654 phosphorylated (**e–h**) peptides in the indicated lipid conditions. Positions of the mode and its 5-95% percentile section are indicated by solid and dashed lines, respectively. See Table I for these values. In (**b–h**), the distribution and the 95% percentile section shown in (**a**) are superimposed (red) for comparison.

Figure 5. Higher-order oligomerization of EGFR TM-JM peptides in the nanodiscs.

Histograms are shown of the total fluorescence intensity of the peptides with Cy3-labeling

at the C-terminus in single nanodiscs containing cholesterol (blue) or not (red). Discs

containing no Cy5 peptide were chosen for measurement to avoid the possible effects of

FRET. Peptides with a non-phosphorylated **(a, c)** or phosphorylated **(b, d)** Thr654 were

reconstituted into nanodiscs in PC **(a, b)** or PC/PS **(c, d)** membranes.

Figure 6. Assembly of EGFR TM regions. (**a**) Representative fluorescence trajectories of Cy3 (green) and Cy5 (red) in nanodiscs containing two Cy3-labeled and one Cy5-labeled peptide. Fluorescence intensities and/or two-step photobleaching dynamics after Cy5 photobleaching indicated that these nanodiscs contained two Cy3 peptides. (**b–i**) Fluorescence intensity histograms of N-terminal-labeled Cy3 peptides before (blue) and after (red) Cy5 photobleaching. Nanodiscs contained non-phosphorylated (**b–e**) and Thr654 phosphorylated (**f–i**) peptides at the indicated lipid conditions. (**j**) Schematic structures indicating proximity between three TM domains before (left) and after (right) Cy5 photobleaching. Note that acceptance of the excitation energy from Cy3 was not saturated for Cy5 under our experimental conditions, even in the presence of two Cy3 molecules.

Figure 7. Assembly of EGFR JM regions. (**a**) Representative fluorescence trajectories of
Cy3 (green) and Cy5 (red) in nanodiscs containing two Cy3-labeled and one Cy5-labeled
peptide at the C-terminus. (**b–i**) Fluorescence intensity histograms of C-terminal-labeled
Cy3 peptides from nanodiscs containing two Cy3 and one Cy5 peptide before (blue) and
after (red) Cy5 photobleaching. Nanodiscs contained non-phosphorylated (**b–e**) and
Thr654 phosphorylated (**f–i**) peptides at the indicated lipid conditions. (**j**) Schematic
structures indicating proximity between three JM domains before (left and middle) and
after (right) Cy5 photobleaching.

Figure 8. Possible configurations of EGFR TM-JM dimers and trimers regulated by
membrane lipids and Thr654 phosphorylation.
810

Figure 9. Thr and Tyr phosphorylation of EGFR. (**a**) Thr654 phosphorylation after EGF
stimulation and PMA pretreatment. (**b, c**) Timecourses of Y1068 phosphorylation for the
wt and T654A mutant of EGFR during EGF stimulation. Typical western blotting results
are indicated (**b, top**) and the average of four independent experiments are shown with
SE (**c**). Phosphorylation levels were normalized to the expression levels of the whole
EGFR molecule (**b, bottom**).
818

Figure 10. Oligomerization and lateral movements of EGFR on the living cell surface.
(**a**) Single molecule imaging of wt EGFR-GFP on the surface of living CHO-K1 cells
with (right) and without (left) PMA pretreatment. Cells were stimulated with (lower) and
without (upper) EGF. Bar, 5 μm. Quantized transitions in the fluorescence intensities
(bottom) indicate a single-molecule resolution of the imaging. (**b**) Oligomer size
distributions of wt (left) and T654A mutant (right) EGFR in cells. The oligomer size ratio
was measured before and 10 min after EGF stimulation. (**c**) Mean square displacement
(MSD) of wt (left) and T654A (right) EGFR spots as a function of the time interval,
indicating lateral mobility. The MSD was calculated before and 10 min after EGF
stimulation. In (**b, c**), cells were pretreated with (blue, green) or without (black, red) PMA
and stimulated (red, green) or not (black, blue) with EGF. (**d**) Diagram of the
oligomerization and immobilization states of wt (left) and T654A mutant (right) EGFR
suggested from single-molecule measurements. Arrows indicate the state transitions after
PMA treatment and EGF stimulation.
834

835
836
837

Figure 11. NanoBiT assay for the EGFR/GRB2 interaction in living cells. (**a**) Schematic
illustration of the NanoBiT assay of EGFR/GRB2 interactions. (**b**) Typical time courses
of chemiluminescence signals generated from the complex formation of large BiT
(LgBiT)-fused EGFR and small BiT (SmBiT)-fused GRB2 after EGF application at time
0. The final concentration of EGF in the culture medium was varied from 0.0001 to 100
nM. Measurements were done in cells with (+) or without (-) PMA pretreatment. (**c**) Dose
dependency of the chemiluminescence intensities at 30 min after EGF stimulation. The
average values from four independent experiments are shown with SE. Lines indicate
fitting with a Hill-equation function. *$p < 0.05$ determined by *t*-test against the signal in
wt cells without PMA. (**d**) A schematic model of the activation and signal transduction
process for EGFR dimers and oligomers. In the oligomers of EGFR, affinity with GRB$_2$
is increased.
850

852 **Table I. Peak values in the E_{FRET} distributions between the TM-JM peptides**

lipid composition	pT654	labelling			
		N-term		C-term	
		mean	95%	mean	95%
PC	-	0.93	(0.89-0.95)	0.81	(0.73-0.87)
PC/PS	-	0.90	(0.85-0.97)	0.83	(0.79-0.89)
PC/chol	-	0.95	(0.93-0.95)	0.93	(0.85-0.97)
PC/PS/chol	-	0.98	(0.97-0.99)	0.94	(0.91-0.97)
PC	+	0.85	(0.83-0.89)	0.68	(0.63-0.89)
PC/PS	+	0.87	(0.87-0.89)	0.36	(0.33-0.43)
PC/chol	+	0.99	(0.99)	0.90	(0.89-0.95)
PC/PS/chol	+	0.97	(0.97)	0.82	(0.81-0.89)

856 Mean and their 95% percentile section of the peak values in Figs 3 and 4 were estimated

857 by bootstrap method with 300-times of resampling. An increase of the resampling times

858 to 1000 did not change the result.