1

Stabilizing genetically unstable simple sequence repeats in the *Campylobacter*

***jejuni* genome by multiplex genome editing: a reliable approach for delineating**

multiple phase-variable genes

Short title: Genetic engineering of multiple hypermutable simple sequence repeats

Shouji Yamamoto*, Sunao Iyoda, and Makoto Ohnishi

Department of Bacteriology I, National Institute of Infectious Diseases, Tokyo, Japan

* Corresponding author

E-mail: yshouji@nih.go.jp (SY)

Abstract

    Hypermutability of simple sequence repeats (SSR) through DNA slippage is a major

mechanism of phase variation in *Campylobacter jejuni*. The presence of multiple SSR-

mediated phase-variable genes encoding enzymes that modify surface structures,

including capsular polysaccharide (CPS) and lipooligosaccharide (LOS), generates high

levels of structural variants within bacterial populations, thereby promoting adaptation

to selective pressures in host environments. Therefore, the phenotypic diversity

generated by phase variation can limit the reproducibility of results with *C. jejuni*;

therefore, researchers need to genetically control the mutability of multiple SSRs. Here,

we show that natural "cotransformation" is an effective method for *C. jejuni* genome

editing. Cotransformation is a trait of naturally competent bacteria that causes uptake

and integration of multiple different DNA fragments, which has been recently adapted

to multiplex genome editing by natural transformation (MuGENT), a method for

introducing multiple scarless mutations into the genomes of these bacteria. We found

that the cotransformation frequencies of antibiotic resistance gene-marked DNA

fragments and unmarked DNA fragments reached ~40% in *C. jejuni*. To examine the

feasibility of MuGENT in *C. jejuni*, we "locked" either different polyG SSR tracts in

strain NCTC11168 (which are located in the biosynthetic CPS and LOS gene clusters)

into either the ON or OFF configurations by interrupting the continuous runs of G

residues without changing the encoded amino acids. This approach, termed "MuGENT-

SSR," enabled the generation of all eight edits within 2 weeks and the identification of a

phase-locked strain with a highly stable type of Penner serotyping, a CPS-based

serotyping scheme. Furthermore, extensive genome editing of this strain by MuGENT-

SSR identified a phase-variable gene that determines the Penner serotype of

44   NCTC11168. Thus, MuGENT-SSR provides a platform for genetic and phenotypic

45   engineering of genetically unstable *C. jejuni*, making it a reliable approach for

46   elucidating the mechanisms underlying phase-variable expression of specific

47   phenotypes.

Author summary

Campylobacter jejuni is the leading bacterial cause of food-borne gastroenteritis in

developed countries and occasionally progresses to the autoimmune disease Guillain–

Barré syndrome. The genetically and phenotypically unstable features of this bacterial

species limit research and development efforts. A relatively large number of

hypermutable simple sequence repeat (SSR) tracts in the *C. jejuni* genome markedly

decreases its phenotypic stability through reversible changes in the ON or OFF

expression states of the genes in which they reside, a phenomenon called phase

variation. Thus, controlling SSR-mediated phase variation can be important for

achieving stable and reproducible research on *C. jejuni*. In this study, we developed a

feasible and effective approach to genetically manipulate multiple SSR tracts in the *C.*

jejuni genome using natural cotransformation, a trait of naturally transformable bacterial

species that causes the uptake and integration of multiple different DNA molecules. This

approach will greatly help to improve the genetic and phenotypic stability of *C. jejuni* to

enable diverse applications in research and development.

Introduction

Simple sequence repeats (SSR) or microsatellites in bacterial genomes are highly mutable because of the potential for DNA-strand slippage during DNA replication [1]. Mispairing DNA by slippage results in the insertion or deletion of one or more repeats within a repetitive DNA element. When an SSR tract is present within an open reading frame (ORF), promoter, or other regulatory sequences, its hypermutability mediates reversible and frequent changes in specific phenotypes through transcriptional or translational genetic switches between ON and OFF states, which is called "phase variation" [2,3]. In several bacterial species, including *Campylobacter jejuni*, the presence of multiple SSR-mediated phase-variable genes (also called contingency genes) per genome generates high levels of phenotypic variants within bacterial populations [4,5].

C. jejuni is the leading bacterial cause of food-borne gastroenteritis in developed countries, primarily depending on its ability to colonize the caeca of chickens and to survive in the food chain by attaching to undercooked chicken meat [6,7]. *C. jejuni* readily colonizes the intestinal mucosa of a wide variety of wild and domestic birds and other animals. Infections in poultry are usually asymptomatic, while human infection can cause significant inflammation and bloody diarrhea, occasionally progressing to the autoimmune disease, Guillain–Barré syndrome [8]. As a commensal bacterium of poultry and a human pathogen, *C. jejuni* needs to rapidly adapt to differences in host environments, such as changing nutrient compositions and immune systems. Additional selective pressures are caused by transmission through genetically and immunologically variable host populations and exposure to bacteriophages. *C. jejuni* likely utilizes phase variation as a major mechanism for adapting to these selective pressures.

A relatively large number of SSR tracts consisting of seven or more G or C bases was

unexpectedly found in the AT-rich genomes of *C. jejuni*. These polyG/C SSR tracts are

mainly involved in phase variation in this species, and a genomic analysis of four *C.*

jejuni strains indicated the presence of 12 to 29 tracts per genome [9]. *C. jejuni*

NCTC11168 encodes 29 polyG/C tracts, 23 of which are in protein-coding regions [10],

suggesting that phase variation in this strain is regulated mainly at the translational

level. The majority of these loci are clustered on the genomic regions predicted to

encode enzymes involved in modifying surface structures, including lipooligosaccharide

(LOS), capsular polysaccharide (CPS), and flagella, but a few are located on genes

encoding cell-surface proteins or restriction enzymes [5]. The ability to change the

antigenicity of surface structures by phase variation strongly suggests that this

phenotypic diversity could affect colonization or pathogenesis in host organisms.

Consistently, it has been reported that phase-variable expression of specific genes

affects pathogenesis in humans and colonization in chickens [11–15]. Comprehensive

studies of multiple phase-variable genes in *C. jejuni* have also demonstrated that

culturing or passage through animal and human hosts results in significant phase

changes in multiple SSR-containing genes, particularly those involved in surface-

structure modifications, which in some cases, are coupled with enhanced colonization

and pathogenesis [9,16–20].

A limitation of studying the contributions of multiple SSR-mediated phase-variable

genes to specific phenotypes is that researchers currently lack genetic tools for

efficiently "locking" SSR into either ON or OFF states at multiple loci in the *C. jejuni*

genome. For example, the use of a multiplexed SSR-editing technology enables

phenotype stabilization and therefore promotes reproducible research activities in this

area. Such phenotypic engineering would also be applicable for stably producing phase-

variable surface antigenic determinants, which can be used for vaccine development and

raising serotyping antisera. To date, multiplex automated genome engineering (MAGE)

using oligonucleotides, a highly efficient λRed recombinase, and clustered regularly

interspaced short palindromic repeats (CRISPR)/CRISPR-associated protein (Cas)

systems have been developed for targeted genome engineering in bacteria [21,22].

However, these methods require not only engineered plasmids or chromosomes for the

respective genome-editing systems (i.e., expressing proteins and/or RNAs involved in

the λRed and CRISPR/Cas systems) but also, in the case of CRISPR/Cas, the selection

of edited genomic sites. Natural competence for transformation is shared by diverse

bacterial species [23,24]. This process involves the uptake of exogenous DNA, followed

by integration of the imported DNA into the genome by homologous recombination.

Because natural transformation does not require any special factors and is dependent on

the recipient bacterium and donor DNA molecules, it has long been used for genetic

engineering in naturally competent bacterial species. *C. jejuni* was shown to be

naturally transformable over 30 years ago [25]. However, natural transformation has not

been conventionally used as a genetic engineering method in this bacterium because it

can be highly transformed with DNA prepared from *C. jejuni*, but not with DNA

propagated in *Escherichia coli* or amplified by the polymerase chain reaction (PCR).

Recently, Beauchamp *et al*. demonstrated that methylation at the RAATTY sequence of

E. coli-derived plasmid DNA or PCR-amplified DNA can efficiently transform *C. jejuni*

[26]. In a study of other transformable species conducted by Dalia *et al*., the authors

established the multiplex genome editing by natural transformation (MuGENT) system,

which is based on "cotransformation," a trait that causes the uptake and integration of

multiple different DNA molecules [27–29]. These two pioneering studies have provided

great potential for performing multiplexed gene modifications in *C. jejuni* and

motivated us to examine this feasibility. Here, we optimized natural transformation and

cotransformation using PCR-amplified donor DNA fragments and demonstrated its

utility as a method for multiplex genome editing in naturally competent *C. jejuni*.

Results and discussion

Optimization of donor DNA for the natural transformation of *C. jejuni*

Data from a previous study by Beauchamp *et al*. demonstrated that DNA methylated at the RA^{m6}ATTY sequence is an efficient substrate for *C. jejuni* transformation [26]. To optimize the natural transformation of this bacterium, we investigated in greater detail the characteristics of DNA substrate generated using PCR products. For this purpose, we amplified various DNA fragments of the *rpsL*^{K88R} allele, which confers streptomycin resistance (Sm^R) [30]. These *rpsL*^{K88R} marker fragments had different lengths of regions homologous to the recombination target sequence (50–2,000 base pairs [bp]) and different numbers of the EcoRI methyltransferase-recognition sequence (GAATTC), which was endogenously present or exogenously added to the PCR primers (Fig 1, n = 1 to 3). Each 0.5 pmol of the amplified DNA was treated with EcoRI methyltransferase and then used in transformation assays to transform the NCTC11168 and 81-176 strains, which are representative strains used for genetic studies of *C. jejuni* [5,31,32] (Table 1). We did not obtain transformants using fragments with short regions of homology (50 or 100 bp), even though these DNA fragments had two methylated GAATTC sites (Fig 1, *rpsL*^{K88R}-1 and *rpsL*^{K88R}-2). In contrast, the NCTC11168 and 81-176 strains were transformed at substantially higher frequencies using fragments with longer regions of homology (500 bp; Fig 1, *rpsL*^{K88R}-5). The highest transformation frequencies were obtained when 1,000- to 2,000-bp homologies were present (Fig 1, *rpsL*^{K88R}-8 and *rpsL*^{K88R}-9). The length of a homologous sequence is a major determinant of the efficiency of RecA-dependent recombination [33]. In the Δ*recA* background, no transformants were detected using a fragment with 2,000 homologous bp (Fig 1, *rpsL*^{K88R}-9), suggesting that transformation in *C. jejuni* is mediated by RecA. We also

confirmed that the lack of the GAATTC sequence or the lack of GAATTC methylation

markedly decreased the transformation frequency (Fig 1, $rpsL^{K88R}$-3 and unmethylated

$rpsL^{K88R}$-5). However, increasing the number of methylated GAATTC sites did

not substantially increase the transformation frequencies (Fig 1, $rpsL^{K88R}$-4 and

$rpsL^{K88R}$-5 or $rpsL^{K88R}$-6, $rpsL^{K88R}$-7, and $rpsL^{K88R}$-8). These results suggest that one

methylated GAATTC sequence is sufficient for transformation, consistent with previous

results obtained using plasmid DNA as a donor [26]. In summary, to maximize the

efficiency of *C. jejuni* transformation, the donor DNA should contain two key structural

elements: (1) a ≥1,000-bp region of homology and (2) at least one methylated GAATTC

site.

Fig 1. Effects of homology length and methylation on natural transformation. *C.*

jejuni cells were transformed using $rpsL^{K88R}$ marker fragments with different lengths of

homologous regions compared to the recombination target sequence (50–2,000 bp,

indicated in parentheses) and different numbers of EcoRI methyltransferase-recognition

sequences (RIs). Each 0.5 pmol of amplified DNA was treated with EcoRI

methyltransferase and then used in the transformation assays. The transformation

frequency was defined as the number of Sm^R colony-forming units (CFUs) divided by

the total number of CFUs. Data from three independent transformations are presented as

the means ± standard deviations. As controls, we used unmethylated DNA (indicated by

"Unmethylation of $rpsL^{K88R}$-5") or assayed for transformation in the absence of donor

DNA (indicated by "No addition of DNA"). "ND" indicates not done. The following

strains were used: NCTC11168 WT, NCTC11168; NCTC11168 Δ*recA*, SYC1004; 81-

176 WT, 81-176; 81-176 Δ*recA*, and SYC2004. The following DNA fragments were

used (S2 Table): $rpsL^{K88R}$-1 and $rpsL^{K88R}$-2 (for both strains); $rpsL^{K88R}$-3 and $rpsL^{K88R}$-

3-1 (for NCTC11168) and $rpsL^{K88R}$-3-2 (for 81-176); $rpsL^{K88R}$-4 and $rpsL^{K88R}$-4-1 (for

NCTC11168) and $rpsL^{K88R}$-4-2 (for 81-176); $rpsL^{K88R}$-5 and $rpsL^{K88R}$-5-1 (for

NCTC11168) and $rpsL^{K88R}$-5-2 (for 81-176); $rpsL^{K88R}$-6 and $rpsL^{K88R}$-6-1 (for

NCTC11168) and $rpsL^{K88R}$-6-2 (for 81-176); $rpsL^{K88R}$-7 and $rpsL^{K88R}$-7-1 (for

NCTC11168) and $rpsL^{K88R}$-7-2 (for 81-176); $rpsL^{K88R}$-8 and $rpsL^{K88R}$-8-1 (for

NCTC11168) and $rpsL^{K88R}$-8-2 (for 81-176); and $rpsL^{K88R}$-9 and $rpsL^{K88R}$-9-1 (for

NCTC11168) and $rpsL^{K88R}$-9-2 (for 81-176).

Table 1 Strains and plasmids used in this study

Strain or plasmid	Characteristics	Source or reference
Strain		
NCTC11168	*C. jejuni* human origin, serotype B (antigenic factor HS2)	[31]
SYC1001	NCTC11168 Δ*flaA*::*cat*	This study
SYC1002	NCTC11168 Δ*flaA*::*kan*	This study
SYC1003	NCTC11168 *rpsL*^{K88R}	This study
SYC1004	NCTC11168 Δ*recA*::*cat*	This study
SYC1006	NCTC11168 *cj1426*::*astA* Δ*flaA*::*kan*	This study
SYC1007	NCTC11168 *cj1426*^{ON}::*astA* Δ*flaA*::*cat*	This study
SYC1008	NCTC11168 *cj1426*^{OFF}::*astA* Δ*flaA*::*kan*	This study
SYC1P000K	NCTC11168 Δ*flaA*::*kan cj1139*^{OFF} *cj1144*^{OFF} *cj1420*^{OFF} *cj1421*^{OFF} *cj1422*^{OFF} *cj1426*^{OFF} *cj1429*^{OFF} *cj1437*^{OFF}	This study
SYC1P000	NCTC11168 *cj1139*^{OFF} *cj1144*^{OFF} *cj1420*^{OFF} *cj1421*^{OFF} *cj1422*^{OFF} *cj1426*^{OFF} *cj1429*^{OFF} *cj1437*^{OFF}	This study
SYC1P001C	NCTC11168 Δ*flaA*::*cat cj1139*^{OFF} *cj1144*^{OFF} *cj1420*^{OFF} *cj1421*^{OFF} *cj1422*^{OFF} *cj1426*^{OFF} *cj1429*^{OFF} *cj1437*^{ON}	This study
SYC1P004K	NCTC11168 Δ*flaA*::*kan cj1139*^{OFF} *cj1144*^{OFF}	This study

	$cj1420^{OFF}$ $cj1421^{OFF}$ $cj1422^{OFF}$ $cj1426^{ON}$ $cj1429^{OFF}$ $cj1437^{OFF}$	
SYC1P005C	NCTC11168 $\Delta flaA::cat$ $cj1139^{OFF}$ $cj1144^{OFF}$ $cj1420^{OFF}$ $cj1421^{OFF}$ $cj1422^{OFF}$ $cj1426^{ON}$ $cj1429^{OFF}$ $cj1437^{ON}$	This study
SYC1P032K	NCTC11168 $\Delta flaA::kan$ $cj1139^{OFF}$ $cj1144^{OFF}$ $cj1420^{ON}$ $cj1421^{OFF}$ $cj1422^{OFF}$ $cj1426^{OFF}$ $cj1429^{OFF}$ $cj1437^{OFF}$	This study
SYC1P033C	NCTC11168 $\Delta flaA::cat$ $cj1139^{OFF}$ $cj1144^{OFF}$ $cj1420^{ON}$ $cj1421^{OFF}$ $cj1422^{OFF}$ $cj1426^{OFF}$ $cj1429^{OFF}$ $cj1437^{ON}$	This study
SYC1P036C	NCTC11168 $\Delta flaA::cat$ $cj1139^{OFF}$ $cj1144^{OFF}$ $cj1420^{ON}$ $cj1421^{OFF}$ $cj1422^{OFF}$ $cj1426^{ON}$ $cj1429^{OFF}$ $cj1437^{OFF}$	This study
SYC1P037C	NCTC11168 $\Delta flaA::cat$ $cj1139^{OFF}$ $cj1144^{OFF}$ $cj1420^{ON}$ $cj1421^{OFF}$ $cj1422^{OFF}$ $cj1426^{ON}$ $cj1429^{OFF}$ $cj1437^{ON}$	This study
SYC1P037	NCTC11168 $cj1139^{OFF}$ $cj1144^{OFF}$ $cj1420^{ON}$ $cj1421^{OFF}$ $cj1422^{OFF}$ $cj1426^{ON}$ $cj1429^{OFF}$ $cj1437^{ON}$	This study
SYC1P255K	NCTC11168 $\Delta flaA::kan$ $cj1139^{ON}$ $cj1144^{ON}$ $cj1420^{ON}$ $cj1421^{ON}$ $cj1422^{ON}$ $cj1426^{ON}$ $cj1429^{ON}$ $cj1437^{ON}$	This study
SYC1P255	NCTC11168 $cj1139^{ON}$ $cj1144^{ON}$ $cj1420^{ON}$ $cj1421^{ON}$ $cj1422^{ON}$ $cj1426^{ON}$ $cj1429^{ON}$ $cj1437^{ON}$	This study
81-176	*C. jejuni* raw milk origin, serotype R (antigenic factors HS23/HS36)	[32]
SYC2001	81-176 $\Delta flaA::cat$	This study
SYC2002	81-176 $\Delta flaA::kan$	This study
SYC2003	81-176 $rpsL^{K88R}$	This study
SYC2004	81-176 $\Delta recA::cat$	This study
Plasmid		
pUCFa	Cloning vector, *bla*	Fasmac
pSYC-*cat*	pUCFa *cat* from *Campylobacter coli*	This study
pSYC-*kan*	pUCFa *kan* from *C. coli*	This study

Occurrence of natural cotransformation in *C. jejuni*

Using *Vibrio cholerae* and *Streptococcus pneumoniae*, Dalia *et al*. developed the MuGENT system using cotransformation, a trait of several naturally competent species [27–29]. During MuGENT, a bacterial culture is incubated with two types of donor DNA fragments: (1) a selected fragment that introduces an antibiotic-resistance gene into the genome and (2) unselected fragments that introduce scarless or transgene-free edits of interest at one or more loci. In *V. cholerae*, the frequencies of cotransformation of these distinct genetic markers can be made to exceed 60% by increasing the length of homology and the concentration of the unselected fragment [27]. To assess natural cotransformation in *C. jejuni*, we used two strains (SYC1003 and SYC2003) as recipients, which harbor the $rpsL^{K88R}$ mutations in the NCTC11168 and 81-176 genomes, respectively, and therefore are resistant to Sm (Fig 2). We also used a $\Delta flaA$::*kan* fragment with 1,000-bp regions of homology to replace the flagellin gene with a kanamycin-resistance (Km^R) marker (selected) and $rpsL^+$ fragments with homologous regions of different sizes (1,000 bp: $rpsL^+$-8, 2,000 b: $rpsL^+$-9) to revert to the wild-type Sm-sensitive (Sm^S) phenotype (unselected). After transforming SYC1003 and SYC2003 with equimolar concentrations of the selected and unselected methylated DNA (mDNA) fragments, we selected Km^R-transformants, 100 of which were then subjected to Sm-sensitivity testing to evaluate cotransformation (Fig 2). Using unselected mDNA with a 1,000-bp region of homology resulted in cotransformation frequencies of 24% and 16% in SYC1003 and SYC2003, respectively, and acquisition of the Km^R and Sm^S phenotypes (Fig 2). Furthermore, increasing the homology of unselected mDNA (2,000 bp) slightly increased the cotransformation frequencies of

both strains (Fig 2). Thus, we recommend the use of unselected mDNA with 2,000-bp

homology for efficient *C. jejuni* cotransformation.

Fig 2. Evaluation of natural *C. jejuni* cotransformation. Natural cotransformation of

C. jejuni was examined using two different PCR fragments, namely Δ*flaA*::*kan* and *rpsL*

$^+$ (S2 Table). The former fragments (Δ*flaA*::*kan*-1 for NCTC11168, or Δ*flaA*::*kan*-2 for

81-176) had 1,000 bp regions of homology (selected), whereas the latter fragments

(*rpsL*$^+$-8-1 and *rpsL*$^+$-9-1 for NCTC11168, or *rpsL*$^+$-8-2 and *rpsL*$^+$-9-2 for 81-176) had

1,000-b or 2,000-b regions of homology (unselected). "Me" indicates a methylated site.

Equimolar quantities (0.5 pmol) of selected and unselected fragments were

independently methylated (indicated by the labeling "Selected mDNA" and "Unselected

mDNA," respectively), mixed, and then purified. After transforming strains carrying the

rpsL^{K88R} allele (SY1003 and SY2003) with these mDNA fragments, the bacteria were

spread onto brain heart infusion (BHI) agar plates containing Km. One hundred Km^R-

transformants were replica-plated onto BHI agar plates containing Sm. The

cotransformation frequency (%) was calculated as follows: $100 \times$ number of Sm^S CFUs

per 100 Km^R CFUs. Data from three independent transformations are presented as the

means ± standard deviations.

Natural cotransformation is thought to reflect the nature of competent bacterial cells,

among which only a subpopulation of cells in a culture is transformable [28]. We

demonstrated for the first time that *C. jejuni* cells are capable of cotransformation. It

will be important to elucidate whether *C. jejuni* cells show variable competence within a

population.

Establishment of MuGENT in *C. jejuni*

MuGENT provides methods for simultaneously generating multiple scarless

mutations and can therefore be broadly applied in diverse research and biotechnology

applications [27]. We wanted to test whether natural cotransformation could be used for

multiplex genome editing of *C. jejuni*. Fig 3 presents a schematic representation of our

strategy. Briefly, multiple unselected mDNA fragments (used for introducing genome

edits of interest) were mixed at equimolar concentrations with a selected mDNA

fragment, and the resulting mixture was used to cotransform *C. jejuni* cells. Because

MuGENT often requires multiple cycles of cotransformation in order to complete the

genome editing, different selectable markers should be used for each cycle [27]. We

swapped different resistance markers at the *flaA* gene at every MuGENT cycle. This

also enabled easy removal of the marker genes and reversion to the wild-type allele by

transformation with *flaA*$^+$ mDNA and subsequent selection of motile clones. Genome

editing was verified by multiplex allele-specific colony (MASC) PCR [34] and

nucleotide sequencing.

Fig 3. Overview of MuGENT in *C. jejuni*. Multiple unselected mDNA fragments

(used for introducing genome edits of interest) were mixed at equimolar concentrations

with a selected mDNA fragment, in order to cotransform *C. jejuni* cells. During

MuGENT, we swapped different selective markers at the *flaA* gene during every cycle,

resulting in the cells becoming nonmotile (Mot$^-$). This also enabled easy removal of

marker genes and reversion to the wild-type allele by transformation with a *flaA*$^+$

mDNA and subsequent selection of motile (Mot$^+$) clones. Genome edits in the

transformants were verified by MASC PCR and nucleotide sequencing.

To demonstrate the efficacy of MuGENT in *C. jejuni*, we targeted the biosynthetic

genes, CPS and LOS. These cell-surface molecules are known to play key roles in

interactions that affect bacteriophage infectivity, chick colonization, invasion of human

epithelial cells, and host immune responses [34–40]. CPS is also the primary antigenic

determinant of the Penner serotyping scheme, a passive slide hemagglutination [41],

although other surface molecules, including LOS, may contribute to serotype specificity

[42]. Because the CPS and LOS gene clusters contain multiple phase-variable genes that

are interrupted by polyG tracts, *C. jejuni* cells can generate structural variations of CPS

and LOS, which aids in evading killing by host immune systems or predation by

bacteriophages [12,43–46]. In addition, phase-variable expression of CPS and LOS

markedly decreases the phenotypic stability of *C. jejuni* cells and, thus, may limit

related research, including basic studies, epidemiological surveillance, and vaccine

development. For example, the NCTC11168 strain (serotype B, antigenic factor HS2)

did not provide reproducible results for Penner serotyping, which frequently changed

between typeable and untypeable during subcultures (S1 Fig). Although this unstable

phenotype may be attributable to ON/OFF switching of the CPS expression due to

phase variation [41,47,48], the mechanism whereby phase variation regulates CPS gene

expression to determine the Penner serotype remains unknown.

There are 29 polyG/C tracts in the NCTC11168 genome, eight of which are located

within the CPS and LOS gene clusters [10]. Six of these phase-variable genes (*cj1420*,

cj1421, *cj1422*, *cj1426*, *cj1429*, and *cj1437*) reside in the CPS cluster, whereas the other

genes (*cj1139* and *cj1144*) reside in the LOS cluster, which theoretically gives rise to

$\sim 2^8$ different combinations of ON (1) /OFF (0) expression states or more specifically

"phasotypes" [4]. For example, a bacterium that has the following expression states —

cj1139^{ON} *cj1144*^{OFF} *cj1420*^{OFF} *cj1421*^{OFF} *cj1422*^{OFF} *cj1426*^{OFF} *cj1429*^{OFF} *cj1437*^{OFF} (in

that order) — would have a binary phasotype coded as 1-0-0-0-0-0-0-0, which can be

converted to decimal 128 format. In this study, we defined one of the 2^8 phasotypes

generated by these eight phase-variable genes as a "Penner phasotype (PPT)." Using

MuGENT, we tried to "lock" all of the eight ORFs into ON or OFF states, where their

polyG tracts were altered to translate into the largest possible ORF (locked-"ON" states)

or a smaller incomplete ORF by frameshifting through -1 deletions from the ON states

(locked-"OFF" states). For example, *cj1139* encodes a glucosyltransferase that mediates

phase variation of LOS epitopes responsible for autoimmunity [13], with G8 being in an

ON state and G7 in an OFF state (Fig 4A). To prevent replicative slippage at the polyG

tracts, we interrupted the continuous run of G residues without changing the translated

amino acids by replacing the last G residue of every G triplet with a different nucleotide

(Fig 4A). As an example of phase-locked mutant construction, we introduced locked-

OFF mutations into *cj1139*, *cj1144*, *cj1420*, *cj1421*, *cj1422*, *cj1426*, *cj1429*, and *cj1437*

by repeating the MuGENT cycle to generate a "PPT0" strain. During MuGENT,

genome editing from 40 transformants per cycle was monitored by MASC PCR. After

the second cycle, we found that 100% of the population had at least one edit (Fig 4B

and S2 Fig). We accomplished all eight edits within the 4th cycle, which took less than

2 weeks to perform (Fig 4B and S2 Fig). Thus, MuGENT works effectively in *C. jejuni*

and is useful for rapidly introducing multiple mutations.

Fig 4. Constructing a strain with all eight phase-variable genes locked into the OFF state by MuGENT against SSR (MuGENT-SSR) sites. The polyG tracts from eight genes (*cj1139*, *cj1144*, *cj1420*, *cj1421*, *cj1422*, *cj1426*, *cj1429*, and *cj1437*) located in the biosynthetic gene clusters for LOS and CPS in NCTC11168 were subjected to MuGENT-SSR to lock them all into OFF states. (A) Example of polyG editing (*cj1139*). The polyG tract was altered to translate into the largest possible ORF (G8: ON state, encoding 303 amino acids [aa]) or a smaller incomplete ORF by introducing frameshifts via -1 deletions (G7: OFF state, encoding 148 amino acids). To prevent replicative slippage at the polyG tract, we interrupted the continuous run of G residues without changing the translated amino acids by replacing the last G residue of every G triplet with a different nucleotide. We also used different nucleotides when locking the genes into ON and OFF states, so that these two different states could be swapped. (B) Distribution of genome edits in the population following successive cycles of MuGENT. This image was based on the MASC PCR data (S2 Fig).

Improving the phenotypic stability during Penner serotyping using MuGENT

Controlling SSR-mediated phase variation is important for achieving reproducible results with *C. jejuni*. Regarding the low reproducibility of Penner serotyping (S1 Fig), we hypothesized that this may be associated with the phase-variable expression of one or more genes that determine antigenicity. To narrow down candidate genes, we sequenced PPT from naturally occurring typeable and untypeable variants of NCTC11168, termed "PPT-Seq." PPT-Seq analysis of 12 variants revealed six PPT subtypes, one of which was typeable (Fig 5 and S1 Fig, PPT37). We constructed a strain locked to PPT37 by MuGENT and confirmed that it was also typeable (Fig 5 and S1

Fig). In contrast to the wild-type strain, the locked PPT37 mutant maintained typeability

during at least five subcultures (S1 Fig). To evaluate the effect of polyG editing by

MuGENT on phase variation, the *astA* gene (encoding arylsulfatase) [49] was fused in

frame to the phase-variable gene *cj1426* (Fig 5), which encodes a methyltransferase that

methylates the heptose of the CPS repeat unit of NCTC11168 [50], and to its phase-

locked derivatives, such that changes in the repeat number would alter arylsulfatase

expression [51]. Single colonies grown on BHI agar plates containing 5-bromo-4-

chloro-3-indolyl sulfate (XS), a chromogenic substrate of arylsulfatase, were used to

measure mutation rates, as described in the Methods section. With the wild-type

cj1426::*astA* fusion, the ON-to-OFF mutation occurred at a rate of 2.8×10^{-2}, whereas

the OFF-to-ON mutation rate was 2.3×10^{-2} (Table 2). In the locked-ON and locked-

OFF constructs, however, switching to different phases was not detectable (Table 2).

These results suggest that MuGENT can serve as a feasible and effective approach for

stabilizing unstable phenotypes generated by phase variation with a defined set of

multiple SSRs.

Fig 5. Penner serotyping and phasotyping of naturally occurring and phase-locked

variants of NCTC11168. After naturally occurring variants (typeable or untypeable by

Penner serotyping; SC1-1 to SC2-6, S1 Fig) were subjected to PPT-Seq, their PPTs were

decoded (indicated by decimal). The numbers with in red boxes indicate the numbers of

repeat polyG tracts in the ON configurations, whereas those in the gray boxes indicate

those in the OFF configurations. The serotyping results for the phase-locked variants are

shown in S3 Fig. The following phase-locked strains were used: Locked-0, SYC1P000;

Locked-255, SYC1P255; Locked-37, SYC1P037; Locked-36, SYC1P036C; Locked-5,

SYC1P005C; Locked-33, SYC1P033C; Locked-4, SYC1P004K; Locked-1,

SYC1P001C; Locked-32, SYC1P032K.

Table 2 Effect of the polyG editing on phase variation of *cj1426*

Reporter gene[a]	Mutation rate[b]	
	ON to OFF	OFF to ON
cj1426::*astA*	2.8×10^{-2}	2.3×10^{-2}
cj1426[ON]::*astA*	$<2.1 \times 10^{-3}$	ND[c]
cj1426[OFF]::*astA*	ND	$<2.2 \times 10^{-3}$

369　[a]The following strains were used: *cj1426*::*astA*, SYC1006; *cj1426*[ON]::*astA*, SYC1007;
370　and *cj1426*[OFF]::*astA*, SYC1008.
371　[b]Mutation rates were calculated using the equation reported previously by Drake [52]
372　(see Methods section).
373　[c]Not done
374

Using MuGENT-SSR to identify a phase-variable gene that determines the Penner

serotype of NCTC11168

Phenotypic engineering using MuGENT-SSR combined with PPT-Seq identified

PPT37 as a PPT subtype typeable for Penner serotyping. In PPT37, *cj1420*, *cj1426*, and

cj1437 were in the ON state, whereas the other five genes were in the OFF state. These

findings suggest that one or more genes of these "ON" genes may determine typeability.

To elucidate the responsible gene(s), we constructed strains locked to various PPT

subtypes by MuGENT-SSR and then performed serotyping. We demonstrated that

cj1426 expression was indispensable and sufficient for serological typing, whereas

cj1420 and *cj1437* expression were not (Fig 5 and S3 Fig). The CPS repeat unit of *C.*

jejuni NCTC11168 consists of β-D-ribose , β -D-N-acetylgalactosamine (GalfNAc), α-

D-glucuronic acid modified with 2-amino-2-deoxyglycerol at C-6, and 3,6-O-methyl-

D-glycero-α-L-gluco-heptose (Hep) [53] (S4 Fig). Examination of the CPS structure by

high-resolution magic-angle spinning nuclear magnetic resonance spectroscopy

revealed highly variable or no modification of the methyl, ethanolamine, aminoglycerol,

and phosphoramidate groups of the repeating unit [38,54]. In particular, changes in

methyl (Me) and phosphoramidate modifications were regulated by at least three phase-

variable genes located in the CPS cluster (*cj1421*, *cj1422*, and *cj1426*). The *cj1426* gene

encodes the methyltransferase involved in 6-O-Me modification of the Hep residue of

NCTC11168 CPS [50] (S4 Fig), suggesting that the presence of this modification may

be an antigenic determinant of the Penner B serotype (or antigenic factor HS2).

However, we observed that some PPT subtypes (including naturally occurring variants

PPT109, PPT111, PPT189, and PPT191, and a phase-locked variant PPT255), where

cj1426 and some or all of the other phase-variable genes (including *cj1144*, *cj1421*,

cj1422, and *cj1429*) were simultaneously switched ON, were untypeable (Fig 5, S1 and

S3 Figs). Of these genes, the *cj1421* and *cj1422* genes have already been demonstrated

to encode O-Me-phosphoramidate (MeOPN) transferases that attach MeOPN to the

GalfNAc and Hep residues, respectively [38]. Thus, one or more CPS modifications

catalyzed by these gene products may interfere with the binding of specific antibodies to

the epitope (S4 Fig). To directly test this possibility, structural analysis using nuclear

magnetic resonance spectroscopy and deeper genetic and immunogenic studies are

required.

Similar results were reported previously regarding the phase-variable CPS of *C.*

jejuni. In a series of studies on the tropism of *Campylobacter* bacteriophages, Sørensen

et al. demonstrated that the MeOPN-modified GalfNAc of the NCTC11168 CPS acts as

a receptor of the F336 phage, but that its receptor function was modulated by the

presence or absence of other CPS modifications [37,45]. In addition, a study of *C. jejuni*

81-176 (Penner serotype R, antigenic factors HS23 and HS36) revealed that phase-

variable MeOPN modifications at the three CPS galactose residues modulated serum

resistance [44]. These previous findings and our current findings suggest that phase-

variable changes in the CPS structure of *C. jejuni* occur at two levels: (1) phase

variations of the receptors and epitopes themselves, and (2) interference with receptor

and epitope functions by chemical modifications at different positions. These

combinatorial effects may aid in rapidly avoiding killing by phages and the host

immune system, while lowering the typeability of Penner serotyping.

Concluding remarks

Here, we demonstrate that MuGENT is applicable for genetic engineering of *C.*

jejuni. In this genetically unstable species, MuGENT specifically provided a feasible

and effective approach for editing multiple hypermutable SSRs. Specifically, MuGENT-

SSR was performed to uncover the contributions of multiple phase-variable genes to

specific phenotypes. By combining MuGENT-SSR with whole-genome SSR analysis

[4,10] may enable comprehensive studies of numerous phase-variable genes (~30

polyG/C-containing genes per genome) in order to decode the "phasotypes" that

determine specific phenotypes and more collective behaviors, such as colonization of

animal hosts [9,16–20].

We also propose that MuGENT-SSR can be utilized to engineer strains suitable for

serological typing and vaccination. In *C. jejuni* NCTC 11168, the majority of polyG/C

SSRs are clustered in genomic regions encoding proteins involved in the biosynthesis of

cell-surface antigenic determinants, including CPS and LOS [5]. Thus, phase variation

makes these antigens less desirable as serodeterminants and vaccine candidates. Penner

serotyping is often utilized to investigate epidemiological associations with Guillain–

Barré syndrome, but its low typeability is currently problematic [55]. The generation of

phase-locked strains by MuGENT-SSR may overcome these defects and stabilize their

antigenicity, thereby increasing the supply of stable serotyping antisera and vaccines.

Furthermore, PPT-decoding of Penner serotypes provides a reliable approach for

identifying antigenic determinant genes and improving or developing DNA-based

typing methods that do not rely on CPS expression [56].

Methods

Bacterial culture conditions

C. jejuni strains used in this study (Table 1) were routinely cultured for 24–48 h at 42°C on BHI (Becton Dickinson) plates containing 1.3% agar (Kyokuto) under microaerophilic conditions using Mitsubishi Anaeropack MicroAero gas generator packs (Mitsubishi Gas). Motility was determined by culturing the cells on BHI plates containing 0.3% agar. For liquid culture, fresh single colonies grown on BHI agar plates were inoculated into 5 ml of BHI broth in 25 ml volumetric test tubes and then cultured overnight at 42°C with reciprocal shaking at 160 rpm in a Taitec Precyto MG-71M-A Obligatory Anaerobe Culture System (Taitec), which can efficiently create a microaerophilic atmosphere by actively aerating gases into each test tube. The aeration conditions in each test tube were set to 5% O_2, 10% CO_2, and 85% N_2 at a constant flow rate (10 ml/min). The antibiotics used were 25 µg/ml Cm, 50 µg/ml Km, and 10 µg/ml Sm. If necessary, XS (Sigma) was added to the BHI agar plates at a final concentration of 100 µg/ml.

DNA manipulation

PCR amplification of DNA was performed using a LifeECO thermal cycler (version 1.04, Bioer Technology) and Quick Taq® HS DyeMix DNA polymerase (Toyobo). Sanger sequencing was performed by Fasmac. PCR products were purified using a High Pure PCR Product Purification Kit (Roche). Customized oligonucleotide primers (S1 Table) were purchased from Fasmac and Hokkaido System Science. Chromosomal and plasmid DNA molecules were extracted with a DNeasy Blood and Tissue Kit (Qiagen) and a High Pure Plasmid Isolation Kit (Roche), respectively.

Construction of pSYC-*cat* and pSYC-*kan*

The pSYC-*cat* and pSYC-*kan* plasmids (Table 1) were constructed by Fasmac as follows: DNA fragments containing the Cm-resistance gene (*cat*) (GenBank accession number M35190.1, nucleotides 1–1,034) and the Km-resistance gene (*kan*) (GenBank accession number M26832.1, nucleotides 1–1,426) from *C. coli* [57,58] were synthesized and cloned into the EcoRV site of pUCFa (Fasmac) to generate pSYC-*cat* and pSYC-*kan*, respectively.

Natural transformation of *C. jejuni* cells using PCR products

Genetically engineered *C. jejuni* strains were constructed by performing natural transformation [26] optimized to use PCR products as donor DNA. Briefly, the procedures include three processes: (1) PCR amplification of donor DNA with methylation sites, (2) methylation of the amplified DNA, and (3) transformation of *C. jejuni* with methylated DNA.

PCR amplification of donor DNA with methylation sites. The donor DNA fragments used for natural transformation were amplified using specific primers and templates. The typical amplified fragment contained an antibiotic-resistance gene (or mutation) with 5′- and 3′-flanking regions homologous to the target site and an EcoRI-recognition sequence (GAATTC) on both sides. DNA was amplified via splicing by overlap extension PCR, which includes a two-step amplification process [59,60]. The first step independently generated two or three fragments with overlapping sequences. In the second step, the splicing by overlap extension PCR fragments amplified in the first step are ligated. The thermal cycling program was as follows: 94°C for 5 min, followed by 30 cycles of 94°C

for 30 s, 55°C for 30 s, and 68°C for 1 min per kb. S2 Table shows the specific

combinations of template DNA and primer pairs used for PCR amplification.

Methylation of donor DNA. DNA methylation was performed using a 25 μl mixture

containing 20 nM DNA, EcoRI methyltransferase reaction buffer (New England Biolabs;

50 mM Tris-HCl, 50 mM NaCl, 10 mM ethylenediaminetetraacetic acid, pH 8.0), 80 μM

S-adenosylmethionine (New England Biolabs), and 40 units of EcoRI methyltransferase

(New England Biolabs). After incubation for 2.5 h at 37°C, the reaction was terminated

by incubation for 20 min at 65°C. Methylated DNA was purified using the High Pure PCR

Product Purification Kit and dissolved in 100 μl H_2O.

Natural transformation. Overnight cultures of *C. jejuni* cells were diluted 1/50 in 5 ml

BHI broth and grown until they reached an optical density and 600 nm (OD_{600}) of

approximately 0.15. Each culture (900 μl) was mixed with methylated DNA (100 μl) and

then statically incubated overnight in a 25 ml volumetric plastic test tube. On the

following day, 10-fold serial dilutions of the culture were spread onto BHI agar plates,

with or without antibiotics. The transformation frequency was defined as the number of

antibiotic-resistant CFUs divided by the total number of CFUs.

Evaluation of natural cotransformation in *C. jejuni*

Natural cotransformation in *C. jejuni* was examined using two different PCR fragments,

Δ*flaA*::*kan* and *rpsL*$^+$ (see S2 Table). The former fragment (Δ*flaA*::*kan*-1 or Δ*flaA*::*kan*-

2) had 1,000-bp regions of homology, whereas the latter had 1,000-bp or 2,000-bp regions

of homology (*rpsL*$^+$-8-1, *rpsL*$^+$-9-1, *rpsL*$^+$-8-2, or *rpsL*$^+$-9-2). The Δ*flaA*::*kan* and *rpsL*$^+$

fragments were independently methylated and then mixed for purification as described

above. After transformation of SY1003 and SY2003 with these methylated DNA

fragments, the reactions were spread onto BHI agar plates containing Km. One hundred

Km^R transformants were replica plated onto BHI agar plates containing Sm. The

cotransformation frequency (%) was calculated as follows: $100 \times$ number of Sm^S CFUs

in 100 Km^R CFUs.

MuGENT-SSR

MuGENT-SSR was performed with *C. jejuni* cells, as follows. After independently

methylating multiple unselected fragments, including eight fragments used for editing of

the eight polyG sites in the biosynthetic LOS and CPS gene clusters in NCTC11168, as

well as a single selected fragment (either Δ*flaA*::*kan*-1 or Δ*flaA*::*cat*-1), a mixture of these

fragments was prepared as described above. Unselected fragments had 1,500–2,000-bp

regions of homology, whereas the selected fragment had a 1,000-bp region of homology

(S2 and S3 Tables). After transformation with the DNA mixture, the bacteria were plated

onto BHI agar plates containing the antibiotic corresponding to the resistance protein

encoded in the selected fragment. Colonies on selective agar plates were used for MASC

PCR (described below). The remaining colonies (3,000–5,000 CFUs) were suspended in

BHI broth and diluted in 5 ml BHI broth to an OD_{600} of 0.05. After the cells were grown

to an OD_{600} of 0.15, the next cycle of MuGENT was performed using unselected

fragments and a selected fragment with a different resistance gene from that used in the

previous cycle. MuGENT cycles were repeated until editing was completed. Genome

editing was ultimately verified by nucleotide sequencing using specific primers (see the

PPT-Seq section below). If necessary, the Δ*flaA* mutation maintained during MuGENT

was reverted to the wild-type allele by transformation using the *flaA⁺* fragment with a

1,000-bp region of homology. After the transformation, colonies showing motility (and

sensitivity to the antibiotic used for selection) were chosen for further study.

MASC PCR

During each cycle of MuGENT, 40 colonies were inoculated in 100 μl of BHI broth

in a 96-wel plate and incubated overnight. Each PCR mixture (20 μl) contained 2 μl of

the bacterial culture, 2 μL of 2.5 μM primer mix (one of four primer mixes, including Mix

ON1, Mix ON2, Mix OFF1, and Mix OFF2; see S4 Table), 6 μl of H_2O, and 10 μl of

Quick Taq® HS DyeMix. The following thermal cycling program was used for MASC

PCR: 94°C for 5 min, followed by 35 cycles of 94°C for 30 s, 62.6°C for 30 s, and 68°C

for 3 min. If necessary, several single colonies were isolated from positive colonies (to

eliminate contamination by unedited clones) and further assessed for genome editing by

MASC PCR.

PPT-Seq

The eight phase-variable genes located in the LOS- and CPS-biosynthetic gene clusters

in NCTC11168 were amplified by PCR, and the repeat numbers of the polyG SSR tracts

were determined by nucleotide sequencing. S5 Table shows the specific combinations of

target genes and primers used for PPT-Seq.

Penner serotyping

A single colony was streaked on the surface of a horse blood agar plate (Kyokuto) and

incubated for 48 h at 42°C to determine the Penner serotype [61]. After the resulting

colonies grown on the plate were suspended in 250 µL of 0.9% NaCl, serotyping was

performed using *Campylobacter* antisera (Denka), including a commercial 25 antisera set,

and a reagent for preparing sensitized blood cells (Denka) according to the manufacturer's

instructions. Phenotypic stability was assessed by repeating the subculture cycle and

serotyping. Serotyping was performed using six single colonies for each cycle.

Measurement of mutation rates

The mutation rates of the phase-variable *cj1426* gene and its phase-locked variants

were determined as described previously [62], with some modifications. *C. jejuni* strains

carrying *astA* to *cj1426* translational fusions were streaked on BHI agar plates containing

XS. During growth on the medium, blue colonies were in the *cj1426*-ON phase, whereas

white colonies were in the *cj1426*-OFF phase. Single blue or white colonies were picked

using a micropipette tip were resuspended in 500 µL BHI, after which 250 µl of 10^4-, 10^5-,

and 10^6-fold dilutions were spread onto BHI agar plates containing XS. The number of

variant colonies that switched to different phases and the total numbers of colonies were

counted. Ten independent single colonies were examined for each strain and phase. To

estimate the mutation rate, the total number of colonies was averaged for all single

colonies tested, and the median value for the frequency of variants per colony was

calculated. The mutation rate was calculated using the following equation: $\mu =$

$0.4343f/log(N\mu)$, where μ is the mutation rate, f is the median frequency, and N is

the average population size [52]. The μ value was determined by solving the equation

using the Goal Seek function in Microsoft Excel.

Acknowledgments

We thank Jiro Mitobe for helpful suggestions.

Funding

This study was supported by AMED (grant numbers 18fk0108065j0201,

19fk0108065j0202, 20fk0108065j0203, and 21fk0108611j0201).

References

1.   Levinson G, Gutman GA. Slipped-strand mispairing: a major mechanism for

DNA sequence evolution. Mol Biol Evol. 1987;4: 203–221.

doi:10.1093/oxfordjournals.molbev.a040442.

2.   Moxon R, Bayliss C, Hood D. Bacterial contingency loci: the role of simple

sequence DNA repeats in bacterial adaptation. Annu Rev Genet. 2006;40: 307–

333. doi:10.1146/annurev.genet.40.110405.090442.

3.   Bayliss CD. Determinants of phase variation rate and the fitness implications of

differing rates for bacterial pathogens and commensals. FEMS Microbiol Rev.

2009;33: 504–520. doi:10.1111/j.1574-6976.2009.00162.x.

4.   Bidmos FA, Bayliss CD. Genomic and Global Approaches to Unravelling How

Hypermutable Sequences Influence Bacterial Pathogenesis. Pathogens. 2014;3:

164–184. doi:10.3390/pathogens3010164.

5.   Parkhill J, Wren BW, Mungall K, Ketley JM, Churcher C, Basham D, et al. The

genome sequence of the food-borne pathogen Campylobacter jejuni reveals

hypervariable sequences. Nature. 2000;403: 665–668. doi:10.1038/35001088.

6.   Hermans D, Van Deun K, Martel A, Van Immerseel F, Messens W, Heyndrickx

M, et al. Colonization factors of Campylobacter jejuni in the chicken gut. Vet

Res. 2011;42: 82. doi:10.1186/1297-9716-42-82.

7.   Young KT, Davis LM, Dirita VJ. Campylobacter jejuni: molecular biology and

pathogenesis. Nat Rev Microbiol. 2007;5: 665–679. doi:10.1038/nrmicro1718.

8.   McCarthy N, Giesecke J. Incidence of Guillain-Barré syndrome following

infection with Campylobacter jejuni. Am J Epidemiol. 2001;153:610–614. doi:

10.1093/aje/153.6.610.

9.   Bayliss CD, Bidmos FA, Anjum A, Manchev VT, Richards RL, Grossier JP, et al.

Phase variable genes of Campylobacter jejuni exhibit high mutation rates and

specific mutational patterns but mutability is not the major determinant of

population structure during host colonization. Nucleic Acids Res. 2012;40: 5876–

5889. doi:10.1093/nar/gks246.

10.   Lango-Scholey L, Aidley J, Woodacre A, Jones MA, Bayliss CD. High

Throughput Method for Analysis of Repeat Number for 28 Phase Variable Loci

of Campylobacter jejuni Strain NCTC11168. PLOS ONE. Gerlach RG, editor.

2016;11: e0159634. doi:10.1371/journal.pone.0159634.

11.   Guerry P, Szymanski CM, Prendergast MM, Hickey TE, Ewing CP, Pattarini DL,

et al. Phase variation of Campylobacter jejuni 81-176 lipooligosaccharide affects

ganglioside mimicry and invasiveness in vitro. Infect Immun. 2002;70: 787–793.

doi:10.1128/iai.70.2.787-793.2002.

12.   Prendergast MM, Tribble DR, Baqar S, Scott DA, Ferris JA, Walker RI, et al. In

vivo phase variation and serologic response to lipooligosaccharide of

Campylobacter jejuni in experimental human infection. Infect Immun. 2004;72:

916–922. doi:10.1128/iai.72.2.916-922.2004.

13.   Linton D, Gilbert M, Hitchen PG, Dell A, Morris HR, Wakarchuk WW, et al.

Phase variation of a β-1,3 galactosyltransferase involved in generation of the

ganglioside GM1-like lipo-oligosaccharide of Campylobacter jejuni. Mol

Microbiol. 2000;37: 501–514. doi:10.1046/j.1365-2958.2000.02020.x.

14.   Ashgar SSA, Oldfield NJ, Wooldridge KG, Jones MA, Irving GJ, Turner DPJ, et

al. CapA, an autotransporter protein of Campylobacter jejuni, mediates

association with human epithelial cells and colonization of the chicken gut. J

Bacteriol. 2007;189: 1856–1865. doi:10.1128/JB.01427-06.

15.   Hendrixson DR. A phase-variable mechanism controlling the Campylobacter

jejuni FlgR response regulator influences commensalism. Mol Microbiol.

2006;61: 1646–1659. doi:10.1111/j.1365-2958.2006.05336.x.

16.   Wanford JJ, Lango-Scholey L, Nothaft H, Hu Y, Szymanski CM, Bayliss CD.

Random sorting of Campylobacter jejuni phase variants due to a narrow

bottleneck during colonization of broiler chickens. Microbiology (Reading).

2018;164: 896–907. doi:10.1099/mic.0.000669.

17.   Jerome JP, Bell JA, Plovanich-Jones AE, Barrick JE, Brown CT, Mansfield LS.

Standing Genetic Variation in Contingency Loci Drives the Rapid Adaptation of

Campylobacter jejuni to a Novel host. PLOS ONE 2011;6: e16399.

doi:10.1371/journal.pone.0016399.

18.   Kim JS, Artymovich KA, Hall DF, Smith EJ, Fulton R, Bell J, et al. Passage of

Campylobacter jejuni through the chicken reservoir or mice promotes phase

variation in contingency genes Cj0045 and Cj0170 that strongly associates with

colonization and disease in a mouse model. Microbiology (Reading). 2012;158:

1304–1316. doi:10.1099/mic.0.057158-0.

19.   Wilson DL, Rathinam VAK, Qi W, Wick LM, Landgraf J, Bell JA, et al. Genetic

diversity in Campylobacter jejuni is associated with differential colonization of

broiler chickens and C57BL/6J IL10-deficient mice. Microbiology (Reading).

2010;156: 2046–2057. doi:10.1099/mic.0.035717-0.

20.   Thomas DK, Lone AG, Selinger LB, Taboada EN, Uwiera RRE, Abbott DW, et

al. Comparative Variation within the Genome of Campylobacter jejuni NCTC

11168 in Human and Murine hosts. PLOS ONE Freitag NE, et al., host. 2014;9:

663   e88229. doi:10.1371/journal.pone.0088229.

664   21.   Wang HH, Isaacs FJ, Carr PA, Sun ZZ, Xu G, Forest CR, et al. Programming

665   cells by multiplex genome engineering and accelerated evolution. Nature.

666   2009;460: 894–898. doi:10.1038/nature08187.

667   22.   Jiang W, Bikard D, Cox D, Zhang F, Marraffini LA. RNA-guided editing of

668   bacterial genomes using CRISPR-Cas systems. Nat Biotechnol. 2013;31: 233–

669   239. doi:10.1038/nbt.2508.

670   23.   Johnston C, Martin B, Fichant G, Polard P, Claverys JP. Bacterial transformation:

671   distribution, shared mechanisms and divergent control. Nat Rev Microbiol.

672   2014;12: 181–196. doi:10.1038/nrmicro3199.

673   24.   Lorenz MG, Wackernagel W. Bacterial gene transfer by natural genetic

674   transformation in the environment. Microbiol Rev. 1994;58: 563–602. DOI:

675   10.1128/MR.58.3.563-602.1994.

676   25.   Wang Y, Taylor DE. Natural transformation in Campylobacter species. J

677   Bacteriol. 1990;172: 949–955. doi:10.1128/jb.172.2.949-955.1990.

678   26.   Beauchamp JM, Leveque RM, Dawid S, DiRita VJ. Methylation-dependent DNA

679   discrimination in natural transformation of Campylobacter jejuni. Proc Natl Acad

680   Sci U S A. 2017;114: E8053–E8061. doi:10.1073/pnas.1703331114.

681   27.   Dalia AB, McDonough E, Camilli A. Multiplex genome editing by natural

682   transformation. Proc Natl Acad Sci U S A. 2014;111: 8937–8942.

683   doi:10.1073/pnas.1406478111.

684   28.   Mell JC, Lee JY, Firme M, Sinha S, Redfield RJ. Extensive cotransformation of

685   natural variation into chromosomes of naturally competent Haemophilus

686   influenzae. G3 Amp. 2014;4;# 4; Genes|Genomes|Genetics: 717–731.

doi:10.1534/g3.113.009597.

29. Erickson RJ, Copeland JC. Congression of unlinked markers and genetic

mapping in the transformation of Bacillus subtilis 168. Genetics. 1973;73: 13–21.

Available: http://www.ncbi.nlm.nih.gov/pubmed/4631599.

30. Olkkola S, Juntunen P, Heiska H, Hyytiäinen H, Hänninen ML. Mutations in the

rpsL gene are involved in streptomycin resistance in Campylobacter coli. Microb

Drug Resist. 2010;16: 105–110. doi:10.1089/mdr.2009.0128.

31. Skirrow MB. Campylobacter enteritis: a "new" disease. Br Med J. 1977;2: 9–11.

doi:10.1136/bmj.2.6078.9.

32. Korlath JA, Osterholm MT, Judy LA, Forfang JC, Robinson RA. A point-source

outbreak of campylobacteriosis associated with consumption of raw milk. J Infect

Dis. 1985;152: 592–596. doi:10.1093/infdis/152.3.592.

33. Watt VM, Ingles CJ, Urdea MS, Rutter WJ. Homology requirements for

recombination in Escherichia coli. Proc Natl Acad Sci U S A. 1985;82: 4768–

4772. doi:10.1073/pnas.82.14.4768.

34. Wang HH, Church GM. Multiplexed genome engineering and genotyping

methods applications for synthetic biology and metabolic engineering. Methods

Enzymol. 2011;498: 409–426. doi:10.1016/B978-0-12-385120-8.00018-8.

35. Burnham PM, Hendrixson DR. Campylobacter jejuni: collective components

promoting a successful enteric lifestyle. Nat Rev Microbiol. 2018;16: 551–565.

doi:10.1038/s41579-018-0037-9.

36. Maue AC, Mohawk KL, Giles DK, Poly F, Ewing CP, Jiao Y, et al. The

polysaccharide capsule of Campylobacter jejuni modulates the host immune

response. Infect Immun. 2013;81: 665–672. doi:10.1128/IAI.01008-12.

37.   Sørensen MCH, van Alphen LB, Harboe A, Li J, Christensen BB, Szymanski

712   CM, et al. Bacteriophage F336 Recognizes the Capsular Phosphoramidate

713   Modification of Campylobacter jejuni NCTC11168. J Bacteriol. 2011;193: 6742–

714   6749. doi:10.1128/JB.05276-11.

715   38.   McNally DJ, Lamoureux MP, Karlyshev AV, Fiori LM, Li J, Thacker G, et al.

716   Commonality and Biosynthesis of the O-Methyl Phosphoramidate Capsule

717   Modification in Campylobacter jejuni. J Biol Chem. 2007;282: 28566–28576.

718   doi:10.1074/jbc.M704413200.

719   39.   van Alphen LB, Wenzel CQ, Richards MR, Fodor C, Ashmus RA, Stahl M, et al.

720   Biological Roles of the O-Methyl Phosphoramidate Capsule Modification in

721   Campylobacter jejuni. PLOS ONE. Skurnik M, et al., editor. 2014;9: e87051.

722   doi:10.1371/journal.pone.0087051.

723   40.   Yuki N, Susuki K, Koga M, Nishimoto Y, Odaka M, Hirata K, et al. Carbohydrate

724   mimicry between human ganglioside GM1 and Campylobacter jejuni

725   lipooligosaccharide causes Guillain-Barre syndrome. Proc Natl Acad Sci U S A.

726   2004;101: 11404–11409. doi:10.1073/pnas.0402391101.

727   41.   Karlyshev AV, Linton D, Gregson NA, Lastovica AJ, Wren BW. Genetic and

728   biochemical evidence of a Campylobacter jejuni capsular polysaccharide that

729   accounts for Penner serotype specificity. Mol Microbiol. 2000;35: 529–541.

730   doi:10.1046/j.1365-2958.2000.01717.x.

731   42.   Moran AP, Penner JL. Serotyping of Campylobacter jejuni based on heat-stable

732   antigens: Relevance, molecular basis and implications in pathogenesis. J Appl

733   Microbiol. 1999;86: 361–377. doi:10.1046/j.1365-2672.1999.00713.x.

43.   Aidley J, Sørensen MCH, Bayliss CD, Brøndsted L. Phage exposure causes

dynamic shifts in the expression states of specific phase-variable genes of

Campylobacter jejuni. Microbiology (Reading). 2017;163: 911–919.

doi:10.1099/mic.0.000470.

44.   Pequegnat B, Laird RM, Ewing CP, Hill CL, Omari E, Poly F, et al. Phase-

Variable Changes in the Position of O-Methyl Phosphoramidate Modifications on

the Polysaccharide Capsule of Campylobacter jejuni Modulate Serum Resistance.

J Bacteriol. 2017;199. doi:10.1128/JB.00027-17.

45.   Sørensen MCH, van Alphen LB, Fodor C, Crowley SM, Christensen BB,

Szymanski CM, et al. Phase Variable Expression of Capsular Polysaccharide

Modifications Allows Campylobacter jejuni to Avoid Bacteriophage Infection in

Chickens. Front Cell Infect Microbiol. 2012;2: 11.

doi:10.3389/fcimb.2012.00011.

46.   Bacon DJ, Szymanski CM, Burr DH, Silver RP, Alm RA, Guerry P. A phase-

variable capsule is involved in virulence of Campylobacter jejuni 81-176. Mol

Microbiol. 2001;40: 769–777. doi:10.1046/j.1365-2958.2001.02431.x.

47.   Linton D, Karlyshev AV, Wren BW. Deciphering Campylobacter jejuni cell

surface interactions from the genome sequence. Curr Opin Microbiol. 2001;4:

35–40. doi:10.1016/S1369-5274(00)00161-2.

48.   Guerry P, Poly F, Riddle M, Maue AC, Chen YH, Monteiro MA. Campylobacter

Polysaccharide Capsules: Virulence and Vaccines. Front Cell Infect Microbiol.

2012;2: 7. doi:10.3389/fcimb.2012.00007.

49.   Yao R, Guerry P. Molecular cloning and site-specific mutagenesis of a gene

involved in arylsulfatase production in Campylobacter jejuni. J Bacteriol.

1996;178: 3335–3338. doi:10.1128/jb.178.11.3335-3338.1996.

50.   Sternberg MJE, Tamaddoni-Nezhad A, Lesk VI, Kay E, Hitchen PG, Cootes A, et

al. Gene Function Hypotheses for the Campylobacter jejuni Glycome Generated

by a Logic-Based Approach. J Mol Biol. 2013;425: 186–197.

doi:10.1016/j.jmb.2012.10.014.

51.   Hendrixson DR, DiRita VJ. Transcription of σ54-dependent but not σ28-

dependent flagellar genes in Campylobacter jejuni is associated with formation of

the flagellar secretory apparatus. Mol Microbiol. 2003;50: 687-702. doi:

10.1046/j.1365-2958.2003.03731.x.

52.   Drake JW. A constant rate of spontaneous mutation in DNA-based microbes. Proc

Natl Acad Sci U S A. 1991;88: 7160–7164. doi:10.1073/pnas.88.16.7160.

53.   St Michael F, Szymanski CM, Li J, Chan KH, Khieu NH, Larocque S, et al. The

structures of the lipooligosaccharide and capsule polysaccharide of

Campylobacter jejuni genome sequenced strain NCTC 11168. Eur J Biochem.

2002;269: 5119–5136. doi:10.1046/j.1432-1033.2002.03201.x.

54.   Szymanski CM, Michael FS, Jarrell HC, Li J, Gilbert M, Larocque S, et al.

Detection of conserved N-linked glycans and phase-variable lipooligosaccharides

and capsules from campylobacter cells by mass spectrometry and high resolution

magic angle spinning NMR spectroscopy. J Biol Chem. 2003;278: 24509–24520.

doi:10.1074/jbc.M301273200.

55.   Pike BL, Guerry P, Poly F. Global Distribution of Campylobacter jejuni Penner

Serotypes: A Systematic Review. PLOS ONE. 2013;8: e67375.

doi:10.1371/journal.pone.0067375.

56.   Poly F, Serichantalergs O, Kuroiwa J, Pootong P, Mason C, Guerry P, et al.

Updated Campylobacter jejuni Capsule PCR Multiplex Typing System and Its

Application to Clinical Isolates from South and Southeast Asia. PLOS ONE.

2015;10: e0144349. doi:10.1371/journal.pone.0144349.

57. Wang Y, Taylor DE. Chloramphenicol resistance in Campylobacter coli:

nucleotide sequence, expression, and cloning vector construction. Gene. 1990;94:

23–28. doi:10.1016/0378-1119(90)90463-2.

58. Trieu-Cuot P, Gerbaud G, Lambert T, Courvalin P. In vivo transfer of genetic

information between gram-positive and gram-negative bacteria. EMBO J.

1985;4: 3583–3587. DOI: 10.1002/j.1460-2075.1985.tb04120.x.

59. Yamamoto S, Izumiya H, Morita M, Arakawa E, Watanabe H. Application of λ

Red recombination system to Vibrio cholerae genetics: Simple methods for

inactivation and modification of chromosomal genes. Gene. 2009;438: 57–64.

doi:10.1016/j.gene.2009.02.015.

60. Horton RM, Hunt HD, Ho SN, Pullen JK, Pease LR. Engineering hybrid genes

without the use of restriction enzymes: gene splicing by overlap extension. Gene.

1989;77: 61–68. doi:10.1016/0378-1119(89)90359-4.

61. Penner JL, Hennessy JN. Passive hemagglutination technique for serotyping

Campylobacter fetus subsp. jejuni on the basis of soluble heat-stable antigens. J

Clin Microbiol. 1980;12: 732–737. doi:10.1128/JCM.12.6.732-737.1980.

62. De Bolle X, Bayliss CD, Field D, van de Ven T, Saunders NJ, Hood DW, et al.

The length of a tetranucleotide repeat tract in Haemophilus influenzae determines

the phase variation rate of a gene with homology to type III DNA

methyltransferases. Mol Microbiol. 2000;35: 211–222. doi:10.1046/j.1365-

2958.2000.01701.x.

Supporting information captions

S1 Fig. Penner serotyping of naturally occurring and phase-locked NCTC11168 variants. Phenotypic stability was examined during successive subcultures, which highlights only the results obtained using anti-serotype B antiserum. Red-numbered colonies were typeable, and black-numbered colonies were untypeable.

S2 Fig. MASC PCR of transformants following successive cycles of MuGENT-SSR.

S3 Fig. Penner serotyping of phase-locked NCTC11168 variants.

S4 Fig. Putative modification patterns of the CPS repeat unit in naturally occurring and phase-locked NCTC11168 variants. Figures were modified from illustrations published in Sternberg et al, J Mol Biol 2013 (425) 186-197.

S1 Table. Primers used in this study.

S2 Table. Specific combinations of template DNA and primers used to amplify donor DNA templates.

S3 Table. Specific combinations of donor DNA molecules and recipient strains used for natural transformation.

S4 Table. Primer mixes used for MASC PCR.

S5 Table. Specific combinations of target genes and primers in PPT-Seq.

S6 Table. Primer sets used for allele-specific PCR and sequencing of *cj1426*::*astA* translational fusions.

S7 Table. Denka antisera used for Penner serotyping.

Fig 1

	Transformation frequency (Sm^R CFUs/total CFUs)			
	NCTC11168		81-176	
	WT	$\Delta recA$	WT	$\Delta recA$
rpsL^{K88R}-1 (50 b)	$<1.5 \times 10^{-8}$	ND	$<2.1 \times 10^{-8}$	ND
rpsL^{K88R}-2 (100 b)	$<1.4 \times 10^{-8}$	ND	$<2.2 \times 10^{-8}$	ND
rpsL^{K88R}-3 (500 b)	$1.8 \pm 2 \times 10^{-8}$	ND	$1.4 \pm 1 \times 10^{-8}$	ND
rpsL^{K88R}-4 (500 b)	$1.2 \pm 1 \times 10^{-6}$	ND	$2.1 \pm 2 \times 10^{-5}$	ND
rpsL^{K88R}-5 (500 b)	$3.2 \pm 3 \times 10^{-6}$	ND	$3.1 \pm 1 \times 10^{-5}$	ND
rpsL^{K88R}-6 (1 kb)	$1.2 \pm 1 \times 10^{-5}$	ND	$4.0 \pm 1 \times 10^{-5}$	ND
rpsL^{K88R}-7 (1 kb)	$9.0 \pm 1 \times 10^{-6}$	ND	$1.3 \pm 1 \times 10^{-4}$	ND
rpsL^{K88R}-8 (1 kb)	$2.0 \pm 1 \times 10^{-5}$	ND	$1.3 \pm 1 \times 10^{-4}$	ND
rpsL^{K88R}-9 (2 kb)	$5.0 \pm 2 \times 10^{-5}$	$< 1.6 \times 10^{-7}$	$1.2 \pm 1 \times 10^{-4}$	$<1.4 \times 10^{-7}$
Unmethylation of rpsL^{K88R}-5	$2.7 \pm 1 \times 10^{-8}$	ND	$1.2 \pm 3 \times 10^{-8}$	ND
No addition of DNA	$<1.2 \times 10^{-8}$	ND	$<1.9 \times 10^{-8}$	ND

Fig 2

Fig 3

Fig 4

A

$cj1139^{ON}$ (encoding 303 aa)

Locked ON
GGCGGAGG
D M G G G G K I D L L C F E A F I T N A K K S I K K L N
GATATGGGTGGGGGGGGTAAAATTGATTTGTTGTGTTTTGAAGCTTTTATTACCAATGCAAAAAAATCAATAAAAAAATTAAAT

-1 G   +1 G

GATATGGGTGGGGGGGTAAAATTGATTTGTTGTGTTTTGAAGCTTTTATTACCAATGCAAAAAAATCAATAAAAAAATTAAAT
D M G G G G V K L I C C V L K L L L P M Q K N Q *
GGAGGCG
Locked OFF

$cj1139^{OFF}$ (encoding 130 aa + extra 18 aa)

B

Fig 5

NCTC11168 variant	Typeability	Number of polyG tract repeats in phase-variable genes								PPT
		cj1139	*cj1144*	*cj1420*	*cj1421*	*cj1422*	*cj1426*	*cj1429*	*cj1437*	
		8	**7**	**9**	**9**	**9**	**10**	**10**	**9**	
SC1-1	No	9	**10**	**9**	10	**9**	**10**	11	**9**	109
SC1-2	No	9	**10**	**9**	10	**9**	**10**	**10**	**9**	111
SC1-3	No	**8**	9	**9**	**9**	**9**	**10**	11	**9**	189
SC1-4	**Yes**	9	9	**9**	10	10	**10**	11	**9**	**37**
SC1-5	No	**8**	9	**9**	**9**	**9**	**10**	**10**	**9**	191
SC1-6	No	9	**10**	**9**	10	**9**	**10**	11	**9**	109
SC2-1	**Yes**	9	9	**9**	10	10	**10**	9	**9**	**37**
SC2-2	**Yes**	9	9	**9**	10	10	**10**	11	**9**	**37**
SC2-3	**Yes**	9	9	**9**	10	10	**10**	11	**9**	**37**
SC2-4	No	9	9	8	10	10	9	11	**9**	1
SC2-5	No	9	9	8	10	10	11	11	**9**	1
SC2-6	**Yes**	9	9	**9**	10	10	**10**	11	**9**	**37**
Locked-0	No	Locked OFF	Locked OFF	Locked OFF	Locked OFF	Locked OFF	Locked OFF	Locked OFF	Locked OFF	0
Locked-255	No	**Locked ON**	**Locked ON**	**Locked ON**	**Locked ON**	**Locked ON**	**Locked ON**	**Locked ON**	**Locked ON**	255
Locked-37	**Yes**	Locked OFF	Locked OFF	**Locked ON**	Locked OFF	Locked OFF	**Locked ON**	Locked OFF	**Locked ON**	**37**
Locked-36	**Yes**	Locked OFF	Locked OFF	**Locked ON**	Locked OFF	Locked OFF	**Locked ON**	Locked OFF	Locked OFF	**36**
Locked-5	**Yes**	Locked OFF	Locked OFF	Locked OFF	Locked OFF	Locked OFF	**Locked ON**	Locked OFF	**Locked ON**	**5**
Locked-33	No	Locked OFF	Locked OFF	**Locked ON**	Locked OFF	Locked OFF	Locked OFF	Locked OFF	**Locked ON**	33
Locked-4	**Yes**	Locked OFF	Locked OFF	Locked OFF	Locked OFF	Locked OFF	**Locked ON**	Locked OFF	Locked OFF	**4**
Locked-1	No	Locked OFF	Locked OFF	Locked OFF	Locked OFF	Locked OFF	Locked OFF	Locked OFF	**Locked ON**	1
Locked-32	No	Locked OFF	Locked OFF	**Locked ON**	Locked OFF	Locked OFF	Locked OFF	Locked OFF	Locked OFF	32