2   <u>*Running title:*</u> Dispersal and biome shifts in *Pterocarpus*

The diversification of *Pterocarpus* (Leguminosae: Papilionoideae)

was influenced by biome-switching and infrequent long-distance dispersal

Rowan J. Schley[1*6], Ming Qin[2], Mohammad Vatanparast[3,4], Panagiota Malakasi[1], Manuel de la

Estrella[5], Gwilym Lewis[1], Bente Klitgård[1]

[1]*Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK*

[2]*South China Botanical Garden, Chinese Academy of Sciences, 1190 Tianyuan Rd, Tianhe, Guangzhou, Guangdong, China*

[3]*Department of Geosciences and Natural Resource Management, University of Copenhagen, 1958, Denmark*

[4]*Department of Botany, National Museum of Natural History, Smithsonian Institution, MRC 166, Washington, DC 20560, USA*

[5]*Departamento de Botánica, Ecología y Fisiología Vegetal, Facultad de Ciencias, Universidad de Córdoba, 14014, Córdoba, Spain*

[6]*University of Exeter, North Park Road, Exeter, EX4 4QE, UK*

*Corresponding author. *E-mail address:* <u>rowan.schley@gmail.com</u> (R. J. Schley).
17

Acknowledgements

All the authors would like to thank the PAFTOL program at RBG, Kew for funding the sequencing of

Pterocarpus clade accessions. MQ's lab visit was funded by the National Natural Science Foundation

of China (grant no. 31670193) and the Science and Technology Planning Project of Guangdong

Province (grant no. 2019B030316020).

<u>Abstract</u>

Aim

Phenotypes which evolved for dispersal over ecological timescales may lead to significant
macroevolutionary consequences, such as infrequent long-distance dispersal and diversification in
novel biomes. We aimed to reconstruct the phylogenetic history of *Pterocarpus* (Leguminosae/
Fabaceae) to assess whether seed dispersal phenotypes and biome switching explain the current
biogeographical patterns of this group.

Location

Pantropical

Taxon

The Pterocarpus clade, particularly *Pterocarpus* (Leguminosae/Fabaceae)

Methods

We sequenced ~300 nuclear loci captured using *Angiosperms*-353, a genomic 'bait set' for flowering
plants, from which we generated a time-calibrated phylogenomic tree. To corroborate this, we also
generated a time-calibrated phylogenetic tree from data-mined Sanger-sequencing data. We then
collated distribution data and fruit dispersal morphology traits to compare trait-dependent and trait-
independent biogeographical models, allowing us to assess whether dispersal traits influenced the
spatio-temporal evolution of *Pterocarpus*. Finally, using the results of these model tests, we estimated
the ancestral ranges and biomes of *Pterocarpus* species to better understand their biogeographical
history, and assessed the degree and direction of biome switching over the course of *Pterocarpus*'
diversification history.

Results

We recovered well-supported phylogenetic relationships within *Pterocarpus,* within which there were
two subclades – one Neotropical and the other Palaeotropical. Our divergence date estimates
suggested that *Pterocarpus* largely diversified from around 12 Ma, during the Miocene.
Trait-dependent biogeographical models were rejected for both range and biome evolution within
Pterocarpus, but models parameterising dispersal were supported. *Pterocarpus'* ancestral node shared
a range across the new-world and old-world tropics, followed by divergence into two clades, one
palaeotropical and one neotropical. Biome switching occurred most frequently into rainforest and
grassland.

Main conclusions

Overall, our analyses suggest that *Pterocarpus* underwent infrequent cross-continental dispersal and
establishment into novel biomes. While this was minimally impacted by fruit dispersal syndromes,
biome switching precipitated by long-distance dispersal and environmental change have played an
important role in diversification within *Pterocarpus* since the Miocene.

<u>*Keywords*</u>: Angiosperms, Biomes, Dispersal, Diversification, Legumes, Phylogenomics, Traits, Tropics

<u>Introduction</u>

Dispersal is a major process structuring the geographic ranges of organisms, both at ecological and
evolutionary timescales (Brown et al., 1996, Sanmartín & Ronquist, 2004). Indeed, adaptations which
evolved to promote dispersal locally may end up being significant evolutionary innovations,
influencing the biogeographical fate of lineages by allowing dispersal across continents and
subsequent diversification over millions of years (Klaus & Matzke, 2020).

While vicariance has a major role in explaining biogeographical patterns, the role of dispersal was
seen as being somewhat less significant (Sanmartín, 2012). However, there is now ample evidence
that dispersal, and especially trans-oceanic dispersal, plays a major role in the biogeographical
dynamics of a great number of terrestrial lineages (De Queiroz, 2005, Gillespie et al., 2012, Harris et
al., 2018). This is especially evident in plants due to the wide range of traits that they have evolved to
promote dispersal of their fruits, seeds and spores (Cousens et al., 2008). Seeds in particular have
developed a wide range of morphological traits which allow dispersal over long distances, including
wings and other aerodynamic structures (Cain et al., 2000), long dormancy periods (van der Pijl,
1982), thick seed coats and air pockets which render the seeds buoyant, thus facilitating dispersal
across oceans (Bellot et al., 2020, Renner, 2004). Multiple recent studies have highlighted that the
biogeographical distributions of modern plant lineages are largely explained by traits which they
evolved for effective dispersal in their ecological context (e.g. Podocarpaceae (Klaus & Matzke,
2020); Annonaceae (Onstein et al., 2019)) using trait-dependent biogeographical modes. Indeed, it has
been well established that a wide range of phenotypic traits have a significant impact on
macroevolutionary dynamics (e.g. Rabosky et al., 2014).

Given their superlative biological diversity, the tropics are one of the most important places to
examine the impact of dispersal on biogeography and diversification. Dispersal is particularly
important for terrestrial organisms in the tropics because they are subject to both geographical and
ecological constraints to their distribution. The tropics are largely separated into three zones
(Neotropical, African and Asian/Australasian) by large areas of ocean, bounded to the north and south
by increasingly unsuitable climates with increasing distance from the equator (Corlett & Primack,
2011, Morley, 2000). There is ample evidence of disjunct ranges across continents within tropical
organisms, many of which likely result from dispersal, particularly in plants (e.g., Arecaceae
(Eiserhardt et al., 2011), Podocarpaceae (Klaus & Matzke, 2020), Leguminosae/Fabaceae (Schrire et
al., 2005, Vatanparast et al., 2013), Vitaceae (Nie et al., 2012)).

Following dispersal, plants may subsequently encounter powerful selective pressures in contrasting habitats to those in which they evolved. This can drive adaptation, and eventually speciation, to these novel environments (Waser & Campbell, 2004). Biomes which span climatic extremes are found in relatively close proximity within the tropics, being defined by their ecological community composition (Pennington & Dick, 2004, Sanmartín, 2014) as well as by the physiognomy of their dominant plant species (Pennington & Ratter, 2006, Woodward et al., 2004). These biomes are often seen as independent evolutionary arenas (Hughes et al., 2013, Nürk et al., 2020, e.g. Pennington et al., 2009, Ringelberg et al., 2020). This biome concept reflects that environmental pressures select for species with similar functional attributes (Echeverría-Londoño et al., 2018), further implying that colonisation of contrasting biomes can promote speciation (Pennington & Dick, 2010). For example, many genera with species typical of seasonally dry tropical forest and savanna also contain species from rainforest (Pennington & Ratter, 2006, Pennington et al., 2000), suggesting that biome switching has driven ecological speciation.

The relative invasibility of certain biomes by immigrants may be higher than in others. Globally, lowland tropical ecosystems such as rainforest and savanna tend to be most permeable to colonisation by taxa from other biomes (Dexter et al., 2015), acting as 'lineage sinks' (Donoghue & Edwards, 2014, Pennington & Hughes, 2014) because plant communities in these environments experience high species turnover through exposure to drought and fire (da Costa et al., 2010, Pennington & Lavin, 2016). Such an imbalance in the direction of biome switching in tropical plant taxa is also likely to be an outcome of the relative size of the adaptive peak to be overcome by immigrants, coupled with the pre-existence of adaptive traits (Donoghue & Edwards, 2014). Again, drought and fire are likely to be modulators of the shape of such adaptive peaks, requiring specialised phenotypes to survive them (De Micco & Aronne, 2012, Midgley & Bond, 2013) and constraining plant evolution in the tropics (Olmstead, 2013).

Pantropical clades which occur in different biomes are an ideal system to study the biogeographical processes outlined above. The legume family (Leguminosae/Fabaceae) is replete with such clades, present in every terrestrial biome in the tropics (Lewis, G. P. et al., 2005) and dominant in many of them (de la Estrella et al., 2017, Gentry, 1988, ter Steege et al., 2013, ter Steege et al., 2020). The genus *Pterocarpus* Jacq. belongs to the largest subfamily of legumes (Papilionoideae) and consists of around 33 species (Klitgård & Lavin, 2005, Lewis, G. P., 1987, Rojo, 1972), many of which are important timber species or are used in traditional medicine (Saslis-Lagoudakis et al., 2011). This genus is diagnosed by either winged or corky (coriaceous) fruits adapted for dispersal by wind or water, respectively, and this is likely one reason why *Pterocarpus* is found pantropically (Klitgård & Lavin, 2005). *Pterocarpus* forms a part of the broader, pantropical 'Pterocarpus clade' which contains 22 genera and ~200 species, of which most are trees (Klitgård et al., 2013).

Given its pantropical distribution, its presence in many contrasting biomes and its diversity of dispersal syndromes, *Pterocarpus* is an ideal system for evolutionary studies of biogeography, biome-switching and dispersal within the tropics (Klitgård et al., 2013). Accordingly, we aim to examine the impact of dispersal traits on the diversification of *Pterocarpus* to provide insights into how trait evolution can structure the biogeographical distribution of species, and so influence the evolution of tropical diversity. We also aim to infer the patterns of biome switching which may result from such

dispersal events, to help clarify how shifts between biomes may accumulate or generate diversity. To
do this, we will reconstruct the relationships between species within the Pterocarpus clade using novel
next-generation sequencing techniques, and infer a time-calibrated phylogeny for *Pterocarpus* using
multiple approaches. Following this, we will test whether trait-dependent or trait-independent
biogeographical models best explain extant biogeographical patterns, both in terms of continental
realm and biome, and assess whether transitions between certain biomes are more frequent than
others. As such, we predict that:

1) Extant biogeographical patterns in *Pterocarpus* were significantly influenced by their seed
147   dispersal traits, and hence that a trait-dependent model will best describe their spatiotemporal
148   evolution.
149   2) Biome shifts will be more frequent into rainforests and savannas, given their ecologically
150   dynamic nature and resulting high invasibility by taxa dispersing into them.

Materials and Methods

Taxon sampling

Novel phylogenomic data were generated using target capture sequencing for 26/33 accepted
Pterocarpus species (75%), which were sampled across the distributional range of the genus. In total,
27 accessions were sampled within *Pterocarpus*, detailed in Supporting Information, Appendix S1,
Table S1.1. A further 22 outgroup taxa from the 21 other genera belonging to the Pterocarpus clade
within which *Pterocarpus* is nested were sampled for phylogenetic analysis. In addition, Sanger
sequence data from one nuclear locus (nrITS) and four plastid loci (*mat*K, *ndh*F-*rpl*32, *rbcl, trn*L-
*trn*F) were downloaded from NCBI GenBank https://www.ncbi.nlm.nih.gov/genbank/ (Appendix S1,
Table S1.2) for comparison between phylogenetic dating methods. These data represented 28/33
Pterocarpus species, along with 8 outgroup species from the Pterocarpus clade.

A species list of all accepted *Pterocarpus* taxa was compiled using IPNI (IPNI, 2020) and a generic
monograph of *Pterocarpus* (Rojo, 1972) to ensure that taxonomically accepted species were included
in analyses, and to account for synonyms and infraspecific taxa. Voucher specimens were examined,
and their determination updated as appropriate. Newly described and recently reinstated species from
within the *P. rohrii* complex were also sampled (Klitgård et al., *in prep*). Leaf material for DNA
extraction was acquired from both silica material (*N*= 1) and herbarium specimens (*N*= 48) collected
from the BM, K and MO herbaria.

Library preparation and sequencing

DNA was extracted from 20 mg of dried leaf material with the CTAB method (Doyle & Doyle, 1987),
following which DNA concentrations were measured using a Quantus fluorometer (Promega,
Wisconsin, USA). Libraries were prepared using the NEBNext® Ultra™ II DNA Kit (New England
Biolabs, Massachusetts, USA). Samples with high molecular weight DNA (>1000 bp, measured with

a TapeStation 4200 (Agilent Technologies, California, USA)) were sheared with a Covaris focussed
ultrasonicator M220 (Covaris, Massachusetts, USA). Following this, end-preparation and Illumina
adaptor ligation were undertaken according to the NEBNext protocol, including a 400bp size-
selection step with Agencourt Ampure XP magnetic beads (Beckman Coulter, California, USA).
Libraries containing a range of insert sizes were then amplified using PCR, following the protocol
outlined in Appendix S1, Table S1.2. Targeted bait capture was performed using the MyBaits protocol
(Arbor Biosciences, Michigan, USA) to target 353 phylogenetically informative nuclear genes with
Angiosperms-353, a bait kit designed for all Angiosperms. The final library pools were sequenced
using a paired-end 150bp run on the Illumina HiSeq platform by Macrogen 154 Inc. (Seoul, South
Korea).

Quality filtering, read assembly and alignment

DNA sequencing reads were quality-checked with FASTQC v0.11.3 (Andrews, 2010) and were
trimmed using TRIMMOMATIC v.0.3.6 (Bolger et al., 2014) to remove adapter sequences and quality-
filter reads. TRIMMOMATIC settings permitted <4 mismatches, a palindrome clip threshold of 30 and a
simple clip threshold of 6. Bases with a quality score <28 and reads shorter than 36 bases long were
removed from the dataset. Following quality-filtering, loci were assembled using SPADES v3.11.1
(Bankevich et al., 2012) by the HybPiper pipeline v1.2 (Johnson et al., 2016) with a coverage cut-off
of 8x. All loci with <50% recovery among taxa were removed, and potentially paralogous loci were
removed using the Python (Python Software Foundation, 2010) script '*paralog_investigator.py*',
distributed with the HybPiper pipeline. Recovery of different loci was visualised using the
'*gene_recovery_heatmap.R*' script distributed with HybPiper.

Following this, targeted loci were aligned by gene region (excluding those with potential paralogs)
using 1,000 iterations in MAFFT (Katoh & Standley, 2013) with the '*—adjustdirectionaccurately*'
option to incorporate reversed sequences. These alignments were then cleaned using the □*automated1*
flag in TRIMAL (Capella-Gutiérrez et al., 2009), following which they were visually inspected for poor
sequence quality using *Geneious* v. 8.1.9 (https://www.Geneious.com). This was done in order to
remove taxa with mostly missing data and to prevent the inclusion of poorly recovered loci into the
dataset, resulting in 303 refined alignments.

The nrITS and four plastid Sanger sequencing loci downloaded from NCBI GenBank were refined to
include 68 sequences belonging to 36 species from nine genera (GenBank numbers for these
sequences are listed in Appendix S1, Table S1.3). These data were aligned using MAFFT with the
same parameters as above. These sequences were also subject to visual data inspection and trimming
in Geneious to minimise missing sequence data where multiple accessions of a species were present.

Phylogenomic inference and divergence dating

Three hundred and three gene trees were inferred using RAxML v.8.0.26 (Stamatakis, 2014) with

1,000 rapid bootstrap replicates and the GTRCAT model of nucleotide substitution. A species tree

was generated based on the best-scoring RAxML trees using ASTRAL v.5.6.1 under the default

parameters, with monophyly not enforced ('-*a*' flag) (Zhang, C. et al., 2018). To compare between

phylogenetic methods, we then concatenated all 303 target capture locus alignments into a single

alignment using AMAS (Borowiec, 2016), and inferred a phylogenetic tree using RAxML HPC2

(Stamatakis, 2014) on the CIPRES web portal ((Miller et al., 2010), https://www.phylo.org/).

Inference was performed using 1,000 rapid bootstrap replicates and the GTRCAT model of nucleotide

substitution.

MCMCTREE, from the PAML package (Yang, 2007), was used to date our phylogenomic tree. This

program allows Bayesian divergence time estimation from large phylogenomic datasets based on a

pre-determined phylogenetic tree (Dos Reis & Yang, 2013). As such, the ASTRAL tree was rooted

with an appropriate outgroup (*Cascaronia astragalina*) and the tree's branch lengths were removed

with the R (R Development Core Team, 2013) package 'PhyTools' (Revell, 2012). This rooted,

branchless tree and the concatenated 303-locus target capture alignment were then used as inputs for

MCMCTREE. Fossil calibrations (listed in Table 1) were added using skew-normal priors as

uncertainty is distributed asymmetrically across the mean age in fossil calibrations (Ho & Phillips,

2009) using the *R* package 'MCMCtreeR' (Puttick, 2019).

The MCMCTREE analysis was run by first calculating the gradient (b) and Hessian (h) parameters

using the HKY substitution model, from which the posteriors of divergence times and rates were

estimated using two independent runs of 20 million generations to ensure MCMC convergence. A

third, identical run with only priors was performed and compared to the runs with data to assess

whether prior settings were adequate. MCMC convergence for posterior estimation was checked using

TRACER (Rambaut et al., 2015), as well as in the R packages 'ape' and 'bppr' (Angelis & Dos Reis,

2015, Paradis & Schliep, 2019). The final time-calibrated phylogenetic tree estimated with

MCMCTREE was then plotted with 'MCMCtreeR'.

Table 1: Fossils used to generate a time-calibrated phylogenomic tree of the Pterocarpus clade in MCMCTREE

and a dated phylogenetic tree in BEAST v.1.8.0. Skewnormal calibration priors were used for fossil ages in

MCMCTREE to provide a minimum age estimate, and the corresponding Lognormal priors were used for the

BEAST analysis. Mean ages are shown for calibrations, with 95% confidence intervals in parentheses. All ages

are in millions of years (Ma).

Calibration point	Age (Ma)	Prior distribution (MCMCTREE)	Prior distribution (BEAST)	Fossil
Pterocarpus clade	37.5 (34-41)	Skewnormal	Lognormal	*Luckowcarpa gunnii* (Martínez, 2018)
Pterocarpus-Riedeliella crown group	21.8 (15.9-27.8)	Skewnormal	Lognormal	*Pterocarpus tertiarius* (Buzek, 1992)

Tipuana crown group	8.2 (7.9-8.5)	Skewnormal	Not used (no suitable taxa)	*Tipuana ecuatoriana* (Burnham, 1995)

The GenBank dataset of five Sanger-sequenced loci was used to perform corroborative divergence time estimation in BEAST v.1.8.0 (Drummond & Bouckaert, 2015) on the CIPRES web portal. Substitution rate models were chosen for each of the five loci using AICc in JMODELTEST (Darriba et al., 2012), and are outlined in Appendix S1, Table S1.4. Tree model choice was based on path sampling analysis output from BEAST, using 10 million generations with 30 path steps. Two fossil calibration points from the MCMCTREE analysis were used to provide date estimates in BEAST (Table1). BEAST comprised three independent runs of 50 million generations which were combined using LOGCOMBINER v.1.8.2 with a burn-in level of 10%, using a birth-death tree model and a lognormal clock model. Convergence of runs was assessed using effective sample size (ESS) of all parameters in TRACER v. 1.6 with a threshold value of 200. Final ultrametric trees were generated in TREEANNOTATOR v1.8.2 and visualised using FIGTREE v1.4.4 (Rambaut, 2014). Further information on BEAST analysis is available in Appendix S1, Supplementary Methods.

Biogeographical analyses

We tested trait-dependent biogeographical models in the R package 'BioGeoBEARS' (Matzke, 2013) to understand the relationships between seed dispersal traits, biome switching and the spatio-temporal evolution of *Pterocarpus*. Fruit morphology traits were collected from the first monograph of *Pterocarpus* (Rojo, 1972), a taxonomic revision circumscribing new species within the *P. rohrii* complex (Klitgård *et al. in prep*), and taxonomic treatments of various *Pterocarpus* species (de Candolle, 1825, Klitgård et al., 2000, Zamora, 2000). Since no fruiting specimen of *P. rohrii* var. *rubiginosus* exists, it was inferred to be winged since it belongs to the *P. rohrii* species complex. Fruit dispersal traits were then plotted on the *Pterocarpus* ASTRAL tree with biome and realm for each species using the *dotplot()* function in the R package 'PhyTools'.

Geographical distribution data for non-cultivated *Pterocarpus* specimens were collected from GBIF (www.GBIF.org (06 November 2019), GBIF Occurrence Download https://doi.org/10.15468/dl.tr3h2y) based on the curated taxonomic name list described above. This dataset was then further refined using the R package 'CoordinateCleaner' ((Zizka et al., 2019); https://github.com/ropensci/CoordinateCleaner) and the input for our *BioGeoBears* analysis was produced from these data with Alex Zizka's 'biogeography in R' scripts (https://github.com/azizka/Using_biodiversity_data_for_biogeography). These records and the specimens underpinning them were then examined in detail by taxonomic experts on the *Pterocarpus* clade (B. B. Klitgaard and G. P. Lewis) to further remove records which were mis-identified.

Based on these data, we assigned species to biogeographical regions using the *wwfLoad()* function in 'speciesgeocodeR' (Töpel et al., 2017), based on the WWF ecoregions of Olson et al. (2001). These ecoregions represented five tropical terrestrial 'realms' (Australasia, Afrotropics, Indo-Malaya, Neotropics and Oceania) and six tropical terrestrial biomes (tropical moist forest (i.e., 'rainforest'),

montane grassland, desert, mangrove, tropical dry forest and tropical grassland (i.e., 'savanna')),
which are displayed in Appendix S1, Fig. S1.1a. The assignments for each species were cross-
referenced with monographs and previous studies of *Pterocarpus* (Rojo, 1972, Saslis-Lagoudakis et
al., 2011, Klitgård et al., in prep), and with the Plants Of the World Online database
(http://powo.science.kew.org/). All traits (fruit morphology, realm assignment and biome assignment)
are then visualised onto the *Pterocarpus* phylogenetic tree in Appendix S1, Fig. S1.1b.

We then compared the fit of trait-independent and trait-dependent biogeographical models outlined by
Klaus and Matzke (2020) for biogeographical realms and biomes in BioGeoBEARS. For realms, we
tested between three default models in BioGeoBEARS, all of which incorporate dispersal, extinction
and range switching. These models were DEC (Ree & Smith, 2008), which additionally accounts for
vicariance and 'subset' sympatric speciation, DIVALIKE (Ronquist, 1997), which also accounts for
vicariance, and BAYAREALIKE (Landis et al., 2013), which additionally accounts for widespread
sympatric speciation but not vicariance. For biomes, we assessed the fit of Markov-k (Mk) models
(Lewis, P. O., 2001) (i.e., BAYAREALIKE a+, d=e=0 in BioGeoBEARS) to accommodate equal
rates of character evolution, after Kriebel et al. (2019). Markov-k more accurately models the
inheritance of ecological preferences in daughter species and avoids bias towards large ancestral
ranges compared to models such as DEC. We also included four trait-dependent models in our
comparisons. The first two models only parameterised trait switching ('*Trait_1Rate*' and
'*Trait_2Rates*'). Then, for realm, a trait-dependent extension of the DEC model was used to include
trait switching and multipliers on dispersal rate (*DEC_t12_t21_m2*), and for biome a trait-dependent
implementation of the Markov-k model was used, which included the same trait switching and
dispersal parameters ('*Markov-k'_t12_t21_m2*).

The best-fit model was chosen using AICc and Akaike weights, from which ancestral ranges were
estimated across *Pterocarpus*. Ancestral ranges were plotted onto the time-calibrated tree from
MCMCTREE using the function '*plot_BioGeoBears_results()*' for realm and biome independently.
Finally, for biome, we counted the number of transitions between the most likely ancestral biomes at
each node, and produced a matrix of biome shifts. This matrix was visualised as a transition plot using
the *plotmat* function in the R package 'diagram' (Soetaert, 2012). Further information regarding
biogeographical data collection and analysis is available in Appendix S1, Supplementary Methods.

<u>Results</u>

Phylogeny and divergence dating

A mean of 17.7% of reads were successfully assembled onto the *Angiosperms*-353 bait sequences,
with a range from 6.9% to 30.5%. Gene recovery is shown as a heatmap in Appendix S1, Fig. S1.2.
Of the 353 loci targeted by *Angiosperms*-353, we discarded 35 loci which were potentially paralogous
and 15 which were assembled poorly or had over 50% missing data across taxa.

Our ASTRAL analysis indicated that most bipartitions were well supported with local posterior
probabilities (LPP) >0.8, with 1 being full support (Fig. 1) and a quartet score of 0.68. Within this
tree, most generic splits are well supported (LPP >0.9) and largely support the monophyly of

Pterocarpus with the exception of one species, *Pterocarpus dubius* (formerly *Etaballia dubia*).

Pterocarpus dubius was resolved in a clade containing *Inocarpus, Maraniona* and *Paramachaerium*,

which was in a sister position to *Pterocarpus*. Within *Pterocarpus*, there were two major subclades,

one containing species from the Neotropics (the neotropical clade) and one containing species from

the Palaeotropics (the palaeotropical subclade). However, this split received a relatively low support

(LPP=0.43, quartet score = 34.48%). The concatenated RAXML analysis of the same 303 loci

(Appendix S1, Fig. S1.3) recovered a nearly identical topology to the ASTRAL species tree (Fig. 1),

and was much better resolved, with nearly all nodes having bootstrap values >95. Moreover, the split

between neotropical and palaeotropical *Pterocarpus* species received full support (BS=100) in the

inference based on the concatenated dataset.

Divergence time estimation carried out in MCMCTREE provided a robust, time-calibrated tree of the

Pterocarpus clade. The two MCMC runs were convergent, with every node showing a suitable

effective sample size (> 200), and posterior estimates fit the priors well as indicated in Appendix S1,

Fig. S1.4a, b and c. The time-calibrated phylogenomic tree in Fig. 1 suggests that the Pterocarpus

clade arose ~35 Ma, during the late Eocene, and diversification of present-day genera occurred during

the Miocene, from 23-5.3 Ma. Of these, *Pterocarpus* (*sensu stricto*) diverged from its sister genera

~12 Ma, and diversified mostly in the late Miocene. It is worth noting that divergence date estimates

are described by larger HPD intervals in the earlier divergence events within the Pterocarpus clade,

and this increased uncertainty is mirrored in Appendix S1, Fig. S1.5. This figure indicates that

uncertainty is greater in the earlier divergence events inferred, as well as being much reduced around

fossil calibration points, as would be expected.

Figure 1: Fossil-calibrated MCMCTREE of the Pterocarpus clade. Node bars show 95% highest posterior

density (HPD) for node ages. Local posterior probabilities (LPP), a measure of topological support, are shown

for each node, and were taken from the ASTRAL analysis which provided the tree topology for the MCMCTREE

analysis. Shading on the background of the phylogram represents geological epochs, which are labelled beneath

the tree. The red and green shading indicates species belonging to the palaeotropical and neotropical subclades

of *Pterocarpus*, respectively, and the extent of the genus *sensu stricto* is shown by the black bar. The dashed

box around *Pterocarpus dubius* highlights its paraphyly with the rest of the genus. Photographs show winged

fruits of *Pterocarpus erinaceus* (© Gwilym Lewis), coriaceous fruits of *Pterocarpus monophyllus* (© Domingos

Cardoso) and winged fruits of *Pterocarpus violaceus* (© Reinaldo Aguilar, CC BY-NC-SA 2.0).

The topology of the BEAST trees (Appendix S1, Fig. S1.6) inferred using Sanger-sequencing data from GenBank recovered a broadly similar topology to the MCMCTREE analysis (Fig. 1). Moreover, the divergence time estimates of this phylogeny were largely congruent with those from Fig. 1, including an origin of the Pterocarpus clade within the late Eocene (~38 Ma), followed by most divergence between genera, and diversification of *Pterocarpus*, within the Miocene (~23-5.3 Ma). However, the age estimate for the initial divergence of *Pterocarpus* is later than that inferred with MCMCTREE, at around 25 Ma, again during the Miocene. In addition, the BEAST analysis displayed

a much broader degree of uncertainty on age estimates, as indicated by the breadth of the 95% HPDs,

and recovered *Pterocarpus dubius* as nested within *Pterocarpus*, whereas in the MCMCTREE analysis

P. dubius is in sister clade position to the rest of *Pterocarpus*.

Biogeography

AICc comparison indicated that the best-fitting BioGeoBEARS model was DIVALIKE (AICc=

52.785) (Appendix S1, Table S1.5). Using DIVALIKE, we estimated a shared ancestral range of both

the Neotropics and Afrotropics for the common ancestor of *Pterocarpus*, with subsequent splitting

into neotropical and palaeotropical clades (Fig. 2a). The neotropical clade diversified only within the

Americas and gave rise to the most species. Within the palaeotropical clade the common ancestor of

P. indicus and *P. dalbergioides* appears to have dispersed into Asia, and subsequently into Australasia

and Oceania, having a shared Afro-Asian range. It is interesting to note that within both neotropical

and palaeotropical clades, there are species which share ranges, being found in both eastern South

America and west Africa (*P. santalinoides* and *P. officinalis*). A shared characteristic of these two

amphi-Atlantic species are thick-walled, wingless, buoyant fruits which allow water-borne dispersal.

The current distribution of *Pterocarpus* species (both in terms of biome and realm), as well as the

distribution of fruit dispersal traits across the ASTRAL tree, are visualised in Appendix S1, Fig.

S1.1b.

For biome evolution, the best-fit model was the trait-independent implementation of Markov-*k*

(AICc= 328.218, Appendix S1, Table S1.5). Moist forest, dry forest and grassland were inferred as

being nearly equally likely as the ancestral biome for the MRCA of *Pterocarpus* (Fig. 2b). Within the

neotropical clade of *Pterocarpus*, there appears to be multiple polymorphic biome preferences, with

species being found in dry forest as well as one other habitat type (desert, grassland or moist forest).

However, for the majority of stem nodes within the neotropical *Pterocarpus* species, the most likely

ancestral biome appears to have been dry forest, with only the clade containing *P. rohrii, P.*

santalinoides, P. amazonum and *P. floribundus* containing species only found in moist forest. Within

the palaeotropical *Pterocarpus* clade, the most probable state for the majority of ancestral nodes was

shared between moist forest and grassland, with most species being found in grassland in addition to

other biomes, mirroring the high polymorphism in neotropical *Pterocarpus* species. It is also of

interest that one of the two *Pterocarpus* species which inhabit both South America and Africa (*P.*

officinalis) occurs both in tropical rainforest as well as mangroves.

We inferred the most shifts between biome states across the *Pterocarpus* tree into moist forests (i.e.,

rainforest) and grassland (i.e., savanna). Specifically, we recovered the most shifts from dry forest to

moist forest (n=4, Fig. 3), followed by shifts from rainforest to grassland (n=3). Transitions also

occurred between drier habitats (e.g. inferred shifts from dry forest to desert and grassland (n=2), or

grassland to dry forest and desert (n=1)), between similar habitats (e.g., grassland into montane

grassland (n=2), from moist forest to mangrove (n=1)) and in the contrasting direction to the most

frequent shifts (e.g. moist forest to dry forest (n=2) and grassland to moist forest (n=1)).

Figure 2a) Biogeographical range estimation of *Pterocarpus*. Pie charts represent likelihoods of ranges derived
from ancestral range estimations run on the MCMCTREE topology, using the *R* package 'BioGeoBEARS'. The
legend represents five biogeographical realms defined by Olson *et al.* (2001) and shown in Appendix S1, Fig.
S1.1a. Shading on the background of the phylogenetic tree represents geological epochs, and within the bottom
bar 'Ple' represents 'Pleistocene', 'Holo.' represents 'Holocene' and 'Quatern.' represents 'Quaternary'. Inset is
a map displaying biogeographical realms as defined by Olson *et al*. (2001).

Figure 2b) Historical biome estimation of *Pterocarpus*. Pie charts represent likelihoods of ranges derived from ancestral biome estimations run on the MCMCTREE topology, using the *R* package 'BioGeoBEARS'. The legend represents six biomes defined by Olson *et al.* (2001) and shown in Appendix S1, Fig. S1.1a. Shading on the background of the phylogenetic tree represents geological epochs, and within the bottom bar 'Ple' represents 'Pleistocene', 'Holo.' represents 'Holocene' and 'Quatern.' represents 'Quaternary'. Inset is a map displaying biomes as defined by Olson *et al.* (2001).

Figure 3) Biome transitions between species within *Pterocarpus*, based on the ancestral biome reconstruction
with the M*k* model shown in Fig. 2b. Arrows on lines show the direction of the biome transition, while numbers
and thickness of the lines represent the number of transitions between each biome. Inset is a map displaying
biomes as defined by Olson *et al*. (2001).

Discussion

Divergence dating indicated that the Pterocarpus clade began to diversify in the late Eocene (~35 Ma),
followed by extensive diversification during the Miocene (23-5.3 Ma). During the Miocene, the
common ancestor of *Pterocarpus* diverged into two clades. One clade was palaeotropical and
diversified largely in Africa with subsequent dispersal into Indo-Malaya, Australasia and Oceania.
The second clade was neotropical, which diversified only in the tropical Americas and was more
speciose than the palaeotropical clade. The diversification of *Pterocarpus* appears to have involved
multiple biome shifts, largely into lowland habitats (namely, savanna and tropical moist forest), and
while dispersal across large geographical areas played a role in range expansion, the fruit dispersal
traits we examined were not significantly associated with macroevolutionary dynamics according to
our model comparisons.

Overall, it appears that traits for hydrochoric dispersal evolved multiple times independently in
Pterocarpus, likely in concert with adaptation to waterlogged habitats such as mangroves. These traits
which evolved to promote dispersal in the ecological theatre may have then led to rare long-distance
dispersal across oceans (as we see in extant species such as *P. santalinoides* and *P. officinalis*). A
similar pattern has been shown in the Bombacoideae (Malvaceae) (Zizka et al., 2020), as well as in
Centrolobium and *Platymiscium*, which are close relatives of *Pterocarpus* (Klitgård, 2005, Pirie et al.,
2009, Saslis□Lagoudakis et al., 2008).

Phylogenetic analysis of the Pterocarpus clade

Most Pterocarpus clade lineages, including *Pterocarpus* itself, arose during the Miocene, between 23-
5.3 Ma (Fig. 1; Appendix S1, Fig. S1.6). This was congruent across methods (MCMCTREE, BEAST)
and across data types (target capture data, Sanger-sequencing data). While our divergence dating
analyses were broadly congruent with each other across methods and data type, they differed in their
date estimates for the origin of *Pterocarpus*. MCMCTREE analysis inferred an origin date of 12Ma
(Fig. 1), whereas the BEAST analysis suggested that *Pterocarpus* arose around 25Ma (Appendix S1,
Fig. S1.6). The most likely reason for this discrepancy is the size of the dataset, since MCMCTREE
was run on a nuclear target capture dataset comprising 303 loci, while BEAST was run on pre-
existing Sanger sequence data for 5 loci, resulting in a much less informative sequence matrix.

It is also likely that the inclusion of plastid loci within the BEAST analysis extended the date estimate
for the origin of *Pterocarpus*, due to deeper divergences within the plastid genome (Wolfe et al.,
1987). This conflicting signal would have also led to the much higher degree of uncertainty on age
estimates in the BEAST analysis, as indicated by the breadth of the 95% HPDs. In addition, the nested
position of *Pterocarpus dubius* within *Pterocarpus*, as shown in the BEAST analysis (Appendix S1,
Fig. S1.6) contrasts with the sister clade position of *P. dubius* to *Pterocarpus* in the MCMCTREE
analysis (Fig. 1). This species was previously circumscribed as *Etaballia dubia* and is highly
divergent in terms of floral morphology, however previous work with Sanger-sequencing loci
resolved this species within *Pterocarpus* (Klitgård et al., 2013). It is possible that the inclusion of
chloroplast loci in the BEAST tree caused this incongruence, due to the lower mutation rate of

chloroplast regions (Wolfe et al., 1987), or through plastid capture resulting from hybridisation

(Naciri & Linder, 2015). While there is undoubtedly useful signal within these plastid loci, further

work using plastomes would help to accurately disentangle these discordances.

Spatio-temporal evolution of Pterocarpus

The Miocene-era diversification of the Pterocarpus clade, and particularly *Pterocarpus*, reflects

patterns shown in other tropical tree genera (e.g., the mahoganies (Meliaceae) (Koenen et al., 2015),

the Brownea clade (Leguminosae) (Schley et al., 2018) and the Daniellia clade (Leguminosae) (Choo

et al., 2020)). This diversification was likely concurrent with major climatic, geological and

ecological changes which occurred in the tropics during the Miocene (Keeley & Rundel, 2005,

Morley, 2000) such as was the Miocene global cooling event (Zachos et al., 2001, 2008). This cooling

was precipitated by the closure of the Tethys seaway and a subsequent change in ocean currents

(Rommerskirchen et al., 2011, Zhang, Z. et al., 2014), leading to a decrease in global temperature,

desiccation of the continents and a retreat of tropical vegetation, especially on the African continent

(Couvreur et al., 2021, Morley & Richards, 1993, Senut et al., 2009). Such cooling likely also

explains the long-distance dispersal events and biome-switching we observed in our ancestral range

reconstructions for *Pterocarpus* (Fig. 2a & 2b), particularly within the palaeotropical clade, as

discussed below.

i.   *The palaeotropical clade*

The palaeotropical clade appears to have diversified largely within Africa over the last ~10Ma, in the

late Miocene and into the early Pliocene (Fig. 2a), with multiple biome shifts from moist forests into

grasslands (n=3, Fig. 3). These events were likely concurrent with the cooling and desiccation of the

African continent which lead to the expansion of grasslands during this period (Couvreur et al., 2021).

This palaeotropical clade was also less speciose than its neotropical counterpart, which may be

explained by higher levels of extinction in rainforest lineages, again precipitated by the expansion of

grassland in Africa's past. This has led to the well-documented 'odd man out' pattern of rainforest

tree diversity, with the Afrotropics being significantly less diverse that other tropical regions

(Couvreur, 2015, Richards, 1973). Moreover, most of the extant *Pterocarpus* species within the

palaeotropical clade are now found in grassland and are wind-dispersed (Fig. 2b; Appendix S1, Fig.

S1.1b), suggesting historical adaptation to novel grassland environments.

Within the Palaeotropical clade, the two Asian sister species (*P. dalbergioides* and *P. indicus*) only

dispersed to the Asian tropics and beyond into Australasia and Oceania around 4Ma, during the

Pliocene (Fig. 2a). It is likely that the divergence of these species was caused by infrequent eastward

seed dispersal, promoted by the east African coastal current (EACC) between Africa and Asia which

formed with the closing of the Tethys seaway in the late Miocene (Scotese, 2004). It is interesting to

note that these species are found in rainforests, as rainforest tree species tend to be excellent

dispersers (Dexter et al., 2017, Pennington & Lavin, 2016), and are often water dispersed. This

mirrors the highly dispersible nature of the other rainforest-dwelling member of the Palaeotropical

clade (*P. officinalis*) which is distributed in both Africa and South America and is also found in

mangroves. Moreover, *P. officinalis* is derived from Miocene African ancestors (Fig. 2a), is present in

the Neotropical fossil record (Graham, 1995) and includes a West African subspecies (subsp. *gilletii*),

further suggesting that the corky exocarp of the fruit, which it evolved for dispersal at an ecological

scale (Appendix S1, Fig. S1.1b), is responsible for rare dispersal across the Atlantic and the

foundation of a Neotropical population (Rivera☐Ocasio et al., 2002).

ii.   *The neotropical subclade*

Within neotropical *Pterocarpus* we inferred multiple events of biome-switching from dry forest into

rainforest (Fig. 2b; n=4, Fig. 3)), with the most likely ancestral biome for this clade being dry forest.

These events mainly occurred between the late Miocene and the Pliocene (~8 - 2 Ma). In particular,

we see the switch from dry forest in stem nodes into clades which are found nearly exclusively in wet

forest towards the tips, mirroring findings in closely related genera (e.g. *Platymiscium*

(Saslis☐Lagoudakis et al., 2008)). Indeed, biome switching between dry forest and moist forest,

which are often interdigitated, is well documented in the Neotropics (Dexter et al., 2018, Ireland et al.,

2010, Klitgård, 2005, Pennington & Dick, 2010, Pezzini et al., 2020).

The majority of Neotropical dry forests formed during the Miocene (Pennington & Ratter, 2006) as

shown by fossil evidence (e.g. Burnham, 1995, Burnham & Carranco, 2004). This was followed by

periods of intense climatic flux and the contraction of dry forests resulting from increasingly wet

Pleistocene interglacial periods, leading to the 'refugia' of dry forest which we observe today within

inter-Andean valleys, Central America and the 'dry arc' south of the Amazon in Brazil and Bolivia

(Pennington et al., 2004). The biome switching we inferred within the Neotropics may be explained

by these oscillations, as the expansion of rainforests would have resulted in a much higher likelihood

of immigration into rainforest by plant species dispersing from dry forest refugia.

iii.   *Asymmetrical biome transitions in* Pterocarpus

Biome transitions across *Pterocarpus* occurred mainly into moist forest (i.e., rainforest) and grassland

(i.e., savanna) (Fig. 3). The higher relative frequency of immigration into both moist forests and

grasslands may be explained by the fact that these communities experience persistent disturbance

(e.g., fire in tropical savannas) or because they contain species which are vulnerable to climatic

perturbation (e.g., drought in rainforests) (da Costa et al., 2010, Pennington & Lavin, 2016). In

rainforests, intermediate disturbance increases the likelihood of seedling establishment across entire

communities due to intense competition for light (Hubbell et al., 1999), and within savannas, seedling

establishment of local species is greatly reduced by frequent fire (Hoffmann, 1996, Setterfield, 2002,

Wakeling et al., 2011). This may promote the competitive success of tree seeds dispersing into such

environments from adjacent habitats. Thus, the highly dispersible nature of *Pterocarpus* fruits could

have promoted immigration into these novel biomes, and the moderate disturbance which occurs to

plant communities within these biomes could have facilitated establishment, adaptation and

speciation.

The frequency of biome shifts into savanna and moist forests also implies that their wide extent
renders them more likely to exchange seeds with adjacent biomes (Donoghue & Edwards 2014). This
is particularly true for the 'donation' of immigrants from restricted biomes which interdigitate with
savannas or rainforest, such as dry forest. Overall, our results highlight that biome switching may play
an important role in diversification of tropical trees and is more common than previously thought, as
recently highlighted across neotropical taxa (Antonelli et al., 2018).

Caveats

Despite the utility of biogeographical models for understanding spatio-temporal evolution, they are
not without their drawbacks. In particular, biogeographical models may be prone to over-estimating
long-distance dispersal when model assumptions are violated (e.g., *DEC+J* in BioGeoBEARS (Ree &
Sanmartín, 2018)), highlighting the importance of appropriate model choice (Matzke, 2014). As far as
is possible, we have attempted to account for these biases by, for example, avoiding the use of the *+J*
parameter in biogeographical analyses.

It is also possible that small sample sizes may be problematic, since with larger datasets parameter
estimates are likely to be more accurate (e.g., in species distribution modelling (Wisz et al., 2008)).
However, as in our study, this can be avoided by having a balanced sampling strategy which accounts
for the nearly all species in studies to <50 taxa (e.g., in Araliaceae (Valcárcel & Wen, 2019), penguins
(Vianna et al., 2020) and cat-eyed snakes (Weinell et al., 2021)). Finally, our specimens were
identified by taxonomic experts on *Pterocarpus* and underwent multiple corroborative phylogenetic
analyses, thereby minimising error as a result of misidentification or phylogenetic uncertainty.
However, even while accounting for such biases, it is important to interpret biogeographical model
outputs cautiously, and to realise that they are approximate estimates of biogeographic history.

Conclusion

Overall, our analyses suggest that while a dispersal trait-dependent model of range macroevolution
was not supported, the biogeographical history of *Pterocarpus* was likely mediated by cross-
continental dispersal, associated range change and diversification into adjacent biomes. Indeed, biome
switching between moist and arid environments, likely precipitated by infrequent long-distance
dispersal events and environmental change, appear to have promoted ecological differentiation and
diversification within *Pterocarpus* since the early Miocene.

Data availability statement

The target capture sequencing data generated for this project are publicly available in the NCBI
sequence read archive under the BioProject number PRJNA728569, with the accession numbers
SAMN19092119 - SAMN19092167.

References

Andrews, S. (2010) FastQC: a quality control tool for high throughput sequence data. Available
online at http://www.bioinformatics.babraham.ac.uk/projects/fastqc.

Angelis, K. & Dos Reis, M. (2015) The impact of ancestral population size and incomplete lineage
sorting on Bayesian estimation of species divergence times. *Current Zoology* **61**(5) 874-885.

Antonelli, A., Zizka, A., Carvalho, F.A., Scharn, R., Bacon, C.D., Silvestro, D. & Condamine, F.L.
(2018) Amazonia is the primary source of Neotropical biodiversity. *Proceedings of the National*
Academy of Sciences of the United States of America **115**(23) 6034-6039.

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S. ... Prjibelski, A.D.
(2012) SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing.
Journal of Computational Biology **19**(5) 455-477.

Bellot, S., Bayton, R.P., Couvreur, T.L., Dodsworth, S., Eiserhardt, W.L., Guignard, M.S. ... Baker,
W.J. (2020) On the origin of giant seeds: the macroevolution of the double coconut (Lodoicea
maldivica) and its relatives (Borasseae, Arecaceae). *The New Phytologist* **228**(3) 1134.

Bolger, A.M., Lohse, M. & Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**(15) 2114-2120.

Borowiec, M.L. (2016) AMAS: a fast tool for alignment manipulation and computing of summary
statistics. *PeerJ* **4**e1660.

Brown, J.H., Stevens, G.C. & Kaufman, D.M. (1996) The geographic range: size, shape, boundaries,
and internal structure. *Annual Review of Ecology and Systematics* **27**(1) 597-623.

Burnham, R.J. (1995) A new species of winged fruit from the Miocene of Ecuador: Tipuana
ecuatoriana (Leguminosae). *American Journal of Botany* **82**(12) 1599-1607.

Burnham, R.J. & Carranco, N.L. (2004) Miocene winged fruits of Loxopterygium (Anacardiaceae)
from the Ecuadorian Andes. *American Journal of Botany* **91**(11) 1767-1773.

Buzek, C. (1992) Fruits of Pterocarpus tertiarius Weyland from the North-Bohemian Basin,
Czechoslovakia. *Advances in legume systematics: Part 4*(ed. by P.S. Herendeen & D.L. Dilcher), pp.
19-31.

Cain, M.L., Milligan, B.G. & Strand, A.E. (2000) Long□distance seed dispersal in plant populations.
American Journal of Botany **87**(9) 1217-1227.

Capella-Gutiérrez, S., Silla-Martínez, J.M. & Gabaldón, T. (2009) trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**(15) 1972-1973.

Choo, L.M., Forest, F., Wieringa, J.J., Bruneau, A. & de la Estrella, M. (2020) Phylogeny and
biogeography of the Daniellia clade (Leguminosae: Detarioideae), a tropical tree lineage largely
threatened in Africa and Madagascar. *Molecular phylogenetics and evolution* **146**106752.

Corlett, R.T. & Primack, R.B. (2011) *Tropical rain forests: an ecological and biogeographical*
*comparison*John Wiley & Sons.

Cousens, R., Dytham, C. & Law, R. (2008) *Dispersal in plants: a population perspective*Oxford
University Press.

Couvreur, T.L. (2015) Odd man out: why are there fewer plant species in African rain forests? *Plant*
Systematics and Evolution **301**(5) 1299-1313.

Couvreur, T.L., Dauby, G., Blach☐Overgaard, A., Deblauwe, V., Dessein, S., Droissart, V. ... Ley,
A.C. (2021) Tectonics, climate and the diversification of the tropical African terrestrial flora and
fauna. *Biological Reviews* **96**(1) 16-51.

da Costa, A.C., Galbraith, D., Almeida, S., Portela, B.T.T., da Costa, M., Junior, João de Athaydes
Silva ... Fisher, R. (2010) Effect of 7 yr of experimental drought on vegetation dynamics and biomass
storage of an eastern Amazonian rainforest. *New Phytologist* **187**(3) 579-591.

Darriba, D., Taboada, G.L., Doallo, R. & Posada, D. (2012) jModelTest 2: more models, new
heuristics and parallel computing. *Nature Methods* **9**(8) 772.

de Candolle, A.P. (1825) *Prodromus Systematis Naturalis Regni Vegetabilis 2. Parisii: Sumptibus*
Sociorum Treuttel et Wu☐rtz.

de la Estrella, M., Forest, F., Wieringa, J.J., Fougere-Danezan, M. & Bruneau, A. (2017) Insights on
the evolutionary origin of Detaroideae, a clade of ecologically dominant tropical African trees. *New*
Phytologist **214**(4) 1722-1735.

De Micco, V. & Aronne, G. (2012) Morpho-anatomical traits for plant adaptation to drought. *Plant*
responses to drought stress(ed. by Anonymous), pp. 37-61. Springer.

De Queiroz, A. (2005) The resurrection of oceanic dispersal in historical biogeography. *Trends in*
Ecology & Evolution **20**(2) 68-73.

Dexter, K.G., Smart, B., Baldauf, C., Baker, T.R., Balinga, M.P., Brienen, R. ... Muledi, J.I. (2015)
Floristics and biogeography of vegetation in seasonally dry tropical regions. *International Forestry*
Review **17**(2) 10-32.

Dexter, K.G., Pennington, R.T., Oliveira-Filho, A.T., Bueno, M.L., Silva de Miranda, Pedro L &
Neves, D.M. (2018) Inserting tropical dry forests into the discussion on biome transitions in the
tropics. *Frontiers in Ecology and Evolution* **6**104.

Dexter, K.G., Lavin, M., Torke, B.M., Twyford, A.D., Kursar, T.A., Coley, P.D. ... Pennington, R.T.
(2017) Dispersal assembly of rain forest tree communities across the Amazon basin. *Proceedings of*
the National Academy of Sciences of the United States of America **114**(10) 2645-2650.

Donoghue, M.J. & Edwards, E.J. (2014) Biome shifts and niche evolution in plants. *Annual Review of*
Ecology, Evolution, and Systematics **45**547-572.

Dos Reis, M. & Yang, Z. (2013) MCMCTree
tutorials. http://abacus.gene.ucl.ac.uk/software/MCMCtree.Tutorials.pdf. .

Doyle, J.J. & Doyle, J.L. (1987) Genomic plant DNA preparation from fresh tissue- CTAB method.
Phytochemical Bulletin **19**(11) 11-15.

Drummond, A.J. & Bouckaert, R.R. (2015) *Bayesian evolutionary analysis with BEAST*Cambridge
University Press.

Echeverría-Londoño, S., Enquist, B.J., Neves, D.M., Violle, C., Boyle, B., Kraft, N.J. ... Sandel, B.
(2018) Plant functional diversity and the biogeography of biomes in North and South America.
Frontiers in Ecology and Evolution **6**219.

Eiserhardt, W.L., Svenning, J., Kissling, W.D. & Balslev, H. (2011) Geographical ecology of the
palms (Arecaceae): determinants of diversity and distributions across spatial scales. *Annals of Botany*
108(8) 1391-1416.

Garamszegi, L.Z. & Møller, A.P. (2011) Nonrandom variation in within-species sample size and
missing data in phylogenetic comparative studies. *Systematic Biology* **60**(6) 876-880.

Gentry, A.H. (1988) Tree species richness of upper Amazonian forests. *Proceedings of the National*
Academy of Sciences **85**(1) 156-159.

Gillespie, R.G., Baldwin, B.G., Waters, J.M., Fraser, C.I., Nikula, R. & Roderick, G.K. (2012) Long-
distance dispersal: a framework for hypothesis testing. *Trends in ecology & evolution* **27**(1) 47-56.

Graham, A. (1995) Diversification of Gulf/Caribbean mangrove communities through Cenozoic time.
Biotropica 20-27.

Harris, A.J., Ickert☐Bond, S. & Rodríguez, A. (2018) Long distance dispersal in the assembly of
floras: A review of progress and prospects in North America. *Journal of Systematics and Evolution*
56(5) 430-448.

Ho, S.Y. & Phillips, M.J. (2009) Accounting for calibration uncertainty in phylogenetic estimation of
evolutionary divergence times. *Systematic Biology* **58(3)**367-380.

Hoffmann, W.A. (1996) The effects of fire and cover on seedling establishment in a neotropical
savanna. *Journal of Ecology* 383-393.

Hubbell, S.P., Foster, R.B., O'Brien, S.T., Harms, K.E., Condit, R., Wechsler, B., Wright, S.J. & De
Lao, S.L. (1999) Light-gap disturbances, recruitment limitation, and tree diversity in a neotropical
forest. *Science* **283**(5401) 554-557.

Hughes, C., Pennington, R.T. & Antonelli, A. (2013) Neotropical plant evolution: assembling the big
picture. *Botanical Journal of the Linnean Society* **171**(1) 1-18.

IPNI. (2020) The international plant names index. **2019**(10 December) .

Ireland, H.E., Kite, G.C., Veitch, N.C., Chase, M.W., Schrire, B., Lavin, M., Linares, J. &
Pennington, R.T. (2010) Biogeographical, ecological and morphological structure in a phylogenetic
analysis of Ateleia (Swartzieae, Fabaceae) derived from combined molecular, morphological and
chemical data. *Botanical Journal of the Linnean Society* **162**(1) 39-53.

Johnson, M.G., Gardner, E.M., Liu, Y., Medina, R., Goffinet, B., Shaw, A.J., Zerega, N.J. & Wickett,
N.J. (2016) HybPiper: Extracting coding sequence and introns for phylogenetics from
high☐throughput sequencing reads using target enrichment. *Applications in Plant Sciences* **4**(7)
1600016.

Katoh, K. & Standley, D.M. (2013) MAFFT multiple sequence alignment software version 7:
improvements in performance and usability
. *Molecular Biology and Evolution* **30**(4) 772-780.

Keeley, J.E. & Rundel, P.W. (2005) Fire and the Miocene expansion of C4 grasslands. *Ecology*
Letters **8**(7) 683-690.

Klaus, K.V. & Matzke, N.J. (2020) Statistical Comparison of Trait-Dependent Biogeographical
Models Indicates That Podocarpaceae Dispersal Is Influenced by Both Seed Cone Traits and
Geographical Distance. *Systematic Biology* **69**(1) 61-75.

Klitgård, B.B. & Lavin, M. (2005) Tribe Dalbergieae sens. lat. *Legumes of the world*(ed. by G.P.
Lewis, B. Schrire, B. Mackinder & M. Lock), pp. 307-335. Kew Publishing, Richmond, Surrey.

Klitgård, B.B., de Queiroz, L.P. & Lewis, G.P. (2000) A remarkable new species of Pterocarpus
(Leguminosae: Papilionoideae: Dalbergieae) from Bahia, Brazil. *Kew Bulletin* 989-992.

Klitgård, B.B. (2005) Platymiscium (Leguminosae: Dalbergieae): Biogeography Systematics,
Morphology, Taxonomy and Uses. *Kew Bulletin* 321-400.

Klitgård, B.B., Forest, F., Booth, T.J. & Saslis-Lagoudakis, C.H. (2013) A detailed investigation of
the Pterocarpus clade (Leguminosae: Dalbergieae): Etaballia with radially symmetrical flowers is
nested within the papilionoid-flowered Pterocarpus. *South African Journal of Botany* **89**128-142.

Koenen, E.J., Clarkson, J.J., Pennington, T.D. & Chatrou, L.W. (2015) Recently evolved diversity and
convergent radiations of rainforest mahoganies (Meliaceae) shed new light on the origins of rainforest
hyperdiversity. *New Phytologist* **207**(2) 327-339.

Kriebel, R., Drew, B.T., Drummond, C.P., González☐Gallegos, J.G., Celep, F., Mahdjoub, M.M. ...
Walker, J.B. (2019) Tracking temporal shifts in area, biomes, and pollinators in the radiation of Salvia
(sages) across continents: leveraging anchored hybrid enrichment and targeted sequence data.
American Journal of Botany **106**(4) 573-597.

Landis, M.J., Matzke, N.J., Moore, B.R. & Huelsenbeck, J.P. (2013) Bayesian analysis of
biogeography when the number of areas is large. *Systematic Biology* **62**(6) 789-804.

Lewis, G.P. (1987) *Legumes of Bahia*Kew Publishing, Richmond, Surrey.

Lewis, G.P., Schrire, B., Mackinder, B. & Lock, M. (2005) *Legumes of the World*Kew Publishing,
Richmond, Surrey.

Lewis, P.O. (2001) A likelihood approach to estimating phylogeny from discrete morphological
character data. *Systematic Biology* **50**(6) 913-925.

Martínez, C. (2018) Dalbergieae (Fabaceae) Samara Fruits from the Late Eocene of Colombia.
International journal of plant sciences **179**(7) 541-553.

Matzke, N.J. (2014) Model selection in historical biogeography reveals that founder-event speciation
is a crucial process in island clades. *Systematic Biology* **63**(6) 951-970.

Matzke, N.J. (2013) BioGeoBEARS: BioGeography with Bayesian (and Likelihood) Evolutionary
Analysis in R Scripts. https://rdrr.io/cran/BioGeoBEARS/.

Midgley, J.J. & Bond, W.J. (2013) Plant adaptations to fire: an evolutionary perspective. *Fire*
Phenomena and the Earth System: An Interdisciplinary Guide to Fire Science(ed. by C. M. Becher),
pp. 125-134. Wiley Online Library.

Miller, M.A., Pfeiffer, W. & Schwartz, T. (2010) Creating the CIPRES Science Gateway for inference
of large phylogenetic trees. *Proceedings of the Gateway Environments Workshop (GCE)* 1-8.

Moore, B.R., Hohna, S., May, M.R., Rannala, B. & Huelsenbeck, J.P. (2016) Critically evaluating the
theory and performance of Bayesian analysis of macroevolutionary mixtures. *Proceedings of the*
National Academy of Sciences of the United States of America **113**(34) 9569-9574.

Morley, R. (2000) *Origin and evolution of tropical rain forests.*John Wiley & Sons.

Morley, R. & Richards, K. (1993) Gramineae cuticle: a key indicator of Late Cenozoic climatic
change in the Niger Delta. *Review of palaeobotany and palynology* **77**(1-2) 119-127.

Naciri, Y. & Linder, H.P. (2015) Species delimitation and relationships: the dance of the seven veils.
Taxon **64**(1) 3-16.

Nie, Z., Sun, H., Manchester, S.R., Meng, Y., Luke, Q. & Wen, J. (2012) Evolution of the
intercontinental disjunctions in six continents in the Ampelopsis clade of the grape family (Vitaceae).
BMC evolutionary biology **12**(1) 1-13.

Nürk, N.M., Linder, H.P., Onstein, R.E., Larcombe, M.J., Hughes, C.E., Piñeiro Fernández, L. ...
Cutts, V. (2020) Diversification in evolutionary arenas—Assessment and synthesis. *Ecology and*
Evolution **10**(12) 6163-6182.

Olmstead, R.G. (2013) Phylogeny and biogeography in Solanaceae, Verbenaceae and Bignoniaceae: a
comparison of continental and intercontinental diversification patterns. *Botanical Journal of the*
Linnean Society **171**(1) 80-102.

Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V., Underwood, E.C. ...
Morrison, J.C. (2001) Terrestrial Ecoregions of the World: A New Map of Life on EarthA new global
map of terrestrial ecoregions provides an innovative tool for conserving biodiversity. *Bioscience*
51(11) 933-938.

Onstein, R.E., Kissling, W.D., Chatrou, L.W., Couvreur, T.L.P., Morlon, H. & Sauquet, H. (2019)
Which frugivory-related traits facilitated historical long-distance dispersal in the custard apple family
(Annonaceae)? *Journal of Biogeography* **46**(8) 1874-1888.

Paradis, E. & Schliep, K. (2019) ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. *Bioinformatics* **35**(3) 526-528.

Pennington, R.T. & Lavin, M. (2016) The contrasting nature of woody plant species in different
neotropical forest biomes reflects differences in ecological stability. *New Phytologist* **210**(1) 25-37.

Pennington, R.T. & Hughes, C.E. (2014) The remarkable congruence of New and Old World savanna
origins. *New Phytologist* **204**(1) 4-6.

Pennington, R.T. & Dick, C.W. (2010) Diversification of the Amazonian flora and its relation to key
geological and environmental events: a molecular perspective. *Amazonia, Landscape and Species*
evolution(ed. by C. Hoorn & F.P. Wesselingh)Blackwell Publishing, Oxford, UK.

Pennington, R.T. & Ratter, J.A. (2006) *Neotropical savannas and seasonally dry forests: plant*
diversity, biogeography, and conservation. CRC press, Boca Raton, Florida, United States.

Pennington, R.T. & Dick, C.W. (2004) The role of immigrants in the assembly of the South American
rainforest tree flora. *Philosophical Transactions of the Royal Society of London Series B: Biological*
sciences **359**(1450) 1611-1622.

Pennington, R.T., Lavin, M. & Oliveira-Filho, A. (2009) Woody plant diversity, evolution, and
ecology in the tropics: perspectives from seasonally dry tropical forests. *Annual Review of Ecology,*
Evolution, and Systematics **40**437-457.

Pennington, R.T., Lavin, M., Prado, D.E., Pendry, C.A., Pell, S.K. & Butterworth, C.A. (2004)
Historical climate change and speciation: neotropical seasonally dry forest plants show patterns of
both Tertiary and Quaternary diversification. *Philosophical Transactions of the Royal Society of*
London.Series B: Biological Sciences **359**(1443) 515-538.

Pennington, R.T., Prado, D.E. & Pendry, C.A. (2000) Neotropical seasonally dry forests and
Quaternary vegetation changes. *Journal of Biogeography* **27**261-273.

Pezzini, F.F., Dexter, K.G., de Carvalho-Sobrinho, J.G., Kidner, C.A., Nicholls, J.A., De Queiroz,
L.P. & Pennington, R.T. (2020) Phylogeny and biogeography of *Ceiba* Mill.(Malvaceae,
Bombacoideae). *Frontiers of Biogeography* .

Pirie, M.D., Klitgaard, B.B. & Pennington, R.T. (2009) Revision and Biogeography of Centrolobium
(Leguminosae-Papilionoideae). *Systematic Botany* **34**(2) 345-359.

Puttick, M.N. (2019) MCMCtreeR: functions to prepare MCMCtree analyses and visualize posterior
ages on trees. *Bioinformatics* **35**(24) 5321-5322.

Python Software Foundation. (2010) Python Language Reference, version 2.7. *http://www.python.org*
.

R Development Core Team. (2013) R: A language and environment for statistical computing.
Available at: http://www.R-project.org/.

Rabosky, D.L., Mitchell, J.S. & Chang, J. (2017) Is BAMM flawed? Theoretical and practical
concerns in the analysis of multi-rate diversification models. *Systematic Biology* .

Rabosky, D.L., Donnellan, S.C., Grundler, M. & Lovette, I.J. (2014) Analysis and visualization of
complex macroevolutionary dynamics: an example from Australian scincid lizards. *Systematic*
Biology **63**(4) 610-627.

Rambaut, A. (2014) FigTree, a graphical viewer of phylogenetic trees (tree.bio.ed.ac.
uk/software/figtree). **1.4.2**.

Rambaut, A., Suchard, M.A., Xie, D. & Drummond, A.J. (2015) Tracer.
http://tree.bio.ed.ac.uk/software/tracer/ **1.6**.

Ree, R.H. & Sanmartín, I. (2018) Conceptual and statistical problems with the DEC J model of
founder□event speciation and its comparison with DEC via model selection. *Journal of Biogeography*
45(4) 741-749.

Ree, R.H. & Smith, S.A. (2008) Maximum likelihood inference of geographic range evolution by
dispersal, local extinction, and cladogenesis. *Systematic Biology* **57**(1) 4-14.

Renner, S. (2004) Plant dispersal across the tropical Atlantic by wind and sea currents. *International*
Journal of Plant Sciences **165**(S4) S23-S33.

Revell, L.J. (2012) phytools: an R package for phylogenetic comparative biology (and other things).
Methods in Ecology and Evolution **3**(2) 217-223.

Richards, P.W. (1973) Africa, the 'odd man out'. *Tropical forest ecosystems in Africa and South*
America: a comparative review.(ed. by B.J. Meggers, E.S. Ayensu, and W.D. Duckworth), pp. 21-26.
Smithsonian Institution Press, Washington, DC, USA.

Ringelberg, J.J., Zimmermann, N.E., Weeks, A., Lavin, M. & Hughes, C.E. (2020) Biomes as
evolutionary arenas: Convergence and conservatism in the trans☐continental succulent biome. *Global*
Ecology and Biogeography **29**(7) 1100-1113.

Rivera☐Ocasio, E., Aide, T.M. & McMillan, W.O. (2002) Patterns of genetic diversity and
biogeographical history of the tropical wetland tree, Pterocarpus officinalis (Jacq.), in the Caribbean
basin. *Molecular ecology* **11**(4) 675-683.

Rojo, J.P. (1972) *Pterocarpus (Leguminosae - Papilionaceae): Revised for the world.*Cramer, Lehre,
Germany.

Rommerskirchen, F., Condon, T., Mollenhauer, G., Dupont, L. & Schefuss, E. (2011) Miocene to
Pliocene development of surface and subsurface temperatures in the Benguela Current system.
Paleoceanography **26**(3) .

Ronquist, F. (1997) Dispersal-vicariance analysis: a new approach to the quantification of historical
biogeography. *Systematic Biology* **46**(1) 195-203.

Sanmartín, I. (2014) Biogeography. *The Tree of Life.*(ed. by P. Vargas and R. Zardoya), pp. 555-576.
Sinauer Associates, Inc., Sunderland, MA, USA.

Sanmartín, I. (2012) Historical Biogeography: Evolution in Time and Space. *Evolution education &*
outreach **5**(4) 555-568.

Sanmartín, I. & Ronquist, F. (2004) Southern hemisphere biogeography inferred by event-based
models: plant versus animal patterns. *Systematic Biology* 216-243.

Saslis☐Lagoudakis, C., Chase, M.W., Robinson, D.N., Russell, S.J. & Klitgaard, B.B. (2008)
Phylogenetics of neotropical Platymiscium (Leguminosae: Dalbergieae): systematics, divergence
times, and biogeography inferred from nuclear ribosomal and plastid DNA sequence data. *American*
Journal of Botany **95**(10) 1270-1286.

Saslis-Lagoudakis, C.H., Klitgaard, B.B., Forest, F., Francis, L., Savolainen, V., Williamson, E.M. &
Hawkins, J.A. (2011) The use of phylogeny to interpret cross-cultural patterns in plant use and guide
medicinal plant discovery: an example from Pterocarpus (Leguminosae). *PloS one* **6**(7) e22275.

Schley, R.J., de la Estrella, M., Pérez-Escobar, O.A., Bruneau, A., Barraclough, T., Forest, F. &
Klitgård, B. (2018) Is Amazonia a 'museum' for Neotropical trees? The evolution of the Brownea
clade (Detarioideae, Leguminosae). *Molecular Phylogenetics and Evolution* **126**279-292.

Schrire, B.D., Lavin, M. & Lewis, G.P. (2005) Global distribution patterns of the Leguminosae:
insights from recent phylogenies. 375-422.

Scotese, C.R. (2004) A continental drift flipbook. *The Journal of geology* **112**(6) 729-741.

Senut, B., Pickford, M. & Ségalen, L. (2009) Neogene desertification of Africa. *Comptes Rendus*
Geoscience **341**(8-9) 591-602.

Setterfield, S.A. (2002) Seedling establishment in an Australian tropical savanna: effects of seed
supply, soil disturbance and fire. *Journal of Applied Ecology* **39**(6) 949-959.

Karline Soetaert(2012) *R Package shape: functions for plotting graphical shapes, colors..., version 1.*
Karline Soetaert.

Stamatakis, A. (2014) RAxML Version 8: A tool for Phylogenetic Analysis and Post-Analysis of
Large Phylogenies. *Bioinformatics* **Open access.**Published online: 10.1093/bioinformatics/btu033.

ter Steege, H., Prado, P.I., de Lima, R.A., Pos, E., de Souza Coelho, L., de Andrade Lima Filho,
Diogenes ... Castilho, C.V. (2020) Biased-corrected richness estimates for the Amazonian tree flora.
Scientific reports **10**(1) 1-13.

ter Steege, H., Pitman, N.C., Sabatier, D., Baraloto, C., Salomao, R.P., Guevara, J.E. ... Silman, M.R.
(2013) Hyperdominance in the Amazonian tree flora. *Science* **342**(6156) 1243092.

Töpel, M., Zizka, A., Calió, M.F., Scharn, R., Silvestro, D. & Antonelli, A. (2017) SpeciesGeoCoder:
fast categorization of species occurrences for analyses of biodiversity, biogeography, ecology, and
evolution. *Systematic Biology* **66**(2) 145-151.

Valcárcel, V. & Wen, J. (2019) Chloroplast phylogenomic data support Eocene amphi□Pacific early
radiation for the Asian Palmate core Araliaceae. *Journal of Systematics and Evolution* **57**(6) 547-560.

van der Pijl, L. (1982) Ecological Developments in Leguminous Fruits. *Principles of Dispersal in*
Higher Plants(ed. by Anonymous), pp. 150-159. Springer.

Vatanparast, M., Klitgård, B.B., Adema, F.A., Pennington, R.T., Yahara, T. & Kajita, T. (2013) First
molecular phylogeny of the pantropical genus *Dalbergia*: implications for infrageneric
circumscription and biogeography. *South African Journal of Botany* **89**143-149.

Vianna, J.A., Fernandes, F.A., Frugone, M.J., Figueiró, H.V., Pertierra, L.R., Noll, D. ... Parker, P.
(2020) Genome-wide analyses reveal drivers of penguin diversification. *Proceedings of the National*
Academy of Sciences **117**(36) 22303-22310.

Wakeling, J.L., Staver, A.C. & Bond, W.J. (2011) Simply the best: the transition of savanna saplings
to trees. *Oikos* **120**(10) 1448-1451.

Waser, N.M. & Campbell, D.R. (2004) Ecological speciation in flowering plants. *Adaptive*
speciation(ed. by Anonymous), pp. 264-277. Cambridge University Press Cambridge.

Weinell, J.L., Barley, A.J., Siler, C.D., Orlov, N.L., Ananjeva, N.B., Oaks, J.R., Burbrink, F.T. &
Brown, R.M. (2021) Phylogenetic relationships and biogeographic range evolution in cat-eyed snakes,
Boiga (Serpentes: Colubridae). *Zoological Journal of the Linnean Society* **192**(1) 169-184.

Wisz, M.S., Hijmans, R.J., Li, J., Peterson, A.T., Graham, C.H., Guisan, A. & NCEAS Predicting
Species Distributions Working Group. (2008) Effects of sample size on the performance of species
distribution models. *Diversity and Distributions* **14**(5) 763-773.

Wolfe, K.H., Li, W.H. & Sharp, P.M. (1987) Rates of nucleotide substitution vary greatly among
plant mitochondrial, chloroplast, and nuclear DNAs. *Proceedings of the National Academy of*
Sciences of the United States of America **84**(24) 9054-9058.

Woodward, F.I., Lomas, M.R. & Kelly, C.K. (2004) Global climate and the distribution of plant
biomes. *Philosophical Transactions of the Royal Society of London.Series B: Biological Sciences*
359(1450) 1465-1476.

Yang, Z. (2007) PAML 4: phylogenetic analysis by maximum likelihood. *Molecular biology and*
evolution **24**(8) 1586-1591.

Zachos, J., Dickens, G.R. & Zeebe, R.E. (2008) An early Cenozoic perspective on greenhouse
warming and carbon-cycle dynamics. *Nature* **451**(7176) 279-283.

Zachos, J., Pagani, M., Sloan, L., Thomas, E. & Billups, K. (2001) Trends, rhythms, and aberrations
in global climate 65 Ma to present. *Science* **292**(5517) 686-693.

Zamora, N. (2000) Nuevas especies y combinaciones en Leguminosas de Mesoamérica. *Novon* 175-
180.

Zhang, C., Rabiee, M., Sayyari, E. & Mirarab, S. (2018) ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. *BMC Bioinformatics* **19**(6) 153.

Zhang, Z., Ramstein, G., Schuster, M., Li, C., Contoux, C. & Yan, Q. (2014) Aridification of the
Sahara desert caused by Tethys Sea shrinkage during the Late Miocene. *Nature* **513**(7518) 401-404.

Zizka, A., Carvalho☐Sobrinho, J.G., Pennington, R.T., Queiroz, L.P., Alcantara, S., Baum, D.A.,
Bacon, C.D. & Antonelli, A. (2020) Transitions between biomes are common and directional in
Bombacoideae (Malvaceae). *Journal of Biogeography* **47**(6) 1310-1321.

Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Duarte Ritter, C., Edler, D. ... Scharn, R. (2019)
CoordinateCleaner: Standardized cleaning of occurrence records from biological collection databases.
Methods in Ecology and Evolution **10**(5) 744-751.

Significance statement

Dispersal involves organisms spreading from one location to another, and is one of the major
processes defining the distribution of organisms. Sometimes, structures which evolved for local
dispersal may lead to rare, long-distance dispersal, and this may allow species to colonise new
landmasses and diversify in novel biomes. As such, here we build an evolutionary tree to understand
relationships between a genus of rosewood species (*Pterocarpus*), and test whether dispersal traits and
biome switching have impacted their evolution. We found that dispersal and biome switching, likely
precipitated by climate change in the Miocene, impacted the modern distribution of Pterocarpus, but
that this was not significantly explained by dispersal structures.

<u>Biosketch</u>

Rowan Schley is a postdoctoral researcher interested in the evolution, biogeography and conservation
of tropical biodiversity. His research is broadly centred around speciation with a particular focus on
the effects of introgression, using phylogenomic and population genomic approaches to address these
questions in tropical trees.

R.J.S. conceived the hypotheses, led data analysis and wrote the manuscript. M.Q. and P.M. led the
phylogenomic data generation. M.V., M.d.l.E. and G.L. provided many comments and background
information on the Legume family for the manuscript. B.K. helped conceive the study and led sample
collection, and provided many comments on the manuscript. All authors contributed to editing the
manuscript.