An epigenomic shift in amygdala marks the transition to maternal behaviors in alloparenting virgin female mice

*Christopher H. Seward[1,2], *Michael C. Saul[2,4], Joseph M. Troy[2], Payam Dibaeinia[3], Huimin Zhang[2], Saurabh Sinha[2,3] and Lisa J. Stubbs[1,2,5]

[1]Pacific Northwest Research Institute, Seattle, WA 98122,
[2] Carl R. Woese Institute for Genomic Biology, and [3]Department of Computer Science, University of Illinois at Urbana-Champaign, Urbana IL 61801

[4] Current address: The Jackson Laboratory, Bar Harbor, ME 04609
[5] Corresponding author, lstubbs@pnri.org
*These authors contributed equally

ABSTRACT

In many species, adults care for young offspring that are not their own, a phenomenon

called alloparenting. However, most nonparental adults must be sensitized by repeated or

extended exposures to newborns before they will robustly display parental-like behaviors. To

capture neurogenomic events underlying the transition to active parental caring behaviors, we

analyzed brain gene expression and chromatin profiles of virgin female mice co-housed with

mothers during pregnancy and after birth. After an initial display of antagonistic behaviors and a

surge of defense-related gene expression, we observed a dramatic shift in the chromatin

landscape specifically in amygdala of the pup-exposed virgin females, accompanied by a

dampening of anxiety-related gene expression. This epigenetic shift coincided with

hypothalamic expression of the oxytocin gene and the emergence of behaviors and gene

expression patterns classically associated with maternal care. The results outline a neurogenomic

program associated with dramatic behavioral changes and suggest molecular networks relevant

to human postpartum mental health.

INTRODUCTION

Interactions between newborn animals and their parents are profoundly important, being

critical to the well-being of the offspring and intensely consequential to the parents as well. In

most mammals, parental care is typically relegated to the female that bears the offspring, with

hormonal shifts that occur during pregnancy and the early postpartum period priming her for this

experience. These dramatic hormonal shifts also alter a mother's morphology, physiology, and

brain structure in ways that persist far beyond the initial parenting experience [1,2]. In addition

to these physical changes [3], mothering also alters a female's behavior, in both the immediate

and the longer-term. In particular, the sight, sounds, and odors of newborns – which may

otherwise be perceived as aversive by adults – become intensely rewarding and motivating to the

mother [4–6]. As with other changes associated with parenting, the shift from aversion to

intense affiliation and reward is coordinated by steroid hormones and a rapid surge in

neuropeptide secretion around the time of birth [3]. Most significantly, a surge of oxytocin,

stored during pregnancy within the paraventricular and supraoptic nuclei of the hypothalamus

[7], is released to target neurons within a brain circuit central to fear, aversion, reward, and the

evaluation of emotional salience [8].

New mothers are not the only individuals that can experience this switch to pup-

affiliative behaviors. For example, although virgin female rats display a clearly aversive

response to pup stimuli, this response can be overcome by the process of sensitization, which

involves a series of repeated interactions; after sensitization, virgin rats will display robust

maternal behaviors – hereafter referred to as maternal behaviors for simplicity – with pups [6].

In contrast, adult virgin female mice do not display any obvious sign of aversion and will instead

spontaneously display certain maternal behaviors shortly after given first access to young pups

[9,10]. Sensitization enhances this response; virgin female mice repeatedly exposed to pups

significantly increases both the range and intensity of maternal behaviors [11]. Intriguingly, it

has been shown virgin female mice continuously co-housed with new mothers will display

maternal behaviors more rapidly under the instruction and encouragement of the mothers, a

process that depends upon the activation of oxytocin neurons [12]. Like mothering itself, this

experience of caring for young that are not one's own, or alloparenting, impacts future behavior.

For example, juvenile female rats that have had the experience of "babysitting" younger siblings

are highly motivated to display maternal behaviors in future encounters with pups [13], and

sensitized adult virgin female mice demonstrate enhanced parenting skills when they have their

offspring of their own [11,14,15]. Indeed, many of the mechanisms that reshape a mother's brain

and behavior also appear to operate in alloparenting females, where intriguingly, they are

activated without the hormonal priming stimulated by pregnancy, parturition, and nursing.

Here, we investigated the functional genomics profile of the brains of co-housed virgin

female mice as they transitioned from pup-naïve to a robust display of alloparenting behaviors

toward pups. To identify genes modulated during this transition, we examined alterations in gene

expression in multiple brain regions over several days of continuous pup exposure. Because

histone modifications have been implicated as central to the behavior of both new mothers and

sensitized virgins [16], we also investigated chromatin accessibility profiles using H3K27Ac

(histone H3 acetylated at lysine 27), a marker of open chromatin, in the same brain regions. The

data reveal defense-related neurogenomic pathways that are silenced, and others that are

activated, across the brains of co-housed alloparenting virgins as they transition to maternal

behaviors and confirm an active role for chromatin remodeling in this behavioral switch,

especially within the amygdala.

RESULTS

Antagonistic behavior, followed by active nurturance in virgins co-housed with mothers

and pups.

In the most common version of rodent pup sensitization experiments, the pups are placed

into the cage of a virgin female for short periods, then removed to be fed, repeatedly over the

course of several consecutive days [11]. However, over years of mouse breeding, we had

observed that nulliparous females co-housed with nursing mouse dams and their litters will

display maternal behaviors toward the pups, suggesting a way to achieve more continuous,

longer-term interaction. Indeed, a recent study has demonstrated that in this context, virgins

respond more quickly to the pups as they are actively instructed and encouraged by the mothers

[12]. This co-housing paradigm provided us with an excellent opportunity to measure the brain's

functional genomic response to pups in the virgin females over time as they transition to

maternal care. To document the timing of this transition, we co-housed four pairs of virgins and

pregnant dams and filmed activity in the cages from early pregnancy though the fourth postnatal

day (**File S1**; Files S1 and S2 available at https://trackhub.pnri.org/stubbs/ucsc/public/allo.html).

For purposes of this study, we were primarily interested in the interactions between virgins and

mothers, on the one hand, and virgins and pups one the other. Therefore, as a primary indicator

of these interactions, we scored the virgins for pup-grooming and mother-grooming behaviors

during 5-minute intervals at the top of each hour; summing the scores in each cage over 6-hour

periods coordinated with the light/dark cycle, providing a useful summary of the overall

behavioral patterns (**Table S1**). Throughout, the two females were most often found together,

interacting or resting in the shared nest, and grooming each other regularly while awake

throughout the observation period. In contrast, although the virgins began to investigate the pups

immediately after birth, they did not begin licking and grooming the pups consistently until

around postnatal day 2 (P2), after which we increasingly observed the virgins engaged in pup

licking/grooming behavior (**Fig. 1A; Table S1**). To test the hypothesis that pup-focused

grooming increased for the virgins while mother-focused grooming did not, we selected data

binned for 6 hours around 12:00 (the beginning of the dark period during lights-out) (**Fig. 1B**).

Pup-focused grooming bouts significantly differed across days P1-P4 (repeated measures

ANOVA, $F_{3,9} = 9.91$, $p = 0.003$), increasing over time, while mother-focused grooming bouts did

not significantly differ across days P1-P4 ($F_{3,9} = 0.67$, $p = 0.59$).

We also observed additional behaviors that are worth noting here. For example, as

described by Carcea and colleagues [12], we observed mothers steering virgin cagemates that

had wandered off to feed or explore back to the nest; often this involved the mothers grabbing

the virgins by the base of the tail and actively pushing them to the nest and pups. Afterward, the

mother would herself typically leave the nest to feed, leaving the virgin to care for the pups. Not

described in the published study but displayed by all virgins we recorded here, we also observed

signs of early antagonism toward the pups. Specifically, during the first two postnatal days we

observed the virgins grabbing pups in their mouths and actively tossing them or pushing them

out of the shared nest (**Table S1, examples of both behaviors in File S2**). By P3, this behavior

was no longer observed, as the virgins spent more time in the nests, licking and grooming the

pups with increasing frequency in classic hunched or prone nursing postures (**Table S1**), similar

to behavior documented for sensitized female rats [17]. Together, these observations suggested

that we could indeed capture the transition from the possibly antagonistic pup-naïve state to

robust pup affiliation between postnatal days 1 and 3 in this continuous-exposure paradigm.

Figure 1. Behaviors exhibited by virgin females co-housed with new mothers and pups over six days beginning the day before birth. (A) Four cages of co-housed mothers and virgin females were recorded over a period of several days before and after birth and grooming behaviors (virgins to pups, plotted in orange; or virgins to mother, blue) were scored (0 or 1) in 5-minute intervals at the beginning of each hour from 12:00 am (0:00) the day before the birth (E18) through the end of the fourth postnatal day (P4), then scores were summed over 6 h periods. To generate the graph, 6 h summed scores were plotted for the four cages as box-and-whisker plots. Times shown mark the end of each 6h period scored. Blue shading in each plot shows the "lights out" periods (12 h beginning at 12:00 pm) for each day. All pups were born within a 6-hour period at the beginning of the light phase on the day designated as P0 for that particular cage, as marked with a bar below each graph. Behaviors plotted and colors used are shown above each graph. Blue arrows below each graph show the times of day that samples were taken from similarly co-housed pairs for gene expression and chromatin analysis. **(B)** The frequency with which co-housed virgins groomed pups (plotted in orange) increased significantly over days (p=0.003), as illustrated by a plot focused on 12:00 pm (start of lights out), while the frequency with which virgins groomed mothers (blue) did not change. Values plotted are mean ± standard error.

Hormone- and neurotransmitter-related genes are dynamically expressed throughout the virgin brain during the first three days of pup interaction. To understand the functional genomic underpinnings of this behavioral transition, we collected RNA from the brains of five virgin females co-housed with a pregnant dam at each of three time points: before birth (2 hr into the dark period of embryonic day 18, or E18) and at the same time during postnatal day 1 (P1)

Figure 2. Numbers of genes up- (orange) or down (blue)-regulated in brains of pup-exposed compared to non-exposed virgin females over time. Numbers represent all genes identified as differentially expressed at fdr ≤ 0.05 in each set of pairwise comparisons. H=hypothalamus; A=amygdala; FC=frontal cortex; S=striatum; P1=postnatal day 1, P3=postnatal day 3.

and P3. We collected and sequenced RNA from four brain regions involved in pup response,

aversion, affiliation, and reward: hypothalamus, amygdala, striatum, and frontal cortex.

At P1, we saw an intense transcriptomic response in amygdala with very little expression

change in other brain regions; by P3, relatively large numbers of differentially expressed genes

(DEGs) were detected in both amygdala and hypothalamus (**Fig. 2; Table S2**). The

transcriptomic response did not correlate simply with expression of immediate early genes

(IEGs) such as *Fos*, which is classically used to mark neuronal activity [18]. However, the IEG

Npas4, which has been implicated specifically in social recognition [19] and reward-related

behaviors [20], was upregulated at P3 in all brain regions tested (**Fig. 3**). Focusing first on the

hypothalamus, genes encoding neuropeptide hormones oxytocin (*Oxt*) and prolactin (*Prl*) were

first up-regulated at P3, when the virgins were beginning to consistently display maternal

behaviors (**Fig. 1, Fig. 3**); these hormones are central to initiation of maternal response in both

mothers and alloparenting virgins [12,21,22]. At P1, dopaminergic (DA) signaling components

including *Drd1* were down-regulated along with related Gene Ontology (GO) and functional

categories such as morphine addiction and behavioral despair. However, *Drd1* returned to pre-

exposure levels in hypothalamus at P3, at the same time that genes related to the activity of

dopaminergic neurons were significantly up-regulated.

Therefore, a switch from repressed to increased dopamine-related gene expression was

coordinated with the increase of *Oxt* and *Prl* expression in the hypothalamus. This pattern is

similar to that observed in mothers at the time of birth and is consistent with the role of DA

signaling in OXT and PRL release [23,24]. It is also consistent with recent observations from

single-cell sequencing that show DA neurons in the hypothalamic preoptic area to be activated in

maternally behaving animals [25]. The data suggested that a shift to a neuropeptide and

neurotransmitter environment favoring stable maternal behavior was developing in the

hypothalamus at P3, concomitant with increased expression of alloparenting behaviors in the

virgin mice. Furthermore, in light of the hypothesized role of histone modifications on maternal

behavior [16,26,27], it is also worth noting the low-level but coordinated up-regulation of genes

encoding chromatin remodeling and binding proteins (*Hdac10, Hdac7, Sirt6,* and *Sirt7*, *Kmt5c*,

Smarcd3, Atrx, L3mbtl1) which we observed in the virgin hypothalamus at P3. This coordinated

shift suggested the existence of a subtle but significant epigenetic response in the hypothalamus

around or before that time (**Table S2A**).

In striking contrast to hypothalamus, components of DA signaling were coordinately *up-*

regulated in the amygdala at P1, along with genes encoding endogenous opioids, proenkephalin

(*Penk*)*,* and prodynorphin (*Pdyn)*. The combined up-regulation of these genes led to P1

enrichment of multiple functional categories indicating that the virgin females were experiencing

stress and anxiety during the first day after the birth of the pups (**Fig. 3; Table S3**). At P3, many

of the anxiety-related amygdala DEGs had returned to pre-exposure levels or were down-

regulated compared to E18 controls, suggesting that the initial P1 surge of transcription for these

genes might be actively silenced. At the same time that the surge of anxiety-related genes was

suppressed, the GO biological process category "maternal behavior" was identified as being

enriched in P3 up-regulated genes (**Fig. 3**).

Figure 3. Enrichment of differentially expressed genes in functional categories. Differentially expressed genes (identified at fdr≤0.05 and with absolute value of fold change ≥1.5) were used to identify enriched functional categories using the ToppCluster tool (Kaimal et al., 2010), as described in Methods. Top panel shows differential expression levels for selected genes, as described in the text. Categories shown are a representative subset of the full report included in Table S3, with up- or down-regulation and category enrichment levels displayed as a heat map. Colored cells denote up- (orange) or down-regulation (blue) for Selected Genes (top panel) or Functional Categories (lower panel) in each brain region/time.

Frontal cortex tracked the amygdala closely in terms of DEGs, direction of change, and enriched functional categories, with a few notable exceptions. In particular, genes related to thyroid hormone activity were uniquely downregulated in the virgin frontal cortex at P3, a finding that is especially interesting given the known role of thyroid hormone in maternal care [28]. Furthermore, in addition to the dopamine-related genes similarly up-regulated in amygdala

and cortex (**Table S2**), a second cadre of genes associated with depressive states, but related to

abnormal thyroid hormone signaling, were down-regulated; the result was that depression-related

functional categories were both up and down-regulated in the frontal cortex DEG set (**Fig. 3;**

Table S3).

Finally, in P3 striatum, down-regulated categories were centered on neuropeptide-related

genes including those encoding vasopression receptor (*Avpr1a*) and prolactin receptor (*Prlr*)

(**Fig. 3**). On the other hand, the gene encoding neuropeptide cholecystokinin (*Cck*), which

positively regulates striatal dopamine signaling in *Drd2*-expressing neurons [29,30] was up-

regulated in striatum at P3 compared to P1. This event is notable, since *Cck* plays a critical role

in the postnatal maintenance of maternal behaviors [31] and mediates responses to anxiety and

reward [32,33]. Together these data indicated that a transcriptomic signature consistent with a

"maternal response" - as it is classically defined by neuropeptide and neurotransmitter gene

expression -was observed in the virgin mice beginning around P3. In particular, the P1 burst of

anxiety-related genes was down-regulated to pre-exposure levels in amygdala and frontal cortex

by this time. In contrast with this response in amygdala, DA signaling was *down-regulated* at

P1, then *up-regulated* at P3 in the hypothalamus of the pup-exposed virgins, concordant with the

onset of maternal behaviors in those mice**.**

Comparison to published datasets.

Parallels to gene expression in brains of new mothers. As referenced above, expression of

several key markers that have been identified in new mothers was also observed in the

alloparenting virgins at the P3 time point. An obvious next question was whether and how gene

expression aligned more globally between maternal and alloparenting virgin brains. Most

published maternal datasets were generated with distinct hypotheses and biological questions in

mind, investigating brain regions and time points very different from ours, complicating direct

comparisons. Nevertheless, two published data series warrant some discussion.

In the first series, Gammie and colleagues used microarrays to compare gene expression

between virgins (not exposed to pups) and nursing females at P7, after maternal behaviors have

been robustly established [34–37]. The same group later completed a meta-analysis of their data

to identify genes that were commonly dysregulated across the maternal brain. Despite the

differences in methods, time points selected, and brain regions examined, we noted that DEGs

identified in the meta-analysis were enriched in similar GO categories, pathways, and disease

associations to those we identified as most pronounced in the pup-exposed virgin brains: neuron

development, addiction, mental health disorders, and pathways involving oxytocin, vasopressin,

prolactin, and opioids [38]. The similarity suggests commonalities between maternal behavior

and alloparenting behavior.

A second published data series examined maternal gene expression over a wide range of

time points pre-and post-partum including P1 and P3, and importantly, used experimental and

statistical methodology very similar to ours [39]. However, cortex (neocortex in the maternal

study, which includes frontal cortex and additional cortical regions) and hypothalamus were the

only brain regions examined commonly in both studies. This similarity allowed us to use formal

statistical techniques to measure the degree of overlap between gene sets from our study and this

previously collected dataset. Using a hypergeometric test to compare gene expression in pup-

exposed virgins and mothers (**Table S2B, S2C**), we found that DEGs up-regulated in the

hypothalamus of P3 virgins correlated positively and most significantly with genes up-regulated

in the maternal hypothalamus at P10 (**Table 1**).

Table 1. Significant correlations between differential gene expression in specific brain regions of pup-exposed virgins and new mothers, or virgins and socially challenged male mice. Hypothalamus (H), Amygdala (A), Frontal cortex (FC) and Neocortex (NC). Highest correlations for all comparisons involving at least 3 overlapping genes are shown, for a full list of comparisons see Table S2.

Virgin dataset/ Maternal Dataset[1]	p value	example genes
HP3-up/HP10-up	1.21E-16	Prl, En1, Slc6a3, Slc10a4, Cryab, Mif, Mfge8
HP3-up/HP10-down	9.12E-10	Egr1, Fos, Junb, Lamb2, Col6a1, Col6a2
HP3-up/HP3-up	2.16E-07	Prl, Nxph4
HP3-up/HP1-up	2.99E-07	Prl, Nxph4
HP1-up/HP10-down	1.23E-06	Magel2, Nr1d1, Slc13a4, Ogn
FCP3-up/NCP1-down	1.59E-25	Arc, Fos, Npas4, Celsr3, Igsf9b, Robo3
FCP3-up/NCP3-down	1.06E-21	Arc, Fos, Npas4, Celsr3, Igsf9b, Robo3
FCP1-up/NCP1-up	9.32E-15	Gpr88, Pde10a, Ppp1r1b, Tac1, Rasd2
FCP3-down/NCP10-down	2.54E-11	Sgk1, Nnat, Calb2, Igsf1

Virgin dataset/ Social challenge dataset[2]	p value	example genes
AP1-up/A120-up	3.67E-54	Drd1, Drd2, Rarb, Grp88, Ppp1r1b, Tac1, Tcf7l2
FCP1-up/FC60-up	1.33E-48	Drd1, Drd2, Gpr88, Ppp1r1b, Rxrg, Tac1, Penk
AP3-down/A120-up	4.16E-36	Cdh1, Ogn, Ccn2, Igf2, Fmod, Sgk1, Grin2b
AP1-down/A120-up	7.16E-35	Avp, Ccn2, Grin2b, Gucy1a2
FCP3-down/FC120-up	1.65E-21	Igsf1, Calb2, Nnat, Trh, Gabrq
AP3-down/A120-dn	3.09E-16	Slc17a7, Tbr1, Nrn1, Sv2b, Lmo3, Tafa1

The overlapping genes included several involved in DA neuron development and function (*En1*, *Slc6a3* and *Slc10a4*), and neuroprotection and neuroinflammatory processes (*Cryab, Mif,* and *Mfge8*). On the other hand, *up-regulated* hypothalamic DEGs from P3 virgins also overlapped with genes that were *down-regulate*d in the maternal hypothalamus at P10 (**Table 1**); IEGs (*Egr1, Fos,* and *Junb)* dominated this list along with genes encoding extracellular matrix (ECM) proteins.

We further identified both positively and negatively-correlated overlaps in comparisons between virgin frontal cortex and maternal neocortex. DEGs *up-regulated* in virgin frontal cortex at P3 overlapped significantly with DEGs *down-regulated* in neocortex of mothers at P1 and P3 (**Table 1**); as in hypothalamus, this group of oppositely regulated genes was dominated by IEGs (*Npas4, Arc*) and genes involved in ECM, and more particularly ECM proteins involved in axon pathfinding (*Celsr3, Igsf9b, Robo3*). Interestingly, there was also significant overlap between DEGs *up-regulated* in virgin P1 frontal cortex and P1 maternal neocortex. This cluster

included genes related to the anxiety-related response that, as noted above, were also up-

regulated in the virgin P1 amygdala (*Adora2a, Gpr88, Pde10a, Rasd2, Ppp1r1b, Tac1, Syndigl1*)

(**Fig. 3; Table S2B, S2C**); this finding suggested the possibility that mothers might also

experience a similar anxiety-related reaction soon after pups were born.

More generally, DEGs across the maternal brain showed enrichment in many of the same

functional categories detected in brains of the alloparenting virgin mice [39]. Although direct

comparison of the same brain regions at similar time points will be required for further

clarification, the data are consistent with the idea that virgin and maternal brains activate many

of the same pathways in response to pups. We note the exception of activation of IEGs and

plasticity-related ECM genes to this general pattern.

Gene expression in P1 virgins closely parallels that of socially challenged males. The

similarities between gene expression in mothers and the P3 virgins fits well with the fact that the

virgins were beginning to exhibit maternal behaviors around this time. However, the molecular

events in the virgin frontal cortex and amygdala around P1 remained something of a puzzle. We

noted some similarities between DEGs in the virgin P1 amygdala and frontal cortex and DEGs

previously identified in the same brain regions taken from of male mice undergoing a territory

threat [40], and a hypergeometric test confirmed a very robust correlation (**Table 1; Table S2D**).

DEGs up-regulated in P1 virgin amygdala -and particularly those down-regulated in P3vP1

comparisons - showed especially high levels of overlap with genes up-regulated in the amygdala

of the socially challenged males; frontal cortex DEGs followed a similar pattern. The

overlapping amygdala genes included those associated with dopamine and cholinergic signaling

(e.g. *Drd1, Drd2*) as well as a large cohort of TF-encoding genes (e.g. *Rarb, Foxp1, Neurod2,*

Tcf7l2) (**Table S2E**). The common up-regulation of these genes in the two social contexts

suggests an especially important and common role. The data are consistent with the

interpretation that at P1, the virgin females are experiencing emotions related to fear and threat,

marked by a genomic response that is remarkably similar to that operating in the brains of males

involved in territory defense. Notably, this threat-related P1 transcriptomic response was largely

extinguished in the virgins at P3, as the females began to display maternal behavior toward the

pups.

DEGs cluster into network modules, suggesting regulatory factors with coordinated roles.

To gain insights into the coordination and interactions of regulatory factors involved in these

brain transcriptomic events, we used a weighted gene correlation network analysis (WGCNA)

approach [41] to generate a co-expression network, including 25 co-expression modules

connected by positive or negative links (**Fig. 4A**, **Table S4A-D).** DEGs from particular brain

regions and time points clustered strongly within certain network modules, indicating the

coordinated regulation of functionally inter-related genes **(Table S4E**). In particular, the threat-

related genes that were up-regulated in the virgin amygdala at P1 and then down-regulated at P3

compared to P1, were especially highly enriched in module 3, with modules 7 and 8 showing a

similar but less robust enrichment pattern (**Fig. 4B**). These modules included most of the genes

that were similarly expressed in P1 virgins and socially challenged males (**Table S2D**). The three

positively correlated modules also included several sets of known interacting genes and DEGs

with related functions. For example, Module 3 includes *Drd1* and *Penk* together with TF genes

Rarb and *Foxp1***,** both of which are important to development and activity of development of

dopaminergic neurons [42,43]. Module 7 includes *Drd5* and TF-encoding DEG *Tcf7l2,* which

has been implicated in fear learning [44]; Module 8 includes *Drd2, Pdyn, Tac1,* and *Rxrg,* the

latter encoding RARB dimerization partner, RXRG. Therefore, the DEGs cluster into modules

with inter-related functions, including TFs with known regulatory interactions.

Other DEG classes clustered into distinct network modules. For example, genes down-

regulated at both P1 and P3 clustered together, especially in Modules 4 and 6 (**Table S4E**); P3

up-regulated genes clustered with especially high concentration in Module 15, including the heat

shock factor regulator, *Hsf1,* a neuroprotective factor involved in adaptation to stressful

experience [45]. Other modules displaying more modest levels of amygdala DEG enrichment

reflect brain expression patterns that are strongly correlated with, or anticorrelated to, Modules 3,

6 or 15 and might thus also include regulatory factors involved in cross-module gene activation

or repressive effects (**Table S4E**).

To identify TFs most central to the pup response, we used GENIE3 [46] to reconstruct a

gene regulatory network (GRN) with these same data (Table S5A). We then identified TFs in the

Figure 4. Weighted gene correlation network representing gene expression in four brain regions of virgin female mice. (A) The network, composed of genes expressed in Amygdala, Frontal Cortex, Hypothalamus, and Striatum of pup-exposed and non-exposed virgin controls, consists of 25 modules, each represented by clusters of different color and joined by lines representing positive (blue) or negative (red) eigengene correlations (in both cases showing only those correlations ≥0.6). Numbers have been added to label modules with particular enrichments in amygdala (A) DEGs as referred to in the text. (B-D) representations of the same network but showing the module membership of up- (blue dots) or down-regulated (red dots) DEGs identified in P1 v E18 (A-P1, B), P3 v E18 (A-P3, C), or P3vP1 (A-P3vP1, D) transcriptomic comparisons. Full details of network structure, membership and correlations are provided in Table S4.

network with target gene sets that were most highly enriched in DEGs from each brain region

and time point (**Table S5B-E**). The data pointed clearly to Module 3 TF *Rarb* as the most

central TF in the amygdala P1 transcriptomic response, whereas Module 6 TF genes *Foxc2,*

Osr1, and *Prdm6,* all three of which are down-regulated in P3 amygdala, dominated the

amygdala P3 transcriptomic response (Table S5B); brain functions of these Module 6 TFs are not

known. Module 15 TF gene *Hsf1*, which is itself up-regulated in hypothalamus at P3, was the

most highly associated with DEGs in that brain region and time point (Table S5C). Of potential

interest, *Snapc4,* a module 15 TF that activates expression of small nuclear RNAs [47,48]

figured prominently in hypothalamus at both time points, suggesting a role for regulation of

RNA splicing in the hypothalamic response.

A dramatic shift in chromatin landscape during the long-term nurturance experience.

Dynamic changes in amygdala chromatin at the P3 time point. The behavioral adaptations that

follow maternal and alloparenting experiences have long been thought to involve epigenetic

factors [23,26]. We therefore expected that histone modifications could play a key role in the

virgins' transition to maternal care. In particular, we hypothesized that the key genes involved in

the threat reaction we observed at P1 might be actively silenced by these mechanisms as the

virgins began to display maternal behaviors at P3. We tested this hypothesis by carrying out

chromatin immunoprecipitation (ChIP) in chromatin from each of the four brain regions from

virgin females co-housed with mothers at E18 and P3. For these ChIP experiments, we used an

antibody specific to histone 3 acetylated at lysine 27 (H3K27Ac), a general marker for accessible

chromatin [49]).

Consistent with our previous results [40], the ChIP profiles revealed tens of thousands of

open-chromatin peaks in every brain region for both pup-exposed and non-exposed females

(**Table S6**). Since differentially accessible peaks (DAPs) offer a unique window into chromatin

dynamics that may drive the brain response, we paid special attention to DAP regions – defined

as genomic regions in which the relative levels of H3K27Ac were at least two-fold higher or

lower in brains collected at P3 compared to E18 consistently in biological replicate samples at

FDR < 0.05 (**Table S7**). Surprisingly although peaks were identified in similar numbers overall

in the each of the four brain regions, DAPs were virtually absent in the chromatin samples from

hypothalamus at P3 and were found in relatively low numbers in frontal cortex and striatum at

this time point as well. In striking contrast, chromatin from the P3 amygdala contained thousands

of DAPs, either increased (5325 DAPs) or decreased in accessibility (7209 DAPs) at P3

compared to E18 (**Fig. 5A, Table S7A**). To maximize the chances of linking DAPs to specific

DEGs, we focused our attention on the smaller number of DAPs located within 5 kb of the TSS

of an annotated gene (called TSS-DAPs). Altogether we found 2738 TSS-DAPs with decreased

H3K27Ac at P3 compared to E18 abbreviated hereafter as P3-closed DAPs), and 1040 TSS-

DAPs with increased levels of H3K27Ac accumulation at P3 compared to E18 (P3-open DAPs).

Interestingly, all genes associated with P3-open or P3-closed TSS-DAPs, respectively, clustered

into network modules that were also enriched for up- or down-regulated DEGs. For example,

amygdala P3-closed DAPs were particularly enriched for linkage genes in modules 3 and 6,

whereas P3-open DAPs were most likely to be associated with genes in Modules 7 and 8 (**Table**

S4F). The TSS-DAPs were associated with 138 amygdala DEGs, including 22 TSS containing

P3-open DAPs, 115 TSS containing P3-closed DAPs, and 1 TSS hosting DAPs of both types

(**Table S6B**). Given that amygdala DEGs were up-and down-regulated in roughly equal numbers

(Fig. 2) the preponderance of down-regulated genes in TSS-DAPs suggested that alterations in

the chromatin landscape at P3 were primarily focused on silencing amygdala genes. The DEGs

associated with P3-closed DAPs were enriched specifically and significantly in network module

3 (hypergeometric *p* = 2.57E-17), suggesting an especially important role for histone de-

acetylation in silencing this coregulated cluster of threat-associated genes.

Figure 5. Dramatic changes in the amygdala chromatin landscape accompanies the transition to maternal-like behaviors in alloparenting virgin mice. (A) Relative numbers of TSS-associated differentially accessible peaks (DAPs), as measured by >2-fold change in detected levels of H3K27Ac, in the four brain regions tested in this study. Positive numbers represent peaks more accessible at P3 than E18 (P3-open DAPs); negative numbers represent less accessible (P3-closed) peaks. (B-E) Examples of DAPs in *Drd1, Rarb1, Rora, and Nkx6-2* genes, showing normalized H3K27Ac profiles in amygdala chromatin at E18 (A_E18, top track) and P3 (A_P3, bottom track) for each gene. (F) An example of a DAP in chromatin from Striatum (S_E18, S_P3) within the *Cck* gene. Red arrows in each panel point to examples of significant differentially accessible peaks. Full ChIP profiles are available online as a UCSC Browser track hub at https://trackhub.pnri.org/stubbs/ucsc/public/allo.html), and data are available in Tables S5 and S6.

Examples include amygdala P3-closed DAPs associated with TSS of Module 3 genes,

Drd1 and *Rarb* (**Fig 4B, 3C**), P3-closed DAPs associated with TSS and upstream promoter

of *Rora* (Module 6 TF gene down-regulated at both P3 and P30 (**Fig. 4D**), and P3-open DAPs

associated with *Nkx6-2* (Module 1 TF gene up-regulated at P3 timepoint) (**Fig. 4E**).

Together the data suggest that activities of many key genes involved in pup response are

regulated by differential chromatin accessibility, specifically in the amygdala.

Although much smaller in numbers, some DAPs associated with DEGs in other brain

regions also deserve some mention. For example, *Rarb* was also down-regulated in frontal cortex

P3 vs. P1 comparisons, and displayed a pattern of DAPs in cortex very similar to that seen in

amygdala (**Table S6A**). Additional DAPs that may be relevant to gene expression were

discovered in brain regions other than amygdala by lowering the fold-change cutoff to 1.5

instead of 2 (1.5X vs 2X), while keeping the same replicate FDR significance threshold (FDR <

0.05). For example, a 2X P3-open DAP in striatal chromatin was identified approximately 25

Kb downstream of *Cck*, and several 1.5X P3-closed DAPs were found within and closer to the

gene (**Table S6A; Fig. 4F**). Since *Cck* was up-regulated in striatum at P3, this chromatin

configuration suggests the possible role for chromatin dynamics in the regulation of this critical

gene.

Enrichment of binding motifs points to mechanistic insights. To obtain further information

regarding the potential activity of TFs in the pup response, we searched for enrichment of known

TF binding motifs (TFBMs) in the P3-open and P3-closed amygdala TSS DAPs. The search

identified REST/NRSF binding motifs as the top enrichment within *P3 closed* TSS-DAPs (E18

enriched compared to P3); although *Rest* itself was not identified as differentially expressed, the

data suggest that REST TFBMs were being closed between E18 and P3 in amygdala of the pup-

exposed virgins. This finding is of interest, because REST is a central regulatory of neuron

differentiation and plasticity [50], and also plays a role in stress resilience in adult brain [51].

The search also identified enrichments for motifs in the P3-closed DAPs that are recognized by

TFs encoded by amygdala DEGs, including P3 down-regulated genes *Mef2c* and *Rora* (Table

394 S2). Consistent with their expression, the TSS of both genes were associated with P3-closed

395 DAPs (**Fig. 4D; Table S7B**). Together these data indicate that histone deacetylation events

396 evident at P3 serve to not only reduce levels of *Mef2c* and *Rora* gene expression, but

397 simultaneously, to reduce the accessibility of both TFs to their target genes. Notably, both TF

398 genes have been associated with deficits in social behavior [52–54] and *Rora* has been

399 implicated in maternal behavior specifically [55], supporting a functional role. Notably, *Mef2c*

400 target genes predicted by GRN analysis were significantly and specifically enriched in pup-

401 driven DEGs in amygdala; MEF2C target genes were also predicted to include a notably high

402 number of other TFs (Table S5B). These data suggest that MEF2C may play a role as one of the

403 central hubs coordinating the amygdala transcriptomic response.

404   Some notable TFBM enrichments were also detected in P3-open DAPs. For example, we

405 noted enrichment of FOX family TFs including the specific TFBM of the protein encoded by

406 DEG, *Foxp1* which was up-regulated in amygdala at P1, then down-regulated between P1 and P3

407 (**Table S7**). However, this FOX motif could potentially also be recognized by other family

408 members including FOXC2; *Foxc2* was down-regulated in the P3 amygdala and was predicted in

409 the GRN to be central to the P3 response (Table 3). Because we did not measure chromatin at

410 P1, the initial timing of these epigenetic events is not discernible. However, the data suggest that

411 while TSS-linked targets of REST, RORA and MEF2C became less accessible, targets of FOX

412 family proteins became more accessible via epigenetic modifications during the postnatal period.

414 **DISCUSSION**

415   With the goal of understanding molecular mechanisms that underlie the transition to

416 maternal behaviors, we investigated the behavioral, transcriptomic, and epigenomic response of

virgin females as they were co-housed with mothers and newborn pups over a period of several

days. A recent study used this same paradigm to show that alloparenting virgins are instructed in

pup care by co-housed mothers [12], and we observed a very similar pattern of behaviors in the

mothers and virgins we tested here. Along the way, although virgin female mice did not display

an obvious aversion toward the pups, we also observed evidence of an initial antagonism during

the first two postnatal days; these antagonistic behaviors gave way to increasing levels of

attention to the pups, with the virgins increasingly licking, grooming and huddling over the pups

by postanatal day 3.

The data presented here reveal a dramatic and dynamic neurogenomic shift that coincides

with successful maternal instruction and the activation of oxytocin neurons in the virgin brain.

In particular, at P1 we observed a striking signal of fear and anxiety in the virgin hypothalamus,

frontal cortex and especially in the amygdala, in the form of a gene expression pattern that

correlated with high significance to that observed in territory-challenged males [40]. Despite the

lack of obvious aversion, these data indicate that indeed - at least within this co-housed paradigm

- virgin female mice do initially perceive the pups as anxiety-inducing, or even possibly

threatening. If the threat signal is related to the aversive response observed in rats and other

species, the data would be consistent with the hypothesis that pup aversion and defensive

behaviors share a common brain circuitry [4], and would suggest a shared molecular mechanism

for diverse types of threat response as well. Several TF genes implicated by gene expression,

network co-expression, chromatin analysis, and/or motif-enrichment analysis were similarly up-

or down regulated in the pup-exposed virgins and socially challenged males, suggesting crucial

roles for these TFs in regulating this shared molecular signature of social threat.

Published data support the roles of several of these TFs in threat/anxiety response. For

example, the RARB:RXRG dimer's activities in amygdala have been linked to expression of

anxiety-related phenotypes [56], and *RORA* is associated with enhanced fear response in humans

[57] and mice [58]. Furthermore, *Rora* mutant mouse mothers do not retrieve, care for, or suckle

their young [59]; the data presented here support further investigation of this gene's role in

maternal amygdala. Furthermore*,* other TF genes implicated in the shared threat signature are

associated with social-behavior phenotypes, including *Tcf7l2*, which is up-regulated in the

amygdala of virgins at P1 as well as in socially challenged males [40] and is important for fear-

learning and adaptation [44]. Our data suggest that these TFs have coordinated roles in the fear

response, with *Rarb* playing a central role.

We hypothesize that these and other networked TFs work together to modulate the

response to pups, and the networks developed from our dataset suggests a robust framework of

positive and negative gene interactions that coordinate this behavioral switch over time. The

expression of the threat-related TF genes was extinguished along with the pulse of dopamine

signaling after the rise of oxytocin, prolactin, and other neuropeptides by P3, paving the way for

a shift in the virgin females' behavior toward the pups; we surmise that this shift was driven, at

least in part, by a substantial level of chromatin remodeling in the amygdala. Many of the genes

that returned to normal expression levels between P1 and P3 in amygdala were associated with

differentially accessible chromatin, consistent with their active epigenetic silencing in that brain

region at P3.

Chromatin remodeling has been implicated in the development of maternal behaviors in

both mothers and alloparenting virgins, although most published studies have focused on the

hypothalamic MPOA as the primary site of this epigenetic response [16,26,60]. These studies

have shown that *suppression* of HDAC activity – and thus *inhibition* of chromatin silencing - in

hypothalamus is key to driving the females' maternal response. Surprisingly therefore, we found

no evidence of chromatin remodeling in the P3 hypothalamus, and the massive chromatin

remodeling we did observe in amygdala was weighted toward *de-acetylation,* or chromatin

closure, along with the silencing of differentially expressed genes. These findings would suggest

that HDAC activity plays a positive role in the acquisition of maternal behavior, although it is

certainly possible that histone deacetylation at earlier time points or in different brain regions

could have been crucial. For example, although deacetylation in amygdala may be critical in

quenching the threat response once established, *suppression* of deacetylation in hypothalamus

(and/or other brain regions) before P1 might have prevented the establishment of the

aversive/fear response in the first place. Possibly relevant to this hypothesis is the coordinated

up-regulation of histone deacetylase and chromatin remodeling genes we detected in the

hypothalamus at P3; this signal could reflect the trace of earlier epigenetic events associated with

the expression of fear and anxiety in the virgins before they transitioned to active pup care.

This is the first study to investigate global gene expression in the amygdala in the context

of alloparental care, and supports the idea of including amygdala in future studies with mothers

as well. Our studies highlight a special role for the amygdala in the switch to alloparenting

behavior in this context, a hypothesis that is consistent with the known functions of amygdala in

maternal behavior and bonding [61,62]. As underscored by human brain imaging studies,

maternal behavior involves a global brain response that unfolds over an extended period of pre-

and postnatal time [63]; in the pup-exposed virgins, we detected just the start of this behavioral

transition during the third postnatal day. Nevertheless, the mechanisms involved in this

transition to intensive pup care could be relevant to a successful transition to motherhood as well.

Although it is not yet possible to determine whether a similar response is activated, or actively

suppressed, at some time around birth in the maternal amygdala, this question is an important

one in the context of maternal bonding and infant care. Especially given the similar up-

regulation of anxiety-related genes we identified in published data from the maternal neocortex,

we speculate that a similar active suppression of a threat/anxiety program may occur in the

amygdala of new mothers, and that dysregulation of this program could underlie the failure of

mother-infant bonding, post-partum anxiety and depression. Addressing this hypothesis will

offer a novel perspective on the causes of these very common, painful and highly consequential

human maladies.

MATERIALS AND METHODS

Mice and behavioral analysis

All work with mice was done under the approval of the IACUC at the University of Illinois,

Urbana Champaign. Mice were housed in a temperature-controlled room in a reverse 12h/ 12h

light-dark cycle. Six-week-old female mice (C57BL/6J x C3HJ F1 hybrids, with an agouti coat

to allow clear distinction with the black-coated virgins) purchased from the Jackson Laboratory

were impregnated, and co-housed with age-matched virgin female C57BL/6J mice during

pregnancy and through the early post-partum pregnancy. To record behavior, four pairs were

filmed in clear-topped cages continuously using a Samsung SCB-2000 CCTV Camera with iSpy

64 v7.2.1.0 CCTV software. Behavior was scored in each cage (0 or 1) in 5-minute snapshots at

the top of each hour from the day before and until the end of the fourth day after birth, with

scores combined over 6 h periods for each cage to generate the illustrative plot in Fig. 1. To test

the hypothesis that pup-grooming behaviors increased over time, while mother-grooming

behaviors did not, we performed one-way repeated measures ANOVA in R (v4.0.4) using the

rstatix package (v0.7.0). The anova_test() function in rstatix automatically assesses repeated

measures data for the assumption of sphericity. Scores for these and other behaviors are also

presented in Table S1. The 5-minute video snapshots are provided as File S1 with video clips of

specific and unusual behaviors noted in the text provided as File S2; additional video is available

on request.

Dissections and RNA preparation

Dissections were performed as described in detail previously [40] with the addition of the

striatum. Briefly, mice were euthanized by cervical dislocation followed by rapid decapitation.

Their brains were removed and sectioned in a coronal slicing mouse brain matrix. A total of

three cuts were made: two cuts separated by 4 mm defined by the rostral and caudal aspects of

the hypothalamus and a third cut bisecting these two cuts. The hypothalamus, frontal cortex,

striatum, and amygdala were dissected from the resultant brain slices (**Supplementary Figure**

S1). Upon completion of these dissections, focal brain regions were placed into cryotubes, snap-

frozen in liquid N2, and stored at -80°C until downstream processing. Samples were prepared for

sequencing from the four dissected brain regions of five mice per condition (E18, P1, P3). RNA

isolation and QC were completed as described previously, with libraries prepared robotically at

the Roy J. Carver Biotechnology Center at University of Illinois, also as described in [40].

Gene expression analysis

Illumina sequencing libraries were generated with the TruSeq Stranded mRNA HT kit (Illumina)

using an Eppindorf ePMotion 5075 robot and were sequenced to a depth 45-60 million reads per

sample on Hi-Seq 2500 instruments at the Roy J. Carver Biotechnology Center at the University

of Illinois. All sequencing data generated in this study have been deposited to the GEO database

under Accession number GSE184549. Pairwise comparisons of E18, P1 and P3 samples were

completed as previously described in detail [40], with results provided in Table S2. For

functional analysis, genes that were found to be differentially expressed at fdr < 0.05 were first

filtered for absolute fold change >1.5, and uploaded to the ToppCluster web analysis tool [64]

using default conditions (Bonferroni correction, fdr <0.05). Selected categories are summarized

in Table 2, with full ToppCluster Results reported in Table S3.

Network Analysis

We used signed WGCNA (Langfelder & Horvath, 2008) to generate networks from the data

from all individuals, brain regions, and time points, as described in depth in our previous study

[40]. Eigengenes calculated for each module were used to generate module correlations; details

of module structure, module gene content, eigengene correlations, and hypergeometric

enrichments are presented in Table S4. After log-transforming our data using voom+limma, we

filtered zero variance genes, selected a soft thresholding coefficient of 3, then used a signed

Pearson correlation analysis with a minimum module size of 30. Images in Figure 3 were

generated using version 3.7.1 of Cytoscape [65].

To reconstruct the GRN, we obtained a list of 1523 potential transcription factors in mouse from

Animal Transcription Factor Database [66]. GENIE3 [46] was applied on the expression data of

37991 genes in 53 conditions consisting of various brain regions (H=hypothalamus,

A=amygdala, FC=frontal cortex, and S=striatum) and different time points (E18, P1-P3) to score

the relative significance of each TF-gene interaction. (Auto-regulatory relationships were

excluded). To construct a GRN, for each gene we collected up to top five TF regulators of that

gene as predicted by GENIE3, additionally requiring that the TF-gene pair have a Spearman's

correlation of at least 0.5 (in absolute value) and a GENIE3 score of at least 0.005. The resulting

GRN included 92717 interactions involving 1400 unique TFs and 21156 genes (Table S5A). To

assess the significance of TF regulons in different brain regions, for each TF we computed the

enrichment of its regulon (gene set predicted to be regulated by the TF) for DEGs from each

brain region, using hypergeometric test (Table S5B-E).

ChIP Tissue Preparation, Chromatin Immunoprecipitation, and Library Preparation

ChIP was performed essentially as described in detail in our previous study [40]. Briefly, brain

tissue dissected from 3 animals was pooled, homogenized, and fixed in PBS with 1%

formaldehyde for 10 minutes. Nuclei were prepared from the fixed cells and stored at -80º C

until use. Thawed nuclei were sonicated using a BiorupterTM UCD-200 (Diagenode, Liège,

Belgium) sonicator, and fragmented chromatin was processed for ChIP with 2 ug histone

H3K27Ac antibody per sample (Abcam ab4729), using one million nuclei for each IP. IPs were

performed in biological replicate, with one pool of 3 samples in each replicate, as previously

described. Libraries were prepared from eluted DNA using KAPA LTP library kits (KK8230)

using Bioo Scientific index adapters, size-selected using AmpureXP beads (Beckman Coulter,

Brea, CA, USA) and quality checked by Qubit 2.0 and Bioanalyzer (Agilent 2100). Samples

were sequenced to a depth of 20-30M reads per replicate on an Illumina HiSeq 2500 sequencer

using a TruSeq SBS sequencing kit, version 4, in single-end format with fragment length of 100

bp. Base calling and demultiplexing into FASTQ files was done using bcl2fastq v1.8.4 software

(Illumina, San Diego, CA, USA).

ChIP-Seq Bioinformatics

ChIP sequencing reads were mapped with Bowtie2 [67] to the UCSC Mus musculus mm9 or

mm10 genome, using default settings and analyzed for peaks using HOMER (Hypergeometric

Optimization of Motif EnRichment) v4.7 [68], as previously described [40]. Differential

chromatin peaks were identified in biological replicates using the HOMER

getDifferentialPeaksReplicates.pl script, looking for any peaks that changed at least two-fold

between conditions with an FDR cutoff of 0.05. Known motif discovery was performed with the

HOMER findMotifsGenome.pl script using default settings with 201bp peak regions extracted

from all histone peaks or only differential histone peaks. Chromatin profiles are available online

as a UCSC Genome Browser track hub at https://trackhub.pnri.org/stubbs/ucsc/public/allo.txt).

ACKNOWLEGEMENT

We thank Gene Robinson, Alison Bell, Dave Zhao for helpful discussions throughout this project

and Alison Bell and Elbert Branscomb for critical comments on the manuscript. This work was

supported by the Simons Foundation (#SfLife 291812), with additional funding provided by the

University of Illinois and the Pacific Northwest Research Institute.

LITERATURE CITED

Adachi, M., Lin, P. Y., Pranav, H., & Monteggia, L. M. (2016). Postnatal Loss of Mef2c Results in Dissociation of Effects on Synapse Number and Learning and Memory. *Biol Psychiatry, 80*(2), 140-148. doi:10.1016/j.biopsych.2015.09.018

Aguggia, J. P., Suarez, M. M., & Rivarola, M. A. (2019). Multiparity Dampened the Neurobehavioral Consequences of Mother-Pup Separation Stress in Dams. *Neuroscience, 416*, 207-220. doi:10.1016/j.neuroscience.2019.07.042

Althammer, F., & Grinevich, V. (2017). Diversity of oxytocin neurons: beyond magno- and parvocellular cell types? *J Neuroendocrinol*. doi:10.1111/jne.12549

Araujo, D. J., Anderson, A. G., Berto, S., Runnels, W., Harper, M., Ammanuel, S., . . . Konopka, G. (2015). FoxP1 orchestration of ASD-relevant signaling pathways in the striatum. *Genes Dev, 29*(20), 2081-2096. doi:10.1101/gad.267989.115

Atzil, S., Touroutoglou, A., Rudy, T., Salcedo, S., Feldman, R., Hooker, J. M., . . . Barrett, L. F. (2017). Dopamine in the medial amygdala network mediates human bonding. *Proc Natl Acad Sci U S A, 114*(9), 2361-2366. doi:10.1073/pnas.1612233114

Bridges, R. S. (2015). Neuroendocrine regulation of maternal behavior. *Front Neuroendocrinol, 36*, 178-196. doi:10.1016/j.yfrne.2014.11.007

Brunton, P. J. (2019). Endogenous opioid signalling in the brain during pregnancy and lactation. *Cell Tissue Res, 375*(1), 69-83. doi:10.1007/s00441-018-2948-1

Calemandrei, G., Keverne, E. B. (1994). Differential Expression of Fos Protein in the Brain of Female Mice Dependent on Pup Sensory Cues and Maternal Experience. *Behav Neuro, 108*(1), 113-120.

Damberg, M. (2005). Transcription factor AP-2 and monoaminergic functions in the central nervous system. *J Neural Transm (Vienna), 112*(10), 1281-1296. doi:10.1007/s00702-005-0325-1

Damberg, M., Garpenstrand, H., Alfredsson, J., Ekblom, J., Forslund, K., Rylander, G., & Oreland, L. (2000). A polymorphic region in the human transcription factor AP-2beta gene is associated with specific personality traits. *Mol Psychiatry, 5*(2), 220-224. doi:10.1038/sj.mp.4000691

den Hoed, J., Sollis, E., Venselaar, H., Estruch, S. B., Deriziotis, P., & Fisher, S. E. (2018). Functional characterization of TBR1 variants in neurodevelopmental disorder. *Sci Rep, 8*(1), 14279. doi:10.1038/s41598-018-32053-6

Driessen, T. M., Eisinger, B. E., Zhao, C., Stevenson, S. A., Saul, M. C., & Gammie, S. C. (2014). Genes showing altered expression in the medial preoptic area in the highly social maternal phenotype are related to autism and other disorders with social deficits. *BMC Neurosci, 15*, 11. doi:10.1186/1471-2202-15-11

Dufford, A. J., Erhart, A., & Kim, P. (2019). Maternal brain resting-state connectivity in the postpartum period. *J Neuroendocrinol, 31*(9), e12737. doi:10.1111/jne.12737

Dulac, C., O'Connell, L. A., & Wu, Z. (2014). Neural control of maternal and paternal behaviors. *Science, 345*(6198), 765-770. doi:10.1126/science.1253291

Dulac, C., O'Connell, L. A., Wu, Z. (2014). Neural control of maternal and paternal behaviors. *Science, 345*(6198), 765-770.

Eisinger, B. E., Driessen, T. M., Zhao, C., & Gammie, S. C. (2014). Medial prefrontal cortex: genes linked to bipolar disorder and schizophrenia have altered expression in the highly social maternal phenotype. *Front Behav Neurosci, 8*, 110. doi:10.3389/fnbeh.2014.00110

Eisinger, B. E., Zhao, C., Driessen, T. M., Saul, M. C., & Gammie, S. C. (2013). Large scale expression changes of genes related to neuronal signaling and developmental processes found in lateral septum of postpartum outbred mice. *PLoS One, 8*(5), e63824. doi:10.1371/journal.pone.0063824

Fleming, A. S., & Rosenblatt, J. S. (1974). Maternal behavior in the virgin and lactating rat. *J Comp Physiol Psychol, 86*(5), 957-972. doi:10.1037/h0036414

Frederic, F., Chianale, C., Oliver, C., & Mariani, J. (2006). Enhanced endocrine response to novel environment stress and lack of corticosterone circadian rhythm in staggerer (Rora sg/sg) mutant mice. *J Neurosci Res, 83*(8), 1525-1532. doi:10.1002/jnr.20843

Funahashi, Y., Ariza, A., Emi, R., Xu, Y., Shan, W., Suzuki, K., . . . Kaibuchi, K. (2019). Phosphorylation of Npas4 by MAPK Regulates Reward-Related Gene Expression and Behaviors. *Cell Rep, 29*(10), 3235-3252 e3239. doi:10.1016/j.celrep.2019.10.116

Gammie, S. C., Driessen, T. M., Zhao, C., Saul, M. C., & Eisinger, B. E. (2016). Genetic and neuroendocrine regulation of the postpartum brain. *Front Neuroendocrinol, 42*, 1-17. doi:10.1016/j.yfrne.2016.05.002

Gandelman, R. (1973). Induction of Maternal Nest Building in Virgin Female Mice by the Presentation of Young. *Horm Behav, 4*, 191-197.

Guastavino, J. M. (1983). Constraint of the mother with pups restores some aspects of the maternal behavior of mutant staggerer mice. *Physiol Behav, 30*(5), 771-774. doi:10.1016/0031-9384(83)90176-2

Guissart, C., Latypova, X., Rollier, P., Khan, T. N., Stamberger, H., McWalter, K., . . . Kury, S. (2018). Dual Molecular Effects of Dominant RORA Mutations Cause Two Variants of Syndromic Intellectual Disability with Either Autism or Cerebellar Ataxia. *Am J Hum Genet, 102*(5), 744-759. doi:10.1016/j.ajhg.2018.02.021

Gutman, D. A., & Nemeroff, C. B. (2002). Neurobiology of early life stress: rodent studies. *Semin Clin Neuropsychiatry, 7*(2), 89-95. doi:10.1053/scnp.2002.31781

Haller, J., Harold, G., Sandi, C., & Neumann, I. D. (2014). Effects of adverse early-life events on aggression and anti-social behaviours in animals and humans. *J Neuroendocrinol, 26*(10), 724-738. doi:10.1111/jne.12182

Hamdan, F. F., Daoud, H., Rochefort, D., Piton, A., Gauthier, J., Langlois, M., . . . Michaud, J. L. (2010). De novo mutations in FOXP1 in cases with intellectual disability, autism, and language impairment. *Am J Hum Genet, 87*(5), 671-678. doi:10.1016/j.ajhg.2010.09.017

Harding, K. M., & Lonstein, J. S. (2016). Extensive juvenile "babysitting" facilitates later adult maternal responsiveness, decreases anxiety, and increases dorsal raphe tryptophan hydroxylase-2 expression in female laboratory rats. *Dev Psychobiol, 58*(4), 492-508. doi:10.1002/dev.21392

Harrington, A. J., Bridges, C. M., Berto, S., Blankenship, K., Cho, J. Y., Assali, A., . . . Cowan, C. W. (2020). MEF2C Hypofunction in Neuronal and Neuroimmune Populations Produces MEF2C Haploinsufficiency Syndrome-like Behaviors in Mice. *Biol Psychiatry, 88*(6), 488-499. doi:10.1016/j.biopsych.2020.03.011

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., . . . Glass, C. K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell, 38*(4), 576-589. doi:10.1016/j.molcel.2010.05.004

Heslin, K., & Coutellier, L. (2018). Npas4 deficiency and prenatal stress interact to affect social recognition in mice. *Genes Brain Behav, 17*(5), e12448. doi:10.1111/gbb.12448

Hon, G. C., Hawkins, R. D., & Ren, B. (2009). Predictive chromatin signatures in the mammalian genome. *Hum Mol Genet, 18*(R2), R195-201. doi:10.1093/hmg/ddp409

Huang, T. N., Chuang, H. C., Chou, W. H., Chen, C. Y., Wang, H. F., Chou, S. J., & Hsueh, Y. P. (2014). Tbr1 haploinsufficiency impairs amygdalar axonal projections and results in cognitive abnormality. *Nat Neurosci, 17*(2), 240-247. doi:10.1038/nn.3626

Kaimal, V., Bardes, E. E., Tabar, S. C., Jegga, A. G., & Aronow, B. J. (2010). ToppCluster: a multiple gene list feature analyzer for comparative enrichment clustering and network-based dissection of biological systems. *Nucleic Acids Res, 38*(Web Server issue), W96-102. doi:10.1093/nar/gkq418

Kinsley, C. H., Bales, K. L., Bardi, M., & Stolzenberg, D. S. (2015). Reproductive experiential regulation of cognitive and emotional resilience. *Neurosci Biobehav Rev, 58*, 92-106. doi:10.1016/j.neubiorev.2015.05.015

Kinsley, C. H., Bardi, M., Karelina, K., Rima, B., Christon, L., Friedenberg, J., & Griffin, G. (2008). Motherhood induces and maintains behavioral and neural plasticity across the lifespan in the rat. *Arch Sex Behav, 37*(1), 43-56. doi:10.1007/s10508-007-9277-x

Kinsley, C. H., & Bridges, R. S. (1990). Morphine treatment and reproductive condition alter olfactory preferences for pup and adult male odors in female rats. *Dev Psychobiol, 23*(4), 331-347. doi:10.1002/dev.420230405

Kitaoka, K., Shimizu, M., Shimizu, N., Chikahisa, S., Nakagomi, M., Shudo, K., . . . Sei, H. (2011). Retinoic acid receptor antagonist LE540 attenuates wakefulness via the dopamine D1 receptor in mice. *Brain Res, 1423*, 10-16. doi:10.1016/j.brainres.2011.09.023

Kombian, S. B., Ananthalakshmi, K. V., Parvathy, S. S., & Matowe, W. C. (2004). Cholecystokinin activates CCKB receptors to excite cells and depress EPSCs in the rat rostral nucleus accumbens in vitro. *J Physiol, 555*(Pt 1), 71-84. doi:10.1113/jphysiol.2003.056739

Kombian, S. B., Ananthalakshmi, K. V., Parvathy, S. S., & Matowe, W. C. (2005). Cholecystokinin inhibits evoked inhibitory postsynaptic currents in the rat nucleus accumbens indirectly through gamma-aminobutyric acid and gamma-aminobutyric acid type B receptors. *J Neurosci Res, 79*(3), 412-420. doi:10.1002/jnr.20349

Krzyzosiak, A., Szyszka-Niagolov, M., Wietrzych, M., Gobaille, S., Muramatsu, S., & Krezel, W. (2010). Retinoid x receptor gamma control of affective behaviors involves dopaminergic signaling in mice. *Neuron, 66*(6), 908-920. doi:10.1016/j.neuron.2010.05.004

Kuemerle, B., Gulden, F., Cherosky, N., Williams, E., & Herrup, K. (2007). The mouse Engrailed genes: a window into autism. *Behav Brain Res, 176*(1), 121-132. doi:10.1016/j.bbr.2006.09.009

Langfelder, P., & Horvath, S. (2008). WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics, 9*, 559. doi:10.1186/1471-2105-9-559

Langfelder, P., & Horvath, S. (2012). Fast R Functions for Robust Correlations and Hierarchical Clustering. *J Stat Softw, 46*(11). Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/23050260

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods, 9*(4), 357-359. doi:10.1038/nmeth.1923

Lepagnol-Bestel, A. M., Maussion, G., Ramoz, N., Moalic, J. M., Gorwood, P., & Simonneau, M. (2007). Nrsf silencing induces molecular and subcellular changes linked to neuronal plasticity. *Neuroreport, 18*(5), 441-446. doi:10.1097/WNR.0b013e328011dc81

Lin, C. H., Hansen, S., Wang, Z., Storm, D. R., Tapscott, S. J., & Olson, J. M. (2005). The dosage of the neuroD2 transcription factor regulates amygdala development and emotional learning. *Proc Natl Acad Sci U S A, 102*(41), 14877-14882. doi:10.1073/pnas.0506785102

Lonstein, J. S., Wagner, C. K., De Vries, G. J. (1999). Comparison of the "Nursing" and Other Parental Behaviors of Nulliparous and Lactating Female Rats. *Horm Behav, 36*, 242-251.

Mann, P. E., Felicio, L. F., & Bridges, R. S. (1995). Investigation into the role of cholecystokinin (CCK) in the induction and maintenance of maternal behavior in rats. *Horm Behav, 29*(3), 392-406. doi:10.1006/hbeh.1995.1027

Mayer, H. S., Helton, J., Torres, L. Y., Cortina, I., Brown, W. M., & Stolzenberg, D. S. (2019). Histone deacetylase inhibitor treatment induces postpartum-like maternal behavior and immediate early gene expression in the maternal neural pathway in virgin mice. *Horm Behav, 108*, 94-104. doi:10.1016/j.yhbeh.2018.02.011

Miller, M. W., Wolf, E. J., Logue, M. W., & Baldwin, C. T. (2013). The retinoid-related orphan receptor alpha (RORA) gene and fear-related psychopathology. *J Affect Disord, 151*(2), 702-708. doi:10.1016/j.jad.2013.07.022

Moffitt, J. R., Bambah-Mukku, D., Eichhorn, S. W., Vaughn, E., Shekhar, K., Perez, J. D., . . . Zhuang, X. (2018). Molecular, spatial, and functional single-cell profiling of the hypothalamic preoptic region. *Science, 362*(6416). doi:10.1126/science.aau5324

Nishi, M., Horii-Hayashi, N., & Sasagawa, T. (2014). Effects of early life adverse experiences on the brain: implications from maternal separation models in rodents. *Front Neurosci, 8*, 166. doi:10.3389/fnins.2014.00166

Noirot, M. (1972). Ultrasounds and Maternal Behavior in Small Rodents. *Dev Psychobiol, 5*(4), 371-387.

Numan, M. (2007). Motivational systems and the neural circuitry of maternal behavior in the rat. *Dev Psychobiol, 49*(1), 12-21. doi:10.1002/dev.20198

Ray, S., Tzeng, R. Y., DiCarlo, L. M., Bundy, J. L., Vied, C., Tyson, G., . . . Arbeitman, M. N. (2015). An Examination of Dynamic Gene Expression Changes in the Mouse Brain During Pregnancy and the Postpartum Period. *G3 (Bethesda), 6*(1), 221-233. doi:10.1534/g3.115.020982

Rosenblatt, J. S., Mayer, A. D., & Giordano, A. L. (1988). Hormonal basis during pregnancy for the onset of maternal behavior in the rat. *Psychoneuroendocrinology, 13*(1-2), 29-46. doi:10.1016/0306-4530(88)90005-4

Rotzinger, S., & Vaccarino, F. J. (2003). Cholecystokinin receptor subtypes: role in the modulation of anxiety-related and reward-related behaviours in animal models. *J Psychiatry Neurosci, 28*(3), 171-181. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/12790157

Russell, J. A., Leng, G., & Douglas, A. J. (2003). The magnocellular oxytocin system, the fount of maternity: adaptations in pregnancy. *Front Neuroendocrinol, 24*(1), 27-61. doi:10.1016/s0091-3022(02)00104-8

Sagar, S. M., Sharp, F. R., & Curran, T. (1988). Expression of c-fos protein in brain: metabolic mapping at the cellular level. *Science, 240*(4857), 1328-1331. doi:10.1126/science.3131879

Sarachana, T., Xu, M., Wu, R. C., & Hu, V. W. (2011). Sex hormones in autism: androgens and estrogens differentially and reciprocally regulate RORA, a novel candidate gene for autism. *PLoS One, 6*(2), e17116. doi:10.1371/journal.pone.0017116

Saul, M. C., Seward, C. H., Troy, J. M., Zhang, H., Sloofman, L. G., Lu, X., . . . Stubbs, L. (2017). Transcriptional regulatory dynamics drive coordinated metabolic and neural response to social challenge in mice. *Genome Res, 27*(6), 959-972. doi:10.1101/gr.214221.116

Savic, D., Distler, M. G., Sokoloff, G., Shanahan, N. A., Dulawa, S. C., Palmer, A. A., & Nobrega, M. A. (2011). Modulation ofTcf7l2 expression alters behavior in mice. *PLoS One, 6*(10), e26897. doi:10.1371/journal.pone.0026897

Sayad, A., Noroozi, R., Omrani, M. D., Taheri, M., & Ghafouri-Fard, S. (2017). Retinoic acid-related orphan receptor alpha (RORA) variants are associated with autism spectrum disorder. *Metab Brain Dis, 32*(5), 1595-1601. doi:10.1007/s11011-017-0049-6

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., . . . Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res, 13*(11), 2498-2504. doi:10.1101/gr.1239303

Singh-Taylor, A., Molet, J., Jiang, S., Korosi, A., Bolton, J. L., Noam, Y., . . . Baram, T. Z. (2018). NRSF-dependent epigenetic mechanisms contribute to programming of stress-sensitive neurons by neonatal experience, promoting resilience. *Mol Psychiatry, 23*(3), 648-657. doi:10.1038/mp.2016.240

Stolzenberg, D. S., & Champagne, F. A. (2016). Hormonal and non-hormonal bases of maternal behavior: The role of experience and epigenetic mechanisms. *Horm Behav, 77*, 204-210. doi:10.1016/j.yhbeh.2015.07.005

Stolzenberg, D. S., & Numan, M. (2011). Hypothalamic interaction with the mesolimbic DA system in the control of the maternal and sexual behaviors in rats. *Neurosci Biobehav Rev, 35*(3), 826-847. doi:10.1016/j.neubiorev.2010.10.003

Stolzenberg, D. S., & Rissman, E. F. (2011). Oestrogen-independent, experience-induced maternal behaviour in female mice. *J Neuroendocrinol, 23*(4), 345-354. doi:10.1111/j.1365-2826.2011.02112.x

Stolzenberg, D. S., Stevens, J. S., & Rissman, E. F. (2012). Experience-facilitated improvements in pup retrieval; evidence for an epigenetic effect. *Horm Behav, 62*(2), 128-135. doi:10.1016/j.yhbeh.2012.05.012

Stolzenberg, D. S., Stevens, J. S., & Rissman, E. F. (2014). Histone deacetylase inhibition induces long-lasting changes in maternal behavior and gene expression in female mice. *Endocrinology, 155*(9), 3674-3683. doi:10.1210/en.2013-1946

Vaccarino, F. J. (1994). Nucleus accumbens dopamine-CCK interactions in psychostimulant reward and related behaviors. *Neurosci Biobehav Rev, 18*(2), 207-214. doi:10.1016/0149-7634(94)90025-6

Wonch, K. E., de Medeiros, C. B., Barrett, J. A., Dudin, A., Cunningham, W. A., Hall, G. B., . . . Fleming, A. S. (2016). Postpartum depression and brain response to infants: Differential amygdala response and connectivity. *Soc Neurosci, 11*(6), 600-617. doi:10.1080/17470919.2015.1131193

Zhao, C., Eisinger, B. E., Driessen, T. M., & Gammie, S. C. (2014). Addiction and reward-related genes show altered expression in the postpartum nucleus accumbens. *Front Behav Neurosci, 8*, 388. doi:10.3389/fnbeh.2014.00388

Zhu, X., Cheng, M., Peng, M., Xiao, X., Yao, S., & Zhang, X. (2008). Basal behavioral characterization of hsf1 deficient mice and its cellular and behavioral abnormalities underlying chronic unpredictable stressors. *Behav Brain Res, 193*(2), 225-229. doi:10.1016/j.bbr.2008.05.024