Chromatin accessibility of primary human cancers ties regional mutational

processes with tissues of origin

Oliver Ocsenas[1,2], Jüri Reimand[1,2,3,@]

1. Computational Biology Program, Ontario Institute for Cancer Research, Toronto, ON, Canada
2. Department of Medical Biophysics, University of Toronto, Toronto, ON, Canada
3. Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada
[@] correspondence: Juri.Reimand@utoronto.ca

ABSTRACT

Background. Regional distribution of somatic mutations in cancer genomes associates with

DNA replication timing (RT) and chromatin accessibility (CA), however normal tissues and cell

lines have contributed these insights while associations with the epigenomes of primary cancers

remain uncharacterized.

Results. Here we model megabase-scale mutation burden in whole cancer genomes using ~900

CA and RT profiles of primary cancers, normal tissues, and cell lines. CA profiles of primary

cancers, rather than normal tissues, predict regional mutagenesis in most cancer types. Regional

mutation burden associates with the CA profiles of matching cancer types, indicating tissue-

specific determinants of mutagenesis. However, mutagenesis in squamous cell and lymphoid

cancers instead associates with RT profiles. Mutational signatures also show tissue-specific

associations with cancer epigenomes, especially for carcinogen-induced and unannotated

signatures. Lastly, while each cancer type includes certain frequently-mutated genomic regions

exceeding epigenome-informed predictions of mutation burden, these regions show a pan-cancer

convergence to biological processes involved in development and cancer. Thus, modelling

excess mutations using epigenomes highlights known cancer driver genes as well as frequently

mutated non-coding regions.

Conclusions. The dominant association of regional mutation burden with cancer epigenomes

suggests that many passenger mutations are determined by the epigenetic landscapes of

transformed cells and may occur later in tumor evolution. CA-informed models help find cancer

genes and pathways with positive selection and highlight regions where additional mutation

burden is contributed by local mutational processes. This study underlines the complex interplay

of mutational processes, genome function and evolution in cancer and tissues of origin.

INTRODUCTION

The cancer genome is a footprint of its evolution and molecular environment that is shaped by somatic mutations such as single nucleotide variants (SNVs) and structural alterations [1,2]. Cancer initiation and progression is caused by a small number of driver mutations that provide cells with selective advantages [3-5], however most mutations are functionally neutral passengers that are caused by various mutational processes [6-8]. Somatic mutations also occur in normal tissues and are frequently observed in known cancer genes [9,10]. Thus, we need to understand mutational processes to decipher tumor etiology and evolution and better characterize driver mutations.

Mutational processes act at different scales of the cancer genome [11,12]. Single base substitution (SBS) signatures affect certain trinucleotide context of DNA and are associated with aging, carcinogen exposures, defects in DNA repair pathways, and cancer therapies [6,13]. At a 100-nucleotide resolution, local mutational processes disproportionately affect certain non-coding genomic elements such as transcription start sites and binding sites of gene-regulatory proteins such as CTCF [14-16]. At the regional, megabase-scale resolution of the genome, mutation burden correlates with DNA replication timing (RT), chromatin accessibility (CA) and transcriptional activity [17-19]. Early-replicating, transcriptionally active regions of open chromatin have fewer mutations than late-replicating, passive regions of heterochromatin, potentially due to increased error rates and decreased mismatch repair later in DNA replication [20-23]. SBS signatures are also distributed asymmetrically with respect to DNA replication origins and timing [24]. Regional mutation burden is associated with epigenetic information of related normal cells, providing evidence of cells of cancer origin contributing to somatic variation [25] and allowing classification of cancers of unknown origin [26]. However, the precise molecular mechanisms driving these mutational processes remain incompletely understood. In particular, cell lines and normal tissues have been used to associate chromatin accessibility and mutation burden in cancer, while the epigenetic landscapes of primary human cancers remain unexplored.

Here we studied cancer epigenomes as determinants of regional mutagenesis in thousands of whole cancer genomes through a diverse collection of CA and RT profiles of cancers, normal tissues, and cell lines. CA profiles of matching cancer types, rather those than normal tissues, are the major determinants of regional mutagenesis and mutational signatures in most cancer types.

63 We found tissue-of-origin effects of CA and RT in most predictions, bespoke deviations in

64 specific cancer types and mutational signatures, and a pan-cancer convergence of excess

65 mutations to cancer driver genes and developmental pathways. Together, these results underline

66 the spatial complexity of regional mutagenesis in cancer genomes and highlight epigenome-

67 informed avenues to discover driver mutations.

Figure 1. Characterizing chromatin accessibility (CA) and replication timing (RT) as determinants of regional mutagenesis in cancer genomes. A. Somatic mutations in cancer genomes (top) and CA and RT datasets of normal tissues and cancers (bottom) were integrated to study regional mutational processes. Somatic single nucleotide variants (SNVs) of 2,517 whole cancer genomes were analyzed with 869 genome-wide profiles, including 773 CA profiles of primary human cancers, normal tissues, and cell lines from ATAC-seq experiments, and 96 RT profiles of and six cell cycle phases in 16 cell lines from RepliSeq experiments. **B.** Genomic regions of one megabase (Mbp) were analyzed. Regional mutation burden was estimated as the number of SNVs per megabase region. The mean values scores per region were derived for CA and RT profiles. **C.** Random forest models were trained using regional mutation burden profiles as the outcome and CA and RT profiles as the predictors. We analyzed the pan-cancer dataset and 17 datasets of specific cancer types with relevant CA and RT profiles available. **D.** To associate regional mutation burden with CA and RT, mutational signatures, and cancer driver genes, the random forest models were evaluated in terms of accuracy, predictor importance, and model residuals.

RESULTS

Chromatin accessibility of primary cancers is a major determinant of regional mutagenesis

To study cancer epigenetic profiles as determinants of regional mutagenesis in cancer genomes,

we collected 773 ATAC-seq profiles of genome-wide CA measurements in primary human

cancers, normal tissues and cell lines from ENCODE3, TCGA, and additional studies [27-34], as

well as 96 RepliSeq profiles of DNA replication timing measurements in 16 cell lines and six

cell cycle phases [35] (**Figure 1A; Supplementary Figure 1, Supplementary Table 1**). As

regional mutation burden, we studied 23 million SNVs in 2,465 highly-mappable genomic

regions of one megabase mapped across 2517 whole cancer genomes of 37 cancer types of the

ICGC/TCGA PCAWG project [1] (**Figure 1B**). The 869 CA and RT profiles were derived as mean

signal intensity values per megabase.

To map the complex non-linear associations of CA and RT profiles with regional mutagenesis,

random forest regression models were trained with mutation burden profiles as outcomes and CA

and RT profiles as predictors (*i.e.*, features) for the 17 cancer types for which both genomic and

relevant epigenomic profiles were available (**Figure 1C**). The most informative predictors were

quantified using statistical analysis and local feature prioritization of random forest models [36]

(**Figure 1D**). As expected, genome-wide profiles of regional mutation burden clustered

according to cancer types (**Supplementary Figure 2**).

We asked whether the CA profiles of cancers or those of normal cells and tissues showed

stronger associations with regional mutational processes in cancer genomes. We predicted

regional mutation burden in pairs of random forest models with matched data splits where the

predictors included either CA profiles of primary cancers or CA profiles of normal cells and

tissues, respectively. RT profiles were also included as predictors in both models to estimate the

relative contributions of CA.

In most cancer types, CA profiles of primary cancers showed stronger associations with regional

mutagenesis than CA profiles of normal cells and tissues (13 of 17, $P < 0.05$) (**Figure 2A**). The

most pronounced signal was observed in breast cancer where the regional mutagenesis

predictions informed by cancer CA profiles were nearly twice as accurate as those informed by

CA profiles of normal tissues (median adj.R^2 0.70 *vs.* 0.38; $P < 0.001$) (**Figure 2B**). Stronger

associations of cancer CA profiles and regional mutagenesis were also found in cancers of the

prostate, uterus, and kidney, and melanoma: the improvement in prediction accuracy was above

10% in those cancer types ($P < 0.001$). Stronger associations with cancer epigenomes were also

confirmed in the pan-cancer analysis across 37 cancer types, with a small but statistically

significant improvement in model accuracy (adj.R^2 0.90 *vs.* 0.88; $P < 0.001$). As the only

exception, regional mutation burden in liver cancer better associated with CA profiles of normal

tissues (adj.R^2 0.85 *vs.* 0.83; $P = 0.044$). The analysis provided inconclusive evidence for four

cancer types including lymphoid cancers (BNHL, CLL) and lung and thyroid adenocarcinomas.

We confirmed that the accuracy of regional mutation burden predictions in individual cancer

types was not significantly correlated with the overall mutation burden or the number of

sequenced genomes per cancer cohort (**Supplementary Figure 3**).

In summary, this analysis shows that in most cancer types, regional mutagenesis is more strongly

associated with chromatin accessibility of primary human cancers rather than normal tissues and

cell lines, even when accounting for DNA replication timing in the comparison. The diverse

collection of epigenomes included as predictors suggests that tissue-specific chromatin features

of individual cancer types, as well as pan-cancer chromatin features of proliferative cells may

contribute to regional mutagenesis.

Figure 2. Chromatin accessibility of primary cancers is a major determinant of regional mutagenesis in cancer genomes. A. Random forest models informed by CA profiles of primary cancers are more accurate predictors of regional mutation burden, compared to models informed by CA of normal tissues. Bar plot shows relative change in prediction accuracy (Δ adjusted R^2) of random model regression models informed by CA profiles of primary cancers, compared to matching models informed by CA of normal tissues. Replication timing (RT) profiles are included in all models as reference. Empirical P-values and 95% confidence intervals from bootstrap analysis are shown. Accuracy values of models informed by cancer CA profiles are listed below the bars (adjusted R^2). **B.** Examples of regional mutation burden predicted using models informed by CA profiles of cancer (top) *vs.* CA profiles of normal tissues (bottom). Scatterplots show model-predicted and observed mutation burden (X *vs.* Y-axis) in one-megabase regions. Prediction accuracy values are shown (bottom right).

Figure 3. Top predictors of regional mutagenesis tie cancer types and sites of origin. A. Quantitative genome-wide associations of regional mutation burden with the most informative CA and RT profiles in random forest models. 2D-density plots show the association CA and RT scores (Y-axis) and Shapley feature importance (SHAP) scores in each genomic window (X-axis) across all cancer types. CA profiles for cancer and normal samples, early RT profiles, and late RT profiles are plotted separately. CA and early RT profiles negatively

correlate with regional mutation burden while late RT profiles correlate positively. Spearman correlation values are shown (top right). **B.** CA profiles of primary cancers are the top predictors of regional mutagenesis in most cancer types. Bar plot shows the importance scores of the five most important predictors of random forest models for 17 cancer types (permutation $P < 0.001$). Error bars show ±1 standard deviation from bootstrap analysis. Brighter colors indicate the predictors where the epigenomic profile (CA or RT) matches the mutation profile of the related cancer type. **C-F.** Top predictors of regional mutation burden in individual cancer types. Shapley additive explanation (SHAP) scores show the impact of a given predictor on the predictions of regional mutation burden (Y-axis) relative to the values of the predictor (CA or RT; color gradient). In CA profiles, higher values (red) primarily associate with lower SHAP scores corresponding to increased mutation rates, while in contrast, higher values in late RT profiles associate with higher SHAP scores. Symbols indicate RT profiles (triangles) and CA profiles of normal tissues (circles).

Top predictors of regional mutagenesis match cancer types and sites of origin

To interpret the determinants of regional mutagenesis, we asked which specific CA and RT

profiles contributed the most to the predictive models when using all 869 cancer and normal

epigenomes as predictors. We selected the five most significant predictors for each cancer type

(permutation $P < 0.001$). These 85 CA and RT profiles were quantified using Shapley additive

explanation (SHAP) scores [36] that measured the directional associations of individual profiles

with the regional mutation burden in all genomic regions. As expected, regional mutation burden

negatively correlated with CA profiles of primary cancers and normal tissues (ρ_{cancer} = -0.74 vs

ρ_{normal} = -0.79; $P < 10^{-16}$) (**Figure 3A**). A dual relationship was apparent in RT profiles: RT

profiles of late cell cycle phases positively correlated with regional mutation burden while RT

profiles of early cell cycle phases correlated negatively (ρ_{late} = 0.75 *vs*. ρ_{early} = -0.84, $P < 10^{-16}$).

The inverse relationships of CA and RT with respect to regional mutation burden are consistent

with previous studies [17-23] and extend here to a diverse collection of epigenomes from primary

cancers and normal tissues.

We examined the most significant predictors of regional mutation burden for each cancer type

(**Figure 3B**). CA profiles of primary cancers dominated among the strongest predictors of

regional mutation burden in 12 of 17 cancer types. Most CA profiles represented the same or

related cancer type where the regional mutation burden was predicted, underlying tissue-specific

interactions of chromatin state and somatic mutagenesis Overall, CA profiles of primary cancers

were enriched among the top predictors (55 of 85 profiles observed *vs.* 41 expected, Fisher's

exact $P = 0.011$), confirming the stronger association with primary cancer epigenomes and

regional mutagenesis. Regional mutation burden measured in the breast, prostate, kidney,

stomach, and thyroid cancer cohorts of the PCAWG WGS dataset associated with the CA

profiles of the matching cancer samples in TCGA (BRCA, PRAD, KIRP, STAD, and THCA,

respectively). For example, in prostate cancer, four CA profiles of primary prostate cancers and

one breast cancer profile associated negatively with regional mutation burden (**Figure 3C**).

Additional associations were apparent at the level of organ systems. CA profiles of stomach and

colorectal cancers were the top predictors of regional mutation burden in biliary and esophageal

cancers (**Figure 3D**), suggesting epigenetic and mutational similarities of cancers of the

gastrointestinal tract. As another example, regional mutagenesis in in breast cancer genomes was

significantly associated with one CA profile of uterine cancer, and a similar association with

breast cancer CA was apparent in uterine cancer genomes, perhaps explained by common

mutational processes in cancers of the female reproductive system. Therefore, regional

mutational processes in individual cancer types have strong tissue-specific interactions with the

epigenomes of these cancer types.

Fewer CA profiles of normal tissues were found among top predictors of regional mutation

burden. The strongest association with normal tissue epigenomes was apparent in liver cancer, as

three CA profiles of normal liver were detected as the highest-ranking predictors of regional

mutation burden (**Figure 3E**). CA profiles of normal tissues were identified as predictors in

seven other cancer types, however their feature importance scores were lower compared to CA

profiles of related primary cancers. As expected, these CA profiles of normal tissues also

matched the cancer types where regional mutagenesis was measured. For example, in thyroid

cancer, one normal thyroid CA profile and four primary cancer CA profiles of the matching

cancer type (THCA) were the top predictors of regional mutation burden.

Replication timing showed the strongest associations in squamous cell cancers (SCC) and

lymphoid cancers, reflecting tissue-specific effects. Mutations in Lung-SCC and Head-SCC

cohorts of PCAWG associated with RT profiles of NHEK cells, a squamous cell line of human

epidermal keratinocytes (**Figure 3F**). Similarly, regional mutation burden in lymphoid cancers

(Lymph-BNHL, Lymph-CLL) strongly associated with RT profiles of B-cells. One potential

explanation of these normal cell lines associating with regional mutagenesis in those cancer

types, rather than CA profiles of primary cancers, is an earlier occurrence of mutagenesis in the

evolution of these cancer types. In the genomes of Lung-SCC and Head-SCC cohorts of

PCAWG, many mutations are associated with signatures of tobacco exposure, while somatic

hypermutation contributes to genome variation in normal B-cells and lymphomas [37].

Most RT predictors of regional mutagenesis we found in the analysis (12/15) represented late-

replicating cell cycle phases G2 and S4. Individual RT profiles positively associated with

regional mutation burden in late-replicating regions (*e.g.*, phase G2 of MCF-7 in breast cancer)

and negatively in early-replicating regions (*e.g.,* phase S1 of HNEK in Head-SCC), consistent

our analysis above (**Figure 3A**) and with earlier observations that elevated regional mutagenesis

is caused by increased DNA damage and decreased repair in late-replicating regions [20].

Fewer RT profiles occurred among top predictors compared to CA profiles in other cancer types.

RT profiles of matching cell lines (MCF-7, HepG2) were found among predictors of regional

mutation burden in breast and liver cancer, respectively, and the latter RT profile was also a

minor but significant predictor in uterine cancer. In general, fewer and less-diverse RT profiles

of cell lines were available for this analysis, and these offer only a limited representation of

mutational processes in different cancer types. In contrast, the larger set of CA profiles

represents more cancer types and provides complementary information to RT. This analysis

extends our findings of tissue-specific CA and RT profiles as the principal predictors of regional

mutagenesis and underlines the effects of cell-of-origin and tumor heterogeneity. Dominance of

cancer CA profiles among top predictors in most cancer types is consistent with our first

observations that CA profiles of primary cancers provide accurate predictions of regional

mutagenesis.

Figure 4. Associations of mutational signatures with chromatin accessibility and replication timing. A. Top predictors of megabase-scale mutation burden of single base substitution (SBS) signatures (top five predictors; *P* < 0.001, permutation test). Colors indicate the predictor type (CA, RT) and its relationship to the cancer type where mutagenesis is predicted (matching site/tissue or other). Brighter colors indicate the predictors where the epigenomic profile (CA or RT) matches the cancer type of the mutation profile. Asterisks indicate CA profiles of CD4-positive T-cells as predictors of SBS1 mutations. **B.** Prediction accuracy of megabase-scale burden of SBS signatures using CA and RT profiles. Signatures of carcinogens, unknown origin, and aging are more accurately predicted by CA and RT profiles than endogenous signatures. *P*-values are computed using F-tests with adjustment for genome-wide signature burden. Median accuracy values are printed. **C-F.** Top predictors of megabase-scale mutation burden in breast cancer quantified using SHAP scores. SHAP scores show the impact of a predictor (*i.e.*, CA or RT profile) on the predictions (Y-axis) and corresponding predictor values (color

gradient). **C.** Regional mutation burden in breast cancer genomes is predicted by CA profiles of primary breast cancers (BRCA) and uterine cancer (UCEC) as well as later replication timing (G2 phase) in a breast cancer cell line (MCF-7). CA profiles mostly negatively associate with mutagenesis while late RT profiles associate positively. **D.** Age-related mutations of SBS5 are predicted by BRCA CA profiles as well as RT profile of MCF-7, similarly to overall SNV burden. **E.** APOBEC-related mutations of SBS13 are also predicted by CA profiles (BRCA, as well as LUSC, TCGT); however, SHAP scores show that SBS13 mutations are positively correlated with CA. **F.** SBS1 mutations related to molecular clock activity are predicted by diverse CA profiles: two breast cancers as well as blood and immune cells (GM18520, CLL1_CD4, Peyer's patch). Symbols indicate RT profiles (triangles) and CA profiles of normal tissues (circles).

Associations of mutational signatures with chromatin accessibility and replication timing

We asked whether the associations of regional mutagenesis with CA and RT are further
explained by mutational signatures. We quantified the megabase-scale mutation burden
separately for the major single base substitution (SBS) signatures based on PCAWG datasets [6]
and predicted their regional distributions using random forest regression. We then selected the
top five CA and RT profiles that most significantly associated with the burden of each
mutational signature and cancer type ($P < 0.001$).

We compared the top CA and RT profiles that associated with total regional mutation burden and
the burden of individual mutational signatures (**Figure 4A**). Top predictors of individual
signatures were often consistent with predictors of bulk regional mutation burden and showed
tissue-specific associations of mutagenesis and chromatin accessibility. Matching CA profiles of
cancers were the top predictors of mutational signatures in breast, kidney, colorectal and stomach
cancers, while RT profiles of matching cell lines associated with mutations in SCCs and
lymphoid cancers. Top predictors of endogenous and exogeneous signatures were also mostly
consistent, indicating that various mutational processes are affected by the epigenetic landscapes
of cancers or their normal cells of origin.

SBS1 mutations showed the most variation in terms of CA profiles, compared to the profiles
predictive of bulk mutations and other SBS signatures. Interestingly, the CA profile of CD4-
positive T-cells from the peripheral blood of a CLL patient (CLL1_CD4) was consistently

detected as a predictor of SBS1 mutation burden in six solid cancer types (liver, kidney, uterus,

esophagus, stomach, head). This CA profile was only specific to SBS1 mutations and was not

associated with bulk mutation burden or any other SBS signatures in the cancer types we studied.

This T-cell CA profile may represent somatic mutations in non-cancerous cells of the immune

system or the tumor microenvironment. As another example, SBS1 mutations in liver cancer

associated with CA profiles of liver cancers, while overall regional mutation burden was

predominantly associated with CA profiles of normal liver tissues. The clock-like SBS1

signature of 5-methylcytosine deamination is associated with cancer patient age and stem cell

division rate, and this signature has been found in the somatic genomes of normal tissues and

adult stem cells [10,38,39]. Thus, SBS1 mutations may represent an earlier timepoint in tumor

evolution or contribution from normal cells that remain convoluted in bulk tissue sequencing.

We asked if our random forest models were equally informative of various mutational signatures.

Six classes of SBS signatures were compared in terms of prediction accuracy: APOBEC/AID,

DNA-repair, carcinogens, two age-related classes (SBS1 and SBS5/40), and signatures of

unknown cause, as predicted via all 869 CA and RT profiles in 17 cancer types. Three classes of

signatures showed stronger associations with CA and RT profiles (**Figure 4B**): random forest

predictions of carcinogenic signatures, signatures of unknown cause, and aging-associated

signatures SBS5 and SBS40 were significantly more accurate than the DNA repair,

APOBEC/AID, and SBS1 signatures combined, when accounting for number of mutations per

signature as covariate of prediction accuracy ($P \leq 10^{-4}$; F-test) (**Supplementary Figure 4**). Thus,

the mutational processes of carcinogen exposures, SBS5/40, and unknown signatures show

stronger interactions with CA and RT in cancer genomes.

We studied the interactions of SBS signatures with CA and RT profiles in breast cancer (**Figure**

4C-F). SBS5 mutations, representing most mutations in the cohort, associated with four CA

profiles of primary breast cancers (BRCA) and one late DNA-replicating profile of the breast

cancer cell line MCF-7, reflecting tissue-specific associations of mutations and chromatin

(**Figure 4D**). Regional mutation burden of all mutations was consistent with SBS5, and similarly

associated with CA and RT profiles of breast cancer (**Figure 4C**). However, an additional

predictive CA profile of uterine cancer was identified, perhaps due to common epigenetic

features of female hormone-driven cancers. Megabase-scale burden of all mutations and the

regional burden individual signatures negatively correlated with CA profiles in most cases, as

expected. In contrast, SBS2 and SBS13 signatures of AID/APOBEC mutagenesis correlated

positively with CA, such that higher SHAP values corresponded to increased chromatin

accessibility (**Figure 4E**). This agrees with prior observations that AID targets epigenetically

active elements and results in kataegis and clustered mutational signatures [6,40,41]. Lastly, SBS1

mutations associated with three CA profiles representing peripheral blood, lymphoid follicles,

and immune cells (**Figure 4F**), perhaps reflecting somatic mutagenesis in tumor-infiltrated

immune cells and other cells in the tumor microenvironment. In summary, individual mutational

signatures also predominantly associate with CA of primary cancers rather than normal tissues.

The complex interactions of CA and RT with regional mutagenesis in certain mutational

signatures may reflect inter-and intra-tumoral heterogeneity and help characterize the

mechanisms of mutational processes.

Figure 5. Excess mutation burden unexplained by CA and RT profiles converges to cancer genes and developmental pathways. A. Additional mutation burden exceeding predictions of CA and RT profiles, quantified via significantly higher model residuals of individual 100-kbps genomic windows. Scatter plot shows the genomic windows with elevated regional mutation burden (FDR < 0.05). Regions with known cancer genes are labeled if *FDR* < 10^{-10}. Colors indicate cancer types (see panel D). X-axis is capped at 10^{-50} and Y-axis is capped at 400. **B-C**. Examples of genomic regions with excess mutation burden and known or putative cancer driver mutations. The plots show the genes in the region (top), mutation burden (SNVs per kbps; observed and expected) (middle), and the top five most significant predictors (*i.e.*, CA and RT profiles; bottom). **B.** The genomic region encoding the driver gene *PIK3CA* in breast cancer. **C.** Super-enhancer region at the *IRF4-DUSP22* locus with elevated mutation burden in breast cancer. **D.** Pathway enrichment analysis of genomic regions with significantly higher regional mutation burden relative to CA and RT-informed predictions. The enrichment map shows significantly enriched biological processes and pathways (*FDR* < 0.05). The nodes represent enriched pathways and processes, the edges connect nodes sharing many genes, and the manually

Excess mutations unexplained by epigenomes converge to cancer driver genes and developmental pathways

To quantify the regional mutagenesis unexplained by CA and RT, we investigated the genomic
regions that were enriched in mutations relative to the mutation burden predicted by random
forest regression. To enable a more detailed, gene-level functional analysis of enriched
mutations, we repeated the predictions of regional mutation burden using a finer 100-kbps
genomic resolution. This revealed 1570 unique genomic regions in 17 cancer types that were
significantly enriched in mutations based on the CA- and RT-informed model residuals (*FDR* <
0.05) (**Figure 5A**). The mutation-enriched regions encoded 900 protein-coding genes including
67 known cancer genes [42], significantly more than expected by chance (33 expected, Fisher's
exact $P = 3.1$ x 10^{-8}). Most driver genes were only found in single cancer types and represented
key disease-specific drivers such as *EGFR* and *TERT* in glioma, *MYC* in BNHL and *APC* in
colorectal cancer (**Supplementary Figure 5**). For example, in breast cancer, the region encoding
PIK3CA was significantly enriched in mutations compared to the expected mutation burden
based on the CA and RT landscape (92 SNVs observed *vs.* 44 expected; FDR = 9.2 x 10^{-4})
(**Figure 5B**). PIK3CA is a major driver gene of breast cancer with hotspot mutations [43], thus
showing that genome-wide statistical models of CA and RT can capture known driver genes.

The models also revealed regions with frequent non-coding mutations. The most prominent
region was found in 12 of 17 cancer types due to unexpectedly frequent mutations, including
breast cancer (112 SNVs observed *vs.* 63 expected; FDR = 4.6 x 10^{-4}) (**Figure 5C;**
Supplementary Figure 6). The region encodes the oncogenic transcriptional regulator *IRF4*
(interferon regulatory factor 4) [44] and *DUSP22* encoding a signaling protein that was recently
described as a network-implicated driver gene due to non-coding mutations [45]. The region also
includes super-enhancers of immune cells [46]. The recurrence of mutations in this region in
multiple cancer types highlights it as a potential pan-cancer region of interest.

We then asked whether the frequently-mutated regions were associated with common biological

pathways and processes. An integrative pathway enrichment analysis that prioritized genomic

regions detected in multiple cancer types revealed 177 significantly enriched pathways and

processes (*FDR* < 0.05, ActivePathways [47]), of which 142 (80%) were detected in more than one

cancer type (**Figure 5D**). These findings converged into several functional themes of similar

pathways and processes. First, developmental processes related to brain and the central nervous

system, reproductive and sensory organs were associated with elevated mutation burden in

multiple cancer types. Second, a group of processes related to synapse organization, olfactory

and GPCR signaling were also identified in most cancer types. Third, cancer-related processes of

cell cycle, hormone response, and signal transduction were also identified, often through pan-

cancer data integration but not in any specific cancer type specifically. Lastly, a major group of

processes related to immune system activation were predominantly detected in BNHL,

potentially reflecting aberrant somatic hypermutation, as well as fewer associations with liver

and stomach cancers.

This analysis shows that although individual frequently-mutated genomic regions are mostly

characteristic of specific cancer types, enriched mutations converge to common pathways and

processes in multiple cancer types. Convergence of these excess mutations to developmental and

cancer-related processes is potentially explained by additional focal mutational processes

targeting epigenetically active regions of the genome that are not captured by our models at the

broader, sub-megabase resolution. Further, the enrichment of known cancer driver genes

suggests that positive selection of functional mutations may also contribute to this additional

mutation burden. This analysis exemplifies the complex interplay of cancer epigenomes, multi-

scale mutational processes and positive selection of cancer genes.

DISCUSSION

Our analysis highlights chromatin accessibility of primary human cancers as a major determinant of regional mutational processes in cancer genomes. Cancer epigenomes are predictive of regional mutation burden of matching cancer types, indicating tissue of origin associations in most cancer types we studied. While these associations are apparent for overall regional mutation burden in cancer genomes, they are also consistent with the regional variations in mutational signature burden. In contrast, the chromatin states of normal tissues and cell lines show only limited associations with regional mutagenesis of cancer genomes, extending the earlier studies that used the epigenetic profiles of cell lines and normal tissues to characterize mutational processes. The transformation of normal cells to cancer cells involves major changes in their epigenetic landscapes as gene-regulatory programs of cancer hallmark pathways are activated. Thus, one potential explanation to this stronger association of cancer epigenomes and regional mutagenesis is that mutational processes have a longer exposure on the somatic genomes shaped by the epigenomes of transformed cells, suggesting that many passenger mutations occur later in cancer evolution after the cells have acquired the epigenetic characteristics of cancer cells.

Replication timing information also associated with regional mutagenesis and confirmed strong effects with cell types related to cancer origin. However, CA profiles of primary human cancers evidently captured a larger fraction of variation of regional mutagenesis compared to RT profiles, apart from squamous cell and lymphoid cancers that strongly associated with relevant cell lines. Fewer RT profiles are used as predictors in our dataset and include mitotic cell lines that offer only limited representation of the diverse disease types in the pan-cancer cohort. As our models also include DNA replication timing profiles of several cell lines as reference, the stronger association with epigenomes of primary cancers shows that cancer epigenomes complement replication timing information with respect to regional mutagenesis. Interestingly, DNA replication has been shown to determine chromatin state [48]. Thus, the informative CA profiles of human cancers may represent a proxy of cancer-specific replication dynamics.

Mutational signature analysis revealed interactions of mutational processes with CA and tissues of origin. Carcinogen signatures, as well as signatures of unknown etiology, were overall better predicted by CA and RT, in contrast to signatures of aging and DNA damage where the genome-

wide predictions were less accurate. The stronger association of carcinogen signatures suggests

that the chromatin environment interacts with DNA damage or repair processes of carcinogen

exposure, for example through elevated mutational processes targeting active genes that are

otherwise protected from mutations through error-free mismatch repair [41]. Early replicating

regions in cells exposed to tobacco mutagens show elevated mutagenesis in transcribed strands

due to differential nucleotide excision repair activity [49]. Based on their stronger interactions with

RT and CA profiles, we speculate that some mutational signatures of currently unknown etiology

may relate to carcinogens. SBS17 mutations show some of the strongest interactions with CA

and RT in stomach and esophageal cancers in our analysis. This signature is currently of

unknown cause, however it has been linked to gastric acid reflux and reactive oxygen species [24].

Further integrative analysis of clinical and lifestyle information with patterns of regional

mutagenesis may shed light to these mutational processes.

We observed a functional convergence to developmental processes and cancer-related pathways

in the genomic regions where mutations were enriched beyond the predictions of our epigenomic

models. On the one hand, these data suggest that additional mutational processes affect distinct

regions with developmental genes and open-chromatin regions in individual cancer types,

however these regions converge to the same molecular pathways across cancer types. Such local

mutational processes are consistent with previous studies. For example, transcription start sites of

highly expressed genes and constitutively-bound binding sites of CTCF are subject to elevated

local mutagenesis in multiple cancer types [16]. Lineage-specific genes are enriched in indel

mutations in solid cancers [50]. Such local mutational processes are complementary to megabase-

scale processes where open chromatin is generally associated with a lower mutation frequency.

On the other hand, the enrichment of cancer genes and pathways in our data suggests that some

mutations unexplained by CA and RT are functional in cancer and their frequent occurrence at

specific genes, non-coding elements and molecular pathways is explained by positive selection [3-

5,45]. We can use this computational framework to find genomic regions with known and putative

driver mutations in coding and non-coding sites. Further study of these regions may deepen our

understanding of mutational processes and refine the catalogues of driver mutations.

This approach enables future studies to decipher the mechanisms and phenotypic associations of

mutational processes. Clinical, genetic, and epigenetic profiles of cancer patients can be

integrated to understand how regional mutational processes and the chromatin landscape are

modulated by clinical variables such as stage, grade or the therapies applied, genetic features

such as somatic driver mutations or inherited cancer risk variants, or lifestyle choices such as

tobacco or alcohol consumption. Complementary insights from sub-clonal reconstruction

analysis of cancer genomes [2,51], as well as single-cell sequencing of genomes and epigenomes

will allow mapping of regional mutagenesis at the level of distinct cell populations contributing

to temporal and spatial variation in mutational processes. As such multimodal datasets grow, we

can learn about early cancer evolution by comparing regional mutagenesis in the genomes of

cancers and normal cells. Understanding the molecular and genetic determinants of regional

mutagenesis and signatures in cancer genomes may help characterize carcinogen exposures and

genetic predisposition, ultimately enhancing early cancer detection and prevention in the future.

Methods

Somatic mutations in whole cancer genomes. We analyzed somatic single nucleotide variants

(SNVs; n = 43,778,859) derived from whole-genome sequencing (WGS) of 2,583 primary

cancer samples that were uniformly mapped to GRCh37/hg19 as part of the ICGC/TCGA Pan-

cancer Analysis of Whole Genomes (PCAWG) project [1]. Indels and variants in sex chromosomes

were excluded. To integrate mutations with epigenetic information, we mapped the SNVs to the

human genome version GRCh38 using the LiftOver function of the rtracklayer package in R (v

1.48) [52]. We removed 66 hypermutated tumors with more than 90,000 mutations (~30 mutations

/ Mbps), resulting in a dataset of 23,215,600 SNVs in 2,517 whole cancer genomes. We analyzed

the genomes of 17 cancer types with at least 25 samples in PCAWG as well as related epigenetic

profiles of normal and cancer tissues, as well as the pan-cancer dataset of all 37 cancer types.

Chromatin accessibility (CA) and replication timing (RT) profiles. Chromatin accessibility

data was derived from several ATAC-seq datasets, including the ENCODE3 project and six

additional studies to maximize the coverage of cancer types included in the PCAWG dataset, as

described below. CA profiles of 196 human cell and tissue types and 9 cancer cell lines at a

single basepair (bp) resolution in GRCh38 were derived from the ENCODE3 project [27]. CA

profiles of 115 normal human brain samples at a single bp resolution in GRCh37 were retrieved

from the study by Fullard *et al*. [29]. CA profiles of 21 normal immune cell types (B-cells, T-cells)

and 34 primary cancers (CLL) at a single base pair resolution in GRCh37 were retrieved from

the study by Rendeiro *et al*. [30]. Four CA profiles of HEK293 embryonic kidney cells at a 10-bp

resolution in GRCh37 were retrieved from the study by Karabacak Calviello *et al*. [31]. CA profiles

for two lymphoma cell lines at a single-bp resolution in GRCh37 were retrieved from the study

by Scharer *et al.* [32]. One CA profile of the normal human melanocytes (NHM1) cell line at a 10-

bp resolution in GRCh37 were retrieved from the study by Fontanals-Cirera *et al*. [33]. CA profiles

of four normal prostate tissues and six primary prostate cancers at a single-bp resolution in

GRCh37 were retrieved from the study by Pomerantz *et al*. [34]. CA profiles of several cancer

types were retrieved from the TCGA ATAC-seq dataset [28] of 410 primary cancer samples,

representing cancers of 404 unique patient donors and 796 genome-wide profiles in total. We

used 381 CA profiles of the TCGA dataset such that technical and biological replicates of

distinct cancer samples were pooled by per-region averaging CA signal. Prior to this averaging,

22 CA profiles for which only one replicate was available were removed, and one CA profile of

a low-grade glioma (LGG) that was an outlier in our initial analyses as also removed. In total,

773 CA profiles were included in the analysis, including 421 cancer profiles, 341 profiles of

normal tissues and cell lines, and 11 profiles of cancer cell lines. Besides these CA profiles, 96

replication timing profiles of 16 cell lines, each with six cell cycle phases, were derived from the

RepliSeq study by Hansen *et al.* [35]. CA and RT profiles were constructed from the BigWig files

of the original studies using mean values of signal intensity per each genomic window. Genomic

coordinates of the GRCh37 reference genome were mapped to the GRCh38 reference genome

using LiftOver. In total, the set of 869 (773 + 96) CA and RT profiles was used.

Integrating regional mutagenesis with CA and RT profiles. We evaluated chromatin

accessibility, replication timing and mutation burden in non-overlapping genomic regions of one

megabase (Mbp; one million base pairs). We excluded a subset of genomic regions with low

mappability ($\leq$80% in the UMAP software [53]) as well as sex chromosomes, resulting in 2,465

regions included in the study. For megabase-scale CA and RT profiles, each genomic region was

assigned the mean value of its epigenetic signal. For megabase-scale somatic mutation burden,

each region was assigned the total mutation count separately for the pan-cancer dataset and each

of the 17 cancer types. In two cohorts (chronic lymphocytic leukemia; B-cell non-Hodgkin

lymphoma), we removed two regions encoding immunoglobulin genes (chr2:89Mbps,

chr22:23Mbps) with known high somatic variation in immune cells, as observed in our initial

analyses.

Random forest regression. Megabase-scale profiles of mutation burden and CA and RT profiles

were analyzed with random forest regression [54] with CA and RT profiles as the predictors (*i.e.*,

features) and mutation burden as the target (*i.e.*, response). Number of trees (1000) and fraction

of predictors at each split (1/3) were used as hyperparameters. Monte-Carlo cross-validation over

1000 data splits considered subsets of genomic regions for model training (80%) and validation

(20%). We used the adjusted R^2 (adj.R^2) metric to evaluate model performance that measures the

variance explained by the model adjusted for model complexity (*i.e.*, the number of CA and RT

profiles used for predictions).

Comparing CA profiles of primary cancers and normal tissues as predictors of regional mutagenesis. First, we compared the overall accuracy of predicting megabase-scale mutation burden using CA profiles of cancers *vs.* normal tissues. Two sets of random forest regression models were run in a joint Monte-Carlo cross-validation procedure that used all CA profiles of normal tissues (M_n) and cancers (M_c) as predictors, respectively. Both models also included the same set of RT profiles as predictors as reference. At each iteration, models were trained on matching subsets of genomic regions (80%) and tested on the remaining genomic regions (20%), and model accuracy (adj.R^2) values as well as the relative change values (Δadj.R^2 = adj.R^2(M_c)-adj.R^2(M_n)) were derived in the corresponding test sets, allowing us to directly compare the two models. For each cancer type, median Δadj.R^2 values and 95% confidence intervals were reported. Empirical P-values were computed as the fraction of cross-validation iterations where the Δadj.R^2 crossed on the opposite side of zero relative to the median value. We also trained the models M_n and M_c on the full set of genomic regions and compared the accuracy of the two sets of models. Observed and model-predicted mutation burden values per region were visualized as scatter plots with local regression (loess) trendlines (span=0.9). Spearman correlation tests were used to evaluate the associations of model accuracy, WGS cohort size and per-megabase mutation burden in different cancer types.

Evaluating CA and RT profiles as predictors of regional mutation burden. We used the incMSE (increase in model mean-squared-error) metric to evaluate the most important features (*i.e.*, CA, RT profiles) in random forest models. incMSE measures the relative change in model prediction accuracy upon permutations of the values of a given feature. We derived incMSE values of CA and RT profiles for the 17 cancer types for which matching CA and/or RT profiles were available. Two additional statistical methods were used to evaluate the significance of incMSE of CA and RT profiles. First, permutation tests were used to detect CA and RT profiles where incMSE values significantly exceeded those of permuted data. We fitted random forest regression models for every cancer type 1,000 times using randomly reassigned megabase-scale mutation burden estimates as null distributions for incMSE values for CA and RT profiles. Specific profiles were considered statistically significant if their observed incMSE values exceeded all 1000 incMSE values from permuted datasets (*i.e.*, empirical $P < 0.001$). Second, we used bootstrapping of random forest regression where the genomic regions with predictor and

response values were sampled randomly with replacement. We repeated this resampling process
1000 times and recorded the incMSE values for all CA and RT profiles to evaluate the
confidence intervals of the derived incMSEs.

Feature importance of CA and RT profiles in predictions of regional mutagenesis. We used
the SHapley Additive exPlanation (SHAP) method [36,55] to interpret the interactions of regional
mutation burden and CA and RT profiles. Here, SHAP scores reflect the importance of each
feature in the random forest model (*i.e.*, CA or RT profile) in predicting a specific observation
(*i.e.*, mutation burden of a certain genomic region), and represent its relative contribution to the
prediction (*i.e.*, effect size) as well as the direction of the prediction (*i.e.*, positive or negative).
SHAP values were computed using models trained all genomic regions and separately for cancer
types, using the python packages shap (0.35.0) [36,55] and scikit-learn (0.23.1) [56] via the R package
reticulate (1.16) [57].

Associating mutational signatures with CA and RT. For mutational signature analysis, we
used single base substitution (SBS) annotations of SNVs derived in the PCAWG project [6]. For
each genomic region, we computed the mutational signature burden probabilistically by adding
the SBS-specific probabilities of all individual SNVs in the region, thus accounting for all
signature exposures rather than top-ranking signatures for each SNV. We filtered lower-
frequency SBS signatures in each cancer type (*i.e.*, <20,000 or <5% of all SNVs). To evaluate
CA and RT profiles as predictors of megabase-scale mutational signature burden, we trained
random forest models where a probabilistic SBS profile was used as model response. We
evaluated model performance, selected top features, and computed SHAP scores similarly to
bulk mutation analysis described above. We grouped the mutational signatures based on their
etiology according to the COSMIC database (version 3.2, downloaded in March 2021):
AID/APOBEC, deficient DNA repair, exogeneous/carcinogen, unknown/other, SBS5/40 and
SBS1. We compared model accuracy values for predicting regional mutational signature burden
of the six classes of signatures using ANOVA analysis and F-tests. We used the covariate of the
average megabase-scale SBS burden to account for a potential of improved predictions in cancer
types with higher overall mutation burden.

Prioritizing highly-mutated genomic regions exceeding CA and RT predictions. To study regional mutation burden unexplained by CA and RT profiles, we prioritized the genomic regions where the random forest predictions significantly underestimated the observed mutation burden. Random forest regression was repeated on 100-kbps regions to improve gene-level interpretation. To score genomic regions, we subtracted the predicted mutation counts from the observed counts to derive residual values. Residuals were Z-transformed and the resulting one-tailed *P*-values were adjusted for multiple testing using Benjamini-Hochberg FDR.

Pathway enrichment analysis of regional mutation variation. To understand the functional importance of excess mutations unexplained by CA and RT profiles, we performed an integrative pathway enrichment analysis across the relevant cancer types using the ActivePathways method [47] (*FDR* < 0.05). Gene sets of biological processes of Gene Ontology and molecular pathways of Reactome were collected from the GMT files provided in the g:Profiler web server [58] (downloaded Feb 23rd, 2021) and were filtered using default settings of ActivePathways. In each cancer type, all protein-coding genes were assigned the P-values reflecting excess mutation burden unexplained by CA and RT in respective regions. The data fusion in ActivePathways prioritized the genes that were frequently mutated in multiple cancer types. Enriched pathways were visualized as an enrichment map [59] and themes were curated manually. We also visualized the genomic regions with excess mutations as a scatter plot of residual values and -log10-transformed FDR values that were capped at 400, and 10^{-50}, respectively. We highlighted known cancer driver genes of the Cancer Gene Census database [42] (downloaded Mar 26th 2021) and computed their enrichment in the list of pathway-associated genes using a Fisher's exact test.

Code availability. Source code for this study is available at
https://github.com/reimandlab/CA2M_v2.

Acknowledgments. We thank Christian A. Lee, Kevin Cheng, Phedias Diamandis, and Anne

Martel for constructive comments on this study. This work was supported by the Canadian

Institutes of Health Research (CIHR) Project Grant to J.R., A New Investigator Award of the

Terry Fox Research Institute (TFRI) to J.R., and the Investigator Award to J.R. from the Ontario

Institute for Cancer Research (OICR). Funding to OICR is provided by the Government of

Ontario. The results shown here are in whole or part based upon data generated by the TCGA

Research Network: https://www.cancer.gov/tcga. We acknowledge the contributions of the many

clinical networks of ICGC and TCGA who provided samples and data to PCAWG. We thank the

patients and their families for their participation in ICGC and TCGA projects.

Author contributions. O.O. analyzed the data and prepared the figures. J.R. and O.O.

interpreted the data and wrote the manuscript. J.R. conceived and supervised the project. The

authors reviewed and edited the manuscript and approved the final version.

References

ICGC-TCGA Pan-Cancer Analysis of Whole Genomes Consortium. Pan-cancer analysis of whole genomes. *Nature* **578**, 82-93, doi:10.1038/s41586-020-1969-6 (2020).

Gerstung, M. *et al.* The evolutionary history of 2,658 cancers. *Nature* **578**, 122-128, doi:10.1038/s41586-019-1907-7 (2020).

Martincorena, I. *et al.* Universal Patterns of Selection in Cancer and Somatic Tissues. *Cell* **171**, 1029-1041 e1021, doi:10.1016/j.cell.2017.09.042 (2017).

Rheinbay, E. *et al.* Analyses of non-coding somatic drivers in 2,693 cancer whole genomes. *Nature* **578**, 102–111 (2020).

Zhu, H. *et al.* Candidate Cancer Driver Mutations in Distal Regulatory Elements and Long-Range Chromatin Interaction Networks. *Mol Cell*, doi:10.1016/j.molcel.2019.12.027 (2020).

Alexandrov, L. B. *et al.* The repertoire of mutational signatures in human cancer. *Nature* **578**, 94-101, doi:10.1038/s41586-020-1943-3 (2020).

Li, Y. *et al.* Patterns of somatic structural variation in human cancer genomes. *Nature* **578**, 112-121, doi:10.1038/s41586-019-1913-9 (2020).

Kumar, S. *et al.* Passenger Mutations in More Than 2,500 Cancer Genomes: Overall Molecular Functional Impact and Consequences. *Cell* **180**, 915-927 e916, doi:10.1016/j.cell.2020.01.032 (2020).

Martincorena, I. *et al.* Tumor evolution. High burden and pervasive positive selection of somatic mutations in normal human skin. *Science* **348**, 880-886, doi:10.1126/science.aaa6806 (2015).

Blokzijl, F. *et al.* Tissue-specific mutation accumulation in human adult stem cells during life. *Nature* **538**, 260-264, doi:10.1038/nature19768 (2016).

Supek, F. & Lehner, B. Scales and mechanisms of somatic mutation rate variation across the human genome. *DNA Repair (Amst)*, 102647, doi:10.1016/j.dnarep.2019.102647 (2019).

Gonzalez-Perez, A., Sabarinathan, R. & Lopez-Bigas, N. Local Determinants of the Mutational Landscape of the Human Genome. *Cell* **177**, 101-114, doi:10.1016/j.cell.2019.02.051 (2019).

Pich, O. *et al.* The mutational footprints of cancer therapies. *Nature genetics* **51**, 1732-1740, doi:10.1038/s41588-019-0525-5 (2019).

Katainen, R. *et al.* CTCF/cohesin-binding sites are frequently mutated in cancer. *Nature genetics* **47**, 818-821, doi:10.1038/ng.3335 (2015).

Sabarinathan, R., Mularoni, L., Deu-Pons, J., Gonzalez-Perez, A. & Lopez-Bigas, N. Nucleotide excision repair is impaired by binding of transcription factors to DNA. *Nature* **532**, 264-267, doi:10.1038/nature17661 (2016).

Lee, C. A., Abd-Rabbo, D. & Reimand, J. Functional and genetic determinants of mutation rate variability in regulatory elements of cancer genomes. *Genome Biol* **22**, 133, doi:10.1186/s13059-021-02318-x (2021).

Lawrence, M. S. *et al.* Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* **499**, 214-218, doi:10.1038/nature12213 (2013).

Schuster-Bockler, B. & Lehner, B. Chromatin organization is a major influence on regional mutation rates in human cancer cells. *Nature* **488**, 504-507, doi:10.1038/nature11273 (2012).

568	19	Stamatoyannopoulos, J. A. *et al.* Human mutation rate associated with DNA replication
569		timing. *Nature genetics* **41**, 393-395, doi:10.1038/ng.363 (2009).
570	20	Supek, F. & Lehner, B. Differential DNA mismatch repair underlies mutation rate
571		variation across the human genome. *Nature* **521**, 81-84, doi:10.1038/nature14173 (2015).
572	21	Zheng, C. L. *et al.* Transcription restores DNA repair to heterochromatin, determining
573		regional mutation rates in cancer genomes. *Cell Rep* **9**, 1228-1234,
574		doi:10.1016/j.celrep.2014.10.031 (2014).
575	22	Woo, Y. H. & Li, W. H. DNA replication timing and selection shape the landscape of
576		nucleotide variation in cancer genomes. *Nat Commun* **3**, 1004, doi:10.1038/ncomms1982
577		(2012).
578	23	Liu, L., De, S. & Michor, F. DNA replication timing and higher-order nuclear
579		organization determine single-nucleotide substitution patterns in cancer genomes. *Nat*
580		*Commun* **4**, 1502, doi:10.1038/ncomms2502 (2013).
581	24	Tomkova, M., Tomek, J., Kriaucionis, S. & Schuster-Bockler, B. Mutational signature
582		distribution varies with DNA replication timing and strand asymmetry. *Genome Biol* **19**,
583		129, doi:10.1186/s13059-018-1509-y (2018).
584	25	Jiao, W. *et al.* A deep learning system accurately classifies primary and metastatic
585		cancers using passenger mutation patterns. *Nat Commun* **11**, 728, doi:10.1038/s41467-
586		019-13825-8 (2020).
587	26	Polak, P. *et al.* Cell-of-origin chromatin organization shapes the mutational landscape of
588		cancer. *Nature* **518**, 360-364, doi:10.1038/nature14221 (2015).
589	27	Consortium, E. P. *et al.* Expanded encyclopaedias of DNA elements in the human and
590		mouse genomes. *Nature* **583**, 699-710, doi:10.1038/s41586-020-2493-4 (2020).
591	28	Corces, M. R. *et al.* The chromatin accessibility landscape of primary human cancers.
592		*Science* **362**, doi:10.1126/science.aav1898 (2018).
593	29	Fullard, J. F. *et al.* An atlas of chromatin accessibility in the adult human brain. *Genome*
594		*research* **28**, 1243-1252, doi:10.1101/gr.232488.117 (2018).
595	30	Rendeiro, A. F. *et al.* Chromatin mapping and single-cell immune profiling define the
596		temporal dynamics of ibrutinib response in CLL. *Nat Commun* **11**, 577,
597		doi:10.1038/s41467-019-14081-6 (2020).
598	31	Karabacak Calviello, A., Hirsekorn, A., Wurmus, R., Yusuf, D. & Ohler, U.
599		Reproducible inference of transcription factor footprints in ATAC-seq and DNase-seq
600		datasets using protocol-specific bias modeling. *Genome Biol* **20**, 42, doi:10.1186/s13059-
601		019-1654-y (2019).
602	32	Scharer, C. D. *et al.* Genome-wide CIITA-binding profile identifies sequence preferences
603		that dictate function versus recruitment. *Nucleic Acids Res* **43**, 3128-3142,
604		doi:10.1093/nar/gkv182 (2015).
605	33	Fontanals-Cirera, B. *et al.* Harnessing BET Inhibitor Sensitivity Reveals AMIGO2 as a
606		Melanoma Survival Gene. *Mol Cell* **68**, 731-744 e739, doi:10.1016/j.molcel.2017.11.004
607		(2017).
608	34	Pomerantz, M. M. *et al.* Prostate cancer reactivates developmental epigenomic programs
609		during metastatic progression. *Nature genetics* **52**, 790-799, doi:10.1038/s41588-020-
610		0664-8 (2020).
611	35	Hansen, R. S. *et al.* Sequencing newly replicated DNA reveals widespread plasticity in
612		human replication timing. *Proc Natl Acad Sci U S A* **107**, 139-144,
613		doi:10.1073/pnas.0912402107 (2010).

36   Lundberg, S. M. *et al.* From Local Explanations to Global Understanding with
Explainable AI for Trees. *Nat Mach Intell* **2**, 56-67, doi:10.1038/s42256-019-0138-9
(2020).
37   Odegard, V. H. & Schatz, D. G. Targeting of somatic hypermutation. *Nat Rev Immunol* **6**,
573-583, doi:10.1038/nri1896 (2006).
38   Alexandrov, L. B. *et al.* Clock-like mutational processes in human somatic cells. *Nature*
genetics **47**, 1402-1407, doi:10.1038/ng.3441 (2015).
39   Abascal, F. *et al.* Somatic mutation landscapes at single-molecule resolution. *Nature*,
doi:10.1038/s41586-021-03477-4 (2021).
40   Wang, Q. *et al.* Epigenetic targeting of activation-induced cytidine deaminase. *Proc Natl*
Acad Sci U S A **111**, 18667-18672, doi:10.1073/pnas.1420575111 (2014).
41   Supek, F. & Lehner, B. Clustered Mutation Signatures Reveal that Error-Prone DNA
Repair Targets Mutations to Active Genes. *Cell* **170**, 534-547 e523,
doi:10.1016/j.cell.2017.07.003 (2017).
42   Futreal, P. A. *et al.* A census of human cancer genes. *Nat Rev Cancer* **4**, 177-183,
doi:10.1038/nrc1299 (2004).
43   Nik-Zainal, S. *et al.* Landscape of somatic mutations in 560 breast cancer whole-genome
sequences. *Nature* **534**, 47-54, doi:10.1038/nature17676 (2016).
44   Iida, S. *et al.* Deregulation of MUM1/IRF4 by chromosomal translocation in multiple
myeloma. *Nature genetics* **17**, 226-230, doi:10.1038/ng1097-226 (1997).
45   Reyna, M. A. *et al.* Pathway and network analysis of more than 2,500 whole cancer
genomes. *Nature Communications* **11**, 729 (2020).
46   Hnisz, D. *et al.* Super-enhancers in the control of cell identity and disease. *Cell* **155**, 934-
947, doi:10.1016/j.cell.2013.09.053 (2013).
47   Paczkowska, M. *et al.* Integrative pathway enrichment analysis of multivariate omics
data. *Nature Communications* **11**, 735 (2020).
48   Klein, K. N. *et al.* Replication timing maintains the global epigenetic state in human
cells. *Science* **372**, 371-378, doi:10.1126/science.aba5545 (2021).
49   Kucab, J. E. *et al.* A Compendium of Mutational Signatures of Environmental Agents.
Cell **177**, 821-836 e816, doi:10.1016/j.cell.2019.03.001 (2019).
50   Imielinski, M., Guo, G. & Meyerson, M. Insertions and Deletions Target Lineage-
Defining Genes in Human Cancers. *Cell* **168**, 460-472 e414,
doi:10.1016/j.cell.2016.12.025 (2017).
51   Dentro, S. C. *et al.* Characterizing genetic intra-tumor heterogeneity across 2,658 human
cancer genomes. *Cell*, doi:10.1016/j.cell.2021.03.009 (2021).
52   Lawrence, M., Gentleman, R. & Carey, V. rtracklayer: an R package for interfacing with
genome browsers. *Bioinformatics* **25**, 1841-1842, doi:10.1093/bioinformatics/btp328
(2009).
53   Karimzadeh, M., Ernst, C., Kundaje, A. & Hoffman, M. M. Umap and Bismap:
quantifying genome and methylome mappability. *Nucleic Acids Res* **46**, e120,
doi:10.1093/nar/gky677 (2018).
54   Ho, T. K. Random decision forests. *IEEE Proceedings of 3rd international conference on*
document analysis and recognition **1**, 278-282, doi:doi:10.1109/ICDAR.1995.598994
(1995).
55   Lundberg, S. M. & Lee, S. A Unified Approach to Interpreting Model Predictions.
Advances in Neural Information Processing Systems (NIPS) **30** (2017).

56   Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine*
Learning Research **12**, 2825-2830 (2011).
57   Ushey, K., Allaire, J. J. & Tang, Y. Reticulate R Package. *GitHub*
https://rstudio.github.io/reticulate/ doi:https://rstudio.github.io/reticulate/ (2021).
58   Reimand, J., Kull, M., Peterson, H., Hansen, J. & Vilo, J. g:Profiler--a web-based toolset
for functional profiling of gene lists from large-scale experiments. *Nucleic acids research*
35, W193-200, doi:10.1093/nar/gkm226 (2007).
59   Reimand, J. *et al.* Pathway enrichment analysis and visualization of omics data using
g:Profiler, GSEA, Cytoscape and EnrichmentMap. *Nat Protoc* **14**, 482-517,
doi:10.1038/s41596-018-0103-9 (2019).