Early Detection of Tau Pathology

Parag Parekh[1], Andrew Badachhape[1], Qingshan Mu[1], Rohan Bhavane[2], Mayank Srivastava[2], Igor Stupin[2], Prajwal Bhandari[1], Laxman Devkota[1], Jason Eriksen[3], Eric Tanifum[4], Ketan Ghaghada[4], and Ananth Annapragada[4,*]

[1]Baylor College of Medicine, Houston, TX, USA, [2]Texas Children's Hospital, Houston, TX, USA, [3] College of Pharmacy, University of Houston, Houston, TX, USA, [4]Texas Children's Hospital/Baylor College of Medicine, Houston, TX, USA

Email: annaprag@bcm.edu/avannapr@texaschildrens.org

Abstract

While a definitive Alzheimer's disease (AD) diagnosis remains a post-mortem exercise, the ATN Research Framework proposed by the National Institute on Aging and the Alzheimer's Association utilizes a score representing the presence of amyloid deposits (A), tau deposits (T) and neuronal degeneration markers (N), with A+T+ necessary for a positive diagnosis. Current detection of tau pathology lags amyloid detection by years and by the time both markers are detected the disease is fairly advanced. We describe the development of a new generation of molecular imaging probes for *in vivo* detection of cells undergoing abnormal phosphorylation representing the initial stages of pTau pathology, potentially enabling a very early stage diagnosis of AD. We describe a novel nanoparticle formulation that binds such abnormally phosphorylating cells in a mouse model of tau pathology, enabling in *vivo* visualization of the hyperphosphorylative state by magnetic resonance imaging. Our results demonstrate the potential of this novel platform to identify a correlative marker signifying the development of future tau pathology, and has implications for early-stage diagnosis of Alzheimer's disease.

Introduction

The microtubule associated protein tau, coded for by the *MAPT* gene, is abundant in the brain and is present in neurons, glia and other cell types. Tau shows immense diversity. It is expressed in six isoforms and has a vast array of post-translational modifications, including glycosylation, glycation, nitration, ubiquitination and more than 80 phosphorylation sites expanding the complexity of its role in health and disease[1,2]. A definitive feature of many neurodegenerative diseases including AD, frontotemporal lobar degeneration (FTLD), Parkinson's disease (PD) is the presence of intracellular aggregated filamentous tau (collectively termed

"tauopathies")[3–6]. This pathogenic insoluble tau been implicated in disease progression in these dysfunctional

brains. The transition from physiological soluble tau is primarily associated with changes in its phosphorylation

state leading to oligomeric tau[7], then tau fibrils known as paired helical fragments (PHF) that collectively form

characteristic neurofibrillary tangles (NFT). The tau aggregates are then capable of "infecting" a healthy cell

inducing further misfolding, aggregation, and neuro-toxicity[8,9]. Studies of intercellular propagation demonstrate

passage through an extracellular phase that progresses throughout the brain [10,11].

The research framework proposed for precise diagnosis of AD by the National Institute of Aging-

Alzheimer's Association (NIA-AA) categorizes extracellular deposits of amyloid beta (A), presence of

intraneuronal hyperphosphorylated tau (T) and markers of neurodegeneration or neuronal injury (N). Each

biomarker is scored either positive or negative[12]. To be on the AD continuum, A+ (Amyloid positive) is required,

while a positive diagnosis of AD requires A+ and T+. Biomarker detection can be by i) positron emission

tomography (PET) imaging of amyloid and tau or, ii) cerebrospinal fluid (CSF) detection of reduced $A\beta_{42}$, and/or

high $A\beta_{40}$/ $A\beta_{42}$ and high phosphorylated tau and total tau, or iii) neuronal injury or degeneration as shown by

structural brain magnetic resonance imaging (MRI)[13]. Tracking of brain pathology in longitudinal studies suggests

that tau pathology may precede the accumulation of $A\beta$, but is undetectable as it is below current biomarker

detection threshold levels, and is amplified catastrophically by independent $A\beta$ deposition [14–16]. ATN research

framework based diagnosis of AD is therefore limited by tau pathology detection, providing an impetus to develop

specific and sensitive tau detection methods for earlier diagnosis.

Other factors to consider in the development of tau detection methods include the invasive nature of

CSF sampling requiring lumbar puncture, and in the case of PET imaging, exposure to ionizing radiation, high

cost, well documented side effects, irregular availability in primary care setting, and uneven geographical

availability of PET scanners and isotopes[17–21]. The short half-life of PET agents also poses challenges for the

detection of intracellular tau in the early stages of tau pathology formation[22]. Blood based markers are very

promising, but only provide an indirect measure that cannot provide information on the localization of tau

pathology in brain[23–26]. Methods to detect early tau pathology that avoid these pitfalls are therefore highly

desirable.

The initiation of tau pathology is marked by abnormal phosphorylation of tau[27,28]. We hypothesize that

hyperphosphorylative conditions in neurons, consistent with an altered balance of kinase-phosphatase activity

resulting in elevated levels of hyperphosphorylated tau species, result in unique surface markers[29–31]. We

envisioned a targeted contrast-enhanced MRI test identifying the earliest stages of tau pathology represented by

cells in the process of abnormally phosphorylating tau. We used an iterative Cell-SELEX process to identify DNA

thioaptamers that specifically bound such cells[32,33]. The use of phosphorothioate modified aptamers enhances

stability under physiological conditions. High T1 relaxivity PEGylated liposomes bearing macrocyclic Gd-

chelates[34] were modified to bear the aptamers on their surface, thus enabling targeting of the particles to the

surface of hyperphosphorylative cells for contrast-enhanced MRI. At the molecular level, we sought the binding

targets of the aptamers, identifying vimentin, a normally intracellular protein that interestingly is specifically

expressed on the surface of cells under hyperphosphorylative conditions, and representing a possible biomarker

of pathological hyperphosphorylation found in AD.

Results

Validation of cell surface changes in hyperphosphorylative conditions

SH-SY5Y, a human neuroblastoma cell line that can be differentiated into neuron-like cells by changes

in culture medium was used to model cell-surface changes under hyperphosphorylative conditions (Fig 1B). We

used retinoic acid (RA) to induce cell differentiation marked by temporal changes in morphology including the

formation and lengthening of neurites and with a strong increase in levels of intracellular tau. Imbalance in the

kinase and phosphatase activity leading to hyperphosphorylation, simulating early stages of tauopathies, was

induced by the use of a cell permeable neurotoxin okadaic acid (30nM, 24 h) (OA)[35], and confirmed by the

increase in phosphorylated tau S396 (Supp. Fig. 1). In parallel experiments, a milder agent, excitotoxin quinolinic

acid (QA) 1μM for 24 h was also used to induce hyperphosphosphorylation[36].

A reverse phase protein array (RPPA) assay was conducted to test the hypothesis that

hyperphosphorylation results in compositional changes reflected on the cell-surface. RPPA analysis

demonstrated marked cell surface changes in hyperphosphorylative cells including over-expression of cell

surface receptors. RPPA assessment with a panel of 221 proteins across different pathways and post-

translational modifications including phosphorylation examined SH SY5Y cell lysates under differentiated and

hyperphosphorylative states. We identified 98 proteins and 36 phosphorylated proteins that showed significant

change in their expression under different conditions (**Fig 1C**). Uniprot [37] protein associations showed 44 cell-

membrane associated proteins, 10 peripheral membrane or 12 single-pass membrane proteins were significantly

altered under hyperphosphorylative conditions.

Screening for aptamers that bind cells in hyperphosphorylative state

Aptamer screening was performed using the cell-SELEX approach on differentiated SH-SY5Y cells in a hyperphosphorylative state (Fig. 2A). A total of 26 cell SELEX cycles were performed. To remove oligonucleotides that bound common cell surface molecules not specific to the hyperphosphorylative state, a negative selection was introduced at cycles 12 and 13 using differentiated, non hyperphosphorylative cells (i.e. without OA treatment). Anticipating that selected aptamers would be systemically delivered as nanoparticle imaging agents, and the primary toxicity driven by anomalous hepatocyte uptake, we conducted another round of negative selection at cycles 20 and 21 using a hepatocyte cell-line THLE-3 to remove oligonucleotides that exhibited enhanced uptake by hepatocytes.

Tau1 and Tau3 aptamers specifically bind hyperphosphorylative cells

Sequencing of all the selected pools was performed using the Ion Torrent sequencing platform[38] and revealed the evolution of families of DNA sequences, with enrichment particularly evident after 10 rounds of SELEX. Negative selection eliminated certain sequences that were not specific to the hyperphosphorylative state, or promoting hepatocyte uptake. However, the relative abundance of key sequences increased steadily throughout the whole process. The 23 most abundant sequences at round 26 were identified and their abundance throughout the SELEX process as calculated using AptaAligner[39] is shown in Fig. 2B . The sequence Tau1 exemplifies this behavior representing a whole 59% of cycle 26. A single base difference from this sequence, Tau3, is the second most represented sequence. The sequences present at the final round were grouped by hierarchical clustering and sequence homology using the multiple sequence alignment code MAFFT [40] showing five distinct families (Fig 2E) also presented as a cladogram (using the Clustal Omega[41]) showing the common ancestry between these five aptamers families (Fig. 2F). Elevated levels of Tau1 binding to the membrane of hyperphosphorylative cells is demonstrated in Figure 2C. The secondary structure of the aptamers Tau1 and Tau3 calculated using mfold[42] is shown in Fig 2D. The apparent equilibrium dissociation constants (Kd_{app}) were measured by serial dilution of aptamer solutions with target hyperphosphorylated and non-target differentiated and undifferentiated SH-SY5Y cells. The affinity of these aptamers was also tested with another immortal neural progenitor stem cell line ReN-VM[43] in hyperphosphorylative and non-hyperphosphorylative conditions . The Kd_{app} for Tau1 and Tau3 with hyperphosphorylated SH SY5Y cells is 0.167 ± 0.015 nM and 0.194 ± 0.032 nM; and for the ReN-VM cells 318.15 ± 46.2 nM and 234.24 ± 38.6 nM respectively (Supp. Fig 2).

TauX nanoparticles for magnetic resonance imaging

For *in vivo* imaging of the hyperphosphorylative state in the brain of live mice, aptamer-targeted nanoparticles were fabricated for use as a molecular MRI contrast agent (TauX). TauX, was formulated as two versions, one using the Tau1 aptamer (TauT1) and another using the Tau3 aptamer (TauT3). Liposomal nanoparticles were synthesized using a lipid mixture that included lipidized Gd-DOTA for MR imaging, cholesterol for liposomal stability; additionally we also incorporated lipidized rhodamine for studying *ex-vivo* microscopic distribution of liposomal nanoparticles in brain tissues using fluorescence microscopy[44,45]. The TauX compositions also included DSPE-mPEG2000 to increase the *in vivo* circulation time[34]. Particles had a hydrodynamic diameter of ~150nm, ~86,000 Gd-chelates per liposomes and ~500 aptamers conjugated to the outer leaflet of each liposomal nanoparticle (Fig.3)

In vivo molecular MRI using TauX for detection of hyperphosphorylative cells

To test if TauX could detect the hyperphosphorylative state *in vivo*, MRI studies were performed in a P301S transgenic mouse model of AD-related tauopathy. Studies were performed in transgenic and age-matched wild type mice at 2-3 months of age. At this young age, transgenic animals do not show frank tau pathology (i.e., neurofibrillary tangles), but practically all will develop tau pathology by around 8 months of age. Animals underwent baseline, pre-contrast MRI. Thereafter, animals were intravenously administered MRI contrast agent (TauT1, TauT3 or non-targeted control stealth liposomes that were not expected to provide signal enhancement as they did not have a targeting aptamer). Delayed post-contrast MRI was performed 4 days later. MR images were acquired using a T1-weighted spin-echo (T1w-SE) sequence and a T1-weighted fast spin echo inversion recovery (FSE-IR) sequence[46]. Transgenic mice administered TauT1 and TauT3 demonstrated signal enhancement in the cortex and the hippocampus regions of the brain (**Fig. 4B**). Wild-type mice (WT) administered TauT1 or TauT3 did not show signal enhancement in the brain. Similarly, transgenic mice administered non-targeted liposomal-Gd contrast agent did not show signal enhancement in cortex or hippocampus. These regions of interest were further analyzed quantitatively and signal-enhancement between the transgenic and wild-type mice were found to be statistically significant ($p < 0.05$) (Fig 4D). A baseline enhancement threshold of ~6% (=2X standard deviation of signal in baseline scans) was used as the classification threshold. Animals that showed signal enhancement above the threshold were identified as positives. Receiver operator characteristic (ROC) curves were generated with a six-point ordinal scale to assess sensitivity and specificity for detecting the

genotype, using TauT1 and TauT3 contrast agents, and constructed over the entire tested group, including

controls. The aptamer-targeted nanoparticle contrast agents, TauT1 and TauT3, showed overall AUC and

accuracy of ~0.95. TauT3 demonstrated higher sensitivity than TauT1.

Post-mortem brain analysis was performed in 2-3 month old transgenic and wild-type mice.

Immunofluorescence analysis using AT8 antibody staining revealed the presence of hyperphosphorylated tau

species in transgenic mice but absent in wild type mice (**Fig. 4F**). A 100% concordance was observed between

AT8 positivity and animal genotype. In summary, *in vivo* studies demonstrated that TauX enabled *in vivo*

molecular MRI of the hyperphosphorylative state months before frank tau pathology i.e. the presence of

neurofibrillary tangles, becomes evident in transgenic mice.

Target identification of aptamers

To characterize the binding target of the aptamers, we performed an aptamer-based pulldown assay,

aptamer-based immunoprecipitation, followed by mass-spectrometry. We performed the assay for both Tau1 and

Tau3, aptamers. A ranking of the abundance scores for identified proteins revealed keratin 6a, Keratin 6b and

Vimentin as possible binding targets (Table 1). The surface expression of Keratin 6a, 6b was similar on wild-type

and transgenic tissue sections whereas the Vim expression was higher in the transgenic mice (Fig 5C). SH-

SY5Y cells under undifferentiated, and differentiated hyperphosphorylative conditions show increasing levels of

vimentin (Fig.5B) further suggesting it is a potential target of aptamers Tau1 and Tau3.

Discussion

The ATN research framework suggests biomarkers to diagnose and classify AD. Under this framework

CSF based detection of Aβ, tau (total, and phosphorylated) have been reported but only at the prodromal stage

of disease, in patients with mild cognitive impairment[47]. Non-invasive neuroimaging tools, such as structural MRI

to diagnose and monitor neurodegeneration[48] show a definitive correlation with cognitive decline, visualizing

atrophic regions that depict neuronal injury in late stage disease[49]. However, a reliable marker of early

presymptomatic stage disease is yet to be identified.

While the role of Aβ and tau in the development of AD and the mechanism of transition from

presymptomatic to symptomatic AD are yet unclear, the time scale of the transition is generally accepted to be

over a period of 10-20 years[14]. Aβ deposits are considered the start of neurodegeneration but recent studies

indicate that tau pathology[50] shows a stronger correlation with disease progression suggesting that the limitation

of current tests is their inability to identify early stage pathological tau[15]. CSF presence of hyperphosphorylated

tau species p-181 and p-217 is associated with Aβ deposition that precedes a positive tau PET[51] but only has a

concordance of 50%-70% [52]. Taken as a whole, the roles of Aβ and tau deposition in disease progression and

the role of Aβ in the spread of initial tau aggregates, strongly suggest that a biomarker of pathological tau at a

presymptomatic stage of the disease is likely to advance detection by several years, and constitutes the

motivation for this work.

Initial tau aggregation is thought to be triggered by an imbalance in cellular homeostasis caused by

dysregulated phosphorylation[53,54]. Several kinases can phosphorylate tau at multiple locations; at least 45 sites

have been observed experimentally[55–57]. Combined with reduced phosphatase activities in AD, the altered

kinase-phosphatase balance yields hyperphosphorylative conditions that cause abnormal hyperphosphorylation

of tau[58]. Disruption of the normal function of tau, modulating microtubule dynamics by lowering its binding

capabilities[59] increases the level of cytosolic free tau leading to aggregation and fibrillization of tau that spreads

throughout the connected brain, seeding the pathology[60,61]. We hypothesized that this initial process of

hyperphosphorylation is associated with changes on the surface of hyperphosphorylative cells. We therefore

sought to identify these surrogate markers of tau hyperphosphorylation that presage future tau pathology.

Using SH-SY5Y cells as a model of neuronal hyperphosphorylation, we used a reverse-phase protein

array (RPPA) analysis to demonstrate elevated levels of surface molecules specific to the hyperphosphorylative

state (Fig 1C). Cell-SELEX capturing the differences between the surface of hyperphosphorylative cells and

normal cells allowed the selection of phosphorothioate modified short DNA aptamers that bound with high affinity

and specificity to hyperphosphorylative cells Fig 2 A. Having identified unique aptamers that bound such markers,

we developed MR molecular imaging contrast agents that recognize the surface of cells in hyperphosphorylative

state. We acknowledge that SH-SY5Y cells are not true neurons, they are a cell line originating in a

neuroblastoma, a tumor of embryological neural crest origin. However, they can be induced to differentiate to a

neuronal phenotype (as in the current work). While primary neuronal culture, or immortalized neuronal cells e.g.

ReN-VM may offer alternative models of neurons, we have functionally tested the aptamer hits from our SELEX

screen in a transgenic mouse model of tau deposition, and validated their performance, supporting our position

that the choice of cell model was adequate to identify suitable markers of tau hyperphosphorylation.

PET is the leading modality for clinical molecular imaging, driven by its high contrast sensitivity, however,

it suffers from poor spatial resolution on the order of 5-10mm, high cost; limited access to radioactive tracers,

and radiation exposure. Nanoparticle-enhanced MR imaging overcomes all these obstacles, but historically has

not achieved high enough sensitivity. We have previously demonstrated liposomal nanoparticles exhibiting large

numbers of Gd chelates in the external bilayer leaflet, with hyper-T1 relaxive properties resulting in contrast

sensitivity that rival nuclear imaging[45,62].

In P301S mice, the earliest reported histopathological studies are at age of 2.5months[63,64], and report

no tau pathology. "Tau seeding" the cell-cell transfer of pathogenic tau aggregates has been reported using brain

homogenates at 1.5 month of age[65]. We therefore chose P301S mice at 2 months of age for our studies when,

tau seeding should be taking place, but frank tau pathology should be absent. The mice were injected with TauX

nanoparticles targeted either by the Tau1 aptamer or the Tau3 aptamer. When imaged by T1-weighted MRI

sequences, designed to optimize signal from the Gd chelate induced T1 relaxation caused by the liposomal-Gd

nanoparticles, signal enhancement was observed in the cortex and hippocampus regions of the brain.

Hyperphosphorylative conditions were confirmed by post-mortem IF staining with AT8 antibody that recognizes

the S202 and T305 pTau species. Signal enhancement was not observed in non-transgenic mice, or in transgenic

mice injected with untargeted nanoparticles, supporting the specificity of Tau1-or Tau3-bearing nanoparticle

binding to target.

We have narrowed down the possible binding targets of the aptamers, and our data suggest that cell

surface vimentin is a likely target. We have confirmed the specific presence of cell surface vimentin on the surface

of SH-SY5Y cells in a hyperphosphorylative state, and on P301S mouse brain sections. Vimentin is an

intermediate filament protein that undergoes constant assembly and/or remodeling and is usually associated with

mesenchymal cells[66]. The assembly state of filaments is linked to their phosphorylation state, phosphorylation

promotes disassembly[67]. Vimentin contains more than 35 phosphorylation sites targeted by multiple kinases and

phosphatases allowing it to adjust IF dynamics dependent on its environment[68,69]. Mechanical, chemical (toxins,

hypoxia), and microbial stresses upregulate vimentin and its phosphorylation that allows cells to adjust their

mechanical properties[70,71]. The balance of different oligomeric forms influence dynamic cell processes including

adhesion, migration, invasion including stress-induced signaling[72]. Vim IF's (~10nm) distributed throughout the

cell by association with microtubules (tubulin, 24nm) regulating cell-migration, and microfilaments(actin, 7nm)

regulating cell-contractility, form the cytoskeletal network and provide mechanical support for the plasma

membrane where it contacts other cells or the extracellular matrix[73]. Interestingly, during the biological process

epithelial to mesenchymal transition wherein non-motile, polar epithelial cells transform to motile invasive non-

polar mesenchymal cells[74], cells also undergo a cytoskeletal reorganization that includes changes in cell-

membrane integrity, disassembly of junction proteins, increased stress-fiber formations, altered cell-surface

protein expression.   Changes in the localization of proteins is a hallmark of this pathologic process. Our

observation that Vimentin is upregulated and translocated to the cell surface[75] in the early stages of tau

hyperphosphorylation suggests a possible role for EMT-related processes at the start of a slow progression

towards AD pathology.

While positron emission tomography (PET) is the mainstay of molecular imaging, and exhibits

remarkable sensitivity, there are several limitations posed by this methodology[62,76]. Access to PET imaging is

limited, even in the relatively well-served US, and is skewed towards high density urban centers. PET costs are

very high due to the need for same-day radiosynthesis, and rapid decay of the isotopes. Longer half-life isotopes

cause higher radiation exposure. This tradeoff between half-life and radiation exposure greatly limit the reach of

PET to a wider patient population. Current PET tau tracers recognize the tau β-sheets in the PHF and NFT

present in tauopathies[22]. This conformation is not unique to tau and the *in vivo* specificity is circumspect limiting

its interpretation. Off-target binding of Flortaucipir, an approved tau PET agent, has been reported since it binds

the MAO-B enzyme in the brain[77,78]. Further, the vast majority of pathological tau is actually intracellular, posing

a significant barrier to PET tracers that must navigate to the site of tau pathology, bind the target, and have all

unbound tracer molecules cleared from the brain before the radioactive signal decays. Our choice of MRI as the

detection modality is based on hyper-T1 relaxive properties of nanoparticles with surface conjugated Gd chelates,

bringing detection sensitivity to the same range as nuclear imaging, and the MRI agent does not suffer from the

rapid signal decay of PET agents, allowing plenty of time for unbound tracer to clear from the brain before

imaging. Our choice of a cell surface surrogate marker of tau hyperphosphorylation avoids the need to bind an

intracellular target. Finally, MRI imaging is already included in AD management and can be adjusted with agents

such as TauX nanoparticles to constitute a highly sensitive and specific test for future tau pathology.

In summary, our work shows that the hyperphosphorylative conditions coinciding with the initiation of a

decades long process culminating in AD result in markers manifested on the cell surface. By imaging their

presence, our approach detects the presence of hyperphosphorylative cells indicative of a presymptomatic stage

of AD. Coupled with early amyloid detection, this test could enable early positive diagnosis using the NIA-AA

research framework.

Materials and Methods

Cell-lines - SH-SY5Y cells (ATCC, Manassas, VA, #CRL-2266™) were obtained from Dr. Jason Shohet's lab at

the Texas Children's Hospital, Houston, immortalized human hepatocytes (THLE-3) were purchased from

American Type Culture Collection (ATCC, Manassas, VA, # CRL-11233™); both were cultured according to the

ATCC instructions. ReN cell™ VM (#SCC008) cultured as per instruction using neural stem cell maintenance

medium (#SCM005) and growth factors EGF (GF001) and bFGF(#GF005) all from Millipore Sigma, Burlington,

MA.

Differentiation - SH-SY5Y cells were exposed to 30 µM *all-trans*-Retionic acid (Sigma-Aldrich, St.Louis, MO, #

R2625) in serum free cell medium for 10 days with medium change every alternate day. ReNcell VM were

differentiated by the removal of growth factors from its culture medium for 10 days.

Hyperphosphorylation - was induced in SH-SY5Y cells by addition of 30 nM okadaic acid (Sigma Aldrich,

St.Louis, MO, # 459620) in growth medium with 30 µM RA for 24 hours. ReNcell VM were hyperphosphorylated

using 100nM Quinolinic Acid (SigmaAldrich, St.Louis, MO, #P63204) in culture media for 24h.

Synthesis of primers and TA DNA library - All primers and Cy5, and amine labelled selected aptamers were

purchased from Integrated DNA Technologies (IDT, Coralville, IA). The ssDNA library used in Cell-SELEX

contained a central randomized sequence of 30 nucleotides flanked by PCR primer regions to enable the PCR

amplification of the sequence (5'-CGCTCGATAGATCGAGCTTCG-$(N)_{30}$-GTCGATCACGCTCTAGAGCACTG-

3'). The chemically synthesized DNA library was converted to a phosphorothioate modified library by PCR

amplification using, dATP (αS), resulting in the DNA sequences where the 3' phosphate of each residue is

substituted with monothiophosphate groups, as described previously in detail [32,79]. The reverse primer was

labeled with biotin to separate the sense strand from the antisense strand by streptavidin-coated sepharose

beads (PureBiotech, Middlesex, NJ, # MSTR0510) for the next selection round. The concentration of the TA

library was determined with a NanoDrop™ 2000 by measuring the UV absorbance at 260 nm.

Cell-SELEX - The procedures of Cell-SELEX were performed according to protocol previously with some

modification [33,80].The initial ssDNA library of 150 pmole was dissolved in binding buffer with a total volume of 350

μl. It was denatured by heating at 95°C for 5 min, then renatured by rapid cooling on ice for 10 min. The treated

SH-SY5Y cells at approximately 90% confluence in a 100-mm culture plate were washed twice with washing

buffer and followed by incubating with the ssDNA library of 150 pmole for 2 hrs at 4°C. Following the incubation,

for positive selections, the supernatant was discarded, and cells were washed three times with washing buffer to

remove any unbound sequences. Then cells were scraped off and transferred to nuclease-free water, following

another three times nuclease-free water washes. Cells in nuclease-free water was centrifuged at 300×g for 5

min. QIAamp DNA Mini and Blood Mini kit (Qiagen, Germantown, MD, # 51104) was introduced to elute cell

membrane fraction. The cell membrane fraction was PCR-amplified to monitor the presence of cell binding

efficacy at each cycle. For negative selections, the supernatant was simply pipetted out of the flask and

processed for the next cycle of selection. The desired compartment were amplified by PCR, and used to prepare

the TA for the next round of selection. Two different negative selections were involved. One was differentiated

treatment only SH-SY5Y cells at cycles #12 and #13. Another was hepatocyte THLE-3 cells at cycles #20 and

#21. A total of 26 cycles of Cell-SELEX were conducted, including two different types of negative selections

mentioned above.

Next-Generation Sequencing (NGS) - At the studied cycles, the membrane fractions were isolated and the

recovered TA sequences were amplified by PCR. Equimolar quantities of the recovered TA sequences over the

range were pooled together and sequenced by Next-Gen DNA sequencing using Ion318 chip (ThermoFisher,

Waltham, MA). A four base sequence was introduced during PCR amplification to serve as unique "barcode" to

distinguish between the studied cycles. Sequencing results were analyzed by the Aptalinger [39] that uses the

markov model probability theory to find the optimal alignment of the sequences.

Aptamer binding studies – were conducted with undifferentiated, differentiated and hyperphosphorylated SH

SY5Y and ReN cell VM grown in 96-wells seeded at 10000 per well. The apparent dissociation constants (Kds)

were measured by the equation Y=Bmax X/ (Kd+X), with GraphPad Prism 9, San Diego, CA, with a saturation

binding experiment; cells were incubated with varying concentrations of Cy5-labeled aptamer in a 100 μl volume

of binding buffer containing cells, incubated for 30minutes, washed twice and resuspended in 100 μl buffer and

analyzed by Molecular probes microplate reader equipped with the appropriate excitation and emission filters.

All data points were collected in triplicate.

Immunocytochemistry - Eight-well glass plate was coated with a solution of 100 μg/ml Collagen Type I

(Thermofisher Scientific, Waltham, MA, # A1064401) dissolved in 0.01N HCl and air dried, then PBS washed

and air dried prior to seeding 20,000 SH-SY5Y cells per well. Aptamer staining at 100 nM was performed with

live cells for 2h at 4ºC in binding buffer and washed twice with washing buffer. Cells were then fixed by incubation

for 15 min in 4% formaldehyde in PBS at room temperature. Non-specific binding was blocked with blocking

buffer (G-Biosciences, St. Louis, MO, # 786195) for 1 hour and overnight incubation at 4°C with the rabbit pTau

primary antibody (1:100) (Santa Cruz Biotechnology, #: sc-101815) was followed by washing with PBS, and 1h

incubation with goat anti-rabbit IgG secondary antibody, Alexa Fluor 488 (Invitrogen, Carlsbad, CA, # A-11008)

for 1 hour at room temperature. Cytoskeletal actin filaments were stained with Alexa Fluor 594 Phalloidin

(Invitrogen, # A12381). The cells were covered with VECTASHIELD hardset mounting medium with DAPI (Vector

Laboratories, Burlingame, CA, # H-1500) for 5 min at room temperature. Images were visualized under Olympus

Fluoview FV1000 confocal microscopy.

TauX nanoparticle synthesis – L-α-phosphatidylcholine, hydrogenated (Hydro Soy PC; HSPC) and Cholesterol

were purchased from Lipoid Inc., Newark NJ, USA. 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-

[methoxy(polyethylene glycol)-2000] (DSPE-mPEG2000) was purchased from Corden Pharma, Liestahl,

Switzerland. DSPE-PEG3400-COOH and Gd-DOTA-DSPE were synthesized *in house*, lis-rhodamine-DHPE

from ThermoFisher Scientific. HSPC, Cholesterol, DSPE-PEG3400-COOH, DSPE-mPEG2000, Gd-DOTA-

DSPE, lis-rhodamine-DHPE at molar proportions 31.4:40:0.5:3:25:0.1 were dissolved in ethanol to achieve a

total concentration of 100 mM. For the non-targeted control stealth liposomes, carboxy terminated PEG was not

included in the lipid mixture. The ethanolic solution of lipids was hydrated with 150mM saline solution at 65 °C

for 30 minutes, allowing multilamellar liposomes to form. The mixture was then extruded in a 10ml Lipex extruder

(Northern Lipids Inc., Burnaby, Canada) using a 400 nm polycarbonate track-etch polycarbonate filter (3 passes)

followed by a 200 nm (3 passes) and finally 100nM filters. The suspension was then diafiltered using a MicroKros

cross-flow diafiltration cartridge (*500 kDa* cutoff) from Repligen, Rancho Dominguez, CA, exchanging the external

buffer for phosphate buffered saline (PBS, pH 7.2) for 15 volume exchanges. To form the aptamer conjugated

liposomes, liposomes with lipid-PEG-COOH were reacted with amine terminated aptamers using carbodiimide

chemistry. The carboxyl groups on the liposomes were activated with 5 mM EDC and 10 mM sulfo-NHS at pH~6

for 5-10 minutes. The activated liposomes were then immediately reacted with the amine terminated aptamers

and the pH was raised to ~7.3 - 7.6 by titrating µl amounts of 5 N NaOH. The final concentration of aptamers

used in reaction is ~140 µM. The reaction was mixed at room temperature for 1 hr following which the reaction

was carried out at 4 deg ^{0}C overnight. The liposomes were then dialyzed against PBS to remove unconjugated

aptamers using a 300 kDa dialysis membrane. The dialysate (external phase) was concentrated using 10 kD

centrifugal separator and washed with PBS to remove residual EDC/s-NHS. The concentrated dialysate was

analysed by NanoDrop Spectrophotometer (ThermoFisher Sci., Waltham, MA, USA) to determine unconjugated

aptamer fraction, and estimate aptamer density per nanoparticle in TauX formulations. Inductively coupled

plasma atomic emission spectroscopy (ICP-AES) was used to measure Gd and phosphorus concentrations of

TauX formulations. The hydrodynamic diameter of liposomal nanoparticles in TauX formulations was determined

using a dynamic light scattering instrument.

Mice – All the procedures were performed with approval from Institutional Animal Care and Use Committee

(IACUC) of Baylor College of Medicine. Mice were kept under a 12 h light/dark cycle, with food and water available

ad libitum. PS19 mice from Jackson Laboratories (Bar Harbor, ME) B6; C3-Tg (Prnp-MAPT*P301S) PS19Vle/J

Stock No: 008169 were used and experiments were conducted at the 2m of age. The transgenic (TG) mice

develop neurofibrillary tangles by 5 months of age[64]. Age-matched non-transgenic wild type (WT) mice were

used as controls.

Magnetic Resonance Imaging (MRI) - MRI was performed on a 1T permanent magnet scanner (M7, Aspect

Imaging, Shoham, Israel). Mice underwent pre-contrast baseline scans. Thereafter, mice were intravenously

administered one of three nanoparticle MR contrast agents (TauT1, TauT3 or non-targeted control liposomes)

via tail vein at a dose of 0.20 mmol Gd/kg of body weight. Delayed post-contrast MRI was performed 4 days after

contrast agent injections. Pre-contrast and delayed post-contrast MR images were acquired using a T1-weighted

spin echo (T1w-SE) sequence and a fast spin echo inversion recovery (FSE-IR) sequence with the following

parameters: _SE parameters_: TR = 600 ms, TE = 11.5 ms, slice thickness = 1.2 mm, matrix = 192 × 192, FOV = 30

mm, slices = 16, NEX = 4; _FSE-IR parameters_: TR = 13500 ms, TE = 80 ms, TI = 2000 ms, slice thickness = 2.4

mm, matrix = 192 × 192, FOV = 30 mm, slices = 6, NEX = 6. Coil calibration, RF calibration, and shimming were

performed at the beginning of study for each subject. The pre-contrast scans provide a baseline for calculation

of signal enhancement from resulting post-contrast scans[46]. Two-standard deviations above the mean variation

within WT control animals was used as the cutoff signal intensity for identifying tau positive animals. Six

transgenic (TG) mice and six wild type mice (WT) were used for testing of each nanoparticle contrast agent

formulation. Receiver operating characteristic (ROC) curves were generated on a six point ordinal scale by

plotting the true positive fraction (TPF) against the false positive fraction (FPF) based on imaging-based

identification of Tau-positive animals using the cutoff signal intensity and then comparing against histological

confirmation of Tau pathology as a gold standard. A fitted curve was then generated against the empirical points

plotted on the graphs. Qualitative and quantitative analysis of MRI images was performed in OsiriX (version 5.8.5,

64-bit, Pixmeo SARL, Geneva, Switzerland) and MATLAB (version 2015a, MathWorks, Natick, MA).

Immunofluorescence – After the final MRI scan, the mice were euthanized and perfused extensively with 0.9%

saline followed by 4% paraformaldehyde for 15 mins. The brains were then immersion-fixed in 4% formaldehyde

for 48h at 4ºC, transferred to 30% sucrose for cryoprotection and embedded in OCT. Phenotypic confirmation for

the presence of phosphorylated tau and vimentin was done on 25µm thick brain sections. Antigen retrieval in

pH=8.5 citrate buffer was executed in a 1200W GE microwave for 15mins. After 15 minutes of cooling 25µL of

1:50 dilution of primary p-tau antibody namely either AT8,AT100 or AT180 that recognize different p-tau species

were incubated in a tray (RPI, Mt. Prospect, IL #248270) designed for microwave enhanced immunostaining

procedures for 3min at power level 3. After a 2 min cooling, sections were washed with PBS and incubated for

3min with a 1:100 dilution of appropriate secondary antibody. DAPI staining proceeded after 2 mins of cooling

and a PBS washing. ProGold Antifade (Invitrogen, Carlsbad, CA, # P36030) was used to mount slides which

were visualized on Olympus Fluoview LV100. Scanning of whole sections was also conducted using a Biotek

Cytation 5 slide scanning microscope. List of Antibodies – AT8 (#MN1020), Vimentin SP20 (#MA516409) both

Thermo Fisher Scientific, Waltham, MA, Vimentin D21H3 (Cell Signaling Technology, Beverly, MA, #5741T,),

Cell-surface vimentin (Abnova, Taipei City, Taiwan, #H00007431-M08J).

Target Identification - The protein targets of Tau-1, Tau-3, Tau-4 and Tau-5 were identified by affinity-pull down

using the selected aptamers as the capturing reagent followed by mass-spectroscopy. A scrambled DNA

sequence, R2, was used as a control. The hyperphosphorylated SH-SY5Y cells, at 90-95% confluence were

washed with cold PBS buffer and incubated with biotinylated selected aptamers with 25mmol/l each at 4°C in

PBS, respectively. After 2 hours of gentle agitation, SH-SY5Y cells were cross-linked with 1% formaldehyde for

10 minutes at room temperature. The formaldehyde cross-linking was quenched with glycine. Cells were scraped

from the plate, washed and lysed with lysing buffer (Thermofisher Scientific, # 87787) and treated with protease

inhibitor mixture. The lysates were freeze-thawed for 30 minutes on ice and cleared by centrifuging at 10,000 ×g

for 2 minutes at 4°C. To pull down the cross-linked proteins, equal amounts of cell lysate were incubated with

prewashed streptavidin magnetic beads for 1 hour at room temperature with continuous rotation. Protein

digestions were performed on the beads to isolate targeted proteins and processed for mass spectrometric

analysis. Each sample was analyzed in triplicates. The raw data files were processed to generate a Mascot

Generic Format with Mascot Distiller and searched against the SwissProt_2012_01 (Human) database using the

licensed Mascot search engine v2.3.02 (Matrix Science, Boston, MA) run on an in-house server.

References

1.   Goedert, M., Wischik, C. M., Crowther, R. A., Walker, J. E. & Klug, A. Cloning and sequencing of

the cDNA encoding a core protein of the paired helical filament of Alzheimer disease: identification

as the microtubule-associated protein tau. *Proc. Natl. Acad. Sci. U. S. A.* **85**, 4051–4055 (1988).

2.   Goedert, M., Spillantini, M. G., Jakes, R., Rutherford, D. & Crowther, R. A. Multiple isoforms of

human microtubule-associated protein tau: sequences and localization in neurofibrillary tangles of

Alzheimer's disease. *Neuron* **3**, 519–526 (1989).

3.   Iqbal, K., Liu, F. & Gong, C. Tau and neurodegenerative disease: the story so far. *Nat. Rev.*

Neurol. **12**, 15–27 (2016).

4.   Guo, T., Noble, W. & Hanger, D. Roles of tau protein in health and disease. *Acta Neuropathol.*

133, 665–704 (2017).

5.   Wang, Y. & Mandelkow, E. Tau in physiology and pathology. *Nat. Rev. Neurosci.* **17**, 22 (2016).

6.   Chong, F. P., Ng, K. Y., Koh, R. Y. & Chye, S. M. Tau Proteins and Tauopathies in Alzheimer's

Disease. *Cell. Mol. Neurobiol.* **38**, 965–980 (2018).

7.   Karikari, T. K. *et al.* Distinct conformations, aggregation and cellular internalization of different tau

strains. *Front. Cell. Neurosci.* **13**, 1–16 (2019).

8.   d'Errico, P. & Meyer-Luehmann, M. Mechanisms of Pathogenic Tau and Aβ Protein Spreading in

Alzheimer's Disease. *Front. Aging Neurosci.* **12**, 1–10 (2020).

9.   Brunello, C. A., Merezhko, M., Uronen, R. L. & Huttunen, H. J. Mechanisms of secretion and

spreading of pathological tau protein. *Cell. Mol. Life Sci.* **77**, 1721–1744 (2020).

10.   Vogel, J. W. *et al.* Spread of pathological tau proteins through communicating neurons in human

Alzheimer's disease. *Nat. Commun.* **11**, 2612 (2020).

11.   Gibbons, G. S., Lee, V. M. Y. & Trojanowski, J. Q. Mechanisms of Cell-to-Cell Transmission of
Pathological Tau: A Review. *JAMA Neurol.* **76**, 101–108 (2019).

12.   Jack Jr, C. R. *et al.* NIA-AA research framework: toward a biological definition of Alzheimer's
disease. *Alzheimer's Dement.* **14**, 535–562 (2018).

13.   Jack, C. R. *et al.* A/T/N: an unbiased descriptive classification scheme for Alzheimer disease
biomarkers. *Neurology* **87**, 539–547 (2016).

14.   Selkoe, D. J. & Hardy, J. The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol.*
Med. **8**, 595–608 (2016).

15.   Mattsson-Carlgren, N. *et al.* Aβ deposition is associated with increases in soluble and
phosphorylated tau that precede a positive Tau PET in Alzheimer's disease. *Sci. Adv.* **6**, (2020).

16.   Leuzy, A. *et al.* Longitudinal tau and metabolic PET imaging in relation to novel CSF tau measures
in Alzheimer's disease. *Eur. J. Nucl. Med. Mol. Imaging* **46**, 1152–1163 (2019).

17.   Cummings, J. L. Biomarkers in Alzheimer's disease drug development. *Alzheimer's Dement.* **7**,
e13–e44 (2011).

18.   Smoler, M., Coceano, G., Testa, I., Bruno, L. & Levi, V. Apparent stiffness of vimentin intermediate
filaments in living cells and its relation with other cytoskeletal polymers. *Biochim. Biophys. Acta -*
Mol. Cell Res. **1867**, 118726 (2020).

19.   Peskind, E., Nordberg, A., Darreh-Shori, T. & Soininen, H. Safety of lumbar puncture procedures
in patients with Alzheimer's disease. *Curr. Alzheimer Res.* **6**, 290–292 (2009).

20.   Wittenberg, R., Knapp, M., Karagiannidou, M., Dickson, J. & Schott, J. M. Economic impacts of
introducing diagnostics for mild cognitive impairment Alzheimer's disease patients. *Alzheimer's*
Dement. Transl. Res. Clin. Interv. **5**, 382–387 (2019).

21.   Ehman, E. C. *et al.* PET/MRI: where might it replace PET/CT? *J. Magn. Reson. Imaging* **46**, 1247–
1262 (2017).

22.   Leuzy, A. *et al.* Tau PET imaging in neurodegenerative tauopathies—still a challenge. *Mol.*

Psychiatry **24**, 1112–1134 (2019).

23.   Jack Jr, C. R. *et al.* Tracking pathophysiological processes in Alzheimer's disease: an updated

hypothetical model of dynamic biomarkers. *Lancet Neurol.* **12**, 207–216 (2013).

24.   Terry, R. D. *et al.* Physical basis of cognitive alterations in Alzheimer's disease: synapse loss is

the major correlate of cognitive impairment. *Ann. Neurol. Off. J. Am. Neurol. Assoc. Child Neurol.*

Soc. **30**, 572–580 (1991).

25.   Hyman, B. T. *et al.* National Institute on Aging–Alzheimer's Association guidelines for the

neuropathologic assessment of Alzheimer's disease. *Alzheimer's Dement.* **8**, 1–13 (2012).

26.   Montine, T. J. *et al.* National Institute on Aging–Alzheimer's Association guidelines for the

neuropathologic assessment of Alzheimer's disease: a practical approach. *Acta Neuropathol.* **123**,

1–11 (2012).

27.   Wang, J.-Z., Xia, Y.-Y., Grundke-Iqbal, I. & Iqbal, K. Abnormal hyperphosphorylation of tau: sites,

regulation, and molecular mechanism of neurofibrillary degeneration. *J. Alzheimer's Dis.* **33**,

S123–S139 (2013).

28.   Grundke-Iqbal, I. *et al.* Abnormal phosphorylation of the microtubule-associated protein tau (tau) in

Alzheimer cytoskeletal pathology. *Proc. Natl. Acad. Sci.* **83**, 4913–4917 (1986).

29.   Delobel, P. *et al.* Analysis of tau phosphorylation and truncation in a mouse model of human

tauopathy. *Am. J. Pathol.* **172**, 123–131 (2008).

30.   Lund, E. T., McKenna, R., Evans, D. B., Sharma, S. K. & Mathews, W. R. Characterization of the

in vitro phosphorylation of human tau by tau protein kinase II (cdk5/p20) using mass spectrometry.

J. Neurochem. **76**, 1221–1232 (2001).

31.   Liu, F., Grundke-Iqbal, I., Iqbal, K. & Gong, C. Contributions of protein phosphatases PP1, PP2A,

PP2B and PP5 to the regulation of tau phosphorylation. *Eur. J. Neurosci.* **22**, 1942–1950 (2005).

32.   Mu, Q. *et al.* Conjugate-SELEX: A High-throughput Screening of Thioaptamer-liposomal

Nanoparticle Conjugates for Targeted Intracellular Delivery of Anticancer Drugs. *Mol. Ther. -*

Nucleic Acids **5**, e382 (2016).

33.   Parekh, P., Tang, Z., Turner, P. C., Moyer, R. W. & Tan, W. Aptamers Recognizing Glycosylated

Hemagglutinin Expressed on the Surface of Vaccinia Virus-Infected Cells. *Anal. Chem.* **82**, 8642–

8649 (2010).

34.   Ayyagari, A. L. *et al.* Long-circulating liposomal contrast agents for magnetic resonance imaging.

Magn. Reson. Med. An Off. J. Int. Soc. Magn. Reson. Med. **55**, 1023–1029 (2006).

35.   Kamat, P. K., Rai, S., Swarnkar, S., Shukla, R. & Nath, C. Molecular and Cellular Mechanism of

Okadaic Acid (OKA)-Induced Neurotoxicity: A Novel Tool for Alzheimer's Disease Therapeutic

Application. *Mol. Neurobiol.* **50**, 852–865 (2014).

36.   Gulaj, E., Pawlak, K., Bien, B. & Pawlak, D. Kynurenine and its metabolites in Alzheimer's disease

patients. *Adv. Med. Sci.* **55**, 204–211 (2010).

37.   The UniProt Consortium. UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids*

Res. **49**, D480–D489 (2021).

38.   Rothberg, J. M. *et al.* An integrated semiconductor device enabling non-optical genome

sequencing. *Nature* **475**, 348–352 (2011).

39.   Lu, E., Elizondo-Riojas, M.-A., Chang, J. T. & Volk, D. E. Aptaligner: automated software for

aligning pseudorandom DNA X-aptamers from next-generation sequencing data. *Biochemistry* **53**,

3523–3525 (2014).

40.   Katoh, K., Rozewicki, J. & Yamada, K. D. MAFFT online service: multiple sequence alignment,

interactive sequence choice and visualization. *Brief. Bioinform.* **20**, 1160–1166 (2019).

41.   Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence alignments

using Clustal Omega. *Mol. Syst. Biol.* **7**, 539 (2011).

42.   Zuker, M. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids*

Res. **31**, 3406–3415 (2003).

43.   Donato, R. *et al.* Differential development of neuronal physiological responsiveness in two human

497   neural stem cell lines. *BMC Neurosci.* **8**, 1–11 (2007).

498   44.   Ghaghada, K., Hawley, C., Kawaji, K., Annapragada, A. & Mukundan Jr, S. T1 relaxivity of core-

499   encapsulated gadolinium liposomal contrast agents—effect of liposome size and internal

500   gadolinium concentration. *Acad. Radiol.* **15**, 1259–1263 (2008).

501   45.   Ghaghada, K. B. *et al.* New Dual Mode Gadolinium Nanoparticle Contrast Agent for Magnetic

502   Resonance Imaging. *PLoS One* **4**, e7628 (2009).

503   46.   Badachhape, A. A. *et al.* Pre-clinical dose-ranging efficacy, pharmacokinetics, tissue

504   biodistribution, and toxicity of a targeted contrast agent for MRI of amyloid deposition in

505   Alzheimer's disease. *Sci. Rep.* **10**, 16185 (2020).

506   47.   Mattsson, N., Lönneborg, A., Boccardi, M., Blennow, K. & Hansson, O. Clinical validity of

507   cerebrospinal fluid Aβ42, tau, and phospho-tau as biomarkers for Alzheimer's disease in the

508   context of a structured 5-phase development framework. *Neurobiol. Aging* **52**, 196–213 (2017).

509   48.   Coch, R. A. & Leube, R. E. Intermediate filaments and polarization in the intestinal epithelium.

510   *Cells* **5**, 32 (2016).

511   49.   Cuingnet, R. *et al.* Automatic classification of patients with Alzheimer's disease from structural

512   MRI: a comparison of ten methods using the ADNI database. *Neuroimage* **56**, 766–781 (2011).

513   50.   Johansson, M. *et al.* Mild behavioral impairment and its relation to tau pathology in preclinical

514   Alzheimer's disease. *Transl. Psychiatry* **11**, 1–8 (2021).

515   51.   Suárez-Calvet, M. *et al.* Novel tau biomarkers phosphorylated at T181, T217 or T231 rise in the

516   initial stages of the preclinical Alzheimer's continuum when only subtle changes in Aβ pathology

517   are detected. *EMBO Mol. Med.* **12**, e12921 (2020).

518   52.   Mattsson, N. *et al.* 18 F-AV-1451 and CSF T-tau and P-tau as biomarkers in Alzheimer's disease

519   . *EMBO Mol. Med.* **9**, 1212–1223 (2017).

53.   Chung, S.-H. Aberrant phosphorylation in the pathogenesis of Alzheimer's disease. *BMB Rep.* **42**,

467–474 (2009).

54.   Oliveira, J., Costa, M., de Almeida, M. S. C., da Cruz e Silva, O. A. B. & Henriques, A. G. Protein

phosphorylation is a key mechanism in Alzheimer's disease. *J. Alzheimer's Dis.* **58**, 953–978

(2017).

55.   Augustinack, J. C., Schneider, A., Mandelkow, E.-M. & Hyman, B. T. Specific tau phosphorylation

sites correlate with severity of neuronal cytopathology in Alzheimer's disease. *Acta Neuropathol.*

103, 26–35 (2002).

56.   Wang, Y. & Mandelkow, E. Tau in physiology and pathology. *Nat. Rev. Neurosci.* **17**, 22–35

(2016).

57.   Wang, J., Grundke-Iqbal, I. & Iqbal, K. Kinases and phosphatases and tau sites involved in

Alzheimer neurofibrillary degeneration. *Eur. J. Neurosci.* **25**, 59–68 (2007).

58.   Sontag, J.-M. & Sontag, E. Protein phosphatase 2A dysfunction in Alzheimer's disease. *Front.*

Mol. Neurosci. **7**, 16 (2014).

59.   Sontag, J. M., Nunbhakdi-Craig, V., White, C. L., Halpain, S. & Sontag, E. The protein

phosphatase PP2A/Bα binds to the microtubule-associated proteins Tau and MAP2 at a motif also

recognized by the kinase Fyn: Implications for tauopathies. *J. Biol. Chem.* **287**, 14984–14993

(2012).

60.   Franzmeier, N. *et al.* Functional brain architecture is associated with the rate of tau accumulation

in Alzheimer's disease. *Nat. Commun.* **11**, 1–17 (2020).

61.   DeVos, S. L. *et al.* Synaptic tau seeding precedes tau pathology in human Alzheimer's disease

brain. *Front. Neurosci.* **12**, 267 (2018).

62.   Dustin, D., Hall, B. M., Annapragada, A. & Pautler, R. G. Neuroimaging in Alzheimer's disease:

preclinical challenges toward clinical efficacy. *Transl. Res.* **175**, 37–53 (2016).

63.   Takeuchi, H. *et al.* P301S mutant human tau transgenic mice manifest early symptoms of human

tauopathies with dementia and altered sensorimotor gating. *PLoS One* **6**, e21050 (2011).

64. Yoshiyama, Y. *et al.* Synapse loss and microglial activation precede tangles in a P301S tauopathy

mouse model. *Neuron* **53**, 337–351 (2007).

65. Holmes, B. B. *et al.* Proteopathic tau seeding predicts tauopathy in vivo. *Proc. Natl. Acad. Sci.*

111, E4376 LP-E4385 (2014).

66. Eriksson, J. E. *et al.* Introducing intermediate filaments: from discovery to disease. *J. Clin. Invest.*

119, 1763–1771 (2009).

67. Sihag, R. K., Inagaki, M., Yamaguchi, T., Shea, T. B. & Pant, H. C. Role of phosphorylation on the

structural dynamics and function of types III and IV intermediate filaments. *Exp. Cell Res.* **313**,

2098–2109 (2007).

68. Snider, N. T. & Omary, M. B. Post-translational modifications of intermediate filament proteins:

mechanisms and functions. *Nat. Rev. Mol. Cell Biol.* **15**, 163–177 (2014).

69. Battaglia, R. A., Delic, S., Herrmann, H. & Snider, N. T. Vimentin on the move: New developments

in cell migration [version 1; referees: 2 approved]. *F1000Research* **7**, 1–10 (2018).

70. Pattabiraman, S. *et al.* Vimentin protects differentiating stem cells from stress. *Sci. Rep.* **10**, 1–15

(2020).

71. Etienne-Manneville, S. Cytoplasmic Intermediate Filaments in Cell Biology. *Annu. Rev. Cell Dev.*

Biol. **34**, 1–28 (2018).

72. Ivaska, J., Pallari, H.-M., Nevo, J. & Eriksson, J. E. Novel functions of vimentin in cell adhesion,

migration, and signaling. *Exp. Cell Res.* **313**, 2050–2062 (2007).

73. Jones, J. C. R. *et al.* Intermediate filaments and the plasma membrane. *Cold Spring Harb.*

Perspect. Biol. **9**, (2017).

74. Kalluri, R. EMT: when epithelial cells decide to become mesenchymal-like cells. *J. Clin. Invest.*

119, 1417–1419 (2009).

75. Ivaska, J. Vimentin: Central hub in EMT induction? *Small GTPases* **2**, 1436–1448 (2011).

76. Hameed, S. *et al.* Role of Fluid Biomarkers and PET Imaging in Early Diagnosis and its Clinical

Implication in the Management of Alzheimer's Disease. *J. Alzheimer's Dis. Reports* **4**, 21–37

(2020).

77. Lemoine, L. *et al.* Comparative binding properties of the tau PET tracers THK5117, THK5351,

PBB3, and T807 in postmortem Alzheimer brains. *Alzheimers. Res. Ther.* **9**, 1–13 (2017).

78. Baker, S. L., Harrison, T. M., Maass, A., La Joie, R. & Jagust, W. J. Effect of off-target binding on

18F-Flortaucipir variability in healthy controls across the life span. *J. Nucl. Med.* **60**, 1444–1451

(2019).

79. Volk, D. E. & Lokesh, G. L. R. Development of phosphorothioate DNA and DNA thioaptamers.

Biomedicines **5**, 41 (2017).

80. Mu, Q. *et al.* Conjugate-SELEX: A high-throughput screening of thioaptamer-liposomal

nanoparticle conjugates for targeted intracellular delivery of anticancer drugs. *Mol. Ther. Acids* **5**,

e382 (2016).

A

Cell Status 1   2   3   4   5   6   Time

- Cytokines
- Receptor
- Tau
- Phosphorylated Tau
- Tau Paired Helical Filaments
- Tau Neurofibrillary Tangles
- β-amyloid
- β-amyloid plaques
- Changes on cell membrane

1. Healthy cells
2. Cells in hyperphosphorylative conditions
3. Changes on cell surface of hyperphosphorylated cells
4. β amyloid fibres; pTau oligomers
5. β amyloid sheets; Tau PHF's
6. β amyloid plaques; Tau NFT's

B

Undifferentiated SH-SY5Y → Retinoic Acid 30uM 10 days → Differentiated SH-SY5Y → Okadaic Acid 30nM 24 hrs → Hyperphosphorylated SH-SY5Y

Figure 1: Hyperphosphorylation and cell surface changes in a cell of neuronal phenotype. (A) _Cell changes during hyperphosphorylation_: Tau hyperphosphorylation is a precursor to tau fibrillation. We hypothesize that cell surface changes occur in parallel with tau hyperphosphorylation. These changes may constitute a marker for early detection of AD. **(B)** _In vitro model of hyperphosphorylation in a cell of neuronal phenotype_: Differentiated SH-SY5Y cells are used to model hyperphosphorylation. Hyperphosphorylation can be induced by either okadaic acid (OA) or quinolinic acid (QA) **(C)** _Protein analysis at different stages of hyperphosphorylation_: Reverse phase protein array analysis of undifferentiated, retinoic acid (RA) differentiated and hyperphosphorylative cells (by Okadaic acid OA or Quinolinic acid QA). Heat map depicts changes in total protein expression. Proteins expressed on cell-membrane are depicted in red, peripheral –membrane proteins in blue and single-pass and multipass membrane proteins in green. Panel i) depicts changes in un-phosphorylated proteins on undifferentiated and ii) differentiated SH-SY5Y cells. Changes due to phosphorylation treatments with OA and QA on differentiated cells are shown in panels iii) undifferentiated and iv) differentiated SH-SY5Y cells

Figure 2: Cell-SELEX to identify biomarkers onset of AD. Okadaic acid treated differentiated SH-SY5Y cells were used as a surrogate for hyperphosphorylative neurons to screen DNA aptamers that specifically recognize the differences between the surfaces of treated and untreated cells, using the cell-SELEX methodology modified to capture membrane binding aptamers. (A) Pictorial representation of the cell-SELEX process (B) Abundance of the top 23 sequences from SELEX cycle 1 - 26 depicting their evolution. Note that the fractions are low until about cycle 10, when they increase sharply. Note that the abundance of Tau 1, Tau 3 continually increase with increasing cycle number. (C) Undifferentiated, differentiated (RA) and hyperphosphorylated (OA) SH-SY5Y cells stained with 50nM Cy5 labelled Tau1 aptamer for 2h at 4ºC. (D) i) Tau 1 and ii) Tau 3 secondary structure using Mfold. (E) Multiple sequence alignment of the top 23 aptamer sequences by MAFFT. Note the existence of five families. (F) Cladograms showing relationship between the Tau aptamer sequences and the aptamer families.

Gd based Imaging moiety

Gd(III)-DOTA-DSPE

Rhodamine

Aptamer (T1/T3)

Lipid bilayer

~ 150 nm

mPEG$_{2000}$-DSPE

Aptamer Conjugated Polyethylene Glycol -Lipid

Carboxy-Polyethylene glycol –DSPE (COOH-PEG-DSPE)

Na$^+$

(OCH$_2$CH$_2$)$_{43}$

5^l Amino Aptamer

NH$_2$

Figure 3: Aptamer-targeted liposomal-Gd nanoparticle TauX contrast agent. The liposomal bilayer incorporates DSPE-DOTA-Gd for MR contrast, DSPE-PEG2000 to enhance circulation half-life, DSPE-PEG3400-aptamer (Tau1/Tau3) for targeting and lissamine rhodamine for fluorescence imaging.

Figure 4: T1-weighted spin-echo (T1w-SE) and (FSE-IR) images demonstrate signal enhancement as TauX binds to hyperphosphorylated tau *in vivo*. (A) In vivo imaging was performed on a 1 T permanent MRI scanner (M7, Aspect Imaging). Two month old mice were injected with 0.2mM Gd dose TauX NP. Pre- and post-contrast images were acquired with the following parameters: Imaging sequence: T1 weighted spin echo (T1w-SE) TR = 600 ms, TE = 11.5 ms, slice thickness = 1.2 mm, matrix = 192x192, FOV = 30 mm, slices = 16, NEX = 4. Animals were euthanized after final imaging session and brains harvested for post-mortem analysis. (**B**) and (**C**) demonstrate signal enhancement in delayed post-contrast scans of transgenic (Tg) P301S mice treated with TauX relative to age-matched controls. Wild type (WT) animal shows no signal enhancement four days after administration of TauX. Tg animal showed high enhancement in cortical (yellow arrow) and hippocampal regions (white arrow) four days after administration of TauX. Transgenic animal shows no signal enhancement four days after injection of untargeted contrast (UC). Scale bar represents 2 mm. All animals are shown on the same color scale. (**D**) Signal enhancement in TG animals relative to WT counterparts and UC-treated Tg animals for both T1w-SE and FSE-IR sequences (*p<0.05 ; **p<0.005). (**E**) The Receiver operating characteristic (ROC) curve demonstrates TauX accuracy in identifying early age Tg animals. ROC curve is generated by fitting observed operating points for true positive fraction versus false positive fraction (green dots) using a six point ordinal scale. A fitted curve (blue) connects the observed operating points. Area under curve (AUC) is calculated using the fitted curve. Sensitivity describes the true positive rate for Tg mice given TauX (n=6) while specificity is the true negative rate for WT mice given TauX (n=6), WT mice given UC Gd nanoparticle contrast (n=6), and Tg mice receiving UC Gd nanoparticle contrast (n=6). (F) The mice selected for TauX injection were genotyped which only revealed the presence of transgene. We confirmed the presence of hyperphosphorylated tau species in the mice by immunofluoscence on the brains sections harvested post MRI scans. Brains were perfused with heparin-PBS and fixed in OCT. IF with antibody AT8 probed for presence of ptau species provided high concordance with genotype confirming the presence of hyperphosphorylative conditions in the mice.

Figure 5: Proteomic analysis for aptamer target identification. A. Procedure for tryptic digest of aptamer bound cell membrane analyzed by LC/MS/MS. The raw data files were processed and searched against the SwissProt 2012 01 (Human) database. Vimentin was identified as possible target. B. Presence of the aptamers binding target on SH SY5Y cells; undifferentiated and hyperphosphorylated OA (24h, 30nM) and QA (24h, 100nM); i) co-stained with Vimentin (D21H3) and ptau (AT100) antibodies or ii) cell surface vimentin (Clone 84-1) antibody and aptamer T1 (50nM); nuclei counterstained with DAPI. C. Expression of Vimentin in P301S TG and WT frozen mouse tissue sections stained with Vimentin (SP20) and ptau (AT100) antibodies and DAPI stained nuclei.

Figure 6: Early detection of AD. Hyperphosphorylative conditions provoke neuronal cells to undergo changes including on the cell surface. Translocation of Vimentin to the cell surface was exploited to generate an aptamer that was developed for *in vivo* use as targeted liposomal nanoparticle TauX for early MRI detection of tau pathology in a mouse model

It is NOT AD. AD is a human disease. This is a model of tau deposition. The scanner shown is a human scanner.

Supplementary Table 1: Target Identification using aptamer immunoprecipitation and mass spectrometry. <mark>The aptamer immunoprecipation tryptic digest</mark> was purified with a ZipTip and analyzed by LC/MS/MS on an LTQ-Orbitrap-XLmass spectrometer (Thermo Fisher) interfaced with an EksigentNano-LC-Ultra-2D plus CHiPLC Nanoflex system (AB SCIEX). The raw data files were processed to generate a Mascot Generic Format with Mascot Distiller and searched against the SwissProt_2012_01 (Human) database using the licensed Mascot search engine v2.3.02(Matrix Science) run on an in-house server. Exponentially modified protein abundance index (emPAI) are reported in the table below that report protein content proportional to protein content in a mixture.

T1 emPAI	Target ID Aptamer T1	T3 emPAI	Target ID Aptamer T3
6.32	60 kDa heat shock protein, mitochondrial OS=Homo sapiens GN=HSPD1 PE=1 SV=2	4.45	Keratin, type II cytoskeletal 1 OS=Homo sapiens GN=KRT1 PE=1 SV=6
3.49	Keratin, type II cytoskeletal 1 OS=Homo sapiens GN=KRT1 PE=1 SV=6	2.92	Keratin, type II cytoskeletal 2 epidermal OS=Homo sapiens GN=KRT2 PE=1 SV=2
1.29	Keratin, type II cytoskeletal 2 epidermal OS=Homo sapiens GN=KRT2 PE=1 SV=2	2.7	Vimentin OS=Homo sapiens GN=VIM PE=1 SV=4
0.38	Keratin, type II cytoskeletal 6B OS=Homo sapiens GN=KRT6B PE=1 SV=5	0.43	Keratin, type II cytoskeletal 5 OS=Homo sapiens GN=KRT5 PE=1 SV=3
0.38	Keratin, type II cytoskeletal 6A OS=Homo sapiens GN=KRT6A PE=1 SV=3	5.32	Keratin, type I cytoskeletal 10 OS=Homo sapiens GN=KRT10 PE=1 SV=6
0.11	Keratin, type II cytoskeletal 5 OS=Homo sapiens GN=KRT5 PE=1 SV=3	0.45	Keratin, type I cytoskeletal 14 OS=Homo sapiens GN=KRT14 PE=1 SV=4
4.67	Keratin, type I cytoskeletal 10 OS=Homo sapiens GN=KRT10 PE=1 SV=6	1.81	T-cell receptor alpha joining 56 (Fragment) OS=Homo sapiens GN=TRAJ56 PE=4 SV=1
1.66	Keratin, type I cytoskeletal 9 OS=Homo sapiens GN=KRT9 PE=1 SV=3	2.62	Actin, cytoplasmic 1 OS=Homo sapiens GN=ACTB PE=1 SV=1
0.97	Keratin, type I cytoskeletal 14 OS=Homo sapiens GN=KRT14 PE=1 SV=4	2.35	Actin, cytoplasmic 2 OS=Homo sapiens GN=ACTG1 PE=1 SV=1
0.69	Keratin, type I cytoskeletal 17 OS=Homo sapiens GN=KRT17 PE=1 SV=2	0.46	Actin, aortic smooth muscle OS=Homo sapiens GN=ACTA2 PE=1 SV=1
5.01	Single-stranded DNA-binding protein, mitochondrial OS=Homo sapiens GN=SSBP1 PE=1 SV=1	0.37	Serum albumin OS=Homo sapiens GN=ALB PE=1 SV=1
2.35	Actin, cytoplasmic 1 OS=Homo sapiens GN=ACTB PE=1 SV=1	1.16	Keratin, type I cytoskeletal 9 OS=Homo sapiens GN=KRT9 PE=1 SV=3
0.13	Vimentin OS=Homo sapiens GN=VIM PE=1 SV=4	0.69	60 kDa heat shock protein, mitochondrial OS=Homo sapiens GN=HSPD1 PE=1 SV=2
3.6	40S ribosomal protein S5 (Fragment) OS=Homo sapiens GN=RPS5 PE=1 SV=1	0.12	Myosin-9 OS=Homo sapiens GN=MYH9 PE=1 SV=4
0.2	Microtubule-associated protein 1B OS=Homo sapiens GN=MAP1B PE=1 SV=2	0.1	Isoform 2 of Myosin-10 OS=Homo sapiens GN=MYH10
1.27	Elongation factor 1-alpha 1 OS=Homo sapiens GN=EEF1A1 PE=1 SV=1	0.72	Nascent polypeptide-associated complex subunit alpha (Fragment) OS=Homo sapiens GN=NACA PE=1 SV=1
1.07	Cofilin 1 (Non-muscle), isoform CRA_a OS=Homo sapiens GN=CFL1 PE=1 SV=1	0.1	Nestin OS=Homo sapiens GN=NES PE=1 SV=2
0.28	Leucine-rich PPR motif-containing protein, mitochondrial OS=Homo sapiens GN=LRPPRC PE=1 SV=3	0.71	Single-stranded DNA-binding protein, mitochondrial OS=Homo sapiens GN=SSBP1 PE=1 SV=1
2.74	Peptidyl-prolyl cis-trans isomerase B OS=Homo sapiens GN=PPIB PE=1 SV=2	0.23	40S ribosomal protein S12 OS=Homo sapiens GN=RPS12 PE=1 SV=3
2.35	Alpha-enolase OS=Homo sapiens GN=ENO1 PE=1 SV=2	0.7	Serpin H1 (Fragment) OS=Homo sapiens GN=SERPINH1 PE=1 SV=2
0.61	Stress-70 protein, mitochondrial OS=Homo sapiens GN=HSPA9 PE=1 SV=2	0.43	Transcription factor BTF3 homolog 4 OS=Homo sapiens GN=BTF3L4 PE=1 SV=1
1.91	ATP synthase subunit O, mitochondrial OS=Homo sapiens GN=ATP5O PE=1 SV=1	0.29	Reticulocalbin-2 OS=Homo sapiens GN=RCN2 PE=1 SV=1
0.73	Serpin H1 OS=Homo sapiens GN=SERPINH1 PE=1 SV=2	0.26	Prelamin-A/C OS=Homo sapiens GN=LMNA PE=1 SV=1
0.53	ATP synthase subunit alpha, mitochondrial OS=Homo sapiens GN=ATP5A1 PE=1 SV=1	0.12	THO complex subunit 4 OS=Homo sapiens GN=ALYREF PE=1 SV=1
0.61	ATP synthase subunit gamma, mitochondrial OS=Homo sapiens GN=ATP5C1 PE=1 SV=1	0.16	Nuclease-sensitive element-binding protein 1 (Fragment) OS=Homo sapiens GN=YBX1 PE=1 SV=1
0.94	Tubulin beta chain OS=Homo sapiens GN=TUBB PE=1 SV=1	0.3	Peptidyl-prolyl cis-trans isomerase B OS=Homo sapiens GN=PPIB PE=1 SV=2
0.97	Nascent polypeptide-associated complex subunit alpha (Fragment) OS=Homo sapiens GN=NACA PE=1 SV=1	0.1	Isoform 2 of Heat shock protein HSP 90-alpha OS=Homo sapiens GN=HSP90AA1
0.25	78 kDa glucose-regulated protein OS=Homo sapiens GN=HSPA5 PE=1 SV=2	0.12	Heat shock protein HSP 90-beta OS=Homo sapiens GN=HSP90AB1 PE=1 SV=4

Supplementary Figure 1: Hyperphosphorylative conditions. SH SY5Y cells differentiated with 30µM Retinoic acid for 10days were exposed to 30nM Okadaic acid for 24hrs. Under these hyperphosphorylative conditions tau is overexpressed and hyperphosphorylated. Figure shows the presence of ptau Ser202/Thr205 stained with antibody AT8 and p-tau 396 in SH SY5Y cells in hyperphosphorylative conditions..

Supplementary Figure 2: Apparent dissociation constants of Aptamer T1 and T3Target. SH SY5Y and Ren cells were grown in 96-well plates, Retinoic acid was used for differentiation. Okadaic acid for SH SY5Y and quinolinic acid for ReN cells was used to generate hyperphosphorylative conditions. Saturation binding curves were generated using Cy5 labelled T1 and T3 using a microplate fluorescent reader. The binding constant Kd apparent was calculated using the <mark>inbuilt modules</mark> in Graph Pad Prism. * *error bars smaller than symbol not visualized* *NAME THE MODULE*