Title: *C. elegans* DSB-3 Reveals Conservation and Divergence among Protein Complexes

Promoting Meiotic Double-Strand Breaks

Authors: Albert W. Hinman[1,2], Hsin-Yi Yeh[3], Baptiste Roelens[1], Kei Yamaya[1], Alexander

Woglar[1], Henri-Marc G. Bourbon[4], Peter Chi[3,5], Anne M. Villeneuve[1,2,*]

[1]Department of Genetics, Stanford University School of Medicine, Stanford, California, USA

[2]Department of Developmental Biology, Stanford University School of Medicine, Stanford,

California, USA

[3]Institute of Biochemical Sciences, National Taiwan University, Taipei, Taiwan

[4]Centre de Biologie Intégrative (CBI), Molecular, Cellular & Developmental biology unit (MCD),

Université Fédérale de Toulouse, Toulouse, France

[5]Institute of Biological Chemistry, Academia Sinica, Taipei, Taiwan

*Correspondence: annev@stanford.edu

Author Contributions:

AH: Designed Research, Performed Research, Analyzed Data, Wrote the Paper

HY: Performed Research, Analyzed Data

BR: Designed Research, Performed Research

KY: Performed Research

AW: Designed Research

HMB: Performed Research, Analyzed Data

PC: Designed Research, Analyzed Data

AV: Designed Research, Analyzed Data, Wrote the Paper

Abstract:

Meiotic recombination plays dual roles in the evolution and stable inheritance of genomes:

recombination promotes genetic diversity by reassorting variants, and it establishes temporary

connections between pairs of homologous chromosomes that ensure for their future

segregation. Meiotic recombination is initiated by generation of double-strand DNA breaks

(DSBs) by the conserved topoisomerase-like protein Spo11. Despite strong conservation of

Spo11 across eukaryotic kingdoms, auxiliary complexes that interact with Spo11 complexes to

promote DSB formation are poorly conserved. Here, we identify DSB-3 as a DSB-promoting

protein in the nematode *Caenorhabditis elegans*. Mutants lacking DSB-3 are proficient for

homolog pairing and synapsis but fail to form meiotic crossovers. Lack of crossovers in *dsb-3*

mutants reflects a requirement for DSB-3 in meiotic DSB formation. DSB-3 concentrates in

meiotic nuclei with timing similar to DSB-1 and DSB-2 (predicted homologs of yeast/mammalian

Rec114/REC114), and DSB-1, DSB-2, and DSB-3 are interdependent for this localization.

Bioinformatics analysis and interactions among the DSB proteins support the identity of DSB-3

as a homolog of MEI4 in conserved DSB-promoting complexes. This identification is reinforced

by colocalization of pairwise combinations of DSB-1, DSB-2, and DSB-3 foci in structured

illumination microscopy images of spread nuclei. However, unlike yeast Rec114, DSB-1 can

interact directly with SPO-11, and in contrast to mouse REC114 and MEI4, DSB-1, DSB-2 and

DSB-3 are not concentrated predominantly at meiotic chromosome axes. We speculate that

variations in the meiotic program that have co-evolved with distinct reproductive strategies in

diverse organisms may contribute to and/or enable diversification of essential components of

the meiotic machinery.

Significance Statement

Faithful inheritance of chromosomes during meiosis depends on the formation and repair of double-strand DNA breaks (DSBs), which are generated through the activity of a topoisomerase-like protein known as Spo11. Spo11 exhibits strong conservation throughout eukaryotes, presumably reflecting constraints imposed by its biochemical activity, but auxiliary proteins that collaborate with Spo11 to promote and regulate DSB formation are less well conserved. Here we investigate a cohort of proteins comprising a complex required for meiotic DSB formation in *Caenorhabditis elegans*, providing evidence for both conservation with and divergence from homologous complexes in other organisms. This work highlights the evolutionary malleability of protein complexes that serve essential, yet auxiliary, roles in fundamental biological processes that are central to reproduction.

Introduction

Meiotic recombination is important for two reasons. It promotes genetic diversity by

reassorting traits, and it is important for creating temporary attachments between pairs of

homologous chromosomes that are necessary for their future segregation at the meiosis I

division. Recombination is initiated by the programmed introduction of DNA double-strand

breaks (DSBs) (1). Some DSBs are repaired by a mechanism that leads to the formation of

crossovers (COs) between homolog pairs, and the remaining DSBs are repaired as

noncrossover products, thereby restoring genome integrity. Although DSBs are required for CO

formation, they may lead to genomic instability if they are not repaired or are repaired

erroneously. Thus, DSB formation in meiotic cells is governed by regulatory and surveillance

mechanisms that function to ensure that enough DSBs are created to guarantee a CO on each

homolog pair while limiting excess DSBs that may endanger the genome (2). Without

appropriate DSB formation and repair, COs may fail to form between homologs during meiotic

prophase, resulting in unattached homologs (univalents) that mis-segregate during the meiotic

divisions, leading to aneuploidy in the resulting progeny.

Meiotic DSB formation is catalyzed by Spo11, a topoisomerase-like protein homologous to

the catalytic A subunit of archaeal class VI topoisomerases that is well conserved across

eukaryotic kingdoms (3–6). The mechanism of DNA breakage involves formation of a covalent

linkage between Spo11 protein DNA, analogous to a key intermediate in the topisomerase

reaction (1). Despite identification of structural and mechanistic conservation between Spo11

and TopVIA more than 20 years ago, however, counterparts of the archaeal TopVIB subunit that

partner with Spo11 in "Spo11 core complexes" were not recognized until much later, reflecting

substantial divergence both from TopVIB and among their eukaryotic orthologs (7–9).

DSB formation also depends on multiple additional factors that play critical roles in

determining the location, timing, levels, and regulation of DSB formation (2). Several of these

auxiliary DSB-promoting factors, including Rec114, Mei4 and Mer2, were originally discovered

through genetic screens in *Saccharomyces cerevisiae* designed to identify genes required for

initiation of recombination (10, 11). In contrast to the high level of conservation observed for

Spo11, but similar to the other subunits of the Spo11 core complex, many auxiliary DSB protein

such Rec114, Mei4 and Mer2 are poorly conserved at the primary sequence level (1). Indeed,

high levels of sequence divergence had prevented identification of Rec114, Mei4 and Mer2

homologs outside of budding yeast until non-standard bioinformatics approaches were applied

(12, 13). Homologs of Rec114 and Mei4 that are required for meiotic recombination have now

been identified in several species, including *Mus musculus* (13–15), *Schizosaccharomyces*

pombe (11, 16, 17), and *Arabidopsis thaliana* (18, 19). Proteins that were independently

discovered based on roles in meiotic recombination in the ascomycete *Sordaria macrospora*

(Asy1) and in the nematode *Caenorhabditis elegans* (DSB-1 and DSB-2) were also

subsequently identified as putative Rec114 homologs (12, 20, 21), but Mei4 homologs were not

yet identified in these organisms.

Several studies have established that DSB auxiliary factors Rec114 and Mei4 work

closely together with each other and with Mer2 to promote meiotic DSB formation. Physical

interactions among these proteins and their orthologs have been demonstrated for several

organisms (13, 22–26), and coimmunoprecipitation experiments in *M.* musculus have further

confirmed that these proteins interact with one another *in vivo* in a meiotic context (15). Recent

biochemical analyses have shown that Rec114 and Mei4 together form individual complexes

with a stoichiometry of 2 Rec114 molecules for every 1 Mei4 molecule and have further

suggested that these complexes may self-assemble into large molecular condensates on

chromatin during meiotic progression (27). In both *S. cerevisiae* and *M. musculus,* all three

proteins have been reported to localize together in foci on meiotic prophase chromosomes (15,

22, 23, 26). Further, mouse REC114 and MEI4 and the Mer2 homolog IHO1 all localize

predominantly at the meiotic chromosome axis (15, 26), contributing to the idea that they act as

an intermediary between chromosome organization and DSB formation. Consistent with this

view, chromatin immunoprecipitation experiments in both *S. cerevisiae* and *S. pombe* have

shown that these proteins interact with both axis-enriched DNA sequences and with DSB sites

(25, 28–30). Additionally, *S. cerevisiae* Rec114 and Mei4 have been found to interact with the

Rec102 and Rec104 subunits that together comprise the TopVIB-like component of the Spo11

core complex (9, 23). Together these findings implicate Rec114-Mei4 in recruiting Spo11 to the

meiotic chromosome axis.

C. elegans DSB-1 and DSB-2, while clearly implicated in meiotic DSB formation, were

difficult to recognize as Rec114 homologs owing to high sequence divergence (12, 20, 21).

Further, *C. elegans* differs from yeast and mice regarding the relationships between DSB

formation and meiotic chromosome organization. Whereas DSB-dependent recombination

intermediates are required to trigger assembly of the synaptonemal complex (SC) between

homologous chromosomes in yeast and mice, *C. elegans* can achieve full synapsis between

aligned homologs even in the absence of DSB formation (6). Thus, there are substantial

differences in the cellular environments in which DSB-promoting complexes have evolved and

function in different organisms.

In our current work, we identify DSB-3 as a protein that partners with DSB-1 and DSB-2

to promote SPO-11-dependent meiotic DSB formation in *C. elegans*. We demonstrate a

requirement for DSB-3 in promoting the DSBs needed for CO formation, and we show that

DSB-3 becomes concentrated in germ cell nuclei during the time when DSBs are formed, in a

manner that is interdependent with DSB-1 and DSB-2. Through a combination of bioinformatics,

interaction data, and colocalization analyses, we identify DSB-3 as a likely Mei4 homolog and

establish DSB-1-DSB-2-DSB-3 as functional counterpart of the Rec114-Mei4 complex. Despite

homology and a shared role in promoting DSB formation, we uncover surprising differences

between the *C. elegans* DSB-1-DSB-2-DSB-3 and the REC114-MEI4 complexes observed in

mice, notably that *C. elegans* DSB-1, DSB-2 and DSB-3 are distributed broadly on chromatin

rather than becoming concentrated preferentially on chromosome axes. This work highlights the

evolutionary malleability of protein complexes that serve essential, yet auxiliary, roles in meiotic

recombination. Rapid diversification of such proteins may reflect a relaxation of constraints

enabled by changes in another aspect of the reproductive program, or alternatively, they may

reflect a capacity of alterations in such proteins to have an immediate impact on reproductive

success.

Results

Identification of *dsb-3* as a gene required for the formation of meiotic crossovers

The *dsb-3(me6ts)* allele was isolated in a genetic screen for mutants exhibiting a <u>h</u>igh

<u>i</u>ncidence of <u>m</u>ales among the progeny of self-fertilizing hermaphrodites, *i.e.,* the "Him"

phenotype, which is indicative of errors in segregation of X chromosomes during meiosis. *me6ts*

mutant hermaphrodites exhibit temperature-sensitive meiotic defects affecting both autosomes

and X chromosomes (Table 1, Figure 1A). Whereas inviable embryos and males (XO) are rare

among the self-progeny of wild-type hermaphrodites (XX), *dsb-3(me6ts)* mutant hermaphrodites

raised at the non-permissive temperature of 25°C produce 28% inviable embryos, and 17% of

their surviving progeny are males. Further, DAPI staining of chromosomes in oocytes at

diakinesis, the last stage of meiotic prophase, revealed a defect in chiasma formation in the

dsb-3(me6ts) mutant, reflecting an underlying defect in crossover formation (see below).

Whereas wild-type oocyte nuclei consistently exhibit 6 pairs of homologous chromosomes

connected by chiasmata (bivalents), oocyte nuclei in the *dsb-3(me6ts)* mutant exhibited a

mixture of bivalents and unattached (achiasmate) chromosomes (univalents), with the incidence

of univalents increasing with time.

Mapping and sequencing identified a missense mutation at genomic position IV:

7758710 (WS279) as the likely causal mutation responsible for the *dsb-3(me6ts)* mutant

phenotype (see Materials and Methods); this mutation results in a Leu165Phe substitution in the

previously uncharacterized protein C46A5.5. CRISPR/Cas9 genome editing was used to

introduce multiple stop codons early into the first exon of *C46A5.5*, thereby creating the null

allele *me115* (Figure 1B). *me115* fails to complement *dsb-3(me6ts)* (Figure 1A)*,* confirming the

identity of *C46A5.5* as *dsb-3*.

Analysis of the *dsb-3(me115)* null mutant indicates that the DSB-3 protein is required for

the formation of meiotic crossovers between all six pairs of homologous chromosomes. *dsb-*

3(me115) mutant hermaphrodites produced 99% inviable embryos, and 25% of their surviving

progeny were male, reflecting mis-segregation of autosomes and X chromosomes (Table 1).

Diakinesis oocytes of *dsb-3(me115)* mutant hermaphrodites exhibited an average of 11.6 ± 0.6

DAPI-stained bodies, indicating of a lack of chiasmata connecting all six homolog pairs (Figure

1A). A severe defect in crossover formation in the *dsb-3(me115)* mutant was also revealed

using GFP::COSA-1 as a cytological marker of crossover-designated sites in late pachytene

nuclei ((31); Figure 1C). Whereas 6 GFP::COSA-1 foci (1 per homolog pair) were consistently

observed in late pachytene nuclei of control worms, GFP::COSA-1 foci were absent from most

late pachytene nuclei in the *dsb-3(me115)* mutant.

Since pairing and assembly of the synaptonemal complex between homologous

chromosomes are prerequisites for the formation of crossovers during *C. elegans* meiosis, we

evaluated whether these features were impaired in the *dsb-3(me115)* mutant.   We assessed

pairing using fluorescence *in-situ* hybridization (FISH) for a 1 Mbp segment of Chromosome II

and immunostaining for HIM-8, a C2H2 zinc-finger DNA-binding protein that concentrates on the

Chromosome X pairing center (32, 33), demonstrating that *dsb-3(me115)* mutants are proficient

for homolog paring (Figure 1D). Further, immunostaining for the axial element protein HTP-3

and the synaptonemal complex central region protein SYP-2 (34, 35) revealed fully synapsed

chromosomes in early pachytene nuclei in the *dsb-3(me115)* mutant, indicating successful SC

assembly (Figure 1E). Together these data indicate that the DSB-3 protein is dispensable for

pairing and synapsis, and point to a role for this protein in the DNA events of recombination.

During *C. elegans* meiosis, failure to form crossover recombination intermediates

between one or more chromosome pairs prolongs the early pachytene stage of meiotic

prophase, reflecting the operation of a "crossover assurance" checkpoint (20, 21, 36).

Consistent with the observed deficit of interhomolog crossovers, *dsb-3(me115)* mutant gonads

display an extended zone of nuclei exhibiting phosphorylation of nuclear envelope protein SUN-

1, a marker of crossover assurance checkpoint activation (Supplemental Figure 1A).

DSB-3 is required for meiotic double-strand break formation

Meiotic recombination is initiated through the formation of DSBs by the conserved

topoisomerase-like protein SPO-11 (3, 6). Following formation, these DSBs are then processed

to enable the loading of the DNA strand-exchange protein RAD-51 (37, 38). RAD-51 foci thus

mark the sites of recombination intermediates that can be assayed as a proxy for successful

initiation of meiotic recombination (35, 39). We observed a strong decrease in the number of

RAD-51 foci in *dsb-3(me115)* mutants relative to wild type (Figure 2A), suggesting that fewer

DSBs are being created in these mutants or that there is a failure to load RAD-51 at DSB sites.

To determine whether fewer endogenous DSBs was the defect responsible for the observed

reduction in RAD-51 foci, we used γ-irradiation to introduce ectopic DSBs to test whether such

breaks are sufficient to restore crossover formation. Similar approaches were taken with other

DSB-defective mutants in *C. elegans,* such as *dsb-1*, *dsb-2*, and *spo-11* (6, 20, 21). Young adult

dsb-3(me115) and *dsb-3(me6ts)* hermaphrodites (alongside wild-type and *spo-11(me44)*

controls) were exposed to 5000 rad of γ-irradiation and subsequently assayed for crossover

formation through DAPI staining of chromosomes in oocyte nuclei. We observed a full rescue of

normal DAPI-body counts after irradiation (Figure 2B), suggesting that the *dsb-3* mutants are

specifically defective in DSB formation.

DSB-3 is concentrated in DSB-competent nuclei and is interdependent with DSB-1 and

DSB-2

Consistent with its role in promoting meiotic DSB formation, we find that DSB-3 localizes to

germ cell nuclei during the time when meiotic DSBs are formed. To assess DSB-3 localization in

the germ line, we generated a transgenic strain that expresses a DSB-3::GFP fusion protein in

the germ line in the *dsb-3(me15)* null mutant background (*meSi7[sun-1p::dsb-*

3::gfp::sun13'UTR] II; *dsb-3(me115)* IV); based on assessment of progeny viability and DAPI

bodies in diakinesis oocytes, we infer that this DSB-3::GFP fusion protein is largely functional in

promoting meiotic recombination (Table 1). Immunolocalization experiments in whole-mount

dissected gonads show that DSB-3::GFP becomes concentrated in germ cell nuclei within the

transition zone, soon after entry into meiotic prophase (Figure 3A). The DSB-3::GFP

immunofluorescence signal is strongest in early pachytene nuclei, then declines sharply in mid-

pachytene, albeit with a few outlier nuclei in the late pachytene region of the gonad retaining a

strong DSB-3 signal. This pattern of appearance and disappearance of DSB-3 from germ cell

nuclei is similar to the patterns observed for the double-strand break promoting proteins DSB-1

and DSB-2 ((20, 21) and Figure 3A), and corresponds to the timing when nuclei are competent

for meiotic DSB formation.

The DSB-3, DSB-2, and DSB-1 proteins are not only abundant in the same nuclei during

early meiotic prophase, but they are also interdependent for this immunolocalization. Previous

studies had demonstrated interdependence for DSB-1 and DSB-2 (20, 21), with loss of the

DSB-2 immunofluorescence signal in a *dsb-1* null mutant (which lacks meiotic DSB-promoting

activity) and reduction of DSB-1 immunofluorescence signal in a *dsb-2* null mutant (which

retains a low residual level of DSB-promoting activity). Similarly, we found that DSB-3::GFP

immunofluorescence signal was abolished in *dsb-1* null mutant germ lines (Figure 3B).

Likewise, DSB-3::GFP immunofluorescence signal was reduced in *dsb-2* null mutant germ lines

and was restricted to a few rows of nuclei in the transition zone and very early pachytene

regions of the gonad (Figure 3B). Conversely, DSB-1 and DSB-2 immunostaining were lost in

dsb-3(me115) mutant germ lines (Figure 3C). Collectively, these data indicate that DSB-3, DSB-

2, and DSB-1 are interdependent for proper localization to germ cell nuclei, indicating that they

function together in promoting meiotic DSB formation.

Together, our data demonstrating 1) a similar requirement in promoting DSB formation, 2)

concentration in the same nuclei, and 3) interdependence for localization and/or abundance in

meiotic nuclei are all consistent with DSB-3 functioning in a protein complex together with DSB-

1 and DSB-2 to promote the formation of SPO-11-dependent meiotic DSBs.

Evidence that DSB-1, DSB-2 and DSB-3 form a complex homologous to the yeast and

mammalian Rec114-Mei4 complexes

Although the initial PSI-BLAST searches conducted for DSB-1 and DSB-2 had not

identified homologs outside of *Caenorhabditis* (20, 21), DSB-1 and DSB-2 were subsequently

identified as likely distant homologs of the Rec114 meiotic DSB-promoting proteins from fungi

and mammals (12). This identification was enabled using an approach involving PSI-BLAST

searches initiated using sequence alignments, in combination with scanning for patterns of

similarity in predicted secondary structure, to identify short signature motifs (SSMs) in poorly

conserved proteins (40). We obtained additional support for the assignment of DSB-1 and DSB-

2 as Rec114 homologs using the Phyre2 structure prediction server (41), which identified

medium confidence (#6 hit, 40.3%) and low confidence (#19 hit, 11.9%) alignments between the

N-terminal domains of DSB-2 and DSB-1 and the N-terminal domain (where the identified SSMs

are located) of the solved structure of mouse Rec114 (42). We therefore used an alignment

driven PSI-BLAST approach similar to that described above to identify DSB-3 as a putative

homolog of Mei4 (see Materials and Methods, Figure 4A, Supplemental Figure 2), which is

required for meiotic DSB formation in yeast and mice and forms a complex with Rec114 (13–15,

22, 24, 43). Based on local amino acid composition and relative position in the protein

sequence, three of the 6 SSMs previously identified in Mei4 homologs from diverse species

(SSMs #1, 4 and 6) are well supported in the *Caenorhabditis* DSB-3 orthologs, while the other

three SSM are less well conserved.

To complement these *in silico* analyses, we used yeast two-hybrid (Y2H) assays to establish a

network of interactions among the DSB-1, DSB-2, and DSB-3 proteins and SPO-11, the protein

that catalyzes DSB formation (Figure 4B, Supplemental Figure 2). Y2H interactions were

detected between DSB-1 and DSB-2 and between DSB-1 and DSB-3, consistent with an ability

of these proteins to form complexes. Homotypic interactions were also detected both for DSB-1

and for DSB-2. In addition to the interactions detected among the putative Rec114 and Mei4

homologs, DSB-1 also interacted with SPO-11 in the Y2H assay.

We note that a truncated version of DSB-1 lacking the N-terminal 33 amino acids loses the

ability to interact with SPO-11 but retains its ability to associate with DSB-2 and DSB-3. This

suggests that the interactions between DSB-1 and SPO-11 and the interactions between DSB-1

and DSB-2 or DSB-3 may be mediated, at least in part, by different parts of the DSB-1 protein.

DSB-3, DSB-2, and DSB-1 colocalize in meiotic nuclei

To complement our genetic, bioinformatic, and Y2H evidence that DSB-3, DSB-2, and

DSB-1 function together as components of conserved protein complexes to promote DSB

formation, we investigated their colocalization using Structured Illumination Microscopy (SIM) on

spread preparations of meiotic nuclei (Figure 5). For most of these analyses, we used a

moderate nuclear spreading protocol (44), coupled with SIM imaging to provide improved spatial

resolution below the limits of standard light microscopy (45). This approach enabled detection of

these proteins as chromosome-associated foci. To facilitate co-staining of protein pairs for these

colocalization analyses, we used CRISPR/Cas9 gene editing to create strains expressing HA or

FLAG tagged versions of the DSB proteins from the endogenous loci (Table 1), and we

detected the proteins using indirect immunofluorescence.

We used the image analysis pipeline outlined in Figure 5A (described in more detail in

Supplemental Figure 3) to identify DSB protein foci and to conduct object-based colocalization

analyses to assess the degree of colocalization detected for pairwise combinations of the

imaged DSB proteins within individual nuclei. As negative controls, we generated virtual nuclei

in which the second channel in each combination was rotated by 90° in XY, resulting in virtual

composite images in which DSB protein foci are modified in location, but numbers, sizes, and

intensity distributions of foci remain unaltered. Collectively, our analyses indicate that the DSB-

3, DSB-2, and DSB-1 proteins strongly colocalize with each other in meiotic prophase nuclei.

Analysis of all three pairwise combinations of DSB-1, DSB-2, and DSB-3 foci are

presented in Figure 5B-D and Supplemental Figure 4. For all three pairs, numbers of foci for the

two channels detected in each nucleus were strongly correlated, consistent with expectations

for components of the same protein complex. Further, substantial colocalization was observed

for each pair. For example, we found that 45 ± 5% of DSB-2 foci colocalized with DSB-3::GFP

foci, and conversely, that 37 ± 7% of DSB-3::GFP foci colocalized with DSB-2 foci. In contrast,

negative control coincidental colocalization values were 10 ± 3% and 8 ± 3%, respectively.

Similarly, 52 ± 8% of 3xFLAG::DSB-3 foci colocalized with 3xHA::DSB-1 foci, and conversely

55% ± 5% of 3xHA::DSB-1 foci colocalized with 3xFLAG::DSB-3 foci. Likewise, 43 ± 11% of

3xHA::DSB-1 foci colocalized with DSB-2 foci, and conversely 41 ± 9% of DSB-2 foci

colocalized with 3xHA::DSB-1 foci.

Although substantial colocalization was observed for all pairwise combination of DSB-1,

DSB-2 and DSB-3 foci, the fraction of colocalization may seem lower than might be anticipated

for proteins comprising the same protein complex. We note, however, that incomplete

colocalization has been similarly observed for the Rec114 and Mei4 proteins in both budding

yeast and mouse meiocytes (22, 26). One possible explanation is that only a subset of these

protein molecules occur together in complexes, while other molecules exist separately within the

nucleus; however, this explanation is not easily reconciled with the observed interdependence

among these proteins. Another possibility is that the observed degree of colocalization reflects

limitations on our ability to detect all of the DSB-1, DSB-2 and DSB-3 target molecules that are

present. *e.g.* because of isoforms that lack epitopes or because the complexes and/or their

components may be organized in a manner that makes some epitopes inaccessible to detection

reagents.

This latter possibility is supported by data from an experiment in which we assessed

colocalization for fluorescent foci representing separate epitopes on the same protein,

3xHA::DSB-2, expressed from the endogenous *dsb-2* locus. Specifically, we used a mouse

monoclonal antibody (mAB) against the HA epitope and rabbit polyclonal (pAB) antibodies

raised against the C-terminal 100 amino acids of the DSB-2 protein. The numbers of foci for the

two channels detected in each nucleus were strongly correlated (Figure 5C), as expected for

foci representing the same target molecule. However, colocalization was again incomplete, in

both directions: 44 ± 5% of DSB-2 pAB foci colocalized with HA mAB foci, and conversely, 48 ±

2% of HA mAB foci colocalized with DSB-2 pAB foci. This incomplete colocalization of HA mAB

and DSB-2 pAB fluorescent signals supports the conclusion that a subset of epitopes on DSB-2

proteins present in the nucleus were not detected in these experiments.

339   For the DSB-2 - DSB-3::GFP combination, we also conducted a colocalization analysis

340 on "super-spread" nuclei, in which chromosomes were dispersed over an area 6-10 times larger

341 than that of an unperturbed nucleus (46) (Materials and Methods and Supplemental Figure S5).

342 The numbers of foci detected using this approach were 3-5 times higher than the numbers

343 observed in our analysis of partial spreads, but a similar degree of colocalization was detected:

344 55 ± 16% of DSB-2 foci colocalized with DSB-3::GFP foci, and conversely, 44 ± 10% of DSB-

345 3::GFP foci colocalized with DSB-2 foci. The observation of larger numbers of foci with a

346 comparable degree of colocalization suggests the possibility that groups of DSB protein

347 complexes may be split into smaller cohorts by the super-spread procedure.

349 The Presence and Colocalization of DSB-3, DSB-2, and DSB-1 Is Not Confined to the

350 Meiotic Chromosomal Axis

351   Previous chromatin immunoprecipitation experiments in *S. cerevisiae* have shown that

352 Mei4 and Rec114 are enriched at DNA sequences that are also enriched for meiosis-specific

353 axis proteins Hop1 and Red1 (28, 30). Moreover, *M. musculus* Mei4 and Rec114 were found to

354 colocalize cytologically on the axes of meiotic prophase chromosomes (14, 15, 26). Based on

355 these observations, it has been proposed that the Rec114-Mei4 complex primarily functions at

356 the chromosome axes. Strikingly, DSB-1, DSB-2, and DSB-3 foci are not enriched at the

357 chromosome axis during *C. elegans* meiosis. Rather, we find that most foci are detected away

358 from the axis, in the associated chromatin loops (Figure 6A). To quantify axis association, we

359 first segmented images by creating axis masks for each nucleus that corresponded to the pixels

360 containing the immunofluorescence signal derived from the axis protein HTP-3 (Supplemental

361 Figure 3B), then for each DSB protein tested, we identified the subset of foci, termed "axis-

362 associated foci" for which some or all of the pixels coincided with the axis mask. This approach

363 indicated that only 15-30% of DSB protein foci detected in our analyses overlapped with

chromosome axis signal, indicating that on spread chromosomes from meiotic nuclei. the

preponderance of DSB protein foci detected were not associated with the meiotic chromosome

axes.

We also assessed whether DSB protein foci associated with the chromosome axis might

exhibit a higher degree of colocalization with their DSB protein partners relative to the level of

colocalization observed for the full set of foci within the nucleus (Figure 6B). However, this

analysis did not reveal any consistent enrichment of colocalization for DSB protein foci that were

linked to the chromosome axis. Thus, while components of the meiotic chromosome axis do

have roles in promoting and regulating SPO-11 dependent DSB forming activity in *C. elegans*

meiosis(47–49)*,* these roles do not appear to be mediated by concentrating DSB-1-DSB-2-DSB-

3 complexes in close proximity to the chromosome axis.

Discussion

Initiation of meiotic recombination by programmed DSB formation is an ancient and

conserved feature of the meiotic program that predates divergence of plants, animals and fungi.

Thus, it is not surprising that Spo11, the protein directly responsible for catalyzing DSB

formation, is strongly conserved across kingdoms, given constraints imposed by its requirement

to interact with and perform chemistry on DNA. However, many additional proteins required for

DSB formation had been identified in the yeast system, but plant and metazoan homologs of

these auxiliary DSB proteins had long eluded detection by standard BLAST analyses. The

barrier to detection of homologs outside fungi was eventually breached using secondary

structure prediction coupled to MAFFT alignment and phylogenomically-oriented PSI-BLAST

searches (13), which identified characteristic SSMs for putative Mei4 and Rec114 homologs;

moreover, the veracity of these predictions was borne out by demonstration of meiotic roles in

mouse mutants (13–15).

Auxiliary proteins involved in DSB formation during *C. elegans* meiosis were identified

independently based on analysis of meiotic mutants ((20, 21), this work). However, recognition

of these proteins as distant homologs of conserved DSB-promoting factors came later, after

their functional importance in DSB formation was already established (12). Identification of DSB-

1 and DSB-2 as Rec114 homologs was further solidified by alignments of the predicted

structure of DSB-1 and DSB-2 with the solved structure of mouse REC114 ((42); this work).

Likewise, our initial identification of *C. elegans* DSB-3 as a factor important for meiotic DSB

formation was similarly based on functional data. The identity of DSB-3 as a putative Mei4/MEI4

ortholog was derived computationally from alignments and collinearity of SSMs among

metazoan homologs, and this identification was reinforced by demonstration of Y2H

interactions, colocalization and interdependence with Rec114 homologs DSB-1 and DSB-2.

Thus, despite a high degree of divergence at the amino acid sequence level, our data

collectively support the conclusion that DSB-1, DSB-2 and DSB-3 together form complexes that

are the functional counterpart of Rec114-Mei4 complexes.

Having established conservation among the auxiliary complexes that promote the DSB-

forming activity of Spo11, our analyses also reveal interesting differences. First, whereas yeast

and mice each have only a single Rec114/REC114 ortholog, nematodes in the Caenorhabditis

genus each have 2 paralogs, indicating duplication and divergence in the parental lineage.

DSB-1 and DSB-2 are neither identical to nor functionally interchangeable with each other, as

DSB formation is strongly reduced in *dsb-2* mutants and eliminated in *dsb-1* mutants.

Interestingly, recent biochemical analyses indicate a 2 Rec114 : 1 Mei4 stoichiometry of the

yeast complex (27). The interdependence of DSB-1 and DSB-2 for nuclear enrichment, in

combination with the colocalization observed in chromosome spreads (this work) suggest that

the *C. elegans* complexes may typically contain one DSB-1 subunit and one DSB-2 subunit

(rather than two identical subunits). However, there is low residual DSB-promoting activity

present in *dsb-2* null mutants, suggesting that complexes with two DSB-1 subunits may form

and be partially functional when DSB-2 is absent. While the data do support functional

diversification of the *C. elegans* Rec114 paralogs, however, how this diversification came about

and/or how and why it persisted remain unknown.

A second apparent distinction between *C. elegans* DSB-1, DSB-2 and DSB-3 and their

mouse counterparts is the observed relationship to meiotic chromosome axes. Mouse REC114

and MEI4 are reported to localize predominantly at chromosome axes in spread preparations of

meiotic prophase chromosomes, in a manner mediated by Mer2 homolog IHO1 (15, 26) . This

association with chromosome axes led to the proposal that a major role for the Rec114 – Mei4

complex is to recruit the Spo11 core complex to the axis to activate its DSB-promoting activity

specifically in close proximity to the axis, where DSB repair predominantly occurs. In *C.*

elegans, meiosis specific HORMAD proteins HTP-1 and HTP-3, which are major building blocks

of the chromosome axis, are implicated in playing important roles in promoting and regulating

meiotic DSB formation (47–49). HTP-3 is strictly required for DSB formation, and HTP-1 is

required for normal levels of DSB activity. However, despite this requirement for axis

components in DSB formation, our analysis here indicates that DSB-1-DSB-2-DSB-3 complexes

are not preferentially enriched adjacent to axes. This may reflect lack of an apparent

Caenorhabditis ortholog of IHO1, which is recruited to mouse meiotic chromosomes through a

direct interaction with HORMAD1.

Given the observed lack of enrichment of DSB-1, DSB-2 and DSB-3 at the axes, the

role(s) of *C. elegans* meiotic HORMAD proteins in promoting DSB formation may not be strictly

limited to recruiting the homologs of Rec114 and Mei4 to chromosomes. One possibility is that

the *C. elegans* HORMADs might be involved in activating the subset of DSB-1-DSB-2-DSB-3

complexes that do occur in close proximity to the axis. Alternatively, the role(s) of the *C. elegans*

axis proteins might be indirect, *e.g.* assembly of the axis might potentially signal successful

formation of chromosome structure that is proficient for meiotic DSB repair, thereby licensing

the nucleus that it is safe to proceed with DSB formation. The possibility of this signaling

scenario is strengthened by prior work demonstrating a role for *C. elegans* HORMAD proteins in

a signaling process that sustains activity of protein kinase CHK-2, a master regulator of multiple

processes during meiosis, including nuclear enrichment of DSB-1 and DSB-2 (20, 21, 36, 50).

It is possible that the observed difference between mice and *C. elegans* regarding axis

enrichment of Rec114-Mei4 complexes may be related to differences in spatial organization of

recombination events in the genome and/or in coupling between DSB repair and homolog

recognition. Meiotic recombination in mice occurs predominantly within 1-2 kb "hotspot' regions,

separated by larger (50-100kb) cold regions where the probability of recombination is very low.

A similar "local hotspot" distribution of recombination events was not observed in *C. elegans* (for

the portion of the genome studied) (51, 52), suggesting that different constraints are operating

to dictate where DSBs may form. Further, there is substantial variation among organisms

regarding their relative dependence on different mechanisms that promote pairwise alignment

and synapsis between homologous chromosomes. In mouse meiosis, formation of early

SPO11-dependent DSB repair intermediates appears to be the main mechanism of homology

verification, required to trigger SC assembly and constraining it to occur strictly between aligned

homologous chromosomes (53, 54). In contrast, in *C. elegans*, local *cis*-acting chromosomal

domains known as pairing centers play a primary role in homolog recognition, and these are

capable of promoting largely successful pairwise synapsis between homologs even in the

absence of recombination (55). We speculate that differences in the constraints governing the

genomic locations of DSBs and/or differences in dependence on DSBs for homology verification

may have either contributed to, or been enabled by, diversification of meiotic DSB auxiliary

protein complexes. We note that high divergence among essential components of key biological

processes is a hallmark not only of the meiotic program, but of reproduction more generally (1,

56, 57). This likely reflects multiple underlying factors, including the potential for changes in

such proteins to have an immediate impact on processes that directly affect fitness by

modulating reproductive success.

A key question regarding the role(s) of the Rec114-Mei4 (or DSB-1-DSB-2-DSB-3)

complexes is how exactly they are functioning to promote Spo11 activity. It was recently

proposed that Rec114-Mei4 complexes function by forming large DNA-dependent biomolecular

condensates that promote DSB activity by causing a high local concentration of Spo11 core

complexes at presumptive DSB sites held adjacent to the chromosome axis (27). This model

was proposed based on: 1) a large segment of the Rec114 protein exhibiting a high probability

of disorder, 2) the ability of purified Rec114-Mei4 complexes to promote formation of DNA-

dependent condensates *in vitro*, and 3) an ability of Rec114-Mei4 complexes to interact with

and recruit the Spo11 core complex. While evaluating potential for *in vitro* condensation is

outside the scope of the current study, we note that some of the above attributes may be shared

with the *C. elegans* DSB-1-DSB-2-DSB-3 complex. First, based on the Y2H data, the DSB-1-

DSB-2-DSB-3 complex is expected to be able to interact with SPO-11 complexes. However, in

yeast, interactions occur between Rec114 and the Rec102 and Rec104 components (which

together correspond to Top6BL) of the Spo11 core complex, whereas in *C. elegans,* a Top6BL

homolog has not yet been identified, and DSB-1 can interact directly with SPO-11 itself in the

Y2H assay. Second, the DSB-1 and DSB-2 proteins have long segments with predicted protein

disorder scores that hover around 0.5 and include short segments scoring >0.5, leaving it

ambiguous whether these might represent *bona fide* disordered regions. Third, the observation

that higher numbers of DSB-2 and DSB-3 foci are detected in super-spread nuclei than in

moderately-spread nuclei raises the possibility that these proteins might normally occur in larger

groups within intact nuclei, potentially analogous to the condensates proposed to occur during

yeast meiosis. In either system, future investigations aiming to test predictions of the

condensation model will need to address the challenge of visualizing complex dynamic behavior

in vivo.

Materials and Methods

<u>*C. elegans* strains</u>

Strains were cultured at 20℃ using standard nematode growth conditions (58) unless

otherwise noted. Strains used in this study:

AV28 *dsb-3(me6ts)* IV

AV776 *spo-11(me44) IV / nT1[qIs51]* (IV;V)

AV818 *meIs8[gfp::cosa-1]* II; *cosa-1(tm3298)* III

AV913 *dsb-3(me6ts)* IV

AV958 *dsb-3(me6ts) dpy-20(e1282)* IV

AV994 *dpy-3(e184) dsb-3(me6ts)* IV

AV995 *dsb-3(me115) IV / nT1 (*IV;V)

* AV1029 *meSi7 [sun1p::dsb-3::gfp::sun-1 3'UTR] II; dsb-3(me115) IV*

AV1045 *meSi7 II; dsb-3(me115) dsb-1(we11) / nT1 IV*

AV1081 *meSi7 dsb-2(me96) / mnC1 II; dsb-3(me115) IV*

AV1095 *dsb-3(me115) / tmC5 [F36H1.3(tmIs1220)] IV*

^ AV1102 *dsb-1(me124[3xha::dsb-1]) dsb-3(me125[3xflag::dsb-3]) IV*

^ AV1115 *dsb-2(me132)[3xha::dsb-2] II*

AV1132 *meIs8[gfp::cosa-1]* II*; cosa-1(tm3298)* III*; dsb-3(me115)* IV */ nT1* (IV;V)

Bristol N2 Wild type

* The transgene allowing expression of the DSB-3::GFP fusion protein was obtained using the

Mos Single Copy Insertion strategy (MosSCI, (59)) using the *ttTi5605* insertion on chromosome

II as a landing site. The donor plasmid, pBR253, was obtained by assembling fragments

carrying the upstream promoter region of the *sun-1* gene, the *sun-1* downstream 3'UTR region,

and the genomic sequence of *dsb-3* (coding exons and introns), together with a DNA fragment

containing a version of GFP optimized for germline expression (60), into pBR49, a derivative of

pCFJ350 modified to enable type IIs restriction/ligation cloning (61). The genomic fragments

were obtained by PCR amplification of wild-type genomic DNA using the following primer pairs.

The primers for the *sun-1* promoter were oBR840

(cgtcgatgcacaatccGGTCTCaCCTGatttccagatttcatcgtcggtttt) and oBR841

(agtggaatgtcagGGTCTCaCATaccgagtagatctggaagtttag). The primers for *dsb-3* CDS were:

oBR836 (cgtcgatgcacaatccGGTCTCaTATGATCGAAATTACCGATGATGAGG)

and oBR837 (agtggaatgtcagGGTCTCaCTCCATTGCTATATCTCTGTTGATTATCTAAAAAC)

The primers for the *sun-1* 3'UTR were oBR842

(cgtcgatgcacaatccGGTCTCaTAAAaaacgccgtattattgttcctgc) and oBR843

(agtggaatgtcagGGTCTCaGTCAttagtaagttaaagctaaagttagcag). The GFP fragment was obtained

by PCR amplification of pCFJ1848 (60), using oBR406

(cgatgcacaatccGGTCTCaGGAGGTGGATCATCCTCCACATCATCCT) and oBR407

(agtggaatgtcagGGTCTCaTTTATGGGGAAGTACCGGATGACG). Correct assembly of all

fragments within the donor plasmids was verified by sequencing.

^ In order to perform pairwise colocalization experiments between DSB-1, DSB-2 and DSB-3,

we created strains expressing endogenously-tagged versions of these proteins so each pair

could be detected using compatible primary antibodies generated in different host organisms.

For these strains, we used direct injection of Cas9 protein (PNAbio) complexed with single-

guide RNA (sgRNA) (Dharmacon) using the protocol of (62). CRISPR targeting (crRNA)

sequences were designed using Benchling (https:// benchling.com/). Small single-stranded

oligonucleotides (< 200 bp) were purchased (Integrated DNA Technologies) and used as the

repair templates to generate the various tags and nonsense alleles. N2 worms (P0) were

injected with the mix together with sgRNA and repair template for the *dpy-10* co-CRISPR

marker (63). Rol F1s (carrying *dpy-10(Rol)* marker) were singled out, and a subset of F2

progeny was fixed and stained with DAPI (see below) to assess the phenotype of diakinesis

nuclei for null alleles. From plates containing worms exhibiting univalents at diakinesis, the new

mutations were recovered from siblings of the imaged worms and balanced by *nT1* IV or *tmc5*

IV. Tagged alleles were confirmed by immunofluorescence staining (see below). All edits were

confirmed by Sanger sequencing of PCR fragments amplified using primers designed to detect

the edit event. The crRNAs used, description of the edits, and PCR sequencing primers used

are included in Supplemental Table 1.

<u>Isolation, mapping, and genomic identification of the *dsb-3(me6ts)* mutation</u>

dsb-3(me6ts) was isolated in a genetic screen for meiotic mutants exhibiting a high incidence of

males as described in (64). After backcrossing (four times) to generate the AV913 strain,

homozygous *me6ts* worms were subjected to whole-genome sequencing. DNA was extracted

from ~8 60mm confluent plates of N2 and AV913 gravid adult worms; worms were rinsed twice

in M9 and resuspended in 10 mM EDTA and 0.1 M NaCl. Worms were then: pelleted; flash

frozen in liquid nitrogen; resuspended in 450 µL of lysis buffer containing 0.1 M Tris, pH 8.5, 0.1

M NaCl, 50 mM EDTA, and 1% SDS plus 40 µL of 10 mg/mL proteinase K in TE (10 mM Tris, 1

mM EDTA), pH 7.4; vortexed; and incubated at 62°C for 45 min. Two successive phenol-

chloroform extractions were performed using the Phase Lock gel tubes from Invitrogen, and

DNA was precipitated with 1 mL of 100% ethanol plus 40 µL of saturated NH_4Ac (5 M) and 1 µL

of 20 mg/mL GlycoBlue. The DNA pellet was washed with 70% ethanol, air dried, and

resuspended in 50 μL of TE, pH 7.4. Paired end libraries were prepared using the Nextera

technology (Illumina), and sequencing was performed on an MiSeq sequencer (2 × 75 bp)

through the Stanford Functional Genomics Facility. To analyze the genomic data, we used an

analysis pipeline adapted from GATK's recommended best practices (65–67). Reads were

mapped to *C. elegans* reference genome (WBcel 235.82) using the Bowtie 2 software (68).

Variant calling was performed using Haplotype Caller software from GATK, and lists from

AV913 and N2 were compared to eliminate non-causal variants. The predicted effects of

variants specific to AV913 were then annotated using SnpEff (69).

Initial genetic mapping experiments had placed *dsb-3(me6ts)* within 2 cM of *unc-5*,

located at 1.78 cM on chromosome IV; the above sequence analysis identified several

candidate mutations within this region. Additional mapping crosses located *dsb-3(me6ts)* to the

left of *dpy-20* (at 5.22 cM) and near or to the left of *unc-24* (at 3.51 cM). Further, we found that

eDf18 (which deletes the region between 3.7- 4.19 cM) complements *dsb-3(me6ts)*. Together,

these experiments identified a G -> A transition at genomic position IV: 7758710 (WS279), in

the second coding exon of the uncharacterized gene *C46A5.5*, as the likely causal mutation

responsible for the *dsb-3(me6ts)* mutant phenotype.

<u>DAPI staining of oocyte chromosomes and Irradiation Assay</u>

Numbers of DNA bodies present in diakinesis oocytes were assessed in intact adult

hermaphrodites of the indicated ages, raised at the indicated temperatures, fixed in ethanol and

stained with 4′,6-diamidino-2-phenylindole (DAPI) as in (70). This method underestimates the

frequency of achiasmate chromosomes, as some univalents lie too close to each other to be

resolved unambiguously.

To test for rescue of bivalent formation by exogenously derived DSBs, worms were exposed to

5,000 rad (50 Gy) of γ-irradiation using a Cs-137 source at 20 h post-L4 stage. Worms were

fixed and stained at 18–20 h post-irradiation, and numbers of DAPI bodies were counted in

oocyte nuclei in the -1 to -3 positions.

<u>Bioinformatic identification of homology between DSB-3 and Mei4</u>

PSI Blast searches using the MPI BLAST server (71) , initiated using an alignment of DSB-3

homologs from diverse roundworm species as the query, identified a putative *Brugia malayi*

DSB-3 homolog. A subsequent round of PSI-BLAST searches, initiated using an alignment with

the putative *B. malayi* homolog as the header sequence and initially focusing on the N-terminal

portion of the protein, led to retrieval of plant and animal Mei4 homologs. Similarity in protein

lengths and patterns of predicted secondary structure were prioritized over E-value

considerations in selection of proteins chosen for the multiple sequence alignment presented in

Supplemental Figure 2, which was generated using MAFFT Version 7.0 with gap opening

penalty parameter set to 2.0 and offset value parament set to 0.125.

<u>Yeast Two-Hybrid Analysis</u>

Full-length DSB-1, DSB-2, DSB-3, and N-terminally truncated SPO-11 (SPO-11Δ1-47),

DSB-1 (DSB-1Δ1-33) ORFs were individually cloned into the *Bam*HI and *Pst*I sites of pBridge,

and the *Bam*HI and *Xho*I sites of pGADT7 (Clontech) to generate fusion proteins with the N-

terminal Gal4 DNA-binding domain (Gal4BD) or activation domain (Gal4AD). The PJ69-4A

yeast strain (*MATa trp1-901 leu2-3,112 ura3-52 his3-200 gal4Δ gal8Δ GAL2-ADE2*

LYS2::GAL1-HIS3 met2::GAL7-lacZ) was co-transformed with the indicated pairs of constructs

encoding Gal4BD and Gal4AD fusion proteins (and/or empty vector negative controls).

Transformed cells expressing Gal4BD and Gal4AD fusion proteins were selected in SD-Leu⁻Trp⁻

, a drop-out medium without leucine and tryptophan. Protein interactions were assayed by

growing transformed cells for 5 days at 30°C on selective media lacking leucine, tryptophan,

histidine, and adenine (SD-Leu⁻Trp⁻His⁻Ade⁻). Three independent repeats of each transformation

were performed for all pairwise combinations. The full-length SPO-11 ORF was excluded from

analysis of combinations as it exhibited autoactivation in negative control experiments.

<u>Immunofluorescence Methods</u>

The following primary antibodies were used: mouse anti-HA (1:1000, Covance 16B12

clone), rabbit anti-FLAG (1:5000, Sigma Aldrich), rabbit anti-DSB-2 (1:5.000, (20)), guinea pig

anti-DSB-1 (1:500, (21)), rabbit anti-GFP (1:200, (31)), guinea pig anti-HIM-8 (1:500, (32)),

chicken anti-HTP-3 (1:400, (33)), rabbit anti-SYP-2 (1:200, (35)), rat anti-RAD-51 (1:500, (20)),

guinea pig anti-SUN-1 S24pi (1:700, (72)), chicken anti-GFP (1:500, (A01694, Genscript)).

Secondary antibodies were Alexa Fluor 488, 555 and 647-conjugated goat antibodies directed

against the appropriate species (1:400, Life Technologies).

For immunofluorescence experiments involving whole mount gonads, dissection of

gonads, fixation, immuno-staining and DAPI counterstaining were performed as in (48) .

For experiments involving nuclear spreads, spreading was performed as in (73). The

gonads of 20–100 adult worms were dissected in 10 µL Dissection solution (75% v/v Hank's

Balanced Salt Solution [HBSS, Life Technology, 24020-117] with 0.1% v/v Tween-20) on an

ethanol-washed plain slide. 50 µL of spreading solution (32 µL of Fixative [4% w/v

Paraformaldehyde and 3.2%–3.6% w/v Sucrose in water], 16 µL of Lipsol solution [1% v/v in

water], 2 µL of Sarcosyl solution [1% w/v of Sarcosyl in water]) were added, and gonads were

immediately distributed over the whole slide using a pipette tip. Slides were then left to dry at

room temperature overnight, washed for 20 minutes in methanol at -20°C and rehydrated by

washing 3 times for 5 minutes in PBS-T. A 20-minute blocking in 1% w/v BSA in PBS-T at room

temperature was followed by overnight incubation with primary antibodies at room temperature

(antibodies diluted in: 1% w/v BSA in PBS-T). Slides were washed 3 times for 5 minutes in PBS-

T before secondary antibody incubation for 2 hours at room temperature. After PBS-T washes,

the samples were mounted in Vectashield (Vector).

To dissect large quantities of *C. elegans* gonads for spreads, we employed an

alternative method for disrupting worms, using a 125V ~ 60Hz drill capable of achieving 1,600

rotations per minute. Briefly, we synchronized worms by using the standard bleaching protocol

(74) and allowed worms grow to adulthood (L4 + 24 hours). The worms were then washed with

dissection solution into a 1.7 mL Eppendorf tube suspended in an ethanol ice bath. The worms

were then disrupted using a 1/64 inch bit on the drill with its maximum power by angling the drill

bit against the Eppendorf tube wall. 3 µL aliquots were taken from the tube every 20 seconds

and monitored microscopically until most of the gonads had been extruded from the worms

during the drill-induced disruption.

<u>FISH experiments</u>

Barcoded Oligopaint probes targeting a 1 Mb segment of chromosome II (genomic

coordinates 11,500,001-12,500,001) were generated as in (46). Gonads from animals at 24

hours after L4 were dissected on a coverslip and fixed in 1% paraformaldehyde for 5 min. A

slide (Superfrost Plus) was then placed on the coverslip and immersed in liquid N_2. The sample

was then incubated in −20°C methanol for 2 minutes and rehydrated by placing in PBST for at

least 10 minutes. Next, the sample was incubated in 0.1 M HCl for 5 minutes and washed in

PBST 3 times for 5 minutes each. The samples were then incubated for 5 minutes each in 2x

SSCT (2x saline sodium citrate with 0.1% Tween) solutions with increasing concentrations of

formamide: 0%, 5%, 10%, 25%, and 50%. The sample was then incubated in a prewarmed

42°C solution of 50% formamide in 2x SSCT for 1 hour. 2 µL of Oligopaint probe (1,000 ng/µL in

dH2O) was diluted into 30 µL of hybridization solution (50% formamide, 10% Dextran Sulfate,

2x SSC, 0.1% Tween-20) for each slide. After 1 hour incubation, slides were taken out of the

50% formamide solution, wiped, and incubated in 95% ethanol for 5 minutes. Then, the probe

hybridization solution was applied to the sample with a coverslip, and the sample was denatured

for 10 minutes at 77°C on a heat block. After denaturing, the sample was incubated with the

probe hybridization solution at 42°C overnight. The next day, samples were washed 2 times in

42°C 50% formamide in 2x SSCT for 30 minutes each, and the coverslip was removed from the

slide. Then, the sample was incubated for 5 minutes each in solutions with decreasing

concentrations of formamide in 2x SSCT: 25%, 10%, and 5%. Samples were then washed 2

times for 10 minutes each in 2x SSCT. The Oligopaint probes were visualized by hybridizing

Cy3-labelled oligos (agctgatcgtggcgttgatg) to the Oligopaint probe barcode sequence. To do

this, the Cy3-labelled probes (diluted 1:1000 in 25% ethylene carbonate in 2x SSC) were

applied to the sample with a coverslip and incubated for 15 minutes. Then, the sample was

washed and the coverslip was removed by incubating in 30% formamide solution in 2x SSCT for

3 min. The samples were then washed twice in 2x SSC and mounted in Vectashield.

For quantification of pairing between FISH signals, gonads were divided into 6 zones.

Zone 1 corresponds to the distal tip region of the gonad with only premeiotic nuclei. The gonad

region extending from the transition zone to the end of the pachytene stage was split up into 5

equally sized regions, Zones 2-6. The stitched image of the gonad was cropped into zones,

peaks of FISH signals were identified using ImageJ plugin 3D Maxima Finder (75). Each

identified peak was manually assigned to a nucleus, and distances between homologous signal

peaks in the same nucleus were calculated.

<u>Image Acquisition</u>

For spread nuclei, imaging, deconvolution, stitching and 3D-SIM reconstruction were

performed as in (73). Spreading results in squashing of *C. elegans* germline nuclei from 5 to 1-2

μm in thickness. 3D-SIM images were obtained as 125 nm spaced Z-stacks, using a 100x NA

1.40 objective on a DeltaVison OMX Blaze microscopy system, 3D-reconstructed and corrected

for registration using SoftWoRx. For display, images were projected using maximum intensity

projection in ImageJ or SoftWoRx.

For imaging of whole-mount gonads. wide field (WF) images were obtained as 200 nm

spaced Z-stacks, using a 100x NA 1.40 objective on a DeltaVison OMX Blaze microscopy

system, deconvolved and corrected for registration using SoftWoRx. Subsequently, gonads

were assembled using the "Grid/Collection" plugin (76) in ImageJ. For display, assembled

gonads were projected using maximum intensity projection in ImageJ.

For display, contrast and brightness were adjusted in individual color channels using

ImageJ.

Quantification of RAD-51 Foci and COSA-1 Foci

For quantification of RAD-51 foci in whole-mount gonads, at least three gonads were

counted per genotype. Gonads were divided into seven zones: the premeiotic zone (PM), which

includes all nuclei prior to the transition zone (where nuclei enter meiotic prophase), and six

consecutive equal-sized zones encompassing the region of the gonad from the transition zone

to the end of the pachytene stage. For the GFP::COSA-1 experiments, foci were counted in

nuclei within the last six cell rows of the gonad.

Identification of DSB Protein Foci and Object-Based Colocalization Analysis

For Figures 5 and 6, images were analyzed using an object-based colocalization

analysis pipeline that combined standard functions available in ImageJ in conjunction with a

custom Python script. A detailed description of the colocalization analysis pipeline is presented

in Supplemental Figure 3. For these analyses, 32-bit Z-stacks of SIM images of

immunofluorescence signals for at least two different antibodies detecting DSB proteins (C1 and

C2). were imported into ImageJ (77, 78) with the Fiji distribution (79). The signal maxima for

each channel, identified as foci by the image analysis pipeline were qualitatively compared to

the original image to verify accurate identification of foci.

For colocalization analysis of DSB-2 and DSB-3::GFP foci on super-spread nuclei

(Supplemental Figure S5), the same pipeline was used, except that foci were analyzed within

3.43 x 3.43 µm square ROIs located entirely within the spread (1-3 ROIs per nucleus).

Data and reagent availability

The original 32-bit individual nucleus ImageJ files, the segmented axis channel files, the

identified peaks, the values used for and the output position files from the 3D Maxima Finder for

each nucleus, the custom python script used to identify colocalization, and the resulting

spreadsheet files showing colocalization data for each nucleus will be made available from the

BioStudies Database, Accession Number S-BSST568.

Strains and primary images used in this research are available on request from A.M.V.

(annev@stanford.edu).

Figures

Figure 1. Identification of *dsb-3* as a gene required for the formation of meiotic

crossovers. (A) Representative images of DAPI-stained diakinesis-stage oocyte nuclei from

adult worms of the indicated genotypes fixed at 1 day post L4. Left: WT nucleus with six DAPI

bodies corresponding to six pairs of homologs connected by chiasmata (bivalents). Middle: *dsb-*

3(me6ts) nucleus with 9 DAPI bodies (3 bivalents and 6 univalents). Right: *dsb-3(me115)*

nucleus with 12 DAPI bodies (all univalents). Below: Graphs show quantification of the mean

number of DAPI bodies/ nucleus; error bars indicate standard deviation, and numbers in

parentheses indicate the numbers of nuclei assayed. Assays for WT and for *dsb-3(me115)*

homozygotes were performed at 20°C; assays for *dsb-3(me6ts)* homozygotes and *dsb-*

3(me6ts)/dsb-3(me115) heterozygotes were performed at 25°C. (B) Schematic showing the

dsb-3 gene structure, with the positions and nature of mutations used in this work; white boxes

represent UTR sequences, black boxes represent exons, lines indicate introns. Scale bar

indicates 100 bp. (C) Top: Whole-mount immunofluorescence images of GFP::COSA-1 foci,

which correspond to the single CO site on each homolog pair, in nuclei at the late pachytene

stage. WT nuclei have 6 GFP::COSA-1 foci, while foci are reduced or absent entirely in the *dsb-*

3(me115) mutant. Below: Stacked bar graphs showing the distribution of GFP::COSA-1 foci

counts in nuclei from WT and *dsb-3(me115)* mutants. Mean numbers of GFP::COSA-1 foci per

nucleus are indicated, with the numbers of nuclei assayed in parentheses. (D) Homolog pairing

assayed by immunofluorescence of X-Chromosome pairing center binding protein HIM-8 (Top)

or fluorescence *in situ* hybridization FISH detecting a 1Mbp segment of chromosome II (Bottom)

in pachytene nuclei of whole-mount gonads. A single focus is observed in each nucleus,

indicating successful pairing between the homologs. Scale bar is 3.2 µm. (E)

Immunofluorescence image of SC components in late pachytene nuclei in a whole-mount gonad

from the *dsb-3(me115)* mutant. Axis protein HTP-3 and SC central region protein SYP-2

colocalize in continuous stretches between chromosome pairs, indicating successful synapsis.

Scale bar is 3.2 µm.

Figure 2. DSB-3 is required for meiotic double-strand break formation. (A) Left:

Immunofluorescence images of RAD-51 foci in early pachytene nuclei in whole-mount gonads.

RAD-51 foci mark sites of processed DSBs and are strongly reduced in the *dsb-3(me115)*

mutant. Right: Quantification of RAD-51 foci in whole-mount gonads (at least three gonads were

scored per genotype). Gonads were divided into seven zones: the premeiotic zone (PM), which

includes all nuclei prior to the transition zone (where nuclei enter meiotic prophase), and six

consecutive equal-sized zones encompassing the region of the gonad from the transition zone

to the end of the pachytene stage. (B) Rescue of chiasma formation in *dsb-3* mutants by γ-

irradiation induced DNA breaks. Graph showing the average numbers of DAPI bodies present

in diakinesis-stage oocytes of worms exposed to 5 kRad of γ-irradiation at 20 hours post L4,

and un-irradiated age-matched controls, fixed and stained with DAPI 18-20 hours post

irradiation.

Figure 3. DSB-3 is concentrated in DSB-competent nuclei and is interdependent with

DSB-1 and DSB-2. (A) Immunofluorescence image of a whole-mount hermaphrodite gonad

(from distal tip to end of pachytene) stained with DAPI and antibodies detecting DSB-3::GFP,

DSB-1, and DSB-2. Meiotic progression proceeds from left to right, as in the rest of the images

for this figure. DSB-3::GFP becomes concentrated in germ cell nuclei within the transition zone

(rectangular inset - top, soon after entry into meiotic prophase. The DSB-3 immunofluorescence

signal is strongest in early pachytene nuclei, then declines sharply in mid-pachytene, albeit with

a few outlier nuclei (square inset) in the late pachytene region of the gonad retaining a strong

DSB-3 signal. This pattern of appearance and disappearance of DSB-3 from germ cell nuclei is

similar to the patterns observed for the double-strand break promoting proteins DSB-1 and

DSB-2. (B) Immunofluorescence images of a whole-mount hermaphrodite gonads stained with

DAPI and antibody detecting DSB-3::GFP. DSB-3::GFP signal is not detected in the *dsb-1(null)*

mutant, and is strongly reduced and limited to a smaller region of the gonad in the *dsb-2(null)*

mutant. (C) Immunofluorescence images of gonads stained for DSB-1 (left) or DSB-2(right),

showing that DSB-1 and DSB-2 immunofluorescence signals are not detected in the *dsb-*

3(me115) mutant. Scale bars = 16.2 µm.

Figure 4. Evidence that DSB-1, DSB-2 and DSB-3 form a complex homologous to the

yeast and mammalian REC114-MEI4 complexes. (A) Left: Schematic diagram depicting the

positions of six short signature motifs (SSMs, blue boxes) previously defined for Mei4 homologs

in diverse species, indicating the three SSMs (1, 4 and 6) that are most strongly supported in *C.*

elegans DSB-3; gray boxes indicate positions that potentially correspond to SSMs 2, 3 and 5

based on a multiple sequence alignment (MSA) with vertebrate and marine invertebrate

homologs of similar size, but are less well conserved. Right: Aligned sequences of SSMs 1, 4

and 6 from *Mus musculus, Homo sapiens, Xenopus tropicalis, Danio rerio, Strongylocentrotus*

purpuratus, Amphimedon queenslandica, Aplysia californica and the following nematodes of the

genus *Caenorhabditis*: *C. nigoni, C. briggsae, C. sinica, C. remanei, C. latens, C. tropicalis, C.*

elegans. SSMs are cropped from an MSA generated in JalView (80) using the ClustalX coloring

scheme. For additional information, see Supplemental Figure 2 and Materials and Methods. (B)

Yeast two-hybrid assay revealing protein-protein interactions among the DSB-1, -2, and -3

proteins and between DSB-1 and SPO-11. Potential interactions between proteins fused with

the GAL4 activation domain (AD) and proteins fused with the GAL4 DNA binding domain (BD)

were assayed by growth on media lacking histidine and adenine. A construct producing an N-

terminal truncation of SPO-11 lacking the first 47 amino acids (Δ1-47) was used for these

analyses, as severe auto-activation was observed for full-length SPO-11. Negative controls

showing lack of auto-activation for the constructs used are presented in Supplemental Figure 2.

In addition to experiments using constructs expressing full-length DSB-1, some experiments

used a construct expressing an N-terminally truncated DSB-1 lacking the first 33 amino acids

(Δ1-33). A schematic summarizing the identified interactions is shown on the bottom right.

Figure 5. DSB-3, DSB-2, and DSB-1 colocalize in meiotic nuclei. (A) Schematic

summarizing the object-based image analysis pipeline used to assess colocalization of foci in

SIM immunofluorescence images of partially spread nuclei (see Materials and Methods and

Supplemental Figure 3 for more details). (B) Representative SIM immunofluorescence images

of individual spread nuclei stained with antibodies targeting the indicated DSB proteins; each

panel depicts a single Z slice from a 3D image stack. The following experiments are

represented in order from left to right: mouse monoclonal HA and rabbit polyclonal DSB-2

antibodies, both detecting the same 3xHA::DSB-2 tagged protein in a *3xha::dsb-2* II strain;

chicken GFP and rabbit DSB-2 antibodies, detecting DSB-3::GFP and DSB-2 in a *meSi7[dsb-*

3::gfp] II; *dsb-3(me115)* IV strain ; mouse HA and rabbit FLAG antibodies, detecting

3xHA::DSB-1 and 3xFLAG::DSB-3 in a *3xflag::dsb-3 3xha::dsb-1* IV strain; and mouse HA and

rabbit DSB-2 antibodies, detecting 3xHA::DSB-1 and DSB-2 in a *3xflag::dsb-3 3xha::dsb-1* IV

strain. Scale bar is 2 μm. (C) Quantification of DSB protein foci for the indicated pairwise

combinations; each data point represents the numbers of foci for the two analyzed channels in a

single nucleus. Spearman R values reported indicate that numbers of the two types of foci are

strongly correlated within nuclei. (D) Graphs showing the fraction of foci of a given type

(indicated by a single colored circle in the denominator in each schematic below the horizontal

axis) that are colocalized with the other type of focus analyzed in that same experiment

(colocalizing foci are represented by two colored circles in the numerator). For each pair of

focus types analyzed, two sets of experimental analyses (represented by colored data points)

and paired negative controls (represented by grey data points) are presented.

Figure 6. The Presence and Colocalization of DSB-3, DSB-2, and DSB-1 are Not Confined

to the Meiotic Chromosomal Axis. (A) Left: representative maximum-intensity projections of

SIM images, representing the middle third of Z-stacks collected for the depicted spread nuclei.

Nuclei were stained with antibodies targeting DSB proteins and the axis component HTP-3.

Scale bar is 2 μm. Right: graph showing quantification of numbers of DSB protein foci that

colocalize with the axis signal, as well as the total numbers of DSB protein foci identified in the

analyzed nuclei. The data indicate that the majority of foci do not colocalize with the axis. (B)

Graphs depicting the fractions of Colocalized Foci / Total Foci for DSB protein foci of the

indicated types (represented as in Figure 5), reported for axis-associated foci and for total

nuclear foci analyzed within the same data sets. The filled dots represent colocalization

fractions for all DSB protein foci of a given type within the nucleus, while dots with crosses

represent the colocalization fractions for the subset of DSB protein foci of the indicated type that

colocalize with the axis.

Table 1. Quantitation of embryo viability, male frequency and diakinesis karyotypes.

Strains used in and/or created for this study were evaluated for the indicated parameters.

Numbers of DAPI bodies were evaluated for worms fixed and stained at 24 hours post L4 stage.

Analysis of AV913 and corresponding controls were conducted at two temperatures based on

the temperature-sensitive nature of the *dsb-3(me6ts)* mutant.

Supplemental Table 1. Summary of information relevant to the CRISPR/Cas9 edits

created for this work. Information includes: a) sequence of the cRNA used in the injection

mixture; b) sequence of the edit created and description of its effect on the gene and encoded

protein; c) sequences of primers used for mutation detection and verification of edits via Sanger

sequencing.

Supplemental Figure 1. Meiotic prophase progression and homolog pairing in the *dsb-*

***3(null)* mutant.** (A) Immunofluorescence images of whole-mount hermaphrodite gonads (from

distal tip to end of pachytene) stained with DAPI and antibodies detecting SUN-1 Ser24 Pi, an

indicator of CHK-2 activity detected from the onset of meiotic prophase through the early

pachytene stage (36, 72). *dsb-3(me115)* mutant germlines shown an extension of this marker

relative to WT, reflecting operation of a crossover assurance checkpoint/surveillance

mechanism that prolongs the early pachytene stage in response to one or more chromosome

pairs lacking crossover-competent recombination intermediates (20, 21, 36).   (B) Quantification

of homolog pairing assayed by FISH. FISH signals from homologous chromosomes were

considered paired if they were separated by ≤ 0.7 μm. Numbers of nuclei scored for WT: zone

1, n=180; zone 2, n=201; zone 3, n=180; zone 4, n=178; zone 5, n=180; zone 6, n=167.

Numbers of nuclei scored for *dsb-3*: zone 1, n=159; zone 2, n=180; zone 3, n=155; zone 4,

n=160; zone 5, n=143; zone 6, n=117

Supplemental Figure 2. Alignment of MEI4 and DSB-3 orthologs, and Y2H controls. (A)

Multiple sequence alignment generated with MAFFT Version 7.0 with the ClustalX coloring

scheme. Protein sequences included in the multiple sequence alignment were from the

following species: Vertebrates: *Mus musculus, Homo sapiens, Xenopus tropicalis, Danio rerio*;

marine invertebrates: *Strongylocentrotus purpuratus, Amphimedon queenslandica, Aplysia*

californica; Nematodes of the genus *Caenorhabditis*: *C. nigoni, C. briggsae, C. sinica, C.*

remanei, C. latens, C. tropicalis, C. elegans. The outlined boxes indicate the positions of the

SSMs that were previously defined for MEI4 orthologs from diverse species. Red boxes indicate

cases where the SSM includes at least 5 amino acid residues that are conserved or exhibit

similar electrophysiological properties in at least 85% of the aligned sequences. Gray boxes

indicate cases where these thresholds are not met. (B) Negative controls for yeast two-hybrid

assays, showing lack of auto-activation for cells containing the indicated constructs.

Supplemental Figure 3. 3D Object-Based Colocalization Analysis Pipeline for SIM images

of DSB protein foci in spread nuclei. (A) A schematic showing the general pipeline used for

colocalization analyses. ImageJ plugins used were 3D Maxima Finder (75) and the 3D Object

Counter (81). (B) Yellow box indicating protocol used in a subset of our analyses in which

images were additionally segmented to identify DSB protein foci that coincided with the axis

signal; the Otsu method (82) was used in our thresholding process for this segmentation.

Supplemental Figure 4. Cumulative distribution plots for the distances from each

Channel 1 DSB protein focus to its nearest neighbor Channel 2 focus. The x-axis

represents the distances between nearest neighbor foci pairs, and the y-axis indicates the

percentage of measurements at or below the given distance on the x-axis. Experimental data

are depicted in green circles and values for the corresponding negative control rotated images

are indicated with purple squares. (A) The cumulative distribution of distances between

3xHA::DSB-2 mAB foci and nearest neighbor DSB-2 pAB foci. (B) The cumulative distribution

of distances between 3xHA::DSB-1 foci and nearest neighbor 3xFLAG::DSB-3 foci. (C) The

cumulative distribution of distances between 3xHA::DSB-1 foci and nearest neighbor DSB-2

pAB foci. (D) The cumulative distribution of distances between DSB-3::GFP foci and nearest

neighbor DSB-2 pAB foci.

Supplemental Figure 5. Colocalization analysis for DSB-2 and DSB-3::GFP on super-

spread nuclei. Images depict a super-spread nucleus (top) and an inset from the same

nucleus (bottom), showing DAPI-stained DNA and immunofluorescence signals corresponding

to DSB-2, DSB-3::GFP, and axis protein HTP-3. The graph shows the fraction of DSB-2 foci

(magenta) within a given ROI that are colocalized with a DSB-3::GFP focus (green), and vice

versa, together with their paired negative controls (represented by grey data points).

Acknowledgments

We are grateful to C. Girard and S. Ramakrishnan for assistance with irradiation experiments

and generation of sequencing libraries, W. Zhang for crosses performed early in the analysis of

the *me6ts* mutant, and David Paul for discussions regarding object-based colocalization

analysis. We acknowledge N. Bhatla for creating the Exon-Intron Graphic Maker (wormweb.org)

used for Figure 1B. We thank A. Dernburg and V. Jantsch for antibodies and the Caenorhabditis

Genetics Center (funded by NIH Office of Research Infrastructure Programs P40 OD010440) for

strains. This work was supported by an American Cancer Society Research Professor Award

(RP-15-209-01-DDC) and NIH grants R01GM53804 and R35GM126964 to AMV, by a Blavatnik

Family Foundation Fellowship and a Stanford Mason Case Fellowship to AH, by an FWF Erwin

Schrödinger Fellowship (J-3676) to AW, by a Stanford Graduate Fellowship to KY, by grants

from the Taiwan Ministry of Science and Technology (107-2923-B-002-001-MY4 and MoST

104-2628-B-002-002-MY3) to PC, and by grant 1S10OD01227601 from the NCRR to the

Stanford Cell Sciences Imaging Facility.

References

1. S. Keeney, "Spo11 and the Formation of DNA Double-Strand Breaks in Meiosis" in *Recombination and Meiosis*, Genome Dynamics and Stability., R. Egel, D.-H. Lankenau, Eds. (Springer Berlin Heidelberg, 2008), pp. 81–123.

2. S. Keeney, J. Lange, N. Mohibullah, Self-Organization of Meiotic Recombination Initiation: General Principles and Molecular Pathways. *Annu. Rev. Genet.* **48**, 187–214 (2014).

3. S. Keeney, C. N. Giroux, N. Kleckner, Meiosis-Specific DNA Double-Strand Breaks Are Catalyzed by Spo11, a Member of a Widely Conserved Protein Family. *Cell* **88**, 375–384 (1997).

4. A. Bergerat, *et al.*, An atypical topoisomerase II from archaea with implications for meiotic recombination. *Nature* **386**, 414–417 (1997).

5. K. S. McKim, Meiotic Synapsis in the Absence of Recombination. *Science* **279**, 876–878 (1998).

6. A. F. Dernburg, *et al.*, Meiotic Recombination in C. elegans Initiates by a Conserved Mechanism and Is Dispensable for Homologous Chromosome Synapsis. *Cell* **94**, 387–398 (1998).

7. T. Robert, *et al.*, The TopoVIB-Like protein family is required for meiotic DNA double-strand break formation. *Science* **351**, 943–949 (2016).

8. N. Vrielynck, *et al.*, A DNA topoisomerase VI-like complex initiates meiotic recombination. *Science* **351**, 939–943 (2016).

9.　C. Claeys Bouuaert, *et al.*, Structural and functional characterization of the Spo11 core
complex. *Nat. Struct. Mol. Biol.* **28**, 92–102 (2021).

10.　T. M. Menees, G. S. Roeder, MEI4, a yeast gene required for meiotic recombination.
Genetics **123**, 675–682 (1989).

11.　R. E. Malone, D. L. Pittman, J. J. Nau, Examination of the Intron in the meiosis-specific
recombination gene REC114 in Saccharomyces. *Mol. Gen. Genet. MGG* **255**, 410–419
(1997).

12.　S. Tessé, *et al.*, Asy2/Mer2: an evolutionarily conserved mediator of meiotic recombination,
pairing, and global chromosome compaction. *Genes Dev.* **31**, 1880–1893 (2017).

13.　R. Kumar, H.-M. Bourbon, B. de Massy, Functional conservation of Mei4 for meiotic DNA
double-strand break formation from yeasts to mice. *Genes Dev.* **24**, 1266–1280 (2010).

14.　R. Kumar, *et al.*, MEI4 – a central player in the regulation of meiotic DNA double-strand
break formation in the mouse. *J. Cell Sci.* **128**, 1800–1811 (2015).

15.　R. Kumar, *et al.*, Mouse REC114 is essential for meiotic DNA double-strand break
formation and forms a complex with MEI4. *Life Sci. Alliance* **1**, e201800259 (2018).

16.　M. Molnar, *et al.*, Characterization of rec7, an early meiotic recombination gene in
Schizosaccharomyces pombe. *Genetics* **157**, 519–532 (2001).

17.　S. Bonfils, A. E. Rozalen, G. R. Smith, S. Moreno, C. Martin-Castellanos, Functional
interactions of Rec24, the fission yeast ortholog of mouse Mei4, with the meiotic
recombination-initiation complex. *J. Cell Sci.* **124**, 1328–1338 (2011).

18.　A. De Muyt, *et al.*, A high throughput genetic screen identifies new early meiotic
recombination functions in Arabidopsis thaliana. *PLoS Genet.* **5**, e1000654 (2009).

19.　A. Ronceret, M.-P. Doutriaux, I. N. Golubovskaya, W. P. Pawlowski, PHS1 regulates
meiotic recombination and homologous chromosome pairing by controlling the transport of
RAD50 to the nucleus. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 20121–20126 (2009).

20.　S. Rosu, *et al.*, The C. elegans DSB-2 Protein Reveals a Regulatory Network that Controls
Competence for Meiotic DSB Formation and Promotes Crossover Assurance. *PLoS
Genet.* **9**, e1003674 (2013).

21.　E. L. Stamper, *et al.*, Identification of DSB-1, a Protein Required for Initiation of Meiotic
Recombination in Caenorhabditis elegans, Illuminates a Crossover Assurance Checkpoint.
PLoS Genet. **9**, e1003679 (2013).

22.　J. Li, G. W. Hooker, G. S. Roeder, Saccharomyces cerevisiae Mer2, Mei4 and Rec114
Form a Complex Required for Meiotic Double-Strand Break Formation. *Genetics* **173**,
1969–1981 (2006).

23.　S. Maleki, M. J. Neale, C. Arora, K. A. Henderson, S. Keeney, Interactions between Mei4,
Rec114, and other proteins required for meiotic DNA double-strand break formation in
Saccharomyces cerevisiae. *Chromosoma* **116**, 471–486 (2007).

24. S. Steiner, J. Kohli, K. Ludin, Functional interactions among members of the meiotic
initiation complex in fission yeast. *Curr. Genet.* **56**, 237–249 (2010).

25. T. Miyoshi, *et al.*, A Central Coupler for Recombination Initiation Linking Chromosome
Architecture to S Phase Checkpoint. *Mol. Cell* **47**, 722–733 (2012).

26. M. Stanzione, *et al.*, Meiotic DNA break formation requires the unsynapsed chromosome
axis-binding protein IHO1 (CCDC36) in mice. *Nat. Cell Biol.* **18**, 1208–1220 (2016).

27. C. Claeys Bouuaert, *et al.*, DNA-driven condensation assembles the meiotic DNA break
machinery. *Nature* **592**, 144–149 (2021).

28. S. Panizza, *et al.*, Spo11-Accessory Proteins Link Double-Strand Break Sites to the
Chromosome Axis in Early Meiotic Recombination. *Cell* **146**, 372–383 (2011).

29. H. Sasanuma, *et al.*, Cdc7-dependent phosphorylation of Mer2 facilitates initiation of yeast
meiotic recombination. *Genes Dev.* **22**, 398–410 (2008).

30. J. A. Carballo, *et al.*, Budding Yeast ATM/ATR Control Meiotic Double-Strand Break (DSB)
Levels by Down-Regulating Rec114, an Essential Component of the DSB-machinery.
PLoS Genet. **9**, e1003545 (2013).

31. R. Yokoo, *et al.*, COSA-1 Reveals Robust Homeostasis and Separable Licensing and
Reinforcement Steps Governing Meiotic Crossovers. *Cell* **149**, 75–87 (2012).

32. C. M. Phillips, *et al.*, HIM-8 Binds to the X Chromosome Pairing Center and Mediates
Chromosome-Specific Meiotic Synapsis. *Cell* **123**, 1051–1063 (2005).

33. A. J. MacQueen, *et al.*, Chromosome Sites Play Dual Roles to Establish Homologous
Synapsis during Meiosis in C. elegans. *Cell* **123**, 1037–1050 (2005).

34. A. F. Severson, L. Ling, V. van Zuylen, B. J. Meyer, The axial element protein HTP-3
promotes cohesin loading and meiotic axis assembly in C. elegans to implement the
meiotic program of chromosome segregation. *Genes Dev.* **23**, 1763–1778 (2009).

35. M. P. Colaiácovo, *et al.*, Synaptonemal Complex Assembly in C. elegans Is Dispensable
for Loading Strand-Exchange Proteins but Critical for Proper Completion of
Recombination. *Dev. Cell* **5**, 463–474 (2003).

36. A. Woglar, *et al.*, Matefin/SUN-1 Phosphorylation Is Part of a Surveillance Mechanism to
Coordinate Chromosome Synapsis and Recombination with Meiotic Progression and
Chromosome Movement. *PLoS Genet.* **9**, e1003335 (2013).

37. F. Pâques, J. E. Haber, Multiple pathways of recombination induced by double-strand
breaks in Saccharomyces cerevisiae. *Microbiol. Mol. Biol. Rev. MMBR* **63**, 349–404
(1999).

38. P. Sung, Catalysis of ATP-dependent homologous DNA pairing and strand exchange by
yeast RAD51 protein. *Science* **265**, 1241–1243 (1994).

39. A. Alpi, P. Pasierbek, A. Gartner, J. Loidl, Genetic and cytological characterization of the
recombination protein RAD-51 in Caenorhabditis elegans. *Chromosoma* **112**, 6–16 (2003).

40. H.-M. Bourbon, Comparative genomics supports a deep evolutionary origin for the large,
four-module transcriptional mediator complex. *Nucleic Acids Res.* **36**, 3993–4008 (2008).

41. L. A. Kelley, S. Mezulis, C. M. Yates, M. N. Wass, M. J. E. Sternberg, The Phyre2 web
portal for protein modeling, prediction and analysis. *Nat. Protoc.* **10**, 845–858 (2015).

42. M. Boekhout, *et al.*, REC114 Partner ANKRD31 Controls Number, Timing, and Location of
Meiotic DNA Breaks. *Mol. Cell* **74**, 1053-1068.e8 (2019).

43. C. Arora, K. Kee, S. Maleki, S. Keeney, Antiviral Protein Ski8 Is a Direct Partner of Spo11
in Meiotic DNA Break Formation, Independent of Its Cytoplasmic Role in RNA Metabolism.
Mol. Cell **13**, 549–559 (2004).

44. A. Woglar, A. M. Villeneuve, Dynamic Architecture of DNA Repair Complexes and the
Synaptonemal Complex at Sites of Meiotic Recombination. *Cell* **173**, 1678-1691.e16
(2018).

45. M. G. L. Gustafsson, Nonlinear structured-illumination microscopy: Wide-field fluorescence
imaging with theoretically unlimited resolution. *Proc. Natl. Acad. Sci.* **102**, 13081–13086
(2005).

46. A. Woglar, *et al.*, Quantitative cytogenetics reveals molecular stoichiometry and
longitudinal organization of meiotic chromosome axes and loops. *PLOS Biol.* **18**, e3000817
(2020).

47. F. Couteau, HTP-1 coordinates synaptonemal complex assembly with homolog alignment
during meiosis in C. elegans. *Genes Dev.* **19**, 2744–2756 (2005).

48. E. Martinez-Perez, HTP-1-dependent constraints coordinate homolog pairing and synapsis
and promote chiasma formation during C. elegans meiosis. *Genes Dev.* **19**, 2727–2743
(2005).

49. W. Goodyer, *et al.*, HTP-3 Links DSB Formation with Homolog Pairing and Crossing Over
during C. elegans Meiosis. *Dev. Cell* **14**, 263–274 (2008).

50. Y. Kim, *et al.*, The chromosome axis controls meiotic events through a hierarchical
assembly of HORMA domain proteins. *Dev. Cell* **31**, 487–502 (2014).

51. T. Kaur, M. V. Rockman, Crossover heterogeneity in the absence of hotspots in
Caenorhabditis elegans. *Genetics* **196**, 137–148 (2014).

52. M. R. Bernstein, M. V. Rockman, Fine-Scale Crossover Rate Variation on the
Caenorhabditis elegans X Chromosome. *G3 Bethesda Md* **6**, 1767–1776 (2016).

53. F. Baudat, K. Manova, J. P. Yuen, M. Jasin, S. Keeney, Chromosome synapsis defects
and sexually dimorphic meiotic progression in mice lacking Spo11. *Mol. Cell* **6**, 989–998
(2000).

54. P. J. Romanienko, R. D. Camerini-Otero, The mouse Spo11 gene is required for meiotic
chromosome synapsis. *Mol. Cell* **6**, 975–987 (2000).

55. K. J. Hillers, V. Jantsch, E. Martinez-Perez, J. L. Yanowitz, Meiosis. *WormBook Online
Rev. C Elegans Biol.* **2017**, 1–43 (2017).

56. W. J. Swanson, V. D. Vacquier, The rapid evolution of reproductive proteins. *Nat. Rev.
Genet.* **3**, 137–144 (2002).

57. A. L. Dapper, B. A. Payseur, Molecular evolution of the meiotic recombination pathway in
mammals. *Evol. Int. J. Org. Evol.* **73**, 2368–2389 (2019).

58. S. Brenner, The genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94 (1974).

59. C. Frøkjaer-Jensen, *et al.*, Single-copy insertion of transgenes in Caenorhabditis elegans.
Nat. Genet. **40**, 1375–1383 (2008).

60. C. Frøkjær-Jensen, *et al.*, An Abundant Class of Non-coding DNA Can Prevent Stochastic
Gene Silencing in the C. elegans Germline. *Cell* **166**, 343–357 (2016).

61. C. Engler, R. Kandzia, S. Marillonnet, A One Pot, One Step, Precision Cloning Method with
High Throughput Capability. *PLoS ONE* **3**, e3647 (2008).

62. A. Paix, A. Folkmann, D. Rasoloson, G. Seydoux, High Efficiency, Homology-Directed
Genome Editing in *Caenorhabditis elegans* Using CRISPR-Cas9 Ribonucleoprotein
Complexes. *Genetics* **201**, 47–54 (2015).

63. J. A. Arribere, *et al.*, Efficient Marker-Free Recovery of Custom Genetic Modifications with
CRISPR/Cas9 in *Caenorhabditis elegans*. *Genetics* **198**, 837–846 (2014).

64. A. M. Villeneuve, A cis-acting locus that promotes crossing over between X chromosomes
in Caenorhabditis elegans. *Genetics* **136**, 887–902 (1994).

65. G. A. Auwera, *et al.*, From FastQ Data to High-Confidence Variant Calls: The Genome
Analysis Toolkit Best Practices Pipeline. *Curr. Protoc. Bioinforma.* **43** (2013).

66. M. A. DePristo, *et al.*, A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nat. Genet.* **43**, 491–498 (2011).

67. A. McKenna, *et al.*, The Genome Analysis Toolkit: A MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

68. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
357–359 (2012).

69. P. Cingolani, *et al.*, A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118;
iso-2; iso-3. *Fly (Austin)* **6**, 80–92 (2012).

70. J. B. Bessler, K. C. Reddy, M. Hayashi, J. Hodgkin, A. M. Villeneuve, A Role for
Caenorhabditis elegans Chromatin-Associated Protein HIM-17 in the Proliferation *vs* .
Meiotic Entry Decision. *Genetics* **175**, 2029–2037 (2007).

71. F. Gabler, *et al.*, Protein Sequence Analysis Using the MPI Bioinformatics Toolkit. *Curr.*
Protoc. Bioinforma. **72**, e108 (2020).

72. A. M. Penkner, *et al.*, Meiotic Chromosome Homology Search Involves Modifications of the
Nuclear Envelope Protein Matefin/SUN-1. *Cell* **139**, 920–933 (2009).

73. D. Pattabiraman, B. Roelens, A. Woglar, A. M. Villeneuve, Meiotic recombination
modulates the structure and dynamics of the synaptonemal complex during C. elegans
meiosis. *PLOS Genet.* **13**, e1006670 (2017).

74. T. Stiernagle, Maintenance of C. elegans. *WormBook* (2006)
https:/doi.org/10.1895/wormbook.1.101.1 (December 20, 2020).

75. J. Ollion, J. Cochennec, F. Loll, C. Escudé, T. Boudier, TANGO: a generic tool for high-
throughput 3D image analysis for studying nuclear organization. *Bioinformatics* **29**, 1840–
1841 (2013).

76. S. Preibisch, S. Saalfeld, P. Tomancak, Globally optimal stitching of tiled 3D microscopic
image acquisitions. *Bioinforma. Oxf. Engl.* **25**, 1463–1465 (2009).

77. C. T. Rueden, *et al.*, ImageJ2: ImageJ for the next generation of scientific image data.
BMC Bioinformatics **18**, 529 (2017).

78. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
analysis. *Nat. Methods* **9**, 671–675 (2012).

79. J. Schindelin, *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat.*
Methods **9**, 676–682 (2012).

80. A. M. Waterhouse, J. B. Procter, D. M. A. Martin, M. Clamp, G. J. Barton, Jalview Version
2--a multiple sequence alignment editor and analysis workbench. *Bioinforma. Oxf. Engl.*
25, 1189–1191 (2009).

81. S. Bolte, F. P. Cordelières, A guided tour into subcellular colocalization analysis in light
microscopy. *J. Microsc.* **224**, 213–232 (2006).

82. N. Otsu, A Threshold Selection Method from Gray-Level Histograms. *IEEE Trans. Syst.*
Man Cybern. **9**, 62–66 (1979).

1101

Strain	Genotype	Incubated Temperature	Average Brood Size ± SD (number of broods)	Percent Dead Eggs	Percent Males	Average DAPI Bodies/Nucleus ± SD (number of oocytes) 24 h post L4	48 h post L4
N2	WT	20°	264 ± 27 (7)	0.4	0	6.0 ± 0.2 (92)	5.9 ± 0.3 (102)
N2	WT	25°	291 ± 22 (7)	2	0	5.9 ± 0.3 (110)	5.9 ± 0.4 (99)
AV913	*dsb-3(me6ts) IV*	20°	274 ± 27 (7)	0.1	0	6.0 ± 0.2 (144)	5.9 ± 0.3 (111)
AV913	*dsb-3(me6ts) IV*	25°	198 ± 47 (7)	28.2	17.3	7.0 ± 1.4 (108)	9.9 ± 1.3 (99)
AV1095	*dsb-3(me115) IV*	20°	156 ± 77 (16)	99.4	24.9	11.6 ± 0.6 (150)	
AV1029	*meSi7[sun1p::dsb-3::gfp::sun1 3'UTR unc-119+] II; dsb-3(me115) IV*	20°	150 ± 23 (8)	5	3.7	6.1 ± 0.4 (207)	
AV1102	*dsb-3(me125)[3xflag::dsb-3] dsb-1(me124)[3xha::dsb-1] IV*	20°	243 ± 34 (7)	2.1	2.5	6.0 ± 0.2 (213)	
AV1115	*dsb-2(me132)[3xha::dsb-2] II*	20°	193 ± 61 (8)	2.2	0.5	6.0 ± 0.2 (150)	

dsb-1(me124) [3xha::dsb-1] IV

crRNA	CACUGGAGUGUCUGCAAUUC
Edit	ATG<u>tatccatacgatgtcccagattacgcttacccatatgacgttccagactatgcctatccatacgatgtcccagattacgct</u>TTTCCTGA**G**TT**A**CA**A**AC**G**CT**T**CA**A**TG
Description	Insertion of 3 N-terminal HA tags underlined and silent mutations introduced to the guide in bold.
Primers	TGTGAATCATTGCTCCCAAG, CCGTCAGCTTCCTGCTATTC

dsb-3(me125) [3xflag::dsb-3] IV

crRNA	GAUCGAAAUUACCGAUGAUG
Edit	ATG<u>gattataaagacgatgacgataagcgtgactacaaggacgacgacgacaagcgtgattacaaggatgacgatgacaag</u>ATCGAAATTACCGATGATGA**A**G
Description	Insertion of 3 N-terminal FLAG tags underlined and a silent PAM mutation in bold.
Primers	TTTTCCCGAAACACGATTCT, TTCGGAGTTACGACATCTGC

dsb-3(me115) IV

crRNA	GAUCGAAAUUACCGAUGAUG
Edit	ATGATCGAAATTACCGATGATGAGGACTG**A**TTCC*GGTCTC*<u>TGACTGACTGA</u>ACTGC~~T~~GTGTCTTAGCGTTTT
Description	*Leu9X* in bold: Insertion of a BsaI restriction site in italics; STOP codons in +1, +2, +3 frame underlined; and a silent PAM mutation in strikethrough.
Primers	ACACACGCCATCAAGAAAAGCA, TGTGAAGGAAACCGAGTTCCC

dsb-2(me132) [3xha::dsb-2] IV

crRNA	UGUAGUACAUCUCAACUUUC
Edit	ATG<u>tatccatacgatgtcccagattacgcttacccatatgacgttccagactatgcctatccatacgatgtcccagattacgct</u>AGTGCACGTGG**A**CT
Description	Insertion of 3 N-terminal HA tags underlined and silent PAM mutation in bold.
Primers	TGAAGGGACCTGCGCGATGTTT, ATTGCGGTGTCCAGCAGGCATC

A

WT | dsb-3(me6ts) | dsb-3(me115)

DAPI

DAPI bodies / nucleus

- WT: 6.0 (150)
- dsb-3(me6ts): 7.1 (108)
- dsb-3(me6ts)/dsb-3(me115): 8.4 (75)
- dsb-3(me115): 11.6 (150)

B

Leu9X*
me115

Leu165Phe
me6ts

C

DAPI GFP::COSA-1

WT | dsb-3(me115)

% of nuclei indicated number of COSA-1 foci

WT: x̄ = 5.8 (n = 107)
dsb-3(me115): x̄ = 0.4 (n = 164)

6, 5, 4, 3, 2, 1, 0

D

WT | dsb-3(me115)

DAPI HIM-8

DAPI FISH

E

dsb-3(me115)

HTP-3 SYP-2

A

Object-Based Colocalization using SIM images for Spread Nuclei

Negative Control: Rotate C2 by 90° and repeat analysis as below.

1. 3D SIM images with two channels (C1 & C2) corresponding to DSB protein foci. Circular ROI around the nucleus.

2. Remove signal outside of the ROI.

3. Identify local maxima inside foci above intensity threshold using ImageJ 3D Maxima Finder and obtain XYZ coordinates using ImageJ 3D Object Finder.

4. Using a python script, identify coloclized pairs of foci between C1 and C2.

B

inset

HA mAB

DSB-2 pAB

3xha::dsb-2
HAmAB DSB-2pAB

DSB-2
DSB-3::GFP

3xHA::DSB-1
3xFLAG::DSB-3

3xHA::DSB-1
DSB-2

C

R = 0.9049

R = 0.8399

R = 0.9178

R = 0.8299

D

A

B

% nuclei with paired FISH signals

WT
dsb-3

Progression through meiotic prophase

A

1. For each nucleus analyzed, a 3D Z-Stack SIM image, with two channels (C1 & C2) corresponding to DSB protein foci, was imported into ImageJ.

2. Circular ROI was drawn around the nucleus, signal outside of the ROI was cleared, and image was converted to 8-bit.

3. A duplicate C2 window was created and image was rotated by 90 degrees in XY to create a negative control image stack (C2-R) to be analyzed in conjunction with unmodified C1.

4. 3D Maxima Finder of the 3D ImageJ Suite v3.03 was used to identify local maxima within the foci in the image stack. Note in example above, the local maxima may be located in a Z-slice above or below the depicted slice.

5. 3D Object Counter v2.0 was used to extract the XYZ coordinates of the identified maxima (threshold was set to the minimum of 1 as background thresholding was addressed at a prior step). These XYZ coordinates were saved, and for each C1 focus, a custom python script was used to calculate the XY and Z distances to the nearest neighboring C2 focus (and vice versa). Negative controls distances were similarly calculated for C1 and C2-R. The script then reports the numbers and identities of C1 foci considered to be colocalized with C2 (or C2-R foci), or vice versa, based on a defined colocalization distance threshold.

B

A subset of our analyses also segmented C1 or C2 signals depending on whether they coincided with a meiotic axis signal. To accomplish this, we took the 3D Z-stack of SIM images of axis fluorescence and thresholded them (ImageJ>Image>Adjust>Threshold) using the Otsu algorithm, stack histogram setting, and black background parameters enabled. We then adjusted the resulting mask for image calculations (ImageJ>Process>Math [Divide...255]), so that pixels within the mask were assigned a value of 1 and pixels outside the mask had a value of 0. We then used the image calculator to multiply each channel (C1 or C2) with the mask/background values (ImageJ>Process>Image Calculator) to identify foci that had pixels that were included within the mask..

Parameters for 3D Maxima Finder: Radius XY and Radius Z were set to the minimum value of 1.00 pixel. The noise threshold was established by measuring the background fluorescence (in a maximum intensity projection) within an ROI, located inside the nucleus, that lacked any visible signal discernible as a focus. The maximum intensity measured within such an ROI was used as the noise parameter.

XY distance between C1 and C2 foci maxima

The custom python script compares the XYZ coordinates of foci detected in two channels from the output statistics of 3D Object Counter. It calculates the distances between nearest neighbor foci from these two channels and flags all combinations with a distance below a defined threshold that indicates colocalization.

In our analysis, we used 120nm as the XY distance threshold and 340nm as the Z distance threshold to indicate colocalization; these are the reported resolution limits of the DeltaVison OMX Blaze SIM microscopy system used in our study. These XY threshold values were corroborated independently through unsupervised empirical scoring or foci pairs as colocalized/not colocalized based on visual inspection of projected images (example above).

DAPI   DSB-3::GFP
HTP-3   DSB-2