Chromosome-scale and haplotype-resolved genome assembly of a tetraploid potato cultivar

Hequan Sun[1,2,†], Wen-Biao Jiao[1,2,3,4,†], José A. Campoy[2], Kristin Krause[1,2], Manish Goel[1,2], Kat Folz-Donahue[5], Christian Kukat[5], Bruno Huettel[6], and Korbinian Schneeberger[1,2,*]

[1]Faculty of Biology, LMU Munich, Großhaderner Str. 2, 82152 Planegg-Martinsried, Germany

[2]Department of Chromosome Biology, Max Planck Institute for Plant Breeding Research, Carl-von-Linné-Weg 10, 50829 Cologne, Germany

[3]Key Laboratory of Horticultural Plant Biology (Ministry of Education), Huazhong Agricultural University, 430070 Wuhan, China

[4]College of Informatics, Huazhong Agricultural University, 430070 Wuhan, China

[5]FACS & Imaging Core Facility, Max Planck Institute for Biology of Ageing, 50931 Cologne, Germany

[6]Max Planck-Genome-center Cologne, Carl-von-Linné-Weg 10, 50829 Cologne, Germany

[†]These authors contributed equally

[*]Corresponding author. Email: schneeberger@mpipz.mpg.de

Abstract

Potato is the most important tuber crop in the world. However, separate

reconstruction of the four haplotypes of its autotetraploid genome remained an

unsolved challenge. Here, we report the 3.1 Gb haplotype-resolved (at 99.6% precision),

chromosome-scale assembly of the potato cultivar 'Otava' using high-quality long reads

coupled with single-cell sequencing of 717 pollen genomes and Hi-C data.

Unexpectedly, almost 50% of the genome were found to be identical-by-descent due to

recent inbreeding, which contrasted by highly abundant structural rearrangements

involving around 20% of the genome. Among 38,214 genes, only 54% were present in

four haplotypes with an average of 3.2 copies per gene. Analyzing the leaf transcriptome

as example, we found that 11% of the genes featured differently expressed alleles in at

least one of the haplotypes, of which 25% are likely regulated through allele-specific

DNA methylation. Our work sheds light on the recent breeding history of potato, the

functional organization of its tetraploid genome and has the potential to strengthen the

future of genomics-assisted breeding.

Potato (*Solanum tuberosum*) is by far the most important tuber crop and is among the

five most produced crops in the world. Globally more than 350 billion kilograms of potato are

produced per year with an increasing trend particularly in developing countries in Asia[1].

Despite the social and economic importance, the breeding success of potato remained low

over the past decades due to its heterozygous, autotetraploid genome and the high levels of

inbreeding depression, which challenge usual breeding strategies commonly applied to inbred,

diploid crops[2,3].

A fundamental tool for modern breeding is the availability of reference sequences. The

reference sequence for potato was generated from a double haploid plant, *DM1-3 516 R44*

(*DM*), and was initially published in 2011[4] and continuously improved over the past years

including a recent update based on long read sequencing[5,6]. Another major advancement in

potato genomics was the recent assembly of a heterozygous diploid potato, *RH89-039-16*

(*RH*)[7]. This haplotype-resolved genome was generated from a variety of different sequencing

technologies and phase information from a genetic map derived from selfed progeny.

However, as of now, there is no haplotype-resolved assembly of a tetraploid potato

cultivar available nor there is a straightforward method that would enable the assembly of the

individual haplotypes of a tetraploid genome. The latest methods for haplotype phasing include

the separation of sequencing reads based on the differences between the parental genomes[8]

or based on haplotype information derived from gamete[9-12] or offspring genomes[7,13]. Similarly,

chromosome conformation capture sequencing (e.g., Hi-C) can help to resolve haplotypes

during or before the assembly[14-18] and has been applied to polyploids already[15-17]. However,

even though straightforward in its application, chromosome conformation capture sequencing

can lead to haplotype switch errors, and requires additional efforts such genetic maps for

correction[7,9,18].

Genome assembly of a tetraploid potato

We generated an assembly of the autotetraploid genome of *S. tuberosum* 'Otava' using

high-quality long PacBio HiFi reads (30x per haplotype) using *hifiasm*[19] (Fig. 1; Data S1; Fig.

S1-2; Methods). The initial assembly consisted of 6,366 contigs with an N50 of 2.1 Mb (Fig.

S3). While the total assembly size of 2.2 Gb was much larger than the estimated haploid

genome size of ~840 Mb, it accounted only for ~65% of the tetraploid genome size (Fig. S2)

indicating that one third of the genome collapsed during the assembly. A sequencing depth

histogram across the contigs featured four distinct peaks, which originated from regions with

either one, two, three, or four (collapsed) haplotype(s) (Fig. 1b). While most of the contigs

represented only one haplotype (referred to as *haplotigs*) and accounted for 1.5 Gb (68%) of

the assembly, contigs representing two, three or even four collapsed haplotypes (referred to

as *diplotigs*, *triplotigs* or *tetraplotigs*) still made up 470 Mb (21%), 173 Mb (8%) or 43 Mb (2%).

Regions with even higher coverages were virtually absent (9.4 Mb, 0.4%).

As there is no straight forward solution to untangle collapsed contigs after the assembly,

we restarted the genome assembly, but this time based on four separated read sets each

derived from one of the four haplotypes. In diploids, such a read separation prior to the

assembly can be performed by sorting the reads according to their similarity to the parental

genomes (trio binning)[8]. But as autotetraploid individuals inherit two haplotypes through both

the maternal and paternal lineages, this cannot be applied for autotetraploid genomes.

Alternatively, the reads can also be separated using the haplotypes found in gamete genomes

(gamete binning)[9]. While this is straightforward with haploid gametes from diploid individuals,

tetraploid potato develops diploid gametes, which again does not separate individual

haplotypes. However, as the pairing of the two haplotypes in a diploid gamete is random in

potato, we speculated that it might be possible to gain information on individual haplotypes

(and thus to separate the reads into four distinct sets) if we sequence a sufficient number of

diploid gametes.

Figure 1. Haplotype-resolved assembly of an autotetraploid potato genome. a. Assembly strategy

(gamete binning) for tetraploid genomes. Long reads are sequenced from somatic DNA and an initial

contig-level assembly is generated. In addition, sequencing data of gamete genomes are generated.

Genetic linkage enables grouping of the contig into clusters, which represent the individual haplotypes.

Long reads are assigned to haplotypes based on their similarity to the contigs. Each haplotype can be
assembled separately and scaffolded to chromosome-scale using Hi-C. (The figure was created with
help of BioRender.com.) **b.** Histogram of sequencing depth within 10 kb windows of the initial assembly
(top) revealed the presence of haplotigs (68.3%), diplotigs (21.4%), triplotigs (7.9%) and tetraplotigs
(2.0%). As a comparison, only one major peak was observed (with increased frequency) in the final
assembly (bottom). **c.** Linkage grouping. Presence/absence patterns (PAP) at each coverage marker
(50 kb region) are defined by the absence or presence of corresponding sequencing reads from each
of the pollen genomes. For instance, the PAP of contig A is "111000", where "1" refers to pollen genomes
with reads that align to the contig and "0" refers to pollen genomes without such reads. PAPs of coverage
markers A and F are highly correlated and can be grouped as {A, F}. Similarly, marker B and D are
grouped as {B, D}, while coverage marker E remains ungrouped as {E}. The diplotig coverage marker
C shows highest correlations to {A, F} and {B, D} as compared to {E}, and therefore extends these two
larger clusters. Final result are the three clusters {A, C, F}, {B, C, D} and {E}. **d**. Assembly sizes of the
haplotypes, which were highly consistent to the DM^5 and RH^7 assemblies. **e**. k-mer based evaluation of
the haplotyping accuracy. Each point represents one haplotype of one individual chromosome. The
values on the x- and y-axes indicate the numbers of k-mers within the haplotype sequence that are
unique to either of the parental genomes 'Hera' or 'Stieglitz'. Overall, 99.6% of the variation were
correctly phased.

To test if gamete binning could be applied for the genome assembly of 'Otava', we
sequenced the genomes from 717 pollen nuclei with Illumina short reads with an average
sequence coverage of 0.18x (Fig. S4) and aligned each of the 717 read sets against the initial
assembly. As defining a high-density SNP list can be difficult in a highly heterozygous
autotetraploid genome, we defined "coverage markers" (using average alignment depth in 50
kb windows) to assess if a genomic region was present in a pollen genome or not (Methods).

A coverage marker will be covered by reads if one of the two haplotypes of a pollen
carries the region of the coverage marker. With this, we could assess the presence/absence
pattern (PAP) of a coverage marker across all the 717 pollen genomes (Fig.1c). Closely linked
markers featured highly similar PAPs, as most pollen genomes carried the same pair of
haplotypes at two neighboring loci. Occasionally, recombination breakpoints, which are
integrated into the pollen genomes during meiosis, change the haplotype within a pollen
genome, and thereby slightly change the PAPs along the chromosome. However, as
recombination is generally rare, closely linked coverage markers still feature highly correlated
PAPs. We therefore could use the similarities between the PAPs to cluster the contigs into 48

groups representing the four haplotypes of all 12 chromosomes (Fig. S5-6). Haplotigs were assigned to single clusters. Diplotigs, triplotigs and tetraplotigs represented multiple haplotypes and were therefore assigned to two, three or four of the clusters (Methods).

Once the contigs were assigned to haplotypes, also the PacBio HiFi reads could be assigned to these haplotypes based on their alignments against the contigs. Reads aligned to diplotigs, triplotigs or tetraplotigs were randomly assigned to one of the respective haplotypes. With this, more than 99.9% of the non-organellar PacBio HiFi reads could be assigned to one of the 48 read sets (Fig. S7; Methods). Assembling the read sets using *hifiasm* resulted in 48 haplotype-resolved assemblies with an average N50 of 7.1 Mb and a total size of 3.1 Gb (92% of the tetraploid genome). Finally, we used Hi-C short read data (130x per haplotype) to scaffold the contigs of each assembly to a chromosome-scale, haplotype-resolved assembly (Fig. S8; Methods). Comparison of the full assembly to whole genome sequencing short reads of 'Otava' using *Merqury*[20] revealed very high base accuracy (QV > 51.7) and completeness (97.3%) of the 'Otava' genome (Methods).

The sizes of the four haplotypes of each chromosome were highly consistent to each other as well as to those of the *DM* and *RH* assemblies[4-7] except for the consistently shorter assemblies of LG10, which indicated the presence of large-scale chromosomal rearrangements between different cultivars, similar to those previously described[21] (Fig. 1d). Apart from the LG10 differences, the 'Otava' assembly was in high synteny to the DM reference sequence suggesting that also the structure of the chromosomes was assembled correctly (Fig. S9-10). To evaluate the haplotyping accuracy of the tetraploid assembly in more depth, we sequenced the parental cultivars of 'Otava', called 'Stieglitz' and 'Hera', with Illumina short reads at 10x coverage per haplotype, as each of the chromosomes was either inherited from 'Stieglitz' or from 'Hera'. Comparing the *k*-mers, which are specific to one of the parental genomes with each of the 48 chromosome assemblies, we found that each chromosome included almost exclusively *k*-mers from one but not the other parent implying a haplotyping accuracy of 99.6% (Fig. 1e; Methods).

Integrating *ab initio* predictions, protein and RNA-seq read[4-7] alignments, we annotated

152,855 gene models across four haplotypes, with an overall *BUSCO*[22] completeness score

of 97.3%, which is highly comparable to the annotations of the *RH* and *DM* assemblies[5,7](Data

S2-3; Methods). In addition, we found comparable amounts of various different types of non-

coding RNA for all haplotype genomes, which in total accounted for 33.9 Mb across the entire

genome (Data S4). Repetitive sequences made up 66% of the assembly with LTR

retrotransposons as the most abundant class and rDNA clusters of up to 600 kb in size, which

were assembled without any gaps (Data S5-6; Methods). The distribution of genes and repeats

along the chromosome followed the typical distribution of mono-centric plant genomes with

high gene and low repeat densities at the distal parts of the chromosome, while in the peri-

centromeric regions the gene densities were low and the repeat densities were high (Fig. 2).

Figure 2. The genomic features of the autotetraploid potato genome. a. Gene density (number of

genes per Mb; red) and percentage of transposable element-related sequence (yellow) within 2 Mb

windows along the four haplotypes of each of the 12 chromosomes. **b**. Landscapes of methylation in

CG (purple), CHG (blue), CHH (gray) context within 2 Mb windows. The links in the center show over

600 structural rearrangements larger than 100 kb found between the four haplotypes of each

chromosome. Light blue and red box refer to the maternally and paternally inherited chromosomes.

The genomic footprints of inbreeding

A histogram of sequence differences within 10kb windows between the haplotypes

revealed two separated peaks implying the presence of highly similar as well as highly different

regions (Fig. 3a). The divergent regions included 1 SNP per 60 bp on average, while the

remaining 50% of the regions were almost without any differences (Fig. 3a). This extreme

similarity between some of the haplotypes suggested that they were recently inherited from a

common ancestor. In fact, the pedigree of many of the cultivated potatoes, including 'Otava',

contain cultivars that occur more than once in their ancestry[23,24] (Fig. S1). Common ancestors

in different lineages of the pedigree leads to inbreeding and results in regions which are

identical-by-descent (IBD) between their haplotypes (Fig. 3b-c; Fig. S11-22; Methods). The

IBD blocks almost perfectly matched the collapsed regions in the initial assembly explaining

the high degree of unresolved regions. To exclude any potential artefacts, we screened the

IBD blocks for collapsed variation (using pseudo-heterozygous variation in read alignments

against the assembly) and found that a maximum of 1 SNP in 72 kb could have been missed

(Data S7), reassuring that highly similar IBD regions do exist in the genome.

Overall, almost 50% of the tetraploid genome of 'Otava' were included in IBD blocks

and were shared by either two, three or in rare cases even by four haplotypes (Fig. 3b-d).

Individual IBD blocks varied in size and reached up to 41.6 Mb, while IBD blocks in the peri-

centromeres were significantly larger as compared to the IBD blocks in the distal parts of the

chromosomes (Fig. S11-23). Even though it is possible that long IDB blocks were recently

introduced and were not broken up by meiotic recombination yet, it is more likely that these

extremely long IBD blocks exist due to local suppression of meiotic recombination in the peri-

centromeres (Fig. 3c). Using the accumulated mutation rates in the IBD blocks as an estimate

of their age showed that long IBD blocks weren't younger as compared to short IBD blocks

(Fig. S23b).

Extreme haplotype differences and their influence on genes

The highly similar IBD blocks were contrasted by high levels of structural

rearrangements between the non-shared regions of the genome (Fig. 3; Fig. S11-22;

Methods). Inversions, duplications, and translocations made up 3.8% to 42.9% of each of the

haplotypes (or 19.3% of the genome) depending on the abundance of IDB blocks in the

respective haplotypes. Duplications and translocations were highly enriched for Gypsy and

Copia retrotransposons near their breakpoints suggesting their dynamic contribution to

genome diversification as it was described for other plant genomes before[25], while inversions

were not enriched for transposable elements (Fig. 3d; Fig. S24; Data S8).

Excluding IBD blocks, structural rearrangements made up 15.0% to 65.8% of each

chromosome. In addition, each haplotype included 11.0% to 42.5% of unique sequence that

could not be aligned to any of the other haplotypes (Fig. 3d). This amount of structural variation

and haplotype-specific sequence was much higher than what has been reported for any other

crop species so far, supporting earlier suggestions that genomic introgressions from wild

relatives were part of the domestication history of potato[26].

Overall, we found 661 structural variations longer than 100 kb which all were supported

by the contiguity of the assembled contigs or Hi-C contact signals, including 220 duplications,

207 translocations and 234 inversions (Fig. 2; Fig. S8,11-22,25; Data S9). While comparable

in number, inversions were much larger than the other types of rearrangements and reached

sizes of up to 12.4 Mb (Fig. 3e-f). Although these large inversions were mostly located in the

peri-centromeric regions where genes occur at low density, they still harbored nearly 5% of all

genes (7,958 out of 152,855). Meiotic crossover events from the pollen genomes were virtually

absent in inversions, indicating that these regions are likely to introduce large segregating

haplotypes among cultivated potato (Fig. 3c).

Pairwise allelic divergence of the genes ranged from 0 to 140 differences per kb and

included identical as well as divergent alleles. The average pairwise difference of the divergent

alleles was 18 differences per kb (Fig. 4a) and only 53.6% of the genes were present in all four

haplotypes. The remaining 46.4% of the genes were present in three (20.0%), two (15.9%) or

even only one (10.5%) of the haplotypes (Fig. 4b) with an average of 3.2 copies per gene. In

addition, the coding sequences of some of these copies were identical to each other. For

example, only 3,066 (15.4%) of the genes with four copies also featured four distinct alleles.

In consequence, even though each gene featured 3.2 copies on average, there were only 1.9

distinct alleles per gene (Fig. 4b).

While it was expected to find identical gene alleles within the IBD blocks, only ~45% of the identical gene alleles were actually within shared regions. To test if the high number of identical alleles between the otherwise different haplotypes was indicative of selection, we tested whether these genes were enriched for specific functions. This revealed a significant enrichment for genes with GO terms involving photosynthesis, chlorophyll binding and translation (Fig. 4c) suggesting a selection-induced reduction of allelic diversity through the optimization of plant performance.

Figure 3. In-depth haplotype analysis of the tetraploid genome. a. SNP density as observed in pairwise comparisons between the haplotypes revealed two separated peaks. The high abundance of highly similar/identical regions suggested the existence of identical-by-descent (IBD) blocks. **b.** IBD blocks (minimum size: 1 Mb) across the genome. Regions shared by two, three or four haplotypes are

colored in red, orange or blue. **c.** A zoom-in on the IBD blocks and structural rearrangements of LG-4.

Large IBD blocks were more likely to occur in peri-centromeric regions with low gene, but high TE

content and suppressed meiotic recombination. (Colors as defined in **b** and **e**) **d.** Average alignment

statistics and structural rearrangements in each chromosome. **e.** Structural rearrangements between

the four haplotypes of each chromosome. **f.** Correlation of the individual size of 220 duplications, 207

translocations and 234 inversions with the respective gene density.

The low number of distinct alleles per gene and a selection-induced reduction of allelic

difference also implied that the tetraploid nature of the genome is not a necessary feature of

the high performance of potato in different environments. However, transforming potato into a

diploid suffers from the random distribution of the non-functional alleles throughout the

individual haplotypes implying that any ploidy reduction would lead to a significant gene loss.

In fact, the *BUSCO* score (indicating completeness) of the annotations of the individual

haplotypes was 89.5% on average, while the score of all four haplotypes combined was 97.3%

(Data S3) evidencing that the individual haplotypes lack genes which are present elsewhere

in the genome. Likewise, the doubled-monoploid *DM*[4,5] and the diploid *RH*[7] genomes, which

both were derived from tetraploid cultivars, feature 5,901 (15.4%) or 3,245 (8.5%), respectively,

less genes as compared to the tetraploid genome. The gene family with highest percentage of

genes with presence/absence variation (45.4%; 316 out of 696 genes) were the NLR

resistance genes (Data S10), which are known for their high intraspecies variability[27,28].

To investigate how genes are expressed in this tetraploid genome, we sequenced 367

million read pairs of the 'Otava' leaf transcriptome in three replicates (Data S1; Methods). The

four haplotype genomes contributed highly similar amounts of RNA, suggesting that none of

the haplotypes was dominant (Fig. 4d; Fig. S26), which is similar to observations in another

autoploid species[15]. Comparable to earlier analyses on the effects of CNVs on gene

expression[29], the number of allelic copies also impacted on gene expression. Genes with more

allelic copies showed a significantly increased gene expression as compared to genes with

fewer allelic copies (Fig. 4e). Although gene expression of the four haplotype genomes were

comparable at genome scale, 10.9% of the genes with four allelic copies featured significant

expression differences between the individual alleles (Fig. 4f).

To understand more about the regulation of allele-specific expression, we sequenced

the DNA methylome of 'Otava' using enzymatic methylome sequencing with three replicates,

each with 277 million read pairs (Data S1; Methods). Overall, we found that DNA methylation

was consistent across all haplotypes while DNA methylation levels in IBD blocks were slightly

higher as compared to the non-shared regions in the other haplotypes (Fig. 2; Fig. S27). Of

the 1,219 genes with significant differences in the allele-specific expression, 304 genes were

significantly correlated with the level of methylation at the up/downstream regions of these

genes (Fig. 4g), suggesting that around 25% of the allele-specific expression are regulated

through DNA methylation.

Figure 4. Impact of haplotype divergence on genes and their expression. a. Pair-wise evaluation

of allelic divergence of genes. **b.** Presence/absence variations of genes. Overall, 53.6%, 20.0%, 15.9%

or 10.5% of the genes showed four, three, two or one allelic copies/copy within the tetraploid genome

with an average of 3.2 allelic copies per gene. Different configurations of divergent alleles were observed

for the sets of genes with four, three or two copies, for instance, within the genes with four allelic copies,

one (22.3%), two (29.8%), three (32.5%) or four (15.4%) divergent allele/s were observed. Overall, this

led to an average of 1.9 distinct alleles per gene. **c.** GO enrichment analysis of genes with four identical

alleles. **d**. Four haplotypes of chromosome 1 showed comparable amounts of expressed transcripts

(FPKM: fragments per kilobase per million reads; for chromosome 2-12 see Fig. S26). **e**. Genes with

more allelic copies are significantly higher expressed than those with fewer copies ('*': *t*-test, *p*-

value<0.01). **f**. Among the four haplotypes, 10.8% (1,219) of the 11,154 genes with four functional alleles

showed allele-specific expression (Data S11). **g**. Among those, allele-specific expression of 304 genes

was correlated with DNA methylation levels, which were measured in the 1 kb up- or downstream

regions surrounding the genes (Data S12).

Discussions

Here we reported the first haplotype-resolved assembly of an autotetraploid potato.

Leveraging high-quality, long reads and single-cell genotyping of diploid gametes, we were

able to reconstruct the sequences of all four haplotypes. The sequence differences between

the haplotypes were much higher as compared to the differences commonly found within

species and were rather reminiscent of the differences found between species. This supports

earlier suggestions that the diverse haplotypes originated from introgressions from divergent

species during domestication[26].

The high level of sequence differences, however, was contrasted by widespread IBD

blocks, which were most likely introduced by related genotypes during breeding, even though

we cannot exclude that some of these blocks might have been formed via double reduction

during meiosis[30]. This similarity of the IBD blocks was the reason for the abundant collapsed

regions in the initial assembly. As these regions were almost identical, it was not possible to

assemble them from the sequence data alone. IBD blocks are a widespread phenomenon in

many crops or livestock in general, though the challenges associated with the high similarity

between haplotypes can be solved by using the power of genetics and analyzing individual

gamete genomes.

The abundance of IBD blocks, in addition to IBD-independent allele sharing, led to the

unexpected observation that the tetraploid genome only included 1.9 diverse allelic copies per

gene. This implied that the maximal allelic diversity that could be included in the tetraploid

genome was not reached, even though the high yield and yield stability of potato is supposed

to be promoted by the effects of heterosis, which itself is based on non-additive interactions of

diverse alleles[31]. Whether the high abundance of shared alleles suggests that the effects of

heterosis could still be optimized by increasing the number of polymorphic alleles or if this

indicates that the limits of heterosis were already reached remains to be seen.

Over the past years, considerable success has been made in re-domesticating potato

from a clonally-propagated, tetraploid crop into a seed-propagated, diploid crop to increase

reproduction rate, decrease costs in storage and transportation, and improve disease

control[2,32-34]. However, the random distribution of loss-of-function alleles in tetraploid potato

can lead to the accelerated manifestation of inbreeding depression in the diploid genomes,

when they are derived from tetraploids[7,35]. Haplotype-resolved assemblies of autotetraploids

like the one presented here have the potential to support the design of optimal haplotypes by

avoiding the combination of known incompatibility alleles[36]. Of course, this new possibility to

assemble autotetraploid genomes does not eliminate all breeding-related problems that result

from the tetraploid nature of potato. However, being able to reconstruct the four haplotypes of

cultivated potato is a breakthrough for modern genomics-assisted breeding strategies, and

ultimately has the power to increase the breeding success of potato in the future.

Methods

Plant material was grown at Max Planck Institute for Plant Breeding Research (Cologne, Germany). The genome of 'Otava' was sequenced with PacBio HiFi Sequel II platform with four SMRTcells. DNA extracted from individual pollen nuclei was prepared with 10x Genomics CNV kits and subsequently sequenced with Illumina sequencing. Barcodes (in single cell sequencing) were corrected using *cellranger* (10x Genomics). Short/long reads were aligned using *bowtie2*[37]/*minimap2*[38]. BAM, VCF file processing and sequencing depth analysis were performed using *samtools*[39] and *bedtools*[40]. Linkage grouping was performed using in-house developed code. PacBio sequence reads were assembled using *hifiasm*[19], and genome annotation was performed following a previous pipeline[9]. Structural variations were identified using *SyRI*[41] based on *minimap2* genome alignments. Expression analysis was performed following the literature[42]. Methylation calling was performed with *bismark* pipeline[43]. More details and other related methods are provided in the Supplementary Information.

References

1. The Food and Agriculture Organization (FAO). *http://www.fao.org/faostat/en/#data/QV* (2021).

2. S. H. Jansky et al. Reinventing potato as a diploid inbred line-based crop. *Crop Sci.* **56**, 1412-1422 (2016).

3. D.S. Douches, D. Maas, K. Jastrzebski, R.W. Chase. Assessment of Potato Breeding Progress in the USA over the Last Century. *Crop Sci*. **36**,1544-1552 (1996).

4. The Potato Genome Sequencing Consortium. Genome sequence and analysis of the tuber crop potato. *Nature* **475**, 189-195 (2011).

5. G.M. Pham, J.P. Hamilton *et al*. Construction of a chromosome-scale long-read reference genome assembly for potato. *GigaScience* **9**, 1-11 (2020).

6. C.D. Hirsch, J.P. Hamilton, K.L. Childs, J. Cepela, E. Crisovan, et al. (2014), Spud DB: A Resource for Mining Sequences, Genotypes, and Phenotypes to Accelerate Potato Breeding. *The Plant Genome* **7**, plantgenome2013.12.0042 (2014).

7. Q. Zhou, D. Tang *et al*. Haplotype-resolved genome analyses of a heterozygous diploid potato. *Nat. Genet.* **52**, 1018-1023 (2020).

8. S. Koren, A. Rhie *et al*. De novo assembly of haplotype-resolved genomes with trio binning. *Nat. Biotechnol.* **36**, 1174-1182 (2018).

9. J.A. Campoy, H. Sun *et al*. *Gamete binning*: chromosome-level and haplotype-resolved genome assembly enabled by high-throughput single-cell sequencing of gamete genomes. *Genome Biol.* **21**, 306 (2020).

10. R. Li, H. Qu *et al*. Inference of chromosome-length haplotypes using genomic data of three or a few more single gametes. *Mol Biol Evol.* **37**, 3684-3698 (2020).

11. E.F. Kirkness *et al*. Sequencing of isolated sperm cells for direct haplotyping of a human genome. *Genome Res.* **23**, 826-832 (2013).

12. D. Shi, J. Wu, H. Tang, H. Yin *et al*. Single-pollen-cell sequencing for gamete-based phased diploid genome assembly in plants. *Genome Res.* **29**, 1889-1899 (2019).

13. C. Zhou, *et al*. Assembly of whole-chromosome pseudomolecules for polyploid plant

genomes using outbred mapping populations. *Nat. Genet*. **52**, 1256-1264 (2020).

14. S. Garg, *et al.* Chromosome-scale, haplotype-resolved assembly of human genomes. *Nat.*
Biotechnol. (2020).

15. J. Zhang, X. Zhang, H. Tang, Q. Zhang *et al*. Allele-defined genome of the autopolyploid
sugarcane *Saccharum spontaneum* L. *Nat. Genet.* **50**, 1565-1573 (2018).

16. H. Chen, Y. Zeng, Y. Yang, L. Huang, B. Tang, H. Zhang. et al. Allele-aware chromosome-
level genome assembly and efficient transgene-free genome editing for the autotetraploid
cultivated alfalfa. *Nat Commun* **11**, *2494* (2020).

17. X. Zhang, S. Zhang, Q. Zhao, R. Ming, H. Tang. Assembly of allele-aware, chromosomal-
scale autopolyploid genomes based on Hi-C data. *Nat. Plants* **5**, 833-845 (2019).

18. G. Linsmith *et al*. Pseudo-chromosome-length genome assembly of a double haploid
"*Bartlett*" pear (*Pyrus communis L.*). *GigaScience* **8**, 1-17 (2019).

19. H. Cheng, G.T. Concepcion, X. Feng, H. Zhang, H. Li. Haplotype-resolved de novo
assembly with phased assembly graphs. *Nat Methods* **18**, 170-175 (2021).

20. A. Rhie, B.P. Walenz, S. Koren, A.M. Phillippy. *Merqury*: reference-free quality,
completeness, and phasing assessment for genome assemblies. *Genome Biol* **21**, 1-27
(2020).

21. L. Comai, K. Amundson, B. Ordonez, X. Zhao, G. Tomaz Braz, J. Jiang, I. Henry. LD-CNV:
rapid and simple discovery of chromosomal translocations using linkage disequilibrium
between copy number variable loci. Preprint at Biorxiv (2021).

22. F.A. Simão and R.M. Waterhouse *et al*. *BUSCO*: Assessing genome assembly and
annotation completeness with single-copy orthologs. *Bioinformatics* **31**, 3210-3212 (2015).

23. R.C.B. Hutten and R.van Berloo. An online potato pedigree database. URL:
http://www.plantbreeding.wur.nl/PotatoPedigree/ (2001).

24. R.van Berloo, R.C.B. Hutten, H.J.van Eck, and R.G.F Visser. An online potato pedigree
database resource. *Potato research* **50**, 45-57 (2007).

25. E.M. Willing, V. Rawat, T. Mandáková et al. Genome expansion of *Arabis alpina* linked with
retrotransposition and reduced symmetric DNA methylation. *Nature Plants* **1**, 14023

(2015).

26. Ma.A. Hardigan, F.P.E. Laimbeer, L. Newton, E. Crisovan, J.P. Hamilton, B. Vaillancourt et al. Genome diversity of tuber-bearing Solanum uncovers complex evolutionary history and targets of domestication in the cultivated potato. *Proc Natl Acad Sci U S A* **114**, E9999-E10008 (2017). doi: 10.1073/pnas.1714380114.

27. A.L. Van de Weyer, F. Monteiro, O.J. Furzer, et al. A Species-Wide Inventory of NLR Genes and Alleles in Arabidopsis thaliana. *Cell*. **178**, 1260-1272 (2019).

28. K. Seong, E. Seo, K. Witek, M. Li, B. Staskawicz. Evolution of NLR resistance genes with noncanonical N-terminal domains in wild tomato species. *New Phytol.* **227**, 1530-1543 (2020).

29. G.M. Pham, L. Newton, K. Wiegert-Rininger, B. Vaillancourt, D.S. Douches, C.R. Buell. Extensive genome heterogeneity leads to preferential allele expression and copy number-dependent expression in cultivated potato. *Plant J.* **92**, 624-637 (2017).

30. P.M. Bourke, R.E. Voorrips, R.G. Visser, C. Maliepaard. The double-reduction landscape in tetraploid potato as revealed by a high-density linkage map. *Genetics*, 201:853-863 (2015).

31. J. Muthoni, H. Shimelis, R. Melis. Production of hybrid potatoes: Are heterozygosity and ploidy levels important? *Australian Journal of Crop Science* **13**, 687-694 (2019).

32. P. Lindhout. *et al*. Towards F1 Hybrid Seed Potato Breeding. *Potato Res.* **54**, 301-312 (2011).

33. M. Ye, Z. Peng et al. Generation of self-compatible diploid potato by knockout of S-RNase. *Nature Plants* **4**, 651-654 (2018).

34. Y. Li, G. Li, C. Li, D. Qu., S. Huang. Prospects of diploid hybrid breeding in potato. *Chin. Potato J.* **27**, 96-99 (2013).

35. C. Zhang, P. Wang, D. Tang et al. The genetic basis of inbreeding depression in potato. *Nat Genet.* **51**, 374-378 (2019).

36. Q. Lian, D. Tang, Z. Bai, J. Qi, F. Lu, S. Huang, C. Zhang. Acquisition of deleterious mutations during potato polyploidization. *J Integr Plant Biol.* **61**, 7-11 (2019).

37. B. Langmead, S.L. Salzberg. Fast gapped-read alignment with *Bowtie* 2. *Nature methods*

9, 357-359 (2012).

38. H. Li. *Minimap2*: Pairwise alignment for nucleotide sequences. *Bioinformatics* **34**(18),

3094-100 (2018).

39. H. Li. et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-

2079 (2009).

40. A.R. Quinlan, I.M. Hall. *BEDTools*: A flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841-842 (2010).

41. M. Goel, H. Sun, W.B. Jiao, K. Schneeberger. *SyRI*: finding genomic rearrangements and

local sequence differences from whole-genome assemblies. *Genome Biol.* **20**, 1-13 (2019).

42. C.W. Law, M. Alhamdoosh, S. Su, X. Dong, L. Tian et al. RNA-seq analysis is easy as 1-

2-3 with limma, Glimma and edgeR. *F1000Research* **5**, ISCB Comm J-1408 (2016).

43. F. Krueger, S.R. Andrews. *Bismark*: a flexible aligner and methylation caller for Bisulfite-

Seq applications. *Bioinformatics* **27**, 1571-2 (2011).

Acknowledgements

The authors would like to thank Christiane Gebhardt (MPI-PZ, Cologne, Germany) and Benjamin Stich (HHU, Düsseldorf, Germany) for helpful discussions, Birgit Walkemeier and Christine Sänger (both MPI-PZ, Cologne, Germany) for plant cultivation, Christine Brandt and Klaus J. Dehmer (both IPK, Groß Lüsewitz, Germany) for providing material, Pádraic J. Flood (WUR, Wageningen, The Netherlands) for comments on the manuscript as well as Saurabh Pophaly (MPI-PZ, Cologne, Germany) for enthusiastic help in data management. We are very grateful to John Hamilton and C. Robin Buell (both University of Georgia, Georgia, USA) for integrating the 'Otava' assembly into Spud DB[6].

Funding

This work was funded by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy – EXC 2048/1– 390686111 (K.S.), and European Research Council (ERC) Grant "INTERACT" (802629) (K.S.). J.A.C was supported by Humboldt Research Fellowship for "Experienced Researchers" (Alexander von Humboldt Foundation) (J.A.C.), Marie Skłodowska-Curie Individual Fellowship PrunMut (789673).

Author contributions

H.S. and K.S. developed the project. J.A.C., H.S. and K.S. coordinated the generation of data with K.K., K.F-D., C.K. and B.H.. H.S., W-B.J., and M.G. performed all data analysis. H.S. and K.S. wrote the manuscript with input from all authors. All authors read and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Data and materials availability

High-throughput sequencing data analyzed in this project are available under NCBI BioProject: PRJNA751899. This Whole Genome Shotgun project (including assembly and annotation) has been deposited at DDBJ/ENA/GenBank under the accessions JAIVGA000000000, JAIVGB000000000, JAIVGC000000000 and JAIVGD000000000. The version described in this paper is version JAIVGA010000000, JAIVGB010000000,

JAIVGC010000000 and JAIVGD010000000. The genome assembly and gene annotation of

'Otava' are also available on Spud DB (http://spuddb.uga.edu/)[6]. Customed code and scripts

supporting this work are available at *github.com/schneeberger-lab/GameteBinning_tetraploid*.

Supplementary Information

Materials and Methods

Figs. S1 to S28

Data S1 to S12